Unbiased intestinal single cell transcriptomics reveals previously uncharacterized enteric nervous
system populations in larval zebrafish

L.E. Kuil[1,2], N. Kakiailatu[1], J.D. Windster[1,3], E. Bindels[4], J.T.M. Zink[4], G. van der Zee[1], R.M.W. Hofstra[1], I.T. Shepherd[5], V.
Melotte[1,6] and M.M. Alves[1,#]

[1]Department of Clinical Genetics, Erasmus University Medical Center, Sophia Children's Hospital, Rotterdam, The
Netherlands.
[2]Division of Psychosocial Research and Epidemiology, The Netherlands Cancer Institute, Amsterdam, The Netherlands.
[3]Department of Pediatric Surgery, Erasmus University Medical Center, Sophia Children's Hospital, Rotterdam, The
Netherlands.
[4]Department of Hematology, Erasmus MC, Rotterdam, The Netherlands.
[5]Department of Biology, Emory University, Atlanta, Georgia, United states of America.
[6]Department of Pathology, GROW-School for Oncology and Developmental Biology, Maastricht University Medical Center,
Maastricht, The Netherlands.
[#]Corresponding author: m.alves@erasmusmc.nl
15

Abstract

The enteric nervous system (ENS) regulates many gastrointestinal functions including
peristalsis, immune regulation and uptake of nutrients. Defects in the ENS can lead to severe
enteric neuropathies such as Hirschsprung disease (HSCR), which is caused by defective
ENS development. Zebrafish have proven to be fruitful in the identification of novel genes
involved in ENS development and HSCR pathology. However, the composition and
specification of enteric neurons and glial subtypes of the larval zebrafish at a single cell level,
remains mainly unexplored. Here, we performed single cell RNA sequencing of zebrafish
ENS at 5 days post-fertilization. We identified both vagal neural crest progenitors and
Schwann cell precursors, as well as four clusters of early differentiated neurons.
Interestingly, since we took an unbiased approach where we sequenced total intestines, an
elavl3+/phox2bb- population of neurons and the presence of *cx43+/phox2bb-* enteric glia
were identified in larval zebrafish. These populations have not been described before.
Pseudotime analysis supported binary neurogenic branching of ENS differentiation, which
happens via a notch-responsive state. Together, our data revealed previously unrecognized
ENS populations and serve as a resource to gain new insights on ENS development and
specification, proving that the zebrafish is a valuable model organism in the quest towards
understanding and treating congenital enteric neuropathies.

34
35   **Keywords: enteric glia; danio rerio; ENS; hirschsprung disease; HSCR**

Introduction

The enteric nervous system (ENS) consists of neurons and glia that are tightly interconnected together, and with cells in their microenvironment. The function of the ENS extends far beyond regulating peristalsis, as it is also involved in secretion, immune regulation and nutrient absorption, via connections with other cell types in the intestine (1). It is well known that dysregulation of ENS development leads to life-threatening congenital enteric neuropathies, of which Hirschsprung disease (HSCR) is the most common disorder, affecting approximately 1 in 5.000 live births (1, 2). ENS development occurs early during embryogenesis with vagal and sacral neural crest contributions. Recently, it has been found in mice that at postnatal stages, the ENS is supplemented by enteric neurons derived from Schwann cell precursors (SCPs) (3). This suggests that there is a dual origin of ENS cells, namely those derived from embryonic (vagal) neural crest during early gut colonization, and those derived postnatally from SCPs (3).

One of the vertebrate animal models that is regularly used to study ENS development, is the zebrafish (4). Zebrafish are highly suitable for genetic manipulation, develop rapidly, ex-utero and are transparent, which makes them extremely valuable for screening novel disease genes and tracing developmental processes (5, 6). However, the precise composition and specification of different neuronal and glial subtypes in the zebrafish ENS, remains unclear. This holds true particularly at larval stages, in which key processes take place to ensure proper gut colonization with enteric neurons and glia. To date, a few immunohistochemistry studies investigating enteric neuronal identities in larval zebrafish, have reported the presence of vasoactive intestinal peptide (VIP), pituitary adenylate cyclase-activating peptide (PACAP), neuronal nitric oxide synthase (nNOS), serotonin (5-hydroxytryptamine; 5HT), calretinin (CR) and calbindin (CB), from 3 days post fertilization (dpf) onwards (7, 8). Recently, it has also been described that the adult zebrafish intestine contains enteric glia, similar to that observed in mamals (9). The study showed the presence of enteric glia presenting with neurogenic properties in the adult intestine, which could be detected by the notch reporter line *her4.3:*GFP (9). However, the existence of enteric glia in larval zebrafish, is still a controversial subject. Three papers reported contradicting findings regarding expression of canonical glial genes such as *gfap*, the traditional marker for enteric glia in human and mouse. Baker *et al.* showed Gfap+ enteric glia in the outer layer of the intestine of 7 and 18 dpf fish, encapsulated by a layer of enteric neurons (10). Transmission electron microscopy showed the presence of granular vesicles and filiform processes wrapping the muscularis and caveolae, which are typical characteristic of glia (10). McCallum *et al.* also showed Gfap+ staining in the larval intestine, but suggested that the immunostaining was aspecific, since it remained in the intestine of the *ret* mutant HSCR

model, which lacks an ENS (9). Moreover, they showed that other typical enteric glial genes

were not expressed in the zebrafish intestine, including *bfabp (fabp7a), sox10* and *s100b* (9).

Such findings were supported by El-Nachef and Bronner, who reported the absence of

enteric glia expressing *sox10, gfap, plp1a* and *s100b* in larval stages (11).

To gain new insights into the exact ENS composition of larval zebrafish, studies at the

single cell transcriptome level are warranted. Previous zebrafish single cell transcriptomic

studies described, did not capture enough neuronal cells for sub-analysis, or were done at

very young embryonic and larval stages, showing limited neuronal specification (12-14).

Here, we report single cell RNA sequencing (scRNA-seq) of 5 dpf zebrafish intestines.

Importantly, we used an unbiased approach, dissecting whole intestines and sequencing all

live cells without enrichment for specific ENS markers, such as *sox10* and/or *phox2bb*. Such

approach allowed detection of previously unrecognized neuronal and glial populations in the

larval intestine, expanding our understanding of the ENS composition and specification in

larval zebrafish.

Results

Vagal derived ENS cells are complemented by Schwann cell precursors (SCPs),
supporting the dual origin of the ENS in zebrafish

To enable capturing of the ENS from 5-day-old tg(*phox2bb*:GFP) larvae (15), 244 intestines

were pooled to perform 10x scRNA-seq. Based on expression of canonical markers, as well

as markers obtained in previous literature (16-18), we selected clusters that most likely

contained neural crest progenitors, enteric neurons and glia (e.g. expression of *phox2bb,*

elval3/4, sox10, slc1a2b) (n = 1369 cells; 15% of total cells) (Fig S1). Subset analysis of

these cells, led to eleven distinct clusters (Fig 1A, 1B). Two of these clusters were

characterized by shared expression of typical neural crest markers such as, *sox10, foxd3*

and *phox2bb* and were therefore, classified as progenitor cells (Fig S2A). However, while

one cluster selectively expressed genes typical for oligodendrocyte precursor cells (OPCs) or

Schwann cell precursors (SCPs), including *clic6, tppp3,* and *anxa1a* (16, 19-22) (n = 50 cells;

Fig 1B, S2B), the other cluster showed specific expression of more traditional (vagal) neural

crest genes, such as *ret, hoxb5b, a*nd *tlx2* (23-26) (n = 181 cells; Fig 1B, S2C). *Mmp17b,*

which has been described in migrating trunk neural crest and in Schwann cells upon injury,

was specifically identified in the SCP cluster (Fig S2B) (20, 27, 28). We then performed

single molecule fluorescent whole mount in-situ hybridization (smFISH) using a probe

targeting *mmp17b*, to localize these cells in 5 dpf zebrafish and determine if they are

specifically present in the gut. As expected, positive cells were present in the spinal cord and

in the axonal motor neuron branches, corresponding to the known localization of SCPs (Fig

1C) (20, 29, 30). In the intestine, *mmp17b* signal was also observed, occasionally co-
localizing with the *tg(phox2bb:*GFP) signal (Fig 1D). This signal was sparse, which is in line
with our scRNA-seq data, where the majority of *mmp17b* positive cells (32 out of 52 cells)
showed only 1 or 2 RNA counts/cell. Therfore, our results confirm the presence of SCPs in
the zebrafish intestine, and support the rare nature of these cells at 5dpf.

The zebrafish intestine contains four types of differentiated neurons at larval stage
Based on our analysis, four types of 'differentiated neurons' were identified. The largest
cluster (n = 269 cells) consisted of inhibitory motor neurons, expressing *vip* and *nos1* (Fig
1B, 2A) (31, 32). This cluster also included cells expressing *slc6a4b*, *tph1b* and *ddc,* which
are genes involved in serotonin transport and production (Fig S3D-F) (33-35). Sensory
intrinsic primary afferent neurons (IPANs) were identified by expression of *nmu*, *vgf*, *tac3a*
and *calb2a* (n = 109 cells; Fig 1B, 2A, S3A) (32, 36). A third cluster expressing *isl2a/b*, *olig2*
and *neurod1*, most likely represents motor neurons (n = 25 cells; Fig 1B, 2A, S3B) (37-39).
The fourth cluster seems to contain a mix of different neuronal subtypes such as,
glutamatergic, GABAergic and IPANs (n = 110 cells), based on their selective expression of
vglut2a (slc17a6b), gad1b, sv2a, neurod6, gad2 and *cdh8* (Fig 2A, 2B, S3C) (32, 40, 41).
Interestingly, this latter cluster did not express the neural crest marker *phox2bb* (Fig 2B).

Identification of a *phox2bb-* population of differentiated enteric neurons
To confirm the presence of *phox2bb-* neurons in the zebrafish intestine, we first showed co-
localization of the tg(*vglut2*:loxp-dsRed-loxP-GFP) and the tg(*gad1b*:GFP) reporter lines in
the intestine (Fig 2C). Subsequently, we crossed the tg(*phox2bb*:GFP) reporter with the
tg(*vglut2*:loxp-dsRed-loxP-GFP) reporter and found no co-localization between *phox2bb* and
vglut2 in vivo, confirming the presence of *phox2bb-/vglut2a+* cells in the zebrafish intestine
(Fig 2D). Our transcriptomic data also showed that these cells express *elavl3* (encoding
HuC; Fig 2B), which led us to perform a HuC/D staining on 5 dpf tg(*phox2bb*:GFP) fish. We
observed that a limited number (between 0 to 25) of HuC/D+;*phox2bb-* cells were present in
the zebrafish intestine, comprising on average 2.5% of the total ENS (Fig 3A). This number,
roughly corresponds to our scRNA-seq data (8% of total ENS cells). Distribution of
HuC/D+;*phox2bb-* cells seems equal along the anterior to posterior axis, indicating that these
cells are evenly distributed along the total length of the intestine and therefore, do not seem
to be region specific (Fig 3A).

Larval zebrafish intestine contains enteric glia
Although recent compelling evidence suggest the presence of enteric glia in the adult
zebrafish intestine (9), previous studies are unambiguous about the existence of these cells
in larval zebrafish. This is mainly due to the absence of expression markers typical for this

cell type in mice and humans. In line with previous reports, our data confirmed the absence
of *gfap+* cells in the larval intestine of the tg(*gfap*:GFP) reporter line (Fig S4C). However, due
to our unbiased approach of sequencing total intestines, we were able to observe a cluster of
cells lacking expression of *phox2bb* and *sox10,* but expressing Hairy/E(spl)-related 4 (*her4*)
genes (n = 30 cells; Fig S4A). Interestingly, this cluster showed highly specific expression of
genes typically found in radial glia in the zebrafish brain, such as *glula, slc1a2b* and *ptn* (16),
and of genes expressed in mammalian enteric glia, such as *cx43*, *s100b, sox2, ptprz1b,* and
fabp7a (Fig 3B, S4B) (42-46). Analysis of tg(*her4:*GFP);tg(*phox2bb*:kaede) larvae showed
that *her4+;phox2bb-* cells are indeed present in the intestine and are located in close
proximity to, or in some cases in direct contact to, *phox2bb+* cells (Fig 3C). To validate the
enteric glial identity of these cells, we performed immunohistochemistry on tg(*phox2bb:*GFP)
larvae at 3, 4, 6 and 10 dpf using an antibody against connexin 43 (Cx43), a known enteric
glia marker in mice that we found expressed in the Cx43+/*phox2bb-* cluster. Based on our
results, Cx43+ cells were detected from 4 dpf onwards, suggesting that enteric glia arise
between 3 and 4 dpf in zebrafish (Fig 4A). At 4 dpf, Cx43+ cells were most often observed in
the middle intestinal segment, with an average of 12 cells per fish (Fig 4B). Although the
location of Cx43+ cells is similar to *phox2bb+* cells, as they were often observed in the same
focal plane in close proximity to each other, these cells were always negative for the
phox2bb reporter.

Progenitor cells become notch-responsive before differentiation towards neuronal and
glial fate
RNA expression of a marker for differentiated neurons, *elavl3,* and a gene specific for early
neural crest/progenitor cells, *sox10*, showed that these genes are expressed exclusively on
the left and right side of the UMAP, respectively. Analysis of key cell fate mediators for pan-
neurogenic fate (e.g. *elavl4* and *insm1b)* or genes expressed in newborn neurons in the
zebrafish brain (e.g. *tubb5* and *tmsb*), showed a similar pattern of expression on the left side
of the UMAP (Fig S5A) (16). This suggests a trajectory of progenitors' differentiation towards
neurons, from right to left in the UMAP. Pseudotime analysis by monocle3 (47) confirmed
this differentiation trajectory and additionally showed that during differentiation, a bifurcation
into two types of early differentiated neurons occurs, branch 1: sensory IPANs versus branch
2: inhibitory motor neurons (Fig 4C). This latter branch also contains a secondary branch
towards serotonergic neurons (see asterix in Fig. 4C). In line with this finding, Morarach et al.
reported a similar bifurcation in the murine ENS differentiation trajectory, with branch A,
forming Vip/Nos positive neurons (32). Expression of "branch A marker genes" *etv1* and
ebf1a, was also found in our dataset, in the *vip+/nos1+* inhibitor motor neuron branch (Fig
4C, S5C). Conservation in expression of various genes at specific differentiation states is

depicted in figure 4 panel C (32). Comparing our data to the dataset from Howard *et al.*
showed that the early differentiation observed in zebrafish from 68-70 hours post fertilization
(hpf), is prominently seen in our dataset (14). For example, co-expression of *slc18a2* and
pbx3b was observed in differentiated neurons including IPANs (Fig 4C).

Our data also showed that neuronal differentiation seems to start in notch-responsive
cells, specifically expressing *notch1a*, *notch3* and notch-responsive *her4* genes (Fig S5B)
(9). In line, our UMAP showed co-expression of the known cell fate mediator gene *sox11a*,
and   notch   (responsive)   genes   (Fig   S5B)   (16).   Live-imaging   of   5   dpf
tg(*her4*:GFP)/tg(8.3*phox2bb*:keade) fish showed that a subset of *phox2bb+* cells is indeed
her4+ (15% of cells in the posterior intestine), confirming the presence of *phox2bb+/her4+*
notch responsive cells *in vivo* (Fig 4D). Thus, progenitor cells seem to become notch
responsive upon initiation of differentiation towards neurons (branch 1 and 2). In addition,
there is a third trajectory emanating progenitors towards clusters containing proliferative
cells, motor neurons and enteric glia, suggesting a separate differentiation route towards a
cycling/enteric glial fate (Fig 4C).

Discussion

Here, we present for the first time a single cell atlas of the ENS of 5 dpf zebrafish. Our results
show the presence of two clusters of progenitor cells, traditional vagal neural crest cells and
Schwann cell precursors (SCPs), confirming a dual origin of the zebrafish ENS. Based on
our dataset, we were able to identify *mmp17b* as a specific marker for SCPs, and were able
to confirm the presence of these cells in the gut, as well as their rare nature *in vivo*. Our data,
also showed the presence of four clusters of enteric neurons and a cluster of enteric glia, at
this developmental stage. Inhibitory motor neurons were confirmed to be major contributors
of the zebrafish ENS (7, 8), but we were also able to identify other differentiated neuronal
subtypes. We can clearly see that early enteric neuronal differentiation occurs as an initial
bifurcation towards two major branches. Differentiation of sensory IPANs seem to develop
via branch 1, whereas *vip+/nos1+* inhibitory motor neurons specify via branch 2. The latter,
seems to contain a secondary branch towards serotonergic enteric neurons. Therefore,
differentiation of enteric inhibitory motor neurons as well as serotonergic neurons, seems to
be conserved between at least, mice and zebrafish (32). Interestingly, due to our unbiased
sequencing approach in which the whole intestine was analysed, we were able to identify
one   cluster   of   *elavl3+/phox2bb-*   differentiated   neurons,   expressing   genes   specific   for
glutamatergic neurons, GABAergic neurons, and others involved in serotonergic signaling.
Based on our live imaging data, *elavl3+/phox2bb-* neurons are located in close proximity or
sometimes even directly adjected, to *phox2bb+* enteric neurons. To our knowledge, such

population has never been defined before, as all enteric neurons were assumed to express
phox2bb. Future lineage tracing experiments should be performed to confirm the neural
crest-origin of these cells, as well as scRNA-seq experiments at older ages, to provide
insights into which neuronal sub-types these *phox2bb-* cells contribute.

Finally, our dataset showed the presence of a cluster of enteric glial cells. In line with
previous studies, we found that the relative contribution of enteric glial to the ENS seems to
be less abundant in zebrafish, compared to that in human and mice. We now show that
enteric glia can be detected already at zebrafish larval stages. We also confirmed that
canonical enteric glial genes such as, *sox10*, and *plp1a* are not expressed in the putative
enteric glial cluster. However, we did detect RNA expression of *s100b* and *fabp7a*, which is
in contrast to previous studies (9, 11). In addition, we showed that canonical glia in larval
zebrafish express *cx43*, *notch3* and *her* genes, but lack *phox2bb* expression. The *her4*-
reporter line previously described, has specifically characterized enteric glia in the adult
zebrafish intestine (9). However, here we show that only the *phox2bb-/her4+* cells, but not
the *phox2bb+/her4+* cells, express enteric glial markers at larval stages. Interestingly, *her4*
expression was not only limited to enteric glia, but was also consistently found in *phox2bb+*
'intermediate cells' undergoing differentiation. A previous study had already shown that
enteric neural crest cells start to express *her4* after migration, and lose its expression upon
differentiation (9). Here, we extended their findings by showing differentiation at a single cell
transcriptional level, from progenitors via a notch responsive state, towards early
specification of enteric neuronal fate. Together, this suggests that Notch signaling plays a
central role in the transition from progenitor to neuronal state or glial differentiation. In line
with this, disruption of Notch signaling in mice (*Pofut1* knockout), was shown to result in the
absence of an ENS, confirming that this signaling pathway is crucial to maintain the neural
crest progenitor pool (48). The Notch pathway has also been recognized in the maintenance
of neuronal stem cells in the brain, but signaling dynamics in neuronal differentiation have yet
to be elucidated (Reviewed by (49)).

Taken together, our results show that the zebrafish ENS has a dual origin of
precursor cells, that follow specification upon notch activation towards either a neuronal fate,
or via a cycling state towards an enteric glial fate. It also shows that using an unbiased
approach in which cells are not selected for a specific reporter construct, can be instrumental
to find new cell clusters. In summary, our data adds to the understanding of healthy ENS
development and offers an essential framework for intra-study, cross-species, and disease
state comparisons.

Acknowlegdements

We want to thank Remco Hoogenboezem for mapping the scRNA-seq data, Emma de Pater

for useful discussions, and the optical imaging center (OIC) of the Erasmus University

Medical Center for assistance with confocal microscopy.

Funding

This work was funded by the friends of Sophia foundation (SSWO WAR-63).

Figure legends

Figure 1. Single cell transcriptomics of 5 dpf zebrafish ENS

A) UMAP of 1369 ENS cells, containing eleven different clusters. B) Dot plot showing

expression of genes highly differentially expressed between clusters. C) smFISH of 5 dpf

tg(phox2bb:GFP) larvae stained for mmp17b (magenta). Scale bar represents 50 µm. D)

Zoom images of smFISH of 5 dpf tg(phox2bb:GFP) larvae showing co-localization of

mmp17b (magenta) and phox2bb (green) in the intestine, outlined with dotted lines. Arrows

highlight cells of interest showing colocalization. Scale bar represents 10 µm.

Figure 2. Cluster of elavl3+;phox2bb- enteric neurons present with excitatory and

inhibitiory gene expression

A) Featureplots of genes defining four clusters of differentiated neurons. B) Featureplots

highlighting the presence of a cluster of cells expressing elavl3, slc17a6b and gad1b, but

lacking expression of phox2bb (phox2bb- differentiated neurons, depicted by the circle). C)

Live imaging of 7 dpf tg(gad1b:GFP);tg(vglut2a-dsRed) larval intestine shows overlap

between the two reporters (arrows) with one gad1b+ cell that is vlgut2- (arrowhead). Scale

bar represents 40 µm. D) Live imaging of 7 dpf tg(phox2bb:GFP);tg(vglut2a-dsRed) larval

intestine shows no overlap between the two reporters. Scale bar represents 10 µm.

Figure 3. One small cluster expresses genes typical for enteric glia in mammalians

A) HuC/D antibody staining shows that most HuC/D+ cells in the intestine express phox2bb,

but also show phox2bb+/HuC/D- cells (progenitors) depicted by arrowheads and phox2bb-

;HuC/D+ cells (differentiated neurons) depicted by arrows. Scale bar represents 40 µm.

Quantification of the relative amount of double and single positive cells are presented in pie

charts (n=9). B) Featureplots showing selective expression of cx43, glula, slc1a2b and s100b

in one specific cluster of enteric glia depicted by the circle. C) Live-imaging of 5 dpf

tg(8.3phox2bb:kaede);tg(her4:GFP) intestines shows phox2bb-;her4+ cell depicted by the

arrowheads that are in close proximity to, or seem to interact with phox2bb+ neurons. Scale
bar represents 20 µm.
286
Figure 4. Pseudotime analysis shows differentiation trajectories from right to left of
the UMAP
A) Immunohistochemistry staining of Cx43 in the tg(phox2bb:GFP) reporter line shows non
overlapping expression in the intestine. Representative images from 4 dpf larvae. Scale bar
represents 40 µm. B) Upper graph showing the percentage of larvae that contained Cx43
cells in their proximal, middle and distal intestine. The lower graph shows the number of
Cx43 cells per larvae in the proximal, middle and distal intestine (n=19). C) Pseudotime
color-coded featureplot showing a bifurcation towards neuronal differentiation (sensory IPAN:
branch1 and inhibitory motor neurons: branch2 containing a secondary branch towards
serotonergic neurons marked with the asterix). D) Live-imaging of 5 dpf
tg(8.3phox2bb:kaede);tg(her4:GFP) intestines shows phox2bb+;her4+ cells, representing
cells undergoing differentiation from progenitor state towards neuronal or glial fate. Scale bar
represents 20 µm.
300
Supplementary figures
Supplementary figure 1. Featureplots showing the clusters selected for subset
analysis of the ENS
A) UMAP featureplots showing the cells that are selected for subset analysis of the ENS in
purple and other non-ENS cells in grey. B) Featureplots showing expression of phox2bb,
GFP, her4.1 and elavl3 in the ENS subset.
307
308 ***Supplementary figure 2. Featureplots showing specific gene expression in various***
309 ***clusters***
310 *A) UMAP featureplots showing expression of genes in the clusters containing A) precursors;*
311 *B) SCPs, as well as C) vagal neural crest cells and their derivatives.*
312
313 ***Supplementary figure 3. Featureplots showing specific expression in differentiated***
314 ***neuronal clusters***
315 *UMAP featureplots showing expression of genes in the clusters containing A) sensory*
316 *IPANs; B) motor neurons; C) phox2bb- neurons (sv2a+), and D) serotonergic neurons (in the*
317 *inhibitory motor neurons cluster).*
318
319 ***Supplementary figure 4. Featureplots showing specific expression in enteric neurons***

A) UMAP featureplots showing the absence of expression of phox2bb and sox10, but presence of her4 expression in the enteric glial cluster depicted by the circle. B) UMAP featureplots showing expression of genes in the enteric glial cluster, depicted by the circle. C) Live-imaging capture of the tg(gfap:GFP) reporter line showing the absence of GFP+ cells in the intestine, which is outlined by dotted lines. Abbreviation: AF = autofluorescence.

Supplementary figure 5. Featureplots showing expression of genes at specific differentiation states

A) UMAP featureplots showing expression of genes typical for cells committed to neuronal differentiation. B) UMAP featureplots showing expression of notch genes, notch responsive her4.2.1. and the cell fate mediator gene sox11a. C) UMAP featureplots showing expression of genes that define differentiation branches previously described (32).

Material and methods

Animal husbandry

The following zebrafish lines were used: transgenic tg(*phox2bb*:GFP)(15),
tg(*8.3phox2bb:keade*)(50), tg(*her4*:GFP)(51), tg(gfap:GFP)(52), tg(*vglut2*:loxp-dsRed-loxP-
GFP)(53), and tg(*gad1b*:GFP) (54). Zebrafish were kept on a 14/10h light/dark cycle.
Embryos and larvae were kept in an incubator at 28.5°C in HEPES-buffered E3 medium. For
imaging experiments, fish were treated from 24 hpf onwards, with 0.2 mM 1-phenyl 2-
thiourea (PTU), to inhibit pigmentation. Animal experiments were approved by the Animal
Experimentation Committee of the Erasmus MC, Rotterdam.

Isolation of zebrafish intestines
Intestines of 5 days post-fertilization (dpf) larvae were isolated as followed: a row of 6-10
larvae anesthetized with 0,0016% Tricaine, were placed on an 1.8% agarose plate.
Intestines were isolated using insect pins under a dissection microscope (Olympus SZX16),
collected with a tweezer and placed in an Eppendorf tube containing phosphate buffered
saline (PBS) with 10% fetal calf serum (FCS), on ice. In total 244 intestines were isolated
and pooled together.

Pre-processing of zebrafish cell suspension for scRNA sequencing
Cells were dissociated using 2.17mg/mL papain dissolved in HBSS, with $CaCl_2$ and $MgCl_2$.
Papain was activated using 2.5µl cysteine (1M) and dissociation was performed in a water
bath at 37°C, for 10 minutes. Cells were then transferred into a FACS tube using a 35 µm
cell strainer. Cells were centrifuged at 700g for 5 minutes at 4°C, the supernatant was
removed and pellets were resuspended in PBS containing 10% FCS. DAPI was added to
mark dead cells (1:1000). All sorts were performed using the FACSAria III sorter, into
eppendorfs containing PBS with 5% FCS.

Single cell RNA sequencing (scRNA-seq)
Single cells were barcoded using a 10x genomics Chromium Controller, and sequenced
using a Novaseq 6000 instrument (Illumina). In total, 9.858 cells were sequenced with mean
reads per cell of 21.106. For scRNA-seq analysis we used Seurat V3 (55). The Seurat
pipeline was used for filtering (nFeature_RNA > 100 & nFeature_RNA <4200 & percent.mito
< 0.05), normalization and downstream analysis for clustering, where we used 50
dimensions with a resolution of 0.8 for the UMAP processing. This led to 49 clusters, which
we annotated based on differential gene expression. Seven clusters expressing neuronal
and/or enteric progenitor markers were identified (cluster 6, 10, 13, 19, 21, 23, 34)(Fig. S1).

These seven clusters were selected for a subset analysis, using 30 dimensions with a
resolution of 0.7 for the clustering and UMAP, resulting in 14 clusters. One cluster was
excluded as it contained leukocytes (*lcp1+, phox2bb-, elavl3-, sox10-*; cluster 12) and
proliferating cells negative for *phox2bb* and *elavl3* were manually excluded. The final set was
analyzed using six dimensions and a resolution of 0.4. This analysis provided us with eleven
clusters, which were annotated based on the differential gene expression and literature
search. If required, a more thorough analysis of the total set of differentially expressed genes
per cluster, or differentially expressed genes between specific subclusters, was performed.
For pseudotime analysis, monocle3 was used (47).
380

Fluorescent imaging
Imaging was performed as previously described (56). For the keade photoconversion
experiments, all *phox2bb+* cells in the total intestines were photoconverted using the 405
nanometer (nm) laser, as described earlier (11). After photoconversion, the green and red
channels were recorded, using a sequential scan with the 488nm and 561nm lasers to
confirm full photoconversion.
387

Immunohistochemistry
Whole mount immunohistochemistry using mouse anti-HuC/D (1:100, molecular probes A-
21271) was performed, as previously described (57). Antibody staining using rabbit anti-
Cx43 (1:200, Cell Signaling Technologies 83649) was performed as published before (58).
To increase signal to noise ratio we made use of monovalent AffiniPure Fab Fragments (111-
167-003 Jackson; Cy™3 AffiniPure Fab Fragment Goat Anti-Rabbit IgG).

Single-molecule whole-mount fluorescent in situ hybridization
Zebrafish were fixed in 4% paraformaldehyde in PBS, overnight. They were then dehydrated
through a series of 25/50/75/100% MeOH in PBST for 5 minutes each, and stored for a
minimum of 1 hour at -20°C. Next, samples were rehydrated through a series of 75/50/25/0%
MeOH in PBST for 5 minutes each, and incubated in prot K for 15 minutes at 20 $^{\circ}$C. They
were rinsed twice with PBST for 5 minutes and re-fixed in 4% PFA in PBS, for 20 minutes at
room temperature. Susequently, samples were rinsed again 5x5 minutes in PBST. After
manual pre-treatment for permeabilization, we continued with the RNAscope Multiplex
Fluorescence Reagent Kit v2 Assay (Advanced Cell Diagnostics, Bio-Techne), according to
the manufacturers' instructions. A custom made probe for dr-*mmp17b* C1 (NPR-0035110)
was used (Advanced Cell Diagnostics, Bio-Techne). Opal 570 dye (Akoya Biosciences) was
used for channel development.
406

References

1.   Niesler B, Kuerten S, Demir IE, Schafer KH. Disorders of the enteric nervous system - a holistic
view. Nat Rev Gastroenterol Hepatol. 2021;18(6):393-410.
2.   Duess JW, Hofmann AD, Puri P. Prevalence of Hirschsprung's disease in premature infants: a
systematic review. Pediatr Surg Int. 2014;30(8):791-5.
3.   Uesaka T, Nagashimada M, Enomoto H. Neuronal Differentiation in Schwann Cell Lineage
Underlies Postnatal Neurogenesis in the Enteric Nervous System. J Neurosci. 2015;35(27):9879-88.
4.   Kuil LE, Chauhan RK, Cheng WW, Hofstra RMW, Alves MM. Zebrafish: A Model Organism for
Studying Enteric Nervous System Development and Disease. Frontiers in Cell and Developmental
Biology. 2021;8(1874).
5.   Kuil L, MacKenzie KC, Tang C, Windster J, Linh Le T, Karim A, et al. Size matters: large copy
number losses reveal novel Hirschsprung disease genes. medRxiv. 2020:2020.11.02.20221481.
6.   Davidson AE, Straquadine NR, Cook SA, Liu CG, Ganz J. A rapid F0 CRISPR screen in zebrafish
to identify regulators of neuronal development in the enteric nervous system. bioRxiv.
2021:2021.07.17.452230.
7.   Uyttebroek L, Shepherd IT, Harrisson F, Hubens G, Blust R, Timmermans J-P, et al.
Neurochemical coding of enteric neurons in adult and embryonic zebrafish (Danio rerio). Journal of
Comparative Neurology. 2010;518(21):4419-38.
8.   Olsson C, Holmberg A, Holmgren S. Development of enteric and vagal innervation of the
zebrafish (Danio rerio) gut. The Journal of comparative neurology. 2008;508(5):756-70.
9.   McCallum S, Obata Y, Fourli E, Boeing S, Peddie CJ, Xu Q, et al. Enteric glia as a source of
neural progenitors in adult zebrafish. Elife. 2020;9.
10.   Baker PA, Meyer MD, Tsang A, Uribe RA. Immunohistochemical and ultrastructural analysis of
the maturing larval zebrafish enteric nervous system reveals the formation of a neuropil pattern. Sci
Rep. 2019;9(1):6941.
11.   El-Nachef WN, Bronner ME. De novo enteric neurogenesis in post-embryonic zebrafish from
Schwann cell precursors rather than resident cell types. Development. 2020;147(13).
12.   Hayot G, Massonot M, Keime C, Faure E, Golzio C. CHD8-associated
gastrointestinal complaints are caused by impaired vagal neural crest development and homeostatic
imbalance. bioRxiv. 2021:2021.10.06.463249.
13.   Willms RJ, Jones LO, Hocking JC, Foley E. A cell atlas of microbe-responsive processes in the
zebrafish intestine. Cell Rep. 2022;38(5):110311.
14.   Howard AGt, Baker PA, Ibarra-Garcia-Padilla R, Moore JA, Rivas LJ, Tallman JJ, et al. An atlas
of neural crest lineages along the posterior developing zebrafish at single-cell resolution. Elife.
2021;10.
15.   Nechiporuk A, Linbo T, Poss KD, Raible DW. Specification of epibranchial placodes in
zebrafish. Development. 2007;134(3):611-23.
16.   Lange C, Rost F, Machate A, Reinhardt S, Lesche M, Weber A, et al. Single cell sequencing of
radial glia progeny reveals the diversity of newborn neurons in the adult zebrafish brain.
Development. 2020;147(1).
17.   Roy-Carson S, Natukunda K, Chou HC, Pal N, Farris C, Schneider SQ, et al. Defining the
transcriptomic landscape of the developing enteric nervous system and its cellular environment.
BMC Genomics. 2017;18(1):290.
18.   Elworthy S, Pinto JP, Pettifer A, Cancela ML, Kelsh RN. Phox2b function in the enteric nervous
system is conserved in zebrafish and is sox10-dependent. Mech Dev. 2005;122(5):659-69.
19.   Xia W, Zhu J, Wang X, Tang Y, Zhou P, Hou M, et al. ANXA1 directs Schwann cells proliferation
and migration to accelerate nerve regeneration through the FPR2/AMPK pathway. FASEB J.
2020;34(10):13993-4005.

20.   Leigh NR, Schupp MO, Li K, Padmanabhan V, Gastonguay A, Wang L, et al. Mmp17b is
essential for proper neural crest cell migration in vivo. PLoS One. 2013;8(10):e76484.
21.   Siems SB, Jahn O, Eichel MA, Kannaiyan N, Wu LMN, Sherman DL, et al. Proteome profile of
peripheral myelin in healthy mice and in a neuropathy model. Elife. 2020;9.
22.   Pooyan P, Karamzadeh R, Mirzaei M, Meyfour A, Amirkhan A, Wu Y, et al. The Dynamic
Proteome of Oligodendrocyte Lineage Differentiation Features Planar Cell Polarity and
Macroautophagy Pathways. Gigascience. 2020;9(11).
23.   Lui VC, Cheng WW, Leon TY, Lau DK, Garcia-Barcelo MM, Miao XP, et al. Perturbation of
hoxb5 signaling in vagal neural crests down-regulates ret leading to intestinal hypoganglionosis in
mice. Gastroenterology. 2008;134(4):1104-15.
24.   Howard AGAt, Nguyen AC, Tworig J, Ravisankar P, Singleton EW, Li C, et al. Elevated Hoxb5b
Expands Vagal Neural Crest Pool and Blocks Enteric Neuronal Development in Zebrafish. Front Cell
Dev Biol. 2021;9:803370.
25.   Borghini S, Bachetti T, Fava M, Di Duca M, Cargnin F, Fornasari D, et al. The TLX2 homeobox
gene is a transcriptional target of PHOX2B in neural-crest-derived cells. Biochem J. 2006;395(2):355-
61.
26.   Borghini S, Di Duca M, Santamaria G, Vargiolu M, Bachetti T, Cargnin F, et al. Transcriptional
regulation of TLX2 and impaired intestinal innervation: possible role of the PHOX2A and PHOX2B
genes. Eur J Hum Genet. 2007;15(8):848-55.
27.   Prendergast A, Linbo TH, Swarts T, Ungos JM, McGraw HF, Krispin S, et al. The
metalloproteinase inhibitor Reck is essential for zebrafish DRG development. Development.
2012;139(6):1141-52.
28.   Arthur-Farraj P, Coleman MP. Lessons from Injury: How Nerve Injury Studies Reveal Basic
Biological Mechanisms and Therapeutic Opportunities for Peripheral Nerve Diseases.
Neurotherapeutics. 2021;18(4):2200-21.
29.   Rosenberg A, Isaacman-Beck J, Franzini-Armstrong C, Granato M. Schwann Cells and Deleted
in Colorectal Carcinoma Direct Regenerating Motor Axons Towards Their Original Path. The Journal
of neuroscience : the official journal of the Society for Neuroscience. 2014;34:14668-81.
30.   Perlin JR, Lush ME, Stephens WZ, Piotrowski T, Talbot WS. Neuronal Neuregulin 1 type III
directs Schwann cell migration. Development. 2011;138(21):4639-48.
31.   Keef KD, Shuttleworth CW, Xue C, Bayguinov O, Publicover NG, Sanders KM. Relationship
between nitric oxide and vasoactive intestinal polypeptide in enteric inhibitory neurotransmission.
Neuropharmacology. 1994;33(11):1303-14.
32.   Morarach K, Mikhailova A, Knoflach V, Memic F, Kumar R, Li W, et al. Diversification of
molecularly defined myenteric neuron classes revealed by single-cell RNA sequencing. Nat Neurosci.
2021;24(1):34-46.
33.   Del Colle A, Israelyan N, Gross Margolis K. Novel aspects of enteric serotonergic signaling in
health and brain-gut disease. Am J Physiol Gastrointest Liver Physiol. 2020;318(1):G130-G43.
34.   Shih DF, Hsiao CD, Min MY, Lai WS, Yang CW, Lee WT, et al. Aromatic L-amino acid
decarboxylase (AADC) is crucial for brain development and motor functions. PLoS One.
2013;8(8):e71741.
35.   Ren J, Isakova A, Friedmann D, Zeng J, Grutzner SM, Pun A, et al. Single-cell transcriptomes
and whole-brain projections of serotonin neurons in the mouse dorsal and median raphe nuclei.
Elife. 2019;8.
36.   Delvalle NM, Dharshika C, Morales-Soto W, Fried DE, Gaudette L, Gulbransen BD.
Communication Between Enteric Neurons, Glia, and Nociceptors Underlies the Effects of Tachykinins
on Neuroinflammation. Cell Mol Gastroenterol Hepatol. 2018;6(3):321-44.
37.   Catala M, Kubis N. Gross anatomy and development of the peripheral nervous system. Handb
Clin Neurol. 2013;115:29-41.
38.   Hutchinson SA, Eisen JS. Islet1 and Islet2 have equivalent abilities to promote motoneuron
formation and to specify motoneuron subtype identity. Development. 2006;133(11):2137-47.

39.   Memic F, Knoflach V, Sadler R, Tegerstedt G, Sundstrom E, Guillemot F, et al. Ascl1 Is
Required for the Development of Specific Neuronal Subtypes in the Enteric Nervous System. J
Neurosci. 2016;36(15):4339-50.
40.   Tong Q, Ma J, Kirchgessner AL. Vesicular glutamate transporter 2 in the brain-gut axis.
Neuroreport. 2001;12(18):3929-34.
41.   Wright CM, Garifallou JP, Schneider S, Mentch HL, Kothakapa DR, Maguire BA, et al. Dlx1/2
mice have abnormal enteric nervous system function. JCI Insight. 2020;5(4).
42.   Grubisic V, McClain JL, Fried DE, Grants I, Rajasekhar P, Csizmadia E, et al. Enteric Glia
Modulate Macrophage Phenotype and Visceral Sensitivity following Inflammation. Cell Rep.
2020;32(10):108100.
43.   McClain J, Grubisic V, Fried D, Gomez-Suarez RA, Leinninger GM, Sevigny J, et al. Ca2+
responses in enteric glia are mediated by connexin-43 hemichannels and modulate colonic transit in
mice. Gastroenterology. 2014;146(2):497-507 e1.
44.   Rao M, Gershon MD. The bowel and beyond: the enteric nervous system in neurological
disorders. Nat Rev Gastroenterol Hepatol. 2016;13(9):517-28.
45.   Young HM, Bergner AJ, Muller T. Acquisition of neuronal and glial markers by neural crest-
derived cells in the mouse intestine. The Journal of comparative neurology. 2003;456(1):1-11.
46.   Heanue TA, Pachnis V. Prospective identification and isolation of enteric nervous system
progenitors using Sox2. Stem Cells. 2011;29(1):128-40.
47.   Trapnell C, Cacchiarelli D, Grimsby J, Pokharel P, Li S, Morse M, et al. The dynamics and
regulators of cell fate decisions are revealed by pseudotemporal ordering of single cells. Nature
Biotechnology. 2014;32(4):381-6.
48.   Okamura Y, Saga Y. Notch signaling is required for the maintenance of enteric neural crest
progenitors. Development. 2008;135(21):3555-65.
49.   Moore RE, Clarke J, Alexandre P. Protrusion-Mediated Signaling Regulates Patterning of the
Developing Nervous System. Front Cell Dev Biol. 2020;8:579073.
50.   Harrison C, Wabbersen T, Shepherd IT. In vivo visualization of the development of the enteric
nervous system using a Tg(-8.3bphox2b:Kaede) transgenic zebrafish. Genesis. 2014;52(12):985-90.
51.   Sang-Yeob Y, MinJung K, Hyung-Seok K, Tae-Lin H, Ajay BC. Fluorescent protein expression
driven by her4 regulatory elements reveals the spatiotemporal pattern of Notch signaling in the
nervous system of zebrafish embryos. Developmental Biology. 2007;301(2):555-67.
52.   Raymond PA, Barthel LK, Bernardos RL, Perkowski JJ. Molecular characterization of retinal
stem cells and their niches in adult zebrafish. BMC Developmental Biology. 2006;6(1):36.
53.   Miyasaka N, Morimoto K, Tsubokawa T, Higashijima S-i, Okamoto H, Yoshihara Y. From the
Olfactory Bulb to Higher Brain Centers: Genetic Visualization of Secondary Olfactory Pathways in
Zebrafish. The Journal of Neuroscience. 2009;29(15):4756-67.
54.   Satou C, Kimura Y, Hirata H, Suster ML, Kawakami K, Higashijima S-i. Transgenic tools to
characterize neuronal properties of discrete populations of zebrafish neurons. Development.
2013;140(18):3927-31.
55.   Stuart T, Butler A, Hoffman P, Hafemeister C, Papalexi E, Mauck WM, III, et al.
Comprehensive Integration of Single-Cell Data. Cell. 2019;177(7):1888-902.e21.
56.   Kuil LE, Oosterhof N, Ferrero G, Mikulášová T, Hason M, Dekker J, et al. Zebrafish
macrophage developmental arrest underlies depletion of microglia and reveals Csf1r-independent
metaphocytes. eLife. 2020;9:e53403.
57.   Heanue TA, Boesmans W, Bell DM, Kawakami K, Vanden Berghe P, Pachnis V. A Novel
Zebrafish ret Heterozygous Model of Hirschsprung Disease Identifies a Functional Role for mapk10 as
a Modifier of Enteric Nervous System Phenotype Severity. PLoS Genet. 2016;12(11):e1006439.
58.   Hyland C, Mfarej M, Hiotis G, Lancaster S, Novak N, Iovine MK, et al. Impaired Cx43 gap
junction endocytosis causes morphological and functional defects in zebrafish. Mol Biol Cell.
2021;32(20):ar13.

A

nos1   *nmu*   *isl2a*   *cdh8*

B

elavl3   *phox2bb*   *slc17a6b*   *gad1b*

C

tg(*gad1b*-GFP)   tg(*vglut2a*-dsRED)

D

tg(*phox2bb*:GFP)   tg(*vglut2a*-dsRED)

A

tg(*phox2bb*:GFP) | HuC/D (Elavl3/4)

Proximal | Middle | Distal

■ % HuC/D+   ■ % *phox2bb*+   ■ % HuC/D+/*phox2bb*+

B

slc1a2b | *s100b* | *cx43* | *glula*

0 1 2 3 4 5 | 0 1 2 3 | 0 1 2 3 4 | 0 1 2 3 4 5

C

merged

tg(*her4*:GFP)

tg(*phox2bb*:kaede)