Vertical growth dynamics of biofilms

Pablo Bravo[1,3*]**, Siu Lung Ng**[2]**, Kathryn A. MacGillivray**[2,3]**, Brian K. Hammer**[2]**, Peter J. Yunker**[1*]

*For correspondence:
pbravo7@gatech.edu (PB);
peter.yunker@gatech.edu (PJY)

[1]School of Physics, Georgia Institute of Technology, Atlanta, Georgia, USA; [2]School of Biological Sciences, Georgia Institute of Technology, Atlanta, Georgia, USA; [3]Interdisciplinary program in Quantitative Biosciences, Georgia Institute of Technology, Atlanta, Georgia, USA

Abstract   During the biofilm life cycle, bacteria attach to a surface then reproduce, forming crowded, growing communities. As these colonies develop, they expand horizontally and vertically. This horizontal expansion, known as the range expansion, occurs at a constant rate, which is often used as a proxy for bacterial fitness. Conversely, the vertical growth of biofilms is much less studied, despite representing a fundamental aspect of bacterial physiology. Many theoretical models of vertical growth dynamics have been proposed; however, difficulties in measuring biofilm height accurately across relevant time and length scales have prevented testing these models or their biophysical underpinnings empirically. Using white light interferometry, we measure the heights of microbial colonies with nanometer precision from inoculation (sub-micron) to their final equilibrium height (hundreds of microns), producing a novel and detailed empirical characterization of vertical growth dynamics. We show that models relying on logistic growth or nutrient depletion fail to capture biofilm height dynamics on short and long time scales. Our empirical results support a simple model inspired by the fact that biofilms only interact with the environment through their interfaces. This interface model captures the growth dynamics from short to long time scales (10 minutes to 14 days) of diverse microorganisms, including prokaryotes like gram-negative and gram-positive bacteria and eukaryotes like aerobic and anaerobic yeast. This model provides heuristic value, highlighting the biophysical constraints that limit vertical growth as well as establishing a quantitative model for biofilm development.

Introduction

Biofilms are surface attached microbial communities composed of cells and extracellular polymeric substance (EPS) (*Watnick and Kolter, 2000*; *Flemming et al., 2016*; *Hall-Stoodley et al., 2004*). Biofilms form when a cell attaches to a surface and reproduces, leading to horizontal and vertical expansion. Understanding the dynamics of this expansion, and the processes responsible for them, is fundamental to understanding biofilm development(*Lambert et al., 2014*; *Nadell et al., 2016*; *Wille and Coenye, 2020*). Many studies have focused on understanding the horizontal "range expansion," detailing the impact of physical interactions, the environment, inter-strain competitive dynamics, and more(*D'Acunto et al., 2015*; *Ghosh and Levine, 2017*; *Wang et al., 2017*; *Echten et al., 2020*; *Fei et al., 2020*; *Xiong et al., 2020*; *Fortune et al., 2022*; *Tam et al., 2022*; *Moore-Ott et al., 2022*). Conversely, less is known about the vertical growth of biofilms, despite its importance for determining access to nutrients and oxygen(*De Beer et al., 1994*; *Stewart et al., 2016*). The initial steps of this process (*Yan et al., 2016*; *Beroz et al., 2018*), as well as some aspects of competitive growth(*Oliveira et al., 2015*; *Rendueles and Ghigo, 2015*; *Coyte et al., 2017*; *Steinbach et al., 2020*), have been studied. However, our knowledge of how vertical growth proceeds over short and long

time scales is limited. In particular, while growth curves in liquid media have been studied for years building on the works of Jacques Monod(*Monod, 2012*; *Allen and Waclaw, 2019*), and the horizontal expansion of biofilms on agar plates is known to proceed at a constant rate(*Cooper et al., 1968*; *Warren et al., 2019*; *Basaran et al., 2022*), we lack detailed measurements and an heuristic understanding of vertical growth. As biofilms play important roles in microbial ecology and human health(*Davey and O'toole, 2000*; *Donlan, 2001*), elucidating this fundamental aspect of microbial physiology is crucial.

The characterization of microbial topographies is a growing field(*Dervaux et al., 2014*; *Larimer et al., 2016*; *Kesel et al., 2017*; *Werb et al., 2017*; *Schiebel et al., 2020*; *Paula et al., 2020*; *Hayta et al., 2021*; *Mahto and Das, 2021*), but with limited focus on the temporal dynamics of the colony. The lack of clarity about vertical growth dynamics is due in part to the experimental difficulty in measuring the height of a biofilm with sufficient precision over many different time scales in a non-destructive manner(*Huang et al., 2020*). Diffraction-limited optical resolution precludes precise measurement of biofilm height at short times. This issue is further confounded by the fact that fluorescent proteins and dyes are specific or localized, and thus do not necessarily report the true height of a biofilm, which is composed of both cells and extracellular matrix. Fluorescent stains can also bleach over time, making it difficult to continuously image the biofilm and the surface it sits on, which is necessary for assessing height. Thus, we lack a clear empirical picture of how vertical growth dynamics proceed over short and long time scales.

Due to the many relevant biological and physical parameters(*Mazza, 2016*; *Ghosh and Levine, 2017*; *Mattei et al., 2018*; *Nguyen et al., 2021*) involved in biofilm modeling, many models are complex, with multiple interacting phases inside and outside of the colony. Often, these models will incorporate a quadratic carrying capacity term to capture the qualitative features of vertical growth dynamics that are initially rapid before eventually saturating. However, while the logistic growth curve captures these qualitative features, it is unclear if logistic growth is quantitatively accurate. To address this issue, many multi-phase, spatial models have been developed (*Alpkvist et al., 2006*; *Zhang et al., 2008*; *Hartmann et al., 2019*; *Warren et al., 2019*; *Srinivasan et al., 2019*; *Fei et al., 2020*; *Jin et al., 2020*), with a smaller subset of these models explicitly addressing vertical growth, or colony thickness (*Wang et al., 2017*; *Tam et al., 2019*; *Hughes, 2020*). Even if these models are highly accurate, reliance on many precisely measured parameters limits the practical and heuristic utility of such models. Thus, developing a functional and quantitative understanding of vertical biofilm growth with reduced dimensionality would go a long way towards elucidating this fundamental aspect of microbial physiology.

Here, we measure the vertical dynamics of growing biofilms. Using white light interferometry, we obtain height topographic profiles with single nanometer resolution. We perform frequent measurements for a period of 48 hours, and also test the long term behavior over two weeks. We observe that carrying capacity and nutrient depletion models poorly fit, and are qualitatively inconsistent with, our empirical measurements. Guided by our empirical results, we test a simple interface-limited growth model, which accounts for the fact that diffusion limits the thickness of the growing zone inside the colony. With just three parameters, this model accurately reproduces the observed asymmetric vertical growth dynamics, and captures steady-state biofilm heights at 14 days.

Results

Using a Zygo Zegage Pro optical profilometer, we measure the growth of multiple microbial colonies, from an initial height of less than a micrometer to large colonies of hundreds of micrometers after 48 hours of growth (*Figure 1*A-C). These measurements allow us to reconstruct the interface of a growing microbial colony with high resolution across short and long time scales, both in the lateral($0.177\mu m$) and vertical direction ($1nm$) (*Figure 1*D). We define colony height h as the mean height in a $2mm$ long, $34.6\mu m$ wide strip around the colony center (*Figure 1* D). Focusing on the center of the colony ensures that we isolate the vertical growth dynamics from the effects of lateral

Figure 1. White light interferometry imaging of developing biofilm topography. (A) Light intensity and (B) surface measurement data from the edge of an *Aeromonas veronii* inoculum on LB-Agar are shown 30 minutes after inoculation. (C) One-dimensional averaged profile of the surface topography is computed from the data in (B). (D) A 24-hour timelapse of averaged profiles from growing *A. veronii* biofilm is shown. The colony expands horizontally (x-axis) and vertically (y-axis), with some of its surface features persisting during development. The scale in the y-axis has been increased to better observe the data. The region shown in panels A-C is marked with a rectangle.

Figure 1–Figure supplement 1. Sub-micrometer spatial growth of an *A. veronii* colony.

Figure 1–Figure supplement 2. Continuous interferometry measurements of growing colony does not affect vertical growth

expansion and the initial conditions from inoculation, such as the coffee-ring effect (*Yanni et al., 2017*; *Agrawal et al., 2020*). We measured prokaryotic and eukaryotic species, including a wide range of gram-negative and gram-positive bacteria, as well as aerobic and aerotolerant anaerobic yeast (*Saccharomyces cerevisiae*) (see *Table 2*). All species and strains show similar qualitative behavior; we thus initially focus on *A. veronii*, and later expand our study to the full cohort of species and strains.

We observe that the vertical growth rate (i.e., change in height per unit time) changes over time; initially increasing before reaching a maximum and then slowing down, behavior that is qualitatively replicated by logistic growth (*Figure 2*A). Immediately after inoculation, the vertical growth rate is very low ($0 - 2\mu m/hr$, see *Figure 2*A inset). This is consistent with the fact that cells are adapting to a new environment(*Bertrand, 2019*; *Hamill et al., 2020*) and that doubling events primarily lead to more horizontal colonization until a monolayer is formed(*Echten et al., 2020*). During the first few hours of growth, the vertical growth rate increases proportionally with the height (*Figure 2*B), consistent with exponential growth. The growth rate eventually reaches a maximum value, and then decreases linearly, approaching zero.

Nutrient depletion

Models of microbial colony growth often directly model the transport and depletion of nutrients. In fact, a prevalent hypothesis is that the biofilm runs out of nutrients, leading to a decrease in growth rate and eventually a final, maximum height. To evaluate the role of nutrient depletion in vertical growth of biofilms, we grew colonies for 48 hours on polycarbonate membranes with pores large enough for nutrients to pass through but small enough to prevent bacteria from passing through, on small agar columns ($3.24cm^2$ in surface area and $0.5cm$ m tall) cut from larger agar plates. These small agar columns limit lateral diffusion of nutrients. On typical large agar plates, nutrients diffuse horizontally and vertically; on these small columns, nutrients can only diffuse vertically up to the colony. After 48 hours, we removed the membrane and colony. We then placed a new membrane on the used column and inoculated, for a second time, in the same location. We found that despite the fact that the first colony nearly reached its saturated height(80%), the second colony grew to 75% of the same height. We then repeated this process on the same plates, i.e., we inoculated at the same location for a third time. Again, we observed that biofilms readily grew, reaching an average height 45% of their previous height. These observations were robust against 12 replicates(*Figure 2*C). Control plates without replacement showed a total growth of $295.4\mu m$, while columns with replacement grew a total of $462.1\mu m$, an excess of 55% (*Figure 2*D). These experiments demonstrate that vertical growth rates can dramatically slow despite the presence of abundant nutrients. Or, equivalently, in laboratory conditions the steady state height biofilms reach after a long time is not due to the absence of nutrients.

Empirical model selection

We can now use these empirical data to guide the selection of an accurate heuristic model. Plotting Δh as a function of h makes it clear that there are two distinct regimes: (I) a linear increase in growth rate with height for $h < L$ and (II) a linear decrease in growth rate with height for $h > L$ (*Figure 2* A-B), where L is a characteristic height. The combination of a linear increase and a linear decrease suggests that growth dynamics cannot be governed by a single, unchanging relationship over time. Instead, models of vertical growth must include a "switch" from one regime to another, and a heuristic model of this phenomenon must justify this empirical observation.

One possible explanation for this "switch" from one regime to another comes from nutrient diffusion, the main method of transport inside microbial colonies(*Stewart, 2003*). The key insight is that biofilms only interact with their environment through their interfaces, which limits the nutrient acquisition and posterior diffusion to the cells inside the colony. Nutrients can only enter biofilms through the biofilm-air interface that supplies oxygen to the colony, and the biofilm-substrate interface, which supplies water, carbon sources, and other macroscopic nutrients(*Xu et al., 1998*;

Figure 2. Biofilm height dynamics and geometric constraints. (A) Biofilm height versus time is shown for a 48-hour period in linear and log scales (inset). Three replicates of *A. veronii* are shown. Height increases even at early times, as seen on the log scale. (B) Change in height is plotted against biofilm height. There are two clear regimes: (I) accelerated and (II) decelerated growth, separated by a characteristic length. (C) The height of microbial colonies grown on small agar columns, thus preventing lateral diffusion of nutrients is shown. In one set of experiments, colonies on the agar columns are replaced every two days (R). In a control set of experiments, colonies are not replaced (NR) and instead are allowed to continue growing. (D) The total height of colonies grown on individual agar columns over a period of 6 days is shown. These results demonstrate that colony height does not saturate due to nutreint depletion. (E) Total nutrient concentration through an interface can be approximated as a simple minimum function, and can be used to model the two different growth regimes.

Figure 2–Figure supplement 1. Change in heights versus time

Figure 2–Figure supplement 2. Total volume growth in agar columns

Stewart and Franklin, 2008). Ideally, a heuristic model of biofilm growth should be nutrient independent, that is, it should apply equally well to all colonies, regardless of the limiting nutrient. We thus propose a simple "interface growth model" that is biophysically motivated and consistent with empirical data.

To model the impact of these interfaces on nutrient concentration, we directly incorporate the distribution of nutrients as a function of distance from the interface. In fact, the concentration of nutrients near the interface, assuming an unlimited surface source, is known (*Figure 2*E):

$$N(h, L) = \frac{-L \cdot e^{-h^2/L^2} + \sqrt{\pi} \cdot h \cdot \text{erfc}\left(\frac{h}{L}\right) + L}{\sqrt{\pi}} \tag{1}$$

Even with the high resolution of optical interferometry, this level of detail is excessive. First, this analytical expression was derived for infinite media, and biofilms are obviously finite. Second, and more importantly, this relationship does not take into account the active uptake of nutrients. Thus, this analytical expression $N(h, L)$ does not directly apply to real biofilms.

Nonetheless, equation 1 provides us with relevant and critical physical insight. Specifically, the nutrient uptake rate of the biofilms should saturate after the colony reaches a critical height, which we label $h = L$. This phenomenon represents the diffusion limit, which prevents biofilms with $h > L$ from growing exponentially. To incorporate these biophysical insights and account for the uncertainty in the exact functional form of $N(h, L)$ for real biofilms, we approximate this relationship as $\min(h, L)$, a simple linear function that reaches its maximum value of L for $h \geq L$. Similar expressions have been previously derived and explored in mathematical models for biofilm growth (*Klapper, 2012*; *Wang et al., 2017*). This simplified expression captures, to the first order in h, the underlying biophysics of diffusion.

We now assemble a heuristic model of vertical growth that takes these empirical and biophysical phenomena into account. These phenomena can be captured with a simple 3-parameter differential equation:

$$\frac{dh}{dt} = \alpha * \min(h, L) - \beta h \tag{2}$$

Where α is the growth rate, β is the decay rate, and L is the diffusion length of the limiting nutrient. This model can be rewritten as a piecewise function in relation to the critical height L:

$$\frac{dh}{dt} = \begin{cases} (\alpha - \beta) \cdot h & x \leq L \\ \alpha \cdot L - \beta \cdot h & x \geq L \end{cases} \tag{3}$$

This expression captures the main biophysical processes governing vertical growth, while remaining simple for any implementation or quick back of the envelope calculations.

Model accuracy

We now test the accuracy of the interface limited model, as well as two minimal models that rely on common alternative approaches to modeling vertical growth: logistic and nutrient depletion models. These reference models do not represent the state of the art, but instead provide a simple implementation of common modeling choices, allowing us to test their accuracy. Specifically, the logistic model is built on the commonly used carrying capacity term that is quadratic in h, and the nutrient depletion model incorporates first-order growth and death terms, with a nutrient field that is consumed over time. We will also consider logistic and interface models that incorporate nutrient fields, to test if doing so results in improved accuracy (for more details on these alternative models, see Materials and Methods).

Over 48 hours of growth at room temperature, the interface model accurately captures colony height across the full time period (*Figure 3*A). This is true both when the auxiliary nutrient field c is or is not included, supporting the idea that nutrient depletion is not the limiting factor for growth,

6 of 21

Figure 3. Interface model outperforms the nutrient and logistic models over 48 hours of growth. (A) The mean height of *A. veronii* colonies, averaged over three replicates, is shown versus time. Error bars represent standard deviation across replicates. Best fit lines for the interface, logistic, and nutrient models are shown. Interface and logistic models expanded to include depletable nutrient fields are shown in solid lines; models without nutrient fields are shown as dashed lines. Note, the two interface models overlap in the figure, but the model values do differ across the data, as can be seen by their different RMSEs. (B) Residuals for the best fit predictions from the above models are shown as a function of time. On average, the nutrient model is off by $15.47 \mu m$, the logistic model by $10.77 \mu m$, and the interface model by $1.24 \mu m$. The gray region corresponds to the standard deviation of the 3 replicates in relation to the mean value at each time. (C) Change in height, Δ Height, is shown as a function of Height for best fit predictions. The two interface models follow the empirical data much more closely than all other models..

Figure 3–Figure supplement 1. Unbounded best fits for the 48h data.

Figure 3–Figure supplement 2. Comparison between the analytical and approximation for the interface constraint

Figure 4. Long term growth of microbial colonies. (A) Measurements of height versus time are shown for three different species. Error bars represent standard deviations across the 2mm homeland region. Solid lines show the best-fit interface model. (B) Change in height is shown versus colony height in the 2-14 day range. The slow linear decrease in growth rate is observed in all measured strains. The solid line represents the best-fit growth rates according to the interface model. (C) Colony heights during the initial 48 hours of growth of plotted against best fit predictions from data taken during the 2-14 day range. The agreement between the data and the model is evident.

Figure 4–Figure supplement 1. Distribution of steady state heights from 48-hour data.

Figure 4–Figure supplement 2. Interface model best fit parameter evolution for an 88 hour *Escherichia coli* timelapse.

and does not need to be directly incorporated for model accuracy. The root-mean-squared error,
RMSE, quantifies the relative agreement of these models: $0.15 \mu m$ for the interface model without a
nutrient field, and $2.71 \mu m$ for the interface model with a nutrient field (*Figure 3*A inset). This good
agreement can also be quantified with the residual as a function of time (*Figure 3*B). The interface-
limited model oscillates around $0 \mu m$, with fluctuations similar to the expected differences from the
3 parallel replicates. Further, the alternative models are poor fits for the height dynamics; these
models grow slower than the experimental data at early times, and then reach their saturated
height too early (*Figure 3*A). For each of the alternative models RMSE> 10 μm.

To determine why the alternative models fail, we examine the change in height as a function
of the current height for all models (*Figure 3*C). The interface limited model directly incorporates
the empirical relationship between Δh and h, and thus accurately fits these data. Conversely, the
alternative models completely fail to capture the observed linear relationships between growth
rate and height.

Behavior on long time scales

We next test the agreement of the interface model on long time scales, measuring h over a period
of two weeks. To do so, we measure the growth of three different species representing a wide
range of microbial growth dynamics: (i) the previously introduced *A. veronii*, (ii) *E. coli*, and (iii) an
aerotolerant anaerobic *S. cerevisiae* mutant. The interface model accurately fits the first 48 hours of
vertical growth for *E. coli* and the anaerobic *S. cerevisiae*, with RMSE of $1.7 \mu m$ and $3.07 \mu m$, respectively.
To test long term accuracy, we inoculate multiple colonies on multiple plates, and then measure all
colonies on a new plate every two days (see Materials and Methods for more details). The interface
model accurately captures the vertical growth dynamics of all three microbes on long time scales
(*Figure 4*A), while exhibiting a linear decrease in growth speed, consistent with the interface model,
although at different heights and time scales (*Figure 4*B).

207 It is important to note that the long time agreement of the interface model does not come at
208 the cost of accuracy on short time scales (*Figure 4*C). We compared the experimental data and the
209 model predictions in the 48-hour period; for all three colonies R^2 was above 0.99. The ability to
210 capture short and long term dynamics, simultaneously highlights that, in fact, vertical growth is
211 accurately described by the simple interface model.

212 We next test the predictive power of the model at short times to determine what governs the
213 dynamics and steady state behavior of long-standing colonies. These three species exhibit differ-
214 ent growth rates, and different heights after the initial 48 hours of growth: $203.83 \pm 4.34, 208.64 \pm 8.07$,
215 and $116.88 \pm 5.33 \mu m$ for *A. veronii*, *E. coli*, and aerotolerant anaerobic *S. cerevisiae*, respectively. The
216 predicted steady state maximum height $h_{\max}$ can be obtained from the model parameters, follow-
217 ing the expression:

$$h_{\max} = \frac{\alpha \cdot L}{\beta} \tag{4}$$

218 This expression correctly predicts that *S. cerevisiae*, will continue to be the tallest colony by a con-
219 siderable amount, and, more interestingly, that *E. coli* will surpass *A. veronii* in height, despite the
220 fact that *A. veronii* is almost twice as tall as *E. coli* at 48 hours. This prediction would not be possible
221 without high resolution data or the accurate model.

222 However, while experimental steady state heights fall in the 95% confidence interval of the
223 interface model fitted using the initial 48 hours of data(*Figure Supplement 1*), the range of this
224 interval varies substantially across these three species ($21.6 \mu m$, $215.4 \mu m$, and $473.23 \mu m$ for *A. veronii*,
225 *E. coli*, and aerotolerant anaerobic *S. cerevisiae*, respectively).

226 The source of this wide range of uncertainties is the difficulty in rapidly measuring β. β is small,
227 on the order of 10 nm/hr. 48 hours is thus not necessarily enough time for an accurate measure-
228 ment, leading to a wide range of effective growth rates at the end of the measurement. A colony
229 that is already close to its steady state height (*A. veronii*) has a much narrower confidence interval
230 than a colony that is still growing rapidly (*S. cerevisiae*). To determine β with narrow confidence
231 intervals, a large amount of data in the $h > L$ regime is necessary.

232 To demonstrate this issue, we performed an 88 hour long timelapse of *E. coli*, and obtained
233 the best fit parameters as a function of time. It is clear that by 40 hours α has already converged
234 (*Figure Supplement 2*), whereas β and L do not converge to their final values until ~ 65 hours.
235 It is important to note that the percentage change of these values varies greatly, with the limiting
236 parameter being β, which at 50 hours is 230% of its final steady state value. On its own, fluctuations
237 in β that range from $10 - 23 nm/hr$ do not seem substantial. However, in light of *Equation 4*, these
238 fluctuations can greatly change the predicted final height of the colony. As a result, $h_m ax$ cannot
239 be accurately predicted until β is accurately measured.

Universality of the vertical growth dynamics of the interface growth model

241 Finally, while we have thus far focused on a small number of species, the interface growth model
242 is built on biophysical limits of diffusion that should readily apply to many species and strains. To
243 test how broadly this model applies, we perform measurements on nine different organisms in-
244 cluding (1) different cell sizes (from $\sim 1 - 10$ µm), shapes (from rods to commas to nearly spherical
245 ellipsoids), (3) different extracellular matrix production (from engineered strains with limited extra-
246 cellular matrix production to wild type *V. cholerae* that are known biofilm formers), (4) prokaryotic
247 and eukaryotic species, (5) gram positive and gram negative species, and (6) aerobic and anaerobic
248 fungi.

249 For this purpose, we measure the vertical growth of these microbes, with three parallel repli-
250 cates, for a period of 48 hours. This length of time is sufficient to capture the two growth regimes
251 delimited by L, and thus fit all interface model parameters. We observe excellent agreement be-
252 tween the interface model and the vertical growth dynamics of all investigated species and strains

Figure 5. Growth of different species and strains over 48 hours. The average height plus or minus the standard deviation across 3 parallel colonies is shown in gray. Error bars represent standard deviation across replicates. The best fit interface model is shown in orange. The model RMSE and the total number of interferometry profiles analyzed are reported in each panel. While these microbial colonies differ in their composition, height, radius, and overall morphologies, the interface model accurately describes the average height dynamics over time for each one.

Figure 5–Figure supplement 1. Topographic profiles after 48 hours of growth

Figure 5–Figure supplement 2. Fit parameters for the interface model

Figure 5–Figure supplement 3. 48 hour growth of multiple microbes in contrast to a simple logistic model

Figure 5–Figure supplement 4. RMSE comparison between the interface and a simple logistic model

Table 1. Interface model parameters for the cohort of microbes. Shown is the best-fit, and 90% confidence interval for the 48 hour data.

Strain	$\alpha[1/hr]$	$\beta[1/hr]$	$L[\mu m]$
Aeromonas veronii	$0.875(0.858 - 0.911)$	$0.051(0.048 - 0.055)$	$13.205(12.277 - 13.878)$
Escherichia coli	$0.339(0.326 - 0.360)$	$0.023(0.012 - 0.036)$	$14.347(11.705 - 17.622)$
Saccharomyces cerevisiae (aa)	$0.345(0.337 - 0.350)$	$0.023(0.011 - 0.028)$	$32.071(27.485 - 33.612)$
Saccharomyces cerevisiae	$0.555(0.538 - 0.579)$	$0.037(0.026 - 0.05)$	$44.018(37.053 - 51.257)$
Vibrio cholerae (wt)	$1.112(0.99 - 1.284)$	$0.038(0.031 - 0.048)$	$7.066(5.559 - 8.861)$
Vibrio cholerae (EPS-)	$0.995(0.886 - 1.232)$	$0.029(0.017 - 0.045)$	$7.783(5.371 - 10.532)$
Klebsiella pneumoniae	$1.316(1.253 - 1.440)$	$0.057(0.05 - 0.064)$	$14.475(12.241 - 16.278)$
Bacillus cereus	$0.881(0.858 - 0.921)$	$0.089(0.041 - 0.107)$	$22.27(18.486 - 25.641)$
Staphylococcus aureus	$0.475(0.464 - 0.488)$	$0.053(0.041 - 0.063)$	$17.025(14.726 - 18.783)$

(see *Figure 5*), despite dramatic differences in colony morphologies and growth rates during these 48 hours of measurement.

We also summarize the allowed ranges for growth parameters in *Table 1*). This resource is useful for future modeling work, with tested and empirical evidence on the accuracy of the model. The parameter range across the sampled cohort is quite large. α ranges by a factor of 4, between $0.3 hr^{-1}$(*E. coli*) to $1.3 hr^{-1}$ (*K. pneumoniae*). L also exhibits a wide range of values, from $8\mu m$ (both *V. cholerae* strains) to $45\mu m$ (*S. cerevisiae)* grown on YPD media). β is more difficult to characterize. Its range of best fit values is consistent across different strains, going from $0.02 hr^{-1}$ to $0.05 hr^{-1}$ for *E. coli* and *B. cereus*, respectively. However, the confidence intervals, relative to the best fit parameter value, are typically larger for β than for α and L. Due to their large sizes, the true β values may either be nearly all $\sim 0.04 hr^{-1}$, or distributed across a range from $0.01 hr^{-1}$ to $0.07 hr^{-1}$.

Discussion

We investigated the vertical growth dynamics of microbial colonies using timelapse interferometry. This technique allows continuous non-disruptive measurements across broad time and length scales with single nanometer resolution, enabling us to characterize the vertical growth dynamics of microbial colonies with unprecedented precision. Using these high precision data, we found that the vertical growth rate initially increases linearly with height; after reaching a critical length scale (L), the growth rate then decreases linearly with height. We show that a simple heuristic model built on the interface-limited diffusion of nutrients accurately captures short and long term dynamics. This minimal model was validated across many different species and strains of bacteria and fungi.

The use of interferometry was essential for the empirical characterization of vertical growth dynamics presented here. For example, high resolution measurements were necessary to identify that growth rate decreases linearly with h for $h > L$, rather than quadratically with h. This empirical relationship provides valuable insight into the development of microbial colonies. Further, β, the linear decay rate, is typically very small, on the order of tens of nanometers per hour. These measurements were possible as interferometry readily provides both the high resolution out-of-plane necessary to capture small changes in height, as well as the fast measurement speed necessary to characterize changes over time.

Interferometry, however, has three key limitations. First, this technique only measures the top surface of a colony, and cannot measure the internal structure. Second, the steepest measureable slope is limited by the numerical aperture of the objective used. When the surface is too steep ($\theta > 28.13°$ for the objective we used) the light path does not return to the objective. This slope threshold may limit the use of interferometry on colonies that are highly wrinkled or buckled(*Dietrich et al., 2013*; *Wilking et al., 2013*; *Yan et al., 2019*). Finally, optical interferometry requires that a surface

be sufficiently reflective. This condition will always be met for the colony-air interface; however, measurements at biofilm-fluid interfaces, while possible, are much more difficult. Despite these limitations, interferometry measurements work in a broad range of surface attached colonies and allow different models of vertical growth dynamic to be tested.

Models of biofilm growth are diverse; they incorporate different biophysical phenomena and use different mathematical approaches. However, the lack of empirical data characterizing vertical growth dynamics prevented these models and their underlying assumptions from being tested. Crucially, the empirical data presented here have the resolution necessary to test the accuracy of existing models, evaluate their underlying assumptions, and then use empirical data to develop an accurate heuristic model.

For example, we find that models built on logistic growth fail to capture the correct relationship between growth rate and colony height. A logistic term is functionally equivalent to a term that decreases growth rate quadratically with height. This term is mathematically appealing; it is the lowest order term that qualitatively describes the changes in growth. Despite its conceptual appeal, it lacks empirical justification (*Figure 3*C). Of course, most of the models developed in the literature are complex, and include more terms and fields in the set of governing equations. However, the evidence presented here suggests that going forward, logistic-like vertical growth should be replaced with empirically-supported expressions.

Similar to logistic growth, we found that nutrient depletion models are not empirically supported, since they do not capture the relationship between growth rate and height(*Figure 3*C). Further, for rich media laboratory experiments, nutrient depletion is not responsible for the eventually cessation of vertical growth (*Figure 2* C-D), and adding nutrient depletion to the interface model did not improve their accuracy(*Figure 3*A). While nutrient depletion may be important to account for near-starvation conditions, it is not a critical component of vertical growth in typical laboratory conditions.

Empirical measurements of growth rate as a function of height demonstrate that three terms are necessary to fully describe vertical growth dynamics. The inclusion of each of these three terms in the interface model we propose is motivated by three fundamental biophysical quantities. L is the length at which growth rate ceases to increase in proportion to biofilm height. This term is inspired by, and consistent with, the length at which diffusion limits the transportation of nutrients through biofilm interfaces. The growth rate α captures all vertical growth, including the growth and doubling of cells, but also secretion of extracellular substances. The decay rate β captures all effects that decrease growth rate, such as mechanical settling, cell death and lysis, cells entering a lag phase, and diminished efficiency in the reabsorption of biomass. In other words, α and β coarse grain all phenomena that lead to an increase or decrease (respectively) in growth rate into two simple terms. Rather than limiting the utility of this model, coarse graining places the emphasis on the net impact of increased height on growth rates, which may be universal, rather than on the microscopic mechanisms that modify growth rates, which will often be specific and heterogeneous. This approach provides a useful heuristic model for vertical growth dynamics, as opposed to more complex alternatives(*Levins, 1966*).

The unique insight provided by high resolution studies of vertical growth dynamics is exemplified through two case studies comparing: (i) the role of respiration in *S. cerevisiae*, and (ii) the role of extracellular matrix production in *Vibrio cholerae*. First, we find that there is no qualitative difference between the vertical growth dynamics of aerobic and obligate anaerobic strains of *S. cerevisiae*. The aerobic yeast strain used in this study is "mixotrophic," capable of both fermentation and aerobic respiration(*De Deken, 1966*) depending on the concentration and types of sugars found in the environment. The anaerobic strain has lost part of its mitochondrial genome and is therefore only capable of fermentation, which is not oxygen-dependent. Multicellular anaerobic yeast have recently been shown to overcome group size constraints that aerobic yeast face due to the oxygen diffusion limitation into the center of a group(*Bozdag et al., 2021*). The aerobic strain grows faster than the anaerobic strain (*Bozdag et al., 2021*), but in each case the vertical growth

339 dynamics proceed following the interface model. In a different vein, we directly compare the ver-
340 tical growth dynamics of strains of *V. cholerae* with and without EPS. Vibrio polysaccharide is an
341 essential component for biofilm formation, and the associated genes are controlled by the mas-
342 ter biofilm regulator, *VpsR*(*Yildiz et al., 2001*; *Fong et al., 2010*). Deletion of *vpsR* (EPS-) prevents
343 biofilm production in *V. cholerae*, leading instead to the growth of "smooth" colonies with much
344 less extracellular matrix than wild type strains. Again we find that the vertical growth dynamics do
345 not change substantially in the presence or absence of EPS. Thus, microbial colony vertical growth
346 dynamics appear relatively universal, even across microbial strains with different compositions.

347   As the interface model is a simple heuristic model for vertical growth, there are many details it
348 does not address. For example, the model predicts the dynamics of the bulk height, and does not
349 distinguish between cells and extracellular matrix within the colony. Relatedly, as the model fo-
350 cuses on the mean total height it does not address spatial fluctuations. Topographic fluctuations
351 likely encode insight into the underlying phenomena behind vertical growth dynamics, as previ-
352 ously demonstrated in homeostatic biofilms (*Kalziqi et al., 2018*, *2019*). In a different vein, the
353 model does not account for more complex environmental conditions, such as limited access to
354 nutrients(*Chénier et al., 2003*), environmental stresses(*Tsagkari et al., 2022*), or heterogeneities in
355 the surface on which the colony grows(*Gralka and Hallatschek, 2019*). Further, in nature biofilms
356 often contain multiple interacting species and subpopulations(*Stacy et al., 2016*), leading to com-
357 plex dynamics and structuring(*Nadell et al., 2016*; *Ghoul and Mitri, 2016*) of the colony that we do
358 not consider. The work presented here provides a foundation on which these detailed questions
359 about vertical growth dynamics can be studied.

360   The study of microbial community interfaces via interferometry presents an easy and inexpen-
361 sive approach to studying the structure of microbial colonies. Interfaces and surface fluctuations
362 have long been studied in physics, both for the information they provide about what occurs be-
363 neath the surface as well as the rich phenomena that surfaces themselves exhibit. Indeed, biol-
364 ogy also has a rich history of studying interfaces, from the surfaces of cells or organs(*Cosgrove,*
365 *2005*; *Abbott et al., 2010*; *Shen et al., 2011*; *Harayama and Riezman, 2018*), to biome-scale inter-
366 faces(*Erdős et al., 2018*). Biological interfaces on meso-scales are traditionally less studied, despite
367 their known importance in a wide range of systems(*Liu et al., 2013*; *Turley et al., 2015*; *Ladoux*
368 *and Mège, 2017*) including bacterial communities (*Bonachela et al., 2011*; *Kassinger and van Hoek,*
369 *2020*). This is likely due to the difficulty in measuring these microscopic, and often sticky, interfaces.
370 Other techniques for characterizing the height and topographies of colonies lack the resolution
371 (e.g., confocal microscopy), are slow and potentially destructive (e.g., atomic force microscopy), or
372 do not readily permit time lapse measurements (e.g., scanning electron microscopy)(*Huang et al.,*
373 *2020*). While these alternative measurement techniques might be modified to overcome their lim-
374 itations(*Raab and Bachelet, 2017*), interferometry is already very well-suited for studying biofilm
375 interfaces, as demonstrated here and in previous studies (*Larimer et al., 2016*; *Kalziqi et al., 2018*,
376 *2019*). These works represent the first attempts in utilizing optical interferometry as a tool for
377 microbiology, an approach that will only get more useful when paired with the development of
378 heuristic models and detailed measurement protocols.

379   Finally, the dynamics of vertical growth represent a fundamental aspect of microbial physiol-
380 ogy. Biofilms are exceedingly common; they are found in many environments (*Ghiorse and Wil-*
381 *son, 1988*; *Horner-Devine et al., 2004*; *Azam and Malfatti, 2007*) and are of key importance from a
382 medical(*Bryers, 2008*; *Zhao et al., 2013*; *Veerachamy et al., 2014*; *Wi and Patel, 2018*; *Sharma et al.,*
383 *2019*) and economic(*Cámara et al., 2022*) perspective. An understanding of how biofilms develop
384 their 3-dimensional structure is thus relevant for diverse questions of microbial ecology, evolution,
385 and human health. In particular, understanding vertical growth dynamics is crucial for assessing
386 the underlying biophysical process regarding biofilm development: the way resources do and do
387 not limit growth, the relative fitness of various biofilm morphologies (e.g., thin biofilms vs. thick
388 biofilms). The vertical growth measurements and model describing them are but one of the many
389 aspects in biofilm development, but necessary ones for proper understanding how these microbial

Table 2. Timelapses overview details.

Strain	Species	Media	Details	Cell shape	Gram
ZOR0001	*Aeromonas veronii*	LB 1.5%	wild type	rod	negative
MG1655	*Escherichia coli*	LB 1.5%	wild type	rod	negative
GOB33	*Saccharomyces cervisiae*	YPD	petite yeast	ellipsoid	-
Y55	*Saccharomyces cervisiae*	YPD	wild type	ellipsoid	-
C6706	*Vibrio cholerae*	LB 1.5%	wild type	comma	negative
C6706	*Vibrio cholerae*	LB 1.5%	Δ*vpsR*	comma	negative
Top52	*Kelbsiella pneumoniae*	LB 1.5%	wild type	rod	negative
SW520	*Bacillus cereus*	LB 1.5%	wild type	rod	positive
SW519	*Staphylococcus aureus*	LB 1.5%	wild type	spherical	positive

390 colonies grow.

Methods and Materials

Measurement details

All inoculations consist of $1.5\mu L$ from an overnight culture grown at $37\,°C$ diluted to $OD_{600} = 1$.
Control plates were left to grow at a temperature of $23.8\,°C$. For interferometry measurements,
we designed an enclosure that allows continuous measurements with minimal media and sample
evaporation. The full enclosure maintains a temperature of $23.8\,°C$ and a relative humidity of 80%
for the full measuring period.

Since colony growth is radially symmetric, we only measure a horizontal strip of the biofilm,
allowing us to achieve better temporal resolution. The measurement proceeds across the entire
colony, and includes uncolonized substrate surface on both sides of the colony. Given the aspect
ratio of our data and the goals of our study, we then lower the dimension of the data by averag-
ing on the Y-axis (*Figure 1*A-C). Then, we utilize the substrate height measurements to obtain the
relative height of the colony to its background.

Column growth

We label and cut columns of LB-agar to dimensions of $18mm \times 18mm \times 5mm$. On each column we
deposited a Polycarbonate Track Etch Membrane with $0.2\mu m$ pore size, $13mm$ diameter (GVS brand).
We inoculate $1.5\mu L$ of OD1 bacterial suspension on top of said membrane. This is repeated over 6
plates (18 columns/colonies). Every 48 hours, up to 3 iterations, we measure the growth on each
column. On 9 of the columns we than removed the membrane and deposited a new membrane
and inoculated on top of the new membrane.

48h timelapses

We inoculated 3 colonies of the measured strain, on two separate petri dishes with LB-Agar. One
plate was measured continuously (replicates A-C) for 48 hours in the profilometer enclosure. Mea-
surements consist of horizontal radial strips of each colony with a Zygo ZeGage Pro optical pro-
filometer with a 50x Mirau objective (NA 0.55), on the 1000x200 @800Hz capture mode. We in-
crease the Field of View of the measurement adaptively, to always capture an uncolonized region
outside the growing colony. Colonies that were continuously measured did not show any signifi-
cant growth differences compared to their sealed counterparts (replicates D-) after the experiment
(*Figure Supplement 2*).

Long time measurements

421 We inoculated 21 plates 7 for ZOR0001, 7 MG1655, and 7 GOB33 (*Table 2*), with 3 separate colonies
422 each. All plates were sealed with parafilm after inoculation. Every two days, we performed radial

423 measurements with a 10x Mirau objective (NA 0.3) 1000x200 @800Hz capture mode. Each plate
424 was discarded after measurement.

Geometric constraint derivation

426 The geometric constraint detailed in *Equation 1* is rooted in a simple diffusion equation. This ex-
427 pression is used in fabrication principles and was taken from *Ghandhi* (*2008*). Starting from Fick's
428 Law for a concentration field c and diffusion coefficient D:

$$\frac{\partial c}{\partial t} = D \cdot \frac{\partial^2 c}{\partial t^2} \tag{5}$$

429 By direct integration we obtain that the distribution of nutrients as follows:

$$c(x, t) = \frac{c_0}{2\sqrt{\pi D t}} \cdot e^{-x^2/4Dt} \tag{6}$$

430 where c_0 is the initial concentration. If we now consider an infinite source with concentration $c = c_0$
431 on one side of a slab of material (in our case, a microbial colony), the concentration on a finite
432 element of width $d\xi$ is:

$$c(x, t) = \frac{c_0}{2\sqrt{\pi D t}} \cdot \int_x^\infty e^{-\xi^2/4Dt} d\xi \tag{7}$$

433 Using the error function

$$\mathrm{erf}(y) = \frac{2}{\sqrt{\pi}} \int_0^\infty e^{-z^2} dz \tag{8}$$

434 we can rewrite the expression as:

$$c(x, t) = \frac{c_0}{2} \cdot \left(1 - \mathrm{erf}\left(\frac{x}{2\sqrt{Dt}}\right)\right) = \frac{c_0}{2} \cdot \mathrm{erfc}\left(\frac{x}{2\sqrt{Dt}}\right) \tag{9}$$

435 where erfc is the complimentary error function. Finally, we can obtain the total amount of nutrients
436 in a finite sized slab of height h by integrating this expression:

$$N(h, L) = \int_0^h c(x, L) dx = \frac{-L \cdot e^{-h^2/L^2} + \sqrt{\pi} \cdot h \cdot \mathrm{erfc}\left(\frac{h}{L}\right) + L}{\sqrt{\pi}} \tag{10}$$

437 where we removed the temporal dependence by a characteristic diffusion length $L = \sqrt{Dt}$.

Growth models and ODE simulations

439 As benchmarks for comparison to the interface model we utilize two intuitive models, a nutrient
440 depletion model and a logistic model.

441 Nutrient depletion model
This model captures the dynamics of how a biomass of an available nutrient c is transformed into
biomass height h (assuming that mass $\propto$ height in a column). Both fields are linearly related; there-
fore a decrease in growth rate has to be due to the lack of nutrient availability.

$$\frac{dh}{dt} = \alpha \cdot h \cdot \frac{c}{K_c + c} - \beta h \tag{11}$$

$$\frac{dc}{dt} = -\epsilon \cdot \alpha \cdot h \cdot \frac{c}{K_c + c} \tag{12}$$

442 Here, α is the growth rate, β the decay rate, $\frac{c}{K_c + c}$ is the Monod kinematic term for nutrient uptake,
443 and ϵ is the inverse conversion rate for the transfer. For our simulations we set $h_0 = 0.01 \mu m$, and
444 $c_0 = 1.0$. This model can not be used without nutrient c dependency, since the equation would then
445 be reduced to the monotonic equation $\dot{h} = (\alpha - \beta)h$.

Table 3. Fitting boundaries for model fitting. Ranges push the boundaries of what makes physical/biological sense by at least an order of magnitude.

Model	$\alpha[1/hr]$	$\beta[1/hr]$	$L[\mu m]$	$K[\mu m]$	$K_c[a.u.]$	ϵ
Nutrient (n)	$[10^{-5}, 10^3]$	$[10^{-5}, 10^2]$	-	-	$[10^{-5}, 10^3]$	$[10^{-5}, 10^5]$
Logistic (n)	$[10^{-5}, 10^3]$	-	-	$[1, 10^5]$	$[10^{-5}, 10^3]$	$[10^{-5}, 10^5]$
Interface (n)	$[10^{-5}, 10^3]$	$[10^{-5}, 10^2]$	$[1, 10^5]$	-	$[]$	$[10^{-5}, 10^5]$
Logistic	$[10^{-5}, 10^3]$	-	-	$[1, 10^5]$	-	-
Interface	$[10^{-5}, 10^3]$	$[10^{-5}, 10^2]$	$[1, 10^5]$	-	-	-

Logistic

The use of a logistic model makes sense in a qualitative manner, since we do observe a behavior that follows "accelerated-decelerated" growth. The issue with these kind of models is that while simple, they associate the negative(−) term in the differential expression to a quadratic term, obscuring the meaning of the variable.

$$\frac{dh}{dt} = \alpha \cdot h \cdot \frac{c}{K_c + c} \cdot \left(1 - \frac{h}{K}\right) \tag{13}$$

$$\frac{dc}{dt} = -\epsilon \cdot \alpha \cdot h \cdot \frac{c}{K_c + c} \tag{14}$$

Variables are the same as in the previous model, with K representing the carrying capacity, or the maximum height the colony can reach. We can also utilize the logistic model without a nutrient dependency, and rewrite the expression as $\dot{h} = \alpha \cdot h(1 - h/K)$.

Parameter estimation

We optimized the multiple differential equations against experimental data with a sum of squared error loss J:

$$J(f(x), y) = \sum_i^n (f(x_i) - y_i)^2 \tag{15}$$

where y is the collection of experimental heights corresponding to measurement times t, $f(x)$ is the evaluation of the differential equation at time t. This allow us to directly compare the simulated predictions with our non equidistant temporal measurements. While this loss function J prioritizes better agreement with the data on higher heights, results do not suggest that compensating with higher weights at early times is needed since residual values are low across all the measurement period.

To get confidence intervals and parameter distributions we utilize a moving block bootstrap across the 3 different replicates, sampling $n = 20$ blocks of size $s = 5$, returning a total of 100 points, roughly the same amount as an individual timelapse. We iterate for $i = 1 : 1000$ times and obtain the best-fit parameters α_i, β_i, L_i. Then utilizing equation (4), discard the outlying 5% values.

All parameter estimation was done in Julia with the DifferentialEquations.jl(*Rackauckas and Nie, 2017*) and DiffEqFlux.jl(*Rackauckas et al., 2020*) packages.

Data availability

Source code for the cleaning, analysis and figures, and clean datasets have been uploaded to a Github repository. Due to file size limits, full unprocessed images are available upon request.

Acknowledgments

P.J.Y. acknowledges funding from the NIH NIGMS (Grant No. 1R35GM138354-01) and NSF Biomaterials (Grant No. BMAT2003721). B.K.H. acknowledges funding from NSF Biomaterials (Grant No. BMAT2003721).

References

Abbott NJ, Patabendige AA, Dolman DE, Yusof SR, Begley DJ. Structure and function of the blood–brain barrier. Neurobiology of disease. 2010; 37(1):13–25.

Agrawal A, Sinha S, Mukherjee R, Mampallil D. Dynamics of bacterial deposition in evaporating drops. Physics of Fluids. 2020; 32(9):093308.

Allen RJ, Waclaw B. Bacterial growth: a statistical physicist's guide. Reports on Progress in Physics. 2019 Jan; 82(1):016601. https://iopscience.iop.org/article/10.1088/1361-6633/aae546, doi: 10.1088/1361-6633/aae546, 00000.

Alpkvist E, Picioreanu C, van Loosdrecht MCM, Heyden A. Three-dimensional biofilm model with individual cells and continuum EPS matrix. Biotechnology and Bioengineering. 2006; 94(5):961–979. Publisher: Wiley Online Library.

Azam F, Malfatti F. Microbial structuring of marine ecosystems. Nature Reviews Microbiology. 2007; 5(10):782–791.

Basaran M, Yaman YI, Yüce TC, Vetter R, Kocabas A. Large-scale orientational order in bacterial colonies during inward growth. Elife. 2022; 11:e72187.

Beroz F, Yan J, Meir Y, Sabass B, Stone HA, Bassler BL, Wingreen NS. Verticalization of bacterial biofilms. Nature Physics. 2018 Sep; 14(9):954–960. https://www.nature.com/articles/s41567-018-0170-4, doi: 10.1038/s41567-018-0170-4, number: 9 Publisher: Nature Publishing Group.

Bertrand RL. Lag phase is a dynamic, organized, adaptive, and evolvable period that prepares bacteria for cell division. Journal of bacteriology. 2019; 201(7):e00697–18.

Bonachela JA, Nadell CD, Xavier JB, Levin SA. Universality in bacterial colonies. Journal of Statistical Physics. 2011; 144(2):303–315.

Bozdag GO, Libby E, Pineau R, Reinhard CT, Ratcliff WC. Oxygen suppression of macroscopic multicellularity. Nature communications. 2021; 12(1):1–10.

Bryers JD. Medical biofilms. Biotechnology and bioengineering. 2008; 100(1):1–18.

Cámara M, Green W, MacPhee CE, Rakowska PD, Raval R, Richardson MC, Slater-Jefferies J, Steventon K, Webb JS. Economic significance of biofilms: a multidisciplinary and cross-sectoral challenge. npj Biofilms and Microbiomes. 2022; 8(1):1–8.

Chénier MR, Beaumier D, Roy R, Driscoll BT, Lawrence JR, Greer CW. Impact of seasonal variations and nutrient inputs on nitrogen cycling and degradation of hexadecane by replicated river biofilms. Applied and Environmental Microbiology. 2003; 69(9):5170–5177.

Cooper A, Dean A, Hinshelwood CN. Factors affecting the growth of bacterial colonies on agar plates. Proceedings of the Royal Society of London Series B Biological Sciences. 1968; 171(1023):175–199.

Cosgrove DJ. Growth of the plant cell wall. Nature reviews molecular cell biology. 2005; 6(11):850–861.

Coyte KZ, Tabuteau H, Gaffney EA, Foster KR, Durham WM. Microbial competition in porous environments can select against rapid biofilm growth. Proceedings of the National Academy of Sciences. 2017; 114(2):E161–E170.

Davey ME, O'toole GA. Microbial biofilms: from ecology to molecular genetics. Microbiology and molecular biology reviews. 2000; 64(4):847–867.

De Beer D, Stoodley P, Roe F, Lewandowski Z. Effects of biofilm structures on oxygen distribution and mass transport. Biotechnology and bioengineering. 1994; 43(11):1131–1138.

De Deken R. The Crabtree effect: a regulatory system in yeast. Microbiology. 1966; 44(2):149–156.

Dervaux J, Magniez JC, Libchaber A. On growth and form of Bacillus subtilis biofilms. Interface Focus. 2014; 4(6). doi: 10.1098/rsfs.2013.0051.

Dietrich LE, Okegbe C, Price-Whelan A, Sakhtah H, Hunter RC, Newman DK. Bacterial community morphogenesis is intimately linked to the intracellular redox state. Journal of bacteriology. 2013; 195(7):1371–1380.

518 **Donlan RM**. Biofilm formation: a clinically relevant microbiological process. Clinical infectious diseases. 2001;
519 33(8):1387–1392.

520 **D'Acunto B**, Frunzo L, Klapper I, Mattei M. Modeling multispecies biofilms including new bacterial species
521 invasion. Mathematical biosciences. 2015; 259:20–26.

522 **Echten DvHt**, Nordemann G, Wehrens M, Tans S, Idema T. Defect dynamics in growing bacterial colonies. arXiv
523 preprint arXiv: 200310509. 2020; p. 1–17. http://arxiv.org/abs/2003.10509, arXiv: 2003.10509.

524 **Erdős L**, Kröel-Dulay G, Bátori Z, Kovács B, Németh C, Kiss PJ, Tölgyesi C. Habitat heterogeneity as a key to high
525 conservation value in forest-grassland mosaics. Biological Conservation. 2018; 226:72–80.

526 **Fei C**, Mao S, Yan J, Alert R, Stone HA, Bassler BL, Wingreen NS, Košmrlj A. Nonuniform growth and sur-
527 face friction determine bacterial biofilm morphology on soft substrates. Proceedings of the National
528 Academy of Sciences. 2020 Apr; 117(14):7622–7632. https://www.pnas.org/doi/10.1073/pnas.1919607117,
529 doi: 10.1073/pnas.1919607117, publisher: Proceedings of the National Academy of Sciences.

530 **Flemming HC**, Wingender J, Szewzyk U, Steinberg P, Rice SA, Kjelleberg S. Biofilms: an emergent form of
531 bacterial life. Nature Reviews Microbiology. 2016 Aug; 14(9):563–575.

532 **Fong JC**, Syed KA, Klose KE, Yildiz FH. Role of Vibrio polysaccharide (vps) genes in VPS production, biofilm
533 formation and Vibrio cholerae pathogenesis. Microbiology. 2010; 156(Pt 9):2757.

534 **Fortune GT**, Oliveira NM, Goldstein RE. Biofilm Growth under Elastic Confinement. Physical Review Let-
535 ters. 2022 Apr; 128(17):178102. https://link.aps.org/doi/10.1103/PhysRevLett.128.178102, doi: 10.1103/Phys-
536 RevLett.128.178102, publisher: American Physical Society.

537 **Ghandhi SK**. VLSI fabrication principles: silicon and gallium arsenide. John Wiley & Sons; 2008.

538 **Ghiorse WC**, Wilson JT. Microbial ecology of the terrestrial subsurface. Advances in applied microbiology. 1988;
539 33:107–172.

540 **Ghosh P**, Levine H. Morphodynamics of a growing microbial colony driven by cell death. Physical Review E.
541 2017; 96(5):1–8. doi: 10.1103/PhysRevE.96.052404.

542 **Ghoul M**, Mitri S. The ecology and evolution of microbial competition. Trends in microbiology. 2016; 24(10):833–
543 845.

544 **Gralka M**, Hallatschek O. Environmental heterogeneity can tip the population genetics of range expansions.
545 Elife. 2019; 8:e44359.

546 **Hall-Stoodley L**, Costerton JW, Stoodley P. Bacterial biofilms: from the natural environment to infectious dis-
547 eases. Nature Reviews Microbiology. 2004 Feb; 2(2):95–108.

548 **Hamill PG**, Stevenson A, McMullan PE, Williams JP, Lewis AD, Stevenson KE, Farnsworth KD, Khroustalyova G,
549 Takemoto JY, Quinn JP, et al. Microbial lag phase can be indicative of, or independent from, cellular stress.
550 Scientific reports. 2020; 10(1):1–20.

551 **Harayama T**, Riezman H. Understanding the diversity of membrane lipid composition. Nature reviews Molec-
552 ular cell biology. 2018; 19(5):281–296.

553 **Hartmann R**, Singh PK, Pearce P, Mok R, Song B, Díaz-Pascual F, Dunkel J, Drescher K. Emergence of three-
554 dimensional order and structure in growing biofilms. Nature Physics. 2019; 15(3):251–256. http://dx.doi.org/
555 10.1038/s41567-018-0356-9, doi: 10.1038/s41567-018-0356-9, publisher: Springer US.

556 **Hayta EN**, Rickert CA, Lieleg O. Topography quantifications allow for identifying the contribution of parental
557 strains to physical properties of co-cultured biofilms. Biofilm. 2021 Dec; 3:100044. https://www.sciencedirect.
558 com/science/article/pii/S2590207521000022, doi: 10.1016/j.biofilm.2021.100044.

559 **Horner-Devine MC**, Carney KM, Bohannan BJ. An ecological perspective on bacterial biodiversity. Proceedings
560 of the Royal Society of London Series B: Biological Sciences. 2004; 271(1535):113–122.

561 **Huang Y**, Chakraborty S, Liang H. Methods to probe the formation of biofilms: applications in foods and related
562 surfaces. Analytical Methods. 2020 Jan; 12(4):416–432. https://pubs.rsc.org/en/content/articlelanding/2020/
563 ay/c9ay02214g, doi: 10.1039/C9AY02214G, publisher: The Royal Society of Chemistry.

Hughes J. A Mathematical Model of Biofilm Growth on Degradable Substratum. SIAM Undergraduate Research Online. 2020; 13. https://www.siam.org/Portals/0/Publications/SIURO/Vol13/M130847PDF.pdf?ver=2020-06-11-163504-750, doi: 10.1137/19S1308475.

Jin X, Marshall JS, Wargo MJ. Hybrid Model of Bacterial Biofilm Growth. Bulletin of Mathematical Biology. 2020 Feb; 82(2):27–27. doi: 10.1007/s11538-020-00701-6, mAG ID: 3003693092.

Kalziqi A, Ng SL, Yanni D, Steinbach G, Hammer BK, Yunker PJ. Viscosity independent diffusion mediated by death and reproduction in biofilms. arXiv preprint arXiv: 190101350. 2019; http://arxiv.org/abs/1901.01350, arXiv: 1901.01350.

Kalziqi A, Yanni D, Thomas J, Ng SL, Vivek S, Hammer BK, Yunker PJ. Immotile Active Matter: Activity from Death and Reproduction. Physical Review Letters. 2018; 120(1):18101. https://doi.org/10.1103/PhysRevLett.120.018101, doi: 10.1103/PhysRevLett.120.018101, publisher: American Physical Society.

Kassinger SJ, van Hoek ML. Biofilm architecture: An emerging synthetic biology target. Synthetic and Systems Biotechnology. 2020 Mar; 5(1):1–10. https://www.sciencedirect.com/science/article/pii/S2405805X20300016, doi: 10.1016/j.synbio.2020.01.001.

Kesel S, Bronk Bv, García CF, Götz A, Lieleg O, Opitz M. Matrix composition determines the dimensions of Bacillus subtilis NCIB 3610 biofilm colonies grown on LB agar. RSC Advances. 2017 Jun; 7(51):31886–31898. https://pubs.rsc.org/en/content/articlelanding/2017/ra/c7ra05559e, doi: 10.1039/C7RA05559E, publisher: The Royal Society of Chemistry.

Klapper I. Productivity and Equilibrium in Simple Biofilm Models. Bulletin of Mathematical Biology. 2012 Dec; 74(12):2917–2934. https://doi.org/10.1007/s11538-012-9791-4, doi: 10.1007/s11538-012-9791-4.

Ladoux B, Mège RM. Mechanobiology of collective cell behaviours. Nature reviews Molecular cell biology. 2017; 18(12):743–757.

Lambert G, Bergman A, Zhang Q, Bortz D, Austin R. Physics of biofilms: The initial stages of biofilm formation and dynamics. New Journal of Physics. 2014; 16. doi: 10.1088/1367-2630/16/4/045005, publisher: IOP Publishing.

Larimer C, Suter JD, Bonheyo G, Addleman RS. In situ non-destructive measurement of biofilm thickness and topology in an interferometric optical microscope. Journal of Biophotonics. 2016; 9(6):656–666. https://onlinelibrary.wiley.com/doi/abs/10.1002/jbio.201500212, doi: 10.1002/jbio.201500212, _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/jbio.201500212.

Levins R. The strategy of model building in population biology. American scientist. 1966; 54(4):421–431.

Liu QX, Doelman A, Rottschäfer V, de Jager M, Herman PM, Rietkerk M, van de Koppel J. Phase separation explains a new class of self-organized spatial patterns in ecological systems. Proceedings of the National Academy of Sciences. 2013; 110(29):11905–11910.

Mahto KU, Das S. Microscopic techniques to evaluate the biofilm formation ability of a marine bacterium Pseudomonas aeruginosa PFL-P1 on different substrata. Microscopy Research and Technique. 2021; 84(10):2451–2461. https://onlinelibrary.wiley.com/doi/abs/10.1002/jemt.23799, doi: 10.1002/jemt.23799, _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1002/jemt.23799.

Mattei MR, Frunzo L, D'Acunto B, Pechaud Y, Pirozzi F, Esposito G. Continuum and discrete approach in modeling biofilm development and structure: a review. Journal of Mathematical Biology. 2018 Mar; 76(4):945–1003. doi: 10.1007/s00285-017-1165-y, mAG ID: 2739410635.

Mazza MG. The physics of biofilms - An introduction. Journal of Physics D: Applied Physics. 2016; 49(20). doi: 10.1088/0022-3727/49/20/203001, arXiv: 1606.01392 Publisher: IOP Publishing.

Monod J. The growth of bacterial cultures. Selected Papers in Molecular Biology by Jacques Monod. 2012; 139.

Moore-Ott JA, Chiu S, Amchin DB, Bhattacharjee T, Datta SS. A biophysical threshold for biofilm formation. eLife. 2022; 11:e76380.

Nadell CD, Drescher K, Foster KR. Spatial structure, cooperation and competition in biofilms. Nature Reviews Microbiology. 2016; 14(9):589–600.

Nguyen H, A Y, Abraham Ybarra, Başağaoğlu H, Shindell O. Biofilm viscoelasticity and nutrient source location control biofilm growth rate, migration rate, and morphology in shear flow. Scientific Reports. 2021 Aug; 11(1):16118. doi: 10.1038/s41598-021-95542-1, mAG ID: 3188109131.

Oliveira NM, Martinez-Garcia E, Xavier J, Durham WM, Kolter R, Kim W, Foster KR. Biofilm formation as a response to ecological competition. PLoS biology. 2015; 13(7):e1002191.

Paula AJ, Hwang G, Koo H. Dynamics of bacterial population growth in biofilms resemble spatial and structural aspects of urbanization. Nature Communications. 2020 Mar; 11(1):1354. https://www.nature.com/articles/s41467-020-15165-4, doi: 10.1038/s41467-020-15165-4, number: 1 Publisher: Nature Publishing Group.

Raab N, Bachelet I. Resolving biofilm topography by native scanning electron microscopy. Journal of Biological Methods. 2017 Apr; 4(2):e70–e70. https://jbmethods.org/jbm/article/view/173, doi: 10.14440/jbm.2017.173, number: 2.

Rackauckas C, Ma Y, Martensen J, Warner C, Zubov K, Supekar R, Skinner D, Ramadhan A, Edelman A. Universal differential equations for scientific machine learning. arXiv preprint arXiv:200104385. 2020; .

Rackauckas C, Nie Q. DifferentialEquations.jl – A Performant and Feature-Rich Ecosystem for Solving Differential Equations in Julia. Journal of Open Research Software. 2017 May; 5(1):15. http://openresearchsoftware.metajnl.com/articles/10.5334/jors.151/, doi: 10.5334/jors.151, number: 1 Publisher: Ubiquity Press.

Rendueles O, Ghigo JM. Mechanisms of competition in biofilm communities. Microbial Biofilms. 2015; p. 319–342.

Schiebel J, Noack J, Rödiger S, Kammel A, Menzel F, Schwibbert K, Weise M, Weiss R, Böhm A, Nitschke J, Elimport A, Roggenbuck D, Schierack P. Analysis of three-dimensional biofilms on different material surfaces. Biomaterials Science. 2020; 8(12):3500–3510. https://pubs.rsc.org/en/content/articlelanding/2020/bm/d0bm00455c, doi: 10.1039/D0BM00455C, publisher: Royal Society of Chemistry.

Sharma D, Misba L, Khan AU. Antibiotics versus biofilm: an emerging battleground in microbial communities. Antimicrobial Resistance & Infection Control. 2019; 8(1):1–10.

Shen L, Weber CR, Raleigh DR, Yu D, Turner JR. Tight junction pore and leak pathways: a dynamic duo. Annual review of physiology. 2011; 73:283.

Srinivasan S, Kaplan CN, Mahadevan L. A multiphase theory for spreading microbial swarms and films. eLife. 2019; 8:8–10. doi: 10.7554/eLife.42697.

Stacy A, McNally L, Darch SE, Brown SP, Whiteley M. The biogeography of polymicrobial infection. Nature Reviews Microbiology. 2016; 14(2):93–105.

Steinbach G, Crisan C, Ng SL, Hammer BK, Yunker PJ. Accumulation of dead cells from contact killing facilitates coexistence in bacterial biofilms. Journal of the Royal Society Interface. 2020; 17(173):20200486.

Stewart PS. Diffusion in Biofilms. Journal of Bacteriology. 2003 Mar; 185(5):1485–1491. https://www.ncbi.nlm.nih.gov/pmc/articles/PMC148055/, doi: 10.1128/JB.185.5.1485-1491.2003, 01179.

Stewart PS, Franklin MJ. Physiological heterogeneity in biofilms. Nature Reviews Microbiology. 2008; 6(3):199–210.

Stewart PS, Zhang T, Xu R, Pitts B, Walters MC, Roe F, Kikhney J, Moter A. Reaction–diffusion theory explains hypoxia and heterogeneous growth within microbial biofilms associated with chronic infections. NPJ biofilms and microbiomes. 2016; 2(1):1–8.

Tam A, Green JEF, Balasuriya S, Tek EL, Gardner JM, Sundstrom JF, Jiranek V, Binder BJ. A thin-film extensional flow model for biofilm expansion by sliding motility. Proceedings of the Royal Society A: Mathematical, Physical and Engineering Sciences. 2019 Sep; 475(2229):20190175. https://royalsocietypublishing.org/doi/10.1098/rspa.2019.0175, doi: 10.1098/rspa.2019.0175, publisher: Royal Society.

Tam AKY, Harding B, Green JEF, Balasuriya S, Binder BJ. Thin-film lubrication model for biofilm expansion under strong adhesion. Physical Review E. 2022 Jan; 105(1):014408. https://link.aps.org/doi/10.1103/PhysRevE.105.014408, doi: 10.1103/PhysRevE.105.014408, publisher: American Physical Society.

Tsagkari E, Connelly S, Liu Z, McBride A, Sloan WT. The role of shear dynamics in biofilm formation. npj Biofilms and Microbiomes. 2022; 8(1):1–10.

Turley SJ, Cremasco V, Astarita JL. Immunological hallmarks of stromal cells in the tumour microenvironment. Nature reviews immunology. 2015; 15(11):669–682.

Veerachamy S, Yarlagadda T, Manivasagam G, Yarlagadda PK. Bacterial adherence and biofilm formation on medical implants: a review. Proceedings of the Institution of Mechanical Engineers, Part H: Journal of Engineering in Medicine. 2014; 228(10):1083–1099.

Wang X, Stone HA, Golestanian R. Shape of the growing front of biofilms. New Journal of Physics. 2017 Dec; 19(12):125007. https://doi.org/10.1088/1367-2630/aa983f, doi: 10.1088/1367-2630/aa983f, publisher: IOP Publishing.

Warren MR, Sun H, Yan Y, Cremer J, Li B, Hwa T. Spatiotemporal establishment of dense bacterial colonies growing on hard agar. eLife. 2019; 8:1–47. doi: 10.7554/eLife.41093.

Watnick P, Kolter R. Biofilm, city of microbes. Journal of Bacteriology. 2000; 182(10):2675–2679. Publisher: Am Soc Microbiol.

Werb M, Falcón García C, Bach NC, Grumbein S, Sieber SA, Opitz M, Lieleg O. Surface topology affects wetting behavior of Bacillus subtilis biofilms. npj Biofilms and Microbiomes. 2017 Apr; 3(1):1–10. https://www.nature.com/articles/s41522-017-0018-1, doi: 10.1038/s41522-017-0018-1, number: 1 Publisher: Nature Publishing Group.

Wi YM, Patel R. Understanding biofilms and novel approaches to the diagnosis, prevention, and treatment of medical device-associated infections. Infectious Disease Clinics. 2018; 32(4):915–929.

Wilking JN, Zaburdaev V, De Volder M, Losick R, Brenner MP, Weitz DA. Liquid transport facilitated by channels in Bacillus subtilis biofilms. Proceedings of the National Academy of Sciences. 2013; 110(3):848–852.

Wille J, Coenye T. Biofilm dispersion: The key to biofilm eradication or opening Pandora's box? Biofilm. 2020; 2(October 2019):100027. doi: 10.1016/j.bioflm.2020.100027.

Xiong L, Cao Y, Cooper R, Rappel WJ, Hasty J, Tsimring L. Flower-like patterns in multi-species bacterial colonies. Elife. 2020; 9:e48885.

Xu KD, Stewart PS, Xia F, Huang CT, McFeters GA. Spatial physiological heterogeneity in Pseudomonas aeruginosa biofilm is determined by oxygen availability. Applied and environmental microbiology. 1998; 64(10):4035–4039.

Yan J, Fei C, Mao S, Moreau A, Wingreen NS, Košmrlj A, Stone HA, Bassler BL. Mechanical instability and interfacial energy drive biofilm morphogenesis. Elife. 2019; 8:e43920.

Yan J, Sharo AG, Stone HA, Wingreen NS, Bassler BL. Vibrio cholerae biofilm growth program and architecture revealed by single-cell live imaging. Proceedings of the National Academy of Sciences. 2016; 113(36):E5337–E5343.

Yanni D, Kalziqi A, Thomas J, Ng SL, Vivek S, Ratcliff WC, Hammer BK, Yunker PJ. Life in the coffee-ring: how evaporation-driven density gradients dictate the outcome of inter-bacterial competition. arXiv preprint arXiv:170703472. 2017; .

Yildiz FH, Dolganov NA, Schoolnik GK. VpsR, a member of the response regulators of the two-component regulatory systems, is required for expression of vps biosynthesis genes and EPSETr-associated phenotypes in Vibrio cholerae O1 El Tor. Journal of bacteriology. 2001; 183(5):1716–1726.

Zhang T, Cogan NG, Wang Q. Phase-Field models for biofilm. I. Theory and 1-D simulations. Siam Journal On Applied Mathematics. 2008; 69(3):641–669.

Zhao G, Usui ML, Lippman SI, James GA, Stewart PS, Fleckman P, Olerud JE. Biofilms and inflammation in chronic wounds. Advances in wound care. 2013; 2(7):389–399.

Figure 1–Figure supplement 1. Interferometry allows accurate measurement of the spatial vertical growth dynamics. An *A. veronii* colony after inoculation. Initially, individual cells can be observed undergoing reproduction through their height profiles.

Figure 1–Figure supplement 2. Colony heights are not affected by continuous measurement under the interferometer. Control samples were inoculated from the same overnight batch on a different plate, which was then left to grow at room temperature, sealed with Parafilm. Comparisons were after each timelapse finished, at 48.6, 87.7, 47.6, 48.6, 49.5, 50.2, 48.3, 47.1, 47.48 hours respectively.

Figure 2–Figure supplement 1. The growth regimes are not immediately evident when plotting their the growth over time. Each color represents a replicate of *A. veronii*, Measurements are the same as in the main figure.

Figure 2–Figure supplement 2. Growth of microbial colonies in a LB-Agar column, with (R) and without (NR) replacement every two days. (D) Total amount of growth supported by an LB-Agar colony over a period of 6 days.

Figure 3–Figure supplement 1. Best fits are shown for the different models while using a free optimization. For the nutrient model the parameters are: [3.3e7, -1.3e-1, 2.7e7, 5.2e6]. Logistic (n): [4.4e12, 5.2e2, 4.5e12, 5.9e11]. Interface (n): [2.0e0, 1.1e-1,1.6e1,1.8e0,-1.4e-1]. Logistic:[4.6e0, 1.6e2]. Interface:[7.5e-1,5e-2,1.5e1]. The parameter lists are ordered as in Table 2.

Figure 3–Figure supplement 2. Best fits values are shown for the *A. veronii* timelapse in the main figure panel A. Parameters for the analytical expression are: $0.55, 0.05, 20.96$, for the approximation: $0.87, 0.05, 13.20$, for α, β, L respectively. The only significant discrepancies occur around the inflection point, but deviations do not exceed $2\mu m$ in magnitude. On the right, we see the difference between these models by plotting the analytical expression minus the approximation.

707

Figure 4–Figure supplement 1. Interface model h_{max} distributions after N=1000 bootstrapped fits are shown. Black circles represent the best fit for the initial 48 hour time period, and orange triangles represent the best fit including the measurements up to 14 days.

708

Figure 4–Figure supplement 2. Best fit, and 90% confidence interval for the the three interface model parameters: α, β, L, and the steady state maximum height h_{max} are shown. On the left y-axis the actual values are reported, while on the right y-axis reports the value relative to the final 88 hour best fit value.

709

Figure 5–Figure supplement 1. Profiles of the measured microbes after 48 hours of growth. Pictured are two different aspect ratios for the x and y axis for better visualization of the surface structures.

Figure 5–Figure supplement 2. Fit parameters for the interface model across different bacteria. This panel demonstrate the range of possible parameter combinations, and similarities between strains.

Figure 5–Figure supplement 3. A simple logistic model fails to describe vertical height dynamics, across the sampled cohort of microbial colonies.

Figure 5–Figure supplement 4. The interface model outperforms the logistic model in all sampled colonies.