Polynuclear ruthenium complexes are effective antibiotics against *Pseudomonas*

aeruginosa

Brent S. Weber[1], Lindsey A. Carfrae[1], Joshua J. Woods[2,3], Kristina Klobucar[1,4], Nicholas

P. Bigham[2], Craig R. MacNair[1], Tracy L. Raivio[5], Justin J. Wilson[2], and Eric D. Brown[1,*]

[1]Department of Biochemistry and Biomedical Sciences, Michael G. DeGroote Institute

for Infectious Disease Research, McMaster University, Hamilton, ON, Canada

[2]Department of Chemistry and Chemical Biology, Cornell University, Ithaca, NY, USA

[3]Present Affiliation: Chemical Sciences Division, Lawrence Berkeley National

Laboratory. Berkeley, CA, USA

[4]Present Affiliation: Bureau of Gastroenterology, Infection and Viral Diseases, Health

Canada, Ottawa, ON, Canada

[5]Department of Biological Sciences, University of Alberta, Edmonton, AB, Canada

[*]Correspondence: ebrown@mcmaster.ca

Abstract

There is an urgent need to develop new antibiotics for the treatment of infections caused by drug-resistant Gram-negative bacteria. In particular, new and diverse chemical classes of antibiotics are needed, as most antibiotics in clinical development are derivatives of existing drugs. Despite a history of use as antimicrobials, metals and metal-based compounds have largely been overlooked as a source of new chemical matter for antibacterial drug discovery. In this work, we identify several ruthenium complexes, ruthenium red, Ru265, and Ru360', that possess potent antibacterial activity against both laboratory and clinical isolates of *Pseudomonas aeruginosa*. Suppressors with increased resistance were sequenced and found to contain mutations in the mechanosensitive ion channel *mscS-1* or the *colRS* two component system. The antibacterial activity of these compounds translated *in vivo* to *Galleria mellonella* larvae and mouse infection models. Finally, we identify strong synergy between these compounds and the antibiotic rifampicin, with a dose-sparing combination therapy showing efficacy in both infection models. Our findings provide clear evidence that these ruthenium complexes are effective antibacterial compounds against a critical priority pathogen and show promise for the development of future therapeutics.

Introduction

The antimicrobial resistance crisis is a major threat to global public health, and our ability to treat serious bacterial infections remains tenuous. The pipeline of new drugs, particularly for critical priority Gram-negative pathogens, is inadequate to meet the realities facing modern healthcare (1, 2). For numerous reasons, including economic and safety concerns, antibiotic development continues to experience challenges in bringing new drugs to market (3). While most new antibiotics under development for the treatment of Gram-negative infections are derivatives of existing drug classes (1), there has been increasing interest in less conventional therapeutics (4), including anti-virulence compounds, host-directed therapeutics, and drug repurposing strategies (5–8). Despite substantial efforts, however, innovations in this space have yet to translate to successful clinical approvals of novel drugs (9).

Metals have played an important role in medicine, including their use as antimicrobial agents. One of the first synthetic antibiotics, Salvarsan, was an arsenic-based compound used to treat syphilis (10), and dressings embedded with silver formulations are used as therapeutics for burn wounds (11, 12). While some metal-based compounds have found success in other clinical applications, such as platinum metals for cancer therapies (13), there are no metal-based systemic therapies for bacterial infections. This is likely due to the historical success of conventional antibacterial therapies and the perceived toxicity concerns with metal compounds. However, with the rise of antibacterial resistance, metal-based antibiotics are drawing increased interest

(14, 15). Indeed, recent work has emphasized that metal compounds are rich sources of

both antibacterial and antifungal agents (16).

Recently, we showed that the metal complex ruthenium red (RuRed) possesses

antibacterial activity against the Gram-negative bacterial pathogen *Klebsiella*

pneumoniae under certain growth conditions, which translated to efficacy in an *in vivo*

rat infection model (17). RuRed and related ruthenium complexes are primarily

recognized as inhibitors of the mitochondrial calcium uniporter (MCU) (18, 19).

Mitochondrial Ca^{2+} plays an important role in diverse cellular processes, and uptake of

Ca^{2+} into mitochondria is mediated by the MCU ion channel (20–22). Several disease

states are associated with dysregulated mitochondrial Ca^{2+} influx, including cystic

fibrosis (CF), which has led to interest in therapeutics and chemical probes targeting the

MCU (23, 24). In this work, we show that RuRed and other similar ruthenium complexes

have potent antibacterial activity against laboratory and clinical isolates of

Pseudomonas aeruginosa. We find that *P. aeruginosa* shows unique susceptibility to

these compounds and identify mutations that decrease RuRed susceptibility. Finally, we

show that these ruthenium complexes are effective as therapeutics in several *in vivo*

infection models and synergize with the Gram-positive antibiotic rifampicin in dose-

sparing combination treatments. Our work provides a foundation for further exploration

of these ruthenium complexes as therapies to treat *P. aeruginosa* infections.

Results

Ruthenium red has potent antibacterial activity against laboratory and clinical isolates of *P. aeruginosa*

Previously, we found that RuRed had antibacterial activity against *K. pneumoniae* in the presence of human serum (17). In that work, tests in conventional lysogeny broth (LB) growth medium revealed RuRed did not inhibit growth of wild-type *K. pneumoniae* at concentrations up to 256 μg/mL, though we identified genetic mutants that were susceptible under those growth conditions (17). Here, we tested the activity of RuRed against two common laboratory strains of *P. aeruginosa*, PAO1 and PA14 (Fig. 1A). Surprisingly, RuRed inhibited the growth of both strains in standard antibiotic testing medium Mueller-Hinton broth (MHB), with a minimum inhibitory concentration (MIC) of 2 μg/mL (Fig. 1B). Like our previous work, growth in serum enhanced the activity of RuRed against *P. aeruginosa* PAO1 (Fig. S1) and tests in LB showed a 4-fold decrease in efficacy against PAO1 (Table S1). Interestingly, we detected an MIC for RuRed against *K. pneumoniae* in MHB (64 μg/mL), but not LB (>256 μg/mL), suggesting a difference in efficacy between LB and MHB (Table S1).

Given the potent activity of RuRed against *P. aeruginosa* PAO1 and PA14, we next tested a collection of 22 clinical isolates of *P. aeruginosa*. All isolates showed similar susceptibility to RuRed in MHB, with a MIC range of 1-4 μg/mL and a median of 2 μg/mL (Fig. 1C and Table S1). Testing of several other Gram-negative and -positive

pathogens showed that, while most species had some level of RuRed susceptibility in

MHB, *P. aeruginosa* strains were uniformly and uniquely highly susceptible to this

compound (Table S1). We note that, while RuRed has been used experimentally for

decades, its purity can vary between suppliers and is often not reported (25). Therefore,

the purity of the commercially available RuRed used in this work was assessed by

ultraviolet-visible (UV−vis) spectroscopy (Fig. S2). Our samples showed strong

absorbance at 533 nm, characteristic of RuRed, and a second smaller peak at 260 nm,

whose identity is not known but is often observed in commercial RuRed samples as a

minor impurity (25). We also observed a minor peak at 360 nm which likely corresponds

to the Ru360, a known impurity of RuRed preparations (26–28).

130   **Fig 1. RuRed is antibacterial against *P. aeruginosa*.** (A) Structure of Ruthenium red.

131   (B) MIC curves for RuRed against laboratory strains of *P. aeruginosa* PAO1 and PA14

132   in MHB. (C) Distribution of RuRed MICs for a panel of *P. aeruginosa* isolates.

**Mutations in PA4394 (MscS-1) or PA4380 (ColS) decrease *P. aeruginosa*
susceptibility to RuRed**

To further characterize the antibacterial activity of RuRed against *P. aeruginosa*, we
isolated spontaneous resistant mutants by plating the wild-type strain on an inhibitory
concentration of RuRed. Mutants recovered using this method had a 4-fold increased
MIC compared to the parental strain (Fig. 2A). Next, we sequenced the genomes of four
mutants and the parental strain to identify any genetic differences. One of the four
suppressor mutants ("suppressor1") contained a mutation in the gene PA4394, which
encodes a predicted mechanosensitive ion channel involved in osmotic tolerance called
MscS-1 (29). This mutation, a point mutant at base 219, would result in a lysine to
asparagine amino acid change (Fig. 2B). The other three mutants (collectively
"suppressor2") had an identical point mutation in the PA4380 gene at position 601,
causing in a single amino acid change (Fig. 2B). PA4380 encodes the sensor kinase
ColS that, together with the response regulator ColR encoded by PA4381, is part of the
ColRS two-component sensor system (30, 31).

To confirm the role of MscS-1 and ColS in *P. aeruginosa* susceptibility to RuRed, the
wild-type (MscS-1^{WT} and ColS^{WT}) and mutant (MscS-1^{K73N} and ColS^{T201P}) alleles were
cloned into pUCP22 plasmid and expressed in the wild-type and mutant strains.
Expression of MscS-1^{K73N}, but not ColS^{T201P}, in the wild-type background increased
resistance of *P. aeruginosa* to RuRed (Fig. 2C). Interestingly, overexpression of the
wild-type MscS-1 allele in the suppressor1 background did not restore susceptibility to

RuRed (Fig. 2D), whereas overexpression of ColS^WT was able to restore sensitivity in

the suppressor2 background (Fig. 2E). All constructs were sequence confirmed and

produced similar levels of recombinant protein (Fig. S3). Thus, these results were not

due to a lack of protein expression but instead suggest a more complex interaction

between the wild-type and mutant alleles for each gene.

Fig 2. Genetic analysis of RuRed suppressors. (A) MICs of wild-type *P. aeruginosa*

PAO1 or 2 independent suppressors. (B) Whole-genome sequencing of suppressors

identified point mutations in *mscS-1* (suppressor1) or *colS* (suppressor2). (C) RuRed

MICs of wild-type PAO1 containing vector control pUCP22 or expressing the mutant

alleles of MscS-1 (pMscS-1^K73N) or ColS (pColS^T201P). (D) RuRed MICs of suppressor1

containing vector control pUCP22 or expressing the wild-type allele of MscS-1 (pMscS-

1^WT). (E) RuRed MICs of suppressor2 containing vector control pUCP22 or expressing

the wild-type allele of ColS (pColS^WT).

To further investigate the interactions of these alleles in RuRed susceptibility, we

obtained insertionally-inactivated mutants in *mscS-1* and *colS* from the *P. aeruginosa*

MPAO1 transposon mutant collection (32). We reasoned that this would allow us to

determine the contribution of the wild-type or mutant alleles in a background devoid of a

functional chromosomal copy of the genes. Unlike the suppressor mutants we isolated,

both *mscS-1* and *colS* transposon mutants had identical RuRed MICs to the parental

MPAO1 strain (Figure 3A). This indicates that the RuRed resistance seen in our

suppressor mutants was not simply due to production of non-functional variants of these

gene products. Expression of MscS-1^{K73N} increased RuRed resistance of both the

MPAO1 wild-type (Figure 3B) and *mscS-1* transposon mutant background (Figure 3C),

whereas expression of MscS-1^{WT} or the vector control had no effect. ColS^{T201P}

expression had no effect in the wild-type background (Figure 3D), but significantly

increased resistance in the *colS* mutant (Figure 3E) while ColS^{WT} had no impact in

either background. Taken together, these results suggest that MscS-1^{K73N} is dominant

to MscS-1^{WT}, and ColS^{WT} is dominant to ColS^{T201P} for the observed RuRed resistance

phenotypes. Given the fact that *mscS-1* and *colS* are not essential for *P. aeruginosa*

viability, and the identified suppressor mutations impart a modest 4-fold increase in

resistance to RuRed, it is unlikely that these gene products are the ultimate target of

RuRed, but instead play an accessory role in susceptibility.

Fig 3. Effect of expression of MscS-1 and ColS alleles in wild-type and mutant

backgrounds on RuRed MIC. (A) RuRed MIC curves of wild-type *P. aeruginosa*

MPAO1 and MscS-1 and ColS transposon mutants. (B) RuRed MIC curves of wild-type

MPAO1 containing vector control pUCP22 or expressing the WT or mutant allele of

MscS-1. (C) RuRed MIC curves of MscS-1 transposon mutant containing vector control

pUCP22 or expressing the WT or mutant allele of MscS-1. (D) RuRed MIC curves of

wild-type MPAO1 containing vector control pUCP22 or expressing the WT or mutant

allele of ColS. (E) RuRed MIC curves of ColS transposon mutant containing vector

control pUCP22 or expressing the WT or mutant allele of ColS.

RuRed analogs are antibacterial

In addition to RuRed, several related ruthenium compounds have been investigated for their ability to inhibit the MCU in eukaryotic cells (33, 34). Given their structural and functional similarity to RuRed, and our ability to synthesize pure compounds in-house, we tested the antibacterial activity of the well-characterized dinuclear ruthenium compounds Ru265, Ru270 and Ru360' (Fig. 4A) (33). All 3 possessed antibacterial activity against *P. aeruginosa* PAO1, with MICs of 8, 64, and 4 µg/mL, respectively (Fig. 4B). In addition, all *P. aeruginosa* clinical isolates tested were susceptible to these compounds, with very similar MICs (Table S1). We also tested 6 mononuclear and 2 additional dinuclear Ru compounds, which had varying levels of antibacterial activity against *P. aeruginosa* PAO1 (Fig. S4). Like RuRed, Ru265, Ru270, and Ru360' were less effective against non-*P. aeruginosa* bacterial species (Table S1). One exception, however, was the activity of Ru360' against *Acinetobacter baumannii*. All tested *A. baumannii*, including common laboratory strains and clinical isolates, were susceptible to Ru360' at concentrations similar to *P. aeruginosa* (Table S1).

Fig 4. RuRed analogs Ru265, Ru270, and Ru360' have antibacterial activity. (A)
Structures of Ru255, Ru270 and Ru360'. (B) MIC curves of the indicated compounds
against *P. aeruginosa* PAO1. (C) Distribution of MICs for a panel of *P. aeruginosa*
isolates for the 3 compounds.

Ruthenium compounds are effective in a *Galleria mellonella* infection model

Given the *in vitro* efficacy of the ruthenium compounds, in particular that of RuRed,
Ru265, and Ru360', we next used a *Galleria mellonella* infection model to assess their
ability to treat infection *in vivo*. *G. mellonella* larvae are a common model system used
to assess antibacterial activity, and activity in this model often translates to rodent
models of infection (35). Groups of 10 larvae were infected with *P. aeruginosa* PAO1
and treated with a single dose of compound 15 min post-infection, then monitored for
survival every 24h for 72h. Treatment of infected larvae with RuRed significantly

enhanced survival of *G. mellonella* larvae in a dose-dependent manner, with 80%

survival at the highest dose tested (Fig. 5A). Control larvae infected and treated with

vehicle solution succumbed to infection by 24h, whereas those treated with the

antibiotic ciprofloxacin survived for the duration of the experiment. Ru265 was similarly

effective, with 90% survival at the highest dose tested (Fig. 5B). Treatment with Ru360'

also significantly extended survival compared to untreated larvae (Fig. 5C). However,

treatment of uninfected larvae with Ru360' revealed that this compound had toxic

effects on its own, with the highest dose of Ru360' resulting in the death of 70% of

larvae (Fig. 5D). In contrast, the highest doses of RuRed and Ru265 had no negative

effects on survival of uninfected larvae (Fig. 5D). Taken together, these results show

that the *in vitro* efficacy of these compounds translates to an invertebrate model of

infection.

Fig 5. Ruthenium compounds are effective in a *G. mellonella* model of infection.

Kaplan-Meier survival curves from *G. mellonella* larvae infected with *P. aeruginosa*

PAO1 and treated with various doses of (A) RuRed, (B) Ru265, (C) Ru360', or

ciprofloxacin (cipro) or vehicle controls. (D) Uninfected larvae were injected with PBS

and subsequently injected with the indicated compounds. In all experiments, survival

was monitored every 24h for a total of 72h. *** indicates significant difference from the

vehicle control. Significance was determined by using the Gehan-Breslow-Wilcoxon test

with a Bonferroni correction for multiple comparisons.

RuRed is effective in a *P. aeruginosa* mouse infection model

Our results with *G. mellonella* prompted us to evaluate RuRed in a mammalian infection

model. We used a systemic mouse infection model, where mice were infected

intraperitoneally with 10^6 CFU/mL *P. aeruginosa* PAO1 and treated 1h and 5h post-

infection with either 2mg/kg RuRed or vehicle control. Additionally, 4 infected mice were

euthanized 1h-post infection to establish bacterial burden (colony forming units, CFUs)

in the spleen, kidney, liver and bloodstream at the time of treatment (Fig. 6). Vehicle-

treated mice had 1000 to 10000-fold increased bacterial CFU burden in the various

organs and blood compared to the burden at the start of treatment (Fig. 6). In contrast,

RuRed treated animals had either the same or slightly reduced bacterial CFUs

compared to the initial burden seen 1h post-infection. Thus, RuRed shows potent

antibacterial activity in a mouse model of systemic *P. aeruginosa* infection.

Fig 6. RuRed is effective in a systemic mouse infection model. Mice infected

intraperitoneally with *P. aeruginosa* PAO1 were euthanized 1h post-infection for CFU

determination (pre-treatment) or treated 1h and 5h post-infection with vehicle or RuRed

(2mg/kg). Treated mice were euthanized 10h post-infection for CFU determination.

Each point represents an individual animal, with the median indicated by the line. *** P

<0.05 compared to vehicle control. Significance was determined using the Mann-

Whitney test.

Ruthenium compounds synergize with antibiotics *in vitro*

Although our *in vivo* studies show that these ruthenium complexes show promise as
treatments for *P. aeruginosa* infection, it is also known that some of these compounds
can have toxic side effects (36, 37). A common approach to ameliorate toxic side effects
is to identify a partner compound that reduces the dose of the toxic molecule required
for efficacy, resulting in a dose-sparing effect (38, 39). We tested combinations of
RuRed with antibiotics in checkerboard assays and found synergy with several
antibiotics normally used against Gram-positive bacteria, including rifampicin,
vancomycin, novobiocin and erythromycin (Fig. S5 A-D). Similarly, Ru265, Ru270, and
Ru360' also synergized with rifampicin (Fig. S5 E and F), with Ru265 showing
particularly strong synergy as evidenced by a fractional inhibitory concentration index
(FICI) of 0.04 (Fig. 7A). Interestingly, this synergy also extended to *E. coli* (Fig. S6A).
One possible explanation for these synergies is that the ruthenium compounds
physically disrupt the outer membrane to allow entry of these antibiotics (40). To
investigate this, we used a lysozyme lysis assay where physical disruption of the *E. coli*
outer membrane allows lysozyme to reach the peptidoglycan layer and lyse the
bacterial cells. Treatment with the control compound SPR741 led to nearly 60% lysis,
whereas RuRed treatment resulted in no change in lysis compared to the no drug
control (Fig. S6B). Thus, although RuRed can potentiate large scaffold antibiotics, the
mechanism of synergy does not seem to be through outer membrane disruption.

The dose-sparing combination of RuRed or Ru265 and rifampicin is effective in vivo

The synergies of RuRed and Ru265 with rifampicin, and their favorable efficacy *in vivo*, encouraged us to test these combinations in the *G. mellonella* infection model. Larvae were infected with a lethal dose of *P. aeruginosa* and treated with the monotherapies or combinations. Treatment with sub-therapeutic levels of RuRed or rifampicin failed to cure infected larvae (Fig. S7). However, the treatment with the combination of RuRed and rifampicin resulted in 100% survival of the infected larvae, consistent with the observed *in vitro* synergies. Similarly, the combination of Ru265 and rifampicin completely cured infected larvae at doses where neither drug was effective on their own (Fig. 7B).

Next, we tested the combination of Ru265 and rifampicin in the systemic mouse infection model. We chose Ru265 due to the strong *in vitro* synergy with rifampicin seen in the checkerboard assays and efficacy in *G. mellonella*. Mice infected with *P. aeruginosa* and treated with the monotherapies had similar bacterial burdens in their blood and organs as vehicle-treated controls (Fig. 7C). Conversely, mice given the combination of Ru265 and rifampicin had a significant reduction in bacterial load in all organs and blood. These results show that addition of rifampicin can have a significant dose-sparing effect for Ru265 treatment, suggesting a promising approach to treat *P. aeruginosa* infection.

Fig 7. The combination of Ru265 and rifampicin is effective in vitro and in vivo. (A)

Checkerboard assay showing synergistic effects (FICI=0.04) of Ru265 and rifampicin

against *P. aeruginosa*. (B) *G. mellonella* larvae were infected with *P. aeruginosa* PAO1

and treated with monotherapies of rifampicin (rif) or Ru265, or the combination of both

(combo). Control larvae were treated with ciprofloxacin (cipro) or vehicle. *** indicates

significant difference from the vehicle control. Significance was determined by using the

Gehan-Breslow-Wilcoxon test with a Bonferroni correction for multiple comparisons. (C)

Systemic mouse infection model**.** Mice infected intraperitoneally with *P. aeruginosa*

PAO1 were treated at 1h and 5h post-infection and were euthanized 10h post-infection

for CFU determination. *** P <0.05 compared to vehicle control determine by Kruskal-

Wallis test.

Discussion

As antibiotic resistance continues to impact patients globally (41), innovative methods
and approaches to drug discovery will be required to maintain the antibacterial pipeline
(9). Recent studies have emphasized that metal complexes, which are often overlooked
as potential antibiotics, can be productive sources of antibacterial and antifungal
chemistry. Frei and colleagues screened nearly 1000 metal compounds for antimicrobial
activity and found a hit-rate more than 10x higher than from conventional organic
compound libraries, with many of these showing no apparent toxicity against
mammalian cells (16). Gallium has shown promise as an antibacterial therapy, including
in preliminary human trials (42, 43). A wide variety of ruthenium complexes have been
investigated for antimicrobial activity, although there is limited *in vivo* efficacy data,
particularly for Gram-negative bacteria (44–47). Our work shows that metal compounds
like RuRed and Ru265 can be effective both *in vitro* and *in vivo* against *P. aeruginosa*, a
notoriously difficult-to-treat Gram-negative bacterial pathogen.

P. aeruginosa showed a unique pattern of susceptibility to the polynuclear ruthenium
compounds tested in this work. The reason(s) for the enhanced susceptibility of *P.
aeruginosa* to the ruthenium compounds described here, compared to other bacteria,
remain to be determined. In our previous work, we found that RuRed had no activity
against *K. pneumoniae* in LB growth media but did have potent activity against *K.
pneumoniae* grown in human serum (17). Curiously, MHB media seems to enhance the
activity of RuRed, as we observed an MIC in this media even for *K. pneumoniae*. This

suggests that growth media formulation influences susceptibility to these compounds.

For example, serum is known to increase membrane permeability which enhances

penetration of large scaffold drugs normally excluded by the outer membrane (48). We

previously identified several *K. pneumoniae* gene knockouts that were susceptible to

RuRed in LB media, such as those involved in outer membrane biogenesis and DNA

repair pathways, and similar pathways have been shown to affect RuRed sensitivity in

E. coli, further supporting the condition-specific antibacterial efficacy of RuRed (17, 49).

It is possible that RuRed uses an entry mechanism into the cell that is constitutively

active in *P. aeruginosa*, but only induced in other species under certain conditions. In

any case, all bacterial species we tested were more resistant to the ruthenium

compounds than *P. aeruginosa*, which indicates that there is some intrinsic property of

P. aeruginosa that sensitizes them to these compounds. The one exception was our

finding that Ru360' had strong antibacterial activity against *A. baumannii* strains while

RuRed, Ru265, and Ru270 had little or none. Overall, this suggests that this family of

ruthenium compounds may have several possible modes of uptake into the cell,

different mechanisms of action, or both. From a structural perspective, our data

suggests that positive charge is important for activity of these ruthenium compounds, as

the negatively charged dinuclear compounds tested were inactive (see Figure S3).

However, a more thorough structure-activity-relationship campaign with a broader range

of compounds should be a priority for future work.

We identified separate mutations in *mscS-1* and *colS* that led to a modest 4-fold

increase in *P. aeruginosa* resistance to RuRed. However, elucidating the precise role

that these loci play in resistance requires further work. It is likely that neither MscS-1 nor

ColS are direct antibacterial targets of RuRed, as knockout mutants in both loci are

viable (32). Experimental evidence has shown that MscS-1 is a mechanosensitive ion

channelthat releases cytoplasmic osmolytes in response to osmotic stress (29). The

MscS-1^{K73N} mutation we identified is unlikely to result in a loss of function; we showed

that a *mscS-1* transposon mutant was just as susceptible as wild-type *P. aeruginosa*,

and expression of MscS-1^{K73N} from a plasmid in the wild-type background led to

resistance, suggesting the mutant allele is dominant. Interestingly, the eukaryotic target

of RuRed and analogs are ion channels like the MCU, although it remains to be seen if

RuRed directly interacts with MscS-1. The *colS* gene encodes the sensor histidine

kinase ColS which, as part of the ColRS two-component system in various

Pseudomonas species, has been implicated in virulence, host colonization, membrane

integrity, and antibiotic resistance in certain genetic contexts (30, 50, 51, 31). The

ColRS system regulates genes directly downstream from the *colRS* cluster, and these

genes, in particular *warA* and *warB* have been shown to be involved in outer membrane

modification and permeability (51–54). Therefore, we speculate that ColS^{T201P} results in

dysregulation of the ColRS system, leading to changes in genes expression which in

turn may prevent RuRed accumulation. Future studies to determine the mechanism of

action of these ruthenium complexes should help uncover their ultimate targets and

shed light on the unique susceptibility of *P. aeruginosa*.

We have shown that RuRed and Ru265 are effective *in vivo* therapies against *P.*

aeruginosa infection in two different models. While the translation from *in vitro* to *in vivo*

activity is clear, one of the major barriers to further developing these ruthenium

compounds as antibiotics will likely be their toxicity. Indeed, we saw evidence of toxicity

of Ru360' in the *G. mellonella* infection model, and both RuRed and Ru265 have shown

dose-dependent toxicity in mice (36, 37). Ideally, analogs could be developed that

reduce toxicity; however, an alternative strategy is to use synergistic combinations of

compounds. Combinations of antibiotics or small molecules that synergize can

significantly reduce toxicity concerns due to the combinations' dose-sparing effect (38,

55, 56). Indeed, we found that synergistic combinations of RuRed or Ru265 with

rifampicin completely cured infected *G. mellonella,* and significantly reduced the

bacterial burden in the mouse model. For the *G. mellonella* infections, the doses of the

ruthenium compounds in the combination therapies were reduced nearly 10-fold

compared to the monotherapy trials, and it is likely that with additional optimization this

could be reduced further. The mechanism for the synergy seen with rifampicin requires

further investigation, though our data suggest that it is not by physical disruption of the

membrane. It is interesting to note that RuRed has been shown to bind to nucleic acids,

which may suggest a possible route to synergy with an antibiotic like rifampicin (57).

While we have not evaluated the toxicity of Ru265, we note that cell-based assays

suggest minimal cytotoxicity of this compound (33), and thus Ru265 may be a promising

lead compound for future development.

Much of the work on RuRed and analogs like Ru265, Ru270, and Ru360' has focused

on their ability to inhibit eukaryotic ion channels like the MCU (19). For antibacterial drug

discovery, a compound with a eukaryotic target is often a liability. However, it was

recently shown that inhibition of the MCU actually improves outcomes in CF models of

P. aeruginosa infection (58). In that work, the authors showed that *P. aeruginosa*

infection caused inflammation and impaired autophagy, which could be remedied by

inhibition of the MCU with a compound called KB-R7943. In their CF mouse model,

mice infected with *P. aeruginosa* and treated with this compound had significantly

decreased bacterial burdens and increased survival. KB-R7493 is an organic compound

that is structurally unrelated to the ruthenium compounds used here, and the authors'

ruled out any direct antibacterial activity caused by the KB-R7493 itself. Thus, it is

tempting to speculate that in a CF model of *P. aeruginosa* infection, administration of

compounds like RuRed or Ru265 could provide a dual benefit: 1) inhibition of the MCU,

resulting in physiological corrections that improve host clearance of *P. aeruginosa* and

2) direct antibacterial activity against *P. aeruginosa*.

Overall, we have uncovered a series of compounds that are effective therapies against

one of the most troublesome Gram-negative pathogens. Our work provides a framework

for future studies that unravel the antibacterial mechanism and full *in vivo* potential of

these promising ruthenium complexes.

Materials and Methods

Bacterial strains, media, and compounds

The bacterial strains used in this work are listed in Table S2. Strains were routinely
cultured and maintained in LB (per liter: 10 g trypticase peptone, 5 g yeast extract, 10 g
NaCl) or LB agar at 37°C. Cation-adjusted Mueller-Hinton II broth (MHB, BD, Cat. #
212322) was used for susceptibility testing. RuRed was purchased from Sigma-Aldrich
(Cat. # R2751), and Ru265, Ru270 and Ru360', and other ruthenium complexes were
synthesized as previously described (33, 34, 59–66). All compounds were dissolved in
water.

Ultraviolet-visible (UV-vis) Spectroscopy of RuRed

Three samples of RuRed, from two different lot numbers but all separately ordered from
the vendor, were analyzed via UV-vis spectroscopy to assess purity. Each RuRed
sample (2.0 mg; 0.0024 mmol) was added to 10 mL 0.1 M ammonium acetate
(NH_4OAc) solution to give a 250 µM concentration, assuming the material was 100%
pure. The solutions were then further diluted 10-fold in 0.1 M NH_4OAc to an assumed
final concentration of 25 µM. UV-vis spectra were taken at 25 °C using a Shimadzu UV-
1900 Spectrophotometer (Shimadzu, Kyoto, Japan) fitted with a temperature-controlled
circulating water bath. Purity of each sample was assessed by the magnitude of
absorbance at 533 nm, characteristic of RuRed (25).

Antibacterial MIC testing and checkerboard assays

Susceptibility MIC testing and checkerboards were performed using a broth

microdilution assay in MHB, unless otherwise noted. Strains were plated on agar and a

loopful of colonies was resuspended in 0.85% saline and adjusted to an OD_{600nm} of 0.5,

or ~$7.5x10^8$ CFU/mL. For MIC testing, 0.67 µL of this suspension was added per mL of

MHB, giving a final concentration of ~$5x10^5$ CFU/mL, and added to 96-well plates

containing the serially diluted compounds of interest to give a final volume of 100 µL.

Baseline OD_{600nm} reads were taken and plates were incubated stationary for 18h at

37°C and then OD_{600nm} was again measured using a plate reader. For MIC dose-

response curves or checkerboard figures, growth was determined by subtracting

baseline reads, and normalizing the no compound, solvent control to 100%. Reported

MICs were determined by visual inspection of plates and reporting the lowest

concentration at which no growth was visible. For checkerboards, the FIC of each

compound was determined to be its MIC in combination with the other compound

divided by its MIC alone. If a compound did not have an MIC, the next highest two-fold

dilution to that tested was used. The reported FIC indices (FICIs) are the sum of FICs

for the two compounds being tested (67). FICI values ≤ 0.5 were considered synergistic.

MICs in serum were done as above but *P. aeruginosa* PAO1 was inoculated into 50%

human serum (BioIVT), diluted using 1x M9 salts with 25 mM bicarbonate and 1x final

concentration Biolog Redox Dye Mix A (Biolog, Cat.# 74221). This was added to 96-well

plates containing the serially diluted compounds of interest to give a final volume of 200

µL, similar to previous work (17). MICs reported are from biological triplicate data.

Suppressor generation and sequencing

Two biological replicates of *P. aeruginosa* PAO1 were grown overnight in LB, a sample

of each was saved for genome sequencing, and 100µL of culture was plated onto LB

agar plates containing 125µg/mL of RuRed. Two colonies from each biological replicate

that grew on the RuRed plates were picked and struck onto LB agar. A single colony

from each was inoculated in liquid culture and grown overnight, then genome prepped

and cryostocked, and also verified for increased MIC to RuRed in MHB. Genomic DNA

from the wild-type PAO1 biological replicates and 2 mutants from each replicate (6

samples total) were genome sequenced at the Microbial Genome Sequencing Center

(MiGS, Pittsburgh, PA) with Illumina NextSeq 2000 platform and variant called using

BreSeq (68) to identify mutations.

Cloning and expression of PA4394 (mscS-1) and PA4380 (colS) alleles

Genomic DNA from wild-type *P. aeruginosa* PAO1 was used to PCR amplify the *mscS-*

1^{WT} and *colS*^{WT} genes, and genomic DNA from the RuRed suppressors was used to

PCR amplify *mscS-1*^{K73N} *and colS*^{T201P}. *mscS-1*^{WT} and *mscS-1*^{K73N} were amplified with

the primer pair 5' atatggatccatggaattgaactacgaccgactgg 3' and 5'

ataaagctttcagtggtgatggtgatgatggtcggccatcgcgccctgc 3', which incorporated BamHI and

HindIII restriction sites and a C-terminal histidine tag, digested and ligated into pUCP22

digested with BamHI and HindIII. *colS*^{WT} and *colS*^{T201P} were amplified with the primer

pair 5' atatggatccatggagtataagcagagcctcg 3' and 5'

atagtcgactcagtggtgatggtgatgatgagcaacatcgagtaaaacttcg 3', which incorporated BamHI

and SalI restriction sites and a C-terminal 6x histidine tag, digested and ligated into

pUCP22 digested with BamHI and SalI. After ligation, the constructs were

electroporated into *E. coli* Top10 cells and plated on LB agar with 30 µg/mL gentamicin.

Transformants were picked from plates, grown in liquid culture, and plasmids prepped

for sequencing and subsequent transformation into *P. aeruginosa* PAO1 strains. Protein

expression in normalized whole cell samples of *P. aeruginosa* strains overexpressing

the proteins was verified by Western blot. Briefly, after SDS-PAGE and transfer to a

nitrocellulose membrane, recombinant proteins were detected using primary anti-His

antibody (ThermoFisher, Cat. # PA1-983B) and an anti-RNAP antibody (BioLegend,

Cat. # 663104) as a loading control, followed by the secondary antibodies IRDye

800CW Goat anti-Rabbit and IRDye 680RD Goat anti-Mouse (LiCor, Cat. #925-32211

and Cat. #925-68070, respectively).

G. mellonella infections

G. mellonella were reared in-house and larvae weighing approximately 300mg were

used for all experiments. 100 µl of an overnight culture of *P. aeruginosa* PAO1 was

added into fresh media and grown to and OD$_{600}$ of 0.8, then washed twice in PBS and

resuspended to a final density of 1-1.5x10^4 CFU/mL. 10 µl of this was injected into the

last proleg of the larvae using a syringe (U-100 Insulin Syringes, Cat. # 329461, BD)

loaded into a syringe pump (KDS-100, KD Scientific) (69), for a total dose of 100-150

CFU of *P. aeruginosa* PAO1 per larvae. Uninfected controls were injected with PBS.

After 15min, larvae were given a second injection of 10 μl containing drug or vehicle

(water) control and incubated at 37°C. Survival was monitored every 24h for a total of

72h, and larvae were recorded as dead if they failed to respond to touch. Statistics were

performed using GraphPad prism. The statistical significance of ruthenium compound

treatments compared to vehicle control was determined using the Gehan-Breslow-

Wilcoxon test with a Bonferroni correction for multiple comparisons. Using a defined

significance threshold of 0.05, the Bonferroni-corrected significance threshold was

calculated by dividing 0.05 by the number of comparisons. In most cases, this was 3,

yielding a significance threshold of 0.017.

Mouse studies

All mouse studies were conducted according to guidelines set by the Canadian Council

on Animal Care using protocols approved by the Animal Review Ethics Board at

McMaster University under Animal Use Protocol #20-12-43. All animal studies were

performed with 6-10-week-old female CD-1 mice (Envigo). Female mice were used

following previously established models, as well as ease of housing and randomization.

No animals were excluded from analyses, and no sample or effect size assumptions

were made. Sample sizes were selected based on results from pilot experiments.

CD-1 mice were infected intraperitoneally with ~1×10^6 CFU/mL of P. aeruginosa PAO1

with 5% porcine mucin. For the monotherapy, treatment with RuRed (2 mg/kg) or a

vehicle solution (water) was administered intraperitoneally at 1h and 5h post-infection.

For the combination therapy experiment, Ru265 (0.25 mg/kg), rifampicin (10 mg/kg),

vehicle (water), or the combination was administered intraperitoneally at 1h post-

infection. Due to the unknown pharmacokinetics of Ru265, a second treatment of

Ru265 (0.25 mg/kg) was administered to the Ru265 monotherapy and combination

therapy animals at 5h post-infection, and the other groups (vehicle and rifampicin

monotherapy) received vehicle (water) injections. Blood was collected in Li heparin

tubes by facial bleed 10h post-infection. Mice were then euthanized and the spleen,

kidneys, and liver were collected into PBS at necropsy. Organs were homogenized

using a high-throughput tissue homogenizer, serially diluted in PBS, and plated onto

solid LB. Plates were incubated overnight at 37°C, and colonies were quantified to

determine organ load. Statistics were performed using GraphPad Prism software.

Significance was determined by the Mann-Whitney tests and a significance threshold of

0.05 for the RuRed experiments, and the Kruskal-Wallis test for the combination therapy

experiments with a significance threshold of 0.05.

Lysozyme assay for outer membrane permeability

Assay was performed as previously described (40) with modifications. *E. coli* BW25113

was grown overnight (~18h) in MHB and subcultured 1:50 in fresh LB media at 37 ºC

with shaking at 250 rpm to mid-log (OD_{600} ~0.4-0.5). Subcultures were centrifuged and

resuspended in 5 mM HEPES pH 7.2 to OD_{600} ~0.5. A volume of 1 mL of cells was

added to microfuge tubes containing the indicated concentrations of compound and 50

µg/mL lysozyme (Sigma-Aldrich). Tubes were gently inverted to mix and left at room

temperature for ~30 min for any cell debris to settle. A volume of 200 µL was transferred

from each tube to a clear, flat bottom 96-well plate, and OD600 was read using a Tecan

Infinite M1000 Pro plate reader. Controls were included to ensure that the drugs alone

did not lyse cells and to find the baseline levels of cell lysis by lysozyme alone.

Acknowledgements

We thank Dr. J. Dennis and Dr. J. McCutcheon (University of Alberta) for providing

plasmid pUCP22 and *P. aeruginosa* PAO1 and PA14 strains, Tim Cho and members of

the Brown and Raivio laboratories for thoughtful suggestions and discussion of this

work, and Dr. Gerry Wright for providing clinical isolates. Work in the laboratory of

E.D.B. was supported by a Tier I Canada Research Chair award, a Foundation grant

from the Canadian Institutes of Health Research (CIHR; FRN 143215), and a grant from

the Ontario Research Fund (RE09-047). Work in the laboratory of J.J. Wilson was

supported by NSF CAREER Award No. CHE-1750295. Work in the laboratory or T.R.

was supported by operating grants from The National Sciences and Engineering

Research Council (NSERC RGPIN-2021-02710) and The Canadian Institutes of Health

Research (CIHR MOP 142347), and a project grant from the AMR-One Health

Consortium, funded by the Major Innovation Fund program of the Ministry of Jobs,

Economy and Innovation, Government of Alberta. B.S.W. was supported by a CIHR

Postdoctoral Fellowship. J.J. Woods was supported by the National Science Foundation

(DGE-1650441) and the American heart Association (20PRE35120390). All research

was performed prior to K.K. beginning employment at Health Canada (HC). HC was not

involved in the research.

References

1. Theuretzbacher U, Bush K, Harbarth S, Paul M, Rex JH, Tacconelli E, Thwaites GE. 2020.

Critical analysis of antibacterial agents in clinical development. Nat Rev Microbiol 18:286–

298.

2. Tacconelli E, Carrara E, Savoldi A, Harbarth S, Mendelson M, Monnet DL, Pulcini C,

Kahlmeter G, Kluytmans J, Carmeli Y, Ouellette M, Outterson K, Patel J, Cavaleri M, Cox

EM, Houchens CR, Grayson ML, Hansen P, Singh N, Theuretzbacher U, Magrini N, WHO

Pathogens Priority List Working Group. 2018. Discovery, research, and development of

new antibiotics: the WHO priority list of antibiotic-resistant bacteria and tuberculosis.

Lancet Infect Dis 18:318–327.

3. Prasad NK, Seiple IB, Cirz RT, Rosenberg OS. 2022. Leaks in the Pipeline: a Failure

Analysis of Gram-Negative Antibiotic Development from 2010 to 2020. Antimicrob Agents

Chemother 66:e0005422.

4. Theuretzbacher U, Piddock LJV. 2019. Non-traditional Antibacterial Therapeutic Options

and Challenges. Cell Host Microbe 26:61–72.

5. Maura D, Ballok AE, Rahme LG. 2016. Considerations and caveats in anti-virulence drug

development. Curr Opin Microbiol 33:41–46.

6.   Chiang C-Y, Uzoma I, Moore RT, Gilbert M, Duplantier AJ, Panchal RG. 2018. Mitigating

the Impact of Antibacterial Drug Resistance through Host-Directed Therapies: Current

Progress, Outlook, and Challenges. mBio 9:e01932-17.

7.   Farha MA, Brown ED. 2019. Drug repurposing for antimicrobial discovery. Nat Microbiol

4:565–577.

8.   Czaplewski L, Bax R, Clokie M, Dawson M, Fairhead H, Fischetti VA, Foster S, Gilmore

BF, Hancock REW, Harper D, Henderson IR, Hilpert K, Jones BV, Kadioglu A, Knowles D,

Ólafsdóttir S, Payne D, Projan S, Shaunak S, Silverman J, Thomas CM, Trust TJ, Warn P,

Rex JH. 2016. Alternatives to antibiotics-a pipeline portfolio review. Lancet Infect Dis

16:239–251.

9.   Brown ED, Wright GD. 2016. Antibacterial drug discovery in the resistance era. Nature

529:336–343.

10.   Hutchings MI, Truman AW, Wilkinson B. 2019. Antibiotics: past, present and future. Curr

Opin Microbiol 51:72–80.

11.   Aziz Z, Abu SF, Chong NJ. 2012. A systematic review of silver-containing dressings and

topical silver agents (used with dressings) for burn wounds. Burns 38:307–318.

12.   Khansa I, Schoenbrunner AR, Kraft CT, Janis JE. 2019. Silver in Wound Care-Friend or

Foe?: A Comprehensive Review. Plast Reconstr Surg Glob Open 7:e2390.

13.   Rottenberg S, Disler C, Perego P. 2021. The rediscovery of platinum-based cancer

therapy. 1. Nat Rev Cancer 21:37–50.

14.   Sierra MA, Casarrubios L, de la Torre MC. 2019. Bio-Organometallic Derivatives of

Antibacterial Drugs. Chemistry 25:7232–7242.

15.   Frei A. 2020. Metal Complexes, an Untapped Source of Antibiotic Potential? Antibiotics
678   (Basel) 9:E90.

679   16.   Frei A, Zuegg J, Elliott AG, Baker M, Braese S, Brown C, Chen F, G Dowson C, Dujardin
680   G, Jung N, King AP, Mansour AM, Massi M, Moat J, Mohamed HA, Renfrew AK, Rutledge
681   PJ, Sadler PJ, Todd MH, Willans CE, Wilson JJ, Cooper MA, Blaskovich MAT. 2020. Metal
682   complexes as a promising source for new antibiotics. Chem Sci 11:2627–2639.

683   17.   Weber BS, De Jong AM, Guo ABY, Dharavath S, French S, Fiebig-Comyn AA, Coombes
684   BK, Magolan J, Brown ED. 2020. Genetic and Chemical Screening in Human Blood Serum
685   Reveals Unique Antibacterial Targets and Compounds against Klebsiella pneumoniae. Cell
686   Rep 32:107927.

687   18.   Moore CL. 1971. Specific inhibition of mitochondrial Ca++ transport by ruthenium red.
688   Biochem Biophys Res Commun 42:298–305.

689   19.   Woods JJ, Wilson JJ. 2020. Inhibitors of the mitochondrial calcium uniporter for the
690   treatment of disease. Curr Opin Chem Biol 55:9–18.

691   20.   De Stefani D, Raffaello A, Teardo E, Szabò I, Rizzuto R. 2011. A forty-kilodalton protein of
692   the inner membrane is the mitochondrial calcium uniporter. 7360. Nature 476:336–340.

693   21.   Pan X, Liu J, Nguyen T, Liu C, Sun J, Teng Y, Fergusson MM, Rovira II, Allen M, Springer
694   DA, Aponte AM, Gucek M, Balaban RS, Murphy E, Finkel T. 2013. The physiological role
695   of mitochondrial calcium revealed by mice lacking the mitochondrial calcium uniporter. 12.
696   Nat Cell Biol 15:1464–1472.

697   22.   Baughman JM, Perocchi F, Girgis HS, Plovanich M, Belcher-Timme CA, Sancak Y, Bao
698   XR, Strittmatter L, Goldberger O, Bogorad RL, Koteliansky V, Mootha VK. 2011.

Integrative genomics identifies MCU as an essential component of the mitochondrial

calcium uniporter. Nature 476:341–345.

23. Arduino DM, Perocchi F. 2018. Pharmacological modulation of mitochondrial calcium

homeostasis. J Physiol 596:2717–2733.

24. Rimessi A, Vitto VAM, Patergnani S, Pinton P. 2021. Update on Calcium Signaling in

Cystic Fibrosis Lung Disease. Front Pharmacol 12:581645.

25. Broekemeier KM, Krebsbach RJ, Pfeiffer DR. 1994. Inhibition of the mitochondrial Ca2+

uniporter by pure and impure ruthenium red. Mol Cell Biochem 139:33–40.

26. Ying WL, Emerson J, Clarke MJ, Sanadi DR. 1991. Inhibition of mitochondrial calcium ion

transport by an oxo-bridged dinuclear ruthenium ammine complex. Biochemistry 30:4949–

4952.

27. Matlib MA, Zhou Z, Knight S, Ahmed S, Choi KM, Krause-Bauer J, Phillips R, Altschuld R,

Katsube Y, Sperelakis N, Bers DM. 1998. Oxygen-bridged dinuclear ruthenium amine

complex specifically inhibits Ca2+ uptake into mitochondria in vitro and in situ in single

cardiac myocytes. J Biol Chem 273:10223–10231.

28. Emerson J, Clarke MJ, Ying WL, Sanadi DR. 1993. The component of "ruthenium red"

responsible for inhibition of mitochondrial calcium ion transport. Spectra, electrochemistry,

and aquation kinetics. Crystal structure of .mu.-O-[(HCO2)(NH3)4Ru]2Cl3. J Am Chem

Soc 115:11799–11805.

29. Çetiner U, Rowe I, Schams A, Mayhew C, Rubin D, Anishkin A, Sukharev S. 2017.

Tension-activated channels in the mechanism of osmotic fitness in Pseudomonas

aeruginosa. J Gen Physiol 149:595–609.

30. Garvis S, Munder A, Ball G, de Bentzmann S, Wiehlmann L, Ewbank JJ, Tümmler B,

Filloux A. 2009. Caenorhabditis elegans Semi-Automated Liquid Screen Reveals a

Specialized Role for the Chemotaxis Gene cheB2 in Pseudomonas aeruginosa Virulence.

PLoS Pathog 5:e1000540.

31. Dekkers LC, Bloemendaal CJ, de Weger LA, Wijffelman CA, Spaink HP, Lugtenberg BJ.

1998. A two-component system plays an important role in the root-colonizing ability of

Pseudomonas fluorescens strain WCS365. Mol Plant Microbe Interact 11:45–56.

32. Held K, Ramage E, Jacobs M, Gallagher L, Manoil C. 2012. Sequence-verified two-allele

transposon mutant library for Pseudomonas aeruginosa PAO1. J Bacteriol 194:6387–

6389.

33. Woods JJ, Nemani N, Shanmughapriya S, Kumar A, Zhang M, Nathan SR, Thomas M,

Carvalho E, Ramachandran K, Srikantan S, Stathopulos PB, Wilson JJ, Madesh M. 2019.

A Selective and Cell-Permeable Mitochondrial Calcium Uniporter (MCU) Inhibitor

Preserves Mitochondrial Bioenergetics after Hypoxia/Reoxygenation Injury. ACS Cent Sci

5:153–166.

34. Nathan SR, Pino NW, Arduino DM, Perocchi F, MacMillan SN, Wilson JJ. 2017. Synthetic

Methods for the Preparation of a Functional Analogue of Ru360, a Potent Inhibitor of

Mitochondrial Calcium Uptake. Inorg Chem 56:3123–3126.

35. Pereira MF, Rossi CC, da Silva GC, Rosa JN, Bazzolli DMS. 2020. Galleria mellonella as

an infection model: an in-depth look at why it works and practical considerations for

successful application. Pathog Dis 78:ftaa056.

36. Amann R, Donnerer J, Maggi CA, Giuliani S, DelBianco E, Weihe E, Lembeck F. 1990. Capsaicin desensitization in vivo is inhibited by ruthenium red. Eur J Pharmacol 186:169–175.

37. Novorolsky RJ, Nichols M, Kim JS, Pavlov EV, J Woods J, Wilson JJ, Robertson GS. 2020. The cell-permeable mitochondrial calcium uniporter inhibitor Ru265 preserves cortical neuron respiration after lethal oxygen glucose deprivation and reduces hypoxic/ischemic brain injury. J Cereb Blood Flow Metab 40:1172–1181.

38. Stokes JM, MacNair CR, Ilyas B, French S, Côté J-P, Bouwman C, Farha MA, Sieron AO, Whitfield C, Coombes BK, Brown ED. 2017. Pentamidine sensitizes Gram-negative pathogens to antibiotics and overcomes acquired colistin resistance. Nat Microbiol 2:17028.

39. MacNair CR, Brown ED. 2020. Outer Membrane Disruption Overcomes Intrinsic, Acquired, and Spontaneous Antibiotic Resistance. mBio 11:e01615-20.

40. Hancock RE, Wong PG. 1984. Compounds which increase the permeability of the Pseudomonas aeruginosa outer membrane. Antimicrob Agents Chemother 26:48–52.

41. Antimicrobial Resistance Collaborators. 2022. Global burden of bacterial antimicrobial resistance in 2019: a systematic analysis. Lancet 399:629–655.

42. Antunes LCS, Imperi F, Minandri F, Visca P. 2012. In vitro and in vivo antimicrobial activities of gallium nitrate against multidrug-resistant Acinetobacter baumannii. Antimicrob Agents Chemother 56:5961–5970.

43. Goss CH, Kaneko Y, Khuu L, Anderson GD, Ravishankar S, Aitken ML, Lechtzin N, Zhou G, Czyz DM, McLean K, Olakanmi O, Shuman HA, Teresi M, Wilhelm E, Caldwell E,

Salipante SJ, Hornick DB, Siehnel RJ, Becker L, Britigan BE, Singh PK. 2018. Gallium

disrupts bacterial iron metabolism and has therapeutic effects in mice and humans with

lung infections. Sci Transl Med 10:eaat7520.

44. Li F, Collins JG, Keene FR. 2015. Ruthenium complexes as antimicrobial agents. Chem

Soc Rev 44:2529–2542.

45. Bolhuis A, Hand L, Marshall JE, Richards AD, Rodger A, Aldrich-Wright J. 2011.

Antimicrobial activity of ruthenium-based intercalators. Eur J Pharm Sci 42:313–317.

46. Li F, Mulyana Y, Feterl M, Warner JM, Collins JG, Keene FR. 2011. The antimicrobial

activity of inert oligonuclear polypyridylruthenium(II) complexes against pathogenic

bacteria, including MRSA. Dalton Trans 40:5032–5038.

47. Smitten KL, Southam HM, de la Serna JB, Gill MR, Jarman PJ, Smythe CGW, Poole RK,

Thomas JA. 2019. Using Nanoscopy To Probe the Biological Activity of Antimicrobial

Leads That Display Potent Activity against Pathogenic, Multidrug Resistant, Gram-

Negative Bacteria. ACS Nano 13:5133–5146.

48. Heesterbeek D a. C, Martin NI, Velthuizen A, Duijst M, Ruyken M, Wubbolts R,

Rooijakkers SHM, Bardoel BW. 2019. Complement-dependent outer membrane

perturbation sensitizes Gram-negative bacteria to Gram-positive specific antibiotics. Sci

Rep 9:1–10.

49. Maeda T, Horinouchi T, Sakata N, Sakai A, Furusawa C. 2019. High-throughput

identification of the sensitivities of an Escherichia coli ΔrecA mutant strain to various

chemical compounds. J Antibiot 72:566–573.

50. Gutu AD, Sgambati N, Strasbourger P, Brannon MK, Jacobs MA, Haugen E, Kaul RK,

Johansen HK, Høiby N, Moskowitz SM. 2013. Polymyxin resistance of Pseudomonas

aeruginosa phoQ mutants is dependent on additional two-component regulatory systems.

Antimicrob Agents Chemother 57:2204–2215.

51. Wiesmann CL, Zhang Y, Alford M, Thoms D, Dostert M, Wilson A, Pletzer D, Hancock

REW, Haney CH. 2021. The ColR/S two-component system is a conserved determinant of

host association across Pseudomonas species. bioRxiv

https://doi.org/10.1101/2021.12.14.472530.

52. McCarthy RR, Mazon-Moya MJ, Moscoso JA, Hao Y, Lam JS, Bordi C, Mostowy S, Filloux

A. 2017. Cyclic-di-GMP regulates lipopolysaccharide modification and contributes to

Pseudomonas aeruginosa immune evasion. Nat Microbiol 2:17027.

53. Wang NR, Wiesmann CL, Melnyk RA, Hossain SS, Chi M-H, Martens K, Craven K, Haney

CH. 2022. Commensal Pseudomonas fluorescens Strains Protect Arabidopsis from

Closely Related Pseudomonas Pathogens in a Colonization-Dependent Manner. mBio

13:e02892-21.

54. de Weert S, Dekkers LC, Bitter W, Tuinman S, Wijfjes AHM, van Boxtel R, Lugtenberg

BJJ. 2006. The two-component colR/S system of Pseudomonas fluorescens WCS365

plays a role in rhizosphere competence through maintaining the structure and function of

the outer membrane. FEMS Microbiol Ecol 58:205–213.

55. Farha MA, Verschoor CP, Bowdish D, Brown ED. 2013. Collapsing the proton motive force

to identify synergistic combinations against Staphylococcus aureus. Chem Biol 20:1168–

1178.

807 56. MacNair CR, Stokes JM, Carfrae LA, Fiebig-Comyn AA, Coombes BK, Mulvey MR, Brown

808   ED. 2018. Overcoming mcr-1 mediated colistin resistance with colistin in combination with

809   other antibiotics. Nat Commun 9:458.

810 57. Karpel RL, Shirley MS, Holt SR. 1981. Interaction of the ruthenium red cation with nucleic

811   acid double helices. Biophys Chem 13:151–165.

812 58. Rimessi A, Pozzato C, Carparelli L, Rossi A, Ranucci S, De Fino I, Cigana C, Talarico A,

813   Wieckowski MR, Ribeiro CMP, Trapella C, Rossi G, Cabrini G, Bragonzi A, Pinton P. 2020.

814   Pharmacological modulation of mitochondrial calcium uniporter controls lung inflammation

815   in cystic fibrosis. Sci Adv 6:eaax9093.

816 59. Woods JJ, Lovett J, Lai B, Harris HH, Wilson JJ. 2020. Redox Stability Controls the

817   Cellular Uptake and Activity of Ruthenium-Based Inhibitors of the Mitochondrial Calcium

818   Uniporter (MCU). Angew Chem Int Ed Engl 59:6482–6491.

819 60. Pell SD, Sherban MM, Tramontano V, Clarke MJ, Hansongnern K, Krause RA. 1989. cis-

820   Tetraamminedihaloruthenium(III) Halide Complexes, p. 65–68. *In* Inorganic Syntheses.

821   John Wiley & Sons, Ltd.

822 61. Evans IP, Everett GW, Sargeson AM. 1976. The stabilization of .alpha.- and .beta.-

823   diimines: hexaammineruthenium(III), -osmium(III), and -platinum(IV) condensed with

824   .alpha.- and .beta.-diketones. J Am Chem Soc 98:8041–8046.

825 62. Allen AD, Senoff CV. 1967. Preparation and infrared spectra of some ammine complexes

826   of ruthenium(II) and ruthenium(III). Can J Chem 45:1337–1341.

827   63.   Allen AD, Bottomley F, Harris RO, Reinsalu VP, Senoff CV, Badger DW. 1970.

828   Pentaammine(nitrogen)ruthenium(II) Salts and Other Ammines of Ruthenium, p. 1–8. *In*

829   Inorganic Syntheses. John Wiley & Sons, Ltd.

830   64.   Durig JR, McAllister WA, Willis JN, Mercer EE. 1966. Far infrared spectra of potassium

831   and ammonium pentahalonitrosylruthenates. Spectrochimica Acta 22:1091–1100.

832   65.   Cleare MJ, Griffith WP. 1970. Polynuclear nitrido-complexes of osmium, ruthenium, and

833   iridium. J Chem Soc A 1117–1125.

834   66.   Clark RJH, Franks ML, Turtle PC. 1977. Vibrational spectra, resonance Raman spectra,

835   and electronic spectra of the .mu.-oxo-decachlorodiruthenium(IV) ion. J Am Chem Soc

836   99:2473–2480.

837   67.   Lambert RJW, Lambert R. 2003. A model for the efficacy of combined inhibitors. J Appl

838   Microbiol 95:734–743.

839   68.   Deatherage DE, Barrick JE. 2014. Identification of mutations in laboratory-evolved

840   microbes from next-generation sequencing data using breseq. Methods Mol Biol

841   1151:165–188.

842   69.   Ramarao N, Nielsen-Leroux C, Lereclus D. 2012. The insect Galleria mellonella as a

843   powerful infection model to investigate bacterial pathogenesis. J Vis Exp e4392.

844