A genome catalogue of lake bacteria across watershed land use gradients at a continental scale

Authors

Rebecca E. Garner[1,4], Susanne A. Kraemer[2,4], Vera E. Onana[1,4], Maxime Fradette[3], Marie-Pierre Varin[3], Yannick Huot[3,4], David A. Walsh[1,4]#

Affiliations

[1]Department of Biology, Concordia University, Montreal, Quebec, Canada

[2]Environment and Climate Change Canada, Montreal, Quebec, Canada

[3]Département de géomatique appliquée, Université de Sherbrooke, Sherbrooke, Quebec, Canada

[4]Groupe de recherche interuniversitaire en limnologie, Quebec, Canada

#Address correspondence to David A. Walsh, david.walsh@concordia.ca.

Abstract

Lakes are heterogenous ecosystems inhabited by a rich microbiome whose genomic diversity is poorly

defined. We present a continental-scale study of metagenomes representing 6.5 million km^2 of the most

lake-rich landscape on Earth. Analysis of 308 Canadian lakes resulted in a metagenome-assembled

genome (MAG) catalogue of 1,008 mostly novel bacterial genomospecies. Lake trophic state was a

leading driver of taxonomic and functional diversity among MAG assemblages, reflecting the responses of

communities profiled by 16S rRNA amplicons and gene-centric metagenomics. Coupling the MAG

catalogue with watershed geomatics revealed terrestrial influences of soils and land use on assemblages.

Agriculture and human population density were drivers of turnover, indicating detectable anthropogenic

imprints on lake bacteria at the continental scale. The sensitivity of bacterial assemblages to human

impact reinforces lakes as sentinels of environmental change. Overall, the LakePulse MAG catalogue

greatly expands the freshwater genomic landscape, advancing an integrative view of diversity across

Earth's microbiomes.

Main

Freshwater bacteria are a diverse component of lake ecosystems and are central to biogeochemical

cycles and water quality[1–3]. Collectively, Earth's millions of lakes exhibit immense environmental

heterogeneity, reflecting a range of influences from regional variations in geology, landform, and climate

to local compositions of catchment runoff[4]. Lakes integrate terrestrial carbon and other materials from the

catchment and airshed, to an extent that lakes are considered an axis of the global carbon cycle[5] and

sentinels of environmental change[6]. The growing number of metagenomic studies is providing important

insights into the ecology and evolution of freshwater bacteria in the context of changing environmental

conditions[7]. However, most studies are focused on a limited number of lakes, restricting our

understanding on how environmental variation influences the structure and function of bacterial

assemblages across lake ecosystems. Elucidating freshwater bacterial genomic diversity is critical given

the myriad anthropogenic pressures on lakes. Lake warming[8], eutrophication[9], deoxygenation[10],

salinization[11], chemical contamination, and other emergent stressors[12] are expected to modify the

freshwater microbiome, but in ways that we are only beginning to understand[13].

Genome-resolved metagenomics enables the reconstruction of composite genomes from

microbial populations[14]. MAGs supply reference genomes for uncultivated taxa, revealing enormous

microbial dark matter[15]. Currently, MAG analysis is a standard approach to link functions to populations in

complex communities. Lake studies applying genome-resolved metagenomics have brought insights into

the evolution[16,17], metabolism[18–20], and succession[21,22] of freshwater bacteria. Recently, the approach has

been applied at larger scale to generate genomic catalogues from marine[23], terrestrial[24], and host-

associated microbiomes[25–27], culminating in a genome catalogue of Earth's microbiomes (GEM)[28].

Although impressive in its collection of habitats, GEM encompasses very few freshwater lakes (<30),

compared to the immense abundance and heterogeneity worldwide. Further efforts have begun to

elucidate lake genome diversity in stratified water columns[29]. An essential next step is to expand the

representation of freshwater lakes in the global genomic catalogue.

Here, we introduce a genome-resolved perspective of freshwater bacterial diversity across 6.5-

million km^2 of Canada, the most lake-rich landscape on Earth[30]. This study is a component of the NSERC

Canadian LakePulse Network[31], a large-scale coordinated effort to assess the health status of lakes using

standardized sampling and analysis methods with a major focus on microbial communities[32–40]. Through

metagenome co-assembly and binning, we reconstructed 1,184 high- and medium-quality MAGs from

308 lakes reflecting expansive gradients in limnological and watershed conditions. The LakePulse MAG

catalogue was then used to investigate the structure and function of lake bacterial assemblages at

continental scale. In addition to clarifying the functional repertoires of freshwater taxa, we pursued a novel

analysis of the metabolic variation among MAG assemblages across lake physicochemical and other

environmental gradients. The consideration of geomatics variables in the LakePulse survey further

allowed us to report on watershed factors influencing bacterial assemblages across the land-water

interface, including detectable land use effects signaling evidence of human impact. In resolving genomic

diversity from the largest representation of lakes across a temperate to subarctic range, the LakePulse

MAG catalogue forms a key resource for investigating the freshwater microbiome on a rapidly changing

planet.

Results

The LakePulse MAG catalogue

To produce the MAG catalogue, we generated metagenomes from the surface waters of 308 lakes in 12

Canadian ecozones (**Fig. 1**; **Fig. S1**; **Table S1**). Lakes represented a wide range of physicochemical

conditions, productivity, morphometry, climate, and land use within watersheds[41] (**Fig. S2***)*. At the

continental scale (43 – 68 °N, 62 – 141 °W), a full range of ultraoligotrophic to hypereutrophic lakes were

present in the survey (**Fig. 1A**). Comparison of lake features by principal component analysis (PCA)

revealed large-scale spatial patterns in lake and watershed characteristics (**Fig. 1B**). Western Canadian

lakes were the deepest and most oligotrophic on average, and often set in watersheds with high

proportions of natural landscapes and harvested forests. Northern Canadian lakes experienced the

coldest climates and lowest land use pressures. Central Canada was comprised of the agriculturally

intensive Prairies and Boreal Plains ecozones; compared to other Canadian regions, central Canadian

lakes were generally shallow, alkaline, nutrient- and ion-rich, and highly productive. Eastern Canadian

lakes had the warmest surface waters and on average the most extensive built environments in their

watersheds. Agriculture was also a common land use in eastern Canada, particularly within the

Mixedwood Plains ecozone. Lakes in watersheds with the largest population centres were in the

Vancouver and Toronto metropolitan areas within the Pacific Maritime and Mixedwood Plains ecozones,

respectively.

We aimed to capture a broad phylogenetic and phenotypic diversity of microorganisms (e.g.,

organism size and lifestyle) in the LakePulse survey. To this end, lake metagenomes were generated

from biomass collected across a wide size range (0.22 – 100 µm). Eleven metagenome co-assemblies

representing the 12 ecozones were generated (Boreal and Taiga Cordilleras lakes were combined).

Though computationally intensive to produce, co-assemblies facilitated differential coverage binning,

ensuring higher-quality MAGs[42]. In total, contig binning of each co-assembly recovered 1,184 MAGs

exceeding MIMAG medium-quality standards[43] through an additional <10% strain heterogeneity criterion

(**Table S2**). Between 28 – 233 MAGs were generated for each ecozone (**Fig. S3**). MAGs were

dereplicated across ecozones (ANI ≥95%), resulting in a genomospecies-level set of 1,008 MAGs. MAGs

were categorized as 136 high- and 872 medium-quality drafts based on MIMAG standards[43], excluding

rRNA gene requirements (**Fig. 1C**). MAGs ranged widely in size (0.35 – 10.39 Mbp), GC content (23.1 –

75.9%), and coding density (78.4 – 96.8%) (**Table S2**). Single metagenome assemblies produced only

563 bins passing our quality control (**Table S1**); these represented just 202 lakes and 476

genomospecies, over one-third of which were already present in the catalogue. This comparison

validated our more intensive co-assembly approach to reconstructing lake bacterial genomic diversity.

To map MAG biogeography across lakes, we calculated the central 80% truncated average

depths of coverage (TAD$_{80}$) from fragment recruitment[21]. Based on TAD$_{80}$, observed richness within lakes

varied between 11 – 172 MAGs, with diversity hotspots tending to occur in the shallow, nutrient-rich lakes

of the Boreal Plains and Prairies ecozones of central Canada (**Fig. 1D**). A species accumulation curve

showed that hundreds of lakes were required to adequately sample genomes across the heterogenous

Canadian landscape, highlighting the importance of analyzing lake metagenomes across large

geographic and environmental gradients to capture the breadth of local lake bacterial diversity and

ecosystem function (**Fig. 1E**). MAGs captured on average 7.1% (range 0.9 – 36.0%) of metagenome

reads per lake (**Fig. 1F**), signifying that the MAG catalogue is a reasonable but incomplete representation

of bacterial genome diversity in the lakes. Currently, the LakePulse MAG catalogue represents a

significant expansion in the availability of freshwater genomes and range of lake systems to date.

Phylogenetic and metabolic diversity of novel lake bacterial species

The LakePulse MAG catalogue represented a phylogenetically rich set of genomes (**Fig. 2A**; **Table S2**)

exhibiting high taxonomic novelty. Overall, 96% of MAGs were novel candidate species and over a third

(349) belonged to novel genera (**Fig. 2B**), substantially expanding the known genomic diversity of lake

bacteria. The largest collection of MAGs was recovered from *Bacteroidota*, a group that plays a major role

in organic particle degradation in marine settings[44] but is much less studied in freshwater systems[3]. A

third of *Bacteroidota* MAGs (80 of 275) represented new genera (**Table S3**), presenting a rich genomic

resource for future targeted research on freshwater *Bacteroidota* ecology and metabolism, including

potential zooplankton associations[45]. *Actinobacteriota* and *Gammaproteobacteria* were the next most

MAG-rich groups and included lineages of common and abundant freshwater bacteria (e.g., *Rhodoluna*,

Polynucleobacter, and *Limnohabitans*). Other freshwater taxa that were well represented in the MAG

catalogue included *Verrucomicrobiota*, *Planctomycetota*, *Patescibacteria*, and *Myxococcota*. Intriguingly,

we identified 38 MAGs from *Bdellovibrionota*, almost all of which represented new genera (**Table S3**).

Bdellovibrionota are described as obligate predators of Gram-negative bacteria[46] and their role in

freshwater food webs warrants further attention.

To assess whether the MAGs captured a representative phylogenetic diversity of lake bacteria,

we compared taxonomy between MAGs and 16S rRNA gene amplicon sequence variants (ASVs). MAGs

and ASVs overlapped in their 10 most diverse phyla, though ASVs exhibited an expected long tail of rare

diversity (**Fig. S4A**). Moreover, the three most MAG-rich orders (*Chitinophagales*, *Burkholderiales*, and

Flavobacteriales) were in the top four among ASVs (**Fig. S4B**). The largest differences in diversity across

datasets were within *Bacteroidota* and *Actinobacteriota*, which were slightly more represented as a

percentage of MAGs. In contrast, *Patescibacteria* exhibited higher ASV diversity and were potentially

underrepresented in the MAGs because of low or divergent marker gene content[47]. Overall, the

LakePulse MAG catalogue encompasses broad phylogenetic diversity reflective of 16S rRNA genes,

making it a useful resource for further taxon-focused research.

Bacteria contribute to lake ecosystem function through a diversity of energy-capturing and carbon

cycling metabolisms that regulate biogeochemical cycles. To provide a broad overview of MAG metabolic

diversity, we explored the distributions of select metabolic modules across the MAG catalogue (**Fig. 2C**;

Table S2). Oxygenic photosynthetic *Cyanobacteria* and anoxygenic photosynthetic *Chlorobia* were

present, while bacteriorhodopsins were identified in *Thermoleophilia*. Photosystem II (PS-II) was the most

widespread phototrophic module in the MAGs, including in numerous *Alphaproteobacteria*,

Gammaproteobacteria, and *Gemmatimonadota* MAGs. In combination, PS-II-encoding MAGs were

identified in 95% of lakes, indicating a diverse bacterial contribution to freshwater phototrophy. We

identified three of the six known autotrophic carbon fixation pathways[48], including the Calvin-Benson-

Bassham (CBB) cycle, the 3-hydroxypropionate/4-hydroxybutyrate (3HP/4HB) cycle, and the glyoxylate

assimilation pathway. The CBB cycle was linked to *Proteobacteria* and *Cyanobacteria* MAGs present in

127 lakes (42%), demonstrating that non-algal carbon fixation is also common in lakes.

Heterotrophic degradation of complex algal and terrestrial organic matter is a central ecological

function of lake bacteria. An abundance of *Bacteroidota* and *Gemmatimonadota* MAGs were equipped

with SusCD-mediated glycan transport systems (**Fig. 2C**; **Table S2**), suggesting broad potential for

polysaccharide transport and breakdown. Aromatic compound degradation pathways associated with the

metabolism of plant-derived lignin were common within *Alphaproteobacteria* and *Gammaproteobacteria*,

but also broadly distributed across the MAG phylogeny (**Fig. 2C**; **Table S2**). In addition to heterotrophy,

many *Burkholderiales* MAGs appeared to augment their energy metabolism through Sox-mediated sulfur

oxidation pathways. We did not find evidence for methane or ammonia oxidation within the MAG

catalogue. However, energy conservation via methanol oxidation and redox cycling of nitrogen oxides

was evident. Overall, these findings highlight that the MAG catalogue captures a broad diversity of

metabolisms for in-depth comparative analysis on functional contributions to lake ecosystems.

Lake conditions shape the structure and function of bacterial assemblages

Biogeographical mapping across Canadian lakes revealed that MAGs occupied on average 20 lakes (7%)

and as many as 190 lakes (63%) (**Fig. 3A**). Thus, the co-assembly-resolved MAG catalogue appeared to

comprise relatively widely distributed bacteria. *Patescibacteria* MAGs were among those with the most

restricted ranges, suggesting relatively specialist lifestyles, perhaps dependent on host ranges[49] or input

from groundwater environments where they thrive[50]. Notable biogeographic patterns were observed when

MAGs were aggregated at the class to phylum levels (**Fig. S5**; **Fig. S6**). Phyla previously determined to

be common in lakes (e.g., *Bacteroidota*, *Proteobacteria*, and *Actinobacteriota*) were broadly distributed

across the continent. Other groups were relatively restricted to distinct regions. For example, *Bacilli* and

Campylobacteria MAGs were more common in Prairies lakes than elsewhere, perhaps in response to

high nutrients or extensive agriculture within watersheds (**Fig. S5**; **Fig. S6**).

We next used the MAG catalogue to investigate biogeographic patterns of taxonomic and

functional diversity across continental-scale environmental gradients. Taxonomic variation among MAG

assemblages was correlated with lake trophic state, indexed by total phosphorus (TP) concentration

(correlations with principal coordinate analysis [PCoA] axes: r_1 = -0.32; r_2 = 0.33) (**Fig. 3B**). ASV

community taxonomic profiles were likewise distinguished by trophic state (r_1 = 0.41) (**Fig. S7A**),

demonstrating that MAGs were reasonably representative of bacterial community variation. To assess

functional variation, we generated lake functional profiles by aggregating the collection of metabolic

KEGG orthologs (KOs) encoded by the MAG assemblage within each lake. Bacterial taxonomic and

functional dissimilarities exhibited a positive, but variable relationship (**Fig. 3C**). PCoA of functional

dissimilarities revealed a distribution of MAG assemblages along axis 1 related to lake trophic state (r_1 =

0.23) (**Fig. 3D**) that reflected the relationship exhibited by metagenome functional profiles (r_1 = 0.40; r_2 = -

0.33; metagenome functional profiles form the basis of a dedicated study [Onana *et al.*, in prep.]) (**Fig.**

S7B). These results suggest that nutrient availability and productivity play leading roles in structuring the

taxonomic and functional composition of lake bacterial assemblages at continental scale.

To further elucidate the factors influencing bacterial diversity across lakes, we evaluated the

importance of lake physicochemistry, morphometry, geography, and climate conditions using generalized

dissimilarity models (GDMs)[51]. GDMs of membership (Jaccard dissimilarities) explained more deviance

compared to structure (Bray-Curtis dissimilarities) (**Table S4**). As membership is more stable within lake

assemblages than structure due to the seasonal succession of bacterial populations, it follows that

membership is better predicted across large spatial scales. Lake physicochemistry explained the largest

amounts of taxonomic (MAGs 43.9%; ASVs 48.3%) and functional (MAGs 19.4%; metagenomes 32.6%)

variation among MAG assemblages and community profiles (**Fig. 3E**; **Table S4**). The consistency in

deviance explained by taxonomic or functional models showed that MAG assemblages reliably reflected

the significant environmental filtering of the broader community. The influence of trophic state was evident

as taxonomic and functional turnover across MAG and broader assemblages was consistently responsive

to chlorophyll-*a* (Chl-*a*) concentration, a measure of algal biomass (**Fig. 3F**; **Fig. S8A**). MAG taxonomic,

but not functional, turnover was significantly related to TP (**Fig. S8A**), suggesting metabolic differences

among bacterial assemblages are driven more by the composition of organic matter produced by

phytoplankton than by direct nutrient availability. Turnover was most rapid at low Chl-*a* and TP,

suggesting rapid shifts in bacterial assemblage composition across oligotrophic to mesotrophic lakes (**Fig.**

3F).

In addition to the effects of trophic state, GDMs provided insights into other physicochemical

factors shaping lake bacterial assemblages at continental scale. MAG and community taxonomic turnover

was significantly associated with pH, and MAG functional turnover was additionally predicted by lake

surface temperature (**Fig. 3G**; **Fig. S8A**). Temperature and pH are often considered master variables

driving large-scale spatial patterns in marine[52] and soil[53] communities, respectively. As aquatic systems

that are heavily influenced by their terrestrial surroundings, it follows that lake bacterioplankton

assemblages are influenced by a combination of temperature and pH.

Remarkably, calcium concentration was the strongest predictor of both taxonomic and functional

turnover across MAG assemblages and community-level taxonomy (**Fig. 3H**; **Fig. S8A**). A shift in

bacterial ecotype diversity along a freshwater calcium gradient was previously reported[54], but how

calcium directly influences assemblage composition is unclear. The effect is perhaps indirectly linked to

interactions among plankton, as low calcium was previously shown to limit zooplankton growth[55],

potentially eliciting trophic cascades through the lake food web. Further investigation of the mechanisms

and implications of the relationship between calcium and bacterial assemblages is warranted, particularly

as lake calcium decline is a legacy effect of emissions-linked acid rain[56], while calcium levels may be

increasing elsewhere through winter road salt application[57].

The demonstration that lake physicochemistry shapes the functional composition of MAG

assemblages prompted a deeper analysis focused on specific metabolic categories. We generated GDMs

for different KEGG metabolic categories implicated in energy and nutrient cycling. As with the full

functional profiles, lake physicochemistry consistently explained more deviance than morphometry,

geography, or climate (**Fig. 4A**). Interestingly, explained deviance varied by metabolic category, and

variation within carbohydrate metabolism was the best predicted by physicochemistry (**Fig. 4A**).

Significant predictors of carbohydrate metabolism included Chl-*a* concentration and lake colour (**Fig. 4B**;

Fig. S8B). Lake colour reflects terrestrial organic matter input from the watershed. Hence, variation in

carbohydrate metabolism across bacterial assemblages appears to be linked to the quantity and

composition of autochthonous (algal) and allochthonous (plant) organic matter in lakes. We note that lipid

metabolism was the only additional metabolic category where turnover was predicted by Chl-*a* and colour

(**Fig. 4B**; **Fig. S8B**). Subsequent PCoAs of carbohydrate and lipid metabolisms in lakes showed strong

relationships with trophic state and large variation across the eutrophic to hypereutrophic spectrum (**Fig.**

4C-D). Overall, these results suggest that a diverse and dynamic organic matter pool characterized by

punctuated inputs or resuspensions of nutrients in eutrophic to hypereutrophic systems may be a key

driver of metabolic diversity within the lake microbiome.

The metabolism of complex carbohydrates of aquatic or terrestrial origin depends on initial

degradation through carbohydrate-active enzymes (CAZymes). MAGs contained >13,000 genes assigned

to 129 glycoside hydrolase (GH) gene families, implicated in cleaving the glycosidic bonds of diverse

polysaccharides. Like carbohydrate metabolism (KEGG category), GH repertoires within bacterial

assemblages were linked to lake physicochemistry (**Fig. 4A**). 41% of GH genes were identified in

Bacteroidota MAGs (**Fig. 4E**) and were often encoded in polysaccharide utilization loci (PULs) (**Fig. S9**),

demonstrating that freshwater *Bacteroidota* are central to complex organic matter degradation

comparable to their role in marine and gut ecosystems[58]. *Verrucomicrobiota* MAGs were also replete with

GH genes (18%), expanding on previous observations that freshwater *Verrucomicrobiota* are central

players in organic matter degradation[59]. PCA of GH distributions across MAGs demonstrated a strong

taxonomic structure for carbohydrate-processing potential in lake bacteria (**Fig. 4F**). *Bacteroidota*

displayed the most distinct GH family repertoires, characterized by an enrichment in GH74 implicated in

cellulose and xyloglucan degradation and GH20 implicated in anhydromuropeptides recycling and chitin

degradation. *Proteobacteria* and *Bdellovibrionota* displayed overlapping GH repertoires distinct from other

taxonomic groups and driven by GH23, GH102, and GH103, which are all lysozymes implicated in

peptidoglycan degradation, aligning with potential predatory activity against other bacteria[60]. The

thousands of GH genes encoded in the LakePulse MAG catalogue represent a rich resource to further

examine organic matter degradation in freshwaters and to expand on previous global explorations of

CAZyme diversity across ecosystems[61,62].

Connecting lake bacterial diversity to watershed characteristics and human impact

While the influence of lake physicochemistry on bacterial ecology has received considerable attention and

is elaborated upon in this study, terrestrial influences from the surrounding watershed are less

understood. The LakePulse survey was complemented with geomatics descriptions of lake watersheds,

including soils and land use[31]. Here, we coupled the geomatics data with the MAG catalogue to provide

continental-scale insights into watershed impacts on lake bacterial assemblages across the land-water

interface. Soil properties explained significant GDM deviance in both taxonomic (MAGs 12.1%; ASVs

23.2%) and functional (MAGs 16.3%; metagenomes 9.2%) turnover (**Fig. 5A**). Notably, across all GDMs

developed in this study, the importance of soil was second only to lake physicochemistry, providing

evidence for significant terrestrial influence on lake bacterial assemblage structure and function. Turnover

across MAG and community profiles was most strongly associated with soil pH and the volumetric fraction

of coarse fragments, a variable describing soil texture (**Fig. 5B**; **Fig. S8A**). Soil texture governs water

retention, while pH determines the mobility of ions and nutrients[63]. Soil attributes thereby direct the

composition and release of catchment runoff, which in turn modifies lake physicochemistry. Hence,

terrestrial inputs from soils appear to influence lake bacterial diversity in a moderately predictable manner

at continental scale.

Next, we investigated if bacterial assemblages could be predicted by human impact on the

watershed. There is accumulating evidence that the taxonomic diversity of lake bacteria is influenced by

land use at large spatial scales[32,64,65], yet effects on function are unclear. We were particularly interested

in how agriculture (**Fig. 5C**) and human population density (**Fig. 5D**) within watersheds may shape

bacterial diversity, as each contribute to the eutrophication and chemical contamination of lakes. Land

use GDMs explained a lesser, but still significant, amount of taxonomic (MAGs 6.8%; ASVs 9.6%) and

functional (MAGs 5.4%) turnover compared to soils. There was no statistically significant response of

metagenome function to land use gradients, and given that MAGs are an incomplete representation of the

broader community, signals in MAG functional turnover should be interpreted cautiously. Based on

evidence from GDMs, MAG assemblages are an informative subset through which to investigate – and

potentially indicate – anthropogenic pressures on watersheds altering lake bacterial assemblages across

the land-water interface. The functional effects of land use on MAG assemblages may reflect the

responses of taxonomic composition, yet the coupling of taxonomy and function in MAGs present a dual-

faceted resource to study human impact.

Of all human impact variables, agriculture exhibited a significant effect on both MAG taxonomic

and functional turnover, while human population density exhibited a significant effect on functional

turnover (**Fig. 5E**). Intriguingly, when analyzing specific metabolic categories, xenobiotic metabolism was

the one most strongly explained by land use (**Fig. 5E**). Consistent with general functional turnover (**Fig.**

5F; **Fig. S8A**), human population density followed by agriculture were the strongest predictors of turnover

within xenobiotic metabolism (**Fig. 5G**; **Fig. S8B**). In both cases, the most rapid turnover occurred where

human populations were densest (1,000 – 3,000 individuals/km^2), corresponding to the few lakes

sampled within the Vancouver and Toronto metropolitan areas (**Fig. 5D**). The underlying effect of

population density appeared complex: bacterial assemblages within highly populated landscapes were

characterized by a lower diversity of KOs linked to xenobiotic metabolism. Moreover, PCoA showed that

although xenobiotic metabolism in lakes within high-population watersheds were distinct from those within

low-population watersheds, there was also large variation in xenobiotic metabolism among lakes near

population centres (**Fig. S10**). As many of the detected KOs within xenobiotic metabolism were involved

in the degradation of aromatic compounds, lake bacteria may be capable of metabolizing aromatic

environmental pollutants, such as agricultural pesticides or industrial wastes[66]. The metabolism of land-

derived pollution by lake bacteria merits further investigation, considering watersheds are increasingly

converted to agriculture and urban development[67], and altered precipitation patterns are increasing

runoff[68]. Overall, these results suggest that the size and activities of human populations have a detectable

and significant influence on the functional diversity encoded in the freshwater microbiome, as represented

in the LakePulse MAG catalogue.

Discussion

The LakePulse MAG catalogue, compiled from a continental-scale sampling of hundreds of lakes,

addresses an urgent need to probe the microbiome underpinning freshwater ecosystems under rapid and

accelerating anthropogenic change. This study has substantially expanded the number and diversity of

genomes from freshwater ecosystems and demonstrated that lake bacteria are equipped to metabolize

diverse substrates originating from both within lakes and their catchments. Trophic state emerged as a

leading driver of taxonomic and functional variation among lake bacterial assemblages at continental

scale, with implications for understanding bacterial responses to ongoing anthropogenic eutrophication of

lakes[9]. The functional responses of MAG assemblages to land use pressures present a useful system for

investigating human impact on the lake microbiome across the land-water interface. The links between

lake bacterial diversity and watershed variables, more stable over short timescales than lake

physicochemical conditions, could be leveraged to predict freshwater assemblages using remote sensing

data to decrease the dependence on *in situ* surveys and monitoring[37,40]. Our findings make clear that lake

ecosystems cannot be studied in isolation of their watersheds in an era of intensifying land use[67].

Through its construction and analysis, the LakePulse MAG catalogue makes strides toward

resolving the rich and spatially heterogeneous freshwater fraction of the global microbiome. Among its

potential future utilities are as an essential resource for mapping the distributions of freshwater bacteria,

tracing evolutionary transitions within the land-ocean continuum, and tracking microbial dispersal and

gene transfer from sources of pollution (e.g., xenobiotics; antibiotic resistance[39]) in human-altered

watersheds. More lake bacterial diversity and ecosystem function remain to be captured in stratified water

columns[20,29], sediments[69,70], across seasons[71], and potentially through deeper metagenome binning[21].

Finally, microdiversity[72] contained in the MAG catalogue may provide insights into intraspecific metabolic

adaptations to the environmental heterogeneity among lakes and watersheds, including under future

scenarios of environmental change.

Methods

Lake sampling

Lake sampling was conducted by the Natural Sciences and Engineering Research Council (NSERC)

Canadian Lake Pulse Network (LakePulse) between 2017 – 2019[31]. Sampling was conducted in late June

to mid-September, when some lakes are thermally stratified. To capture the natural and human-mediated

heterogeneity in lake and watershed conditions, lakes were selected across three lake area categories

(small: $0.1 - 0.5$ km^2, medium: $0.5 - 5$ km^2, and large: $5 - 100$ km^2) and human impact categories (low,

moderate, and high as determined by land use types and coverage of the watershed) in 12 terrestrial

ecozones of Canada.

Integrated surface water samples were collected over the euphotic zone up to 2 m below the

surface at the site of maximum lake depth using an integrated tube sampler[73]. The euphotic zone was

estimated to be twice the Secchi disk depth. Water samples were stored in chilled coolers until they were

filtered onshore on the same day. Water was pre-filtered through 100 μm nylon mesh and vacuum-filtered

through 0.22 μm Durapore membranes in glass funnels at a maximum pressure of 8 inHg. 500 mL of

water was filtered or until the filter nearly clogged. All sampling equipment was acid-washed and rinsed

with lake water prior to use. Filters were stored in sterile cryovials at -80 °C.

Environmental data

Six categories of environmental explanatory variables were selected for analysis: (1) geography, (2) lake

morphometry, (3) surface water physicochemistry, (4) watershed surface soil, (5) land use, and (6)

climate. Geography variables included latitude, longitude, and altitude. Lake morphometry variables

included lake area, circularity, volume, maximum depth, discharge, water residence time, watershed area,

lake-to-watershed area ratio, and watershed slope within 100 m of the shoreline (data on volume,

discharge, residence time, and slope were accessed from HydroLakes v. 1.0[30]). Shoreline length and

average depth were strongly colinear with other morphometry variables ($r ≥ 0.7$) and therefore excluded

from the analysis. Surface water physicochemical variables included temperature, pH, colour, and

concentrations of Chl-*a*, DIC, DOC, TN, TP, calcium, chloride, magnesium, potassium, sodium, and

sulfate. Missing physicochemical data were replaced with ecozone medians. Watershed soil variables

estimated for the top 0 – 5 cm soil depth interval were accessed from SoilGrids250m[74] and included bulk

density of the fine earth fraction, cation exchange capacity, volumetric fraction of coarse fragments,

proportions of clay, sand, and silt particles in the fine earth fraction, total nitrogen, pH, soil organic carbon

content in the fine earth fraction, and organic carbon density. Land use variables were calculated as

fractions of watershed area not covered by water and included crop agriculture, pasture, forestry, built

development, human population density, livestock density, and poultry density. Climate variables

measured over the seven days prior to lake sampling were accessed from ERA5-Land hourly reanalysis[75]

and included mean air temperature, total precipitations, mean net solar radiation, mean wind speed, and

ice disappearance day for the year of sampling. Lake trophic state was categorized by TP concentrations

according to the Canadian Water Quality guidelines: ultraoligotrophic (<4 µg/L), oligotrophic (4 – 10 µg/L),

mesotrophic (10 – 20 µg/L), mesoeutrophic (20 – 35 µg/L), eutrophic (35 – 100 µg/L), and hypereutrophic

(>100 µg/L)[76].

DNA extraction and metagenome sequencing

DNA was extracted from filters using the DNeasy PowerWater kit (QIAGEN) according to the

manufacturer's instructions supplemented by the optional addition of 1 µL ribonuclease A and 30 min

incubation at 37 °C. DNA was submitted to Genome Quebec for library preparation using the NEBNext

Ultra II DNA Library Prep Kit (New England Biolabs) and 150 bp paired-end shotgun sequencing on an

Illumina NovaSeq 6000 platform.

Metagenome assembly and binning

Adapter-clipping and quality-trimming of raw reads were performed in Trimmomatic v. 0.36 using a Phred

quality score of 33 and default settings[77]. We used a co-assembly approach to leverage the differential

depth of coverage across contigs and metagenomes and thus maximize the recovery of genomes

common to the same ecozone. Metagenomes were co-assembled in MEGAHIT v. 1.2.7 using k-mer

lengths 27, 37, 47, 57, 67, 77, 87 and a minimum k-mer multiplicity filter of two[78]. Because only three

lakes were sampled in the Taiga Cordillera and these bordered the Boreal Taiga, an ecozone with similar

climate, landform, and vegetation, metagenomes from these two ecozones were combined into one co-

assembly.

Local alignment of metagenome reads to co-assemblies was performed with the Burrows-

Wheeler Aligner (BWA) v. 0.7.17 using the BWT-Smith-Waterman (BWTSW) index algorithm and BWA-

Maximal Exact Matches[79]. Binning was performed in MetaBAT v. 2.12.1 using the default minimum

scaffold length of 2,500 bp and based on the average differential coverage of each contig determined by

the jgi_summarize_bam_contig_depths script implementing a 97% sequence identity threshold[80]. Bin

completeness, contamination, and strain heterogeneity were estimated in CheckM v. 1.0.7[81]. Bins scoring

≥50% completeness, <10% contamination, and <10% strain heterogeneity were retained as MAGs.

MAGs were dereplicated at average nucleotide identity (ANI) ≥95% and alignment coverage ≥10% in

dRep v. 3.2.0 implementing gANI as the second clustering algorithm[82]. In cases where more than one

MAG belonged to the same genomospecies, the highest-quality MAG was selected as the

genomospecies representative to be included in the final MAG catalogue and subsequent analysis. MAGs

were categorized according to MIMAG standards by which high-quality drafts display completeness

>90%, contamination <5%, and ≥18/20 tRNA genes (16S, 23S, and 5S rRNA gene presence criteria were

omitted) and medium-quality drafts display completeness ≥50% and contamination <10%[43].

Functional analysis

Protein-coding genes were annotated in Prokka v. 1.12[83] which implemented Prodigal v. 2.6.3 for gene

prediction[84]. Ribosomal RNA genes were predicted in Infernal v. 1.1.2[85] against Rfam v. 14.2[86]. Gene

functions were annotated in KofamScan using default settings and a bitscore-to-threshold ratio of 0.7[87].

Transporters were predicted in BLASTp against the Transporter Classification Database downloaded on

January 27, 2021[88] using a maximum e-value of 10^{-20}. CAZymes were predicted in hmmsearch from

HMMER v. 3.1b2[89] against dbCAN2 HMMdb v. 9[90]. GH family 109 was omitted from CAZyme analysis

due to nonspecificity of the hidden Markov model.

Functions were assigned based on the following set of rules. Oxygenic photosynthesis required

evidence of at least one marker gene from both photosystems I (psaA-F: K02689, K02690, K02691,

K02692, K02693, or K02694) and II (psbA-F: K02703, K02704, K02705, K02706, K02707, or K02708).

For anoxygenic photosynthesis, evidence of photosystem II required the presence of both pufL (K08928)

and pufM (K08929); evidence of photosystem I required the presence of all four proteins (pscA-D:

K08940, K08941, K08942, K08943). Bacteriorhodopsin assignment required evidence of bop (K04641).

Autotrophic carbon fixation via three independent pathways was evidenced by rbcL (K01601) and prkB

(K00855) for the CBB cycle, mct (K14470) for the glyoxylate assimilation pathway, and abfD (K14534) for

the 3HP/4HB cycle. Evidence of glycan import mediated by the susCD protein complex was determined

by the presence of both susC (K21573) and susD (K21572). Aromatic compound degradation was

determined by the presence of either a funneling or ring fission pathway: evidence of catabolic funneling

was identified by ferulate to vanillin conversion (evidenced by fcs (K12508)), vanillin to protocatechuate

conversion (evidenced by vanA (K03862)), or protocatechuate ring opening (evidenced by pcaG

(K00448) or ligA (K04100)); ring fission was identified by gallate ring opening (evidenced by galA

(K04099)), catechol ring opening (evidenced by K00446 or K03381), salicylate to gentisate conversion

(evidenced by K00480 or K18242 or K18243), or gentisate ring opening (evidenced by K00450).

Methanol oxidation was evidenced by xoxF (K23995). Nitrogen fixation was evidenced by nifD (K02586)

or nifH (K02588) or nifK (K02591). Dissimilatory nitrate reduction was evidenced by narG (K00370) and

narH (K00371) or napA (K02567) and napB (K02568). Evidence of nitrite reduction was determined by

the presence of 1) nirB (K00362) and nirD (K00363), 2) nrfA (K03385) and nrfH (K15876), 3) nirK

(K00368), or 4) nirS (K15864). Nitric oxide reduction was evidenced by norB (K04561) and norC

(K02305). Nitrous oxide reduction was evidenced by nosZ (K00376). Evidence of sulfur oxidation via SOX

was determined by the presence of 1) soxA (K17222) and soxB (K17224) and soxX (K17223) and soxY

(K17226) and soxZ (K17227), or 2) soxC (K17225) and soxD (K22622). Dissimilatory sulfate reduction

was evidenced by 1) aprA (K00394) and aprB (K00395), or 2) dsrA (K11180) and dsrB (K11181).

PULs were delineated following the algorithm described in Terrapon *et al.*[91]. First, adjacent SusC

and SusD genes located on the same DNA strand (termed tandem susCD pairs) were identified. Tandem

susCD pairs within five loci of each other were considered a single susCD-containing locus. Next,

operons structured around tandem susCD pairs were delineated by including genes within intergenic

distances of ≤200 bp. PUL boundaries were iteratively extended to neighbouring operons when a

CAZyme was detected within the first five genes. The selected intergenic distance threshold exceeded

the empirically-derived 102 bp cut-off proposed by Terrapon *et al.*[91] so as to include functionally-

associated genes other than CAZymes that would otherwise be excluded. Only CAZymes implicated in

glycan degradation (glycoside hydrolases, polysaccharide lyases, carbohydrate-binding modules, and

carbohydrate esterases) were included in conditional PUL extension. Finally, only PULs containing

specified CAZyme types were retained.

MAG read recruitment and coverage normalization

To avoid recruitment to conserved regions, rRNA and tRNA gene sequences were masked in BEDTools

v. 2.26.0[92] prior to mapping. To determine the coverage of MAGs across all lake metagenomes, trimmed

metagenome reads were initially recruited to MAGs at a sequence identity threshold of 95% in BBMap v.

37.76[93], then filtered with a hard cut-off of 96% identity using a custom Python script. The selection of a

95% alignment similarity cut-off reflected a prokaryotic genome ANI species delineation[94]. Mapping files

were converted from SAM to sorted BAM format in Samtools[95]. Horizontal coverage was calculated

following the method of Rodriguez-R *et al.*[21] as TAD_{80}, the average sequencing depth truncated to the

central 80% of mapped positions normalized by the number of genome equivalents. Genome equivalents

in each metagenome were estimated in MicrobeCensus v. 1.1.0[96]. A positive TAD_{80} value signified that

≥10% of MAG positions were mapped by metagenome reads; the ≥10% sequencing breadth threshold

has been shown to detect species presence at a confidence of >95%[97].

Taxonomic classification and phylogenetic analysis

MAG taxonomic classification was performed in GTDB-Tk v. 1.3.0 by phylogenetic placement into

bacterial reference tree v. r95[98]. A *de novo* phylogenetic tree was constructed in GTDB-Tk using the

alignment of 120 marker genes based on the JTT model of evolution and subsequently plotted with the R

package ggtree[99].

Metagenome analysis

Single metagenome assemblies were generated in MEGAHIT[78] using k-mer lengths 23, 43, 63, 83, 103,

123 or metaSPAdes[100] v. 3.13.0 using k-mer lengths 21, 33, 55, 77, 99, 127 and mapped in BWA v.

0.7.17 using the BWTSW index algorithm and BWA-Maximal Exact Matches[79]. Assemblies and coverage

files were submitted to the DOE-JGI Metagenome Annotation Pipeline v. 4.16.6, 5.0.20, or 5.0.23[101].

Metagenome functional diversity was quantified by relative gene counts, calculated as KO counts

normalized by total contig average depth of coverage.

16S rRNA gene amplicon sequencing and analysis

A ~300 bp fragment of the V4 region of the 16S rRNA gene was amplified using the primer set 515F (5'-

GTGCCAGCMGCCGCGGTAA-3') and 806R (5'-GGACTACHVGGGTWTCTAAT-3')[102]. Each PCR

mixture consisted of 5 µL Phusion High Fidelity Buffer, 0.5 µL dNTPs (10 mM), 1.8 µL of each primer (5

µM), 0.25 µL Phusion polymerase, 13.65 µL H_2O, and 10 ng in a 2 µL volume of template DNA. PCR

conditions were 30 s of 98 °C, followed by 22 cycles of 98 °C for 20 s, 54 °C for 35 s, 72 °C for 30 s, and

a final elongation at 72 °C for 60 s. Sequencing of 250 bp paired-end reads was performed on an Illumina

MiSeq platform.

Primer sequences were trimmed in Cutadapt v. 3.1[103]. ASVs were inferred in DADA2 v. 1.16[104],

which performed trimming of low-quality 3'-end positions from forward and reverse reads, quality retention

of reads 160 – 200 bp in length, ASV inference within individual samples informed by machine-learned

sequencing error rates, merging of paired reads, and removal of consensus chimeric ASVs. ASVs <250

or >256 bp in length were removed along with ASVs not assigned at the kingdom level and ASVs

assigned to archaea, eukaryotes, or chloroplasts. ASV data from two lakes were discarded because they

did not meet the threshold 10,000 sequences. ASVs were classified against GTDB r95 16S rRNA gene

sequences formatted for DADA2[105].

Ecological analysis

PCA of lake features was performed in the R package vegan v. 2.5.6[106] on scaled and centred variables;

latitude was passively plotted. Biogeographical analysis was conducted on MAG assemblages in 300

freshwater and oligosaline lakes, identified as having conductivity <8 mS/cm and total major ions <4,000

mg/L. To build MAG functional assemblage matrices, we assumed that a function was implied when an

encoding MAG was mapped to a lake metagenome. PCoA was performed in the R package ape v. 5.4[107]

on pairwise Jaccard dissimilarities between MAG or ASV taxonomic assemblage presence-absence data

or Bray-Curtis dissimilarities between metagenome gene count data normalized by total depth of

coverage. GDMs fitting assemblage turnover to environmental gradients were constructed in the R

package gdm v. 1.4.2 using 100 permutations during each step of backward variable elimination[108]. PCA

of MAG GH composition was performed on Hellinger-transformed data. Data analysis and visualization

were performed in R v. 4.0.1[109] using the tidyverse v. 1.3.0 suite of packages[110].

Code availability

535 Scripts associated with this study are available at https://github.com/rebeccagarner/lakepulse_mags.

Data availability

Raw metagenome reads were archived in the European Nucleotide Archive under study accession

PRJEB29238 (https://www.ebi.ac.uk/ena/browser/view/PRJEB29238). Metagenome co-assemblies were

deposited and annotated at the Joint Genome Institute (JGI) Genomes OnLine Database (GOLD) under

study accession Gs0136026 and analysis projects Ga0495746 (Boreal/Taiga Cordilleras), Ga0495744

(Montane Cordillera), Ga0495745 (Pacific Maritime), Ga0495743 (Taiga Plains), Ga0485099 (Semi-Arid

Plateaux), Ga0485102 (Boreal Plains), Ga0485100 (Prairies), Ga0364548 (Mixedwood Plains),

Ga0373103 (Boreal Shield), Ga0372599 (Atlantic Highlands), and Ga0372598 (Atlantic Maritime). MAGs

from co-assemblies and annotations were deposited in Dryad (https://doi.org/10.5061/dryad.zkh1893fs).

Acknowledgements

This study was funded by the NSERC Canadian LakePulse Network (Strategic Network Grant NETGP-

479720) and Canada Research Chairs held by DAW and YH. REG and VEO received scholarships from

the NSERC CREATE ÉcoLac Training Program in Lake and Fluvial Ecology. REG also received support

from a *Fonds de recherche du Québec – Nature et technologies* Doctoral Research Scholarship and the

Stephen Bronfman Graduate Scholarship in Environmental Studies. The pan-Canadian field campaigns

were made possible through the immense commitment and efforts of the LakePulse sampling crews and

support teams, and the cooperation and assistance of Indigenous groups, municipal and park employees,

lake associations, and landowners. We thank the LakePulse researchers involved in the generation,

management, and quality control of the dataset, and acknowledge contributions from Jelena Juric, Paul

MacKeigan, Cindy Paquette, Marieke Beaulieu, Katherine Griffiths, Geneviève Potvin, Bruno Cremella,

Gabriel Diab, Vincent Fugère, Jihyeon Kim, Anaïs Oliva, and Katherine Velghe. We thank Wentworth

Brookes, Shawn Simpson, and Compute Canada for bioinformatics technical support.

References

1.   Newton, R. J., Jones, S. E., Eiler, A., McMahon, K. D. & Bertilsson, S. A Guide to the Natural

History of Freshwater Lake Bacteria. *Microbiol. Mol. Biol. Rev.* **75**, 14–49 (2011).

2.   Pernthaler, J. Competition and niche separation of pelagic bacteria in freshwater habitats. *Environ.*

Microbiol. **19**, 2133–2150 (2017).

3.   Bertilsson, S. & Mehrshad, M. Diversity and Dynamics of Bacterial Communities in Freshwater

Lakes. in *Encyclopedia of Inland Waters* 601–615 (Elsevier, 2022).

4.   Kratz, T. K., MacIntyre, S. & Webster, K. E. Causes and Consequences of Spatial Heterogeneity

in Lakes. in *Ecosystem Function in Heterogeneous Landscapes* (eds. Lovett, G. M., Jones, C.,

Turner, M. G. & Weathers, K. C.) 329–347 (Springer New York, 2005).

5.   Tranvik, L. J. *et al*. Lakes and reservoirs as regulators of carbon cycling and climate. *Limnol.*

Oceanogr. **54**, 2298–2314 (2009).

6.   Williamson, C. E., Dodds, W., Kratz, T. K. & Palmer, M. A. Lakes and streams as sentinels of

environmental change in terrestrial and atmospheric processes. *Front. Ecol. Environ*. **6**, 247–254

(2008).

7.   Grossart, H., Massana, R., McMahon, K. D. & Walsh, D. A. Linking metagenomics to aquatic

microbial ecology and biogeochemical cycles. *Limnol. Oceanogr.* **65**, 1–19 (2020).

8.   O'Reilly, C. M. *et al.* Rapid and highly variable warming of lake surface waters around the globe.

Geophys. Res. Lett. **42**, 10773–10781 (2015).

9.   Smith, V. H. & Schindler, D. W. Eutrophication science: where do we go from here? *Trends Ecol.*

Evol. **24**, 201–207 (2009).

10. Jane, S. F. *et al.* Widespread deoxygenation of temperate lakes. *Nature* **594**, 66–70 (2021).

11. Dugan, H. A. *et al.* Lakes at Risk of Chloride Contamination. *Environ. Sci. Technol.* **54**, 6639–6650

(2020).

12. Reid, A. J. *et al.* Emerging threats and persistent conservation challenges for freshwater

biodiversity. *Biol. Rev.* **94**, 849–873 (2019).

13. Sheridan, E. A. *et al*. Plastic pollution fosters more microbial growth in lakes than natural organic

matter. *Nat. Commun*. **13**, 4175 (2022).

14. Tyson, G. W. *et al*. Community structure and metabolism through reconstruction of microbial

genomes from the environment. *Nature* **428**, 37–43 (2004).

15. Hug, L. A. *et al*. A new view of the tree of life. *Nat. Microbiol.* **1**, 16048 (2016).

16. Cabello-Yeves, P. J. *et al.* Genomes of Novel Microbial Lineages Assembled from the Sub-Ice
Waters of Lake Baikal. *Appl. Environ. Microbiol.* **84**, 1–21 (2018).

17. Okazaki, Y., Nakano, S., Toyoda, A. & Tamaki, H. Long-Read-Resolved, Ecosystem-Wide
Exploration of Nucleotide and Structural Microdiversity of Lake Bacterioplankton Genomes.
mSystems **7**, e00433-22. (2022).

18. Mehrshad, M. *et al.* Hidden in plain sight—highly abundant and diverse planktonic freshwater
Chloroflexi. *Microbiome* **6**, 176 (2018).

19. Arora-Williams, K. *et al.* Dynamics of microbial populations mediating biogeochemical cycling in a
freshwater lake. *Microbiome* **6**, 165 (2018).

20. Tran, P. Q. *et al.* Depth-discrete metagenomics reveals the roles of microbes in biogeochemical
cycling in the tropical freshwater Lake Tanganyika. *ISME J.* **15**, 1971–1986 (2021).

21. Rodriguez-R, L. M., Tsementzi, D., Luo, C. & Konstantinidis, K. T. Iterative subtractive binning of
freshwater chronoseries metagenomes identifies over 400 novel species and their ecologic
preferences. *Environ. Microbiol.* **22**, 3394–3412 (2020).

22. Kavagutti, V. S. *et al.* High-resolution metagenomic reconstruction of the freshwater spring bloom.
Microbiome **11**, 15 (2023).

23. Tully, B. J., Graham, E. D. & Heidelberg, J. F. The reconstruction of 2,631 draft metagenome-
assembled genomes from the global oceans. *Sci. Data* **5**, 170203 (2018).

24. Anantharaman, K. *et al.* Thousands of microbial genomes shed light on interconnected
biogeochemical processes in an aquifer system. *Nat. Commun.* **7**, 13219 (2016).

25. Almeida, A. *et al.* A new genomic blueprint of the human gut microbiota. *Nature* **568**, 499–504
(2019).

26. Nayfach, S., Shi, Z. J., Seshadri, R., Pollard, K. S. & Kyrpides, N. C. New insights from
uncultivated genomes of the global human gut microbiome. *Nature* **568**, 505–510 (2019).

27. Pasolli, E. *et al.* Extensive Unexplored Human Microbiome Diversity Revealed by Over 150,000
Genomes from Metagenomes Spanning Age, Geography, and Lifestyle. *Cell* **176**, 649-662.e20
(2019).

28. Nayfach, S. *et al.* A genomic catalog of Earth's microbiomes. *Nat. Biotechnol.* **39**, 499–509 (2021).

29. Buck, M. *et al.* Comprehensive dataset of shotgun metagenomes from oxygen stratified freshwater
lakes and ponds. *Sci. Data* **8**, 131 (2021).

30. Messager, M. L., Lehner, B., Grill, G., Nedeva, I. & Schmitt, O. Estimating the volume and age of
water stored in global lakes using a geo-statistical approach. *Nat. Commun.* **7**, 13603 (2016).

31. Huot, Y. *et al.* The NSERC Canadian Lake Pulse Network: A national assessment of lake health
providing science for water management in a changing climate. *Sci. Total Environ.* **695**, 133668
(2019).

32. Kraemer, S. A. *et al.* A large-scale assessment of lakes reveals a pervasive signal of land use on
bacterial communities. *ISME J.* **14**, 3011–3023 (2020).

33. Wight, J., Varin, M.-P., Robertson, G. J., Huot, Y. & Lang, A. S. Microbiology in the Field:
Construction and Validation of a Portable Incubator for Real-Time Quantification of Coliforms and
Other Bacteria. *Front. Public Heal*. **8**, 1–11 (2020).

34. Garner, R. E., Gregory-Eaves, I. & Walsh, D. A. Sediment Metagenomes as Time Capsules of
Lake Microbiomes. *mSphere* **5**, e00512-20 (2020).

35. Garner, R. E. *et al.* Protist Diversity and Metabolic Strategy in Freshwater Lakes Are Shaped by
Trophic State and Watershed Land Use on a Continental Scale. *mSystems* **7**, (2022).

36. MacKeigan, P. W. *et al.* Comparing microscopy and DNA metabarcoding techniques for identifying
cyanobacteria assemblages across hundreds of lakes. *Harmful Algae* **113**, 102187 (2022).

37. Oliva, A., Garner, R. E., Walsh, D. & Huot, Y. The occurrence of potentially pathogenic fungi and
protists in Canadian lakes predicted using geomatics, in situ and satellite-derived variables:
Towards a tele-epidemiological approach. *Water Res.* **209**, 117935 (2022).

38. Griffiths, K. *et al.* Pervasive changes in algal indicators since pre-industrial times: A
paleolimnological study of changes in primary production and diatom assemblages from ~200
Canadian lakes. *Sci. Total Environ.* **838**, 155938 (2022).

39. Kraemer, S. A., Barbosa da Costa, N., Oliva, A., Huot, Y. & Walsh, D. A. A resistome survey
across hundreds of freshwater bacterial communities reveals the impacts of veterinary and human
antibiotics use. *Front. Microbiol*. **13**, 1–13 (2022).

40. Oliva, A. *et al*. Geospatial analysis reveals a hotspot of fecal bacteria in Canadian prairie lakes
linked to agricultural non-point sources. *Water Res*. **231**, 119596 (2023).

41. Sánchez Schacht, J. R., MacKeigan, P. W., Taranu, Z. E., Huot, Y. & Gregory-Eaves, I.
Agricultural and urban land use explain substantial variation in lake water quality across national-
scale physiographic and climatic gradients. Preprint at https://doi.org/10.1101/2022.08.29.505280
(2022).

42. Albertsen, M. *et al*. Genome sequences of rare, uncultured bacteria obtained by differential
coverage binning of multiple metagenomes. *Nat. Biotechnol*. **31**, 533–538 (2013).

43. Bowers, R. M. *et al.* Minimum information about a single amplified genome (MISAG) and a
metagenome-assembled genome (MIMAG) of bacteria and archaea. *Nat. Biotechnol.* **35**, 725–731
(2017).

44. Fernández-Gómez, B. *et al.* Ecology of marine Bacteroidetes: a comparative genomics approach.
ISME J. **7**, 1026–1037 (2013).

45. De Corte, D. *et al.* Metagenomic insights into zooplankton-associated bacterial communities.
Environ. Microbiol. **20**, 492–505 (2018).

46. Ezzedine, J. A., Desdevises, Y. & Jacquet, S. Bdellovibrio and like organisms: current
understanding and knowledge gaps of the smallest cellular hunters of the microbial world. *Crit.*

667   *Rev. Microbiol.* **48**, 428–449 (2022).

668 47. Brown, C. T. *et al.* Unusual biology across a group comprising more than 15% of domain Bacteria.

669   *Nature* **523**, 208–211 (2015).

670 48. Berg, I. A. Ecological Aspects of the Distribution of Different Autotrophic CO_2 Fixation Pathways.

671   *Appl. Environ. Microbiol.* **77**, 1925–1936 (2011).

672 49. He, C. *et al.* Genome-resolved metagenomics reveals site-specific diversity of episymbiotic CPR

673   bacteria and DPANN archaea in groundwater ecosystems. *Nat. Microbiol.* **6**, 354–365 (2021).

674 50. Tian, R. *et al.* Small and mighty: adaptation of superphylum Patescibacteria to groundwater

675   environment drives their genome simplicity. *Microbiome* **8**, 51 (2020).

676 51. Ferrier, S., Manion, G., Elith, J. & Richardson, K. Using generalized dissimilarity modelling to

677   analyse and predict patterns of beta diversity in regional biodiversity assessment. *Divers. Distrib.*

678   **13**, 252–264 (2007).

679 52. Sunagawa, S. *et al.* Structure and function of the global ocean microbiome. *Science* **348**,

680   1261359–1261359 (2015).

681 53. Delgado-Baquerizo, M. *et al.* A global atlas of the dominant bacteria found in soil. *Science* **359**,

682   320–325 (2018).

683 54. Hahn, M. W., Huemer, A., Pitt, A. & Hoetzinger, M. Opening a next-generation black box:

684   Ecological trends for hundreds of species-like taxa uncovered within a single bacterial >99% 16S

685   rRNA operational taxonomic unit. *Mol. Ecol. Resour.* **21**, 2471–2485 (2021).

686 55. Azan, S. S. E. & Arnott, S. E. The impact of calcium decline on population growth rates of

687   crustacean zooplankton in Canadian Shield lakes. *Limnol. Oceanogr.* **63**, 602–616 (2018).

688 56. Weyhenmeyer, G. A. *et al.* Widespread diminishing anthropogenic effects on calcium in

689   freshwaters. *Sci. Rep.* **9**, 10450 (2019).

690 57. Kaushal, S. S. *et al.* Freshwater salinization syndrome on a continental scale. *Proc. Natl. Acad.*

691   *Sci.* **115**, E574–E583 (2018).

692 58. McKee, L. S. *et al.* Polysaccharide degradation by the Bacteroidetes: mechanisms and

693   nomenclature. *Environ. Microbiol. Rep.* **13**, 559–581 (2021).

694 59. He, S. *et al.* Ecophysiology of Freshwater Verrucomicrobia Inferred from Metagenome-Assembled

695   Genomes. *mSphere* **2**, e00277-17 (2017).

696 60. Harding, C. J. *et al.* A lysozyme with altered substrate specificity facilitates prey cell exit by the

697   periplasmic predator Bdellovibrio bacteriovorus. *Nat. Commun.* **11**, 4817 (2020).

698 61. Berlemont, R. & Martiny, A. C. Glycoside Hydrolases across Environmental Microbial

699   Communities. *PLOS Comput. Biol.* **12**, e1005300 (2016).

700 62. Garron, M.-L. & Henrissat, B. The continuing expansion of CAZymes and their families. *Curr.*

701   *Opin. Chem. Biol.* **53**, 82–87 (2019).

702 63. Delgado, A. & Gómez, J. A. The Soil. Physical, Chemical and Biological Properties. in *Principles of*

703   *Agronomy for Sustainable Agriculture* (eds. Villalobos, F. J. & Fereres, E.) 15–26 (Springer

International Publishing, 2016).

64. Marmen, S. *et al*. The Role of Land Use Types and Water Chemical Properties in Structuring the
Microbiomes of a Connected Lake System. *Front. Microbiol*. **11**, 89 (2020).

65. Sperlea, T. *et al*. The relationship between land cover and microbial community composition in
European lakes. *Sci. Total Environ*. **825**, 153732 (2022).

66. Phale, P. S., Sharma, A. & Gautam, K. Microbial degradation of xenobiotics like aromatic
pollutants from the terrestrial environments. in *Pharmaceuticals and Personal Care Products:*
Waste Management and Treatment Technology 259–278 (Elsevier, 2019).

67. Foley, J. A. *et al*. Global Consequences of Land Use. *Science* **309**, 570–574 (2005).

68. Kaushal, S. S. *et al*. Land Use and Climate Variability Amplify Carbon, Nutrient, and Contaminant
Pulses: A Review with Management Implications. *JAWRA J. Am. Water Resour. Assoc.* **50**, 585–
614 (2014).

69. Salcher, M. M., Schaefle, D., Kaspar, M., Neuenschwander, S. M. & Ghai, R. Evolution in action:
habitat transition from sediment to the pelagial leads to genome streamlining in Methylophilaceae.
ISME J. **13**, 2764–2777 (2019).

70. Biessy, L., Pearman, J. K., Waters, S., Vandergoes, M. J. & Wood, S. A. Metagenomic insights to
the functional potential of sediment microbial communities in freshwater lakes. *Metabarcoding and*
Metagenomics **6**, 59–74 (2022).

71. Linz, A. M. *et al*. Freshwater carbon and nutrient cycles revealed through reconstructed population
genomes. *PeerJ* **6**, e6075 (2018).

72. Neuenschwander, S. M., Ghai, R., Pernthaler, J. & Salcher, M. M. Microdiversification in genome-
streamlined ubiquitous freshwater Actinobacteria. *ISME J*. **12**, 185–198 (2018).

73. NSERC Canadian Lake Pulse Network. NSERC Canadian Lake Pulse Network field manual 2017
– 2018 - 2019 surveys (eds. Varin, M.-P., Beaulieu, M. & Huot, Y.).
https://doi.org/10.17118/11143/18662 (Université de Sherbrooke, 2021).

74. Hengl, T. *et al*. SoilGrids250m: Global gridded soil information based on machine learning. *PLoS*
One **12**, e0169748 (2017).

75. Muñoz Sabater, J. ERA5-Land hourly data from 1981 to present. *Copernicus Climate Change*
Service (C3S) Climate Data Store (CDS) https://doi.org/10.24381/cds.e2161bac (2019).

76. Canadian Council of Ministers of the Environment. *Canadian water quality guidelines for the*
protection of aquatic life: Phosphorus: Canadian Guidance Framework for the Management of
Freshwater Systems. Canadian environmental quality guidelines (2004).

77. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina sequence data.
Bioinformatics **30**, 2114–2120 (2014).

78. Li, D. *et al*. MEGAHIT v1.0: A fast and scalable metagenome assembler driven by advanced
methodologies and community practices. *Methods* **102**, 3–11 (2016).

79. Li, H. & Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics **25**, 1754–1760 (2009).

80.   Kang, D. D. *et al.* MetaBAT 2: an adaptive binning algorithm for robust and efficient genome
reconstruction from metagenome assemblies. *PeerJ* **7**, e7359 (2019).

81.   Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM: assessing
the quality of microbial genomes recovered from isolates, single cells, and metagenomes.
Genome Res. **25**, 1043–1055 (2015).

82.   Olm, M. R., Brown, C. T., Brooks, B. & Banfield, J. F. dRep: a tool for fast and accurate genomic
comparisons that enables improved genome recovery from metagenomes through de-replication.
ISME J. **11**, 2864–2868 (2017).

83.   Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068–2069 (2014).

84.   Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site identification.
BMC Bioinformatics **11**, 119 (2010).

85.   Nawrocki, E. P. & Eddy, S. R. Infernal 1.1: 100-fold faster RNA homology searches. *Bioinformatics*
29, 2933–2935 (2013).

86.   Kalvari, I. *et al.* Rfam 14: expanded coverage of metagenomic, viral and microRNA families.
Nucleic Acids Res. **49**, D192–D200 (2021).

87.   Aramaki, T. *et al.* KofamKOALA: KEGG Ortholog assignment based on profile HMM and adaptive
score threshold. *Bioinformatics* **36**, 2251–2252 (2020).

88.   Saier, M. H. et al. The Transporter Classification Database (TCDB): recent advances. *Nucleic*
Acids Res. **44**, D372–D379 (2016).

89.   Finn, R. D., Clements, J. & Eddy, S. R. HMMER web server: interactive sequence similarity
searching. *Nucleic Acids Res.* **39**, W29–W37 (2011).

90.   Zhang, H. *et al.* dbCAN2: a meta server for automated carbohydrate-active enzyme annotation.
Nucleic Acids Res. **46**, W95–W101 (2018).

91.   Terrapon, N., Lombard, V., Gilbert, H. J. & Henrissat, B. Automatic prediction of polysaccharide
utilization loci in Bacteroidetes species. *Bioinformatics* **31**, 647–655 (2015).

92.   Quinlan, A. R. & Hall, I. M. BEDTools: a flexible suite of utilities for comparing genomic features.
Bioinformatics **26**, 841–842 (2010).

93.   Bushnell, B. BBMap. *https://sourceforge.net/projects/bbmap/* (2015).

94.   Jain, C., Rodriguez-R, L. M., Phillippy, A. M., Konstantinidis, K. T. & Aluru, S. High throughput ANI
analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat. Commun.* **9**, 5114
(2018).

95.   Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079
(2009).

96.   Nayfach, S. & Pollard, K. S. Average genome size estimation improves comparative
metagenomics and sheds light on the functional ecology of the human microbiome. *Genome Biol.*
16, 51 (2015).

97.   Castro, J. C. *et al*. imGLAD: accurate detection and quantification of target organisms in
779   metagenomes. *PeerJ* **6**, e5882 (2018).
780   98.   Chaumeil, P.-A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk: a toolkit to classify
781   genomes with the Genome Taxonomy Database. *Bioinformatics* **36**, 1925–1927 (2019).
782   99.   Yu, G., Smith, D. K., Zhu, H., Guan, Y. & Lam, T. T. ggtree: an R package for visualization and
783   annotation of phylogenetic trees with their covariates and other associated data. *Methods Ecol.*
784   *Evol.* **8**, 28–36 (2017).
785   100.   Nurk, S., Meleshko, D., Korobeynikov, A. & Pevzner, P. A. metaSPAdes: a new versatile
786   metagenomic assembler. *Genome Res*. **27**, 824–834 (2017).
787   101.   Huntemann, M. *et al*. The standard operating procedure of the DOE-JGI Metagenome Annotation
788   Pipeline (MAP v.4). *Stand. Genomic Sci*. **11**, 17 (2016).
789   102.   Caporaso, J. G. *et al*. Global patterns of 16S rRNA diversity at a depth of millions of sequences
790   per sample. *Proc. Natl. Acad. Sci*. **108**, 4516–4522 (2011).
791   103.   Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
792   *EMBnet.journal* **17**, 10–12 (2011).
793   104.   Callahan, B. J. *et al*. DADA2: High-resolution sample inference from Illumina amplicon data. *Nat.*
794   *Methods* **13**, 581–583 (2016).
795   105.   Alishum, A. DADA2 formatted 16S rRNA gene sequences for both bacteria & archaea. (2020)
796   https://doi.org/10.5281/zenodo.3951383.
797   106.   Oksanen, J. *et al.* vegan: an R package for community ecologists. (2020).
798   107.   Paradis, E. & Schliep, K. ape 5.0: an environment for modern phylogenetics and evolutionary
799   analyses in R. *Bioinformatics* **35**, 526–528 (2019).
800   108.   Fitzpatrick, M. C. *et al.* gdm: Generalized dissimilarity modeling. (2021).
801   109.   R Core Team. R: a language and environment for statistical computing. *R Foundation for*
802   *Statistical Computing, Vienna, Austria* (2021).
110.   Wickham, H. *et al.* Welcome to the Tidyverse. *J. Open Source Softw.* **4**, 1686 (2019).

Figure legends

Figure 1. The LakePulse MAG catalogue: geographical and environmental scope, quality, and patterns of
local-scale diversity. **(A)** Map of 308 sampled lakes from across a broad trophic state gradient (shown as
point colours) in 12 Canadian ecozones (shown as base map colours). **(B)** Principal component analysis
illustrating the variation in environmental features among sampled lakes. The letter symbols (see panel A
for legend) and colours of each point represent ecozones and geographic regions of Canada,
respectively. Arrows show the contributions of environmental variables to PC loadings, except for latitude
which was passively plotted. **(C)** Distributions of MAG characteristics for medium- and high-quality MAGs.
(D) MAG richness within lakes. Point colours represent richness, which is the number of MAGs detected

at each site based on positive TAD_{80} values (see panel A for ecozone colour legend). **(E)** MAG

accumulation curve (i.e. the number of new genomes detected) in a random ordering of lakes. **(F)**

Fraction of LakePulse metagenomes captured within MAGs (measured as the percentage of each

metagenome recruited to the MAGs). Point colours represent lake trophic states and letters represent

ecozones (see legend of panel A).

Figure 2. Taxonomic, phylogenetic, and functional diversity of the LakePulse MAG catalogue. **(A)**

Number of MAGs assigned to each phylum and phylogenetic tree of MAGs constructed from 120

bacterial marker genes. **(B)** Taxonomic novelty of the LakePulse MAG catalogue. Bar plots are divided by

colour to show the number of novel (i.e. unclassified) and classified MAGs from class to species level. **(C)**

Functional diversity across taxonomic orders. The lower heat map shows the number of MAGs within

each class containing genetic markers for a selection of metabolic functions (grey = no evidence; point

colours next to class names indicate the phylum-level affiliation of MAGs shown in panel A). The upper

heat map shows the number of lakes containing MAGs with evidence of metabolic functions.

Figure 3. Niche breadth and environmental filtering of MAGs based on taxonomic and functional

biogeography. **(A)** Frequencies of MAG incidence (i.e. the number of lakes occupied by each MAG).

Colours show MAG phylum-level assignments. **(B)** Principal coordinate analysis (PCoA) showing the

variation in MAG taxonomic composition among lakes based on Jaccard dissimilarities. Point colours

represent lake trophic state and letter symbols represent ecozones. **(C)** Relationship between functional

and taxonomic diversity among MAG assemblages in different lakes. The colour gradient shows the

density of pairwise comparisons. **(D)** PCoA showing the variation in functional gene content among MAG

assemblages based on Jaccard dissimilarities. Point colours and letter symbols are described in the

legend of panel B. **(E)** Percent deviance explained by generalized dissimilarity models (GDMs) fitting the

responses of taxonomic or functional MAG assemblages or community profiles to physicochemical,

morphometric, geographic, or climatic gradients. (**F-H**) Partial effects of **(F)** total phosphorus (TP) and

chlorophyll-*a* (Chl-*a*), **(G)** temperature and pH, and **(H)** calcium and lake colour on the taxonomic and

functional turnover of MAG assemblages across lakes. Curve shapes represent the rate of turnover and

the maximum height of curves represents the magnitude of turnover across environmental gradients. Line

colours represent different MAG assemblage response data (red = taxonomic turnover, blue = functional

turnover) and line types (solid or dashed) represent different environmental explanatory variables.

Figure 4. Diversity of specific bacterial functions among lake assemblages and individual MAGs. **(A)**

Percent deviance explained by generalized dissimilarity models (GDMs) fitting the responses of specific

MAG assemblage functions to physicochemical, morphometric, geographic, or climatic gradients. **(B)**

Partial effects of chlorophyll-*a* and lake colour on the carbohydrate and lipid metabolism turnover of MAG

assemblages across lakes. **(C)** PCoA showing the variation in carbohydrate metabolism among MAG

assemblages. **(D)** PCoA showing the variation in lipid metabolism among MAG assemblages. Point

colours in **C** and **D** represent lake trophic state and letter symbols represent ecozones. **(E)** Percentage

and number of glycoside hydrolase (GH) genes distributed across MAG phyla. **(F)** PCA showing the

variation in GH gene families among MAGs. Point colours represent MAG phyla and letter symbols

represent order-level taxonomy within phyla.

Figure 5. Influences of watershed soil and land use on lake bacterial assemblages. **(A)** Percent deviance

explained by generalized dissimilarity models (GDMs) fitting the responses of taxonomic or functional

MAG assemblages or community profiles to watershed soil characteristics. **(B)** Partial effects of soil pH

and volumetric fraction of coarse fragments on the taxonomic and functional turnover of MAG

assemblages across lakes. **(C)** Map of proportion of land used for crop agriculture in lake watersheds. **(D)**

Map of human population densities in lake watersheds. **(E)** Percent deviance explained by generalized

dissimilarity models (GDMs) fitting the responses of taxonomic or functional MAG assemblages or

community profiles to watershed land use characteristics. Partial effects of crop agriculture and human

population density **(F)** on the taxonomic and functional turnover of MAG assemblages across lakes and

(G) on the turnover of xenobiotics biodegradation and metabolism within MAG assemblages across lakes.

Supplementary information

Figure S1. Project workflow illustrating the generation and analysis of the LakePulse MAG catalogue.

Figure S2. Distributions of **(A)** geography, **(B)** lake morphometry, **(C)** watershed soil, **(D)** land use, **(E)**

climate, and **(F)** surface water physicochemistry variables. Colours represent ecozones and ecozone

medians are indicated by dashed lines.

Figure S3. Number of MAGs generated within each ecozone co-assembly.

Figure S4. Comparison of taxonomic diversity in MAG and 16S rRNA gene ASV datasets: percentages of

MAGs and ASVs assigned to **(A)** phyla and **(B)** orders.

Figure S5. Phylum-level biogeographic distributions of MAGs.

Figure S6. Hierarchical clustering (Ward's linkage) of MAG phylum-level taxonomic assemblages. Letter

symbols above each MAG assemblage (scaled to relative TAD_{80}) represent ecozones and coloured

squares represent lake chlorophyll-*a* concentration.

Figure S7. PCoAs of **(A)** the taxonomic variation among ASV assemblages based on Jaccard

dissimilarities and **(B)** the variation in metabolic gene content among metagenomes based on Bray-Curtis

dissimilarities.

Figure S8. Significant predictors of **(A)** taxonomic and functional turnover across MAG assemblages and

community profiles and **(B)** turnover in specific metabolic functions across MAG assemblages. The height

of bars represents the relative importance of predictors within the generalized dissimilarity model. MAG

assemblage turnover is shown above zero and community (ASV or metagenome) turnover is shown

below zero.

Figure S9. Gene maps of polysaccharide utilization loci (PULs) identified across *Bacteroidota* MAGs.

Arrows indicate gene directions. Colours represent gene types (SusCD, CAZymes, tRNA genes, other

KOs).

Figure S10. Principal coordinate analysis (PCoA) showing the variation in xenobiotics biodegradation and

metabolism among MAG assemblages.

Table S1. Summary of metagenome information: lake names, accession information, sampling

coordinates, and assembly characteristics.

Table S2. Summary of MAG accession information, quality, genome characteristics, taxonomy,

associated file names, marker gene content, and TAD_{80} across 300 freshwater to oligosaline lakes.

Table S3. Number of novel MAGs in each phylum.

Table S4. Generalized dissimilarity modelling results for MAG assemblages and community (ASV and

metagenome) profiles across lakes based on taxonomic and functional composition.

A

B
Novel Classified

Class
Order
Family
Genus
Species

No. MAGs

No. MAGs

C

No. lakes
299
3

No. MAGs

Rows (top to bottom):
Bacteroidia, Rhodothermia, Kapabacteria, Chlorobia, Gemmatimonadetes, Verrucomicrobiae, Chlamydiia, Phycisphaerae, Gammaproteobacteria, Alphaproteobacteria, UBA796, Bradimonadia, UBA727, Myxococcia, Polyangia, Binatia, Bacteriovoracia, Acidimicrobiia, Actinomycetia, Thermoleophilia, Anaerolineae, UBA2235, Ellin6529, Eremiobacteria, Cyanobacteriia, Vampirovibrionia, Campylobacteria, Marinamargulisbacteria, Leptospirae

Columns (left to right):
Oxygenic photosynthesis, Anoxygenic photosynthesis (psII), Anoxygenic photosynthesis (psI), Rhodopsin, Calvin cycle, Glyoxylate assimilation, 3HP/4HB cycle, TonB-dependent transport (susCD), Aromatic funneling, Aromatic ring fission, Methanol oxidation, Nitrogen fixation, Dissimilatory nitrate reduction, Nitrite reduction (nirBD), Nitrite reduction (nrf), Nitrite reduction (nirK), Nitrite reduction (nirS), Nitric oxide reduction, Nitrous oxide reduction, Sulfur oxidation (soxABXYZ), Sulfur oxidation (soxCD), Dissimilatory sulfate reduction (apr), Dissimilatory sulfate reduction (dsr)

Panel A x-axis categories:
Bacteroidota, Gammaproteobacteria, Alphaproteobacteria, Actinobacteriota, Verrucomicrobiota, Planctomycetota, Patescibacteria, Bdellovibrionota, Cyanobacteria, Firmicutes, Myxococcota, Chloroflexota, Armatimonadota, Acidobacteriota, Gemmatimonadota, Spirochaetota, Campylobacterota, Dependentiae, Margulisbacteria, UBP7, Desulfobacterota, Eremiobacterota, Fusobacteriota