TITLE: Resolving marine–freshwater transitions by diatoms through a fog of gene tree

discordance and hemiplasy

RUNNING HEAD: Phylogenomics of marine and freshwater diatoms

AUTHORS:

Wade R. Roberts[1], Elizabeth C. Ruck[1], Kala M. Downey[1], Andrew J. Alverson[1,*]

[1]Department of Biological Sciences, University of Arkansas, 1 University of Arkansas,

Fayetteville, AR, 72701, USA

[*]Corresponding author: Andrew J. Alverson, Department of Biological Sciences, University of

Arkansas, 1 University of Arkansas, Fayetteville, AR, 72701, USA; aja@uark.edu

17
18

ABSTRACT

Despite the obstacles facing marine colonists, most lineages of aquatic organisms have colonized

and diversified in freshwaters repeatedly. These transitions can trigger rapid morphological or

physiological change and, on longer timescales, lead to increased rates of speciation. Diatoms are

a lineage of ancestrally marine microalgae that have diversified throughout freshwater habitats

worldwide. We generated a phylogenomic dataset of genomes and transcriptomes for 59 species

to resolve freshwater transitions in one diatom lineage, the Thalassiosirales. Although most parts

of the species tree were consistently resolved with strong support, we had difficulties resolving a

Paleocene radiation, which affected the placement of one freshwater lineage. This and other parts

of the tree were characterized by high levels of gene tree discordance caused by incomplete

lineage sorting and low phylogenetic signal. Despite differences in species trees inferred from

concatenation versus summary methods and codons versus amino acids, traditional methods of

ancestral state reconstruction supported six transitions into freshwaters, two of which led to

subsequent species diversification. However, simulations suggested as few as two independent

transitions when accounting for hemiplasy, transitions occurring on branches in gene trees not

shared with the species tree. This suggested that transitions across the salinity divide were

originally facilitated by alleles already present in the ancestral marine populations. Accounting

for differences in evolutionary outcomes, in which some taxa became locked into freshwaters

while others were able to return to the ocean or become salinity generalists, might help

distinguish between the ancestral changes that opened the door to freshwaters versus the

subsequent, lineage-specific adaptations that allowed them to stay and thrive.

KEYWORDS: Gene concordance, Phylogenomics, Salinity, Site concordance, Thalassiosirales

INTRODUCTION

From bacteria to animals, the salinity gradient separating marine and freshwater

environments poses a significant barrier to the distributions of many organisms (Lozupone and

Knight 2007; McCairns and Bernatchez 2010; Kenny et al. 2019). Identifying how different

lineages cross the salinity divide will improve our understanding of lineage diversification

(Dittami et al. 2017) and the adaptive potential of species to climate change (Dickson et al. 2002;

L E E et al. 2022). Diatoms are a diverse lineage of microalgae that occur throughout marine and

freshwaters, and despite the numerous obstacles facing marine colonists (Kirst 1990, 1996;

Nakov et al. 2020), ancestrally marine diatoms have successfully colonized and diversified in

freshwaters repeatedly throughout their history (Nakov et al. 2019). These patterns are based on

phylogenetic analyses of a small number of molecular markers, however, so they lack the

insights of phylogenomic approaches, which can resolve large-scale macroevolutionary patterns

and, at the same time, uncover key processes at play during important evolutionary transitions.

Although phylogenomic datasets have helped resolve historically recalcitrant nodes

across the tree of life, they have also revealed how discordance in the evolutionary histories of

different genes can confound inferences of species relationships. Gene tree discordance can be

caused by biological sources, such as incomplete lineage sorting (ILS), hybridization, and

compositional heterogeneity (Maddison 1997; Foster 2004; Degnan and Rosenberg 2006), or

methodological sources, such as character sampling and gene tree error (Philippe et al. 2011; Xi

et al. 2015; Molloy and Warnow 2018). Each of these challenge our ability to resolve species

relationships and impact downstream analyses, such as estimation of divergence times (Smith et

al. 2018). Multiple strategies have been proposed to overcome various sources of error, such as

excluding third-codon positions from DNA datasets (Sanderson et al. 2000), using site-

heterogeneous models for amino acid data (Wang et al. 2018), and identifying the conditions

under which concatenation (Edwards 2009) or gene tree summary approaches (Mirarab et al.

2014; Liu et al. 2015) more accurately resolve species relationships.

Discordance between gene and species trees can confound inferences of trait evolution as

well (Hahn and Nakhleh 2016), particularly for complex traits that appear to have evolved

convergently. Focusing on the species tree alone, without considering discordant gene trees, can

lead to artifactual inferences of molecular convergence (Mendes et al. 2016). This failure occurs

when a trait is determined by genes with topologies that do not match the species topology, a

condition known as hemiplasy (Avise and Robinson 2008; Hahn and Nakhleh 2016; Storz 2016).

Hemiplasy has been identified as a likely explanation for patterns of character incongruence in

amino acid substitutions in columnar cacti (Copetti et al. 2017), flower and fruit traits in

Jaltomata (Solanaceae) (Wu et al. 2018), and dietary specialization in *Dysdera* spiders (Vizueta

et al. 2019). High levels of gene tree discordance were detected in all of these cases.

Adaptation to low salinity is a complex trait involving many genes and pathways (Jones

et al. 2012; Artemov et al. 2017; Hughes et al. 2017; Paver et al. 2018), so the genomic changes

associated with successful freshwater colonizations are multifaceted (Cabello-Yeves and

Rodriguez-Valera 2019; Rogers et al. 2021) and generally involve mutations in multiple genes

(DeFaveri et al. 2011; Terekhanova et al. 2019; Chen et al. 2021). To better understand the

pattern, timing, and process of marine–freshwater transitions by diatoms, we assembled a dataset

of 45 genomes and 42 transcriptomes—most of them newly sequenced—to resolve species

relationships, explore the causes and consequences of gene tree discordance, and determine

whether different freshwater transitions were influenced by hemiplasy.

MATERIALS & METHODS

Detailed methods are provided in Supplementary File S1. Briefly, fresh diatom cultures

were isolated from natural plankton or acquired from the National Center for Marine Algae and

Microbiota (NCMA) or Roscoff Culture Collection (RCC). Collection data, culture conditions,

and voucher information are available in Supplementary Table S1. For genome and

transcriptome sequencing, we extracted total DNA and RNA from diatom cultures, constructed

sequencing libraries, and sequenced them on the Illumina platform. Based on a large multigene

phylogeny of diatoms (Nakov et al. 2018), we included *Coscinodiscus,* Lithodesmiales, and

Eunotogramma as outgroups. Accession numbers for reads and assemblies are provided in

Supplementary Table S2.

We used OrthoFinder (Emms and Kelly 2019) to cluster amino acid sequences from all

genomes and transcriptomes into orthogroups, then aligned orthogroups containing ≥20% of the

taxa with MAFFT (Katoh and Standley 2013). For each alignment, we identified the best-fit

substitution model using ModelFinder (Kalyaanamoorthy et al. 2017) and estimated gene trees

with IQ-TREE (Minh et al. 2020b) or FastTree (Price et al. 2010). We then filtered and trimmed

the gene trees using the Rooted Ingroup method to produce final ortholog sets (Yang and Smith

2014). This filtering and trimming procedure was performed twice. We used PAL2NAL

(Suyama et al. 2006) to reconcile nucleotide coding sequence alignments against amino acid

alignments. We used Degen (Regier et al. 2010; Zwick et al. 2012) to replace synonymous sites

in the full coding sequence alignments with degenerate nucleotides. In total, we analyzed

datasets consisting of amino acids (AA), the first and second codon positions (CDS12), and

degenerate codons (DEGEN). We generated final ortholog alignments and inferred trees as

described above, using 1000 ultrafast bootstrap replicates to estimate branch support (Minh et al.

2013). We generated summary statistics for the alignments and gene trees using AMAS

(Borowiec 2016) and PhyKit (Steenwyk et al. 2021) (Supplementary Table S3). Correspondence

analysis of amino acid frequencies across all taxa was performed using GCUA (McInerney

1998).

Species trees were inferred using maximum-likelihood analysis of a concatenated

supermatrix with IQ-TREE and the summary quartet approach implemented in ASTRAL-III

(Zhang et al. 2018). We performed the matched-pairs test of symmetry in IQ-TREE to identify

and remove partitions that violated assumptions of stationarity, reversibility, and homogeneity

(SRH; Naser-Khdour et al. 2019). For the IQ-TREE analysis, we partitioned supermatrices by

gene, used ModelFinder to select the best-fit substitution model for each partition, and estimated

branch support with 10,000 ultrafast bootstrap replicates. For ASTRAL, we used the ortholog

trees as input, collapsing branches with low bootstrap support (<33) to help mitigate gene tree

error (Sayyari and Mirarab 2016; Simmons and Gatesy 2021). We estimated branch support

using local posterior probability (Sayyari and Mirarab 2016). In addition to heterogeneity in gene

histories, substitutional heterogeneity can also make tree inference difficult at deep time scales

(Lartillot and Philippe 2004). To account for this possibility, we also estimated a species tree

from the concatenated amino acid matrix using the Posterior Mean Site Frequency (PMSF)

model implemented in IQ-TREE (Wang et al. 2018).

We calculated the Robinson-Foulds distance between each pair of species trees and

visualized the results with a multidimensional scaling plot made with the R package *treespace*

(Jombart et al. 2017). We characterized discordance using gene and site concordance factors

(Minh et al. 2020a) and quartet concordance factors (Pease et al. 2018). We tested the support for

competing backbone topologies in our species tree using the approximately unbiased (AU) test

(Shimodaira 2002) implemented in IQ-TREE. Relative gene tree support for the same set of

backbone topologies was further evaluated using gene genealogy interrogation method (Arcila et

al. 2017). Finally, we performed the polytomy test on the ASTRAL species tree to test whether

any of the unstable backbone branches were better represented as a polytomy (Sayyari and

Mirarab 2018).

Divergence times were estimated using MCMCtree (Yang 2007; Reis and Yang 2011),

using five fossil-based calibrations (Supplemental File S1), autocorrelated rates, and the

approximate likelihood approach. We performed marginal ancestral state reconstruction for

marine and freshwater habitat across the species tree using hidden state speciation and extinction

(SSE) models in the R package *hisse* (Beaulieu and O'Meara 2016). Lastly, we explored the

probability of hemiplasy in our habitat reconstructions using the models and approaches

implemented in the R package *pepo* and the program HeIST (Guerrero and Hahn 2018; Hibbins

et al. 2020). Due to the computational demands of gene tree simulations, HeIST simulations

were based on a reduced 16-taxon tree.

Assembled genomes and transcriptomes have been deposited at NCBI under BioProject

PRJNA825288. Predicted proteomes, phylogenomic datasets, and scripts have been deposited in

the Dryad Digital Repository: ____________.

RESULTS

We combined 42 newly sequenced draft genomes and 50 newly sequenced

transcriptomes with publicly available genome or transcriptome data for a final dataset of 87 taxa

representing 59 distinct species. A total of 17 transcriptomes were used directly in phylogenomic

analyses, and the other 33 were used for genome annotation. From the combined dataset of

genomes and transcriptomes, we generated alignments and gene trees for 6262 orthologs, with

each taxon represented in an average of 3275 (52.3%) orthologs.

Compositional heterogeneity and dataset construction

167   Relative Composition Variability (Phillips and Penny 2003) indicated greater

compositional heterogeneity in third codon positions compared to amino acids and first and

second codon positions (Supplementary Fig. S1). We also found substantial variation in GC

content of third codon positions across taxa and genes (Supplementary Fig. S2), ranging from an

average proportion of 0.40 ± 0.07 in *Cyclotella kingstonii* to 0.76 ± 0.14 in *Shionodiscus*

oestrupii. To minimize potentially misleading signal in the nucleotide data, we removed third

codon positions and, to examine the effects of saturation and GC heterogeneity in coding

sequences (CDS), we replaced synonymous sites by recoding CDS sequences with degenerate

nucleotides (Zwick et al. 2012). Based on these results, we created three datasets to estimate

phylogenetic relationships: amino acids (AA), first and second codon positions (CDS12), and

degenerate codons (DEGEN).

178   Dataset and alignment characteristics are summarized in Table 1 and detailed in

Supplementary Table S3. Each dataset initially contained 6262 orthologs (Table 1). Gene trees

constructed from all datasets were well supported ($80 \pm 7\%$ average bootstrap; Supplementary

Table S3). To reduce systematic errors due to model misspecification, we removed orthologs that

failed assumptions of stationarity, reversibility, and homogeneity (SRH; $P < 0.05$), which

retained 5522 (AA), 3259 (CDS12), and 3788 (DEGEN) loci in the datasets (hereafter referred to

as the "complete" datasets; Table 1). We additionally subset the complete datasets to maximize

signal or minimize missing data. To maximize signal and reduce stochastic error, we sorted

orthologs by the percentage of parsimony informative (PI) sites and retained the top 25% ranked

orthologs ("top-PI"; Table 1). To minimize the amount of missing data and maximize taxon

occupancy, we sorted complete datasets by the number of taxa and subset these to include the top

25% ranked orthologs with highest taxon occupancy ("top-Taxa"; Table 1). Orthologs in the top-

Taxa datasets contained an average of $76 \pm 8\%$ of the total taxa.

Table 1. Summary of datasets used in the study.

Data type	Dataset name	Num loci	Total sites	Parsimony informative sites (%)	Num loci passing SRH[1]	*Thalassiosira* grade topology	
					$P < 0.05$	IQ-TREE	ASTRAL
Amino acids (AA)	complete	6262	3,777,062	60	5522	Topology 4	Topology 1
	top-PI	1588	996,027	76	1314	Topology 4	Topology 1
	top-Taxa	1574	808,477	64	1374	Topology 5	Topology 1
	top-PI-top-Taxa + PMSF[2] model	488	246,300	77	488	Topology 5	Topology 1
1st + 2nd codon positions (CDS12)	complete	6262	7,554,124	55	3259	Topology 4	Topology 5
	top-PI	1570	1,950,228	71	661	Topology 4	Topology 3
	top-Taxa	1574	1,616,954	59	782	Topology 5	Topology 3
Degenerate codons (DEGEN)	complete	6262	11,311,186	33	3788	Topology 4	Topology 3
	top-PI	1569	2,884,728	44	771	Topology 4	Topology 2
	top-Taxa	1574	2,425,431	36	903	Topology 5	Topology 3

193   [1] Matched-pairs test of Symmetry, Reversibility, and Homogeneity.

194   [2] Posterior Mean Site Frequency model.

Species tree inference and placement of freshwater clades

We initially estimated 18 species trees using amino acids, codon positions 1 and 2, and

degenerate codon sequences with different cutoffs for taxon occupancy or proportion of

parsimony informative sites, using both concatenation and summary quartet approaches (Table

1). Correspondence analysis of amino acid frequencies separated taxa principally by habitat

(marine vs. freshwater) rather than phylogeny (Supplementary Fig. S3), which led us to explore

whether substitutional heterogeneity affected phylogenetic reconstructions. To do so, we

estimated an additional species tree using the PMSF mixture model, which can accommodate

heterogeneity in the amino acid substitution process between sites (Wang et al. 2018). Due to the

computational demands of implementing this model, we applied it only to a reduced amino acid

dataset with orthologs that met both the top-PI and top-Taxa filtering criteria ("AA-top-PI-top-

Taxa"; Table 1). Gene trees from this dataset were also used as input to ASTRAL.

Previous phylogenetic analyses resolved freshwater taxa into two main clades: the genus

Cyclotella, which also includes several marine and brackish species, and the "cyclostephanoids",

comprised of several stenohaline genera confined exclusively in freshwaters (Alverson et al.

2011). Given the potential implications for uncovering the mechanisms of freshwater adaptation,

we were primarily interested in the placements of these two clades. Gross differences among

data types and methods were evident in an ordination of species trees based on pairwise

Robinson-Foulds distances, which showed a clear separation between IQ-TREE and ASTRAL

topologies, with further separation of IQ-TREE trees estimated from datasets that maximized

signal (top-PI) or minimized missing data (top-Taxa) (Supplementary Fig. S4). The phylogenetic

position of *Cyclotella* was strongly supported and robust to differences in data type (codons vs.

amino acids) and analysis (IQ-TREE vs. ASTRAL) (Fig. 1). The cyclostephanoids were placed

consistently within a large clade of marine *Thalassiosira* and relatives (Fig. 1), but the

arrangements of five main subclades—*Thalassiosira* I–IV and the freshwater

cyclostephanoids—varied depending on data type and analysis (Table 1; Fig. 2a). We refer to

this part of the tree as the *Thalassiosira* grade.

223   One resolution of the *Thalassiosira* grade (topology 1), recovered only by ASTRAL

analysis of amino acid gene trees, placed the freshwater cyclostephanoids as sister to a clade of

Thalassiosira I–IV (Table 1; Fig. 2a). All other species trees placed cyclostephanoids as sister to

Thalassiosira III (Table 1; Fig. 2a). ASTRAL analyses of codon-based gene trees (CDS12 and

DEGEN) alone recovered topology 3, which placed cyclostephanoids and *Thalassiosira* III as

sister to the remaining *Thalassiosira* (Table 1; Fig. 2a). Topologies 4 and 5 were recovered by

both data types, but only topology 5 was robust to both data type and analysis, having been

recovered by IQ-TREE analysis of both amino acid and codon alignments, and ASTRAL

analysis of codon-based gene trees (Table 1; Fig. 2a). Moreover, topology 5 was also recovered

by IQ-TREE analysis with the PMSF model, with almost all branches in the *Thalassiosira* grade

receiving maximum support (Fig. 1, branches A–E). Notably, the PMSF analysis also recovered

a monophyletic *Stephanodiscus*, which matches expectations based on morphology (Theriot et

al. 1987). *Stephanodiscus* was paraphyletic, with strong support, in relation to *Cyclostephanos* in

18 of the 20 species trees. Based on all of these results, we chose topology 5 from the PMSF

analysis as the reference species tree (Fig. 1).

Figure 1. Phylogram based on maximum likelihood analysis of amino acids using the posterior

mean site frequency (PMSF) model and a dataset of 488 loci with the highest proportions of taxa

and informative sites ("AA-top-PI-top-Taxa" dataset; Table 1). Backbone nodes of the

Thalassiosira grade are indicated by the letters A–E. All branches had bootstrap support (BS)

244   values of 100 except for those with black circles which had BS = 90. Pie charts on backbone

245   nodes show the proportion of gene trees that support the clade (gCF), the proportion that support

246   both discordant topologies (gDF1, gDF2), and the proportion that are discordant due to

247   polyphyly (gDFP). The size of the pie charts is only for aesthetics.

249   *Discordance underlies topological uncertainty*

250   Discordance among genes and sites is an important factor impacting phylogenetic

reconstruction (Degnan and Rosenberg 2006; Mallet et al. 2016) and multiple tools now exist for

identifying and characterizing discordance (Salichos et al. 2014; Smith et al. 2015; Pease et al.

2018). We characterized discordance by calculating gene, site, and quartet concordance factors

for each branch (Figs. 1 and 2a; Supplementary Figs. S5 and S6). Gene (gCF) and site

concordance factors (sCF) represent the proportion of genes or sites that are in agreement with a

particular branch in the species tree (Minh et al. 2020a). Gene concordance factors range from 0

to 100, and site concordance factors typically range from one-third (33) to 100, with values near

33 indicative of no signal for that branch (Minh et al. 2020a). Quartet concordance factors (QC)

provide an alternative, likelihood-based estimate of the relative support at each branch for the

three possible resolutions of four taxa (Pease et al. 2018). Quartet concordance factors range

from -1 to 1, with positive values showing support for the focal branch, negative values

supportive for an alternate quartet, and values of zero indicating equal support among the three

possible quartets (Pease et al. 2018).

264   All three concordance factors were high for most branches in the tree, indicating that a

majority of genes, sites, and quartets supported those relationships (Supplementary Figs. S5 and

S6). For example, the branch subtending the *Skeletonema* clade had high concordance

(gCF/sCF/QC = 93/75/0.87; Supplementary Figs. S5 and S6). Despite having maximum

bootstrap and local posterior probability support, however, concordance factors were low for

many of the backbone branches (Fig. 1; Supplementary Figs. S5 and S6). These included

backbone branches within the *Thalassiosira* grade (Fig. 1, nodes A–E) that affected placement of

the freshwater cyclostephanoids (Fig. 2). In this part of the tree, concordance was generally low

for the backbone branches (gCF = 4–26; sCF = 32–42; QC = –0.04–0.36) (Figs. 1 and 2;

Supplementary Figs. S5 and S6). Gene concordance factors were lowest for branches C and D,

the only two branches in the species tree with <100% bootstrap support (Fig. 1). Gene

concordance factors were only slightly higher for branches A, B, and E (Fig. 1). These had low

site concordance (Fig. 2a) and near-zero quartet concordance factors (Supplementary Fig. S6),

consistent with expectations for little to no signal. Site concordance factors did not change

appreciably when we used the CDS12 data. However, repeating quartet concordance factor

calculation with the CDS12 data revealed a minor switch in support for branches C ($QC_C = 0.004$

$\rightarrow$ –0.006) and E ($QC_E = -0.04 \rightarrow 0.07$) (Supplementary Fig. S6). These patterns of shifting

support combined with the lowest site concordance factor for branch E ($sCF_E = 32$) suggest that

very few sites support the sister relationship of *Thalassiosira* III and cyclostephanoids, despite its

recovery in 16 of the 20 inferred species trees (Figs. 1 and 2a).

Although bootstrap and posterior probability values were uniformly high across our trees,

traditional measures of branch support often fail to capture the underlying agreement or

disagreement among genes or sites for a given tree topology (Minh et al. 2020a). For many of the

branches on our tree, there were more discordant than concordant genes and sites (Fig. 1;

Supplementary Figs. S5 and S6), suggesting that more genes and sites supported an alternative

relationship. This was the case with *Stephanodiscus*, for example, where more genes supported

paraphyly than monophyly, though more sites supported monophyly (Supplementary Figs. S5

and S6). Across the tree, support for alternative relationships was neither strong nor consistent

among genes or sites. Illustrative of this, discordance in more than one-third of the tree (29 of 83

branches) was due to polyphyly (Supplementary Fig. S5), indicative of a widespread lack of

signal in our gene trees. Taken together, these analyses highlighted extensive gene and site

discordance, some well-supported and much of it not, across Thalassiosirales.

Figure 2. **a)** Phylogenetic hypotheses of the *Thalassiosira* grade inferred using concatenation and

summary methods on the amino acid (AA), codon positions 1 and 2 (CDS12), and recoded

codon (DEGEN) datasets. Nodes are labeled with the percentage of amino acid sites concordant

with the branch (site concordance factor). The principal clade of interest, the freshwater

cyclostephanoids, is colored orange and labeled 'FW'. The four focal clades of marine

Thalassiosira and allies are labeled I–IV. Below, results of gene genealogy interrogation tests of

alternative hypotheses of relationships within the *Thalassiosira* grade. These tests used datasets

filtered to include only the top 25% of orthologs based on the percentage of parsimony

informative sites (top-PI) for **b)** amino acids, **c)** codon positions 1 and 2, and **d)** recoded codons.

Lines correspond to the cumulative number of genes (x-axis) supporting topology hypotheses

with the highest probability and their *P* values (y-axis) from the Approximately Unbiased (AU)

topology tests. Values above the dashed line indicate topological hypotheses that are

significantly better than the alternatives ($P < 0.05$). For example, the green line in (b) shows that

there were a total of 88 genes that best supported topology 3, while only four of those genes were

above the dotted line and were significantly better supported than the other four alternative

topologies.

315   There are both biological (e.g., ILS) and technical causes (e.g., gene tree error) of gene

discordance, but the proportion attributable to each factor can be difficult to discern (Morales-

Briones et al. 2020; Cai et al. 2021). To better assess the importance of gene tree error in our

dataset—whether genes with low phylogenetic signal produced inaccurate gene trees—we

recalculated gene and site concordance using just the 1588 amino acid orthologs with the highest

percentage of parsimony informative sites (top-PI dataset; Table 1). Average gene concordance

increased modestly, from 56.6% to 62.1%, but site concordance was unchanged (Supplemental

Fig. S7). Gene concordance factors for branches A, B, and E in the *Thalassiosira* grade increased

by 5–7% but were largely unchanged for branches C and D (Supplemental Fig. S7). These

increases in gene concordance when using the most signal-rich genes suggest that errors in gene

tree estimation contributed to the lack of resolution in several critical branches (Chan et al. 2020;

Vanderpool et al. 2020). Deeper nodes in the tree may be more prone to technical errors caused

by long-branch attraction, poor alignments, or model misspecification, despite our attempts to

minimize these during dataset construction (Supplementary File S1). We found slight negative

correlations between node age and both gene concordance ($R^2 = 0.10$) and site concordance ($R^2 = $

0.26) (Supplementary Fig. S8), which suggests that older branches more likely suffered from

saturation due to recurrent substitutions.

Placement of the freshwater cyclostephanoids

We used two additional tree-based methods to test the relative support for competing

topologies within the *Thalassiosira* grade. Approximately unbiased (AU) tests on the

concatenated alignment for each dataset in Table 1 were used to assess the relative statistical

support for topologies 1–5 (Fig 2a). Like the original species tree inferences, results of AU tests

largely reflected data type (Table 1), with amino acid characters supporting topology 1 ($P = 0.01$,

AU test) and the codon and degenerate codon datasets supporting topology 4 ($P = 0$, AU test;

Supplementary Table S4). Although concordance factors suggested that most gene trees were

uninformative for relationships in the *Thalassiosira* grade, we looked for secondary signal for

one or more of the five competing topologies using gene genealogy interrogation (GGI; Arcila et

al. 2017). To do this, we performed constrained gene tree searches on the most information-rich

(top-PI) orthologs and compared their likelihoods using the AU test. The GGI test assumes

monophyly of the tested clades, so using our time-calibrated tree, we converted branch lengths

(in millions of years) to coalescent time units using a range of plausible effective population

sizes and generation times for diatoms (Supplementary File S1). The estimated stem branch

lengths for the five clades in the *Thalassiosira* grade were all >5 coalescent units, suggesting

sufficient time to reach monophyly (Rosenberg 2003). After ranking likelihood scores from the

AU tests and selecting constraint topologies with the best score (rank 1 trees), no single topology

was strongly favored in a majority of constrained gene trees across the three datasets, implying

similar levels of support (Fig. 2b–d; Supplementary Table S5). Support from the amino acid

dataset was split between topologies 1 and 4, which were recovered by both summary and

concatenation methods (Table 1; Fig. 2b; Supplementary Table S5). The most frequent best-fit

topology for the codon and degenerate codon datasets corresponded to topology 4 (Fig. 2c,d),

one originally recovered by concatenation only (Table 1).

357   Gene genealogy interrogation can also be used to explore the effects of gene tree error on

summary quartet methods by filtering the input to ASTRAL to include only the highest ranking

constrained genes (Arcila et al. 2017; Mirarab 2017). For each of the three character types, we

performed two ASTRAL analyses using as input either all the top scoring (rank 1) constrained

gene trees ($n_{AA} = 1588$, $n_{CDS12} = 1570$, $n_{DEGEN} = 1569$) or just the subset that had statistical

support ($P < 0.05$) above the AU-based rank 2 topology ($n_{AA} = 142$, $n_{CDS12} = 56$, $n_{DEGEN} = 71$). In

all six cases, the inferred trees were consistent with topology 5, despite it being best supported

(rank 1) in just 13–16% of the constrained gene trees (Fig. 2b–d; Supplementary Table S5). We

originally chose topology 5 as the reference species tree (Fig. 1) because it was recovered by

both amino acids and codons, ASTRAL and IQ-TREE analysis with the PMSF model, and

because it recovered monophyly of *Stephanodiscus*. Coalescent theory predicts that in severe

cases of ILS, short internal branches can produce gene trees that conflict with the species tree

more often than they agree, creating the so-called "anomaly zone" (Degnan and Rosenberg

2006). Our recovery of a species tree topology here that is not the most frequent among the GGI

gene trees could indicate that the backbone of the *Thalassiosira* grade lies in the anomaly zone.

Despite this, polytomy tests in ASTRAL using each dataset rejected the null hypothesis that any

of these branches is a polytomy ($P < 0.05$).

The temporal sequence of marine–freshwater transitions

Divergence time estimates dated the crown Thalassiosirales to the upper Cretaceous,

around 113 Ma (96–120, 95% CI; Fig. 3a; Supplementary Fig. S9). One of the two main

freshwater lineages, *Cyclotella*, originated in the late Cretaceous (79 Ma [66–86, 95% CI]) and

the other freshwater lineage, the cyclostephanoids, originated later in the Eocene (43.5 Ma [36–

48, 95% CI]) (Fig. 3a; Supplementary Fig. S9). Radiation of the *Thalassiosira* grade lineages

occurred during the Paleocene, from 57 Ma [48–63, 95% CI] to 73 Ma [61–80, 95% CI] (Fig. 3a;

Supplementary Fig. S9). The overlapping confidence intervals allow for the possibility that these

lineages diverged in much more rapid succession than suggested by the mean ages.

Hemiplasy impacts marine–freshwater transitions

We estimated the number of marine–freshwater transitions in two ways. A traditional

trait reconstruction method, HiSSE, was used to estimate the maximum number of independent

marine–freshwater transitions under a "homoplasy-only" model. A second model allowed us to

estimate how many of the transitions were non-independent due to hemiplasy, i.e., transitions to

freshwater occurring on branches of discordant gene trees that are not found in the species tree

(Hibbins et al. 2020). The best-fit HiSSE model was a character independent model (CID-4, AIC

= 535.3; Supplementary Table S6), which suggests that shifts in diversification rate occurred

independently of marine–freshwater transitions. Using parameter estimates from the CID-4

model, we inferred six transitions from marine to freshwater habitats (Fig. 2a). With this dataset,

the most recent common ancestor of *Cyclotella* was marine, with the transition to freshwater

occurring shortly thereafter (Fig. 2a). To assess the possible role of hemiplasy in our trait

reconstructions, we calculated hemiplasy risk factors (HRF), which are the ratio of the

probabilities of hemiplasy to homoplasy in different parts of the species tree (Guerrero and Hahn

2018). This analysis revealed an increased chance of character state transitions due to hemiplasy

along most of the backbone branches of the species tree (Fig. 3b). Gene trees simulated from a

reduced 16-taxon species tree using HeIST returned 1133 loci with a simulated site pattern that

matched the marine–freshwater character states (i.e. freshwater taxa shared the same mutation).

With this approach, the most likely scenario is the one supported by the largest number of gene

trees. In our dataset, the scenarios could range from strict homoplasy (n = 6 marine–freshwater

transitions) to strict hemiplasy (n = 1 transition). A total of 732 of the 1133 simulated gene trees

(64.6%) corresponded to a scenario of combined hemiplasy and homoplasy, with two

independent transitions into freshwaters (Supplementary Table S7). Under this model, freshwater

taxa grouped into one of two clades in a simulated gene tree—each clade representing a

freshwater transition—where at least one transition was discordant with the species tree. Just

23% of the simulated trees supported a hemiplasy-only model with a single freshwater transition,

and there was no support for a homoplasy-only history of six independent marine–freshwater

transitions (Supplementary Table S7).

Figure 3. **a)** Divergence times and ancestral state reconstruction of marine and freshwater habitat

in the Thalassiosirales. Conspecific taxa were removed prior to ancestral state reconstruction,

leaving one tip per species. Tip labels are colored according to their habitat (blue=freshwater,

orange=euryhaline, grey=marine). Pie charts denote the probability of each node reconstructed as

either marine (grey) or freshwater (blue) using parameters estimated from the HiSSE CID–4

model. Euryhaline taxa were coded as marine for the purposes of ancestral state reconstruction.

Divergence times for the full set of taxa can be found in Supplementary Fig. S9. **b)** Hemiplasy

Risk Factors (HRF) on internal branches show increased values on most short internal branches,

demonstrating an increased potential for hemiplasy to influence trait reconstruction. Branch

lengths in coalescent units were inferred using ASTRAL.

DISCUSSION

Transitions to freshwaters

Marine–freshwater transitions have been key events in the diversification of lineages

across the tree of life (Logares et al. 2009; e.g., Tedesco et al. 2017), including diatoms where

freshwater taxa have experienced significantly increased rates of both speciation and extinction

compared to their marine ancestors (Nakov et al. 2019). Our study focused on one model clade,

the Thalassiosirales, which is among the most abundant and diverse lineages in the marine and

freshwater plankton and where genetic and genomic resources are readily available (Armbrust et

al. 2004; Nawaly et al. 2020; Roberts et al. 2020). The 42 draft genomes presented here greatly

expand the phylogenetic and ecological diversity of sequenced genomes for Thalassiosirales, and

diatoms as a whole, and will greatly facilitate efforts to identify the genomic basis of freshwater

adaptation in diatoms.

We identified six different marine–freshwater transitions in Thalassiosirales, more than

previous studies (Alverson et al. 2007). We tested whether these transitions were fully

independent, owing to separate mutations (homoplasy) in each of the six freshwater lineages, or

whether some of the transitions were attributable to hemiplasy, i.e., shared ancestral

polymorphisms in discordant gene trees (Avise and Robinson 2008). Hemiplasy simulations

supported a scenario of as few as two independent transitions at the genic level, a result that

followed naturally from the extensive history of gene tree discordance across the backbone of the

tree. Long considered an ecological "Rubicon" that is rarely crossed (Mann 1999), phylogenetic

studies have shown that diatoms move much more frequently across the salinity divide than was

assumed historically (Alverson et al. 2007; Nakov et al. 2019). The discovery of hemiplasy,

which reduced the number of independent freshwater transitions in Thalassiosirales by a factor

of three, suggests that transitions across steep environmental gradients might occur more

commonly than expected when colonists can leverage ancestral polymorphisms rather than

relying exclusively on new mutations.

With hundreds of species, a crown age of roughly 100 Mya, and freshwater transitions

dating as far back as 80 Mya, the breadth and depth of the Thalassiosirales phylogeny far

exceeds other, much younger lineages with traits that have been impacted by hemiplasy (Copetti

et al. 2017; Wu et al. 2018). Relatively deep, and short, backbone branches that predated

freshwater transitions were the ones identified as most susceptible to hemiplasy. The coalescent

branch lengths were short in these parts of the tree (Fig. 3b), and the substitution- (Fig. 1) and

clock-based branch lengths (Fig. 3a) are also shorter than suggested by our trees due to

incomplete taxon sampling (Alverson et al. 2007). Although many of the conditions for

hemiplasy were clear (high levels of gene tree discordance coinciding with a rapid radiation,

particularly along the *Thalassiosira* grade), other properties of the habitat transitions were

indicative of homoplasy: ancestral state reconstructions were unambiguous, the two main

freshwater transitions were not paraphyletic but were aligned with phylogenetic relationships,

and gene concordance was high on the long internal branches subtending the two main

freshwater lineages (Hahn and Nakhleh 2016; Wu et al. 2018).

Given the sheer length of time that has elapsed since the transitions occurred, the shared

ancestral polymorphisms that opened the door to freshwaters for Thalassiosirales will be difficult

to distinguish from the augmenting, lineage-specific adaptive mutations that followed (Zou and

Zhang 2015; Mendes et al. 2016). A large number of genes and pathways have been implicated

in the response to low salinity (Nakov et al. 2020; Downey et al. 2022; Pinseel et al. 2022), so an

adaptive allele in one of the many possible target genes might have made it possible simply to

survive in freshwaters initially. In the tens of millions of years since then, the hemiplasious

alleles could have been overwritten in one or more freshwater lineages, leaving the shared

mechanism itself (e.g., enhanced transport of sodium ions) as the only remaining direct evidence

of the hemiplasy. The divergent evolutionary outcomes might also help distinguish hemiplasy

from homoplasy. The genus *Cyclotella* includes freshwater, secondarily marine, and generalist

euryhaline species that can tolerate a wide range of salinities (Guillard and Ryther 1962; Nakov

et al. 2020; Downey et al. 2022). Similarly, most freshwater transitions at the tips of the tree

(Fig. 1) involved species with populations that also grow in marine habitats (*Conticribra*

weissflogii and *Cyclotella nana*) or can tolerate slightly brackish water (*Thalassiosira lacustris*

and *Skeletonema potamos*). The cyclostephanoids are, by contrast, stenohaline specialists found

exclusively in freshwaters, suggestive of a different genetic trajectory into freshwaters.

Although we have treated salinity as a categorical variable, salinity varies along a

continuum from freshwater to marine and even hypersaline habitats. Moreover, salinity

fluctuations are common in brackish and marine systems, such as coastlines influenced by

precipitation and river discharge. Adaptation to variable environmental conditions may be more

likely linked to modifications of gene expression (Wray 2007). In diatoms, exposure to

freshwater induces profound changes gene expression related to cellular metabolism, ion

transport, photosynthesis, and storage compound biosynthesis (Bussard et al. 2017; Nakov et al.

2020; Downey et al. 2022; Pinseel et al. 2022). Diatoms exhibit high levels of inter- and

intraspecific variation in gene expression in response to reduced salinity (Nakov et al. 2020;

Pinseel et al. 2022), providing numerous targets for natural selection to optimize gene expression

for a particular salinity environment (López-Maury et al. 2008; Bedford and Hartl 2009; Gomez-

Mestre and Jovani 2013). Finally, differences in codon usage (Prabha et al. 2017), nucleotide

substitution rates (Mitterboeck et al. 2016), transposable element activity (Yuan et al. 2018),

epigenetic responses (Artemov et al. 2017), and epistatic interactions (Stern et al. 2022) have

been implicated in adaptation to freshwaters in other groups. The genomic resources and

phylogenetic framework presented here represent an important advance towards identifying the

genes and evolutionary processes underpinning freshwater adaptation by diatoms.

The impact of discordance on placement of freshwater clades

502   Comparative analyses require a strong phylogenetic framework (Felsenstein 1985), so a

major goal of this study was to establish a robust phylogenetic hypothesis for Thalassiosirales.

Our primarily genome-based analysis of 6262 nuclear orthologs provided better resolution and,

superficially, increased support across most of the tree compared to previous analyses based on a

few genes (Alverson et al. 2007). Across character types, methods of inference, and different

criteria for including characters or taxa, the placement of one of the two principal freshwater

clades, *Cyclotella*, was consistent across species trees, with an estimated origin in the late

Cretaceous. The placement of the second major freshwater lineage, the cyclostephanoids, was

less certain.

511   The freshwater cyclostephanoid clade was placed within a grade of marine species, most

of which belong to the polyphyletic genus *Thalassiosira*. These marine *Thalassiosira* were

divided among four clades, but the arrangement of these clades and the freshwater

cyclostephanoids varied across analyses. Uncertainty in the backbone relationships for this part

of the tree was likely caused by a combination of gene tree error and high levels of incomplete

lineage sorting. First, many individual genes contained too little information to confidently

resolve deep splits separated by short branch lengths—a finding that is not unique to this dataset

(Chan et al. 2020; Arcila et al. 2021). Divergence time estimates suggest that these splits

occurred in relatively quick succession, as few as 5 million years. Although many of these

bipartitions had consistently weak support, gene concordance factors increased when we

analyzed orthologs with the highest phylogenetic signal, implicating gene tree error as one of the

sources of instability (Chan et al. 2020; Vanderpool et al. 2020). In other phylogenomic studies

impacted by low phylogenetic signal and high gene tree error, gene genealogy interrogation

(GGI) has been used to overcome noise and identify majority gene support for a single

hypothesis (Hughes et al. 2018; Tea et al. 2021). In our case, no single resolution was supported

by a majority of genes, indicative of nodes that are difficult to resolve even with hundreds of

genes (Nesi et al. 2021). The best-supported constraint tree identified by GGI differed between

amino acid and codon-based gene trees, highlighting conflicting signal even within genes.

Second, in addition to gene tree error, the large number of alternative topologies among gene

trees along the *Thalassiosira* grade is also consistent with ILS (Arcila et al. 2017). Gene tree

summary methods such as ASTRAL outperform concatenation when ILS is the major source of

discordance (Kubatko and Degnan 2007; Roch and Warnow 2015), but summary methods

perform poorly when gene tree error is high (Roch and Warnow 2015; Xi et al. 2015). Following

Arcila et al. (2017), we attempted to eliminate some of the noise in our dataset by restricting

ASTRAL analyses to the top-ranked constrained gene trees and in doing so recovered a

backbone topology congruent with one of the few originally recovered by both gene tree

summary and concatenation methods (topology 5; Table 1 and Fig. 2). Taken together, these

results suggest that gene tree error negatively impacted our ASTRAL analyses. After identifying

and removing some of that error, we were able to recover a stronger hypothesis for the placement

of the freshwater cyclostephanoids within the marine *Thalassiosira* grade.

The anomaly zone describes an especially vexing phylogenetic problem in which short

branch lengths are unresolvable, resulting in gene trees that differ from the species more

frequently than they agree (Degnan and Rosenberg 2006). Within the *Thalassiosira* grade, the

most common GGI gene tree topologies either did not match the reference species tree or were

uninformative for these short branches. This implies that unresolved gene trees (i.e., those with

polytomies) are more probable than resolved ones in the *Thalassiosira* grade, as branch lengths

that exceed the boundaries of the anomaly zone should produce resolved gene trees (Huang and

Knowles 2009). Gene tree error like that identified in our dataset can lead to underestimation of

coalescent branch lengths, which define the anomaly zone boundaries (Linkem et al. 2016;

Forthman et al. 2022), resulting in the mistaken identification of an anomaly zone where none

exists. Although not entirely clear in our case, the *Thalassiosira* grade appears to fall outside of

the anomaly zone, so the large number of conflicting gene histories is more likely due to ILS and

gene tree error. Challenges remain in anomaly zone detection due to the limited application and

computational costs of multispecies coalescent methods applied to large genomic datasets and

the inclusion of additional biological factors other than ILS (Flouri et al. 2018, 2020).

Factors that are more poorly known in diatoms might also have contributed to the high

levels of discordance. We attempted to minimize technical sources of gene tree error during

dataset construction (Supplemental File S1), but other sources are more difficult to discern (Cai

et al. 2021). These include hybridization and polyploidy, gene duplication and loss, or

recombination. Although hybridization is not well documented in diatoms (Casteleyn et al. 2009;

Koester et al. 2010; Tanaka et al. 2015), there is evidence for an ancient allopolyploidy event

early on in the evolutionary history of Thalassiosirales (Parks et al. 2018). Many methods that

identify hybridization based on gene trees (Edelman et al. 2019; Vanderpool et al. 2020) or site

patterns in an alignment (Blischak et al. 2018) can be confounded by ancestral population

structure, substitutional saturation, or unsampled ghost lineages (Slatkin and Pollack 2008;

Tricou et al. 2022), all of which are poorly characterized in diatoms. High GC content regions

have been linked to higher recombination rates (Kent et al. 2012; Lartillot 2013) and can lead to

increased discordance (Pease and Hahn 2013). Despite high levels of variation in GC content

across Thalassiosirales, intralocus recombination was detected in a vanishingly small (<0.001%)

percent of the 6262 orthologs in our analysis (results available on Dryad). In addition to showing

whether any of these factors contribute to gene tree discordance, a more thorough exploration of

each one will fill important gaps in our understanding of diatom evolution.

Codon bias and amino acid composition in freshwater diatoms

Thalassiosirales includes divergence times across a timescale ranging from thousands

(Theriot et al. 2006) to tens of millions of years ago (Fig. 3a), which led us to explore the utility

of both amino acid and nucleotide characters for resolving phylogenetic relationships. Amino

acids are less susceptible to saturation and useful for resolving deep relationships (Philippe et al.

2011; Rota-Stabelli et al. 2012), whereas nucleotides contain more information to resolve more

recent divergences (Simmons et al. 2002; Townsend et al. 2008). Both data types recovered the

vast majority of relationships consistently and with strong support, while at the same time

revealing similar patterns of discordance along the backbone of the tree. In many cases, however,

they differed in their resolutions of the most recalcitrant parts of the tree. Disagreements between

character types within the same dataset, such as those within the *Thalassiosira* grade here, have

been found in other groups as well (Gillung et al. 2018; Skinner et al. 2020).

Almost every analysis of the amino acid dataset—including species trees, concordance

factors, and AU tests—supported the placement of cyclostephanoids as sister to the remaining

Thalassiosira clades, but nucleotide analyses placed them with *Thalassiosira* III (Fig. 2).

Discordance caused by codon usage bias and differences in amino acid composition might

account for this discrepancy. An association between codon bias and ecology has been

demonstrated in a broad diversity of microbes, where species that share an ecological niche have

similar codon usage, independent of phylogeny (Botzman and Margalit 2011; Roller et al. 2013;

Arella et al. 2021). Similar patterns of codon usage within marine and freshwater habitats have

been described in prokaryotes (Cabello-Yeves and Rodriguez-Valera 2019), and we discovered

differences in both codon usage and amino acid composition between marine and freshwater

diatoms. The amino acid compositions of distantly related freshwater lineages might be

sufficiently similar to cause the amino acid characters to support the "sister to the rest"

placement of cyclostephanoids. Protein sites with different structural, functional, or selective

constraints can lead to differences in amino acid composition between species (Villar and

Kauvar 1994; Youssef et al. 2021), something that is not accounted for by standard empirical

protein models and may have led to artifacts in our gene and species tree inferences (Wang et al.

2018). When we applied the PMSF model, which accounts for substitution heterogeneity in

amino acid sites, cyclostephanoids were placed as sister to *Thalassiosira* III, in agreement with

the codon datasets. The similarity in codon usage and amino acid composition between distantly

related freshwater diatoms merits further study into the causes and functional significance, if

any.

Conclusions

The vast differences between marine and freshwaters result in strong selective pressures

on freshwater colonists. Indeed, some of these transitions have become some of our most

powerful model systems for studying convergent evolution (Elmer and Meyer 2011). Low

salinity provokes a broad range of physiological and metabolic responses in diatoms (Nakov et

al. 2020; Downey et al. 2022; Pinseel et al. 2022), but the current genetic architectures of

freshwater adaptation reflect tens of millions of years of optimization and change since the

earliest transitions. As a result, it may not be possible to distinguish the derived alleles that

currently allow these diatoms to thrive in freshwaters from the ancestral ones that made the first

transitions possible. Nevertheless, the phylogenomic analyses presented here highlighted a key

role for hemiplasy in charting the genetic trajectory for freshwater colonizations in this group of

diatoms, indicating that some of the necessary alleles were already present in the ancestral

marine populations. Some of the subsequent mutations that built upon this scaffold of hemiplasy

might be identifiable by the different evolutionary outcomes in which some taxa became

irreversibly locked into freshwaters (e.g., cyclostephanoids) while others were able to transition

back to the ocean or became salinity generalists (e.g., *Cyclotella*). The vast new genomic

resources and phylogenetic framework presented here represent an important step forward in

addressing these types of questions to better understand how diatoms have made this complex

ecological transition appear to be so simple superficially.

SUPPLEMENTAL MATERIAL

Datasets and scripts are available from the Dryad Digital Repository:

http://dx.doi.org/10.5061/dryad.[NNNN]

FUNDING

This work was supported by the National Science Foundation (DEB 1651087 to A.J.A).

This research used resources available through the Arkansas High Performance Computing

Center, which is funded through multiple NSF grants and the Arkansas Economic Development

Commission.

ACKNOWLEDGEMENTS

We thank Ed Theriot for sharing several culture strains, and we thank Matt Ashworth and

Jeffery Stone for help with scanning electron microscopy.

LITERATURE CITED

Alverson A.J., Beszteri B., Julius M.L., Theriot E.C. 2011. The model marine diatom
     Thalassiosira pseudonana likely descended from a freshwater ancestor in the genus
     Cyclotella. BMC Evol. Biol. 11:125.

Alverson A.J., Jansen R.K., Theriot E.C. 2007. Bridging the Rubicon: Phylogenetic analysis
     reveals repeated colonizations of marine and fresh waters by thalassiosiroid diatoms. Mol.
     Phylogenet. Evol. 45:193–210.

Arcila D., Hughes L.C., Meléndez-Vazquez B., Baldwin C.C., White W.T., Carpenter K.E.,
     Williams J.T., Santos M.D., Pogonoski J.J., Miya M., Ortí G., Betancur-R R. 2021. Testing
     the Utility of Alternative Metrics of Branch Support to Address the Ancient Evolutionary
     Radiation of Tunas, Stromateoids, and Allies (Teleostei: Pelagiaria). Syst. Biol. 70:1123–
     1144.

Arcila D., Ortí G., Vari R., Armbruster J.W., Stiassny M.L.J., Ko K.D., Sabaj M.H., Lundberg J.,
     Revell L.J., Betancur-R. R. 2017. Genome-wide interrogation advances resolution of
     recalcitrant groups in the tree of life. Nature Ecology & Evolution. 1:1–10.

Arella D., Dilucca M., Giansanti A. 2021. Codon usage bias and environmental adaptation in
     microbial organisms. Mol. Genet. Genomics. 296:751–762.

Armbrust E.V., Berges J.A., Bowler C., Green B.R., Martinez D., Putnam N.H., Zhou S., Allen
     A.E., Apt K.E., Bechner M., Brzezinski M.A., Chaal B.K., Chiovitti A., Davis A.K.,
     Demarest M.S., Detter J.C., Glavina T., Goodstein D., Hadi M.Z., Hellsten U., Hildebrand
     M., Jenkins B.D., Jurka J., Kapitonov V.V., Kröger N., Lau W.W.Y., Lane T.W., Larimer

F.W., Lippmeier J.C., Lucas S., Medina M., Montsant A., Obornik M., Parker M.S., Palenik

B., Pazour G.J., Richardson P.M., Rynearson T.A., Saito M.A., Schwartz D.C.,

Thamatrakoln K., Valentin K., Vardi A., Wilkerson F.P., Rokhsar D.S. 2004. The genome

of the diatom *Thalassiosira pseudonana*: Ecology, evolution, and metabolism. Science.

306:79–86.

Artemov A.V., Mugue N.S., Rastorguev S.M., Zhenilo S., Mazur A.M., Tsygankova S.V.,

Boulygina E.S., Kaplun D., Nedoluzhko A.V., Medvedeva Y.A., Prokhortchouk E.B. 2017.

Genome-Wide DNA Methylation Profiling Reveals Epigenetic Adaptation of Stickleback to

Marine and Freshwater Conditions. Mol. Biol. Evol. 34:2203–2213.

Avise J.C., Robinson T.J. 2008. Hemiplasy: A new term in the lexicon of phylogenetics. Syst.

Biol. 57:503–507.

Beaulieu J.M., O'Meara B.C. 2016. Detecting Hidden Diversification Shifts in Models of Trait-

Dependent Speciation and Extinction. Syst. Biol. 65:583–601.

Bedford T., Hartl D.L. 2009. Optimization of gene expression by natural selection. Proc. Natl.

Acad. Sci. U. S. A. 106:1133–1138.

Blischak P.D., Chifman J., Wolfe A.D., Kubatko L.S. 2018. HyDe: A Python Package for

Genome-Scale Hybridization Detection. Syst. Biol. 67:821–829.

Borowiec M.L. 2016. AMAS: a fast tool for alignment manipulation and computing of summary

statistics. PeerJ. 4:e1660.

Botzman M., Margalit H. 2011. Variation in global codon usage bias among prokaryotic

organisms is associated with their lifestyles. Genome Biol. 12:R109.

Bussard A., Corre E., Hubas C., Duvernois-Berthet E., Le Corguillé G., Jourdren L., Coulpier F.,

Claquin P., Lopez P.J. 2017. Physiological adjustments and transcriptome reprogramming

are involved in the acclimation to salinity gradients in diatoms. Environ. Microbiol. 19:909–

925.

Cabello-Yeves P.J., Rodriguez-Valera F. 2019. Marine-freshwater prokaryotic transitions require

extensive changes in the predicted proteome. Microbiome. 7:117.

Cai L., Xi Z., Lemmon E.M., Lemmon A.R., Mast A., Buddenhagen C.E., Liu L., Davis C.C.

2021. The Perfect Storm: Gene Tree Estimation Error, Incomplete Lineage Sorting, and

Ancient Gene Flow Explain the Most Recalcitrant Ancient Angiosperm Clade,

Malpighiales. Syst. Biol. 70:491–507.

Casteleyn G., Adams N.G., Vanormelingen P., Debeer A.-E., Sabbe K., Vyverman W. 2009.

Natural hybrids in the marine diatom *Pseudo-nitzschia pungens* (Bacillariophyceae):

genetic and morphological evidence. Protist. 160:343–354.

Chan K.O., Hutter C.R., Wood P.L. Jr, Grismer L.L., Brown R.M. 2020. Target-capture

phylogenomics provide insights on gene and species tree discordances in Old World

treefrogs (Anura: Rhacophoridae). Proc. Biol. Sci. 287:20202102.

Chen M.-Y., Teng W.-K., Zhao L., Hu C.-X., Zhou Y.-K., Han B.-P., Song L.-R., Shu W.-S.

2021. Comparative genomics reveals insights into cyanobacterial evolution and habitat

adaptation. ISME J. 15:211–227.

Copetti D., Búrquez A., Bustamante E., Charboneau J.L.M., Childs K.L., Eguiarte L.E., Lee S.,

Liu T.L., McMahon M.M., Whiteman N.K., Wing R.A., Wojciechowski M.F., Sanderson

M.J. 2017. Extensive gene tree discordance and hemiplasy shaped the genomes of North

American columnar cacti. Proc. Natl. Acad. Sci. U. S. A. 114:12003–12008.

DeFaveri J., Shikano T., Shimada Y., Goto A., Merilä J. 2011. Global analysis of genes involved

in freshwater adaptation in threespine sticklebacks (*Gasterosteus aculeatus*). Evolution.

65:1800–1807.

Degnan J.H., Rosenberg N.A. 2006a. Discordance of species trees with their most likely gene

trees. PLoS Genet. 2:e68.

Dickson B., Yashayaev I., Meincke J., Turrell B., Dye S., Holfort J. 2002. Rapid freshening of

the deep North Atlantic Ocean over the past four decades. Nature. 416:832–837.

Dittami S.M., Heesch S., Olsen J.L., Collén J. 2017. Transitions between marine and freshwater

environments provide new clues about the origins of multicellular plants and algae. J.

Phycol. 53:731–745.

Downey K.M., Judy K.J., Pinseel E., Alverson A.J., Lewis J.A. 2022. The dynamic response to

hypoosmotic stress reveals distinct stages of freshwater acclimation by a euryhaline diatom.

bioRxiv.:2022.06.24.497401.

Edelman N.B., Frandsen P.B., Miyagi M., Clavijo B., Davey J., Dikow R.B., García-Accinelli

G., Van Belleghem S.M., Patterson N., Neafsey D.E., Challis R., Kumar S., Moreira G.R.P.,

Salazar C., Chouteau M., Counterman B.A., Papa R., Blaxter M., Reed R.D., Dasmahapatra

K.K., Kronforst M., Joron M., Jiggins C.D., McMillan W.O., Di Palma F., Blumberg A.J.,

Wakeley J., Jaffe D., Mallet J. 2019. Genomic architecture and introgression shape a

725   butterfly radiation. Science. 366:594–599.

Edwards S.V. 2009. Is a new and general theory of molecular systematics emerging? Evolution.

63:1–19.

Elmer K.R., Meyer A. 2011. Adaptation in the age of ecological genomics: insights from

parallelism and convergence. Trends Ecol. Evol. 26:298–306.

Emms D.M., Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative

genomics. Genome Biol. 20:238.

Felsenstein J. 1985. Phylogenies and the Comparative Method. Am. Nat. 125:1–15.

Flouri T., Jiao X., Rannala B., Yang Z. 2018. Species Tree Inference with BPP Using Genomic

Sequences and the Multispecies Coalescent. Mol. Biol. Evol. 35:2585–2593.

Flouri T., Jiao X., Rannala B., Yang Z. 2020. A Bayesian Implementation of the Multispecies

Coalescent Model with Introgression for Phylogenomic Analysis. Mol. Biol. Evol.

37:1211–1223.

Forthman M., Braun E.L., Kimball R.T. 2022. Gene tree quality affects empirical coalescent

branch length estimation. Zool. Scr. 51:1–13.

Foster P.G. 2004. Modeling compositional heterogeneity. Syst. Biol. 53:485–495.

Gillung J.P., Winterton S.L., Bayless K.M., Khouri Z., Borowiec M.L., Yeates D., Kimsey L.S.,

Misof B., Shin S., Zhou X., Mayer C., Petersen M., Wiegmann B.M. 2018. Anchored

phylogenomics unravels the evolution of spider flies (Diptera, Acroceridae) and reveals

744 discordance between nucleotides and amino acids. Mol. Phylogenet. Evol. 128:233–245.

745 Gomez-Mestre I., Jovani R. 2013. A heuristic model on the role of plasticity in adaptive

746 evolution: plasticity increases adaptation, population viability and genetic variation. Proc.

747 Biol. Sci. 280:20131869.

748 Guerrero R.F., Hahn M.W. 2018. Quantifying the risk of hemiplasy in phylogenetic inference.

749 Proc. Natl. Acad. Sci. U. S. A. 115:12787–12792.

750 Guillard R.R.L., Ryther J.H. 1962. Studies of marine planktonic diatoms: Guillard R. R. L. and J.

751 H. Ryther, 1962. I. *Cyclotella nana* Hustedt, and *Detonula confervacea* (Cleve) Gran. Can J

752 Microbiol. 8:229–240.

753 Hahn M.W., Nakhleh L. 2016. Irrational exuberance for resolved species trees. Evolution. 70:7–

754 17.

755 Hibbins M.S., Gibson M.J., Hahn M.W. 2020. Determining the probability of hemiplasy in the

756 presence of incomplete lineage sorting and introgression. Elife. 9.

757 Huang H., Knowles L.L. 2009. What is the danger of the anomaly zone for empirical

758 phylogenetics? Syst. Biol. 58:527–536.

759 Hughes L.C., Ortí G., Huang Y., Sun Y., Baldwin C.C., Thompson A.W., Arcila D., Betancur-R

760 R., Li C., Becker L., Bellora N., Zhao X., Li X., Wang M., Fang C., Xie B., Zhou Z., Huang

761 H., Chen S., Venkatesh B., Shi Q. 2018. Comprehensive phylogeny of ray-finned fishes

762 (Actinopterygii) based on transcriptomic and genomic data. Proc. Natl. Acad. Sci. U. S. A.

763 115:6249–6254.

Hughes L.C., Somoza G.M., Nguyen B.N., Bernot J.P., González-Castro M., Díaz de Astarloa

J.M., Ortí G. 2017. Transcriptomic differentiation underlying marine-to-freshwater

transitions in the South American silversides *Odontesthes argentinensis* and *O. bonariensis*

(Atheriniformes). Ecol. Evol. 7:5258–5268.

Jombart T., Kendall M., Almagro-Garcia J., Colijn C. 2017. treespace: Statistical exploration of

landscapes of phylogenetic trees. Mol. Ecol. Resour. 17:1385–1392.

Jones F.C., Grabherr M.G., Chan Y.F., Russell P., Mauceli E., Johnson J., Swofford R., Pirun

M., Zody M.C., White S., Birney E., Searle S., Schmutz J., Grimwood J., Dickson M.C.,

Myers R.M., Miller C.T., Summers B.R., Knecht A.K., Brady S.D., Zhang H., Pollen A.A.,

Howes T., Amemiya C., Broad Institute Genome Sequencing Platform & Whole Genome

Assembly Team, Baldwin J., Bloom T., Jaffe D.B., Nicol R., Wilkinson J., Lander E.S., Di

Palma F., Lindblad-Toh K., Kingsley D.M. 2012. The genomic basis of adaptive evolution

in threespine sticklebacks. Nature. 484:55–61.

Kalyaanamoorthy S., Minh B.Q., Wong T.K.F., von Haeseler A., Jermiin L.S. 2017.

ModelFinder: fast model selection for accurate phylogenetic estimates. Nat. Methods.

14:587–589.

Katoh K., Standley D.M. 2013. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. Mol. Biol. Evol. 30:772–780.

Kenny N.J., Plese B., Riesgo A., Itskovich V.B. 2019. Symbiosis, Selection and Novelty:

Freshwater Adaptation in the Unique Sponges of Lake Baikal. Mol. Biol. Evol.

Kent C.F., Minaei S., Harpur B.A., Zayed A. 2012. Recombination is associated with the

evolution of genome structure and worker behavior in honey bees. Proc. Natl. Acad. Sci. U.

S. A. 109:18012–18017.

Kirst G.O. 1990. Salinity Tolerance of Eukaryotic Marine Algae. Annu. Rev. Plant Physiol. Plant

Mol. Biol. 41:21–53.

Kirst G.O. 1996. Osmotic Adjustment in Phytoplankton and MacroAlgae. In: Kiene R.P.,

Visscher P.T., Keller M.D., Kirst G.O., editors. Biological and Environmental Chemistry of

DMSP and Related Sulfonium Compounds. Boston, MA: Springer US. p. 121–129.

Koester J.A., Swalwell J.E., von Dassow P., Armbrust E.V. 2010. Genome size differentiates co-

occurring populations of the planktonic diatom *Ditylum brightwellii* (Bacillariophyta). BMC

Evol. Biol. 10:1.

Kubatko L.S., Degnan J.H. 2007. Inconsistency of phylogenetic estimates from concatenated

data under coalescence. Syst. Biol. 56:17–24.

Lartillot N. 2013. Phylogenetic patterns of GC-biased gene conversion in placental mammals and

the evolutionary dynamics of recombination landscapes. Mol. Biol. Evol. 30:489–502.

Lartillot N., Philippe H. 2004. A Bayesian mixture model for across-site heterogeneities in the

amino-acid replacement process. Mol. Biol. Evol. 21:1095–1109.

L E E C.E., Downey K., Colby R.S., Freire C.A., Nichols S., Burgess M.N., Judy K.J. 2022.

Recognizing salinity threats in the climate crisis. Integr. Comp. Biol.

Linkem C.W., Minin V.N., Leaché A.D. 2016. Detecting the Anomaly Zone in Species Trees

and Evidence for a Misleading Signal in Higher-Level Skink Phylogeny (Squamata:

Scincidae). Syst. Biol. 65:465–477.

Liu L., Wu S., Yu L. 2015. Coalescent methods for estimating species trees from phylogenomic

data. J. Syst. Evol. 53:380–390.

Logares R., Bråte J., Bertilsson S., Clasen J.L., Shalchian-Tabrizi K., Rengefors K. 2009.

Infrequent marine-freshwater transitions in the microbial world. Trends Microbiol. 17:414–

422.

López-Maury L., Marguerat S., Bähler J. 2008. Tuning gene expression to changing

environments: from rapid responses to evolutionary adaptation. Nat. Rev. Genet. 9:583–

593.

Lozupone C.A., Knight R. 2007. Global patterns in bacterial diversity. Proc. Natl. Acad. Sci. U.

S. A. 104:11436–11440.

Maddison W.P. 1997. Gene Trees in Species Trees. Syst. Biol. 46:523–536.

Mallet J., Besansky N., Hahn M.W. 2016. How reticulated are species? Bioessays. 38:140–149.

Mann D.G. 1999. Crossing the Rubicon : the effectiveness of the marine/freshwater interface as

a barrier to the migration of diatom germplasm. Proceedings of the 14th International

Diatom Symposium, Koenigstein, 1999.:1–21.

McCairns R.J.S., Bernatchez L. 2010. Adaptive divergence between freshwater and marine

sticklebacks: insights into the role of phenotypic plasticity from an integrated analysis of

candidate gene expression. Evolution. 64:1029–1047.

McInerney J.O. 1998. GCUA: general codon usage analysis. Bioinformatics. 14:372–373.

Mendes F.K., Hahn Y., Hahn M.W. 2016. Gene Tree Discordance Can Generate Patterns of

Diminishing Convergence over Time. Mol. Biol. Evol. 33:3299–3307.

Minh B.Q., Hahn M.W., Lanfear R. 2020a. New Methods to Calculate Concordance Factors for

Phylogenomic Datasets. Mol. Biol. Evol. 37:2727–2733.

Minh B.Q., Nguyen M.A.T., von Haeseler A. 2013. Ultrafast Approximation for Phylogenetic

Bootstrap. Mol. Biol. Evol. 30:1188–1195.

Minh B.Q., Schmidt H.A., Chernomor O., Schrempf D., Woodhams M.D., von Haeseler A.,

Lanfear R. 2020b. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic

Inference in the Genomic Era. Mol. Biol. Evol. 37:1530–1534.

Mirarab S. 2017. Phylogenomics: Constrained gene tree inference. Nat Ecol Evol. 1:56.

Mirarab S., Reaz R., Bayzid M.S., Zimmermann T., Swenson M.S., Warnow T. 2014. ASTRAL:

genome-scale coalescent-based species tree estimation. Bioinformatics. 30:i541–i548.

Mitterboeck T.F., Chen A.Y., Zaheer O.A., Ma E.Y.T., Adamowicz S.J. 2016. Do saline taxa

evolve faster? Comparing relative rates of molecular evolution between freshwater and

marine eukaryotes. Evolution. 70:1960–1978.

Molloy E.K., Warnow T. 2018. To Include or Not to Include: The Impact of Gene Filtering on

Species Tree Estimation Methods. Syst. Biol. 67:285–303.

Morales-Briones D.F., Kadereit G., Tefarikis D.T., Moore M.J., Smith S.A., Brockington S.F.,

Timoneda A., Yim W.C., Cushman J.C., Yang Y. 2020. Disentangling Sources of Gene

Tree Discordance in Phylogenomic Data Sets: Testing Ancient Hybridizations in

Amaranthaceae s.l. Syst. Biol. 70:219–235.

Nakov T., Beaulieu J.M., Alverson A.J. 2018. Accelerated diversification is related to life history

and locomotion in a hyperdiverse lineage of microbial eukaryotes (Diatoms,

Bacillariophyta). New Phytol. 219:462–473.

Nakov T., Beaulieu J.M., Alverson A.J. 2019. Diatoms diversify and turn over faster in

freshwater than marine environments. Evolution. 73:2497–2511.

Nakov T., Judy K.J., Downey K.M., Ruck E.C., Alverson A.J. 2020. Transcriptional Response of

Osmolyte Synthetic Pathways and Membrane Transporters in a Euryhaline Diatom During

Long-term Acclimation to a Salinity Gradient. J. Phycol. 56:1712–1728.

Naser-Khdour S., Minh B.Q., Zhang W., Stone E.A., Lanfear R. 2019. The Prevalence and

Impact of Model Violations in Phylogenetic Analysis. Genome Biol. Evol. 11:3341–3352.

Nawaly H., Tsuji Y., Matsuda Y. 2020. Rapid and precise genome editing in a marine diatom,

Thalassiosira pseudonana by Cas9 nickase (D10A). Algal Research. 47:101855.

Nesi N., Tsagkogeorga G., Tsang S.M., Nicolas V., Lalis A., Scanlon A.T., Riesle-Sbarbaro

S.A., Wiantoro S., Hitch A.T., Juste J., Pinzari C.A., Bonaccorso F.J., Todd C.M., Lim

B.K., Simmons N.B., McGowen M.R., Rossiter S.J. 2021. Interrogating Phylogenetic

Discordance Resolves Deep Splits in the Rapid Radiation of Old World Fruit Bats

(Chiroptera: Pteropodidae). Syst. Biol. 70:1077–1089.

Parks M.B., Nakov T., Ruck E.C., Wickett N.J., Alverson A.J. 2018. Phylogenomics reveals an

extensive history of genome duplication in diatoms (Bacillariophyta). Am. J. Bot. 105:330–

347.

Paver S.F., Muratore D., Newton R.J., Coleman M.L. 2018. Reevaluating the Salty Divide:

Phylogenetic Specificity of Transitions between Marine and Freshwater Systems.

mSystems. 3.

Pease J.B., Brown J.W., Walker J.F., Hinchliff C.E., Smith S.A. 2018. Quartet Sampling

distinguishes lack of support from conflicting support in the green plant tree of life. Am. J.

Bot. 105:385–403.

Pease J.B., Hahn M.W. 2013. More accurate phylogenies inferred from low-recombination

regions in the presence of incomplete lineage sorting. Evolution. 67:2376–2384.

Philippe H., Brinkmann H., Lavrov D.V., Littlewood D.T.J., Manuel M., Wörheide G., Baurain

D. 2011. Resolving difficult phylogenetic questions: why more sequences are not enough.

PLoS Biol. 9:e1000602.

Phillips M.J., Penny D. 2003. The root of the mammalian tree inferred from whole mitochondrial

genomes. Mol. Phylogenet. Evol. 28:171–185.

Pinseel E., Nakov T., Van den Berge K., Downey K.M., Judy K.J., Kourtchenko O., Kremp A.,

Ruck E.C., Sjöqvist C., Töpel M., Godhe A., Alverson A.J. 2022. Strain-specific

transcriptional responses overshadow salinity effects in a marine diatom sampled along the

Baltic Sea salinity cline. ISME J.

Prabha R., Singh D.P., Sinha S., Ahmad K., Rai A. 2017. Genome-wide comparative analysis of

codon usage bias and codon context patterns among cyanobacterial genomes. Mar.

Genomics. 32:31–39.

Price M.N., Dehal P.S., Arkin A.P. 2010. FastTree 2--approximately maximum-likelihood trees

for large alignments. PLoS One. 5:e9490.

Regier J.C., Shultz J.W., Zwick A., Hussey A., Ball B., Wetzer R., Martin J.W., Cunningham

C.W. 2010. Arthropod relationships revealed by phylogenomic analysis of nuclear protein-

coding sequences. Nature. 463:1079–1083.

Reis M. dos, Yang Z. 2011. Approximate Likelihood Calculation on a Phylogeny for Bayesian

Estimation of Divergence Times. Mol. Biol. Evol. 28:2161–2172.

Roberts W.R., Downey K.M., Ruck E.C., Traller J.C., Alverson A.J. 2020. Improved Reference

Genome for *Cyclotella cryptica* CCMP332, a Model for Cell Wall Morphogenesis, Salinity

Adaptation, and Lipid Production in Diatoms (Bacillariophyta). G3 . 10:2965–2974.

[dataset]* Roberts W.R., Ruck E.C., Downey K.M., Alverson A.J. 2022. Resolving marine–

freshwater transitions by diatoms through a fog of discordance and hemiplasy. Dryad.

http://dx.doi.org/10.5061/dryad.[NNNN].

Roch S., Warnow T. 2015. On the Robustness to Gene Tree Estimation Error (or lack thereof) of

Coalescent-Based Species Tree Methods. Syst. Biol. 64:663–676.

Rogers R.L., Grizzard S.L., Titus-McQuillan J.E., Bockrath K., Patel S., Wares J.P., Garner J.T.,

Moore C.C. 2021. Gene family amplification facilitates adaptation in freshwater unionid

bivalve Megalonaias nervosa. Mol. Ecol. 30:1155–1173.

Roller M., Lucić V., Nagy I., Perica T., Vlahovicek K. 2013. Environmental shaping of codon

usage and functional adaptation across microbial communities. Nucleic Acids Res.

41:8842–8852.

Rosenberg N.A. 2003. The shapes of neutral gene genealogies in two species: probabilities of

monophyly, paraphyly, and polyphyly in a coalescent model. Evolution. 57:1465–1477.

Rota-Stabelli O., Lartillot N., Philippe H., Pisani D. 2012. Serine Codon-Usage Bias in Deep

Phylogenomics: Pancrustacean Relationships as a Case Study. Syst. Biol. 62:121–133.

Salichos L., Stamatakis A., Rokas A. 2014. Novel information theory-based measures for

quantifying incongruence among phylogenetic trees. Mol. Biol. Evol. 31:1261–1271.

Sanderson M.J., Wojciechowski M.F., Hu J.M., Khan T.S., Brady S.G. 2000. Error, bias, and

long-branch attraction in data for two chloroplast photosystem genes in seed plants. Mol.

Biol. Evol. 17:782–797.

Sayyari E., Mirarab S. 2016. Fast Coalescent-Based Computation of Local Branch Support from

Quartet Frequencies. Mol. Biol. Evol. 33:1654–1668.

Sayyari E., Mirarab S. 2018. Testing for Polytomies in Phylogenetic Species Trees Using

Quartet Frequencies. Genes . 9.

Shimodaira H. 2002. An approximately unbiased test of phylogenetic tree selection. Syst. Biol.

51:492–508.

Simmons M.P., Gatesy J. 2021. Collapsing dubiously resolved gene-tree branches in

phylogenomic coalescent analyses. Mol. Phylogenet. Evol. 158:107092.

Simmons M.P., Ochoterena H., Freudenstein J.V. 2002. Amino acid vs. nucleotide characters:

challenging preconceived notions. Mol. Phylogenet. Evol. 24:78–90.

Skinner R.K., Dietrich C.H., Walden K.K.O., Gordon E., Sweet A.D., Podsiadlowski L.,

Petersen M., Simon C., Takiya D.M., Johnson K.P. 2020. Phylogenomics of

Auchenorrhyncha (Insecta: Hemiptera) using transcriptomes: examining controversial

relationships via degeneracy coding and interrogation of gene conflict. Syst. Entomol.

45:85–113.

Slatkin M., Pollack J.L. 2008. Subdivision in an ancestral species creates asymmetry in gene

trees. Mol. Biol. Evol. 25:2241–2246.

Smith S.A., Brown J.W., Walker J.F. 2018. So many genes, so little time: A practical approach

to divergence-time estimation in the genomic era. PLoS One. 13:e0197433.

Smith S.A., Moore M.J., Brown J.W., Yang Y. 2015. Analysis of phylogenomic datasets reveals

conflict, concordance, and gene duplications with examples from animals and plants. BMC

Evol. Biol. 15:150.

Steenwyk J.L., Buida T.J., Labella A.L., Li Y., Shen X.-X., Rokas A. 2021. PhyKIT: a broadly

applicable UNIX shell toolkit for processing and analyzing phylogenomic data.

Bioinformatics.

Stern D.B., Anderson N.W., Diaz J.A., Lee C.E. 2022. Genome-wide signatures of synergistic

epistasis during parallel adaptation in a Baltic Sea copepod. Nat. Commun. 13:4024.

Storz J.F. 2016. Causes of molecular convergence and parallelism in protein evolution. Nat. Rev.

Genet. 17:239–250.

Suyama M., Torrents D., Bork P. 2006. PAL2NAL: robust conversion of protein sequence

alignments into the corresponding codon alignments. Nucleic Acids Res. 34:W609–W612.

Tanaka T., Maeda Y., Veluchamy A., Tanaka M., Abida H., Maréchal E., Bowler C., Muto M.,

Sunaga Y., Tanaka M., Yoshino T., Taniguchi T., Fukuda Y., Nemoto M., Matsumoto M.,

Wong P.S., Aburatani S., Fujibuchi W. 2015. Oil accumulation by the oleaginous diatom

Fistulifera solaris as revealed by the genome and transcriptome. Plant Cell. 27:162–176.

Tea Y.-K., Xu X., DiBattista J.D., Lo N., Cowman P.F., Ho S.Y.W. 2021. Phylogenomic

Analysis of Concatenated Ultraconserved Elements Reveals the Recent Evolutionary

Radiation of the Fairy Wrasses (Teleostei: Labridae: *Cirrhilabrus*). Syst. Biol. 71:1–12.

Tedesco P.A., Paradis E., Lévêque C., Hugueny B. 2017. Explaining global-scale diversification

patterns in actinopterygian fishes. J. Biogeogr. 44:773–783.

Terekhanova N.V., Barmintseva A.E., Kondrashov A.S., Bazykin G.A., Mugue N.S. 2019.

Architecture of Parallel Adaptation in Ten Lacustrine Threespine Stickleback Populations

from the White Sea Area. Genome Biol. Evol. 11:2605–2618.

Theriot E.C., Fritz S.C., Whitlock C., Conley D.J. 2006. Late Quaternary rapid morphological

evolution of an endemic diatom in Yellowstone Lake, Wyoming. Paleobiology. 32:38–54.

Theriot E., Stoermer E., Håkansson H. 1987. Taxonomic interpretation of the rimoportula of

freshwater genera in the centric diatom family Thalassiosiraceae. Diatom Res. 2:251–265.

Townsend J.P., López-Giráldez F., Friedman R. 2008. The phylogenetic informativeness of

nucleotide and amino acid sequences for reconstructing the vertebrate tree. J. Mol. Evol.

67:437–447.

Tricou T., Tannier E., de Vienne D.M. 2022. Ghost Lineages Highly Influence the Interpretation

of Introgression Tests. Syst. Biol.

Vanderpool D., Minh B.Q., Lanfear R., Hughes D., Murali S., Harris R.A., Raveendran M.,

Muzny D.M., Hibbins M.S., Williamson R.J., Gibbs R.A., Worley K.C., Rogers J., Hahn

M.W. 2020. Primate phylogenomics uncovers multiple rapid radiations and ancient

interspecific introgression. PLoS Biol. 18:e3000954.

Villar H.O., Kauvar L.M. 1994. Amino acid preferences at protein binding sites. FEBS Lett.

349:125–130.

Vizueta J., Macías-Hernández N., Arnedo M.A., Rozas J., Sánchez-Gracia A. 2019. Chance and

predictability in evolution: The genomic basis of convergent dietary specializations in an

adaptive radiation. Mol. Ecol. 28:4028–4045.

Wang H.-C., Minh B.Q., Susko E., Roger A.J. 2018. Modeling Site Heterogeneity with Posterior

Mean Site Frequency Profiles Accelerates Accurate Phylogenomic Estimation. Syst. Biol.

67:216–235.

Wray G.A. 2007. The evolutionary significance of cis-regulatory mutations. Nat. Rev. Genet.

8:206–216.

Wu M., Kostyun J.L., Hahn M.W., Moyle L.C. 2018. Dissecting the basis of novel trait evolution

in a radiation with widespread phylogenetic discordance. Mol. Ecol.

Xi Z., Liu L., Davis C.C. 2015. Genes with minimal phylogenetic information are problematic

for coalescent analyses when gene tree estimation is biased. Mol. Phylogenet. Evol. 92:63–

71.

Yang Y., Smith S.A. 2014. Orthology inference in nonmodel organisms using transcriptomes

and low-coverage genomes: improving accuracy and matrix occupancy for phylogenomics.

Mol. Biol. Evol. 31:3081–3092.

Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. Mol. Biol. Evol.

24:1586–1591.

Youssef N., Susko E., Roger A.J., Bielawski J.P. 2021. Shifts in amino acid preferences as

proteins evolve: A synthesis of experimental and theoretical work. Protein Sci. 30:2009–

2028.

Yuan Z., Liu S., Zhou T., Tian C., Bao L., Dunham R., Liu Z. 2018. Comparative genome

analysis of 52 fish species suggests differential associations of repetitive elements with their

living aquatic environments. BMC Genomics. 19:141.

Zhang C., Rabiee M., Sayyari E., Mirarab S. 2018. ASTRAL-III: polynomial time species tree

reconstruction from partially resolved gene trees. BMC Bioinformatics. 19:153.

Zou Z., Zhang J. 2015. Are Convergent and Parallel Amino Acid Substitutions in Protein

Evolution More Prevalent Than Neutral Expectations? Mol. Biol. Evol. 32:2085–2096.

Zwick A., Regier J.C., Zwickl D.J. 2012. Resolving discrepancy between nucleotides and amino

acids in deep-level arthropod phylogenomics: differentiating serine codons in 21-amino-

acid models. PLoS One. 7:e47450.

FIGURE LEGENDS

Figure 1. Phylogram based on maximum likelihood analysis of amino acids using the posterior

mean site frequency (PMSF) model and a dataset of 488 loci with the highest proportions of taxa

and informative sites ("AA-top-PI-top-Taxa" dataset; Table 1). Backbone nodes of the

Thalassiosira grade are indicated by the letters A–E. All branches had bootstrap support (BS)

values of 100 except for those with black circles which had BS = 90. Pie charts on backbone

nodes show the proportion of gene trees that support the clade (gCF), the proportion that support

both discordant topologies (gDF1, gDF2), and the proportion that are discordant due to

polyphyly (gDFP). The size of the pie charts is only for aesthetics.

1014

Figure 2. **a)** Phylogenetic hypotheses of the *Thalassiosira* grade inferred using concatenation and

summary methods on the amino acid (AA), codon positions 1 and 2 (CDS12), and recoded

codon (DEGEN) datasets. Nodes are labeled with the percentage of amino acid sites concordant

with the branch (site concordance factor). The principal clade of interest, the freshwater

cyclostephanoids, is colored orange and labeled 'FW'. The four focal clades of marine

Thalassiosira and allies are labeled I–IV. Below, results of gene genealogy interrogation tests of

alternative hypotheses of relationships within the *Thalassiosira* grade. These tests used datasets

filtered to include only the top 25% of orthologs based on the percentage of parsimony

informative sites (top-PI) for **b)** amino acids, **c)** codon positions 1 and 2, and **d)** recoded codons.

Lines correspond to the cumulative number of genes (x-axis) supporting topology hypotheses

with the highest probability and their *P* values (y-axis) from the Approximately Unbiased (AU)

topology tests. Values above the dashed line indicate topological hypotheses that are

significantly better than the alternatives ($P < 0.05$). For example, the green line in (b) shows that

there were a total of 88 genes that best supported topology 3, while only four of those genes were

above the dotted line and were significantly better supported than the other four alternative

topologies.

1031

Figure 3. **a)** Divergence times and ancestral state reconstruction of marine and freshwater habitat

in the Thalassiosirales. Conspecific taxa were removed prior to ancestral state reconstruction,

leaving one tip per species. Tip labels are colored according to their habitat (blue=freshwater,

orange=euryhaline, grey=marine). Pie charts denote the probability of each node reconstructed as

either marine (grey) or freshwater (blue) using parameters estimated from the HiSSE CID–4

model. Euryhaline taxa were coded as marine for the purposes of ancestral state reconstruction.

Divergence times for the full set of taxa can be found in Supplementary Fig. S9. **b)** Hemiplasy

Risk Factors (HRF) on internal branches show increased values on most short internal branches,

demonstrating an increased potential for hemiplasy to influence trait reconstruction. Branch

lengths in coalescent units were inferred using ASTRAL.

a)

Porosira pseudodenticulata
Porosira glacialis
Lauderia annulata
Roundia cardiophora
Thalassiosira livingstoniorum
Thalassiosira lacustris
Conticribra guillardii
Conticribra weissflogii
Cyclotella nana
Cyclotella kingstonii
Cyclotella distinguenda
Cyclotella cryptica
Cyclotella meneghiniana
Cyclotella baltica
Cyclotella choctawhatcheeana
Cyclotella atomus
Mediolabrus comicus
Skeletonema potamos
Skeletonema dohrnii
Skeletonema marinoi
Skeletonema menzelii
Skeletonema japonicum
Skeletonema tropicum
Skeletonema grethae
Thalassiosira oceanica
Thalassiosira profunda
Thalassiosira exigua
Thalassiosira mediterranea
Thalassiosira sp
Bacterosira constricta
Thalassiosira sp
Thalassiosira ordinaria
Thalassiosira hispida
Thalassiosira allenii
Discostella pseudostelligera
Discostella stelligera
Discostella stelligeroides
Stephanodiscus triporus
Stephanodiscus minutulus
Cyclostephanos invisitatus
Cyclostephanos tholiformis
Thalassiosira sp
Thalassiosira pacifica
Detonula pumila
Detonula confervacea
Thalassiosira minuscula
Thalassiosira antarctica
Shionodiscus bioculatus
Shionodiscus oestrupii
Thalassiosira tumida
Thalassiosira punctigera
Thalassiosira delicatula
Thalassiosira gravida

● freshwater
● euryhaline
● marine

100   80   60   40   20   0
Million Years Ago (Ma)

b)

Hemiplasy
Risk
Factor
1.00
0.75
0.50
0.25
0.00

coalescent
units

1