TITLE: Resolving marine–freshwater transitions by diatoms through a fog of discordant gene

trees

RUNNING HEAD: Phylogenomics of marine and freshwater diatoms

AUTHORS:

Wade R. Roberts[1], Elizabeth C. Ruck[1], Kala M. Downey[1], Eveline Pinseel[1], Andrew J.

Alverson[1,*]

[1]Department of Biological Sciences, University of Arkansas, 1 University of Arkansas,

Fayetteville, AR, 72701, USA

[*]Corresponding author: Andrew J. Alverson, Department of Biological Sciences, University of

Arkansas, 1 University of Arkansas, Fayetteville, AR, 72701, USA; aja@uark.edu

18
19

ABSTRACT

Despite the obstacles facing marine colonists, most lineages of aquatic organisms have colonized

and diversified in freshwaters repeatedly. These transitions can trigger rapid morphological or

physiological change and, on longer timescales, lead to increased rates of speciation and

extinction. Diatoms are a lineage of ancestrally marine microalgae that have diversified

throughout freshwater habitats worldwide. We generated a phylogenomic dataset of genomes

and transcriptomes for 59 diatom taxa to resolve freshwater transitions in one lineage, the

Thalassiosirales. Although most parts of the species tree were consistently resolved with strong

support, we had difficulties resolving a Paleocene radiation, which affected the placement of one

freshwater lineage. This and other parts of the tree were characterized by high levels of gene tree

discordance caused by incomplete lineage sorting and low phylogenetic signal. Despite

differences in species trees inferred from concatenation versus summary methods and codons

versus amino acids, traditional methods of ancestral state reconstruction supported six transitions

into freshwaters, two of which led to subsequent species diversification. Evidence from gene

trees, protein alignments, and diatom life history together suggest that habitat transitions were

largely the product of homoplasy rather than hemiplasy, a condition where transitions occur on

branches in gene trees not shared with the species tree. Nevertheless, we identified a small set of

putatively hemiplasious genes, many of which have been associated with shifts to low salinity,

indicating that hemiplasy played a small but potentially important role in freshwater adaptation.

Accounting for differences in evolutionary outcomes, in which some taxa became locked into

freshwaters while others were able to return to the ocean or become salinity generalists, might

help further distinguish different sources of adaptive mutation in freshwater diatoms.

KEYWORDS: Gene concordance, Phylogenomics, Salinity, Site concordance, Thalassiosirales

INTRODUCTION

From bacteria to animals, the salinity gradient separating marine and freshwater

environments poses a significant barrier to the distributions of many organisms (Lozupone and

Knight 2007; McCairns and Bernatchez 2010; Kenny et al. 2019). Identifying how different

lineages cross the salinity divide will improve our understanding of lineage diversification

(Dittami et al. 2017) and the adaptive potential of species to climate change (Dickson et al. 2002;

Lee et al. 2022). Diatoms are a diverse lineage of microalgae that occur throughout marine and

freshwaters, and despite the numerous obstacles facing marine colonists (Kirst 1990, 1996;

Nakov et al. 2020), ancestrally marine diatoms have successfully colonized and diversified in

freshwaters repeatedly throughout their history (Nakov et al. 2019). These patterns are based on

phylogenetic analyses of a small number of molecular markers, however, so they lack the

insights of phylogenomic approaches, which can resolve large-scale macroevolutionary patterns

and, at the same time, uncover key processes at play during important evolutionary transitions

(Pease et al. 2016).

Although phylogenomic datasets have helped resolve historically recalcitrant nodes

across the tree of life, they have also revealed how discordance in the evolutionary histories of

different genes can confound inferences of species relationships. Systematic errors that lead to

gene tree discordance can be caused by biological sources, such as incomplete lineage sorting

(ILS) and hybridization (Maddison 1997; Degnan and Rosenberg 2006a), or methodological

sources, such as character sampling, compositional heterogeneity, and gene tree error (Foster

2004; Philippe et al. 2011; Xi et al. 2015; Molloy and Warnow 2018). Each of these challenges

our ability to resolve species relationships and impacts downstream analyses, such as estimation

of divergence times (Smith et al. 2018). Multiple strategies have been proposed to overcome

different sources of error, such as excluding third-codon positions from DNA datasets

(Sanderson et al. 2000), using site-heterogeneous models for amino acid data (Wang et al. 2018),

and identifying the conditions under which concatenation (Edwards 2009) or gene tree summary

approaches (Mirarab et al. 2014; Liu et al. 2015) more accurately resolve species relationships.

Discordance between gene and species trees can confound inferences of trait evolution as

well (Hahn and Nakhleh 2016), particularly for complex traits that appear to have evolved

convergently. Focus on the species tree alone, without considering discordant gene trees, can

lead to artifactual inferences of molecular convergence (Mendes et al. 2016). This failure occurs

when a trait is determined by genes with tree topologies that do not match the species topology, a

condition known as hemiplasy (Avise and Robinson 2008; Hahn and Nakhleh 2016; Storz 2016).

Hemiplasy has been identified as a likely explanation for patterns of character incongruence in

amino acid substitutions in columnar cacti (Copetti et al. 2017), flower and fruit traits in

Jaltomata (Solanaceae) (Wu et al. 2018), and dietary specialization in *Dysdera* spiders (Vizueta

et al. 2019). In conditions where decreased gene concordance is coupled with short branch

lengths and shallow time scales, hemiplasy is expected to have a higher impact on trait

reconstruction (Hahn and Nakhleh 2016).

Adaptation to low salinity is a complex trait, so the genomic changes associated with

successful freshwater colonizations are multifaceted (Artemov et al. 2017; Cabello-Yeves and

Rodriguez-Valera 2019; Rogers et al. 2021) and generally involve mutations in multiple genes

and pathways (Jones et al. 2012; Terekhanova et al. 2019; Chen et al. 2021). To better

understand the pattern, timing, and process of marine–freshwater transitions by diatoms, we

assembled a dataset of 45 genomes and 42 transcriptomes—most of them newly sequenced—to

resolve species relationships, explore the causes and consequences of gene tree discordance, and

provide insight into how discordance impacted trait reconstruction.

MATERIALS & METHODS

Detailed methods are provided in Supplementary File S1. Briefly, diatom cultures were

isolated from natural plankton or acquired from the National Center for Marine Algae and

Microbiota (NCMA) or Roscoff Culture Collection (RCC). Collection data, culture conditions,

and voucher information are available in Supplementary Table S1. For genome and

transcriptome sequencing, we extracted total DNA and RNA from diatom cultures, constructed

sequencing libraries, and sequenced them on the Illumina platform. Based on a large multigene

phylogeny of diatoms (Nakov et al. 2018), we included *Coscinodiscus,* Lithodesmiales, and

Eunotogramma as outgroups. Accession numbers for reads and assemblies are provided in

Supplementary Table S2.

We used OrthoFinder (Emms and Kelly 2019) to cluster amino acid sequences from all

genomes and transcriptomes into orthogroups, then aligned orthogroups containing ≥20% of the

taxa with MAFFT (Katoh and Standley 2013). For each alignment, we identified the best-fit

substitution model using ModelFinder (Kalyaanamoorthy et al. 2017) and estimated gene trees

with IQ-TREE (Minh et al. 2020b) or FastTree (Price et al. 2010). We then filtered and trimmed

the gene trees using the Rooted Ingroup method to produce final ortholog sets (Yang and Smith

2014). This filtering and trimming procedure was performed twice. We used PAL2NAL

(Suyama et al. 2006) to reconcile nucleotide coding sequence (CDS) alignments against amino

acid alignments. To account for possible saturation at third-codon positions, we removed them

from the CDS alignments and, separately, used Degen (Regier et al. 2010; Zwick et al. 2012) to

recode synonymous sites in the full coding sequence alignments with the nucleotide ambiguity

codes. For example, all leucine codons (CTN, TTR) were degenerated and replaced with YTN.

In total, we analyzed datasets consisting of amino acids (AA), first and second codon positions

(CDS12), and degenerate codons (DEGEN). We generated final ortholog alignments and

inferred trees as described above, using 1000 ultrafast bootstrap replicates to estimate branch

support (Minh et al. 2013). We generated summary statistics for final alignments and gene trees

with AMAS (Borowiec 2016) and PhyKit (Steenwyk et al. 2021) (Supplementary Table S3).

Correspondence analysis of amino acid frequencies across all taxa was performed using GCUA

(McInerney 1998).

Species trees were inferred using gene-partitioned maximum likelihood analysis of a

concatenated supermatrix with IQ-TREE and the summary quartet approach implemented in

ASTRAL-III (Zhang et al. 2018). We performed the matched-pairs test of symmetry in IQ-TREE

to identify and remove gene partitions that violated assumptions of stationarity, reversibility, and

homogeneity (SRH; Naser-Khdour et al. 2019). Briefly, stationarity refers to the assumption of

constant amino acid or nucleotide frequencies across time, reversibility implies that substitutions

between amino acids or nucleotides occur equally, and homogeneity implies that the

instantaneous substitution rates are constant along a tree or branch (Felsenstein 2004). For the

IQ-TREE analysis, we partitioned supermatrices by gene, used ModelFinder to select the best-fit

substitution model for each partition, and estimated branch support with 10,000 ultrafast

bootstrap replicates. For ASTRAL, we used the ortholog trees as input, collapsing branches with

low bootstrap support (<33) to help mitigate gene tree error (Sayyari and Mirarab 2016;

Simmons and Gatesy 2021). We estimated branch support using local posterior probability

(Sayyari and Mirarab 2016). In addition to heterogeneity in gene histories, compositional

heterogeneity can also make species tree inferences difficult at deep time scales (Lartillot and

Philippe 2004). To account for this possibility, we estimated a species tree from the concatenated

amino acid matrix using the Posterior Mean Site Frequency (PMSF) model implemented in IQ-

TREE (Wang et al. 2018).

We calculated the Robinson-Foulds distance between each pair of species trees and

visualized the results with a multidimensional scaling plot made with the R package *treespace*

(Jombart et al. 2017). We characterized discordance using gene and site concordance factors

(Minh et al. 2020a) and quartet concordance factors (Pease et al. 2018). We tested the support for

competing backbone topologies in our species tree using the approximately unbiased (AU) test

(Shimodaira 2002) implemented in IQ-TREE. Relative gene tree support for the same set of

backbone topologies was further evaluated using gene genealogy interrogation method (Arcila et

al. 2017). Finally, we performed the polytomy test on the ASTRAL species tree to test whether

any of the unstable backbone branches were better represented as a polytomy (Sayyari and

Mirarab 2018).

Divergence times were estimated using MCMCtree (Yang 2007; Reis and Yang 2011),

using five fossil calibrations (Supplemental File S1), autocorrelated rates, and the approximate

likelihood approach. We performed marginal ancestral state reconstruction for marine and

freshwater habitat across the species tree using hidden state speciation and extinction models in

the R package *hisse* (Beaulieu and O'Meara 2016). Lastly, we explored the potential for

hemiplasy in our habitat reconstructions by calculating: (1) hemiplasy risk factors using the R

package *pepo* (Guerrero and Hahn 2018); (2) the differences between the number of marine–

freshwater transitions optimized on unconstrained versus species-tree-constrained gene trees; and

(3) the number of alignment site changes that differed between unconstrained and species-tree-

constrained gene trees.

Phylogenetic trees were plotted using the R Bioconductor packages *ggtree* and *treeio* (Yu

et al. 2017; Wang et al. 2019). Assembled genomes and transcriptomes have been deposited at

NCBI under BioProject PRJNA825288. Supplementary Files, Figures, and Tables are available

from the Dryad Digital Repository: https://doi.org/10.5061/dryad.7m0cfxpxp. Voucher images,

proteomes, alignments, trees, log files, and code have been deposited in Zenodo:

https://doi.org/10.5281/zenodo.7713227.

RESULTS

We combined 42 newly sequenced draft genomes and 50 newly sequenced

transcriptomes with publicly available genome or transcriptome data for a final dataset of 87 taxa

representing 59 species. A total of 17 transcriptomes were used directly in phylogenomic

analyses, and the other 33 were used for genome annotation. From the combined dataset of

genomes and transcriptomes, we generated alignments and gene trees for 6262 orthologs, with

each taxon represented in an average of 3275 (52%) orthologs.

Compositional heterogeneity and dataset construction

Relative Composition Variability (Phillips and Penny 2003) indicated greater

compositional heterogeneity in third codon positions compared to amino acids and first and

second codon positions (Supplementary Fig. S1). We also found substantial variation in GC

content of third codon positions across taxa and genes (Supplementary Fig. S2), ranging from an

average proportion of 0.40 ± 0.07 in *Cyclotella kingstonii* to 0.76 ± 0.14 in *Shionodiscus*

oestrupii. In addition, nucleotide alignments including all three codon positions had significantly

higher levels of saturation compared to the amino acids (Supplementary Fig. S3). To minimize

potentially misleading signal in the nucleotide data, we removed third codon positions and, to

examine the effects of saturation and GC heterogeneity in coding sequences (CDS), we replaced

synonymous signal by recoding CDS sequences with degenerate codons (Zwick et al. 2012).

Based on these results, we created three datasets to estimate phylogenetic relationships: amino

acids (AA), first and second codon positions (CDS12), and degenerate codons (DEGEN).

Dataset and alignment characteristics are summarized in Table 1 and detailed in

Supplementary Table S3. Each dataset initially contained 6262 orthologs (Table 1). Gene trees

constructed from all datasets were well supported ($80 \pm 7\%$ average bootstrap; Table 1;

Supplementary Table S3). To reduce systematic errors due to model misspecification, we

removed orthologs that failed SRH assumptions ($P < 0.05$), which retained 5522 (AA), 3259

(CDS12), and 3788 (DEGEN) genes in the datasets (hereafter referred to as the "complete"

datasets; Table 1). We then subsetted the complete datasets to maximize phylogenetic signal or

minimize missing data. To maximize signal and reduce stochastic error, we sorted orthologs by

the percentage of parsimony informative (PI) sites and retained the top 25% ranked orthologs

("top-PI"; Table 1). To minimize the amount of missing data and maximize taxon occupancy, we

sorted complete datasets by the number of taxa and subsetted these to include the top 25%

ranked orthologs with highest taxon occupancy ("top-Taxa"; Table 1). Orthologs in the top-Taxa

datasets contained an average of $76 \pm 8\%$ of the total taxa.

Roberts et al.

205   Table 1. Summary of datasets used in the study.

Data type	Dataset name	# genes	Total sites	Parsimony informative sites (%)	Avg bootstrap support (%)	# genes passing SRH[1]	*Thalassiosira* grade topology	
						$P < 0.05$	IQ-TREE	ASTRAL
Amino acids (AA)	complete	6262	3,777,062	60	80	5522	Topology 4	Topology 1
	top-PI	1588	996,027	76	84	1314	Topology 4	Topology 1
	top-Taxa	1574	808,477	64	82	1374	Topology 5	Topology 1
	top-PI-top-Taxa + PMSF[2] model	488	246,300	77	85	488	Topology 5	Topology 1
1st + 2nd codon positions (CDS12)	complete	6262	7,554,124	55	80	3259	Topology 4	Topology 5
	top-PI	1570	1,950,228	71	82	661	Topology 4	Topology 3
	top-Taxa	1574	1,616,954	59	82	782	Topology 5	Topology 3
Degenerate codons (DEGEN)	complete	6262	11,311,186	33	81	3788	Topology 4	Topology 3
	top-PI	1569	2,884,728	44	83	771	Topology 4	Topology 2
	top-Taxa	1574	2,425,431	36	83	903	Topology 5	Topology 3

206   [1] Matched-pairs test of Symmetry, Reversibility, and Homogeneity.

207   [2] Posterior Mean Site Frequency model.

Species tree inference and placement of freshwater clades

We initially estimated 18 species trees using amino acids, codon positions 1 and 2, or

degenerate codon sequences, with different cutoffs for taxon occupancy or proportion of

parsimony informative sites and using both concatenation and summary quartet approaches

(Table 1). Correspondence analysis of amino acid frequencies separated taxa principally by

habitat (marine versus freshwater) rather than phylogeny (Supplementary Fig. S4), which led us

to explore whether compositional heterogeneity affected phylogenetic reconstructions. To do so,

we estimated an additional species tree using the PMSF mixture model, which can accommodate

heterogeneity in amino acid composition between species at a particular site (Wang et al. 2018).

Due to the computational demands of implementing this model, we applied it only to a reduced

amino acid dataset with orthologs that met both the top-PI and top-Taxa filtering criteria ("AA-

top-PI-top-Taxa"; Table 1).

Previous phylogenetic analyses of this group resolved freshwater taxa into two main

clades: the genus *Cyclotella*, which also includes several marine and brackish species, and the

"cyclostephanoids", comprised of several stenohaline genera confined exclusively to freshwaters

(Alverson et al. 2011). Given the potential implications for uncovering the mechanisms of

freshwater adaptation, we were primarily interested in the placements of these two clades. Gross

differences among data types and methods were evident in an ordination of species trees based

on pairwise Robinson-Foulds distances, which showed a clear separation between IQ-TREE and

ASTRAL topologies, with further separation of IQ-TREE trees estimated from datasets that

maximized signal (top-PI) or minimized missing data (top-Taxa) (Supplementary Fig. S5). The

phylogenetic position of *Cyclotella* was strongly supported and robust to differences in data type

(codons versus amino acids) and analysis (IQ-TREE versus ASTRAL) (Fig. 1). The

Roberts et al.

cyclostephanoids were placed consistently within a large clade of marine *Thalassiosira* and

relatives (Fig. 1), but the arrangements of five main subclades—*Thalassiosira* I–IV and the

freshwater cyclostephanoids—varied depending on data type and analysis (Table 1; Fig. 2a). We

refer to this part of the tree as the *Thalassiosira* grade.

One resolution of the *Thalassiosira* grade (topology 1) was recovered only by ASTRAL

analysis of amino acid gene trees and placed the freshwater cyclostephanoids as sister to a clade

of *Thalassiosira* I–IV (Table 1; Fig. 2a). All other species trees placed cyclostephanoids as sister

to *Thalassiosira* III (Table 1; Fig. 2a). ASTRAL analyses of codon-based gene trees (CDS12 and

DEGEN) alone recovered topology 3, which placed cyclostephanoids and *Thalassiosira* III as

sister to the remaining *Thalassiosira* (Table 1; Fig. 2a). Topologies 4 and 5 were recovered by

both data types, but only topology 5 was robust to both data type and analysis, having been

recovered by IQ-TREE analysis of amino acid and codon alignments, and ASTRAL analysis of

codon-based gene trees (Table 1; Fig. 2a). Moreover, topology 5 was also recovered by IQ-

TREE analysis with the PMSF model, with almost all branches in the *Thalassiosira* grade

receiving maximum support (Fig. 1, branches A–E). Notably, the PMSF analysis also recovered

a monophyletic *Stephanodiscus*, which matches expectations based on morphology (Theriot et

al. 1987). *Stephanodiscus* was paraphyletic in relation to *Cyclostephanos* in 18 of the 20 species

trees. Based on all of these results, we chose topology 5 from the PMSF analysis as the reference

species tree (Fig. 1).

Figure 1. Phylogram based on maximum likelihood analysis of amino acids using the posterior

mean site frequency (PMSF) model and a dataset of 488 loci with the highest proportions of taxa

and informative sites ("AA-top-PI-top-Taxa" dataset; Table 1). Backbone nodes of the

Thalassiosira grade are indicated by the letters A–E. All branches had bootstrap support (BS)

values of 100 except for those with black circles which had BS = 90. Pie charts on backbone

nodes show the proportion of gene trees that support the clade (gCF), the proportion that support

both discordant topologies (gDF1, gDF2), and the proportion that are discordant due to

paraphyly (gDFP). Size of the pie charts is for clarity only.

Discordance underlies topological uncertainty

Incongruence among genes trees and alignment sites is an important factor impacting

phylogenetic reconstruction (Degnan and Rosenberg 2006b; Mallet et al. 2016). We

characterized discordance by calculating gene, site, and quartet concordance factors for each

branch (Figs. 1 and 2a; Supplementary Figs. S6 and S7). Gene and site concordance factors (gCF

and sCF) represent the proportion of genes or sites that are in agreement with a particular branch

in the species tree (Minh et al. 2020a). Gene concordance factors range from 0 to 100, and site

concordance factors typically range from 33 to 100, with values near 33 indicative of no signal

for that branch (Minh et al. 2020a). Quartet concordance factors (QC) provide a likelihood-based

estimate of the relative support at each branch for the three possible resolutions of four taxa

(Pease et al. 2018). Quartet concordance factors range from -1 to 1, with positive values showing

support for the focal branch, negative values supportive for an alternate quartet, and values of

zero indicating equal support among the three possible quartets (Pease et al. 2018).

All three concordance factors were high for most branches in the tree, indicating that a

majority of genes, sites, and quartets supported those relationships (Supplementary Figs. S6 and

S7). Despite having maximum bootstrap and local posterior probability support, however,

concordance factors were low for many of the backbone branches (Fig. 1; Supplementary Figs.

S6 and S7), including ones within the *Thalassiosira* grade (Fig. 1, nodes A–E) that affected

placement of the freshwater cyclostephanoids (Fig. 2). In this part of the tree, concordance was

generally low for the backbone branches (gCF = 4–26; sCF = 32–42; QC = -0.04–0.36) (Figs. 1

and 2; Supplementary Figs. S6 and S7). Gene concordance factors were lowest for branches C

and D, the only two branches in the species tree with <100% bootstrap support (Fig. 1). Gene

concordance factors were only slightly higher for branches A, B, and E (Fig. 1), which had low

site concordance (Fig. 2a) and near-zero quartet concordance factors (Supplementary Fig. S7),

consistent with expectations for little to no signal. Site concordance factors did not change

appreciably with the CDS12 dataset. However, repeating quartet concordance factor calculations

with the CDS12 data revealed minor switches in support for branches C ($QC_C = 0.004 \rightarrow -0.006$)

and E ($QC_E = -0.04 \rightarrow 0.07$) (Supplementary Fig. S7). These patterns of shifting support, though

minor, combined with the lowest site concordance factor for branch E ($sCF_E = 32$) suggest that

very few sites support the sister relationship of *Thalassiosira* III and cyclostephanoids, despite its

recovery in 16 of the 20 inferred species trees (Figs. 1 and 2a).

For some branches on the species tree, there were more discordant than concordant genes

and sites (Fig. 1; Supplementary Figs. S6 and S7), suggesting that more genes and sites

supported an alternative relationship. This was the case with *Stephanodiscus*, for example, where

more genes supported paraphyly than monophyly, though more sites supported monophyly

(Supplementary Figs. S6 and S7). Across the tree, support for alternative relationships was

neither strong nor consistent among genes or sites. Illustrative of this, discordance in more than

one-third of the tree (29 of 83 branches) was due to paraphyly (Supplementary Fig. S6),

indicative of widespread lack of signal in gene trees. Taken together, these analyses highlighted

extensive gene and site discordance, some well-supported and much of it not, across

Thalassiosirales.

Roberts et al.

Figure 2. **a)** Phylogenetic hypotheses of the *Thalassiosira* grade inferred using concatenation and

summary methods on the amino acid (AA), codon positions 1 and 2 (CDS12), and recoded

codon (DEGEN) datasets. Nodes are labeled with the percentage of amino acid sites concordant

with the branch (site concordance factor). The principal clade of interest, the freshwater

cyclostephanoids, is colored orange and labeled 'FW'. The four focal clades of marine

Thalassiosira and allies are labeled I–IV. Panels b–d show results of gene genealogy

interrogation tests of alternative hypotheses of relationships within the *Thalassiosira* grade.

These tests used datasets filtered to include only the top 25% of orthologs based on the

percentage of parsimony informative sites (top-PI) for **b)** amino acids, **c)** codon positions 1 and

2, and **d)** recoded codons. Lines correspond to the cumulative number of genes (x axis)

supporting topology hypotheses with the highest probability and their *P* values (y axis) from the

Approximately Unbiased (AU) topology tests. Values above the dashed line indicate topological

hypotheses that are significantly better than the alternatives ($P < 0.05$). For example, the green

line in panel b shows that there were a total of 88 genes that best supported topology 3, though

only four of those genes were above the dotted line and were significantly better supported than

the other four alternative topologies.

There are both biological (e.g., ILS) and technical causes (e.g., gene tree error) of gene

discordance, but the proportion attributable to each factor can be difficult to discern (Morales-

Briones et al. 2020; Cai et al. 2021). To better assess the importance of gene tree error in our

dataset—whether genes with low phylogenetic signal produced inaccurate gene trees—we

recalculated gene and site concordance using just the 1588 amino acid orthologs with the highest

percentage of parsimony informative sites (top-PI dataset; Table 1). Average gene concordance

increased modestly, from 56.6% to 62.1%, but site concordance was unchanged (Supplemental

Fig. S8). Gene concordance factors for branches A, B, and E in the *Thalassiosira* grade increased

by 5–7% but were largely unchanged for branches C and D (Supplemental Fig. S8). These

increases in gene concordance when using the most signal-rich genes suggest that errors in gene

tree estimation contributed to the lack of resolution in several critical branches (Chan et al. 2020;

Vanderpool et al. 2020). Deeper nodes in the tree may be more prone to technical errors caused

by long-branch attraction, poor alignments, or model misspecification, despite our attempts to

minimize these during dataset construction (Supplementary File S1). We found slight negative

correlations between node age and both gene concordance ($R^2 = 0.10$, $P < 0.01$) and site

concordance ($R^2 = 0.26$, $P < 0.001$) (Supplementary Fig. S9), which suggests that older branches

more likely suffered from saturation due to recurrent substitutions.

Placement of the freshwater cyclostephanoids

We used two additional tree-based methods to assess the relative support for topologies

1–5 within the *Thalassiosira* grade (Fig. 2a). Like the original species tree inferences, results of

AU tests on concatenated alignments for each dataset in Table 1 largely reflected data type

(Table 1), with amino acid characters supporting topology 1 ($P = 0.01$, AU test) and the codon

and degenerate codon datasets supporting topology 4 ($P = 0$, AU test) (Supplementary Table S4).

We then performed gene genealogy interrogation (GGI) to look for secondary signal supporting

one or more of the five competing topologies (Arcila et al. 2017). To do this, we performed

constrained gene tree searches on the most information-rich (top-PI) orthologs and compared

their likelihoods using the AU test. The GGI test assumes monophyly of the tested clades, so

using our time-calibrated tree, we converted branch lengths (in millions of years) to coalescent

time units using a range of plausible effective population sizes and generation times for diatoms

(Supplementary File S1). The estimated stem branch lengths for the five clades in the

Thalassiosira grade were all >5 coalescent units, suggesting sufficient time to reach monophyly

(Rosenberg 2003). After ranking likelihood scores from the AU tests and selecting constraint

topologies with the best score (rank 1 trees), no single topology was strongly favored in a

majority of constrained gene trees across the three datasets, implying similar levels of support

(Fig. 2b–d; Supplementary Table S5). Support from the amino acid dataset was split between

topologies 1 and 4, which were recovered by both summary and concatenation methods (Table 1;

Fig. 2b; Supplementary Table S5). The most frequent best-fit topology for the codon and

degenerate codon datasets corresponded to topology 4 (Fig. 2c, d), which was originally

recovered by concatenation only (Table 1).

Gene genealogy interrogation can also be used to explore the effects of gene tree error on

summary quartet methods by filtering the input trees for ASTRAL to include only the highest

ranking constrained genes (Arcila et al. 2017; Mirarab 2017). For each of the three datasets, we

performed two ASTRAL analyses using as input either all the top scoring (rank 1) constrained

gene trees (n_{AA} = 1588, n_{CDS12} = 1570, n_{DEGEN} = 1569) or just the subset that had statistical

support ($P < 0.05$) above the AU-based rank 2 topology (n_{AA} = 142, n_{CDS12} = 56, n_{DEGEN} = 71). In

all six cases, the inferred trees were consistent with topology 5, despite it being best supported

(rank 1) in just 13–16% of the constrained gene trees (Fig. 2b–d; Supplementary Table S5). We

originally chose topology 5 as the reference species tree (Fig. 1) because it was recovered by

both amino acids and codons, ASTRAL and IQ-TREE analysis with the PMSF model, and

because it recovered monophyly of *Stephanodiscus*.

Coalescent theory predicts that in severe cases of ILS, short internal branches can

produce gene trees that conflict with the species tree more often than they agree, creating a so-

called "anomaly zone" (Degnan and Rosenberg 2006a). Our recovery of a species tree topology

that is not the most frequent among the GGI gene trees could indicate that the backbone of the

Thalassiosira grade lies in the anomaly zone. Despite this, polytomy tests in ASTRAL using

each dataset rejected the null hypothesis that any of these branches is a polytomy ($P < 0.05$).

The temporal sequence of marine–freshwater transitions

Divergence time estimates dated the crown Thalassiosirales to the upper Cretaceous,

around 113 Ma (95% CI: 96–120 Ma) (Fig. 3a; Supplementary Fig. S10). One of the two main

freshwater lineages, *Cyclotella*, originated in the late Cretaceous (95% CI: 66–86 Ma) and the

other freshwater lineage, the cyclostephanoids, originated later in the Eocene (95% CI: 36–48

Roberts et al.

Ma) (Fig. 3a; Supplementary Fig. S10). Radiation of the *Thalassiosira* grade lineages occurred

during the Paleocene, from 57 Ma (95% CI: 48–63 Ma) to 73 Ma (95% CI: 61–80 Ma) (Fig. 3a;

Supplementary Fig. S10). The overlapping confidence intervals allow for the possibility that

these lineages diverged in much more rapid succession than suggested by their mean ages.

Marine–freshwater transitions on gene versus species trees

We used HiSSE to estimate the number of independent marine–freshwater transitions on

the time-calibrated species tree (Fig. 3a). The best-fit HiSSE model was a character independent

model (CID-4; Supplementary Table S6), which indicates that shifts in diversification rate

occurred independently of marine–freshwater transitions. Using parameter estimates from the

CID-4 model, we inferred a total of six transitions from marine to freshwaters (Fig. 3a). Given

the large number of discordant gene trees (Fig. 1), we next sought to determine whether the

reconstructions were impacted by hemiplasy, i.e., marine–freshwater transitions that occurred on

branches in a gene tree not shared with the species tree. Narrowly defined, hemiplasy describes

discordance due to ILS (Avise and Robinson 2008), but we use the term hemiplasy broadly to

describe discordant gene trees with fewer overall habitat transitions than the species tree,

regardless of the underlying cause (see Hahn and Nakhleh 2016).

To assess the potential for hemiplasy in our trait reconstructions, we calculated hemiplasy

risk factors (HRF), which are the ratio of the probabilities of hemiplasy to homoplasy in different

parts of the species tree (Guerrero and Hahn 2018). HRFs were calculated using 12 different sets

of coalescent branch lengths that represent a range of scenarios with different effective

population size (N_e) and generation times (Supplementary Fig. S11). The risk of hemiplasy

ranged from low to nonexistent in nearly all scenarios involving realistic generation times for

diatoms (1 day, 7 days, and 1 month), whereas elevated hemiplasy risk was found only under

scenarios of unrealistically long generation times (6 months) (Supplementary Fig. S11).

We next assessed whether there was any empirical evidence for hemiplasy in the

alignments and gene trees. For each ortholog, we separately optimized, using parsimony, the

number of marine–freshwater transitions on the estimated gene tree (GT) and the gene tree

constrained to match the species tree topology (ST). We calculated the difference between the

number of transitions on the ST versus GT and found a small percentage of orthologs (13.7%)

with at least one instance of hemiplasy, broadly defined to include all instances with fewer

transitions on the gene versus species tree topology (GT < ST; Fig. 3b). This set of orthologs

likely includes topologies affected by hemiplasy due to ILS (Avise and Robinson 2008),

homoplasy (convergent or parallel substitutions), and gene tree error. Manual inspection of these

gene trees showed that many of them were paraphyletic and occurred between *Thalassiosira*

lacustris, *Conticribra weissflogii*, *C. nana*, and the remaining *Cyclotella* taxa, whereas a smaller

number were polyphyletic and involved *Cyclotella, Skeletonema potamos,* and the

cyclostephanoid clade (Supplemental Table S7). The remaining majority of orthologs either had

identical numbers of marine–freshwater transitions (GT = ST), matching the species tree, or had

more transitions on the gene tree (GT > ST), which we attribute to error (Fig. 3b).

To determine whether the set of putatively hemiplasious (GT < ST) genes might be

related to salinity adaptation, we compared them against a set of genes that were differentially

expressed in a low-salinity acclimation experiment using the marine/euryhaline diatom

Skeletonema marinoi (Pinseel et al. 2022). A total of 532 of the 658 rooted orthologs with GT <

ST were also present in the *S. marinoi* genome, and 203 of these (~38%) were differentially

expressed in response to salinity change (Supplementary Table S8). This included genes

Roberts et al.

involved in key pathways and functions important in acclimation to low salinity, such as

carotenoid biosynthesis, the xanthophyll cycle, light-harvesting activities, protein chaperones,

nitrogen assimilation, regulation of transcription and translation, and the metabolism of

polyamines, amino acids, and fatty acids (Supplementary Table S8).

For the same set of 658 orthologs, we then calculated the difference in the number of

amino acid transitions that occurred at each alignment site using the inferred GT versus

constrained ST topologies (Fig. 3c). Although agnostic to marine–freshwater transitions, this

approach measured the impact of gene tree discordance, irrespective of the underlying cause, to

overall molecular homoplasy. Using this approach, we found that an average of 18% of amino

acid sites within an ortholog had fewer transitions on the GT versus the ST (ST < GT; Fig. 3c).

Figure 3. **a)** Divergence times and ancestral state reconstruction of marine and freshwater habitat

in Thalassiosirales. Conspecific taxa were removed prior to ancestral state reconstruction,

leaving one tip per species. Pie charts denote the probability of each node reconstructed as either

marine (grey) or freshwater (blue) using parameters estimated from the HiSSE CID-4 model.

Euryhaline taxa were coded as marine for the purposes of ancestral state reconstruction.

Divergence times for the full set of taxa can be found in Supplementary Fig. S10. **b)** Summary of

the difference between the number of parsimony optimized marine–freshwater transitions on

rooted orthologs with the estimated gene tree topology (GT) versus the topology constrained to

match the species tree (ST). Bar colors denote whether a GT had greater (purple), equal (green),

or fewer (yellow) numbers of transitions relative to the ST. Numbers above bars are the number

of rooted orthologs in each bar. **c)** Summary of the percentage of aligned amino acid sites that

have greater (purple), equal (green), or fewer (yellow) numbers of state transitions on the GT

versus ST**.**

DISCUSSION

Transitions to freshwaters

Marine–freshwater transitions have been key events in the diversification of lineages

across the tree of life (Jamy et al. 2022), including diatoms where freshwater taxa have

experienced increased rates of both speciation and extinction compared to their marine ancestors

(Nakov et al. 2019). Our study focused on a model clade, the Thalassiosirales, which is among

the most abundant and diverse lineages in the marine and freshwater plankton and where genetic

and genomic resources are readily available (Armbrust et al. 2004; Nawaly et al. 2020; Roberts

et al. 2020). Data from the 42 genomes presented here greatly expand the phylogenetic and

Roberts et al.

ecological diversity of sequenced genomes for Thalassiosirales, and diatoms as a whole, and will

greatly facilitate efforts to identify the genomic basis of freshwater adaptation in diatoms.

We identified six marine–freshwater transitions in Thalassiosirales and tested whether

these transitions were fully independent, owing to separate parallel or convergent mutations

(homoplasy) or whether some of the transitions were attributable to hemiplasy, i.e., the sorting of

shared ancestral polymorphisms in discordant gene trees (Avise and Robinson 2008). Even in the

face of extensive gene tree discordance, the probability of hemiplasy influencing trait

reconstructions was virtually nonexistent across the backbone of the tree (Supplementary Fig.

S11). The risk of hemiplasy was only elevated under scenarios with unrealistically long

generation times for diatoms (Supplementary Fig. S11), many of which can undergo daily cell

divisions (Smayda 1969; Tuchman et al. 1984). The large effective population sizes of some

microbes, which increase the probability of hemiplasy, may be offset by their rapid generation

times, reducing the risk of hemiplasy and highlighting the need for better empirical estimates of

N_e and μ in groups like diatoms (Guerrero and Hahn 2018). The relatively few empirical studies

of hemiplasy so far have focused on vascular plants or animals with generation times that are

years or decades in length (Copetti et al. 2017; Wu et al. 2018; Vizueta et al. 2019). These

studies also focused on much younger lineages, making it more likely that shared ancestral

polymorphisms could still be detected.

Longer branches subtending separate clades with a shared trait make it more likely that

homoplasy, rather than hemiplasy, has occurred (Hahn and Nakhleh 2016). In our case, each of

the clades with freshwater taxa have sufficiently long coalescent branch lengths to have achieved

monophyly and have had multiple intervening speciation events leading to non-freshwater taxa

(Fig. 3a). Other properties of the habitat transitions were also indicative of homoplasy rather than

hemiplasy: ancestral state reconstructions were unambiguous, the six freshwater transitions were

not paraphyletic, and gene concordance was high on the long internal stem branches subtending

the two main freshwater lineages (Hahn and Nakhleh 2016; Wu et al. 2018).

Empirical analyses of gene trees and alignments revealed relatively small numbers of

genes and sites with putative hemiplasies (Fig. 3b,c). We identified approximately three-fold

fewer amino acid sites with potential hemiplasies than was found in a much younger (10 Ma)

radiation of saguaro cacti (Copetti et al. 2017). Given the length of time that has elapsed since

the two major freshwater transitions occurred, the probability that shared ancestral

polymorphisms persist decreases and the probability that lineage-specific adaptive mutations

arise increases (Suh et al. 2015; Zou and Zhang 2015; Mendes et al. 2016). A large number of

genes and pathways have been implicated in the response to low salinity (Nakov et al. 2020;

Downey et al. 2022; Pinseel et al. 2022), so an adaptive allele in one of the many possible target

genes might have made it possible simply to survive in freshwaters initially. In the tens of

millions of years since then, any hemiplasious alleles were most likely overwritten in the extant

freshwater descendents, leaving a shared ancestral phenotype (e.g., enhanced transport of sodium

ions) as the only remaining evidence of the hemiplasy. The divergent evolutionary outcomes also

suggest a greater role for homoplasy over hemiplasy. The genus *Cyclotella* includes freshwater,

secondarily marine, and generalist euryhaline species that can tolerate a wide range of salinities

(Guillard and Ryther 1962; Nakov et al. 2020; Downey et al. 2022). Similarly, most freshwater

transitions at the tips of the tree (Fig. 1) involve species with populations that also grow in

marine habitats (*Conticribra weissflogii* and *Cyclotella nana*) or can tolerate slightly brackish

water (*Thalassiosira lacustris* and *Skeletonema potamos*). The cyclostephanoids are, by contrast,

narrowly adapted stenohaline specialists found exclusively in freshwaters, suggestive of a

different genetic trajectory into freshwaters.

Although we have treated salinity as a categorical variable, salinity varies along a

continuum from freshwater to marine and even hypersaline. Moreover, salinity fluctuations are

common in brackish and marine systems, such as coastlines influenced by precipitation and river

discharge. Nevertheless, we found that a substantial set of the putatively hemiplasious genes

identified here are involved in acclimation to low salinity conditions in one of the species in our

analysis, *S. marinoi* (Pinseel et al. 2022), suggesting a possible role for hemiplasy in freshwater

adaptation. In addition, adaptation to low or changing salinity may be linked to modifications of

gene expression (Bussard et al. 2017; Nakov et al. 2020; Downey et al. 2022; Pinseel et al.

2022), codon usage (Prabha et al. 2017), nucleotide substitution rates (Mitterboeck et al. 2016),

transposable element activity (Yuan et al. 2018), epigenetic responses (Artemov et al. 2017), and

epistatic interactions (Stern et al. 2022). The genomic resources and phylogenetic framework

presented here represent an important advance towards identifying the genes and processes

underpinning freshwater adaptation by diatoms.

The impact of discordance on placement of freshwater clades

Comparative analyses require a strong phylogenetic framework (Felsenstein 1985), so a

major goal of this study was to establish a robust phylogenetic hypothesis for Thalassiosirales.

Our primarily genome-based analysis of 6262 nuclear orthologs provided better resolution and,

superficially, increased support across most of the tree compared to previous analyses based on a

few genes (Alverson et al. 2007). Across character types, methods of inference, and different

criteria for including characters or taxa, the placement of one of the two principal freshwater

clades, *Cyclotella*, was consistent across species trees, with an estimated origin in the late

Cretaceous. The placement of the second major freshwater lineage, the cyclostephanoids, was

less certain.

The cyclostephanoid clade was placed within a grade of marine species, most of which

belong to the polyphyletic genus *Thalassiosira*. These marine *Thalassiosira* were divided among

four clades, but the arrangement of these clades and the freshwater cyclostephanoids varied

across datasets and analyses. Uncertainty in the backbone relationships for this part of the tree

was likely caused by a combination of gene tree error and IL. First, many individual genes

contained too little information to confidently resolve deep splits separated by short branch

lengths—a finding that is not unique to this dataset (Chan et al. 2020; Arcila et al. 2021).

Divergence time estimates suggest that these splits occurred in relatively quick succession, as

few as 5 million years. Although many of these bipartitions had consistently weak support, gene

concordance factors increased when we analyzed orthologs with the highest phylogenetic signal,

implicating gene tree error as a source of instability (Chan et al. 2020; Vanderpool et al. 2020).

In other phylogenomic studies impacted by low phylogenetic signal and high gene tree error,

gene genealogy interrogation (GGI) has been used to identify majority gene support for a single

hypothesis (Hughes et al. 2018; Tea et al. 2021). In our case, no single resolution was supported

by a majority of genes, indicative of nodes that are difficult to resolve even with hundreds of

genes (Nesi et al. 2021). The best-supported constraint tree identified by GGI differed between

amino acid and codon-based gene trees, highlighting conflicting signal even within genes.

Second, in addition to gene tree error, the large number of alternative topologies among gene

trees along the *Thalassiosira* grade is also consistent with ILS (Arcila et al. 2017). Gene tree

summary methods such as ASTRAL outperform concatenation when ILS is the major source of

discordance (Kubatko and Degnan 2007; Roch and Warnow 2015), but summary methods

perform poorly when gene tree error is high (Roch and Warnow 2015; Xi et al. 2015). Following

Arcila et al. (2017), we attempted to eliminate some of the noise in our dataset by restricting

ASTRAL analyses to the top-ranked constrained gene trees and in doing so recovered a

backbone topology congruent with one of the few originally recovered by both summary and

concatenation methods (topology 5; Table 1 and Fig. 2). Taken together, these results suggest

that gene tree error negatively impacted our ASTRAL analyses. After identifying and removing

some of that error, we recovered a stronger hypothesis for the placement of freshwater

cyclostephanoids within the marine *Thalassiosira* grade.

The anomaly zone describes an especially vexing phylogenetic problem in which short

branch lengths are unresolvable, resulting in gene trees that differ from the species more

frequently than they agree (Degnan and Rosenberg 2006a). Within the *Thalassiosira* grade, the

most common GGI gene tree topologies either did not match the reference species tree or were

uninformative for these short branches. This implies that unresolved or weakly supported gene

trees are more probable than resolved ones in the *Thalassiosira* grade. Under ILS, branch lengths

that exceed the boundaries of the anomaly zone should produce resolved gene trees (Huang and

Knowles 2009). Gene tree error like that identified in our dataset can lead to underestimation of

coalescent branch lengths by summary methods like ASTRAL (Sayyari and Mirarab 2016;

Forthman et al. 2022). These coalescent branch lengths define the anomaly zone boundaries

(Degnan and Rosenberg 2006b; Linkem et al. 2016), and underestimates could result in mistaken

identification of an anomaly zone where none exists. The *Thalassiosira* grade appears to fall

outside of the anomaly zone because our coalescent branch length estimates exceed the

theoretical boundaries (Degnan and Rosenberg 2006b).

Factors that are more poorly known in diatoms might also have contributed to the high

levels of discordance. We attempted to minimize technical sources of gene tree error during

dataset construction (Supplemental File S1), but other sources are more difficult to discern (Cai

et al. 2021). These include hybridization and polyploidy, gene duplication and loss, or

recombination. Although hybridization does occur in diatoms (Casteleyn et al. 2009; Koester et

al. 2010; Tanaka et al. 2015; Çiftçi et al. 2022), it is not especially well studied. There is

evidence for an ancient allopolyploidy event early on in the evolutionary history of

Thalassiosirales (Parks et al. 2018). Many methods that identify hybridization based on gene

trees (Edelman et al. 2019; Vanderpool et al. 2020) or site patterns (Blischak et al. 2018) can be

confounded by ancestral population structure, substitutional saturation, or ghost lineages (Slatkin

and Pollack 2008; Tricou et al. 2022), all of which are poorly characterized in diatoms. High GC

content regions have been linked to higher recombination rates (Kent et al. 2012; Lartillot 2013)

and can lead to increased discordance (Pease and Hahn 2013). Despite high levels of variation in

GC content across Thalassiosirales, intralocus recombination was detected in a vanishingly small

(<0.001%) percent of the 6262 orthologs in our analysis (results available on Zenodo). In

addition to showing whether any of these factors have contributed to gene tree discordance, a

more thorough exploration of each one will fill important gaps in our understanding of diatom

evolution.

Codon bias and amino acid composition in freshwater diatoms

Thalassiosirales includes divergence times across a timescale ranging from thousands

(Theriot et al. 2006) to tens of millions of years ago (Fig. 3a), which led us to explore the utility

of both amino acid and nucleotide characters for resolving phylogenetic relationships. Amino

Roberts et al.

acids are less susceptible to saturation and useful for resolving deep relationships (Philippe et al.

2011; Rota-Stabelli et al. 2012), whereas nucleotides contain more information to resolve recent

divergences (Simmons et al. 2002; Townsend et al. 2008). Both data types recovered the vast

majority of relationships consistently and with strong support, while at the same time revealing

similar patterns of discordance along the backbone of the tree. In many cases, however, they

differed in their resolutions of the most recalcitrant parts of the tree. Different schemes to

mitigate the effects of saturation in the codon datasets also produced disagreements within the

Thalassiosira grade. Disagreements between character types within the same dataset, such as

those within the *Thalassiosira* grade here, have been found in other groups as well (Gillung et al.

2018; Skinner et al. 2020).

Almost every analysis of the amino acid dataset—including species trees, concordance

factors, and AU tests—supported the placement of cyclostephanoids as sister to the remaining

Thalassiosira clades, but nucleotide analyses placed them with *Thalassiosira* III (Fig. 2).

Discordance caused by codon usage bias and differences in amino acid composition might

account for this discrepancy. An association between codon bias and ecology has been

demonstrated in a broad diversity of microbes, where species that share an ecological niche have

similar codon usage, independent of phylogeny (Botzman and Margalit 2011; Roller et al. 2013;

Arella et al. 2021). Differences in codon usage between marine and freshwater prokaryotes have

been described (Cabello-Yeves and Rodriguez-Valera 2019), and we discovered differences in

both codon usage and amino acid composition between marine and freshwater diatoms. The

amino acid compositions of distantly related freshwater lineages might be sufficiently similar to

cause amino acid characters to support the "sister to the rest" placement of cyclostephanoids.

Protein sites with different structural, functional, or selective constraints can lead to differences

in amino acid composition between species (Villar and Kauvar 1994; Youssef et al. 2021),

something that is not accounted for by standard empirical protein models and may have led to

artifacts in our gene and species tree inferences (Wang et al. 2018). When we applied the PMSF

model, which accounts for compositional heterogeneity in amino acid sites, cyclostephanoids

were placed as sister to *Thalassiosira* III, in agreement with the codon datasets. The similarity in

codon usage and amino acid composition between distantly related freshwater diatoms merits

further study into the causes and functional significance, if any.

Conclusions

The vast differences between marine and freshwaters result in strong selective pressures

on freshwater colonists. Low salinity provokes a broad range of physiological and metabolic

responses in diatoms (Nakov et al. 2020; Downey et al. 2022; Pinseel et al. 2022), but the current

genetic architectures of freshwater adaptation reflect tens of millions of years of optimization

and change since the earliest transitions. As a result, it may be difficult to identify specific

alleles—whatever the process that generated them—that currently allow these diatoms to thrive

in freshwaters. Nevertheless, the phylogenomic analyses presented here suggest that convergent

or parallel substitutions likely played a more important role than hemiplasy in facilitating

freshwater colonizations in this group of diatoms. The vast new genomic resources and

phylogenetic framework presented here represent an important step forward in addressing these

types of questions to better understand how diatoms have made this complex ecological

transition appear to be so superficially simple.

Roberts et al.

SUPPLEMENTAL MATERIAL

Supplementary Files, Figures, and Tables are available from the Dryad Digital

Repository: https://doi.org/10.5061/dryad.7m0cfxpxp. Voucher images, proteomes, alignments,

trees, log files, and code have been deposited in Zenodo:

https://doi.org/10.5281/zenodo.7713227.

FUNDING

This work was supported by the National Science Foundation (DEB 1651087 to A.J.A)

and the Simons Foundation (725407 to E.P.). This research used resources available through the

Arkansas High Performance Computing Center, which is funded through multiple NSF grants

and the Arkansas Economic Development Commission.

ACKNOWLEDGEMENTS

We thank Ed Theriot for sharing several culture strains, and we thank Matt Ashworth and

Jeffery Stone for help with scanning electron microscopy.

LITERATURE CITED

Alverson A.J., Beszteri B., Julius M.L., Theriot E.C. 2011. The model marine diatom

Thalassiosira pseudonana likely descended from a freshwater ancestor in the genus

Cyclotella. BMC Evol. Biol. 11:125.

Alverson A.J., Jansen R.K., Theriot E.C. 2007. Bridging the Rubicon: Phylogenetic analysis

reveals repeated colonizations of marine and fresh waters by thalassiosiroid diatoms. Mol.

Phylogenet. Evol. 45:193–210.

Arcila D., Hughes L.C., Meléndez-Vazquez B., Baldwin C.C., White W.T., Carpenter K.E.,

Williams J.T., Santos M.D., Pogonoski J.J., Miya M., Ortí G., Betancur-R R. 2021. Testing

the Utility of Alternative Metrics of Branch Support to Address the Ancient Evolutionary

Radiation of Tunas, Stromateoids, and Allies (Teleostei: Pelagiaria). Syst. Biol. 70:1123–

1144.

Arcila D., Ortí G., Vari R., Armbruster J.W., Stiassny M.L.J., Ko K.D., Sabaj M.H., Lundberg J.,

Revell L.J., Betancur-R. R. 2017. Genome-wide interrogation advances resolution of

recalcitrant groups in the tree of life. Nature Ecology & Evolution. 1:1–10.

Arella D., Dilucca M., Giansanti A. 2021. Codon usage bias and environmental adaptation in

microbial organisms. Mol. Genet. Genomics. 296:751–762.

Armbrust E.V., Berges J.A., Bowler C., Green B.R., Martinez D., Putnam N.H., Zhou S., Allen

A.E., Apt K.E., Bechner M., Brzezinski M.A., Chaal B.K., Chiovitti A., Davis A.K.,

Demarest M.S., Detter J.C., Glavina T., Goodstein D., Hadi M.Z., Hellsten U., Hildebrand

M., Jenkins B.D., Jurka J., Kapitonov V.V., Kröger N., Lau W.W.Y., Lane T.W., Larimer

Roberts et al.

F.W., Lippmeier J.C., Lucas S., Medina M., Montsant A., Obornik M., Parker M.S., Palenik

B., Pazour G.J., Richardson P.M., Rynearson T.A., Saito M.A., Schwartz D.C.,

Thamatrakoln K., Valentin K., Vardi A., Wilkerson F.P., Rokhsar D.S. 2004. The genome

of the diatom *Thalassiosira pseudonana*: Ecology, evolution, and metabolism. Science.

306:79–86.

Artemov A.V., Mugue N.S., Rastorguev S.M., Zhenilo S., Mazur A.M., Tsygankova S.V.,

Boulygina E.S., Kaplun D., Nedoluzhko A.V., Medvedeva Y.A., Prokhortchouk E.B. 2017.

Genome-Wide DNA Methylation Profiling Reveals Epigenetic Adaptation of Stickleback to

Marine and Freshwater Conditions. Mol. Biol. Evol. 34:2203–2213.

Avise J.C., Robinson T.J. 2008. Hemiplasy: A new term in the lexicon of phylogenetics. Syst.

Biol. 57:503–507.

Beaulieu J.M., O'Meara B.C. 2016. Detecting Hidden Diversification Shifts in Models of Trait-

Dependent Speciation and Extinction. Syst. Biol. 65:583–601.

Blischak P.D., Chifman J., Wolfe A.D., Kubatko L.S. 2018. HyDe: A Python Package for

Genome-Scale Hybridization Detection. Syst. Biol. 67:821–829.

Borowiec M.L. 2016. AMAS: a fast tool for alignment manipulation and computing of summary

statistics. PeerJ. 4:e1660.

Botzman M., Margalit H. 2011. Variation in global codon usage bias among prokaryotic

organisms is associated with their lifestyles. Genome Biol. 12:R109.

Bussard A., Corre E., Hubas C., Duvernois-Berthet E., Le Corguillé G., Jourdren L., Coulpier F.,

Claquin P., Lopez P.J. 2017. Physiological adjustments and transcriptome reprogramming

are involved in the acclimation to salinity gradients in diatoms. Environ. Microbiol. 19:909–

925.

Cabello-Yeves P.J., Rodriguez-Valera F. 2019. Marine-freshwater prokaryotic transitions require

extensive changes in the predicted proteome. Microbiome. 7:117.

Cai L., Xi Z., Lemmon E.M., Lemmon A.R., Mast A., Buddenhagen C.E., Liu L., Davis C.C.

2021. The Perfect Storm: Gene Tree Estimation Error, Incomplete Lineage Sorting, and

Ancient Gene Flow Explain the Most Recalcitrant Ancient Angiosperm Clade,

Malpighiales. Syst. Biol. 70:491–507.

Casteleyn G., Adams N.G., Vanormelingen P., Debeer A.-E., Sabbe K., Vyverman W. 2009.

Natural hybrids in the marine diatom *Pseudo-nitzschia pungens* (Bacillariophyceae):

genetic and morphological evidence. Protist. 160:343–354.

Chan K.O., Hutter C.R., Wood P.L. Jr, Grismer L.L., Brown R.M. 2020. Target-capture

phylogenomics provide insights on gene and species tree discordances in Old World

treefrogs (Anura: Rhacophoridae). Proc. Biol. Sci. 287:20202102.

Chen M.-Y., Teng W.-K., Zhao L., Hu C.-X., Zhou Y.-K., Han B.-P., Song L.-R., Shu W.-S.

2021. Comparative genomics reveals insights into cyanobacterial evolution and habitat

adaptation. ISME J. 15:211–227.

Çiftçi O., Alverson A.J., van Bodegom P., Roberts W.R., Mertens A., Van de Vijver B., Trobajo

R., Mann D.G., Pirovano W., van Eijk I., Gravendeel B. 2022. Phylotranscriptomics reveals

the reticulate evolutionary history of a widespread diatom species complex. J. Phycol.

58:643–656.

Copetti D., Búrquez A., Bustamante E., Charboneau J.L.M., Childs K.L., Eguiarte L.E., Lee S.,

Liu T.L., McMahon M.M., Whiteman N.K., Wing R.A., Wojciechowski M.F., Sanderson

M.J. 2017. Extensive gene tree discordance and hemiplasy shaped the genomes of North

American columnar cacti. Proc. Natl. Acad. Sci. U. S. A. 114:12003–12008.

Degnan J.H., Rosenberg N.A. 2006a. Discordance of species trees with their most likely gene

trees. PLoS Genet. 2:e68.

Degnan J.H., Rosenberg N.A. 2006b. Discordance of species trees with their most likely gene

trees. PLoS Genet. 2:e68.

Dickson B., Yashayaev I., Meincke J., Turrell B., Dye S., Holfort J. 2002. Rapid freshening of

the deep North Atlantic Ocean over the past four decades. Nature. 416:832–837.

Dittami S.M., Heesch S., Olsen J.L., Collén J. 2017. Transitions between marine and freshwater

environments provide new clues about the origins of multicellular plants and algae. J.

Phycol. 53:731–745.

Downey K.M., Judy K.J., Pinseel E., Alverson A.J., Lewis J.A. 2022. The dynamic response to

hypoosmotic stress reveals distinct stages of freshwater acclimation by a euryhaline diatom.

Mol. Ecol.:2022.06.24.497401.

Edelman N.B., Frandsen P.B., Miyagi M., Clavijo B., Davey J., Dikow R.B., García-Accinelli

G., Van Belleghem S.M., Patterson N., Neafsey D.E., Challis R., Kumar S., Moreira G.R.P.,

Salazar C., Chouteau M., Counterman B.A., Papa R., Blaxter M., Reed R.D., Dasmahapatra

K.K., Kronforst M., Joron M., Jiggins C.D., McMillan W.O., Di Palma F., Blumberg A.J.,

Wakeley J., Jaffe D., Mallet J. 2019. Genomic architecture and introgression shape a

butterfly radiation. Science. 366:594–599.

Edwards S.V. 2009. Is a new and general theory of molecular systematics emerging? Evolution.

63:1–19.

Emms D.M., Kelly S. 2019. OrthoFinder: phylogenetic orthology inference for comparative

genomics. Genome Biol. 20:238.

Felsenstein J. 1985. Phylogenies and the Comparative Method. Am. Nat. 125:1–15.

Felsenstein J. 2004. Inferring phylogenies. Sinauer associates Sunderland, MA.

Forthman M., Braun E.L., Kimball R.T. 2022. Gene tree quality affects empirical coalescent

branch length estimation. Zool. Scr. 51:1–13.

Foster P.G. 2004. Modeling compositional heterogeneity. Syst. Biol. 53:485–495.

Gillung J.P., Winterton S.L., Bayless K.M., Khouri Z., Borowiec M.L., Yeates D., Kimsey L.S.,

Misof B., Shin S., Zhou X., Mayer C., Petersen M., Wiegmann B.M. 2018. Anchored

phylogenomics unravels the evolution of spider flies (Diptera, Acroceridae) and reveals

discordance between nucleotides and amino acids. Mol. Phylogenet. Evol. 128:233–245.

Guerrero R.F., Hahn M.W. 2018. Quantifying the risk of hemiplasy in phylogenetic inference.

Proc. Natl. Acad. Sci. U. S. A. 115:12787–12792.

Guillard R.R.L., Ryther J.H. 1962. Studies of marine planktonic diatoms: Guillard R. R. L. and J.

H. Ryther, 1962. I. *Cyclotella nana* Hustedt, and *Detonula confervacea* (Cleve) Gran. Can J

Microbiol. 8:229–240.

Hahn M.W., Nakhleh L. 2016. Irrational exuberance for resolved species trees. Evolution. 70:7–

17.

Huang H., Knowles L.L. 2009. What is the danger of the anomaly zone for empirical

phylogenetics? Syst. Biol. 58:527–536.

Hughes L.C., Ortí G., Huang Y., Sun Y., Baldwin C.C., Thompson A.W., Arcila D., Betancur-R

R., Li C., Becker L., Bellora N., Zhao X., Li X., Wang M., Fang C., Xie B., Zhou Z., Huang

H., Chen S., Venkatesh B., Shi Q. 2018. Comprehensive phylogeny of ray-finned fishes

(Actinopterygii) based on transcriptomic and genomic data. Proc. Natl. Acad. Sci. U. S. A.

115:6249–6254.

Jamy M., Biwer C., Vaulot D., Obiol A., Jing H., Peura S., Massana R., Burki F. 2022. Global

patterns and rates of habitat transitions across the eukaryotic tree of life. Nat Ecol Evol.

6:1458–1470.

Jombart T., Kendall M., Almagro-Garcia J., Colijn C. 2017. treespace: Statistical exploration of

landscapes of phylogenetic trees. Mol. Ecol. Resour. 17:1385–1392.

Jones F.C., Grabherr M.G., Chan Y.F., Russell P., Mauceli E., Johnson J., Swofford R., Pirun

M., Zody M.C., White S., Birney E., Searle S., Schmutz J., Grimwood J., Dickson M.C.,

Myers R.M., Miller C.T., Summers B.R., Knecht A.K., Brady S.D., Zhang H., Pollen A.A.,

Howes T., Amemiya C., Broad Institute Genome Sequencing Platform & Whole Genome

Assembly Team, Baldwin J., Bloom T., Jaffe D.B., Nicol R., Wilkinson J., Lander E.S., Di

Palma F., Lindblad-Toh K., Kingsley D.M. 2012. The genomic basis of adaptive evolution

in threespine sticklebacks. Nature. 484:55–61.

Kalyaanamoorthy S., Minh B.Q., Wong T.K.F., von Haeseler A., Jermiin L.S. 2017.

ModelFinder: fast model selection for accurate phylogenetic estimates. Nat. Methods.

14:587–589.

Katoh K., Standley D.M. 2013. MAFFT multiple sequence alignment software version 7:

improvements in performance and usability. Mol. Biol. Evol. 30:772–780.

Kenny N.J., Plese B., Riesgo A., Itskovich V.B. 2019. Symbiosis, Selection and Novelty:

Freshwater Adaptation in the Unique Sponges of Lake Baikal. Mol. Biol. Evol.

Kent C.F., Minaei S., Harpur B.A., Zayed A. 2012. Recombination is associated with the

evolution of genome structure and worker behavior in honey bees. Proc. Natl. Acad. Sci. U.

S. A. 109:18012–18017.

Kirst G.O. 1990. Salinity Tolerance of Eukaryotic Marine Algae. Annu. Rev. Plant Physiol. Plant

Mol. Biol. 41:21–53.

Kirst G.O. 1996. Osmotic Adjustment in Phytoplankton and MacroAlgae. In: Kiene R.P.,

Visscher P.T., Keller M.D., Kirst G.O., editors. Biological and Environmental Chemistry of

DMSP and Related Sulfonium Compounds. Boston, MA: Springer US. p. 121–129.

Koester J.A., Swalwell J.E., von Dassow P., Armbrust E.V. 2010. Genome size differentiates co-

occurring populations of the planktonic diatom *Ditylum brightwellii* (Bacillariophyta). BMC

Evol. Biol. 10:1.

Kubatko L.S., Degnan J.H. 2007. Inconsistency of phylogenetic estimates from concatenated

data under coalescence. Syst. Biol. 56:17–24.

Lartillot N. 2013. Phylogenetic patterns of GC-biased gene conversion in placental mammals and

the evolutionary dynamics of recombination landscapes. Mol. Biol. Evol. 30:489–502.

Lartillot N., Philippe H. 2004. A Bayesian mixture model for across-site heterogeneities in the

amino-acid replacement process. Mol. Biol. Evol. 21:1095–1109.

Lee C.E., Downey K., Colby R.S., Freire C.A., Nichols S., Burgess M.N., Judy K.J. 2022.

Recognizing salinity threats in the climate crisis. Integr. Comp. Biol.

Linkem C.W., Minin V.N., Leaché A.D. 2016. Detecting the Anomaly Zone in Species Trees

and Evidence for a Misleading Signal in Higher-Level Skink Phylogeny (Squamata:

Scincidae). Syst. Biol. 65:465–477.

Liu L., Wu S., Yu L. 2015. Coalescent methods for estimating species trees from phylogenomic

data. J. Syst. Evol. 53:380–390.

Lozupone C.A., Knight R. 2007. Global patterns in bacterial diversity. Proc. Natl. Acad. Sci. U.

S. A. 104:11436–11440.

Maddison W.P. 1997. Gene Trees in Species Trees. Syst. Biol. 46:523–536.

Mallet J., Besansky N., Hahn M.W. 2016. How reticulated are species? Bioessays. 38:140–149.

McCairns R.J.S., Bernatchez L. 2010. Adaptive divergence between freshwater and marine

sticklebacks: insights into the role of phenotypic plasticity from an integrated analysis of

candidate gene expression. Evolution. 64:1029–1047.

McInerney J.O. 1998. GCUA: general codon usage analysis. Bioinformatics. 14:372–373.

Mendes F.K., Hahn Y., Hahn M.W. 2016. Gene Tree Discordance Can Generate Patterns of

Diminishing Convergence over Time. Mol. Biol. Evol. 33:3299–3307.

Minh B.Q., Hahn M.W., Lanfear R. 2020a. New Methods to Calculate Concordance Factors for

Phylogenomic Datasets. Mol. Biol. Evol. 37:2727–2733.

Minh B.Q., Nguyen M.A.T., von Haeseler A. 2013. Ultrafast Approximation for Phylogenetic

Bootstrap. Mol. Biol. Evol. 30:1188–1195.

Minh B.Q., Schmidt H.A., Chernomor O., Schrempf D., Woodhams M.D., von Haeseler A.,

Lanfear R. 2020b. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic

Inference in the Genomic Era. Mol. Biol. Evol. 37:1530–1534.

Mirarab S. 2017. Phylogenomics: Constrained gene tree inference. Nat Ecol Evol. 1:56.

Mirarab S., Reaz R., Bayzid M.S., Zimmermann T., Swenson M.S., Warnow T. 2014. ASTRAL:

genome-scale coalescent-based species tree estimation. Bioinformatics. 30:i541–i548.

Mitterboeck T.F., Chen A.Y., Zaheer O.A., Ma E.Y.T., Adamowicz S.J. 2016. Do saline taxa

evolve faster? Comparing relative rates of molecular evolution between freshwater and

marine eukaryotes. Evolution. 70:1960–1978.

Molloy E.K., Warnow T. 2018. To Include or Not to Include: The Impact of Gene Filtering on

Species Tree Estimation Methods. Syst. Biol. 67:285–303.

Morales-Briones D.F., Kadereit G., Tefarikis D.T., Moore M.J., Smith S.A., Brockington S.F.,

Timoneda A., Yim W.C., Cushman J.C., Yang Y. 2020. Disentangling Sources of Gene

847    Tree Discordance in Phylogenomic Data Sets: Testing Ancient Hybridizations in

848    Amaranthaceae s.l. Syst. Biol. 70:219–235.

849  Nakov T., Beaulieu J.M., Alverson A.J. 2018. Accelerated diversification is related to life history

850    and locomotion in a hyperdiverse lineage of microbial eukaryotes (Diatoms,

851    Bacillariophyta). New Phytol. 219:462–473.

852  Nakov T., Beaulieu J.M., Alverson A.J. 2019. Diatoms diversify and turn over faster in

853    freshwater than marine environments. Evolution. 73:2497–2511.

854  Nakov T., Judy K.J., Downey K.M., Ruck E.C., Alverson A.J. 2020. Transcriptional Response of

855    Osmolyte Synthetic Pathways and Membrane Transporters in a Euryhaline Diatom During

856    Long-term Acclimation to a Salinity Gradient. J. Phycol. 56:1712–1728.

857  Naser-Khdour S., Minh B.Q., Zhang W., Stone E.A., Lanfear R. 2019. The Prevalence and

858    Impact of Model Violations in Phylogenetic Analysis. Genome Biol. Evol. 11:3341–3352.

859  Nawaly H., Tsuji Y., Matsuda Y. 2020. Rapid and precise genome editing in a marine diatom,

860    *Thalassiosira pseudonana* by Cas9 nickase (D10A). Algal Res. 47:101855.

861  Nesi N., Tsagkogeorga G., Tsang S.M., Nicolas V., Lalis A., Scanlon A.T., Riesle-Sbarbaro

862    S.A., Wiantoro S., Hitch A.T., Juste J., Pinzari C.A., Bonaccorso F.J., Todd C.M., Lim

863    B.K., Simmons N.B., McGowen M.R., Rossiter S.J. 2021. Interrogating Phylogenetic

864    Discordance Resolves Deep Splits in the Rapid Radiation of Old World Fruit Bats

865    (Chiroptera: Pteropodidae). Syst. Biol. 70:1077–1089.

866  Parks M.B., Nakov T., Ruck E.C., Wickett N.J., Alverson A.J. 2018. Phylogenomics reveals an

extensive history of genome duplication in diatoms (Bacillariophyta). Am. J. Bot. 105:330–

347.

Pease J.B., Brown J.W., Walker J.F., Hinchliff C.E., Smith S.A. 2018. Quartet Sampling

distinguishes lack of support from conflicting support in the green plant tree of life. Am. J.

Bot. 105:385–403.

Pease J.B., Haak D.C., Hahn M.W., Moyle L.C. 2016. Phylogenomics Reveals Three Sources of

Adaptive Variation during a Rapid Radiation. PLoS Biol. 14:e1002379.

Pease J.B., Hahn M.W. 2013. More accurate phylogenies inferred from low-recombination

regions in the presence of incomplete lineage sorting. Evolution. 67:2376–2384.

Philippe H., Brinkmann H., Lavrov D.V., Littlewood D.T.J., Manuel M., Wörheide G., Baurain

D. 2011. Resolving difficult phylogenetic questions: why more sequences are not enough.

PLoS Biol. 9:e1000602.

Phillips M.J., Penny D. 2003. The root of the mammalian tree inferred from whole mitochondrial

genomes. Mol. Phylogenet. Evol. 28:171–185.

Pinseel E., Nakov T., Van den Berge K., Downey K.M., Judy K.J., Kourtchenko O., Kremp A.,

Ruck E.C., Sjöqvist C., Töpel M., Godhe A., Alverson A.J. 2022. Strain-specific

transcriptional responses overshadow salinity effects in a marine diatom sampled along the

Baltic Sea salinity cline. ISME J.

Prabha R., Singh D.P., Sinha S., Ahmad K., Rai A. 2017. Genome-wide comparative analysis of

codon usage bias and codon context patterns among cyanobacterial genomes. Mar.

Genomics. 32:31–39.

Price M.N., Dehal P.S., Arkin A.P. 2010. FastTree 2--approximately maximum-likelihood trees

for large alignments. PLoS One. 5:e9490.

Regier J.C., Shultz J.W., Zwick A., Hussey A., Ball B., Wetzer R., Martin J.W., Cunningham

C.W. 2010. Arthropod relationships revealed by phylogenomic analysis of nuclear protein-

coding sequences. Nature. 463:1079–1083.

Reis M. dos, Yang Z. 2011. Approximate Likelihood Calculation on a Phylogeny for Bayesian

Estimation of Divergence Times. Mol. Biol. Evol. 28:2161–2172.

Roberts W.R., Downey K.M., Ruck E.C., Traller J.C., Alverson A.J. 2020. Improved Reference

Genome for *Cyclotella cryptica* CCMP332, a Model for Cell Wall Morphogenesis, Salinity

Adaptation, and Lipid Production in Diatoms (Bacillariophyta). G3 . 10:2965–2974.

Roch S., Warnow T. 2015. On the Robustness to Gene Tree Estimation Error (or lack thereof) of

Coalescent-Based Species Tree Methods. Syst. Biol. 64:663–676.

Rogers R.L., Grizzard S.L., Titus-McQuillan J.E., Bockrath K., Patel S., Wares J.P., Garner J.T.,

Moore C.C. 2021. Gene family amplification facilitates adaptation in freshwater unionid

bivalve *Megalonaias nervosa*. Mol. Ecol. 30:1155–1173.

Roller M., Lucić V., Nagy I., Perica T., Vlahovicek K. 2013. Environmental shaping of codon

usage and functional adaptation across microbial communities. Nucleic Acids Res.

41:8842–8852.

Rosenberg N.A. 2003. The shapes of neutral gene genealogies in two species: probabilities of

monophyly, paraphyly, and polyphyly in a coalescent model. Evolution. 57:1465–1477.

Rota-Stabelli O., Lartillot N., Philippe H., Pisani D. 2012. Serine Codon-Usage Bias in Deep

Phylogenomics: Pancrustacean Relationships as a Case Study. Syst. Biol. 62:121–133.

Sanderson M.J., Wojciechowski M.F., Hu J.M., Khan T.S., Brady S.G. 2000. Error, bias, and

long-branch attraction in data for two chloroplast photosystem genes in seed plants. Mol.

Biol. Evol. 17:782–797.

Sayyari E., Mirarab S. 2016. Fast Coalescent-Based Computation of Local Branch Support from

Quartet Frequencies. Mol. Biol. Evol. 33:1654–1668.

Sayyari E., Mirarab S. 2018. Testing for Polytomies in Phylogenetic Species Trees Using

Quartet Frequencies. Genes . 9.

Shimodaira H. 2002. An approximately unbiased test of phylogenetic tree selection. Syst. Biol.

51:492–508.

Simmons M.P., Gatesy J. 2021. Collapsing dubiously resolved gene-tree branches in

phylogenomic coalescent analyses. Mol. Phylogenet. Evol. 158:107092.

Simmons M.P., Ochoterena H., Freudenstein J.V. 2002. Amino acid vs. nucleotide characters:

challenging preconceived notions. Mol. Phylogenet. Evol. 24:78–90.

Skinner R.K., Dietrich C.H., Walden K.K.O., Gordon E., Sweet A.D., Podsiadlowski L.,

Petersen M., Simon C., Takiya D.M., Johnson K.P. 2020. Phylogenomics of

Auchenorrhyncha (Insecta: Hemiptera) using transcriptomes: examining controversial

relationships via degeneracy coding and interrogation of gene conflict. Syst. Entomol.

45:85–113.

Slatkin M., Pollack J.L. 2008. Subdivision in an ancestral species creates asymmetry in gene

trees. Mol. Biol. Evol. 25:2241–2246.

Smayda T.J. 1969. Experimental observations on the influence of temperature, light, and salinity

on cell division of the marine diatom, *Detonula confervacea* (Cleve) Gran. J. Phycol.

5:150–157.

Smith S.A., Brown J.W., Walker J.F. 2018. So many genes, so little time: A practical approach

to divergence-time estimation in the genomic era. PLoS One. 13:e0197433.

Steenwyk J.L., Buida T.J., Labella A.L., Li Y., Shen X.-X., Rokas A. 2021. PhyKIT: a broadly

applicable UNIX shell toolkit for processing and analyzing phylogenomic data.

Bioinformatics.

Stern D.B., Anderson N.W., Diaz J.A., Lee C.E. 2022. Genome-wide signatures of synergistic

epistasis during parallel adaptation in a Baltic Sea copepod. Nat. Commun. 13:4024.

Storz J.F. 2016. Causes of molecular convergence and parallelism in protein evolution. Nat. Rev.

Genet. 17:239–250.

Suh A., Smeds L., Ellegren H. 2015. The Dynamics of Incomplete Lineage Sorting across the

Ancient Adaptive Radiation of Neoavian Birds. PLoS Biol. 13:e1002224.

Suyama M., Torrents D., Bork P. 2006. PAL2NAL: robust conversion of protein sequence

alignments into the corresponding codon alignments. Nucleic Acids Res. 34:W609–W612.

Tanaka T., Maeda Y., Veluchamy A., Tanaka M., Abida H., Maréchal E., Bowler C., Muto M.,

Sunaga Y., Tanaka M., Yoshino T., Taniguchi T., Fukuda Y., Nemoto M., Matsumoto M.,

Wong P.S., Aburatani S., Fujibuchi W. 2015. Oil accumulation by the oleaginous diatom

Fistulifera solaris as revealed by the genome and transcriptome. Plant Cell. 27:162–176.

Tea Y.-K., Xu X., DiBattista J.D., Lo N., Cowman P.F., Ho S.Y.W. 2021. Phylogenomic

Analysis of Concatenated Ultraconserved Elements Reveals the Recent Evolutionary

Radiation of the Fairy Wrasses (Teleostei: Labridae: Cirrhilabrus). Syst. Biol. 71:1–12.

Terekhanova N.V., Barmintseva A.E., Kondrashov A.S., Bazykin G.A., Mugue N.S. 2019.

Architecture of Parallel Adaptation in Ten Lacustrine Threespine Stickleback Populations

from the White Sea Area. Genome Biol. Evol. 11:2605–2618.

Theriot E.C., Fritz S.C., Whitlock C., Conley D.J. 2006. Late Quaternary rapid morphological

evolution of an endemic diatom in Yellowstone Lake, Wyoming. Paleobiology. 32:38–54.

Theriot E., Stoermer E., Håkansson H. 1987. Taxonomic interpretation of the rimoportula of

freshwater genera in the centric diatom family Thalassiosiraceae. Diatom Res. 2:251–265.

Townsend J.P., López-Giráldez F., Friedman R. 2008. The phylogenetic informativeness of

nucleotide and amino acid sequences for reconstructing the vertebrate tree. J. Mol. Evol.

67:437–447.

Tricou T., Tannier E., de Vienne D.M. 2022. Ghost Lineages Highly Influence the Interpretation

of Introgression Tests. Syst. Biol.

Tuchman M.L., Theriot E., Stoermer E.F. 1984. Effects of low level salinity concentrations on

the growth of *Cyclotella meneghiniana* Kütz. (Bacillariophyta). Archiv für Protistenkunde.

128:319–326.

Vanderpool D., Minh B.Q., Lanfear R., Hughes D., Murali S., Harris R.A., Raveendran M.,

Muzny D.M., Hibbins M.S., Williamson R.J., Gibbs R.A., Worley K.C., Rogers J., Hahn

M.W. 2020. Primate phylogenomics uncovers multiple rapid radiations and ancient

interspecific introgression. PLoS Biol. 18:e3000954.

Villar H.O., Kauvar L.M. 1994. Amino acid preferences at protein binding sites. FEBS Lett.

349:125–130.

Vizueta J., Macías-Hernández N., Arnedo M.A., Rozas J., Sánchez-Gracia A. 2019. Chance and

predictability in evolution: The genomic basis of convergent dietary specializations in an

adaptive radiation. Mol. Ecol. 28:4028–4045.

Wang H.-C., Minh B.Q., Susko E., Roger A.J. 2018. Modeling Site Heterogeneity with Posterior

Mean Site Frequency Profiles Accelerates Accurate Phylogenomic Estimation. Syst. Biol.

67:216–235.

Wang L.-G., Lam T.T.-Y., Xu S., Dai Z., Zhou L., Feng T., Guo P., Dunn C.W., Jones B.R.,

Bradley T., Zhu H., Guan Y., Jiang Y., Yu G. 2019. Treeio: An R Package for Phylogenetic

Tree Input and Output with Richly Annotated and Associated Data. Mol. Biol. Evol.

37:599–603.

Wu M., Kostyun J.L., Hahn M.W., Moyle L.C. 2018. Dissecting the basis of novel trait evolution

in a radiation with widespread phylogenetic discordance. Mol. Ecol.

Xi Z., Liu L., Davis C.C. 2015. Genes with minimal phylogenetic information are problematic

987 for coalescent analyses when gene tree estimation is biased. Mol. Phylogenet. Evol. 92:63–

988 71.

989 Yang Y., Smith S.A. 2014. Orthology inference in nonmodel organisms using transcriptomes

990 and low-coverage genomes: improving accuracy and matrix occupancy for phylogenomics.

991 Mol. Biol. Evol. 31:3081–3092.

992 Yang Z. 2007. PAML 4: phylogenetic analysis by maximum likelihood. Mol. Biol. Evol.

993 24:1586–1591.

994 Youssef N., Susko E., Roger A.J., Bielawski J.P. 2021. Shifts in amino acid preferences as

995 proteins evolve: A synthesis of experimental and theoretical work. Protein Sci. 30:2009–

996 2028.

997 Yuan Z., Liu S., Zhou T., Tian C., Bao L., Dunham R., Liu Z. 2018. Comparative genome

998 analysis of 52 fish species suggests differential associations of repetitive elements with their

999 living aquatic environments. BMC Genomics. 19:141.

1000 Yu G., Smith D.K., Zhu H., Guan Y., Lam T.T.-Y. 2017. Ggtree : An r package for visualization

1001 and annotation of phylogenetic trees with their covariates and other associated data.

1002 Methods Ecol. Evol. 8:28–36.

1003 Zhang C., Rabiee M., Sayyari E., Mirarab S. 2018. ASTRAL-III: polynomial time species tree

1004 reconstruction from partially resolved gene trees. BMC Bioinformatics. 19:153.

1005 Zou Z., Zhang J. 2015. Are Convergent and Parallel Amino Acid Substitutions in Protein

1006 Evolution More Prevalent Than Neutral Expectations? Mol. Biol. Evol. 32:2085–2096.

Zwick A., Regier J.C., Zwickl D.J. 2012. Resolving discrepancy between nucleotides and amino

acids in deep-level arthropod phylogenomics: differentiating serine codons in 21-amino-

acid models. PLoS One. 7:e47450.

[dataset]* Roberts W.R., Ruck E.C., Downey K.M., Alverson A.J. 2022. Resolving marine–

freshwater transitions by diatoms through a fog of discordance and hemiplasy. Dryad.

http://dx.doi.org/10.5061/dryad.[NNNN].

FIGURE LEGENDS

Figure 1. Phylogram based on maximum likelihood analysis of amino acids using the posterior

mean site frequency (PMSF) model and a dataset of 488 loci with the highest proportions of taxa

and informative sites ("AA-top-PI-top-Taxa" dataset; Table 1). Backbone nodes of the

Thalassiosira grade are indicated by the letters A–E. All branches had bootstrap support (BS)

values of 100 except for those with black circles which had BS = 90. Pie charts on backbone

nodes show the proportion of gene trees that support the clade (gCF), the proportion that support

both discordant topologies (gDF1, gDF2), and the proportion that are discordant due to

paraphyly (gDFP). Size of the pie charts is for clarity only.

1022

Figure 2. **a)** Phylogenetic hypotheses of the *Thalassiosira* grade inferred using concatenation and

summary methods on the amino acid (AA), codon positions 1 and 2 (CDS12), and recoded

codon (DEGEN) datasets. Nodes are labeled with the percentage of amino acid sites concordant

with the branch (site concordance factor). The principal clade of interest, the freshwater

cyclostephanoids, is colored orange and labeled 'FW'. The four focal clades of marine

Thalassiosira and allies are labeled I–IV. Panels b–d show results of gene genealogy

interrogation tests of alternative hypotheses of relationships within the *Thalassiosira* grade.

These tests used datasets filtered to include only the top 25% of orthologs based on the

percentage of parsimony informative sites (top-PI) for **b)** amino acids, **c)** codon positions 1 and

2, and **d)** recoded codons. Lines correspond to the cumulative number of genes (x axis)

supporting topology hypotheses with the highest probability and their *P* values (y axis) from the

Approximately Unbiased (AU) topology tests. Values above the dashed line indicate topological

hypotheses that are significantly better than the alternatives ($P < 0.05$). For example, the green

line in panel b shows that there were a total of 88 genes that best supported topology 3, though

only four of those genes were above the dotted line and were significantly better supported than

the other four alternative topologies.

1039

Figure 3. **a)** Divergence times and ancestral state reconstruction of marine and freshwater habitat

in Thalassiosirales. Conspecific taxa were removed prior to ancestral state reconstruction,

leaving one tip per species. Pie charts denote the probability of each node reconstructed as either

marine (grey) or freshwater (blue) using parameters estimated from the HiSSE CID-4 model.

Euryhaline taxa were coded as marine for the purposes of ancestral state reconstruction.

Divergence times for the full set of taxa can be found in Supplementary Fig. S10. **b)** Summary of

the difference between the number of parsimony optimized marine–freshwater transitions on

rooted orthologs with the estimated gene tree topology (GT) versus the topology constrained to

match the species tree (ST). Bar colors denote whether a GT had greater (purple), equal (green),

or fewer (yellow) numbers of transitions relative to the ST. Numbers above bars are the number

of rooted orthologs in each bar. **c)** Summary of the percentage of aligned amino acid sites that

have greater (purple), equal (green), or fewer (yellow) numbers of state transitions on the GT

versus ST**.**