***Drosophila* pVALIUM10 TRiP RNAi lines cause undesired silencing of Gateway-based transgenes**

Dimitrije Stanković[1], Gábor Csordás[2], Mirka Uhlirova[1#]

[1]Institute for Genetics and Cologne Excellence Cluster on Cellular Stress Responses in Aging-Associated

Diseases (CECAD), University of Cologne, Cologne, Germany

[2]Institute of Genetics, Biological Research Centre of the Eötvös Loránd Research Network, Szeged,

Hungary

[#] Address correspondence to Mirka Uhlirova, mirka.uhlirova@uni-koeln.de

Running title: pVALIUM10 TRiP lines silence Gateway-based transgenes

Keywords: *Drosophila*, RNAi, dsRNA, pValium10, Gateway recombination, TRiP lines

SUMMARY

Post-transcriptional gene silencing using double-stranded RNA has revolutionized the field of functional

genetics, allowing fast and easy disruption of gene function in various organisms. In *Drosophila*, many

transgenic RNAi lines have been generated in large-scale efforts, including the *Drosophila* Transgenic RNAi

Project (TRiP), to facilitate *in vivo* knockdown of virtually any *Drosophila* gene with spatial and temporal

resolution. The available transgenic RNAi lines represent a fundamental resource for the fly community,

providing an unprecedented opportunity to address a vast range of biological questions relevant to basic

and biomedical research fields. However, caution should be applied regarding the efficiency and

specificity of the RNAi approach. Here, we demonstrate that pVALIUM10-based RNAi lines, representing

~13% of the total TRiP collection (1,808 out of 13,410 TRiP pVALIUM-based RNAi lines), cause unintended

off-target silencing of transgenes expressed from Gateway destination vectors generated via site-specific

recombination. The silencing is mediated by targeting attB1 and attB2 sequences generated in the

recombination reaction and included in the transcribed mRNA. Deleting these attB sites from the Gateway

expression vector prevents silencing and restores expected transgene expression.

INTRODUCTION

Double-stranded RNA (dsRNA)-mediated post-transcriptional gene silencing, also known as RNA

interference (RNAi), represents an ancient anti-viral defense mechanism that has been harnessed as a

powerful tool for reverse functional genetics in various cultured cells, plant and animal model systems

(Kennerdell & Carthew, 1998; Lohmann *et al*, 1999; Wargelius *et al*, 1999; Wianny & Zernicka-Goetz, 2000;

Zamore *et al*, 2000). The dsRNA silencing relies on a highly conserved RNase III enzyme called Dicer that

recognizes and processes dsRNA into small 21-23-bp segments (siRNA). Upon loading into a multiprotein

RNA-induced silencing complex (RISC), the guide strand facilitates pairing with complementary on-target

RNAs, which are subsequently degraded by the action of the Argonaut proteins (Tomari & Zamore, 2005).

In *Drosophila*, spatio-temporally controlled RNAi knockdown can be achieved by expressing dsRNA using

transgenic binary expression systems, such as Gal4/UAS (Brand & Perrimon, 1993). The initial transgenic

RNAi lines were generated using plasmids with ligated inverted repeats transcribed from one (Kennerdell

& Carthew, 1998) or two opposite promoters (Giordano *et al*, 2002). However, these vectors were

unstable and not well tolerated by the bacterial strains commonly used for cloning (Hagan & Warren,

1983; Piccin *et al*, 2001), a problem later solved by the introduction of an intron spacer between the

repeats (Bao & Cagan, 2006; Lee & Carthew, 2003). This step-by-step progress led to the development of

the pVALIUM (Vermilion-attB-LoxP-Intron-UAS-MCS), a universal vector system for transgenic RNAi library

generation. The pVALIUM RNAi vectors contain an eye-color selection marker (*vermillion*), an attB site

enabling phiC31-mediated site-directed integration into predefined genomic locations, *white* or *ftz*

introns promoting the processing of dsRNA, and a cassette of two UAS pentamers, one of which is flanked

by loxP sites to allow tuning of the dsRNA expression levels, as well as a multiple cloning site (MCS) (Ni *et*

al, 2008).

The highly efficient pVALIUM10-based transgenic RNAi stocks produced within the *Drosophila* Transgenic

RNAi Project (TRiP) at Harvard Medical School (HMS) (https://fgr.hms.harvard.edu) represent an

invaluable resource for the *Drosophila* community (Perkins *et al*, 2015). At the moment of writing, 1,808

transgenic TRiP RNAi stocks based on the pVALIUM10 are available at the Bloomington *Drosophila* stock

center (BDSC, https://bdsc.indiana.edu/index.html) (Perrimon *et al*, 2010). The pVALIUM10 vector carries

gypsy insulators that markedly enhance knockdown efficiency, along with two inverted "attR1-selection-

attR2" cassettes that facilitate cloning of dsRNA fragments via site-specific Gateway *in vitro*

recombineering (Ni *et al*, 2009; Reece-Hoyes & Walhout, 2018). A desired single fragment, 400 - 600 bp

in length and flanked by attL1 and attL2 recombination sites, can be easily moved from an entry vector

into a pVALIUM10 destination vector via recombination between attL and attR sites catalyzed by LR

clonase, producing a vector for UAS-driven expression of dsRNA (Ni *et al.*, 2009).

The Gateway cloning technology is also routinely used to generate transgenic constructs for expression of

epitope-tagged or untagged proteins, taking advantage of the *Drosophila* Gateway™ plasmid collection at

the *Drosophila* Genomics Resource Center (https://dgrc.bio.indiana.edu/). Importantly, as with

pVALIUM10 dsRNA plasmids, production of Gateway expression vectors relies on the recombination

between attL and attR sites resulting in the formation of attB1 and attB2 sites.

Here, we demonstrate that the pVALIUM10-derived RNAi lines cause undesirable knockdown of

transgenic reporters and overexpression lines generated with the help of the Gateway cloning technology.

They do so by targeting the attB sites that are part of the transcribed mRNA. Deletion of the attB1 and

attB2 sequences from the Gateway expression destination vector abrogates this effect and restores

expected transgene expression levels.

RESULTS

Unexpected, off-target silencing of the RNAse H1::GFP reporter by multiple pVALIUM10-RNAi lines

In vivo RNAi-based genetic screens have proved very successful in identifying novel genes involved in

developmental and pathologic processes in a variety of *Drosophila* somatic tissues when expressed using

tissue-specific Gal4 driver lines (Graca *et al*, 2021; Lesch *et al*, 2010; Neely *et al*, 2010; Pletcher *et al*, 2019;

Port *et al*, 2011; Rotelli *et al*, 2019; Rylee *et al*, 2022; Saj *et al*, 2010; Zeng *et al*, 2015; Zhou *et al*, 2019).

Hence, we embarked on a candidate genetic screen aimed at identifying factors controlling genome

stability. During the pre-screening of potential candidates we observed that expression of pVALIUM10-

based *BuGZ[RNAi[TRiP.JF02830]]* RNAi line in the pouch region of the developing third instar wing discs, using the

nubbin-Gal4, UAS-mRFP (*nub>mRFP*) driver line, efficiently suppressed the multi-epitope tagged BuGZ

transgene (*BuGZ^FlyFos[v318366]*) (Sarov *et al*, 2006) (Figure 1A, B, D). A similar reduction was observed using

an independent *BuGZ^RNAi[KK104498]* line from the Vienna *Drosophila* Resource Center (VDRC) KK RNAi library

(Figure 1C, D). Encouraged by these results, both BuGZ RNAi lines were included in the candidate genetic

screen that was based on expression of selected dsRNAs with the *nub>mRFP* driver. As readout for

genome stability, we used the ubiquitously driven RNase H1::GFP that was generated by recombining the

RNase H1 coding sequence into the pUWG vector from the *Drosophila* Gateway™ plasmid collection. This

resulted in the expression of a C-terminally GFP-tagged RNase H1 under the control of the poly-ubiquitin

promoter (Figure 2A, B). The RNase H1 enzyme recognizes and resolves R-loops, DNA:RNA hybrids, that

emerge during transcription and represent a threat to genome stability when unresolved (Crossley *et al*,

2019; Santos-Pereira & Aguilera, 2015). Unexpectedly, we observed reduction of RNase H1::GFP signal

when expressing *BuGZ^RNAi[TRiP.JF02830]* but not *BuGZ^RNAi[KK104498]* line (Figure 2C, D, H). Strikingly, similar

downregulation of RNase H1::GFP was caused by pVALIUM10-based *myc^RNAi[TRiP.JF01761]* but not

myc^RNAi[TRiP.HMS01538] or *myc^RNAi[GD2948]* derived from pVALIUM20 and pMF3 vectors, respectively (Figures 1D

and 2E-H). All three myc-RNAi lines have been shown effective by others (Aughey *et al*, 2016; Kim-Yip &

Nystul, 2018; Ledru *et al*, 2022; Rust *et al*, 2018). These results indicated that the tested pVALIUM10 TRiP

RNAi lines but not pVALIUM20, KK or GD lines may cause undesirable silencing of the RNase H1::GFP

transgene. We further speculated that dsRNA expressed from pVALIUM10 contains segments unrelated

to the target gene-specific sequences that trigger this unexpected off-target effect.

pVALIUM10 RNAi and Gateway expression plasmids share attB recombination sites

To identify the potential target for dsRNA-mediated knockdown of the RNAse H1::GFP transgene by

pVALIUM10 TRiP RNAi, we compared the plasmid sequences. We found that attB1 and attB2 sites are

shared between the Gateway-based expression vectors, including pUWG, and pVALIUM10 TRiP RNAi

plasmids. AttB1 and attB2 sites are 25 bp in length and generated by LR clonase-mediated recombination

between attL and attR sequences. Given the repetition of attB1 and attB2 sites, flanking the passenger

and guide target sequence fragments in pVALIUM10 TRiP RNAis, these sequences are transcribed and

could contribute to the dsRNA stem-loop substrate for Dicer. Based on these results we hypothesized that

dsRNAs generated from attB1 and attB2 could target, and thus silence, Gateway-based transgenes via

attB sites (Figure 2I).

pVALIUM10 RNAi lines silence Gateway transgenes via attB sites

To test our hypothesis, we cloned the *mCherry* coding sequence into the pUWG Gateway expression

vector (pUWG-*mCherry*) allowing ubiquitous expression of mCherry transcript flanked by the attB1 and

attB2 sites. The pUWG-*mCherry* vector then served as a template for overlap PCR to generate the

pUWG$^{\Delta attB}$-*mCherry* construct that lacks attB sites (Figure 3A). Both plasmids were used to establish

transgenic fly lines through a standard *Drosophila* germline transformation method (Rubin & Spradling,

1982). The susceptibility or resistance of *ubi-mCherry* (pUWG-*mCherry*) and *ubi$^{\Delta attB}$-mCherry* (pUWG$^{\Delta attB}$-

mCherry) transcripts to RNAi-mediated silencing was then tested by driving various RNAi lines under the

control of the *nub-Gal4* driver. Immunostaining of the third instar wing imaginal discs using an anti-RFP

antibody revealed an identical pattern of ubi-mCherry and ubi$^{\Delta attB}$-mCherry transgenes in the control

(*nub>*) background (Figure 3B, C). In contrast, levels of ubi-mCherry but not ubi$^{\Delta attB}$-mCherry transgenic

protein were markedly reduced in the wing pouch region relative to the rest of the tissue following

overexpression of *BuGZ* and *myc* pVALIUM10 TRiP RNAi lines (Figure 3D, E, H, I, J, M) while no significant

changes were observed when *BuGZ* and *myc* were knocked down using KK and pVALIUM20 RNAi lines,

respectively (Figure 3F, G, H, K, L, M). Importantly, the attB site-dependent silencing of ubi-mCherry was

recapitulated using an additional set of pVALIUM10 TRiP RNAi lines targeting *Atf3*, *Lip4*, *yki*, *DH44*, *ftz-f1*,

mago, *brat*, and *Actβ* genes (Supplementary Figure S1).

Finally, we tested whether pVALIUM10 TRiP RNAi lines may also interfere with UAS-inducible attB-

containing transcripts, such as transgenic fly open reading frames (ORFs) expressed from the Gateway

destination vector pGW-HA.attB (Bischof *et al*, 2014). To this end, we assessed levels of the HA-tagged

SmD3 FlyORF protein in the presence of pVALIUM10 or KK BuGZ-RNAi line. In contrast to an enrichment

of SmD3::3xHA protein and positive punctae in the wing pouch cells of control (*nub>*) and *BuGZ^RNAi[KK104498]*

expressing wing discs, the overall levels and the number of positive punctae were noticeably reduced in

nub>BuGZ^RNAi[TRiP.JF02830] cells (Figure 3N-Q). These results suggest that the dsRNA hairpin produced from

pVALIUM10 RNAi vectors generates attB1- and attB2-siRNAs, which guide destruction of transcripts

containing such target sites. This off-target silencing via attB sites is specific to RNAi lines based on

pVALIUM10. Absence of attB1 and attB2 sites renders the Gateway-derived transcript resistant to

pVALIUM10 TRiP RNAi.

DISCUSSION

The use of transgenic RNAi to achieve developmental stage- and tissue-specific knockdown of candidate

genes and to perform genome-wide screens in *Drosophila* has become commonplace in the past two

decades (Blake *et al*, 2017; Dietzl *et al*, 2007; Hu *et al*, 2017; Port *et al.*, 2011). Both our knowledge on the

RNAi mechanism, and the tools to generate new types of RNAi constructs efficiently have considerably

evolved in recent years. As a result, several independent transgenic RNAi stock collections were

established, including Vienna *Drosophila* stock center (VDRC, 23,411 RNAi lines,

https://stockcenter.vdrc.at/control/main), Bloomington *Drosophila* Stock Center (BDSC, 13,698 RNAi

lines, https://bdsc.indiana.edu/), and the National Institute of Genetics (NIG, 12,365 RNAi lines,

https://shigen.nig.ac.jp/fly/nigfly/) at the time of writing. Yet, in spite of being a broadly used technique,

transgenic manipulation of gene expression can still have unforeseen consequences. A large portion of

the RNAi lines from the VDRC KK collection were found to produce phenotypes due to a secondary

insertion and consequent ectopic expression of the *tiptop* gene (Green *et al*, 2014; Vissers *et al*, 2016).

More recently, van der Graaf and colleagues reported that transgenes inserted in the commonly used

attP40 landing site on the second chromosome can decrease the expression of Msp300, a member of the

LINC complex, and hence lead to nuclear size, shape and positioning phenotypes (van der Graaf *et al*,

2022).

Here, we demonstrate that the pVALIUM10-based RNAi lines cause off-target silencing of ORFs expressed

from Gateway expression vectors or potentially any transcripts containing attB1 and attB2 sites. We show

that the removal of these sites abrogates the undesirable knockdown. Given the popularity of the

Gateway expression vectors to generate transgenic protein-reporters and tagged overexpression

constructs, our results suggest that simultaneous use of pVALIUM10 TRiP RNAi may lead to experimental

artefacts unrelated to the investigated biological process.

Of note, in the original publication of pVALIUM10 vectors, the authors tested for potential off-target

effects by comparing the hairpin sequence against the *Drosophila* genome (Ni *et al.*, 2009), which does

not contain endogenous 21nt stretches equivalent to the attB1 and attB2 sequences. However, as the

transgenic toolkit in *Drosophila* is ever expanding, each additional transgene has the possibility to produce

a synthetic effect with the existing RNAi lines. Hence, we suggest that the researchers validate the type

of TRiP RNAi line or the expression vector sequence before commencing their experiments. As an

example, we do not recommend combining RNAi based on pVALIUM10 with the 3xHA-tagged

overexpression lines from FlyORF (https://flyorf.ch) (Bischof *et al.*, 2014) or the fly-FUCCI (Fluorescent

Ubiquitination-based Cell Cycle Indicator) lines, which allow real-time visualization of the cell cycle

dynamics by the expression two fluorescently tagged degrons (Zielke et al., 2014). In both cases, the

transgenes were generated via Gateway reaction and contains attB sites flanking the inserted ORF (Bischof

et al., 2014; Zielke *et al*, 2014), which can lead to off-target silencing of the transcript. Furthermore, this

finding reinforces the idea that RNAi experiments should adhere to the standards established in the past

decades, including investigating possible off-target effects to eliminate knockdown of the unintended

gene, using multiple independent RNAi lines to recapitulate phenotypes, and confirming the knockdown

of the targeted gene products (Echeverri *et al*, 2006; Green *et al.*, 2014; Kaya-Çopur & Schnorrer, 2016).

MATERIALS AND METHODS

Fly stocks and Husbandry

The following *Drosophila* strains were used: w^{1118} (BDSC; RRID: BDSC_3605), *nubbin-Gal4, UAS-myr-mRFP*

(*nub>mRFP*), *BuGZ^{FlyFos[v318366]}* (VDRC), *BuGZ^{RNAi[TRiP.JF02830]}* (BDSC; RRID:BDSC_27996), *BuGZ^{RNAi[KK104498]}*

(VDRC), *Lip4^{RNAi[TRiP.HMO5136]}* (BDSC; RRID:BDSC_28925), *myc^{RNAi[TRiP.JF01761]}* (BDSC; RRID:BDSC_25783),

myc^{RNAi[TRiP.HMS01538]} (BDSC; RRID:BDSC_36123), *myc^{RNAi[GD2948]}* (VDRC), *Actβ^{RNAi[TRiP.JF03276]}* (BDSC;

RRID:BDSC_29597), *ftz-f1^{RNAi[TRiP.JF02738]}* (BDSC; RRID:BDSC_27659), *yki^{RNAi[TRiP.JF03119]}* (BDSC;

RRID:BDSC_31965), *Atf3^{RNAi[TRiP.JF02303]}* (BDSC; RRID:BDSC_26741), *DH44^{RNAi[TRiP.JF01822]}* (BDSC;

RRID:BDSC_25804), *brat^{RNAi[TRiP.HM05078]}* (BDSC; RRID:BDSC_28590), *mago^{RNAi[TRiP.HM05142]}* (BDSC;

RRID:BDSC_28931), *SmD3::3xHA^{[FlyORF F003987]}* (FlyORF), *nub>SmD3::3xHA^{[FlyORF F003987]}* (Erkelenz *et al*, 2021),

ubi-RNase H1::GFP, *ubi-mCherry*, *ubi^{ΔattB}-mCherry* (this study). All *Drosophila* stocks are listed in

Supplementary Table S1. All crosses were set up and maintained at 25 °C, unless specified otherwise, on

a diet consisting of 0.8% agar, 8% cornmeal, 1% soymeal, 1.8% dry yeast, 8% malt extract, and 2.2% sugar-

beet syrup, which was supplemented with 0.625% propionic acid and 0.15% Nipagin. w^{1118} line was used

as control.

Generation of plasmids and transgenic lines

The open reading frame of *rnh1* (FBgn0023171) was amplified from *Drosophila melanogaster* cDNA (w^{1118})

using primers harboring NotI and SalI restriction sites. The stop codon was excluded to allow C-terminal

tagging of the protein. The ORF was cloned into the Gateway pENTR4 Dual Selection Vector (Thermo

Fisher Scientific; #A10465). The ORF was then recombined into the pUWG vector (DGRC; #1284) using the

LR Clonase II enzyme mix according to manufacturer's instructions (Thermo Fisher Scientific; #11791020).

The coding sequence of *mCherry* (GenBank: AY678264.1) was amplified with specific primers, cloned into

the pENTR 4 Dual Selection vector (Invitrogen, Cat. Nr. #A10465) including a stop codon and subsequently

recombined into the pUWG vector (pUWG-*mCherry*). Overlap PCR strategy was employed to generate the

pUWG$^{\Delta attB}$-*mCherry* construct lacking attB sites resulting from the LR recombination. Three separate PCR

reactions were performed to produce fragments for assembly using the original pUWG-*mCherry* plasmid

as a template. The forward primer of the first primer pair contained a XhoI restriction site, while the

reverse primer contained a XbaI restriction site. Partial complementarity of the reverse and forward

primers used to synthesize the downstream fragment enabled the exclusion of attB sites in the final

product. The final PCR product was ligated into the pUWG-*mCherry* plasmid digested with XbaI and XhoI.

Transgenic fly lines with random insertions of the constructs were generated by standard P-element-

mediated transformation using w^{1118} *Drosophila* strain. All primers and plasmids are listed in

Supplementary Table S2.

Tissue dissection and immunostaining

Wing imaginal discs dissected from third instar *Drosophila* larvae (7 days after egg laying (AEL)) were fixed

for 25 minutes with 4% paraformaldehyde in PBS containing 0.1% Triton X (PBS-T) at room temperature.

Fixed tissues were washed 3 times with PBS-T. Primary antibodies were diluted in blocking buffer (PBS-T

with 0.3% BSA) and tissues were stained overnight at 4 °C. The following primary antibodies were used:

rabbit anti-mRFP (1:500, MBL International Cat# PM005, RRID:AB_591279), goat anti-GFP (1:500, Abcam

Cat# ab6673, RRID:AB_305643), and rabbit-anti HA (1:1000, Abcam Cat# ab9110, RRID:AB_307019). After

washing, the samples were incubated with the corresponding Alexa Fluor 488 (1:2000, Thermo Fisher

Scientific Cat# A-11034, RRID:AB_2576217), Cy2 (1:2000, Jackson ImmunoResearch Labs Cat# 705-225-

147, RRID:AB_2307341) or Cy3 (Jackson ImmunoResearch Labs Cat# 711-165-152, RRID:AB_2307443)

conjugated secondary antibodies overnight at 4 °C, washed and counterstained with DAPI (1:1000 dilution

of 5 mg/ml stock, 6335.1, Carl Roth GmbH) to visualize nuclei. Tissues were mounted on glass slides in

Dabco-Mowiol 4-88 (Sigma-Aldrich Cat# D2522 and Cat# 81381).

Image acquisition and processing

Confocal images and stacks were acquired with Olympus FV1000 confocal microscope equipped with 20X

UPlan S-Apo (NA 0.85), 40X UPlan FL (NA 1.30) and 60X UPlanApo (NA1.35) objectives. Maximum Z-

projections were generated from consecutive sections taken at 1.4 µm steps using Fluoview 1000

Software (Olympus) (RRID: SCR_014215) and Fiji (https://fiji.sc/) (RRID:SCR_002285). Final image

processing, including panel assembly, brightness and contrast adjustments were performed in Adobe

Photoshop CC (Adobe Systems, Inc.) (RRID:SCR_014199).

Quantification of mCherry, RNase H1::GFP, and SmD3::3xHA signal

Fluorescent signal intensity was quantified with Fiji (https://fiji.sc/) (RRID:SCR_002285). For mCherry and

RNase H1::GFP, a square selection was created in the middle of the pouch region, and the „Mean

intensity" was measured, then the same selection was moved to the middle of the notum and the

measurement was repeated. The two values were divided to produce a pouch/notum intensity ratio for

each wing disc. For SmD3::3xHA, the „Mean intensities" of SmD3::3xHA and mRFP signals in the pouch

region were measured and divided to produce a normalized value for each wing disc. Zoom images of the

nubbin domain of independent WDs were used to count the number of SmD3::3xHA-positive punctae per

1000 μm^2 area. Statistical significance was determined by one-way ANOVA with Tukey's multiple

comparison test (signal intensity) or Kruskall-Wallis test (puncta) in GraphPad Prism (RRID:SCR_002798).

ACKNOWLEDGEMENTS

We thank Norbert Perrimon for discussions and comments on the manuscript. We thank the Bloomington

Drosophila Stock Center (BDSC, Bloomington, IN, USA), the Vienna *Drosophila* Resource Center (VDRC,

Vienna, Austria), *Drosophila* Genomic Resource Center (DGRC, supported by NIH grant 2P40OD010949,

Bloomington, IN, USA), the Zurich ORFeome Project (FlyORF, Zurich, Switzerland) for fly stocks and

plasmids. We are grateful to Tina Bresser for generation of transgenic lines and Nils Teuscher for fly stock

maintenance and technical assistance.

REFERENCES

Aughey GN, Grice SJ, Liu J-L (2016) The Interplay between Myc and CTP Synthase in Drosophila. *PLOS*
Genetics 12: e1005867

Bao S, Cagan R (2006) Fast cloning inverted repeats for RNA interference. *RNA* 12: 2020-2024

Bischof J, Sheils EM, Björklund M, Basler K (2014) Generation of a transgenic ORFeome library in
Drosophila. *Nature protocols* 9: 1607-1620

Blake AJ, Finger DS, Hardy VL, Ables ET (2017) RNAi-Based Techniques for the Analysis of Gene Function
in Drosophila Germline Stem Cells. *Methods in molecular biology (Clifton, NJ)* 1622: 161-184

Brand AH, Perrimon N (1993) Targeted gene expression as a means of altering cell fates and generating
dominant phenotypes. *Development* 118: 401-415

Crossley MP, Bocek M, Cimprich KA (2019) R-Loops as Cellular Regulators and Genomic Threats. *Mol Cell*
73: 398-411

Dietzl G, Chen D, Schnorrer F, Su KC, Barinova Y, Fellner M, Gasser B, Kinsey K, Oppel S, Scheiblauer S *et*
al (2007) A genome-wide transgenic RNAi library for conditional gene inactivation in Drosophila. *Nature*
448: 151-156

Echeverri CJ, Beachy PA, Baum B, Boutros M, Buchholz F, Chanda SK, Downward J, Ellenberg J, Fraser
AG, Hacohen N *et al* (2006) Minimizing the risk of reporting false positives in large-scale RNAi screens.
Nature Methods 3: 777-779

Erkelenz S, Stanković D, Mundorf J, Bresser T, Claudius A-K, Boehm V, Gehring NH, Uhlirova M (2021)
Ecd promotes U5 snRNP maturation and Prp8 stability. *Nucleic Acids Research*

Giordano E, Rendina R, Peluso I, Furia M (2002) RNAi triggered by symmetrically transcribed transgenes
in Drosophila melanogaster. *Genetics* 160: 637-648

Graca FA, Sheffield N, Puppa M, Finkelstein D, Hunt LC, Demontis F (2021) A large-scale transgenic RNAi
screen identifies transcription factors that modulate myofiber size in Drosophila. *PLOS Genetics* 17:
e1009926

Green EW, Fedele G, Giorgini F, Kyriacou CP (2014) A Drosophila RNAi collection is subject to dominant
phenotypic effects. *Nat Methods* 11: 222-223

Hagan CE, Warren GJ (1983) Viability of palindromic DNA is restored by deletions occurring at low but
variable frequency in plasmids of Escherichia coli. *Gene* 24: 317-326

Hu Y, Comjean A, Roesel C, Vinayagam A, Flockhart I, Zirin J, Perkins L, Perrimon N, Mohr SE (2017)
FlyRNAi.org-the database of the Drosophila RNAi screening center and transgenic RNAi project: 2017
update. *Nucleic Acids Res* 45: D672-D678

Kaya-Çopur A, Schnorrer F (2016) A Guide to Genome-Wide In Vivo RNAi Applications in Drosophila.
Methods Mol Biol 1478: 117-143

Kennerdell JR, Carthew RW (1998) Use of dsRNA-mediated genetic interference to demonstrate that
frizzled and frizzled 2 act in the wingless pathway. *Cell* 95: 1017-1026

Kim-Yip RP, Nystul TG (2018) Wingless promotes EGFR signaling in follicle stem cells to maintain self-
renewal. *Development* 145: dev168716

Ledru M, Clark CA, Brown J, Verghese S, Ferrara S, Goodspeed A, Su TT (2022) Differential gene
expression analysis identified determinants of cell fate plasticity during radiation-induced regeneration
in Drosophila. *PLoS Genet* 18: e1009989

Lee YS, Carthew RW (2003) Making a better RNAi vector for Drosophila: use of intron spacers. *Methods*
30: 322-329

Lesch C, Jo J, Wu Y, Fish GS, Galko MJ (2010) A targeted UAS-RNAi screen in Drosophila larvae identifies
wound closure genes regulating distinct cellular processes. *Genetics* 186: 943-957

Lohmann JU, Endl I, Bosch TC (1999) Silencing of developmental genes in Hydra. *Dev Biol* 214: 211-214

Neely GG, Kuba K, Cammarato A, Isobe K, Amann S, Zhang L, Murata M, Elmen L, Gupta V, Arora S *et al*
(2010) A global in vivo Drosophila RNAi screen identifies NOT3 as a conserved regulator of heart
function. *Cell* 141: 142-153

Ni JQ, Liu LP, Binari R, Hardy R, Shim HS, Cavallaro A, Booker M, Pfeiffer BD, Markstein M, Wang H *et al*
(2009) A Drosophila resource of transgenic RNAi lines for neurogenetics. *Genetics* 182: 1089-1100

Ni JQ, Markstein M, Binari R, Pfeiffer B, Liu LP, Villalta C, Booker M, Perkins L, Perrimon N (2008) Vector
and parameters for targeted transgenic RNA interference in Drosophila melanogaster. *Nat Methods* 5:
49-51

Perkins LA, Holderbaum L, Tao R, Hu Y, Sopko R, McCall K, Yang-Zhou D, Flockhart I, Binari R, Shim HS *et*
al (2015) The Transgenic RNAi Project at Harvard Medical School: Resources and Validation. *Genetics*
201: 843-852

Perrimon N, Ni JQ, Perkins L (2010) In vivo RNAi: today and tomorrow. *Cold Spring Harb Perspect Biol* 2:
a003640

Piccin A, Salameh A, Benna C, Sandrelli F, Mazzotta G, Zordan M, Rosato E, Kyriacou CP, Costa R (2001)
Efficient and heritable functional knock-out of an adult phenotype in Drosophila using a GAL4-driven
hairpin RNA incorporating a heterologous spacer. *Nucleic Acids Research* 29

Pletcher RC, Hardman SL, Intagliata SF, Lawson RL, Page A, Tennessen JM (2019) A Genetic Screen Using
the Drosophila melanogaster TRiP RNAi Collection To Identify Metabolic Enzymes Required for Eye
Development. *G3 (Bethesda)* 9: 2061-2070

Port F, Hausmann G, Basler K (2011) A genome-wide RNA interference screen uncovers two p24 proteins
as regulators of Wingless secretion. *EMBO Rep* 12: 1144-1152

Reece-Hoyes JS, Walhout AJM (2018) Gateway Recombinational Cloning. *Cold Spring Harb Protoc* 2018:
pdb.top094912

Rotelli MD, Bolling AM, Killion AW, Weinberg AJ, Dixon MJ, Calvi BR (2019) An RNAi Screen for Genes
Required for Growth of Drosophila Wing Tissue. *G3 Genes|Genomes|Genetics* 9: 3087-3100

Rubin GM, Spradling AC (1982) Genetic transformation of Drosophila with transposable element vectors.
Science 218: 348-353

Rust K, Tiwari MD, Mishra VK, Grawe F, Wodarz A (2018) Myc and the Tip60 chromatin remodeling
complex control neuroblast maintenance and polarity in Drosophila. *EMBO J* 37

Rylee J, Mahato S, Aldrich J, Bergh E, Sizemore B, Feder LE, Grega S, Helms K, Maar M, Britt SG *et al*
(2022) A TRiP RNAi screen to identify molecules necessary for Drosophila photoreceptor differentiation.
G3 Genes|Genomes|Genetics: jkac257

Saj A, Arziman Z, Stempfle D, van Belle W, Sauder U, Horn T, Dürrenberger M, Paro R, Boutros M,
Merdes G (2010) A Combined Ex Vivo and In Vivo RNAi Screen for Notch Regulators in Drosophila
Reveals an Extensive Notch Interaction Network. *Developmental Cell* 18: 862-876

Santos-Pereira JM, Aguilera A (2015) R loops: new modulators of genome dynamics and function. *Nat*
Rev Genet 16: 583-597

Sarov M, Schneider S, Pozniakovski A, Roguev A, Ernst S, Zhang Y, Hyman AA, Stewart AF (2006) A
recombineering pipeline for functional genomics applied to Caenorhabditis elegans. *Nat Methods* 3:
839-844

Tomari Y, Zamore PD (2005) Perspective: machines for RNAi. *Genes Dev* 19: 517-529

van der Graaf K, Srivastav S, Singh P, McNew JA, Stern M (2022) The Drosophila attP40 docking site and
derivatives are insertion mutations of MSP300. *bioRxiv*: 2022.2005.2014.491875

Vissers JH, Manning SA, Kulkarni A, Harvey KF (2016) A Drosophila RNAi library modulates Hippo
pathway-dependent tissue growth. *Nat Commun* 7: 10368

Wargelius A, Ellingsen S, Fjose A (1999) Double-stranded RNA induces specific developmental defects in
zebrafish embryos. *Biochem Biophys Res Commun* 263: 156-161

Wianny F, Zernicka-Goetz M (2000) Specific interference with gene function by double-stranded RNA in
early mouse development. *Nat Cell Biol* 2: 70-75

Zamore PD, Tuschl T, Sharp PA, Bartel DP (2000) RNAi: double-stranded RNA directs the ATP-dependent
cleavage of mRNA at 21 to 23 nucleotide intervals. *Cell* 101: 25-33

Zeng X, Han L, Singh SR, Liu H, Neumuller RA, Yan D, Hu Y, Liu Y, Liu W, Lin X *et al* (2015) Genome-wide
RNAi screen identifies networks involved in intestinal stem cell regulation in Drosophila. *Cell Rep* 10:
1226-1238

Zhou J, Xu L, Duan X, Liu W, Zhao X, Wang X, Shang W, Fang X, Yang H, Jia L *et al* (2019) Large-scale RNAi
screen identified Dhpr as a regulator of mitochondrial morphology and tissue homeostasis. *Science
Advances* 5: eaax0365

Zielke N, Korzelius J, van Straaten M, Bender K, Schuhknecht GFP, Dutta D, Xiang J, Edgar BA (2014) Fly-
FUCCI: A versatile tool for studying cell proliferation in complex tissues. *Cell Rep* 7: 588-598
372

Figure 1. pVALIUM10 and KK RNAi lines cause efficient silencing.

(A-C) Expression of pVALIUM10 *BuGZ^RNAi[TRIP.JF02830]* (B, B') and KK-based *BuGZ^RNAi[KK104498]* transgenic RNAi

lines (C, C') using the *nubbin-Gal4, UAS-myr-mRFP* driver (*nub>mRFP*, light blue) causes a marked

reduction of a BuGZ^FlyFos[v318366] transgenic protein in the pouch region of wing discs (WDs) relative to the

hinge and notum, and control WD (A, A'). Micrographs show projections of multiple confocal sections of

WDs dissected from third instar larvae 7 days AEL. The multiple-tagged BuGZ^FlyFos [v318366] protein was

visualized by immunostaining with an anti-GFP antibody (white). Nuclei were counterstained with DAPI

(magenta). Scale bars: 100 μm. **(D)** Schematic representation of various vectors used for cloning and

overexpression of dsRNA. In the absence of the target-specific antibody, the RNAi efficacy can be assessed

with the help of FlyFos transgenes expressing multiple-tagged proteins under the control of the

endogenous regulatory sequences.

Figure 2. pVALIUM10-based lines cause off-target silencing of the RNAse H1::GFP transgenic reporter.

(A) Schematic representation of the transgenic *pUWG-RNase H1* reporter construct expressing the C-terminally-tagged RNase H1::GFP protein from the ubiquitous (*ubi*) promoter. The *RNase H1* coding sequence is flanked by attB1 and attB2 sites generated via Gateway-mediated recombination catalyzed by the LR clonase. **(B-G)** Immunostaining with a GFP-specific antibody revealed a marked reduction of the RNase H1::GFP levels in the nubbin domain (mRFP, light blue), where pVALIUM10-based *BuGZ^RNAi[TRIP.JF02830]* (C) and *myc^RNAi[TRiP.JF01761]* (E) RNAi lines were expressed using the *nubbin-Gal4, UAS-myr-mRFP* driver (*nub>mRFP*). No such downregulation of the RNase H1::GFP signal was observed in control WD (B) and following expression of KK-based *BuGZ^RNAi[KK104498]* (D), pVALIUM20-based *myc^RNAi[TRiP.HMS01538]*

(F) or GD-based *myc^RNAi[GD2948]* (G). Micrographs show projections of multiple confocal sections of WDs

dissected from third instar larvae 7 days AEL. Nuclei were counterstained with DAPI (magenta). Scale bars:

100 μm. **(H)** Quantification of the RNase H1::GFP signal depicted as relative intensity ratio between the

pouch and notum region. Statistical significance was determined by one-way ANOVA with Tukey's

multiple comparison test; n ≥ 3; **p<0.01, ****p<0.0001, n.s. non-significant. **(I)** Generation of

pVALIUM10 RNAi lines relied on a LR clonase-mediated recombination of dsRNA fragment from an Entry

vector to the pVALIUM10 destination vector (Ni *et al.*, 2009). The dsRNA hairpin produced from the

pVALIUM10 dsRNA expression vector is the source of siRNA species targeting the transcript of interest

(target ORF) but also attB1 and attB2 sites present in any of the Gateway expression vectors generated

via the same recombination mechanisms.

Figure 3

Figure 3. pVALIUM10 RNAi lines silence Gateway transgenes via attB1 and attB2 sites.

(A) Schematic of the transgenic *pUWG-mCherry* and mutant *pUWG^{ΔattB}-mCherry* reporter constructs. The mCherry is expressed from the poly-ubiquitin (*ubi*) promoter. **(B, C)** *ubi-mCherry* and mutant *ubi^{ΔattB}-mCherry* reporters showed similar expression pattern in the wing imaginal discs. **(D-M)** *ubi-mCherry* was efficiently downregulated by pVALIUM10 *BuGZ$^{RNAi[TRiP.JF02830]}$* and *myc$^{RNAi[TRiP.JF01761]}$* expressed under the control of the *nubbin-Gal4* driver (*nub>*) relative to the rest of the WD (D, I, H, M). In contrast, absence of attB1 and attB2 sequences rendered the *ubi^{ΔattB}-mCherry* reporter resistant to pVALIUM10 RNAi-mediated silencing (E, H, J, M). Neither KK- (*BuGZ$^{RNAi[KK104498]}$*) nor pVALIUM20-based (*myc$^{RNAi[TRiP.HMS01538]}$*) RNAi lines affected mCherry levels expressed from the unmodified or mutant reporter constructs (F-H, K-M). Micrographs show projections of multiple confocal sections of WDs dissected from third instar larvae 7 days AEL that were immunostained with an anti-mRFP antibody (white). Nuclei were counterstained with DAPI (magenta). Scale bars: 100 µm. Quantification of mCherry signal depicted as relative intensity ratio between the pouch and notum region (H, M). Superscript text in each group denotes the type of RNAi used. Statistical significance was determined by one-way ANOVA with Tukey's multiple comparison test; n ≥ 3; **p<0.01, ****p<0.0001, n.s. non-significant. **(N-Q)** Expression of pVALIUM10 *BuGZ$^{RNAi[TRiP.JF02830]}$* (O) in the wing pouch using *nubbin-Gal4, UAS-mRFP* driver (*nub>mRFP*) reduces SmD3::3xHA levels compared to control (N) and KK-based *BuGZ$^{RNAi[KK104498]}$* (P). Note foci of SmD3::3xHA protein in control (N") and *BuGZ$^{RNAi[KK104498]}$* cells (P") cells that are not present in *BuGZ$^{RNAi[TRiP.JF02830]}$* expressing cells (O"). Micrographs show projections of multiple or single (N"-P") confocal sections of WDs dissected from third instar larvae 7 days AEL that were immunostained with an anti-HA antibody (white, N'-P'; red, N"-P"). Nuclei were counterstained with DAPI (magenta). Scale bars: 100 µm or 5 µm (N"-P"). Quantification of the SmD3::3xHA signal intensity (Q, left) and SmD3::3xHA-positive punctae (Q, right) in the pouch region. Statistical significance was determined by one-way ANOVA with Tukey's multiple comparison test and Kruskall-Wallis test, respectively; n ≥ 6; **p<0.01, ***p<0.001, n.s. non-significant.