2   **Propagation of viral genomes by replicating ammonia-oxidising archaea during**

     soil nitrification

     Sungeun Lee[1], Ella T. Sieradzki[1], Graeme W. Nicol[1]* and Christina Hazard[1]*

     [1] Univ Lyon, CNRS, INSA Lyon, Université Claude Bernard Lyon 1, Ecole Centrale de

8   Lyon, Ampère, UMR5005, 69134 Ecully, France

10   * Corresponding authors. Email: graeme.nicol@ec-lyon.fr

                                       christina.hazard@ec-lyon.fr

12 **Abstract**

Ammonia-oxidising archaea (AOA) are a ubiquitous component of microbial

14 communities and dominate the first stage of nitrification in some soils. While we are

beginning to understand soil virus dynamics, we have no knowledge of the

16 composition or activity of those infecting nitrifiers or their potential to influence

processes. This study aimed to characterise viruses having infected autotrophic AOA

18 in two nitrifying soils of contrasting pH by following transfer of assimilated CO_2-derived

^{13}C from host to virus via DNA stable-isotope probing and metagenomic analysis.

20 Incorporation of ^{13}C into low GC mol% AOA and virus genomes increased DNA

buoyant density in CsCl gradients but resulted in co-migration with dominant non-

22 enriched high GC mol% genomes, reducing sequencing depth and contig assembly.

We therefore developed a hybrid approach where AOA and virus genomes were

24 assembled from low buoyant density DNA with subsequent mapping of ^{13}C isotopically

enriched high buoyant density DNA reads to identify activity of AOA. Metagenome-

26 assembled genomes were different between the two soils and represented a broad

diversity of active populations. Sixty-four AOA-infecting viral operational taxonomic

28 units (vOTUs) were identified with no clear relatedness to previously characterised

prokaryote viruses. These vOTUs were also distinct between soils, with 42% enriched

30 in ^{13}C derived from hosts. The majority were predicted as capable of lysogeny and

auxiliary metabolic genes included an AOA-specific multicopper oxidase suggesting

32 infection may augment copper uptake essential for central metabolic functioning.

These findings indicate virus infection of AOA may be a frequent process during

34 nitrification with potential to influence host physiology and activity.

Main text

Microbially mediated oxidation of ammonia to nitrate during nitrification is a central component of the global nitrogen (N) cycle. It is also responsible for major losses of

applied fertiliser N in soil, generating atmospheric pollution via direct and indirect production of nitrous oxide (N_2O) as well as nitrate (NO_3^-) pollution of groundwater [1].

Autotrophic ammonia-oxidising archaea (AOA) of the class *Nitrososphaeria* are a ubiquitous component of soil microbial communities and often dominate ammonia

oxidation and nitrification-associated N_2O emissions when ammonia is supplied at low rates via organic matter mineralisation [2], slow-release fertilisers [3] or in acidic soils

[4]. Integrated temperate viruses (proviruses) and other virus-associated protein-encoding genes are found in most AOA genomes suggesting frequent interaction (see

Supplementary Text). While viruses infecting marine AOA have been characterised through metagenomic approaches [5] and cultivation [6], those infecting soil AOA or

other nitrifier groups are currently uncharacterised.

         Virus infection can influence biogeochemical cycling by augmenting host

activity or causing cell mortality and subsequent release of nutrients [7]. Recent advances have demonstrated that soil virus communities are dynamic in a wide range

of soils [e.g. 8, 9] and augmenting virus loads modulate C and N fluxes [10, 11]. Nevertheless, identifying active interactions with specific populations or functional

groups in soil remains challenging due to structural complexity and the vast diversity of hosts and viruses. Recent use of stable-isotope approaches has investigated whole

community host-virus dynamics [12, 13] or interactions between individual host-virus populations specific to a functional process and substrate [14]. The aim of this study

was to utilise the latter approach with $^{13}CO_2$-based DNA-SIP to focus on nitrification-associated interactions and to test the hypothesis that viruses are a dynamic

component of soil AOA activity.

 Hybrid analysis of GC mol% fractionation and ^{13}C-DNA-SIP for identifying active AOA hosts and viruses

Microcosms were established (see Supplementary Text) with two sandy loam soils of contrasting pH (4.5 or 7.5), which were characterised previously as containing distinct

AOA communities [15]. As ammonia from hydrolysed urea is used by AOA in these and other soils [1, 3], nitrification was fuelled by addition of urea (two pulses of 100 µg

N g^{-1} soil (dry weight)) to stimulate AOA growth and potential enrichment of AOA-infecting viruses. Microcosms were incubated at 25°C for 30 days (Supplementary

Figure 1) with a 5% ^{12}C- or ^{13}C-CO_2 headspace to target viruses of autotrophs via transfer of assimilated carbon. Extracted DNA was subjected to isopycnic

centrifugation in CsCl gradients, in which DNA migrates as a function of GC mol% and isotopic enrichment, before fractionation and analysis of genomic DNA of different

buoyant densities [14]. Quantification of AOA genomic DNA distribution across CsCl gradients demonstrated growth of AOA populations (Figure 1A; Supplementary Figure

2) and genomic DNA in high buoyant density (HBD) fractions (>1.719 g ml^{-1}) was then pooled for each replicate microcosm and metagenomes sequenced (Supplementary

Table 1). While ^{13}C-incorporation clearly separated DNA from growing vs non-growing AOA populations, ^{13}C-enriched AOA genomes (average contig GC mol% content of

38%) co-migrated within the CsCl gradient with more abundant community ^{12}C genomes with higher GC mol% (>57%), reducing AOA host and virus sequencing

depth, contig assembly and binning. Consequently, only one medium-quality AOA metagenome-assembled genome (MAG) was recovered from HBD DNA and no AOA-

associated virus contigs were identified amongst 290 viral operational taxonomic units (vOTUs) ≥10 kb (Figure 1B).

To facilitate analysis of low GC mol% genomes while also identifying activity, we developed a hybrid approach combining metagenomic analysis of low buoyant density (LBD) ^{12}C DNA with read-mapping of sequenced HBD DNA (Figure 1; Supplementary Figure 2). The AOA and virus genomes were assembled from metagenomic libraries of ^{12}C LBD DNA only (<1.699 g ml^{-1}) representing both dormant and active populations. Individual sequence reads from HBD DNA libraries from both ^{12}C- and ^{13}C-incubations were then mapped to identify activity with a significantly greater relative abundance in ^{13}C-incubated samples demonstrating isotope incorporation. For taxonomically defined contigs of hosts or host-linked viruses, in comparison to HBD, assemblies from LBD libraries increased the relative proportion of contigs from the AOA-containing *Thermoproteota* phylum (GTDB taxonomy; *Nitrososphaerota* in ICNP taxonomy and *Thaumarchaeota* in NCBI taxonomy) and *Thermoproteota*-infecting virus contigs from 0.3 to 2.8% and 0 to 22.3%, respectively (Supplementary Figure 3). In LBD metagenomes, 123 medium- and high-quality MAGs from both soils were assigned taxonomy (Supplementary Table 2) including nine AOA MAGs representing six genera (GTDB classification [16]) within the *Nitrososphaerales* and *Nitrosopumilales* orders and one nitrite-oxidising bacterium (NOB) *Nitrobacter* MAG (MAG 15) (Figure 1B, Supplementary Figure 4), contributing to the oxidation of nitrite (NO_2^-) to NO_3^- during the second step of canonical nitrification. Mapping of HBD sequence reads revealed that eight of the ten MAGs were derived from active autotrophic populations (Figure 1C). Consistent with previous studies characterising the distribution of soil AOA from different taxa [17], this included four *Nitrosotalea* genus populations (MAGs 9, 61, 77 and 107) from pH 4.5 soil and three *Nitrososphaera* genus populations (MAGs 12, 52 and 96) from pH 7.5 soil. No evidence of autotrophic growth was observed for populations represented by a further two MAGs within the *Nitrososphaerales*. Both are from genera with no cultivated

114   representatives with MAG 112 from genus TA-21 (*amoA*-gene lineage NS-β-2 [17])

and MAG 63 (*amoA*-gene lineage NS-δ-2.1) representing a novel genus (designated

116   here as CS63). The latter belongs to the *amoA* gene-defined NS-δ lineage that while

widely found in soil, is rarely implicated in contributing to ammonia oxidation. Although

118   these two MAGs may simply be from AOA populations not active under the incubation

conditions used, they are also consistent with microorganisms possessing

120   physiologies distinct from cultivated AOA.

122   *AOA viruses*

After predicting contigs of virus origin using established bioinformatic tools, those from

124   AOA-infecting viruses were identified with a custom database of hallmark genes from

AOA proviruses and analysis of other shared homologues (see Supplementary Text).

126   Specifically, proviruses were identified in 184 (87%) of 212 AOA genome sequences

downloaded from GTDB. Capsid, terminase, portal and integrase protein sequences

128   from these proviruses were subsequently used as search queries. While previous

analysis of 12 total soil metagenome or virome libraries from the same soil samples

130   did not enable identification of any AOA-associated virus contigs (average of 152

million quality filtered reads and 1,947 contigs ≥10 kb per library) [14], this targeted

132   approach using LBD DNA from nitrifying microcosms yielded 64 contigs predicted as

derived from AOA-infecting viruses and each representing a unique vOTU. In addition,

134   viruses infecting other nitrifier populations were identified with five and three vOTUs

derived from viruses infecting ammonia-oxidising bacteria (AOB) and NOB,

136   respectively. Virus and host genome GC mol% correlate [18] and AOA-infecting virus

contigs possessed a similar average content of 36.3% which was considerably lower

138   than averages of 46.6% and 67.0% of all viral contigs recovered from LBD and HBD

metagenomes, respectively. As with their hosts, AOA virus communities were distinct

140   between the two pH soils (Figure 2A) as also observed previously for total virus

communities [19] with 29 (45%) and 34 (53%) vOTUs significantly more relatively

142   abundant in pH 4.5 and 7.5 LBD libraries compared to the other soil, respectively.

Comparison of the relative abundance of HDB sequence reads from both ^{12}C and ^{13}C

144   libraries revealed that 27 contigs from both soils (42%) were enriched in ^{13}C and thus

confirming infection during incubation and transfer of C from AOA hosts to viruses.

146   Soil AOA viruses were distinct from characterised archaeal and bacterial

viruses. Gene-sharing network analysis using vConTACT v2.0 [20] did not identify any

148   associations with reference sequences or recently described viruses of marine AOA

[6] and genome-wide sequence comparisons [21] placed 60 of the 64 soil AOA viral

150   contigs within a unique proteomic grouping (Figure 2B). The majority were predicted

as capable of lysogeny with 55 encoding an integrase. While it can be difficult to

152   determine virulent versus temperate capability from partial genomes, only 6.8 and

8.9% of all virus contigs from LBD and HBD metagenomes possessed a predicted

154   integrase. Although suggesting a higher proportion of AOA viruses may be temperate,

it should be noted that this could be a consequence of using AOA provirus-associated

156   genes to identify viral contigs, biasing towards identification of those from temperate

viruses. These results also highlight that while DNA-SIP of total metagenomes

158   enables identification of ^{13}C incorporation into virus genomes, it is potentially limited

in facilitating confident bioinformatic prediction of free temperate virus particles versus

160   proviruses present from a previous infection of a currently active host. While the

integration status of a temperate virus can be predicted bioinformatically, using

162   targeted "viromes" metagenomes of filtered virus particles will provide greater

confidence of identifying genomes from free virus particles.

164   Eleven contigs contained large terminase (TerL) sub-unit gene sequences and
were similar to those recovered from other predicted AOA virus or provirus genomes
166   only (e-value <10^{-5}) (Figure 2C). In addition to viral hallmark genes encoding portal,
endonuclease, tail and capsid proteins, contig 053596 contained a type 1 multicopper
168   oxidase (MCO1) homologue found exclusively in AOA genomes [22] (Figure 2D).
Copper (Cu) is essential for ammonia oxidation and electron transport in AOA and
170   MCO1 is upregulated under Cu limitation and potentially involved in Cu sequestration
from the environment [23]. This virus-encoded MCO1 homologue therefore represents
172   an auxiliary metabolic gene that may contribute to a core process in AOA physiology.

174   *Summary*

These results from two soils indicate that virus infection of AOA during nitrification is
176   likely a common and dynamic process, with the potential to influence nitrification rates
in soil. Although viruses were only identified from contigs representing partial viral
178   genomes, analysis of individual marker genes and genome-wide similarity
comparisons revealed that those infecting AOA are highly divergent from known
180   viruses. While one *Nitrobacter* MAG was obtained and both AOB and NOB viruses
were also predicted, the hybrid approach developed in this study targeted low GC
182   mol% DNA and was specifically tailored for the analysis of AOA populations and their
viruses. A broader analysis of all viruses infecting nitrifying prokaryotic populations
184   would therefore likely require a more comprehensive sequencing effort of all fractions
throughout the CsCl gradient using quantitative SIP metagenomics [24] or alternative
186   approaches, such as virus-targeted metagenomic viromes.

188   *Data availability*

Metagenome sequence data are deposited in NCBI's GenBank under BioProject

190   accession nos. PRJNA621418-PRJNA621429 (^{13}C data) and PRJNA868779 (^{12}C

data) with MAGs available under accession numbers SAMN30436125-

192   SAMN30436246.

194   *Conflict of interest*

The authors declare that they have no conflict of interest.

Acknowledgements

198   This work was funded by an AXA Research Chair in Microbial Ecology awarded to

GWN, Agence Nationale de la Recherche grant 'CONSERVE' (ANR-22-CE02-0006-

200   01) awarded to GWN and CH, and an EU Horizon Europe MSCA

Postdoctoral Fellowship 'DIVOBIS' awarded to ETS. Sequencing data generated

202   under JGI Community Science Program Proposal No. 503702 awarded to GWN and

CH was conducted by the US DOE JGI, a DOE Office of Science User Facility, and

204   supported by the Office of Science of the US DOE under Contract No. DE-AC02-

05CH11231. The authors would like to thank Dr. Willm Martens-Habbena (University

206   of Florida) for continued discussions on soil AOA ecophysiology and taxonomy, Dr.

Melina Kerou (University of Vienna) for guidance on AOA MCO diversity, Dr. Gary

208   Trubl for comments on the manuscript and Dr. Robin Walker (SRUC, Aberdeen) for

access to the much-lamented Craibstone Estate Woodlands Field long term

210   experiment.

212   *Author contributions*

SL, GWN and CH conceived and designed the experiment with additional input from

214   ETS. SL performed all laboratory work and analysed the data with input from GWN in

phylenetic analyses. SL, GWN and CH drafted the manuscript text and figures with

216   all authors contributing to the final paper.

218   **References**

1. Prosser JI, Hink L, Gubry-Rangin C, Nicol GW. Nitrous oxide production by
220   ammonia oxidizers: Physiological diversity, niche differentiation and potential
    mitigation strategies. Glob Chang Biol. 2020;26:103–118.

222   2. Huang L, Chakrabarti S, Cooper J, Perez A, John SM, Daroub SH, et al. Ammonia-
    oxidizing archaea are integral to nitrogen cycling in a highly fertile agricultural soil.
224   ISME Commun. 2021;1:19.

3. Hink L, Gubry-Rangin C, Nicol GW, Prosser JI. The consequences of niche and
226   physiological differentiation of archaeal and bacterial ammonia oxidisers for
    nitrous oxide emissions. ISME J. 2018;12:1084–1093.

228   4. Li Y, Chapman SJ, Nicol GW, Yao H. Nitrification and nitrifiers in acidic soils. Soil
    Biol Biochem. 2018;116:290–301.

230   5. Ahlgren NA, Fuchsman CA, Rocap G, Fuhrman JA. Discovery of several novel,
    widespread, and ecologically distinct marine Thaumarchaeota viruses that encode
232   *amoC* nitrification genes. ISME J. 2019;13:618–631.

6. Kim J-G, Kim S-J, Cvirkaite-Krupovic V, Yu W-J, Gwak J-H, López-Pérez M, et al.
234   Spindle-shaped viruses infect marine ammonia-oxidizing thaumarchaea. Proc
    Natl Acad Sci U S A. 2019;116:15645–15650.

236   7. Emerson JB. Soil Viruses: A New Hope. mSystems. 2019;4:e00120-19.

8. Santos-Medellín C, Estera-Molina K, Yuan M, Pett-Ridge J, Firestone MK,
238   Emerson JB. Spatial turnover of soil viral populations and genotypes overlain by
    cohesive responses to moisture in grasslands. Proc Natl Acad Sci U S A.
240   2022;119:e2209132119.

9. Wu R, Davison MR, Gao Y, Nicora CD, Mcdermott JE, Burnum-Johnson KE, et al. Moisture modulates soil reservoirs of active DNA and RNA viruses. Commun Biol. 2021;4:992.

10. Braga LPP, Spor A, Kot W, Breuil M-C, Hansen LH, Setubal JC, et al. Impact of phages on soil bacterial communities and nitrogen availability under different assembly scenarios. Microbiome. 2020;8:52.

11. Albright MBN, Gallegos-Graves LV, Feeser KL, Montoya K, Emerson JB, Shakya M, et al. Experimental evidence for the impact of soil viruses on carbon cycling during surface plant litter decomposition. ISME Commun. 2022;2:1–8.

12. Starr EP, Shi S, Blazewicz SJ, Koch BJ, Probst AJ, Hungate BA, et al. Stable-isotope-informed, genome-resolved metagenomics uncovers potential cross-kingdom interactions in rhizosphere soil. mSphere. 2021;6:e0008521.

13. Trubl G, Kimbrel JA, Liquet-Gonzalez J, Nuccio EE, Weber PK, Pett-Ridge J, et al. Active virus-host interactions at sub-freezing temperatures in Arctic peat soil. Microbiome. 2021;9:208.

14. Lee S, Sieradzki ET, Nicolas AM, Walker RL, Firestone MK, Hazard C, et al. Methane-derived carbon flows into host–virus networks at different trophic levels in soil. Proc Natl Acad Sci U S A. 2021;118:e2105124118.

15. Nicol GW, Leininger S, Schleper C, Prosser JI. The influence of soil pH on the diversity, abundance and transcriptional activity of ammonia oxidizing archaea and bacteria. Environ Microbiol. 2008;10:2966–2978.

16. Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit to classify genomes with the Genome Taxonomy Database. Bioinformatics. 2020;36:1925–1927.

17. Alves RJE, Eloy Alves RJ, Minh BQ, Urich T, von Haeseler A, Schleper C. Unifying
    the global phylogeny and environmental distribution of ammonia-oxidising
    archaea based on amoA genes. Nature Commun. 2018;9:1517.

18. Cardinale DJ, Duffy S. Single-stranded genomic architecture constrains optimal
    codon usage. Bacteriophage. 2011;1:219–224.

19. Lee S, Sorensen JW, Walker RL, Emerson JB, Nicol GW, Hazard C. Soil pH
    influences the structure of virus communities at local and global scales. Soil Biol
    Biochem. 2022;166:108569.

20. Bin Jang H, Bolduc B, Zablocki O, Kuhn JH, Roux S, Adriaenssens EM, et al.
    Taxonomic assignment of uncultivated prokaryotic virus genomes is enabled by
    gene-sharing networks. Nat Biotechnol. 2019;37:632–639.

21. Nishimura Y, Yoshida T, Kuronishi M, Uehara H, Ogata H, Goto S. ViPTree: the
    viral proteomic tree server. Bioinformatics. 2017;33:2379–2380.

22. Kerou M, Offre P, Valledor L, Abby SS, Melcher M, Nagler M, et al. Proteomics
    and comparative genomics of *Nitrososphaera viennensis* reveal the core genome
    and adaptations of archaeal ammonia oxidizers. Proc Natl Acad Sci U S A.
    2016;113:7937–7946.

23. Reyes C, Hodgskiss LH, Kerou M, Pribasnig T, Abby SS, Bayer B, et al. Genome
    wide transcriptomic analysis of the soil ammonia oxidizing archaeon
    Nitrososphaera viennensis upon exposure to copper limitation. ISME J.
    2020;14:2659–2674.

24. Sieradzki ET, Greenlon A, Nicolas AM, Firestone MK, Pett-Ridge J, Blazewicz SJ,
    et al. Functional succession of actively growing soil microorganisms during
    rewetting is shaped by precipitation history. bioRxiv. 2022;2022.06.28.498032.

 Figure legends

Figure 1. Contigs and MAGs from low GC mol% DNA and *in situ* growth of AOA in pH 4.5 and 7.5 soils determined from hybrid analysis of ^{12}C- and ^{13}C-enriched DNA. **A** Distribution of total prokaryotic and AOA genomes in CsCl gradients determined from the relative abundance of 16S rRNA genes and *amoA* genes, respectively. Example profiles are shown for pH 4.5 soil only (see Supplementary Figure 2 for pH 7.5 profiles). Vertical error bars are the standard error of the mean relative abundance and horizontal bars (mostly smaller than the symbol size) the standard error of the mean buoyant density of individual fractions from three independent CsCl gradients, each representing an individual microcosm. Genomic DNA in fractions highlighted in blue or pink areas were pooled for each replicate microcosm for metagenomic sequencing. **B** Distribution of assembled contigs ($\geq$5 kb) from metagenomic libraries prepared from LBD or HBD DNA in sequence coverage vs. GC mol% plots. Contigs binned into ten medium- and high-quality nitrifier MAGs and predicted viruses of nitrifiers ($\geq$10 kb) are highlighted, with number in parenthesis beside each MAG identifier giving estimated completeness and contamination (%). **C** Mean and standard error of the relative proportion of metagenomic reads from ^{12}C- and ^{13}C-enriched HBD DNA mapped onto contigs of nine AOA and one *Nitrobacter* MAGs. A significantly greater relative proportion of reads in ^{13}C-derived libraries are indicated with * (p <0.05, two-sample Student's t-test or Welsch's t-test when variances were not homogenous). GTDB genus [16] and *amoA* gene lineage-affiliation [17] are given.

Figure 2. Identification and determination of activity of viruses infecting AOA in pH 4.5 and 7.5 soils using a hybrid analysis of ^{12}C- and ^{13}C-enriched DNA. **A** Heatmap showing the relative abundance of 64 vOTUs identified in LBD DNA libraries from pH 4.5 and 7.5 soil ($\geq$1× coverage, $\geq$75% contig breadth). To identify activity, reads from

316   ^{12}C- and ^{13}C-enriched HBD DNA were mapped onto LBD-derived contigs with a

significantly greater relative proportion of reads in ^{13}C-derived libraries indicated with

318   * (p <0.05, two-sample Student's t-test or Welsch's t-test when variances were not

homogenous). **B** Proteomic tree showing genome-wide sequence similarities between

320   AOA viral contigs and 1,981 curated virus genomes. Values at dotted circles represent

a distance metric based on normalized tBLASTx scores plotted on a log scale. **C**

322   Maximum likelihood phylogenetic analysis of derived large sub-unit terminase (TerL)

protein sequences (336 unambiguously aligned positions, LG substitution model)

324   identified in this and other metagenomic studies or in proviruses of cultivated AOA.

NCBI accession numbers are given in parenthesis. Circles at nodes represent

326   percentage bootstrap support from 1000 replicates, scale bar denotes an estimated

0.05 changes per position and colour-coding describes the comparative relative

328   abundance in each soil. **D** Genetic map of AOA-infecting virus-derived contig 053596

containing a type 1 multicopper oxidase (MCO1) gene. Maximum likelihood

330   phylogenetic tree describes placement of MCO1 within the AOA MCO/NirK family of

cultivated AOA following designations of Kerou *et al.* [22] (206 unambiguously aligned

332   positions, LG substitution model, invariant and gamma distributed sites). NCBI

accession numbers are given in parenthesis. Circles at nodes represent percentage

334   bootstrap support from 1000 replicates and the scale bar denotes an estimated 0.05

changes per position.

A

Prokaryotic 16S rRNA genes

LBD DNA — HBD DNA

AOA *amoA* genes

LBD DNA — HBD DNA

○ ^{12}C
● ^{13}C

□ ^{12}C reads: assembly

□ $^{12}C/^{13}C$ reads: mapping

B

HBD DNA

LBD DNA

- ● MAG 9 (AOA: 62,1)
- ● MAG 12 (AOA: 73,3)
- ● MAG 15 (NOB: 51,2)
- ● MAG 52 (AOA: 89,2)
- ● MAG 61 (AOA: 88,2)
- ● MAG 63 (AOA: 79,1)
- ● MAG 77 (AOA: 65,7)
- ● MAG 96 (AOA: 91,2)
- ● MAG 107 (AOA: 86,3)
- ● MAG 112 (AOA: 63,1)
- ● Nitrifier virus contigs
- ● Non-nitrifier-binned contigs

C

pH 4.5: HBD ^{12}C (pink), HBD ^{13}C (red)

pH 7.5: HBD ^{12}C (light blue), HBD ^{13}C (blue)

AOA

MAG 107 *Nitrosotalea*, NT-α-1 *

MAG 9 *Nitrosotalea*, NT-α-1 *

MAG 77 *Nitrosotalea*, NT-α-1 *

MAG 61 *Nitrosotalea*, NT-α-1 *

MAG 63 CS63, NS-β-2

MAG 112 TA-21, NS-δ-2.1

MAG 96 TH1177, NS-ε-2.2 *

MAG 12 *Nitrososphaera*, NS-α-3.2.3 *

MAG 52 *Nitrososphaera*, NS-α-3.2.3 *

NOB

MAG 15 *Nitrobacter* *

Relative abundance (x 10^{-3} %)

A

pH 4.5 | pH 7.5

| | LBD | ¹²C HBD | ¹³C HBD | | LBD | ¹²C HBD | ¹³C HBD |

AOA viral contigs (n=64)

Relative abundance (ln transformed)

-2.5   0   2.5   5

B

0.5
0.1
0.05
0.01
0.005
0.001

Viruses of soil AOA

Reference virus sequences

C

027541 TerL
Nitrososphaeraceae MAG (WP_162477508)
Nitrososphaeraceae MAG (PWU79996)
Nitrososphaeraceae MAG (WP_162476599)
Nitrososphaeraceae MAG (NWG38094)
Nitrososphaera viennensis (FR725450)
Nitrososphaerales MAG (PIN82302)
Uncultured marine virus (AKH48315)
099384 TerL
Nitrososphaeraceae MAG (MPZ07264)
Nitrososphaeraceae MAG (MPZ07268)
053162 TerL
063910 TerL
061863 TerL
002326 TerL
019310 TerL
056586 TerL
066305 TerL
016734 TerL
Nitrososphaeraceae MAG (MCD6036164)
094402 TerL
Ca. Nitrosocaldaceae cavascurensis (SPC33876)

0.05
● >0.90
◐ >0.75
○ >0.50
>100x in pH 4.5
>100x in pH 7.5

D

HNH endonuclease
Zn-finger
MCO1
tail
tail
HNH endonuclease
portal
capsid
2OG-Fe(II)

contig 053596 (16.3 kb)

unaffiliated
non-archaea
non-AOA archaea
AOA
virus

MCO1 (adjacent permease)

MCO1 (no adjacent permease)

Nitrososphaera viennensis EN76 (NVIE_000600)
Ca. Nitrososphaera gargensis Ga9.2 (Ngar_c23340)
Nitrosopumilus maritimus SCM1 (Nmar_1663)
Ca. Nitrosocosmicus oleophilus MY3 (NMY3_03220)
Nitrososphaera sp. AFS (WP_162477813.1)
AOA-infecting virus contig 053596
Ca. Nitrososphaera gargensis Ga9.2 (Ngar_c24040)
Ca. Nitrosocosmicus hydrocola (WP_231129092)
Ca. Nitrosocosmicus oleophilus MY3 (NMY3_01033)
Nitrososphaera viennensis EN76 (NVIE_019250)
Nitrosopumilus maritimus SCM1 (Nmar_1354)
Nitrososphaera viennensis EN76 (NVIE_017730)
Ca. Nitrosocosmicus franklandianus C13 (NFRAN_2798)

MCO4

Ca. Nitrososphaera gargensis Ga9.2 (Ngar_c0426)
Nitrososphaera viennensis EN76 (NVIE_000530)
Ca. Nitrosocosmicus franklandianus C13 (NFRAN_1409)
Ca. Nitrosotalea devanaterrae ND1 (NDEV_1607)
Nitrosopumilus maritimus SCM1 (Nmar_1667)

NirK

0.05
● >0.95
◐ >0.90
○ >0.70