Title: *Long term maintenance of heterologous symbiont association in Acropora palmata on*

natural reefs

Running Head: Symbiont community dynamics in out-planted Acropora palmata

Holland Elder[1]*

Wyatt C. Million[1]

Erich Bartels[2]

Cory Krediet[3]

Erinn Muller[4]

Carly D. Kenkel[1]

1. University of Southern California, 3616 Trousdale Parkway, AHF 231, Los Angeles, CA 90089

2. Mote Marine Laboratory, 24244 Overseas Hwy, Summerland Key, FL 33042

3. Eckerd College, James Center for Molecular and Life Sciences 124, 4200 54th Ave., St.

Petersberg, FL 33711

4. Mote Marine Laboratory, 1600 Ken Thompson Parkway, Sarasota, FL 34236

*corresponding author

760-622-9116

hollandelder@gmail.com

Keywords: coral reefs, symbiosis, symbiont type, symbiont shift, symbiont switching, symbiont

community, restoration

Paper type: Brief Communication

Abstract

The sensitivity of reef-building coral to elevated temperature is a function of their

symbiosis with dinoflagellate algae in the family Symbiodiniaceae. Changes in the composition

of the endosymbiont community in response to thermal stress can increase coral thermal

tolerance. Consequently, this mechanism is being investigated as a human-assisted intervention

for rapid acclimation of coral in the face of climate change. Successful establishment of novel

symbioses that increase coral thermal tolerance have been demonstrated in laboratory conditions,

however it is unclear how long these heterologous relationships persist in nature. Here, we show

that Acropora palmata can form a novel symbiosis with Durusdinium spp. when reared in land-

based aquaria. We tested the stability of this heterologous relationship by outplanting clonal

replicates (ramets) of five A. palmata host genotypes to natural reefs in the lower Florida Keys.

Amplicon sequencing analysis of ITS2-type profiles revealed that the majority of surviving

ramets remained dominated by Durusdinium spp. two years after transplantation. However, 15%

of ramets, including representatives of all genotypes, did exhibit some degree of symbiont

shuffling or switching at six of eight sites, including complete takeover by site-specific strains of

the native symbiont, Symbiodinium fitti. The predominant long-term stability of the novel

symbiosis supports the potential effectiveness of symbiont modification as a management tool.

However, the finding that 6–7 year-old coral can alter symbiont community composition in the

absence of bleaching indicates that Symbiodiniaceae communities are indeed capable of greater

flexibility under ambient conditions.

Manuscript

Increasingly frequent and severe thermal stress events are the predominant threat to reef

ecosystems in the Anthropocene. Thermal stress perturbs the photosynthetic mechanism and can

result in loss or expulsion of the dinoflagellate algae in the process known as bleaching (Hoegh-

Guldberg & Smith, 1989). While persistent bleaching can result in coral death due to the loss of

autotrophic carbon contributed by the symbionts (Glynn, 1993), bleaching can also precipitate an

acclimatory change in the composition of the endosymbiont community that can subsequently

increase coral thermal tolerance (Berkelmans & van Oppen, 2006a; Cunning, Gillette, et al.,

2015). For coral hosting multiple symbiont types, changes in their relative abundance, or

'shuffling', can facilitate survival and recovery (Berkelmans & van Oppen, 2006a). The ability

for adult coral to acquire completely new symbiont types following bleaching was also recently

confirmed (Scharfenstein et al., 2022). Although this capacity for flexibility is not universal

(Goulet, 2006), and is influenced by environmental cues and physiological trade-offs (Jones &

Berkelmans, 2010), symbiont shuffling and/or switching has been proposed as a key target for

human-assisted evolution (van Oppen et al., 2015). Successful establishment of novel symbioses

that increase coral bleaching tolerance have been demonstrated in laboratory conditions (Buerger

et al., 2020), however it is unclear how long these heterologous relationships persist, especially

in natural reef environments. In a key test of the ultimate utility of this approach, we investigate

the long-term stability of a heterologous coral symbiosis in the context of reef restoration.

Acropora palmata were once one of the dominant habitat builders of shallow Caribbean

reefs (Kuffner & Toth, 2016). Unprecedented demographic declines since the mid-1970's have

resulted in extensive propagation efforts in both land and ocean-based nursery systems to both

maintain existing genetic diversity and produce biomass for outplanting efforts aimed at

restoring ecosystem structure and function (Young et al., 2012). Naturally occurring populations

of *A. palmata* are highly specific in their symbiosis with *Symbiodinium fitti*, formerly ITS2 type

A3, (Baums et al., 2014; Reich et al., 2021). Single nucleotide polymorphism (SNP) genotyping

of *ex situ* nursery-reared lines of *A. palmata*, however, revealed an almost exclusive association

with *Durusdinium spp.* (Fig S1A, Supplementary Methods). This provided a unique opportunity

to test the maintenance of this heterologous symbiotic relationship after outplanting back to the

reef environment.

In April 2018, we transplanted three clonal replicates (ramets) of five exclusively *ex situ*

nursery-reared *A. palmata* genotypes to nine reefs in the lower Florida Keys (n=27 ramets per

genet) to evaluate the long-term stability of the *Durusdinium* association on natural reefs using

periodic phenotypic monitoring and ITS2 amplicon sequencing for symbiont typing using the

SymPortal pipeline (Hume et al., 2019), (Supplemental methods, Figure 1). A random sub-

sampling of 10-12 ramets per genet confirmed that no *Symbiodinium fitti* were detectable prior to

outplanting (T0, Supplemental methods, Figure 2A). Four genets were exclusively dominated by

a *Durusdinium* ITS2 type, whereas genet 13-XK showed a consistent co-infection with a

Cladocopium ITS2 type composed of three co-occurring C1 amplicon sequence variants (Figure

2A). A PCoA using Bray-Curtis dissimilarity of T0 samples confirmed a significant effect of

genotype ($p < 0.001$) with 13-XK driving divergence (Figure 2B).

No bleaching was observed during quarterly site visits, but site-specific mortality was

evident, with 100% ramet mortality at Bahia Honda following 18 months (Supplemental

Methods, Table S2). *Durusdinium* remained the dominant ITS2 profile after two years of

outplanting (T24, Fig. 2B, C). Surviving co-infected 13-XK ramets also shuffled to *Durusdinium*

dominance across sites (Fig. 2C) and a PERMANOVA confirmed genotype was no longer a

significant factor (p = 0.764). Uptake of novel symbionts was detected, however, with site

specific patterns (PERMANOVA, p < 0.031, Fig. 2C). Acquisition of the homologous symbiont,

Symbiodinium fitti (A3) was observed at four sites, with ITS2 type profiles indicating potential

sub-species or strain-level differences between Dave's Ledge, Eastern Dry Rocks, Looe Key,

and Maryland Shoals (Fig. 2C). Two ramets of 13-X7 acquired *Cladocopium* types, including

one apparently novel strain (C66) although their relative abundance remained low (Figure 2C). A

Gerakladium (G) type profile was also detected in ramets of two different genets at two different

sites, also at low relative abundance (Fig. 2C).

The general long-term stability of the heterologous symbiosis between *A. palmata* and

Durusdinium spp. on natural reefs supports the potential effectiveness of symbiont modification

as a management intervention. Infection of coral larvae from mass spawning events with

thermally tolerant symbionts and outplanting them to degraded reefs could also be a restoration

tool (Randall et al., 2020). However, the occasional acquisition of homologous, site-specific

strains of *Symbiodinium fitti* indicates that the temporal dynamics of symbiosis establishment in

coral may be more complicated than previously thought. While corals can take up symbionts in

their larval phase, the current consensus is that the majority of symbionts are taken up after

settlement. New recruits and juvenile corals can form a variety of novel symbiont associations

(Bay et al., 2011), but this diversity is generally pruned to one numerically dominant taxon per

genus over the first few years of life (Abrego et al., 2009). Subsequent changes to the dominant

taxon occur in response to bleaching-level stress (Berkelmans & van Oppen, 2006b; Cunning,

Silverstein, et al., 2015; Scharfenstein et al., 2022).

The *A. palmata* lines used in this study were produced using individual larvae from batch

crosses in 2013/14. Although physically small, at the time of outplanting, experimental ramets

were 4-5 years old, which under normal field conditions is the age of sexual maturity for this

species (Chamberland et al., 2016). The finding that adult corals can switch their symbionts *in*

situ in the absence of bleaching suggests that acquisition and winnowing may be size-dependent

rather than age-dependent process. It is also possible that Symbiodiniaceae communities are

indeed capable of greater flexibility under ambient conditions (Silverstein et al., 2012). However,

the stability of the heterologous symbiosis in most individuals suggests that infection with a

novel symbiont type as a recruit and maintenance till maturity could produce a stable symbiosis.

Works Cited

Abrego, D., van Oppen, M. J. H., & Willis, B. L. (2009). Highly infectious symbiont dominates
initial uptake in coral juveniles. *Molecular Ecology*, *18*(16), 3518–3531.
https://doi.org/10.1111/j.1365-294X.2009.04275.x

Baums, I. B., Devlin-Durante, M. K., & Lajeunesse, T. C. (2014). New insights into the
dynamics between reef corals and their associated dinoflagellate endosymbionts from
population genetic studies. *Molecular Ecology*, *23*(17), 4203–4215.
https://doi.org/10.1111/mec.12788

Bay, L. K., Cumbo, V. R., Abrego, D., Kool, J. T., Ainsworth, T. D., & Willis, B. L. (2011).
Infection Dynamics Vary between Symbiodinium Types and Cell Surface Treatments
during Establishment of Endosymbiosis with Coral Larvae. *Diversity*, *3*, 356–374.
https://doi.org/10.3390/d3030356

Berkelmans, R., & van Oppen, M. J. H. (2006a). The role of zooxanthellae in the thermal
tolerance of corals: a "nugget of hope" for coral reefs in an era of climate change.
Proceedings. Biological Sciences / The Royal Society, *273*(1599), 2305–2312.
https://doi.org/10.1098/rspb.2006.3567

Berkelmans, R., & van Oppen, M. J. H. (2006b). The role of zooxanthellae in the thermal tolerance of corals: A "nugget of hope" for coral reefs in an era of climate change. *Proceedings of the Royal Society B: Biological Sciences*, *273*(1599), 2305–2312. https://doi.org/10.1098/rspb.2006.3567

Buerger, P., Alvarez-Roa, C., Coppin, C. W., Pearce, S. L., Chakravarti, L. J., Oakeshott, J. G., Edwards, O. R., & van Oppen, M. J. H. (2020). Heat-evolved microalgal symbionts increase coral bleaching tolerance. *Sciences Advances*, *6*. https://doi.org/eaba2498

Chamberland, V. F., Petersen, D., Latijnhouwers, K., Snowden, S., Mueller, B., & Vermeij, M. (2016). Four-year-old Caribbean Acropora colonies reared from field-collected gametes are sexually mature. *Bulletin of Marine Science*, *92*(2). https://doi.org/10.5343/105

Cunning, R., Gillette, P., Capo, T., Galvez, K., & Baker, A. C. (2015). Growth tradeoffs associated with thermotolerant symbionts in the coral Pocillopora damicornis are lost in warmer oceans. *Coral Reefs*, *34*(1), 155–160. https://doi.org/10.1007/s00338-014-1216-4

Cunning, R., Silverstein, R. N., & Baker, A. C. (2015). Investigating the causes and consequences of symbiont shuffling in a multi-partner reef coral symbiosis under environmental change. *Proceedings of the Royal Society B*, *282*, 20141725.

Glynn, P. W. (1993). Coral reef bleaching: ecological perspectives. *Coral Reefs*, *12*, 1–17. https://doi.org/10.1007/BF00303779

Goulet, T. L. (2006). Most corals may not change their symbionts. *Marine Ecology Progress Series*, *321*, 1–7.

Hoegh-Guldberg, O., & Smith, J. G. (1989). The effect of sudden changes in temperature, light and salinity on the population density and export of zooxanthellae from the reef corals Stylophora pistillata (Esper) and Seriatopora hysterix (Dana). *Journal of Experimental Marine Biology and Ecology*, *129*, 279–303.

Hume, B. C. C., Smith, E. G., Ziegler, M., Warrington, H. J. M., Burt, J. A., LaJeunesse, T. C., Wiedenmann, J., & Voolstra, C. R. (2019). SymPortal: A novel analytical framework and platform for coral algal symbiont next-generation sequencing ITS2 profiling. *Molecular Ecology Resources*, *19*(4), 1063–1080. https://doi.org/10.1111/1755-0998.13004

Jones, A., & Berkelmans, R. (2010). Potential Costs of Acclimatization to a Warmer Climate : Growth of a Reef Coral with Heat Tolerant vs . Sensitive Symbiont Types. *PloS One*, *5*(5), e10437. https://doi.org/10.1371/journal.pone.0010437

Kuffner, I. B., & Toth, L. T. (2016). A geological perspective on the degradation and conservation of western Atlantic coral reefs. In *Conservation biology : the journal of the Society for Conservation Biology* (Vol. 30, Issue 4, pp. 706–715). Blackwell Publishing Inc. https://doi.org/10.1111/cobi.12725

Randall, C. J., Negri, A. P., Quigley, K. M., Foster, T., Ricardo, G. F., Webster, N. S., Bay, L. K., Harrison, P. L., Babcock, R. C., & Heyward, A. J. (2020). Sexual production of corals

for reef restoration in the Anthropocene. *Marine Ecology Progress Series*, *635*(February),
203–232. https://doi.org/10.3354/MEPS13206

Reich, H. G., Kitchen, S. A., Stankiewicz, K. H., Devlin-Durante, M., Fogarty, N. D., & Baums,
I. B. (2021). Genomic variation of an endosymbiotic dinoflagellate (Symbiodinium 'fitti')
among closely related coral hosts. *Molecular Ecology*, *30*(14), 3500–3514.
https://doi.org/10.1111/mec.15952

Scharfenstein, H. J., Chan, W. Y., Buerger, P., Humphrey, C., & van Oppen, M. J. H. (2022).
Evidence for de novo acquisition of microalgal symbionts by bleached adult corals. *ISME*
Journal. https://doi.org/10.1038/s41396-022-01203-0

Silverstein, R. N., Correa, A. M. S., & Baker, A. C. (2012). Specificity is rarely absolute in
coral–algal symbiosis: Implications for coral response to climate change. *Proceedings of the*
Royal Society B: Biological Sciences, *279*(1738), 2609–2618.
https://doi.org/10.1098/rspb.2012.0055

van Oppen, M. J. H., Oliver, J. K., Putnam, H. M., & Gates, R. D. (2015). Building coral reef
resilience through assisted evolution. *Proceedings of the National Academy of Sciences*,
112(8), 2307–2313. https://doi.org/10.1073/pnas.1422301112

Young, C. N., Schopmeyer, S. A., & Lirman, D. (2012). A review of reef restoration and Coral
propagation using the threatened genus Acropora in the Caribbean and western Atlantic. In
Bulletin of Marine Science (Vol. 88, Issue 4, pp. 1075–1098).
https://doi.org/10.5343/bms.2011.1143

Figure 1. Experimental setup showing genotypes subsampled at time point 0 (T0), the locations

of outplanting sites, the locations of water quality monitoring stations, and the genotype array

strategy at each site. All surviving corals were sampled at time point 24 (T24).

Figure 2. Relative abundance and PCoA plots showing that symbiont community did not

significantly change after two years of outplanting. (**A.**) The relative abundance plot shows the

ITS2 symbiont profiles of each individual coral grouped by genotype that was subsampled at T0.

Each bar represents an individual and each is labeled with either their ex situ nursery frag label

or their outplant site ID. Color represents symbiont type. (**B.**) The relative abundance plot shows

the ITS2 symbiont profiles of each individual coral grouped by genotype that was sampled at

T24. Each bar represents an individual. They are grouped by site, then genotype, and then frag

id. Asterisks (*) identify individuals that were also sampled at T0. Color represents symbiont

type. (**C.**) A PcoA plot that shows that symbiont profiles did not significantly change after two

214   years of being outplanted. Color represents coral genotype and shape represents site. The ex situ

215   nursery and T0 is included as a site.