A pan-cancer analysis of the oncogenic role of COVID-19 risk gene Leucine Zipper Transcription Factor-Like Protein 1 (LZTFL1) in human tumors

Jihao Mo[1]*, Zhenzhen Zhang[2]*,Daping Wang[2,3,4], Meimei Wang[5#], Lei Wang[6#]

Department of Medical Laboratory, Luoyang Orthopedic Hospital of Henan Province, Orthopedic Institute of Henan Province, Henan, 471000, China

Department of Biomedical Engineering, Southern University of Science and Technology, Shenzhen, 518055, China

Department of Orthopedics, Shenzhen Intelligent Orthopaedics and Biomedical Innovation Platform, Guangdong Artificial Intelligence Biomedical Innovation Platform, Shenzhen Second People's Hospital, The First Affiliated Hospital of Shenzhen University Health Science Center, Shenzhen, 518035, China

Guangzhou Medical University, Guangzhou, Guangdong Province, 511436, China

Department of Pathophysiology, School of Basic Medical Science, Anhui Medical University, No. 81, Mei-Shan Road, Hefei, 230032, Anhui, China

Department Physiology and Biophysics, University of Miami Miller School of Medicine, Miami, Florida, 33136, US

* These authors contributed equally to the project.

Co-responding author.

Abstract

Population-based studies showed that COVID-19 infection causes higher death rate in cancer patients. However, the molecular mechanism of COVID-19 with cancer is still largely unknown. Here we analyzed the Leucine Zipper Transcription Factor-Like Protein 1 (LZTFL1) which is the most significant gene associated with COVID-19. First, we explored the potential oncogenic roles of LZTFL1 through transcriptome data from The Cancer Genome Atlas (TCGA) and Gene Expression Omnibus (GEO) database. LZTFL1 is significantly low expressed in 11 of 34 kinds of cancers we analyzed. Consistent with the mRNA expression data, the protein expression of LZTFL1 in lung adenocarcinoma (LUAD), clear cell renal cell carcinoma (ccRCC), Uterine corpus endometrial carcinoma (UCEC), and ovarian cancer (OV) patients are significantly decreased compared to healthy tissues. The survival analysis from the Kidney renal clear cell carcinoma (KIRC), Rectum adenocarcinoma (READ), and Uveal Melanoma (UVM), the LZTFL1 high expression group have a significantly higher survival rate compared to the low expression group. Taken together, LZTFL1 acts as a cancer suppressor gene for several cancers. Moreover, LZTFL1 expression was associated with the cancer-associated fibroblast infiltration in several tumors including Bladder Urothelial Carcinoma (BLCA), Breast invasive carcinoma (BRCA), Esophageal carcinoma (ESCA), Head and Neck squamous cell carcinoma (HNSC), Lung squamous cell carcinoma (LUSC), and Pancreatic adenocarcinoma (PAAD). Gene ontology analysis showed that cilium organization, positive regulation of establishment of protein localization to telomere and SRP-dependent cotranslational protein targeting to the membrane were involved in the function mechanisms related to LZTFL1. Our studies offer a relatively comprehensive understanding of the oncogenic roles of LZTFL1 across different kinds of tumors.

Keywords:

LZTFL1, cancer, COVID-19, transcriptome, GWAS

Introduction

As of Aug 15, 2022, 590 million COVID-19 infection cases and 6.4 million deaths reported worldwide. The SPIKE protein of COVID-19 binds to the ACE2, then cleavaged by TMPRSS2 to infect the mammalian respiratory cells(Lan et al., 2020, Hoffmann et al., 2020). The new discovered COVID-19 target protein named Neuropilin 1, might handing the key to a new door for the virus to replicate in the human airway cells(Cantuti-Castelvetri et al., 2020). A survey published by the American Cancer Society Cancer Action Network showed that COVID-19 pandemic challenges more to cancer patients and survivors, including delays and cancellations of health care services, economic challenges affecting their ability to pay and so on(Network, 2020). Recent publications showed that the immunosuppressed status of some cancer patients increases their risk of infection and result in serious complications compared with the general population(Al-Quteimat and Amer, 2020). The genomics mutations related to cancer are well-studied compared to other diseases. However, how the genetic mutations related to COVID-19 is still largely unknown.

Genome wide Association Study of Severe and critical illness in COVID-19 patient samples indicated that Leucine Zipper Transcription Factor-Like Protein 1 (LZTFL1), tyrosine kinase 2 (TYK2), dipeptidyl peptidase 9 (DPP9), Interferon Alpha and Beta Receptor Subunit 2 (IFNAR2), C-C Motif Chemokine Receptor 2 (CCR2), and Alpha 1-3-N-Acetylgalactosaminyltransferase and Alpha 1-3-Galactosyltransferase (ABO) genes are closely associated with severe COVID-19(Severe Covid et al., 2020, Pairo-Castineira et al., 2021). There is an urgent need to investigate the role of these genomics mutations in cancer. Among all these genes, LZTFL1 is the most significant gene associated with COVID-19. LZTFL1 is known to interact with Bardet-Biedl Syndrome (BBS) proteins(Marion et al., 2012). Recent evidence has shown that LZTFL1 was a novel cancer suppressor gene associated with certain kinds of cancers(Wei et al., 2010). Immunohistochemical (IHC) staining on 311 paired normal/cancer tissue arrays suggested that LZTFL1 suppresses gastric cancer cell migration(Wang et al., 2014). Evidence from breast cancer patients indicated that microRNA-21 promotes breast cancer proliferation and metastasis by suppressing LZTFL1(Wang et al.,

2019). Besides, LZTFL1 suppresses lung tumorigenesis by maintaining differentiation of lung

epithelial cells(Wei et al., 2016). However, there is still no pan-cancer evidence on the

relationship between LZTFL1 and all the tumour types based on bid clinical data analysis.

Here we conduct a pan-cancer analysis of COVID-19 associated gene LZTFL1 using TCGA

and GEO databases. We investigated the potential molecular mechanism of LZTFL1 in the

pathogenesis and clinical prognosis of different cancers by analysing the gene expression,

survival status, DNA methylation, genetic alteration, and relevant cellular pathway. To study

the function of LZTFL1 as a transcription factor, we predicted the top 100 downstream genes

of LZTFL1 and analysed their expression in COVID-19 related RNA-sequencing data set.

Furthermore, we analysed the function, localization, and microRNA target of LZTFL1

downstream genes.

Materials and methods

Genetic alteration analysis

The genetic alteration characteristics of LZTFL1 in the TCGA database were analysed using

the cBioPortal (https://www.cbioportal.org/)(Gao et al., 2013). Briefly, "LZTFL1" was entered

for queries of genetic alteration characteristics. The mutated sites were labelled on the

localization on the amino acid position on protein coding region with several mutations

observed. The alteration frequency from TCGA data set was calculated with mutation,

amplification, and deep deletion information.

Structure prediction

The structure of LZTFL1 was predicted based on the NCBI Reference sequence

NP_065080.1 using SWISS-MODEL (https://swissmodel.expasy.org/)(Waterhouse et al.,

2018). The structure of amino acid sequence 1-64 was built based on template SMTL ID:

5wo3.1(Horowitz et al., 2016). The structure of amino acid sequence 148-222 was built based

on template SMTL ID: 3o0z.1(Tu et al., 2011).

mRNA expression analysis

The systematical analysis of LZTFL1 mRNA expressions between tumour and adjacent

normal tissues in TCGA database were conducted using "Gene_DE" module of tumor

immune estimation resource, version 2 (TIMER2, http://timer.cistrome.org/)(Li et al., 2017).

The box plot with tumour and adjacent normal tissues was labeled gray. The statistical

significance computed by the Wilcoxon test is annotated by the number of stars (*: p-value <

0.05; **: p-value < 0.01; ***: p-value < 0.001).

Protein expression analysis

The systematical analysis of LZTFL1 expressions between tumour and adjacent normal

tissues in TCGA database were conducted using Clinical Proteomic Tumour Analysis

Consortium (CPTAC) of UALCAN portal (http://ualcan.path.uab.edu/analysis-prot.html)(Chen

et al., 2019). The LZTFL1 expression of breast cancer, ovarian cancer, Colon cancer, Clear

cell renal cell carcinoma, Uterine corpus endometrial carcinoma, and lung adenocarcinoma

tumor and normal tissues were analyzed, respectively. The expression of LZTFL1 in each

individual cancer stages cancer type with further investigated.

Survival analysis

The survival analysis of LZTFL1 between high and low expression groups in TCGA database

was conducted using "survival map" module of GEPIA2 (http://gepia2.cancer-

pku.cn/#survival)(Tang et al., 2019). The "overall survival" was used as the method. The

group cutoff was set as the median. The cutoff- high and cutoff-low were set as 50%. The

log-rank test with a 95% confidence interval was used in the analysis of hypothesis test.

Immune infiltration analysis

The Immune infiltration analysis of LZTFL1 in EPIC, MCPCOUNTER, XCELL, and TIDE

database was conducted using "Immune_Gene" module of TIMER2(Li et al., 2017).The

cancer-associated fibroblasts were selected for analysis. The P value and correlation value

were calculated by Spearman's rank after purity adjustment. The heatmap and scatter plot

were used to visualize the data.

Gene ontology analysis

The proteins which interacted with LZTFL1 in *homo sapiens* were investigated using the

STRING tool (https://string-db.org/)(Szklarczyk et al., 2019). The edges indicate both

functional and physical protein associations and the line thickness indicates the strength of

data support. The number of interactors to show for the first and second shell were set as no

more than 20. The top 100 genes that interact with LZTFL1 based on co-expression were

analysed with Harmonizome (https://maayanlab.cloud/Harmonizome/)(Rouillard et al., 2016).

The list of genes was further analysed in COVID-19 related GEO data sets using the Enrichr

tool (https://maayanlab.cloud/Enrichr/)(Chen et al., 2013). The biological function, subcellular

localization, and miRNA regulation of LZTFL1 were visualized using a heatmap.

Results

Genetic alteration of LZTFL1 in cancer

In this study, we analysed the mutation site of LZTFL1 in the amino acid sequence. As shown

in Fig. 1A, there are totally of 41 mutations observed in the LZTFL1 coding region. The most

observed mutation point is R19. R19C was detected in rectal adenocarcinoma (READ) patient

and R19H were detected in uterine endometrioid carcinoma (UCEC) or mucinous

adenocarcinoma of the colon and rectum (COAD) patient based on the TCGA database.

Other mutations observed near this location is R22G/H, K24N, and R28K/S. The Leucine

Zipper (Leu_zip) domain locates at 20-294 amino acids. These mutations located at 19, 22,

24, and 28 are the starting point of the Leu_zip domain. Another location with multiple

mutations accumulated at the binding region of the zipper structure. These mutations are

E148K, E160D, E162D, and E172D/K. All the mutations above shortened the length of side

chain. The most possible explanation is that these mutations altered the formation of homo

dimer of the leucine zipper in the beginning and the closely localized zipper position, further

affected the function of LZTFL1 as a transcription factor (Fig 1B&C).

We further analysed the genetic alteration of LZTFL1 in different tumour samples from TCGA

database. Most of the LZTFL1 mutations are deep deletions. The highest alteration frequency

of LZTFL1 (around 4%) appears for lymphoid neoplasm diffuse large B-cell lymphoma (DLBC)

patients (Fig 1D). Only deep deletions are found in DLBC, ccRCC (clear cell renal cell

carcinoma), sarcoma, mesothelioma, thymoma, and cervical patients. Only mutations are

found uterine, adenoid cystic carcinoma (ACC), papillary renal cell carcinoma (pRCC), and

lung adeno patients. Only amplification found in coulrophobe renal cell carcinoma (RCC) and

pheochromocytoma/paraganglioma (PCPG) patients.

Gene and protein expression of LZTFL1 in cancer

We further analyzed the expression of LZTFL1 in different types of cancers and paired non-

tumor tissues in TCGA database. As shown in Fig.2A, LZTFL1 shows significantly increased

expression in cholangiocarcinoma (CHOL), esophageal carcinoma (ESCA), head and neck

squamous cell carcinoma (HNSC)-HPV+, and STAD (stomach adenocarcinoma) tumor

groups. However, LZTFL1 shows significantly decreased expression in breast invasive

carcinoma (BRCA), cervical squamous cell carcinoma and endocervical adenocarcinoma

(CESC), colon adenocarcinoma (COAD), kidney Chromophobe (KICH), kidney renal clear cell

carcinoma (KIRC), kidney renal papillary cell carcinoma (KIRP), liver hepatocellular

carcinoma (LIHC), lung adenocarcinoma (LUAD), lung squamous cell carcinoma (LUSC),

prostate adenocarcinoma (PRAD), skin cutaneous melanoma (SKCM),   thyroid carcinoma

(THCA), and UCEC (uterine Corpus Endometrial Carcinoma) tumor groups. To conclude, the

mRNA expression of LZTFL1 plays different roles in different cancer types.

We further analysed the protein expression data of LZTFL1 in CPTAC database. As shown in

Fig 2B, the protein expression of LZTFL1 is significantly decreased in LUAD, Clear cell RCC,

and UCEC patient samples. But the protein expression of LZTFL1 is significantly increased in

ovarian cancer patient samples. To further investigate which stages of these cancers are

correlated with LZTFL1 protein expression, we separated the data based on the stages (Fig.

2C). Patients with LUAD and clear cell RCC have decreased LZTFL1 protein expression in

stages 1-3. Patients with UCEC have decreased LZTFL1 protein expression in stages 1 and

3. Patients with ovarian cancer have increased LZTFL1 protein expression in stage 3. These

data indicated that the LZTFL1 protein expression is differentially associated with the

prognosis of patients with different cancer types.

Survival analysis of LZTFL1 in cancer

To investigate the correlation between the expression of LZTFL1 and the prognosis of

patients with different kinds of tumours, we analysed the survival by dividing the cancer cases

into high and low-LZTFL1 expression groups. The datasets from TCGA were used in this

analysis. Based on the log rank P-value, patients with high LZTFL1 expression were linked to

a better prognosis of overall survival for cancers of KIRC ($P < 0.0001$), READ ($P = 0.0039$),

and UVM ($P = 0.0180$). patients with low LZTFL1 expression were linked to a better prognosis

for cancers of Brain Lower Grade Glioma (LGG, $P = 0.0100$) (Fig. 3).

Immune infiltration analysis of LZTFL1 in cancer

To analysis the immune infiltration of LZTFL1 across all kinds of tumours, the immune cells of

cancer-associated fibroblasts were selected. The XCELL, MCPCOUNTER, TIDE and EPIC

algorithms were applied to investigate the correlation between the cancer-associated

fibroblasts and LZTFL1 gene expression. The statistical positive correlation of LZTFL1

expression and infiltration value of cancer-associated fibroblasts for the tumours of BLCA,

BRCA, BRCA-basal, ESCA, HNSC, HNSC-HPV⁻, LUSC, and PAAD were observed (Fig. 4A).

The scatterplot showed the related type of tumour using one algorithm (Fig. 4B).

Enrichment and downstream gene analysis of LZTFL1-related genes

To further investigate the molecular mechanism of the LZTFL1 gene in tumorigenesis, we

analysed the LZTFL1 related genes by protein-protein interaction. Top 50 LZTFL1 related

genes were enriched based on the co-localization, function, and co-expression analysis using

STRING (Fig. 5A). The top 50 genes are enriched in cilium organization, SRP-dependent co-

translational protein targeting to membrane, and positive regulation of establishment of

protein localization to the telomere. The target genes of LZTFL1 are found in transcriptome

database of Down-regulated by SARS-CoV-2 in A549-ACE2 treated cells, Down-regulated

genes from COVID-19 infected bronchoalveolar lavage from patients, MERS-CoV 38

Literature-Associated Genes from Geneshot, SARS perturbation Down Genes PBMCs,

Down-regulated by SARS-CoV-2 infection in Vero E6, SARS Perturbation Down Genes

Mouse Lung, SARS coronavirus endoRNAse from Virus-Host, MERS-Cov 6 Predicted

Kinases Genes, Down Genes from SARS infected Mouse Lung, and SARS-CoV perturbation

Down Genes Vero E6 (Fig.5B). The enriched pathways of LZTFL1 downstream genes are

Collecting duct acid secretion, Vibrio cholerae infection, Synaptic vesicle cycle, Epithelial cell

signalling in Helicobacter pylori infection, Vitamin digestion and absorption, Rheumatoid

arthritis, RNA degradation, Oxidative phosphorylation, Sphingolipid metabolism, and mTOR

signalling pathway (Fig.5C). The enriched cellular organelles of LZTFL1 downstream genes

are Golgi trans cisterna, endosome membrane, filopodium, lysosomal membrane, tertiary

granule membrane, integral component of ER, specific granule membrane, actin cytoskeleton,

centrosome, and integral component of Golgi (Fig.5D). The top 10 enriched miRNAs are hsa-

miR-5683, hsa-miR-519d-3p, hsa-miR-20b-5p, hsa-miR-758-5p, hsa-miR-526b-3p, hsa-miR-

7852-3p, hsa-miR-106a-5p, hsa-miR-17-5p, hsa-miR-20a-5p, and hsa-miR-4640-3p (Fig.5E).

Discussion

LZTFL1 have been known as Bardet-Biedl Syndrome (BBS) proteins and smoothened

trafficking regulator for around ten years(Marion et al., 2012). LZTFL1 interacts with a BBS

protein complex and regulates ciliary trafficking of this complex. A homozygous 5 bp deletion

in LZTFL1 was identified in BBS patients, which causes no LZTFL1 transcript or protein could

be detected in the patient's fibroblasts(Marion et al., 2012). LZTFL1 locals in cytoplasm

helping with the regulation of BBS protein complex ciliary trafficking.

How LZTFL1 works in each kind of cancer types are still largely unknown. The current

hypothesis suggested that by interacting with E-cadherin and the actin cytoskeleton, LZTFL1

regulating the transition of epithelial cells to mesenchymal cells(Li et al., 2019). Moreover,

Wang et al reported that LZTFL1 suppresses gastric cancer cell migration and invasion

through regulating nuclear translocation of β-catenin(Wang et al., 2014). By compare the

protein expression level with normal tissues by immunohistochemistry, Further studies

showed that overexpress LZTFL1 inhibits anchorage-independent cell growth *in vivo* and *in*

vitro(Wei et al., 2016). Our studies showed that LZTFL1 protein expression was closely

related to lung adenocarcinoma, clear cell RCC, UCEC, and ovarian cancer. However,

different from the ICH results, LZTFL1 protein expression was not changed in lung cancer

patients, but significantly increased in ovarian cancer patients' samples compare to normal

tissues. Since the Qun et al's studies use epithelial cell of ovarian from 10 patients and lung

epithelial cell samples from 9 patients, the CPTAC database have 125 samples from breast

and 100 samples from whole ovarian cancer samples, the different results might come from

the difference of cell type and sample numbers.

As a transcription factor, the binding region of LZTFL1 is vitally important for regulating

downstream gene expression. Mutations in the binding region of LZTFL1 have been reported

related to different cancer types as well as COVID-19. Recent studies investigated that the

rs17713054 risk allele of LZTFL1 generates a CCAAT/enhancer binding protein beta

motif(Downes et al., 2021). SNP allele of LZTFL1, rs17713054G>A, regulates the expression

of LZTFL1 in lung samples by interacting with the LZTFL1 promoter. Further studies from lung

samples of COVID-19 patients showed that a signal of epithelia-mesenchymal transition,

which is essential for the innate immune response in lung epithelial cells(Downes et al., 2021).

These findings are consistent with our results that most of the downstream genes regulated

by LZTFL1 were down regulated in SARS-CoV-2 infected lung cells, Vero cells, mouse lung

samples, and bronchoalveolar lavage samples from COVID-19 patients. Our hypothesis is

that LZTFL1 serves as a protective role against the COVID-19 infection as well as the survival

of KIRC, READ, UVM cancer patients (Fig.6).

Acknowledgments

This work was funded by Shenzhen Fund for Guangdong Provincial High level Clinical Key

Specialties (No. SZGSP007; SZSM201412015).

Referecnces

AL-QUTEIMAT, O. M. & AMER, A. M. 2020. The Impact of the COVID-19 Pandemic on Cancer Patients. *Am J Clin Oncol,* 43**,** 452-455.

CANTUTI-CASTELVETRI, L., OJHA, R., PEDRO, L. D., DJANNATIAN, M., FRANZ, J., KUIVANEN, S., VAN DER MEER, F., KALLIO, K., KAYA, T., ANASTASINA, M., SMURA, T., LEVANOV, L., SZIROVICZA, L., TOBI, A., KALLIO-KOKKO, H., OSTERLUND, P., JOENSUU, M., MEUNIER, F. A., BUTCHER, S. J., WINKLER, M. S., MOLLENHAUER, B., HELENIUS, A., GOKCE, O., TEESALU, T., HEPOJOKI, J., VAPALAHTI, O., STADELMANN, C., BALISTRERI, G. & SIMONS, M. 2020. Neuropilin-1 facilitates SARS-CoV-2 cell entry and infectivity. *Science,* 370**,** 856-860.

CHEN, E. Y., TAN, C. M., KOU, Y., DUAN, Q., WANG, Z., MEIRELLES, G. V., CLARK, N. R. & MA'AYAN, A. 2013. Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool. *BMC Bioinformatics,* 14**,** 128.

CHEN, F., CHANDRASHEKAR, D. S., VARAMBALLY, S. & CREIGHTON, C. J. 2019. Pan-cancer molecular subtypes revealed by mass-spectrometry-based proteomic characterization of more than 500 human cancers. *Nat Commun,* 10**,** 5679.

DOWNES, D. J., CROSS, A. R., HUA, P., ROBERTS, N., SCHWESSINGER, R., CUTLER, A. J., MUNIS, A. M., BROWN, J., MIELCZAREK, O., DE ANDREA, C. E., MELERO, I., CONSORTIUM, C. O.-M.-O. B. A., GILL, D. R., HYDE, S. C., KNIGHT, J. C., TODD, J. A., SANSOM, S. N., ISSA, F., DAVIES, J. O. J. & HUGHES, J. R. 2021. Identification of LZTFL1 as a candidate effector gene at a COVID-19 risk locus. *Nat Genet,* 53**,** 1606-1615.

GAO, J., AKSOY, B. A., DOGRUSOZ, U., DRESDNER, G., GROSS, B., SUMER, S. O., SUN, Y., JACOBSEN, A., SINHA, R., LARSSON, E., CERAMI, E., SANDER, C. & SCHULTZ, N. 2013. Integrative analysis of complex cancer genomics and clinical profiles using the cBioPortal. *Sci Signal,* 6**,** pl1.

HOFFMANN, M., KLEINE-WEBER, H., SCHROEDER, S., KRUGER, N., HERRLER, T., ERICHSEN, S., SCHIERGENS, T. S., HERRLER, G., WU, N. H., NITSCHE, A., MULLER, M. A., DROSTEN, C. & POHLMANN, S. 2020. SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor. *Cell,* 181**,** 271-280 e8.

HOROWITZ, S., SALMON, L., KOLDEWEY, P., AHLSTROM, L. S., MARTIN, R., QUAN, S.,

AFONINE, P. V., VAN DEN BEDEM, H., WANG, L., XU, Q., TRIEVEL, R. C., BROOKS, C. L.,

3RD & BARDWELL, J. C. 2016. Visualizing chaperone-assisted protein folding. *Nat Struct*

Mol Biol, 23**,** 691-7.

LAN, J., GE, J., YU, J., SHAN, S., ZHOU, H., FAN, S., ZHANG, Q., SHI, X., WANG, Q.,

ZHANG, L. & WANG, X. 2020. Structure of the SARS-CoV-2 spike receptor-binding domain

bound to the ACE2 receptor. *Nature,* 581**,** 215-220.

LI, S., LI, J. & YU, Z. 2019. Tumor suppressive functions of LZTFL1 in hepatocellular

carcinoma. *Onco Targets Ther,* 12**,** 5537-5544.

LI, T., FAN, J., WANG, B., TRAUGH, N., CHEN, Q., LIU, J. S., LI, B. & LIU, X. S. 2017.

TIMER: A Web Server for Comprehensive Analysis of Tumor-Infiltrating Immune Cells.

Cancer Res, 77**,** e108-e110.

MARION, V., STUTZMANN, F., GERARD, M., DE MELO, C., SCHAEFER, E.,

CLAUSSMANN, A., HELLE, S., DELAGUE, V., SOUIED, E., BARREY, C., VERLOES, A.,

STOETZEL, C. & DOLLFUS, H. 2012. Exome sequencing identifies mutations in LZTFL1, a

BBSome and smoothened trafficking regulator, in a family with Bardet--Biedl syndrome with

situs inversus and insertional polydactyly. *J Med Genet,* 49**,** 317-21.

NETWORK, A. C. S. C. A. 2020. COVID-19 Pandemic Ongoing Impact on Cancer Patients

and Survivors: Survey Findings Summary.

PAIRO-CASTINEIRA, E., CLOHISEY, S., KLARIC, L., BRETHERICK, A. D., RAWLIK, K.,

PASKO, D., WALKER, S., PARKINSON, N., FOURMAN, M. H., RUSSELL, C. D., FURNISS,

J., RICHMOND, A., GOUNTOUNA, E., WROBEL, N., HARRISON, D., WANG, B., WU, Y.,

MEYNERT, A., GRIFFITHS, F., OOSTHUYZEN, W., KOUSATHANAS, A., MOUTSIANAS, L.,

YANG, Z., ZHAI, R., ZHENG, C., GRIMES, G., BEALE, R., MILLAR, J., SHIH, B., KEATING,

S., ZECHNER, M., HALEY, C., PORTEOUS, D. J., HAYWARD, C., YANG, J., KNIGHT, J.,

SUMMERS, C., SHANKAR-HARI, M., KLENERMAN, P., TURTLE, L., HO, A., MOORE, S. C.,

HINDS, C., HORBY, P., NICHOL, A., MASLOVE, D., LING, L., MCAULEY, D.,

MONTGOMERY, H., WALSH, T., PEREIRA, A. C., RENIERI, A., GEN, O. I.,

INVESTIGATORS, I. C., INITIATIVE, C.-H. G., ANDME, I., INVESTIGATORS, B., GEN, C. I.,

SHEN, X., PONTING, C. P., FAWKES, A., TENESA, A., CAULFIELD, M., SCOTT, R.,

ROWAN, K., MURPHY, L., OPENSHAW, P. J. M., SEMPLE, M. G., LAW, A., VITART, V.,

WILSON, J. F. & BAILLIE, J. K. 2021. Genetic mechanisms of critical illness in COVID-19.

Nature, 591**,** 92-98.

ROUILLARD, A. D., GUNDERSEN, G. W., FERNANDEZ, N. F., WANG, Z., MONTEIRO, C.

D., MCDERMOTT, M. G. & MA'AYAN, A. 2016. The harmonizome: a collection of processed

datasets gathered to serve and mine knowledge about genes and proteins. *Database*

(Oxford), 2016.

SEVERE COVID, G. G., ELLINGHAUS, D., DEGENHARDT, F., BUJANDA, L., BUTI, M.,

ALBILLOS, A., INVERNIZZI, P., FERNANDEZ, J., PRATI, D., BASELLI, G., ASSELTA, R.,

GRIMSRUD, M. M., MILANI, C., AZIZ, F., KASSENS, J., MAY, S., WENDORFF, M.,

WIENBRANDT, L., UELLENDAHL-WERTH, F., ZHENG, T., YI, X., DE PABLO, R.,

CHERCOLES, A. G., PALOM, A., GARCIA-FERNANDEZ, A. E., RODRIGUEZ-FRIAS, F.,

ZANELLA, A., BANDERA, A., PROTTI, A., AGHEMO, A., LLEO, A., BIONDI, A.,

CABALLERO-GARRALDA, A., GORI, A., TANCK, A., CARRERAS NOLLA, A., LATIANO, A.,

FRACANZANI, A. L., PESCHUCK, A., JULIA, A., PESENTI, A., VOZA, A., JIMENEZ, D.,

MATEOS, B., NAFRIA JIMENEZ, B., QUEREDA, C., PACCAPELO, C., GASSNER, C.,

ANGELINI, C., CEA, C., SOLIER, A., PESTANA, D., MUNIZ-DIAZ, E., SANDOVAL, E.,

PARABOSCHI, E. M., NAVAS, E., GARCIA SANCHEZ, F., CERIOTTI, F., MARTINELLI-

BONESCHI, F., PEYVANDI, F., BLASI, F., TELLEZ, L., BLANCO-GRAU, A., HEMMRICH-

STANISAK, G., GRASSELLI, G., COSTANTINO, G., CARDAMONE, G., FOTI, G., ANELI, S.,

KURIHARA, H., ELABD, H., MY, I., GALVAN-FEMENIA, I., MARTIN, J., ERDMANN, J.,

FERRUSQUIA-ACOSTA, J., GARCIA-ETXEBARRIA, K., IZQUIERDO-SANCHEZ, L.,

BETTINI, L. R., SUMOY, L., TERRANOVA, L., MOREIRA, L., SANTORO, L., SCUDELLER,

L., MESONERO, F., ROADE, L., RUHLEMANN, M. C., SCHAEFER, M., CARRABBA, M.,

RIVEIRO-BARCIELA, M., FIGUERA BASSO, M. E., VALSECCHI, M. G., HERNANDEZ-

TEJERO, M., ACOSTA-HERRERA, M., D'ANGIO, M., BALDINI, M., CAZZANIGA, M.,

SCHULZKY, M., CECCONI, M., WITTIG, M., et al. 2020. Genomewide Association Study of

Severe Covid-19 with Respiratory Failure. *N Engl J Med,* 383**,** 1522-1534.

SZKLARCZYK, D., GABLE, A. L., LYON, D., JUNGE, A., WYDER, S., HUERTA-CEPAS, J.,

SIMONOVIC, M., DONCHEVA, N. T., MORRIS, J. H., BORK, P., JENSEN, L. J. & MERING,

C. V. 2019. STRING v11: protein-protein association networks with increased coverage,

supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res,*

47**,** D607-D613.

TANG, Z., KANG, B., LI, C., CHEN, T. & ZHANG, Z. 2019. GEPIA2: an enhanced web server

for large-scale expression profiling and interactive analysis. *Nucleic Acids Res,* 47**,** W556-

W560.

TU, D., LI, Y., SONG, H. K., TOMS, A. V., GOULD, C. J., FICARRO, S. B., MARTO, J. A.,

GOODE, B. L. & ECK, M. J. 2011. Crystal structure of a coiled-coil domain from human

ROCK I. *PLoS One,* 6**,** e18080.

WANG, H., TAN, Z., HU, H., LIU, H., WU, T., ZHENG, C., WANG, X., LUO, Z., WANG, J.,

LIU, S., LU, Z. & TU, J. 2019. microRNA-21 promotes breast cancer proliferation and

metastasis by targeting LZTFL1. *BMC Cancer,* 19**,** 738.

WANG, L., GUO, J., WANG, Q., ZHOU, J., XU, C., TENG, R., CHEN, Y., WEI, Q. & LIU, Z. P.

2014. LZTFL1 suppresses gastric cancer cell migration and invasion through regulating

nuclear translocation of beta-catenin. *J Cancer Res Clin Oncol,* 140**,** 1997-2008.

WATERHOUSE, A., BERTONI, M., BIENERT, S., STUDER, G., TAURIELLO, G.,

GUMIENNY, R., HEER, F. T., DE BEER, T. A. P., REMPFER, C., BORDOLI, L., LEPORE, R.

& SCHWEDE, T. 2018. SWISS-MODEL: homology modelling of protein structures and

complexes. *Nucleic Acids Res,* 46**,** W296-W303.

WEI, Q., CHEN, Z. H., WANG, L., ZHANG, T., DUAN, L., BEHRENS, C., WISTUBA, II,

MINNA, J. D., GAO, B., LUO, J. H. & LIU, Z. P. 2016. LZTFL1 suppresses lung tumorigenesis

by maintaining differentiation of lung epithelial cells. *Oncogene,* 35**,** 2655-63.

WEI, Q., ZHOU, W., WANG, W., GAO, B., WANG, L., CAO, J. & LIU, Z. P. 2010. Tumor-

suppressive functions of leucine zipper transcription factor-like 1. *Cancer Res,* 70**,** 2942-50.

Figures and Legends

Figure 1. Mutation feature of LZTFL1 in different tumors. (A) The mutation site of LZTFL1 in tumors. The prediction of Leucine Zipper (Leu_zip) domain (B) and the binding region (C) structure of LZTFL1. The mutation type of LZTFL1 in different tumors.

Figure 2. The mRNA and protein expression of LZTFL1 in different tumors and pathological stages. (A) The mRNA expression of LZTFL1 in different cancers or specific cancer subtypes was analyzed through TIMER2. (B&C) The protein expression of LZTFL1 in between normal tissue and primary tissue of breast cancer, lung adenocarcinoma, clear cell RCC, UCEC, and ovarian cancer by the main pathological stages.

Figure 3. The correlation between LZTFL1 gene expression (A) and survival prognosis (B) of cancers in TCGA.

Figure 4. The correlation analysis between LZTFL1 expression and immune infiltration of cancer-associated fibroblasts. Different algorithms were used to explore the potential correlation between the expression level of the LZTFL1 gene and the infiltration level of cancer-associated fibroblasts across all types of cancer.

Figure 5. The LZTFL1 related enrichment and downstream gene function analysis. (A) The LZTFL1 related enrichment analysis. The downstream of LZTFL1 genes enriched in SARS-CoV-2 related GEO dataset (B) and human metabolism pathway (C). (D) The subcellular localization of LZTFL1 downstream gene. (E) The top 10 downstream miRNA regulated by LZTFL1.

Figure 6. The schematic diagram shows the role of LZTFL1 in COVID-19 and cancer.

432   Figure 2

433
434
435

436   Figure 3

437
438

439   Figure 4

442   Figure 5

445   Figure 6

446