Reliable energy-based antibody humanization and stabilization

Ariel Tennenhouse[1], Lev Khmelnitsky[2], Noa Yeshaya[2], Ashish Noronha[3], Moshit Lindzen[3], Razi Khalaila[3], Ira Zaretsky[4], Yael Fridmann Sirkis[5], Yael Galon-Wolfenson[5], Jakub Abramson[3], Yosef Yarden[3], Deborah Fass[2], Sarel J. Fleishman[1*]

1 Weizmann Institute of Science, Department of Biomolecular Sciences
2 Weizmann Institute of Science, Department of Chemical and Structural Biology
3 Weizmann Institute of Science, Department of Immunology and Regenerative Biology
4 Weizmann Institute of Science, Antibody Engineering Unit
5 Weizmann Institute of Science, Protein Analysis Unit
*correspondence: sarel@weizmann.ac.il

Abstract

Humanization is an essential step in developing animal-derived antibodies into therapeutics, and approximately 40% of FDA-approved antibodies have been humanized. Conventional humanization approaches graft the complementarity-determining regions (CDRs) of the animal antibody onto several homologous human frameworks. This process, however, often drastically lowers stability and antigen binding, demanding iterative mutational fine-tuning to recover the original antibody properties. Here, we present Computational hUMan AntiBody design (CUMAb), a web-accessible method that starts from an experimental or model antibody structure, systematically grafts the animal CDRs on thousands of human frameworks, and uses Rosetta atomistic simulations to rank the designs by energy and structural integrity (http://CUMAb.weizmann.ac.il). CUMAb designs of three independent antibodies exhibit similar affinities to the parental animal antibody, and some designs show marked improvement in stability. Surprisingly, nonhomologous frameworks are often preferred to the highest-homology ones, and several CUMAb designs relying on different human frameworks and encoding dozens of mutations from one another are functionally equivalent. Thus, CUMAb presents a general and streamlined approach to optimizing antibody stability and expressibility while increasing humanness.

Introduction

Antibodies are the largest segment of protein-based therapeutics, with over 100 antibodies in clinical use or under regulatory review[1]. As many as 40% of these antibodies were isolated from an animal source (often a mouse) and were humanized prior to clinical application. Antibody humanization is essential to achieve three important therapeutic goals: recruiting the immune system through Fc effector functionality; increasing blood circulation half-life; and reducing or eliminating immunogenicity in case the antibody is destined for long-term treatments[2-4].

Despite its critical role, antibody humanization has typically relied on an iterative trial-and-error process[5-7]. The first step chimerizes the animal variable fragment (Fv) with human constant domains (Fig. 1A). Next, the Fv, which comprises more than 200 amino acids and is divided into light and heavy chains, is humanized. In this step, the Fv is mutated to resemble natural human antibodies, ideally without compromising antibody binding function or stability. This step is

complicated by the fact that the Fv contains the so-called complementarity-determining regions (CDRs), hypervariable segments which are responsible for antigen recognition and are therefore very sensitive to mutation (Fig. 1A). Thus, the mainstream humanization strategy grafts the parental CDRs from the animal antibody on a human framework, typically leading to an Fv with >80% sequence identity to the human germline (compared to 50-70% identity for a mouse Fv, Supplementary Table 1). The frameworks are conventionally selected according to sequence[5] or structural homology to the animal antibody[8]. Some other humanization methods humanize predicted immunogenic segments in the parental framework[9] instead of the entire framework, graft residues from human frameworks into the animal antibody in solvent-exposed positions[10], or use structure-based calculations to remove humanizing mutations that may destabilize the Fv[11]. Despite these innovations, Fv humanization typically leads to substantial, sometimes orders-of-magnitude, decreases in expression levels, stability, and affinity[5,12–14]. These undesirable outcomes reduce efficacy, increase production costs, and complicate drug delivery and formulation[15]. Therefore, as a rule, a third step of "back mutation", in which positions in the humanized antibody are mutated to their animal identities, is often necessary, requiring lengthy design/experiment iterations[6,13].

The loss in affinity and stability often caused by humanization stems from structure and energy incompatibilities between the animal CDRs and the human framework[16,17]. Structural analyses singled out the importance of a region of the framework called the vernier zone, which underlies the CDRs. This zone comprises approximately 30 sequence determinants that vary even among homologous frameworks, impacting CDR structural integrity and stability[16,17]. Thus, most back mutation attempts select mutations that reconstitute some of the vernier-zone positions seen in the animal antibody[5]. This process can regain the parental antibody's affinity and stability, though at the cost of lower humanness and lengthy iterations.

Here, we test whether energy-based ranking is more effective than homology in designing humanized antibodies. Starting from experimentally determined or computed model structures, we use automated Rosetta all-atom calculations to design humanized variants and rank them solely by their energy. Crucially, automation allows us to systematically expand the options of unique humanized designs from a few dozen per antibody in conventional humanization to more than 20,000. We validate the method by demonstrating that some of the top-ranked designs for three unrelated antibodies exhibit stability and binding properties on par and sometimes greater than the parental animal antibodies without requiring back mutation or iterative design/experiment cycles. Remarkably, in some cases, the experimentally best-performing humanized design derives from human framework genes that do not exhibit high homology to the animal antibody, suggesting that energy-based humanization may provide a different and more effective solution than conventional homology-based humanization. The algorithm, which we call CUMAb for Computational hUMan AntiBody design, is general, computationally efficient, and enabled for academic use through http://CUMAb.weizmann.ac.il.

Results

Systematic energy-based ranking of humanized antibodies

We describe a general computational workflow for modeling and energy ranking humanized Fv designs. Starting from an experimental or computational structure of the animal Fv, we replace all the framework regions with compatible human ones. The framework is encoded in two gene segments, V and J, on both the light and heavy chains (Fig. 1A)[18]. Recombining all human V and J segments on both light and heavy chains yields tens of thousands of unique frameworks (63,180 that comprise kappa light chains and 48,600 comprising lambda light chains)[19]. Our working hypothesis is that the energy-based ranking of a diverse set of human frameworks may yield humanized designs that are more stable and functional than those selected from conventional, homology-based humanization.

We define the CDRs based on visual inspection of representative Fv structures and guided by previous definitions[20,21] (Fig. 1A, Supplementary Table 2). The first step combines all human V and J gene sequences obtained from the ImMunoGeneTics (IMGT) database[19] (Fig. 1B) to generate acceptor framework sequences. D genes are not considered as they are fully encompassed by CDR H3, which is held fixed in all calculations. Depending on the light-chain class of the animal antibody, either kappa or lambda light chain genes are used. We exclude human genes that contain Asn-Gly or Asn-X-Ser/Thr (where X is not Pro) sequence motifs as these may lead to undesirable post-translational modifications, including N-linked glycosylation[5]. Additionally, we exclude any gene that exhibits more than two cysteines outside the CDRs to reduce the chances of aggregation or disulfide shuffling. These restrictions retain a large fraction of possible combinations of human genes resulting in >20,000 unique human acceptor frameworks per antibody. We next concatenate the mouse CDR sequences with the human framework regions to generate humanized sequences (Fig. 1C).

Using the parental antibody Fv structure as a template, each humanized sequence is modeled using Rosetta all-atom calculations[22]. The resulting model structure is relaxed through cycles of Rosetta combinatorial sidechain packing and constrained whole-protein minimization in the entire Fv. If an experimentally determined structure of the antigen-antibody complex is available, the antigen-binding residues on the antibody are held fixed in all design calculations. Each model is ranked using the Rosetta all-atom energy function 2015 (ref2015), which is dominated by van der Waals interactions, hydrogen bonding, electrostatics, and implicit solvation[23]. Designs in which any of the CDR backbone conformations deviate by more than 0.5 Å from the parental conformation are eliminated to ensure the structural integrity of the CDRs in their humanized context. To select a diverse set of sequences for experimental testing, we cluster the top-ranked designs according to V-gene subgroups[19], which are defined according to sequence homology (7, 6, and 10 subgroups for heavy V, kappa, and lambda V genes, respectively). Clustering yields a shortlist of diverse, low-energy models for experimental testing. Of note, unlike conventional CDR-grafting methods, our approach is agnostic to homology between the animal and human framework and is scalable, designing and ranking 20,000 different humanized constructs on our computational cluster within a few hours.

Figure 1: Key steps in energy-based antibody humanization using CUMAb. (**A**) Schematic of antibody domains. (**B**) CUMAb begins by combining human germline heavy and light V and J genes to generate more than 20,000 human frameworks per antibody. (**C**) CDRs from the parental antibody are grafted onto each human acceptor framework. The humanized designs are then modeled in Rosetta and clustered, yielding low-energy cluster representatives for experimental screening.

One-shot humanization matches parental antibody affinity on multiple frameworks

As the first target for humanization, we chose a murine monoclonal antibody MAb492.1 (mαQSOX1) that inhibits the enzymatic activity of human Quiescin Sulfhydryl Oxidase 1 (QSOX1)[24]. This antibody is under development as a potential cancer therapeutic due to its ability to block the contribution of QSOX1 to extracellular matrix support of tumor growth and metastasis[25]. This antibody presents a stringent test for humanization because chimerizing its mouse Fv with a human IgG1 constant region led to a complete loss of expressibility in HEK293 cells[14]. We previously addressed this challenge with the AbLIFT method, which uses atomistic design calculations to improve the molecular interactions between the Fv light and heavy domains[14]. One of the designs, AbLIFT18, rescued expression in HEK293 cells and QSOX1 inhibition of the chimeric antibody composed of human IgG1 constant regions and murine Fv. However, this design was an incompletely humanized antibody (66.3 % and 57.3 % V gene sequence identity to the nearest human germline in the light and heavy chains, respectively) and was not suitable for therapeutic applications in humans.

To simulate a realistic humanization scenario, CUMAb was performed directly on the parental mouse antibody mαQSOX1. We produced expression vectors encoding the five top-ranked CUMAb designs formatted as separate light and heavy chains and experimentally tested all 15 unique pairs of these light and heavy chains. Remarkably, 12 pairs showed comparable expression levels to AbLIFT18 by dot-blot analysis (Supplementary Fig. 1A), while no expression was detected for the chimeric construct comprising the mouse Fv and human constant domains[14]. Furthermore, electrophoretic-mobility analysis after purification indicated that seven designs showed comparable expression levels to AbLIFT18 without obvious misfolding or aggregation (Fig. 2A). The expressible designs were purified and screened for QSOX1 inhibition, and many of the designs were functional (Supplementary Fig. 1B). Two highly expressing, functional designs (hαQSOX1.1, hαQSOX1.2) exhibited similar inhibition constants to that of the parental antibody (Fig. 2B, Supplementary Fig. 1C). We noted that CUMAb ranked highly multiple genes from the VH3 subgroup from which hαQSOX1.1 derives, each pairing with a different kappa light chain gene. We therefore tested two additional constructs (hαQSOX1.3, hαQSOX1.4) finding that they exhibited similar inhibition constants to that of the parental antibody as well (Fig. 2B). All four designs have similar sequence identity to the human V gene germline in both the light chain (78-81%) and the heavy chain (80-86%). These V gene sequence identities are significantly higher than those for the mouse antibody and AbLIFT18, which both have 66% sequence identity (light chain) and 57% (heavy chain, Fig. 2C). Additionally, these V gene sequence identities are in the range of those among FDA-approved humanized antibodies, which have a mean of 84% sequence identity in the light chain and 81% in the heavy chain (Supplementary Table 1). Thus, the four designs recapitulated the parental mouse expression levels and activity using different human Fv frameworks without requiring back mutations.

We measured the melting temperatures of the four designs as well as the parental antibody with nano-differential scanning fluorimetry (nano-DSF) and found all five to be above 70℃ (Supplementary Fig. 1D). To probe the stability of the strongest-inhibiting antibody in a therapeutic setting, we incubated hαQSOX1.1 and mαQSOX1 with plasma from two human donors at 37 degrees and measured the inhibition of native QSOX1 as a function of time. We found that both antibodies nearly completely inhibited QSOX1 for over 60 hours (Fig. 2D), indicating that the humanized designs retained the function and stability properties of the parental mouse antibody. hαQSOX1.1, hαQSOX1.2, and hαQSOX1.3 differ from one another in either the light or heavy chain subgroup and have 33-56 mutations from one another (Fig. 2E). Due to these mutations, they have strikingly different patterns of surface charge. Thus, CUMAb produces antibodies that are functionally nearly identical but have distinct surface properties. As surface properties have been associated with changes in the propensity of antibodies to self-associate or form non-specific interactions[26,27], CUMAb may provide a path to improving properties that are critical for therapeutics development while improving humanness.

Finally, we determined a co-crystal structure of hαQSOX1.4 bound to the oxidoreductase fragment of human QSOX1 and found that the design and parental antibody are strikingly similar, with only a 0.7 Å Cα RMSD despite 51 mutations between the parental and humanized

antibodies (Fig. 2F, Supplementary Fig. 1E, Supplementary Table 3). These results verify CUMAb's atomic accuracy and its ability to rapidly produce functionally similar and stable humanized designs even in a case that defied previous humanization efforts.

Figure 2: Humanization of an anti-QSOX1 antibody. (**A**) Electrophoretic analysis of 15 CUMAb designs (all combinations of the five light chains and three heavy chains from the top five clustered designs) as well as the most successful AbLIFT design (AbLift18) as a positive control[14]. CUMAb designs are labeled according to heavy chain and then by light chain, with the number referring to the IMGT subgroup from which the designed chain derives. If there are two chains from the same subgroup, they are arbitrarily labeled "a" or "b". Amino acid sequences of each antibody are found in Supplementary Table 4. (**B**) Ki values obtained from measurements of relative QSOX1 activity at different antibody concentrations (Fits shown in Supplementary Fig. 1C). Error bars represent standard deviations of three independent repeats. (**C**) Sequence identity to the nearest human germline as calculated by IgBLAST for four CUMAb designs as well as the parental mouse antibody. (**D**) Top: Schematic of the experimental workflow. Blood plasma from human donors is incubated with an antibody at 37°C. At various time points, samples are incubated with dithiothreitol

(DTT). After an incubation period to enable any uninhibited QSOX1 to oxidize the DTT,5,5′-dithiobis-(2-nitrobenzoic acid) (DTNB) was added to quantify remaining thiol groups. The colorimetric output of this assay is detected at 412 nm. Bottom: Absorbance at 412 nm at different timepoints of human blood plasma incubation with hαQSOX1.1 (pink), mαQSOX1 (gray), and PBS (blue) for two human blood donors, as well as PBS incubated with mαQSOX1 ("no plasma", black). Both hαQSOX1.1 and mαQSOX1 completely inhibit human QSOX1 in the blood plasma at all time points. (**E**) Surface electrostatic charge of four CUMAb designs as well as the parental mouse antibody, showing that each design has a very different electrostatic surface and that all four differ from the mouse antibody. Surface electrostatics were generated from CUMAb models using the PyMOL APBS Electrostatics plugin. (**F**) Alignment of crystal structure of the Fv of hαQSOX1.4 (magenta; PDB entry 8AON) to the crystal structure of the parental mouse antibody (gray; PDB entry 4IJ3), showing that the structures are nearly identical despite 51 mutations between them. Mutations are shown in magenta spheres. Fv structures were extracted from co-crystal structures of the antibodies with human QSOX1 oxidoreductase fragment (QSOX1trx). The complex structure is shown in Supplementary Fig. 1E.

Successful humanization based on an Fv model structure

Experimental structure determination is a lengthy and uncertain process, and obtaining an experimentally determined structure for each isolated antibody in an antibody development campaign is unrealistic. Recently, structure-prediction methods have reached the point at which the antibody Fv structure can be predicted to nearly atomic accuracy except in the solvent-exposed parts of CDR H3, where models are still insufficiently reliable[28]. While H3 is a critical determinant of antigen recognition, most of its solvent-exposed region does not form direct interactions with the framework region, which is the target of humanization. Therefore, we hypothesized that despite the demand for stringent accuracy by atomistic design calculations, Fv model structures might be sufficiently accurate for the purposes of humanization design.

Dedicated Fv structure-prediction methods yield accurate structure models[28–33]. Given the recent revolution in deep-learning-based protein structure prediction methods, however, we chose to use AlphaFold-Multimer to predict the Fv structure of our target antibody[34,35]. Starting from the top-ranked AlphaFold model, all design calculations are identical to the CUMAb workflow above, except that we do not eliminate humanized models in which the backbone deviates from the starting model structure, thus allowing for modeling inaccuracies.

To test the ability of CUMAb ability to humanize antibodies without recourse to experimentally determined structures, we targeted an antibody that was raised in mice immunized against the human AXL receptor tyrosine kinase (mαAXL). This antibody has a unique framework mutation from the mouse germline that interacts with CDR H3 (Fig. 3A). To account for the uncertainty regarding this mutation, we included five designs with this mutation in addition to our five CUMAb designs. The ten designs were screened for expression, and four designs exhibited high expression levels (three designs with the additional mutation expressed but at lower levels, Fig. 3B). All ten designs were assayed for binding to AXL by immunoprecipitation, and one design showed much stronger binding than the others. This design (hαAXL) had 40 mutations from the parental antibody and 84% (light chain) and 89% (heavy chain) V gene sequence identity to the closest human germline, compared to 62% (light chain) and 79% (heavy chain) in the mouse antibody (Fig. 3C, D).

We also measured the melting temperature of the best design and the parental antibody, revealing that the design was 3°C more stable (Fig. 3E). Finally, we measured the affinity of both the design and the parental antibody to AXL using surface plasmon resonance and found

that the affinities were comparable (3.2 nM for the design vs. 0.95 nM for the parental, Fig. 3F). We conclude that despite the uncertainty when using Fv model structures rather than experimentally determined ones, CUMAb can rapidly generate stable, well-expressed high affinity humanized antibodies, even in a case where the parental mouse antibody had an unusual framework mutation.

Figure 3: Humanization of an anti-human AXL antibody based on an AlphaFold2 model structure. (**A**) Unusual mutation from germline in the framework of parental mouse antibody (pink), which is predicted to interact with CDR H3 (blue). (**B**) Clear cell extracts were subjected to immunoprecipitation with media containing humanized AXL antibody (or control, only media) and immunoprecipitates were immunoblotted (IB) as indicated. Marked are the 10 CUMAb designs (6-10 include unusual framework mutation from A, the protein ladder (L), and the AXL protein used as input (I). (**C**) Alignment of AlphaFold2 model of mouse anti-AXL antibody (gray) and CUMAb design 1 (green) with mutations indicated by green spheres. (**D**) Sequence identity to nearest human germline as calculated by IgBLAST for the parental mouse antibody (mαAXL, gray) as well as the best-performing CUMAb design (hαAXL, green). (**E**) SPR kinetic analysis of mαAXL (gray) and hαAXL (green) to AXL (kinetic fits shown in black). Antibodies were covalently attached to a CM5 chip, and AXL was injected at 7, 5, 2, 1, and 0.8 nM. mαAXL exhibited k_a = 5.5 * 10^5 $M^{-1}s^{-1}$, k_d = 5.24 * 10^{-4} s^{-1}, and K_D = 0.95 nM. hαAXL exhibited k_a = 1.2 * 10^6 $M^{-1}s^{-1}$, k_d = 3.9 * 10^{-3} s^{-1}, and K_D = 3.2 nM. (**F**) Thermal denaturation of mαAXL (gray) and hαAXL (green) using nano-DSF. Shown is average of two repeats.

Increased humanness through specificity-determining residue grafting

The CUMAb CDR grafting procedure typically raises the sequence identity to the human germline to 80-88% (Supplementary Table 3), in line with the levels observed for antibodies in clinical use (Supplementary Table 1). To increase humanness even further, an alternative strategy, called specificity-determining residue (SDR) grafting, was proposed[36,37]. SDR grafting maintains only the amino acid positions that directly contact the antigen, whereas the remainder of the antibody, including CDR regions that do not interact with the antigen, is humanized. We next extend the CUMAb energy-based strategy to SDR grafting. To prevent undesirable changes to the CDR backbone conformations in SDR grafting, we use only human germline genes that exhibit CDRs that match the lengths of the CDRs in the parental antibody, with the exception of H3 (see Methods). Furthermore, because the heavy-chain J gene segment encodes a part of H3, we cluster the resulting humanized designs according to their V gene subgroups as well as their heavy-chain J segment. Thus, unlike CDR grafting above, CUMAb SDR grafting may select designs that differ from one another only in their heavy-chain J segments. We limit the application of SDR grafting to experimentally determined antigen-bound

structures because it demands accurate determination of antigen-binding amino acid positions, including the solvent-accessible region in H3.

We applied SDR grafting to the murine antibody D44.1 (mαHEWL), which targets hen-egg white lysozyme. The co-crystal structure (PDB entry: 1MLC)[38] reveals that 28 of 63 CDR positions in D44.1 interact with the antigen. Thus, in this case, SDR grafting would lead to roughly 90% (heavy chain) and 94% (light chain) V gene sequence identity to the human germline, compared to 86% (heavy chain) and 79-88% (light chain) using the CDR grafting procedure. In an initial experimental screen, we formatted six designs as single-chain variable fragments (scFvs) and tested lysozyme binding using yeast-surface display (Supplementary Fig. 2A)[39]. This preliminary screen indicated that five designs exhibited high expression levels and that designs one and six bound lysozyme, with design one markedly superior. This qualitative single-concentration experiment indicated that design one may bind lysozyme as strongly as the parental D44.1.

To obtain quantitative binding data, we formatted design 1 (hαHEWL) as a human IgG1 Fab and expressed it and D44.1 (as mouse IgG1 Fab) in HEK293 cells. Design 1 has 64 mutations relative to the parental mouse antibody and substantially increases the antibody's humanness in both the light and heavy chains (Fig. 4A, B). Both antibodies expressed well, with the design showing noticeably higher purification yield when the Fabs were purified using an anti-His tag column (Fig. 4C). Furthermore, the melting temperature of the designed variant was 5°C higher than that of the mouse (Fig. 4D). We measured the affinity of the mouse and humanized Fabs, and found that the humanized design has an affinity of 44 nM as compared to 6.5 nM for the parental mouse antibody (Fig. 4E). Thus, in this case, the greater level of humanization enabled by SDR grafting may have come at the cost of some loss in affinity, though as with CDR grafting, expressibility and stability have increased.

Figure 4: SDR-grafting of anti-hen egg-white lysozyme (HEWL) antibody. (A) Alignment of crystal structure of mouse anti-HEWL antibody (gray, PDB entry: 1MLC) and CUMAb design 1 (blue) with mutations indicated by blue spheres. (B) Sequence identity to nearest human germline as calculated by IgBLAST for the parental mouse antibody (gray, mαHEWL) as well as the best-performing CUMAb design (blue, hαHEWL). (C) Electrophoretic mobility analysis of mαHEWL and hαHEWL. Equal volumes of each protein were analyzed after purification with anti-His tag column. (D) Thermal denaturation of mαHEWL (gray) and hαHEWL (blue) using nano-DSF. Shown is average of two repeats. (E) SPR kinetic analysis of mαHEWL (gray) and hαHEWL (blue) to AXL (kinetic fits shown in black). Antibody Fabs were covalently attached to a CM5 chip, and HEWL was injected at 20, 15, 10, 5, and 3.75 nM. mαHEWL exhibited k_a = 2.0 * 10^6 M^{-1}s^{-1}, k_d = 1.4 * 10^{-2} s^{-1}, and K_D = 7.4 nM. hαAXL exhibited k_a = 3.9 * 10^6 M^{-1}s^{-1}, k_d = 1.6 * 10^{-1} s^{-1}, and K_D = 42 nM.

Successful energy-based humanization exploits nonhomologous frameworks

Conventional antibody humanization strategies use the human germline genes that are closest to those of the sequence or structure of the animal antibody. CUMAb's energy-based strategy ignores sequence or structure homology and focuses instead on the structural integrity and energy of the design models. Energy-based humanization vastly expands the scope of potential frameworks from a few dozen high-homology ones to tens of thousands. We analyzed the three test cases presented here to understand the role of energy versus sequence homology in successful antibody humanization.

Strikingly, in two of the three cases, the best-performing humanized designs do not derive from the best-matched human ones according to sequence identity (Table 1). In the anti-QSOX1 antibody, none of the best designs derive from the closest human V gene subgroup for either the light or heavy chain, and in the anti-lysozyme antibody, only the heavy chain matches. Furthermore, the anti-lysozyme designs based on the same subgroup of human V genes were significantly inferior to the best-performing humanized design. As an additional comparison with conventional humanization strategies, we attempted to humanize the anti-QSOX1 antibody using the "consensus" approach. In this humanization approach, the framework is taken from a sequence-based consensus of V gene subgroups[40] (in this case, IGKV1 and IGHV4). The designed antibody, however, failed to express, and we could not test its binding to QSOX1 (Fig. 5A), providing another demonstration that energy-based humanization may be superior to conventional humanization strategies.

Thus, grafting CDRs from an animal antibody onto the closest (or consensus) human germline often fails to recapitulate their stability and functional properties, as has been observed in decades of antibody engineering[12,13]. By contrast, some of the top-ranked designs from energy-based humanization exhibit stability and binding properties indistinguishable from those of the animal antibody, and these designs typically do not derive from germline gene subgroups that are homologous to those of the animal antibody.

We next ask what are the structural underpinnings of successful design using nonhomologous frameworks. Our analysis carries significant uncertainty because the designs diverge from one another by dozens of mutations across the framework, and thus, claims on which mutations determine success or failure must be tentative. Yet, comparing the designs that use homologous frameworks with the best-performing designs shows striking structural differences that may account for the higher stability and activity of the latter (Supplementary Figure 3). For instance, humanization of the anti-QSOX1 antibody based on the consensus framework of the VH4 subgroup, the highest homology subgroup according to IgBLAST, results in two mutations from the parental antibody in the framework regions. One of these mutations eliminates a buried hydrogen bond network that connects the framework and CDRs 1 and 2 (Fig. 5A), and another leads to steric overlaps with a position on CDR H1 (Fig. 5B). Additionally, all successful CUMAb designs introduced light-chain mutation Ser43Pro, which is also seen in the AbLIFT18 design and is likely to stabilize the antibody (Fig. 5C)[14]. Humanizing the anti-lysozyme antibody with the highest-homology VK3 family would result in a His34Ala mutation in the light chain, disrupting a

hydrogen-bond network among the light chain CDRs (Fig. 5D). Strikingly, although (mαHEWL) introduces a mutation in this same critical position (His34Asn), this mutation is predicted to retain the hydrogen-bond network. We conclude that despite the high homology within V-gene subgroups, critical long-range interactions between the framework and the CDRs are often abrogated by conventional homology-based grafting. By contrast, low-energy ranked designs often retain the very same identities seen in the parental antibody and sometimes mutate them while nevertheless conserving critical stabilizing interactions.

Figure 5: Structural determinants of successful humanization by nonhomologous frameworks. (A-C). Crystal structure of mouse anti-QSOX1 (gray, PDB entry: 4IJ3) aligned with CUMAb model of successful design hαQSOX1.2 (red) and CUMAb model of unsuccessful humanization onto consensus framework IGKV1-IGHV4 (yellow). Amino acid numbering according to Kabat[20]. **(A)** Lys71 in the heavy chain framework of the mouse anti-QSOX1 antibody makes three stabilizing hydrogen bonds with backbone carbonyls of CDRs H1 and H2. This Lys is conserved in the successful design but mutated to Val in the consensus framework IGHV4, likely destabilizing the H1 and H2 loops. **(B)** Val78 in the heavy chain framework of mouse anti-QSOX1 is mutated to Phe in the consensus framework IGHV4. This likely strains the molecule due to a steric overlap with Leu29 in CDR H1. **(C)** Ser43 in the light chain of mouse anti-QSOX1 is mutated to Pro in the successful design, likely stabilizing the molecule by rigidifying the loop. Mutation to Ala in the consensus IGKV1 framework likely has little effect. **(D)** Crystal structure of mαHEWL (gray, PDB entry 1MLC) aligned to CUMAb model of successful design hαHEWL (blue) and CUMAb model of unsuccessful design hαHEWL.5 (yellow). Position 34 in the light chain CDR L1 makes two key hydrogen bonds to the side chain of Ser91 in CDR L3 and to the backbone amide of Tyr50 in L2. Because this residue is not in the interface with the antigen, it is mutated in SDR grafting even though it is in a CDR. In the consensus framework IGKV1, this position is mutated to Ala, likely destabilizing CDRs L2 and L3 due to the loss of the hydrogen bonds. Strikingly, in successful CUMAb design 1, this position is mutated to Asn, which is likely to retain the hydrogen bonds.

Discussion

Instead of using only a few dozen homologous frameworks as in the majority of conventional humanization strategies, CUMAb uses all combinations of possible human gene segments (>20,000 for each antibody) and ranks them by energy. Although genes belonging to a single subgroup are homologous, they contain mutations, including in vernier positions that may stabilize the specific CDRs of the parental antibody. Structural analysis shows that low-energy designs retain critical framework-CDR interactions that may be eliminated in homology-based humanization. Consequently, the lowest-energy designs from this large space of possible frameworks are more likely than the highest homology ones to retain stability, expression yields, and binding affinity and even improve stability and expressibility relative to the parental antibody. CUMAb may thus offer a one-shot strategy to improve stability and expressibility, including of human-sourced antibodies, while maintaining or increasing humanness.

Crucially, CUMAb eliminates iterative back mutation even when starting from an Fv model structure. Thus, this approach increases the speed and scope of antibody humanization, in principle, to any animal antibody without the traditional bottlenecks of obtaining crystallographic

structures to enable accurate structure-based humanization[41]. Furthermore, CUMAb is automated and requires experimental screening of fewer than a dozen constructs, substantially reducing time and cost. It may therefore be applied at scale to dozens of antibodies in parallel. Unlike the results from conventional homology-based humanization methods, CUMAb generates several unique humanization alternatives that exhibit stability and binding properties that are on a par with those of the animal antibody. Since these alternative designs encode dozens of surface mutations from one another, they are likely to exhibit differences in critical developability properties, such as nonspecificity and solution properties at high antibody concentrations[26,27].

Finally, we note that CUMAb is based on the fundamental insight that antibody stability and activity are determined by the entire Fv, including the framework positions on which the CDRs rest[16,42] and the interfaces between the light and the heavy chains[14,43]. This insight may help address other important challenges in antibody engineering, leading to general, reliable, and rapid antibody design strategies.

Acknowledgments

We thank Adva Mechaly and Peter Tessier for critical reading and Ron Diskin and Olga Khersonsky for advice. Research in the Fleishman lab was supported by the European Research Council through a Consolidator Award (815379), the Dr. Barry Sherman Institute for Medicinal Chemistry, and a donation in memory of Sam Switzer. Research in the Fass lab was supported by the European Research Council through a Proof of Concept grant (825076). We acknowledge the European Synchrotron Radiation Facility for the provision of beam time on ID30B, and we thank Andrew McCarthy for his assistance.

Author contributions

AT and SJF wrote the manuscript with input from all authors. AT developed the humanization algorithm and designed all humanized sequences. LK performed experimental work related to anti-QSOX1 antibodies, LK, NY, and DF performed crystallography data collection, and LK and DF analyzed the data. AN performed IP assay for anti-AXL antibody. ML and IZ expressed and purified anti-AXL antibodies. AT and LK expressed and purified anti-HEWL antibodies. AT, YFS, and YGW performed nano-DSF and SPR characterization of anti-AXL antibodies and analyzed the data together with SJF, ML, and YY. AT, YSF, and YGW performed nano-DSF and SPR characterization of anti-HEWL antibodies and analyzed the data together with SJF. RK and JA collected and analyzed data.

Conflicts of interest

AT and SJF are named inventors in a patent filing on the CUMAb method. AT, SJF, LK, and DF are named inventors in a patent filing on humanized anti-QSOX1 designs.

Methods

Computational Methods

A database of antibody germline sequences

Antibody germline sequences were retrieved from the IMGT reference database[19] (downloaded July 29, 2020). For each gene, only the first allele that was annotated as functional was taken. Partial or reverse-complementary genes were pruned. If allele one contains more than two cysteines, a different allele was taken with two cysteines if available. This resulted in 54 heavy chain V gene sequences, six heavy chain J gene sequences, 39 light chain kappa V gene sequences, five light chain kappa J gene sequences, 30 light chain lambda V gene sequences, and five light chain kappa J gene sequences. We then combined these genes all-against-all (V and J for both light and heavy chains) for kappa and lambda light chains separately, resulting in 63,180 sequences for kappa light chains and 48,600 for lambda light chains.

Computational CDR grafting

We defined the CDRs based on previous definitions[20,21] and visual inspection of antibody crystal structures. Starting from a molecular structure of the parental antibody, we use HMMer to identify the sequence segments corresponding to the variable region and classify the light chain as kappa or lambda[44]. For each germline sequence corresponding to the light chain classification, the CDRs of the germline sequence are replaced with the CDRs of the parental (in the current study, mouse) antibody. Any sequence that contains an Asn-Gly or Asn-X-Ser/Thr (where X is not Pro) outside of the CDRs is removed. This typically results in >20,000 unique sequences per parental antibody.

Energy ranking of CDR-grafted sequences

If provided a crystal structure, as a first step, the structure of the parental antibody is relaxed through cycles of sidechain and harmonically restrained backbone minimization and combinatorial sidechain packing in the entire Fv (see Relax.xml RosettaScript[45]). If a bound structure is provided, residues in the interface between the Fv and the antigen are identified using Rosetta and held fixed during the relaxation (see Interface.xml). Additionally, if a bound structure is provided, the entire antibody Fv-antigen complex is relaxed.

Each CDR-grafted germline sequence is then threaded onto the relaxed Fv structure and relaxed in the same manner with fewer cycles (see CUMAb.xml). Sequences are ranked according to the all-atom energy ref2015[23]. Any model that exhibits a Cα-carbonyl O RMSD greater than or equal to 0.5 Å in any of the CDRs is excluded from further consideration. Sequences are clustered according to V gene subgroup as defined by IMGT, meaning that we take only one sequence from each V gene combination. Sequences were visually inspected, and, in some cases, the highest ranking representative for a cluster was replaced with a slightly lower ranking one in order to re-use sequences in different clusters and thus minimize cloning.

When starting from a model rather than an experimental structure, the pipeline is identical, except that sequences are not excluded for deviating from the model structure.

Generation of model structure of Fv for anti-AXL antibody

The model structure was predicted using ColabFold[34] with default settings. The antibody was formatted as a heterodimer, and thus AlphaFold multimer[35] was used to predict the structure of the antibody. ColabFold designs were ranked by pTM score (pLDDT score would have given an identical ranking in the cases examined here), and the highest ranked design was chosen.

Computational SDR grafting

The parental antibody is classified as having a kappa or lambda light chain as described above. We use Rosetta to identify residues in the interface between the antibody Fv and the antigen (see interface.xml). Antibody germline sequences are selected using the following criteria: The sequences must have the same CDR length in all CDRs, excluding CDR H3. Additionally, the sequences must have an H3 length that is equal to or shorter than the H3 length compared to the parental antibody. If the H3 length is shorter than the parental antibody, a number of residues equal to the difference in length of the two H3s from the parental antibody are inserted into the germline sequence. The germline sequences are then threaded and relaxed as described above. Sequences are clustered according to V gene and heavy J gene subgroup.

Evaluation of humanness of antibody sequences

Antibody heavy or light chain sequences are submitted to the IgBLAST[46] web server, and the sequence identity to the top hit is taken.

Generation of surface electrostatic charge maps for anti-QSOX1 antibodies

Charge maps were generated using the APBS Electrostatic plug-in[47] in PyMOL using default settings, with the exception of the Molecular Surface Visualization range, which was set to +/-3.50.

Experimental Methods

Expression and purification of anti-human QSOX1 antibodies

Expression plasmids for CUMAb light and heavy chains were constructed on the basis of the AbLIFT18 chimeric antibody[14] using restriction-free procedures to replace the variable regions. Plasmids were purified from the *E. coli* XL-1 strain. Proteins were produced by transient transfection of plasmids into HEK293F cells. Transfection was done using the PEI Max reagent (Polysciences Inc.) with a 1:3 ratio (w/w) of DNA to PEI at a concentration of 1 million cells per ml. Co-transfections were done with a 1:1 ratio of light- and heavy-chain DNA, while the total amount of DNA remained the same: 1 μg DNA per ml HEK culture. Six days after transfection,

the culture medium was collected and centrifuged for 10 min at 500 g to pellet cells. The supernatant was then centrifuged for 10 min at 3200 g to pellet any remaining particles. The supernatant was filtered through a 0.22 µm filter, and the antibodies were purified by protein G affinity chromatography (GE Healthcare). Antibodies were then dialyzed into PBS pH 8 overnight.

Expression and purification of human QSOX1 proteins

Construction of pET15b expression vectors for human QSOX1, used for activity assays (residues 34–546 of HsQSOX1), and of its oxidoreductase fragment (QSOX1trx, residues 36-275 of HsQSOX1), used for co-crystallization with a Fab fragment of a CUMAb design, were previously described[48]. Both include an amino-terminal His$_6$ tag. The His$_6$ tag of QSOX1trx is cleavable by a downstream thrombin cleavage site and was removed by thrombin cleavage prior to crystallization. QSOX1 proteins were purified by nickel-nitrilotriacetic acid (Ni-NTA) chromatography (GE Healthcare).

QSOX1 activity/inhibition assays

QSOX1 activity on Zymogen granule membrane protein 16 (ZG16) for initial antibody screening: ZG16 contains two cysteines in a flexible, accessible CXXC motif that can be oxidized in vitro by QSOX1[49]. ZG16 fused to maltose binding protein (MBP, ~45 kDa, no cysteines) was used to assay QSOX1 activity in the presence of supernatants from antibody-producing HEK293F cells. Prior to the assay, MBP-ZG16 was reduced by incubation with DTT, excess DTT was removed on a PD-10 column (GE Healthcare), and reactions of the reduced MBP-ZG16 and QSOX1 (50 nM) were initiated in the presence of the antibody variants, supplied by adding clarified HEK293F culture supernatants diluted into the reaction mixture. The mixture contained 8 microliters HEK293F supernatant, 2.8 microliters reduced MBP-ZG16 (2.6 micromolar final concentration), and 1.2 microliters QSOX1 (50 nM final concentration). The reactions were stopped by adding maleimide-functionalized polyethylene glycol of molecular weight 5000 Da (PEG-mal 5000, Iris biotech), which covalently modifies any residual free thiols, resulting in slower migration in SDS-PAGE. Loading buffer was added to the reaction mixture and the whole mixture was run on an SDS-PAGE gel. Increased amounts of modified MBP-ZG16 relative to the uninhibited reaction indicated inhibition in this screen.

Plasma-QSOX1 inhibition:
Human blood collected with heparin as an anticoagulant was centrifuged for 15 min at 300 g. The plasma was collected and used as a native QSOX1 source in QSOX1 inhibition reactions conducted in 100 µl reactions as follows: 10 µl plasma was incubated at 37° C with 25 nM of the CUMAb antibody hαQSOX1.1 in 50 mM potassium phosphate buffer, pH 7.5, 65 mM NaCl, and 1 mM EDTA. The reaction was initiated by the addition of dithiothreitol (DTT) to a final concentration of 300 µM. After 30 min at room temperature, 50 µl were removed and added to 725 µl of the above buffer and 25 µl of 2 mM 5,5′-dithiobis-(2-nitrobenzoic acid) (DTNB), which reacts with residual DTT and produces the chromogenic product 5-thio-2-nitrobenzoate (TNB) with absorption at 412 nm.

QSOX1 inhibition constant determination

A Clark-type oxygen electrode (Hansatech Instruments) was used to monitor changes in dissolved oxygen concentration as a measure of QSOX1 activity. QSOX1 (25 nM) and various concentrations (1-100 nM) of purified antibody were assayed in 50 mM potassium phosphate buffer, pH 7.5, 65 mM NaCl and 1 mM EDTA. Reactions were started by injection of DTT to a concentration of 200 µM in the electrode reaction chamber.

Two independent progress curves were collected for each antibody concentration, and initial slopes were calculated. The background decrease in oxygen concentration due to the presence of DTT and antibody was measured and subtracted from the initial slopes to obtain the rates of QSOX1 activity in the presence of various concentrations of each antibody. The ratios of the initial rates of QSOX1 in the presence and absence of inhibitor were plotted as a function of inhibitor concentration. The data were fit to the Morrison equation for a tight binding inhibitor (GraphPad Prism version 8.4.3 for Windows, GraphPad Software, San Diego, California USA, www.graphpad.com) to determine the inhibition constants of the antibodies.

Purification and crystallization of the hQSOX1trx/hαQSOX1.4 complex

Purified QSOX1trx was concentrated to 10 mg/ml . The purified CUMAb antibody hαQSOX1.4 was concentrated to 1.5 mg/mL in PBS and digested at 37 °C with papain (Sigma) at a 1:20 papain:hαQSOX1.4 molar ratio. Prior to use, papain was activated by dissolving in PBS supplemented with 20 mM EDTA and 20 mM cysteine. Digestion was stopped after 4 h using leupeptin, and the digested antibody was dialyzed against PBS, pH 8. The Fab fragment of hαQSOX1.4 was purified by protein A affinity chromatography (Cytiva). Purified hαQSOX1.4 Fab was incubated for 1 h at 4 °C with a 2- to 3-fold excess of QSOX1trx, and the complex was isolated by size-exclusion chromatography in 10 mM Tris, pH 7.5, and 100 mM NaCl. The complex was concentrated using a centrifugal concentrator to 10 mg/mL. Crystals were grown by hanging-drop vapor diffusion at 293 K by mixing 1 µl of protein complex solution with 1 µl of well solution and suspending over well solution. Initial branched, feather-like crystals were observed when well conditions were 14% w/v polyethylene glycol 4 kDa and 0.1 M ammonium phosphate dibasic. These crystals were seeded into drops in similar conditions (11-14% w/v polyethylene glycol 4 kDa and 0.05-0.15 M ammonium phosphate dibasic) to produce individual crystal rods. These crystals were soaked in a similar solution with an addition of 20% glycerol and flash frozen in liquid nitrogen.

X-ray data collection and model building

X-ray data were collected at a wavelength of 0.9763 Å on the ID30B beamline at the European Synchrotron Radiation Facility, using a helical collection mode. Molecular replacement was performed using Phenix[50] with the hQSOX1trx structure from PDB entry 4IJ3 and an Alphafold prediction model for the hαQSOX1.4 Fab structure as input models. The initial complex model was then iteratively rebuilt and refined using Coot[51] and Phenix[50], respectively. The refined structure was deposited to the PDB with entry code 8AON.

Secreted Fab (anti-HEWL) expression and purification

The variable regions of the different heavy and light chains were cloned separately, upstream of IgG1 human Ab Fab scaffolds, into p3BNC plasmids. Antibody Fabs were produced by transient transfection of plasmids into HEK293F cells. Transfection was done using the PEI Max reagent (Polysciences Inc.) with a 1:3 ratio (w/w) of DNA to PEI at a concentration of 1 million cells per ml. Co-transfections were done with a 1:1 ratio of light- and heavy-chain DNA, while the total amount of DNA remained the same: 1 μg DNA per ml HEK culture. Six days after transfection, the culture medium was collected and centrifuged for 10 min at 500 g to pellet cells. The supernatant was then centrifuged for 10 min at 3200 g to pellet any remaining particles. The supernatant was filtered through a 0.22 μm filter, and the antibodies were purified by nickel-nitrilotriacetic acid (Ni-NTA) chromatography (GE Healthcare). Antibodies were then dialyzed into PBS pH8 overnight.

Expression and purification of humanized anti-human AXL

Anti-hAXL was generated as a human IgG1/IgK subclass. The sequences of the variable regions of the heavy and light chains were synthesized by Twist Bioscience, polymerase chain reaction (PCR)–amplified, and cloned into mammalian expression vectors (purchased from Oxford genetics) in frame with the constant regions of the human IgG1 heavy chain or human kappa light chain, respectively. Plasmid sequences were validated by direct sequencing (Life Sciences Core Facility, Weizmann Institute of Science). To produce antibodies, antibody heavy and light chain expression vectors were transiently transfected into Expi293 cells (Thermo Fisher Scientific). The secreted antibodies in the cell supernatant were purified using protein G Sepharose 4 Fast Flow Resin (GE Healthcare). Purified antibodies were dialyzed in phosphate-buffered saline and sterile-filtered (0.22 μm). Purity was assessed by SDS-PAGE and Coomassie staining and was estimated to be >90% (data not shown).

Immunoprecipitation of anti-AXL antibodies

Immunoprecipitation assays used protein G beads. Following gel electrophoresis, proteins were transferred onto a nitrocellulose membrane. The membrane was blocked in TBST buffer (0.02 M Tris-HCl (pH 7.5), 0.15 M NaCl, and 0.05% Tween 20) containing albumin (3%), blotted with a primary antibody overnight at 4° C, washed in TBST and incubated for 30 min with a secondary antibody conjugated to horseradish peroxidase.

Apparent Tm measurements of anti-HEWL and anti-AXL antibodies

The apparent melting temperature of the antibodies was determined by the Prometheus NT. 48 instrument (NanoTemper Technologies). Full length IgGs (anti-AXL) or Fabs (anti-HEWL) were diluted to 0.5 mg/ml (anti-AXL) or 0.1 mg/mL (anti-HEWL) in 1X PBS. The temperature was ramped from 20°C to 95°C at 1.0°C/min, and Tm was measured. Calculations were done using Prometheus NT. 48 default software. All antibodies were repeated.

Surface-plasmon resonance of anti-AXL antibodies

Surface plasmon resonance (SPR) experiments on the anti-AXL antibodies were carried out on a Biacore S200 instrument (GE Healthcare, Sweden) at 25°C with PBST (1X PBS + 0.05% Tween-20). For binding analysis, approximately 1,000 response units (RU) of AXL (R&D systems) were conjugated to a CM5 sensor chip using NHS/EDC chemistry. Samples of different antibody concentrations were injected over the surface at a flow rate of 30 µL/min for 240 s, and the chip was washed with buffer for 180 s. After each injection, surface regeneration was performed with a 30 s injection of 1 mM NaOH at a flow rate of 30 uL/min. The first channel contained no immobilized ligand, and its sensogram was subtracted from the channel used to measure binding. The acquired sensograms were analyzed using the device's software version 1.1, and kinetic fits to 1:1 binding model were performed.

Yeast surface display of anti-HEWL antibodies

Yeast-display experiments were conducted essentially as described[39]. Briefly, yeast cells were grown in selective medium SDCAA overnight at 30°C. The cells were then resuspended in 1 ml of induction medium and incubated at 20°C for 20 h. Roughly 10^7 cells were then used for yeast-cell surface display experiments: cells were subjected to primary antibody (mouse monoclonal IgG1 anti-c-Myc (9E10) sc-40, Santa Cruz Biotechnology) for expression monitoring and biotinylated ligand at 90 nM lysozyme (GeneTex) in PBS-F for 30 min at room temperature. The cells then underwent a second staining with fluorescently labeled secondary antibody (AlexaFluor488—goat-anti-mouse IgG1 (Life Technologies) for scFv labeling, Streptavidin-APC (SouthernBiotech) for ligand labeling) for 10 min at 4°C. Next, the cell fluorescence was measured. Data was analyzed using FlowJo.

Surface-plasmon resonance of anti-HEWL antibodies

Surface plasmon resonance experiments on the anti-HEWL Fabs were carried out on a Biacore S200 instrument (GE Healthcare, Sweden) at 25°C with HBS-N EP+ [10 mM Hepes, 150 mM NaCl, 3 mM EDTA, 0.005% vol/vol surfactant P20 (pH 7.4)]. For binding analysis, approximately 300 response units (RU) of Fab were conjugated to a CM5 sensor chip using NHS/EDC chemistry. Samples of different HEWL (Sigma) concentrations were injected over the surface at a flow rate of 30 µL/min for 120 s, and the chip was washed with buffer for 280 s (mouse Fab) or 130 s (humanized Fab). The first flow cell contained no Fab and was subtracted from the channel used to measure binding as a reference. The acquired data were analyzed using the device's software, and kinetic fits to 1:1 binding model were performed.

References

(1) *Antibody therapeutics approved or in regulatory review in the EU or US*. https://www.antibodysociety.org/resources/approved-antibodies (accessed 2021-07-20).
(2) Schroff, R. W.; Foon, K. A.; Beatty, S. M.; Oldham, R. K.; Morgan, A. C., Jr. Human Anti-Murine Immunoglobulin Responses in Patients Receiving Monoclonal Antibody

Therapy. *Cancer Res.* **1985**, *45* (2), 879–885.

(3)   Shawler, D. L.; Bartholomew, R. M.; Smith, L. M.; Dillman, R. O. Human Immune Response to Multiple Injections of Murine Monoclonal IgG. *J. Immunol.* **1985**, *135* (2), 1530–1535.

(4)   Hwang, W. Y. K.; Foote, J. Immunogenicity of Engineered Antibodies. *Methods* **2005**, *36* (1), 3–10.

(5)   Kuramochi, T.; Igawa, T.; Tsunoda, H.; Hattori, K. Humanization and Simultaneous Optimization of Monoclonal Antibody. In *Human Monoclonal Antibodies: Methods and Protocols*; Steinitz, M., Ed.; Springer New York: New York, NY, 2019; pp 213–230.

(6)   Saldanha, J. W.; Martin, A. C.; Léger, O. J. A Single Backmutation in the Human kIV Framework of a Previously Unsuccessfully Humanized Antibody Restores the Binding Activity and Increases the Secretion in Cos Cells. *Mol. Immunol.* **1999**, *36* (11-12), 709–719.

(7)   Jones, P. T.; Dear, P. H.; Foote, J.; Neuberger, M. S.; Winter, G. Replacing the Complementarity-Determining Regions in a Human Antibody with Those from a Mouse. *Nature* **1986**, *321* (6069), 522–525.

(8)   Baca, M.; Presta, L. G.; O'Connor, S. J.; Wells, J. A. Antibody Humanization Using Monovalent Phage Display. *J. Biol. Chem.* **1997**, *272* (16), 10678–10684.

(9)   Lazar, G. A.; Desjarlais, J. R.; Jacinto, J.; Karki, S.; Hammond, P. W. A Molecular Immunology Approach to Antibody Humanization and Functional Optimization. *Mol. Immunol.* **2007**, *44* (8), 1986–1998.

(10)   Padlan, E. A. A Possible Procedure for Reducing the Immunogenicity of Antibody Variable Domains While Preserving Their Ligand-Binding Properties. *Mol. Immunol.* **1991**, *28* (4-5), 489–498.

(11)   Choi, Y.; Hua, C.; Sentman, C. L.; Ackerman, M. E.; Bailey-Kellogg, C. Antibody Humanization by Structure-Based Computational Protein Design. *MAbs* **2015**, *7* (6), 1045–1057.

(12)   Kipriyanov, S. M.; Moldenhauer, G.; Martin, A. C.; Kupriyanova, O. A.; Little, M. Two Amino Acid Mutations in an Anti-Human CD3 Single Chain Fv Antibody Fragment That Affect the Yield on Bacterial Secretion but Not the Affinity. *Protein Eng.* **1997**, *10* (4), 445–453.

(13)   Safdari, Y.; Farajnia, S.; Asgharzadeh, M.; Khalili, M. Antibody Humanization Methods - a Review and Update. *Biotechnol. Genet. Eng. Rev.* **2013**, *29*, 175–186.

(14)   Warszawski, S.; Borenstein Katz, A.; Lipsh, R.; Khmelnitsky, L.; Ben Nissan, G.; Javitt, G.; Dym, O.; Unger, T.; Knop, O.; Albeck, S.; Diskin, R.; Fass, D.; Sharon, M.; Fleishman, S. J. Optimizing Antibody Affinity and Stability by the Automated Design of the Variable Light-Heavy Chain Interfaces. *PLoS Comput. Biol.* **2019**, *15* (8), e1007207.

(15)   Shire, S. J.; Shahrokh, Z.; Liu, J. Challenges in the Development of High Protein Concentration Formulations. *J. Pharm. Sci.* **2004**, *93* (6), 1390–1402.

(16)   Foote, J.; Winter, G. Antibody Framework Residues Affecting the Conformation of the Hypervariable Loops. *J. Mol. Biol.* **1992**, *224* (2), 487–499.

(17)   Makabe, K.; Nakanishi, T.; Tsumoto, K.; Tanaka, Y.; Kondo, H.; Umetsu, M.; Sone, Y.; Asano, R.; Kumagai, I. Thermodynamic Consequences of Mutations in Vernier Zone Residues of a Humanized Anti-Human Epidermal Growth Factor Receptor Murine Antibody, 528. *J. Biol. Chem.* **2008**, *283* (2), 1156–1166.

(18)   Janeway, C. A.; Travers, P.; Walport, M.; Shlomchik, M. J. *Immunobiology: The Immune System in Health and Disease*, 6th editio.; Garland Science: New York, 2005.

(19)   Lefranc, M. P. Nomenclature of the Human Immunoglobulin Kappa (IGK) Genes. *Exp. Clin. Immunogenet.* **2001**, *18* (3), 161–174.

(20)   MacCallum, R. M.; Martin, A. C.; Thornton, J. M. Antibody-Antigen Interactions: Contact Analysis and Binding Site Topography. *J. Mol. Biol.* **1996**, *262* (5), 732–745.

(21)   Dondelinger, M.; Filée, P.; Sauvage, E.; Quinting, B.; Muyldermans, S.; Galleni, M.;

Vandevenne, M. S. Understanding the Significance and Implications of Antibody Numbering and Antigen-Binding Surface/Residue Definition. *Front. Immunol.* **2018**, *9*, 2278.

(22)   Leaver-Fay, A.; Tyka, M.; Lewis, S. M.; Lange, O. F.; Thompson, J.; Jacak, R.; Kaufman, K.; Renfrew, P. D.; Smith, C. A.; Sheffler, W.; Davis, I. W.; Cooper, S.; Treuille, A.; Mandell, D. J.; Richter, F.; Ban, Y.-E. A.; Fleishman, S. J.; Corn, J. E.; Kim, D. E.; Lyskov, S.; Berrondo, M.; Mentzer, S.; Popović, Z.; Havranek, J. J.; Karanicolas, J.; Das, R.; Meiler, J.; Kortemme, T.; Gray, J. J.; Kuhlman, B.; Baker, D.; Bradley, P. ROSETTA3: An Object-Oriented Software Suite for the Simulation and Design of Macromolecules. *Methods Enzymol.* **2011**, *487*, 545–574.

(23)   O'Meara, M. J.; Leaver-Fay, A.; Tyka, M. D. M.; Stein, A.; Houlihan, K.; DiMaio, F.; Bradley, P.; Kortemme, T.; Baker, D.; Snoeyink, J.; Kuhlman, B. A Combined Covalent-Electrostatic Model of Hydrogen Bonding Improves Structure Prediction with Rosetta. *J. Chem. Theory Comput.* **2015**, *11* (2), 609–622.

(24)   Grossman, I.; Alon, A.; Ilani, T.; Fass, D. An Inhibitory Antibody Blocks the First Step in the Dithiol/disulfide Relay Mechanism of the Enzyme QSOX1. *J. Mol. Biol.* **2013**, *425* (22), 4366–4378.

(25)   Feldman, T.; Grossman-Haham, I.; Elkis, Y.; Vilela, P.; Moskovits, N.; Barshack, I.; Salame, T. M.; Fass, D.; Ilani, T. Inhibition of Fibroblast Secreted QSOX1 Perturbs Extracellular Matrix in the Tumor Microenvironment and Decreases Tumor Growth and Metastasis in Murine Cancer Models. *Oncotarget* **2020**, *11* (4), 386–398.

(26)   Starr, C. G.; Makowski, E. K.; Wu, L.; Berg, B.; Kingsbury, J. S.; Gokarn, Y. R.; Tessier, P. M. Ultradilute Measurements of Self-Association for the Identification of Antibodies with Favorable High-Concentration Solution Properties. *Mol. Pharm.* **2021**, *18* (7), 2744–2753.

(27)   Makowski, E. K.; Wu, L.; Desai, A. A.; Tessier, P. M. Highly Sensitive Detection of Antibody Nonspecific Interactions Using Flow Cytometry. *MAbs* **2021**, *13* (1), 1951426.

(28)   Almagro, J. C.; Teplyakov, A.; Luo, J.; Sweet, R. W.; Kodangattil, S.; Hernandez-Guzman, F.; Gilliland, G. L. Second Antibody Modeling Assessment (AMA-II). *Proteins: Struct. Funct. Bioinf.* **2014**, *82* (8), 1553–1562.

(29)   Sircar, A.; Kim, E. T.; Gray, J. J. RosettaAntibody: Antibody Variable Region Homology Modeling Server. *Nucleic Acids Res.* **2009**, *37* (Web Server issue), W474–W479.

(30)   Dunbar, J.; Krawczyk, K.; Leem, J.; Marks, C.; Nowak, J.; Regep, C.; Georges, G.; Kelm, S.; Popovic, B.; Deane, C. M. SAbPred: A Structure-Based Antibody Prediction Server. *Nucleic Acids Res.* **2016**, *44* (W1), W474–W478.

(31)   Leem, J.; Dunbar, J.; Georges, G.; Shi, J.; Deane, C. M. ABodyBuilder: Automated Antibody Structure Prediction with Data–driven Accuracy Estimation. *MAbs* **2016**, *8* (7), 1259–1268.

(32)   Norn, C. H.; Lapidoth, G.; Fleishman, S. J. High-Accuracy Modeling of Antibody Structures by a Search for Minimum-Energy Recombination of Backbone Fragments. *Proteins* **2017**, *85* (1), 30–38.

(33)   Kodali, P.; Schoeder, C. T.; Schmitz, S.; Crowe, J. E., , Jr; Meiler, J. RosettaCM for Antibodies with Very Long HCDR3s and Low Template Availability. *Proteins* **2021**, No. prot.26166. https://doi.org/10.1002/prot.26166.

(34)   Mirdita, M.; Schütze, K.; Moriwaki, Y.; Heo, L.; Ovchinnikov, S.; Steinegger, M. ColabFold - Making Protein Folding Accessible to All. https://doi.org/10.1101/2021.08.15.456425.

(35)   Evans, R.; O'Neill, M.; Pritzel, A.; Antropova, N.; Senior, A.; Green, T.; Žídek, A.; Bates, R.; Blackwell, S.; Yim, J.; Ronneberger, O.; Bodenstein, S.; Zielinski, M.; Bridgland, A.; Potapenko, A.; Cowie, A.; Tunyasuvunakool, K.; Jain, R.; Clancy, E.; Kohli, P.; Jumper, J.; Hassabis, D. Protein Complex Prediction with AlphaFold-Multimer. *bioRxiv*, 2022, 2021.10.04.463034. https://doi.org/10.1101/2021.10.04.463034.

(36)   Kashmiri, S. V. S.; De Pascalis, R.; Gonzales, N. R.; Schlom, J. SDR Grafting - A New Approach to Antibody Humanization. *Methods* **2005**, *36* (1), 25–34.

(37)   De Pascalis, R.; Iwahashi, M.; Tamura, M.; Padlan, E. A.; Gonzales, N. R.; Santos, A. D.; Giuliano, M.; Schuck, P.; Schlom, J.; Kashmiri, S. V. S. Grafting of "Abbreviated" Complementarity-Determining Regions Containing Specificity-Determining Residues Essential for Ligand Contact to Engineer a Less Immunogenic Humanized Monoclonal Antibody. *The Journal of Immunology* **2002**, *169* (6), 3076–3084.

(38)   Braden, B. C.; Souchon, H.; Eiselé, J. L.; Bentley, G. A.; Bhat, T. N.; Navaza, J.; Poljak, R. J. Three-Dimensional Structures of the Free and the Antigen-Complexed Fab from Monoclonal Anti-Lysozyme Antibody D44.1. *J. Mol. Biol.* **1994**, *243* (4), 767–781.

(39)   Chao, G.; Lau, W. L.; Hackel, B. J.; Sazinsky, S. L.; Lippow, S. M.; Wittrup, K. D. Isolating and Engineering Human Antibodies Using Yeast Surface Display. *Nat. Protoc.* **2006**, *1* (2), 755–768.

(40)   Knappik, A.; Ge, L.; Honegger, A.; Pack, P.; Fischer, M.; Wellnhofer, G.; Hoess, A.; Wölle, J.; Plückthun, A.; Virnekäs, B. Fully Synthetic Human Combinatorial Antibody Libraries (HuCAL) Based on Modular Consensus Frameworks and CDRs Randomized with Trinucleotides. *J. Mol. Biol.* **2000**, *296* (1), 57–86.

(41)   Apgar, J. R.; Mader, M.; Agostinelli, R.; Benard, S.; Bialek, P.; Johnson, M.; Gao, Y.; Krebs, M.; Owens, J.; Parris, K.; St. Andre, M.; Svenson, K.; Morris, C.; Tchistiakova, L. Beyond CDR-Grafting: Structure-Guided Humanization of Framework and CDR Regions of an Anti-Myostatin Antibody. *MAbs* **2016**, *8* (7), 1302–1318.

(42)   Baran, D.; Pszolla, M. G.; Lapidoth, G. D.; Norn, C.; Dym, O.; Unger, T.; Albeck, S.; Tyka, M. D.; Fleishman, S. J. Principles for Computational Design of Binding Antibodies. *Proc. Natl. Acad. Sci. U. S. A.* **2017**, *114* (41), 10900–10905.

(43)   Borenstein-Katz, A.; Warszawski, S.; Amon, R.; Eilon, M.; Cohen-Dvashi, H.; Leviatan Ben-Arye, S.; Tasnima, N.; Yu, H.; Chen, X.; Padler-Karavani, V.; Fleishman, S. J.; Diskin, R. Biomolecular Recognition of the Glycan Neoantigen CA19-9 by Distinct Antibodies. *J. Mol. Biol.* **2021**, *433* (15), 167099.

(44)   Eddy, S. R. Profile Hidden Markov Models. *Bioinformatics* **1998**, *14* (9), 755–763.

(45)   Fleishman, S. J.; Leaver-Fay, A.; Corn, J. E.; Strauch, E.-M.; Khare, S. D.; Koga, N.; Ashworth, J.; Murphy, P.; Richter, F.; Lemmon, G.; Meiler, J.; Baker, D. RosettaScripts: A Scripting Language Interface to the Rosetta Macromolecular Modeling Suite. *PLoS One* **2011**, *6* (6), e20161.

(46)   Ye, J.; Ma, N.; Madden, T. L.; Ostell, J. M. IgBLAST: An Immunoglobulin Variable Domain Sequence Analysis Tool. *Nucleic Acids Res.* **2013**, *41* (Web Server issue), W34–W40.

(47)   Baker, N. A.; Sept, D.; Joseph, S.; Holst, M. J.; McCammon, J. A. Electrostatics of Nanosystems: Application to Microtubules and the Ribosome. *Proc. Natl. Acad. Sci. U. S. A.* **2001**, *98* (18), 10037–10041.

(48)   Alon, A.; Grossman, I.; Gat, Y.; Kodali, V. K.; DiMaio, F.; Mehlman, T.; Haran, G.; Baker, D.; Thorpe, C.; Fass, D. The Dynamic Disulphide Relay of Quiescin Sulphydryl Oxidase. *Nature* **2012**, *488* (7411), 414–418.

(49)   Javitt, G.; Kinzel, A.; Reznik, N.; Fass, D. Conformational Switches and Redox Properties of the Colon Cancer-associated Human Lectin ZG16. *FEBS J.* **2021**.

(50)   Adams, P. D.; Afonine, P. V.; Bunkóczi, G.; Chen, V. B.; Davis, I. W.; Echols, N.; Headd, J. J.; Hung, L.-W.; Kapral, G. J.; Grosse-Kunstleve, R. W.; McCoy, A. J.; Moriarty, N. W.; Oeffner, R.; Read, R. J.; Richardson, D. C.; Richardson, J. S.; Terwilliger, T. C.; Zwart, P. H. PHENIX: A Comprehensive Python-Based System for Macromolecular Structure Solution. *Acta Crystallogr. D Biol. Crystallogr.* **2010**, *66* (Pt 2), 213–221.

(51)   Emsley, P.; Lohkamp, B.; Scott, W. G.; Cowtan, K. Features and Development of Coot. *Acta Crystallogr. D Biol. Crystallogr.* **2010**, *66* (Pt 4), 486–501.

Supplement to: Reliable energy-based antibody humanization and stabilization

Ariel Tennenhouse, Lev Khmelnitsky, Noa Yeshaya, Ashish Noronha, Moshit Lindzen, Razi Khalaila, Ira Zaretsky, Yael Fridmann Sirkis, Yael Galon-Wolfenson, Jakub Abramson, Yosef Yarden, Deborah Fass, Sarel J. Fleishman

Antibody name	Light chain V gene Sequence Identity (%)	Heavy chain V gene sequence identity (%)	Full Fv sequence identity (%)
Alemtuzumab	86.3	73	79.2
Atezolizumab	86.3	86.7	86.5
Belantamab	90.5	83.7	86.9
Benralizumab	87.4	78.6	82.7
Bevacizumab	87.4	72.4	79.4
Brolucizumab	87.6	79.4	83.3
Certolizumab	85.3	76.5	80.7
Crizanlizumab	86.9	81.6	84.1
Dostarlimab	85.3	93.8	89.7
Eculizumab	84.2	83.7	83.9
Elotuzumab	84.2	81.6	82.8
Emicizumab 1	80	75.5	77.6
Emicizumab 2	80	87.8	84.2
Eptinezumab	86.2	81.4	83.8
Fremanezumab	85.3	85.7	85.5
Galcanezumab	87.4	80.6	83.8
Gemtuzumab	77.8	72.9	75.3
Ibalizumab	92.9	76.5	84.3
Idarucizumab	88	82.3	85.0
Inebilizumab	79.8	83.7	81.8

Inotuzumab	73	82.7	78.0
Ixekizumab	89	82.7	85.8
Loncastuximab	66	74.5	70.6
Mepolizumab	91.1	74.7	82.7
Mogamulizumab	81	83.7	82.4
Natalizumab	80.9	83.7	82.4
Naxitamab	78.9	72.2	75.3
Obinutuzumab	86	84.7	85.3
Ocrelizumab	86.5	80.6	83.3
Omalizumab	86.9	78.6	82.6
Palivizumab	81.9	87.8	85.0
Pembrolizumab	80.8	79.6	80.2
Pertuzumab	84.2	77.6	80.7
Polatuzumab	85.9	75.5	80.6
Ranibizumab	86.3	71.4	78.3
Ravulizumab	84.2	81.6	82.8
Reslizumab	87.4	76.3	81.6
Risankizumab	80	76.3	78.0
Romosozumab	89.5	87.8	88.6
Sacituzumab	80	85.7	83.0
Satralizumab	82.1	76.5	79.2
Sutimlimab	78.1	89.8	84.2
Tafasitamab	71	73.5	72.3
Tildrakizumab	85.3	81.6	83.4
Tocilizumab	89.5	82.7	85.9
Trastuzumab	85.3	81.6	83.3
Vedolizumab	84	84.7	84.4

median	85.3	81.6	82.8
mean	83.9	80.5	82.1

Supplementary Table 1: Sequence identity to the human germline of 47 FDA approved humanized antibodies[1]. % identity computed using IgBLAST[2]. Antibody sequences were taken from Thera-SabDab[3].

	L1	**L2**	**L3**	**H1**	**H2**	**H3**
CUMAb	24-34	46-55	89-97	24-35B	47-58	93-102
Kabat	24-34	50-56	89-97	31-35B	50-65	95-102
AHO	25-32	50-61	91-96	24-35B	51-66	95-101

Supplementary Table 2. CUMAb CDR definitions compared to Kabat and AHO schemes[4,5].

Data collection

Space group	$P2_1$
Cell dimensions	
a, b, c (Å)	55.58, 98.28, 63.12
α, β, γ (°)	90, 101.15, 90
Asymmetric unit	1 Fab-QSOX1trx complex
Resolution (Å)	49.14 - 2.10 (2.18 - 2.10)*
Measured reflections	102,949 (10,755)
Unique reflections	37,687 (3817)
Completeness (%)	97.0 (98.6)
Redundancy	2.7 (2.8)
< I/σI >	6.8 (0.6)
R_{merge}	0.105 (1.637)
R_{pim}	0.075 (1.127)

Refinement

Resolution (Å)	49.14 - 2.10 (2.12 - 2.10)
Reflections in working set	69,556 (2257)
Reflections in test set	7728 (254)
R_{work}/R_{free}	0.198/0.259
Number of protein atoms	5,061
Number of water molecules	211
Mean B-factor	46
Root mean square deviations	
Bond length (Å)	0.007
Bond angle (°)	0.908
Ramachandran plot	
Favored regions (%)	97.1
Additional allowed regions (%)	2.9
Disallowed regions (%)	0

Supplementary Table 3: Crystallographic data and refinement statistics for the complex between the Fab fragment of hαQSOX1.4 and QSOX1trx (PDB code 8AON). Values in parentheses are for the highest resolution bins.

Antibody	Closest Heavy V Germline Match	Closest Light V Germline Match
mɑQSOX1	IGHV4-4 (57%)	IGKV1-33 (66%)
hɑQSOX1.1	IGHV3-33 (80%)	IGKV4-1 (81%)
hɑQSOX1.2	IGHV2-5 (86%)	IGKV4-1 (81%)
hɑQSOX1.3	IGHV3-49 (81%)	IGKVD-29 (78%)
hɑQSOX1.4	IGHV3-49 (81%)	IGKV4-1 (81%)
Consensus H4-K1	IGHV4-4 (85%)	IGKV1-NL1 (84%)
mɑAXL	IGHV3-23 (79%)	IGKV1-16 (62%)
hɑAXL	IGHV3-23 (89%)	IGKV1D-13 (84%)
mɑHEWL	IGHV1-46 (67%)	IGKV3-15 (64%)
hɑHEWL	IGHV1-69-2 (90%)	IGKV1-39 (96%)

Supplementary Table 5: Sequence identity between parental antibodies (first in each block) and designs to the human germline. % identity computed using IgBLAST[2].

Supplementary Figure 1: (**A**) Dot-blot analysis of 15 CUMAb designs. 12 designs exhibit comparable expression levels to the AbLIFT18 design of the anti-QSOX1 antibody[6]. CUMAb designs are labeled according to heavy chain V-gene subgroup and then by light chain subgroup according to IMGT. If there are two chains from the same subgroup, they are labeled as "a" or "b" arbitrarily. "Mock" refers to a control transfection where no DNA was added. "Prev" refers to a construct from a prior version of CUMAb. (**B**) Functional screen of 15 CUMAb designs, labeled as in A. Presence of an upper band, corresponding to MBP-ZG16 modified with PEG-mal 5000, indicates remaining unoxidized MBP-ZG16 as a result of inhibited QSOX1 activity. As evident from the "no QSOX1" lane, the MBP-ZG16 used in the experiment was not fully reduced, but the reduced population was sufficient to distinguish between the mock-transfected sample, which lacks antibody, and the antibody-containing samples, which all show some PEG-mal-modified species indicative of QSOX1 inhibition. Designs marked with * were further analyzed. (**C**) Titration of relative QSOX1 activity at different antibody concentrations for mαQSOX1 and four CUMAb designs (aggregate data shown in main Figure 2B). Shown is the average of two repeats. (**D**) Thermal denaturation of mαQSOX1 and four CUMAb designs using nano differential scanning fluorimetry. Shown is the average of two repeats. (**E**) Crystal structure of oxidoreductase fragment of human QSOX1 in complex with parental mouse antibody (QSOX1 in dark gray, parental mouse antibody Fv in light gray, PDB entry 4IJ3) aligned with crystal structure of oxidoreductase fragment of human QSOX1 in complex with hαQSOX1.4 (QSOX1 in purple, hαQSOX1.4 Fv in magenta, PDB entry 8AON. Structures are nearly identical (0.7 Å) despite 51 Fv mutations.

Supplementary Figure 2: (**A**) Expression levels of negative control (the G6 anti-VEGF[7] antibody), mouse anti-HEWL antibody (mαHEWL), and 6 CUMAb designs, all formatted as scFv. (**B**) Binding levels to HEWL of the eight antibodies.

Supplementary Figure 3: Dot-blot analysis of anti-QSOX1 CDRs grafted onto most homologous consensus frameworks IGKV1-IGHV4 compared to a mock transfection and well-expressed AbLIFT design 3[6].

Supplemental references

(1) *Antibody therapeutics approved or in regulatory review in the EU or US*. The Antibody Society. https://www.antibodysociety.org/resources/approved-antibodies/ (accessed 2022-08-02).

(2) Ye, J.; Ma, N.; Madden, T. L.; Ostell, J. M. IgBLAST: An Immunoglobulin Variable Domain Sequence Analysis Tool. *Nucleic Acids Res.* **2013**, *41* (Web Server issue), W34–W40.

(3) Raybould, M. I. J.; Marks, C.; Lewis, A. P.; Shi, J.; Bujotzek, A.; Taddese, B.; Deane, C. M. Thera-SAbDab: The Therapeutic Structural Antibody Database. *Nucleic Acids Res.* **2020**, *48* (D1), D383–D388.

(4) MacCallum, R. M.; Martin, A. C.; Thornton, J. M. Antibody-Antigen Interactions: Contact Analysis and Binding Site Topography. *J. Mol. Biol.* **1996**, *262* (5), 732–745.

(5) Dondelinger, M.; Filée, P.; Sauvage, E.; Quinting, B.; Muyldermans, S.; Galleni, M.; Vandevenne, M. S. Understanding the Significance and Implications of Antibody Numbering and Antigen-Binding Surface/Residue Definition. *Front. Immunol.* **2018**, *9*, 2278.

(6) Warszawski, S.; Borenstein Katz, A.; Lipsh, R.; Khmelnitsky, L.; Ben Nissan, G.; Javitt, G.; Dym, O.; Unger, T.; Knop, O.; Albeck, S.; Diskin, R.; Fass, D.; Sharon, M.; Fleishman, S. J. Optimizing Antibody Affinity and Stability by the Automated Design of the Variable Light-Heavy Chain Interfaces. *PLoS Comput. Biol.* **2019**, *15* (8), e1007207.

(7) Liang, W.-C.; Wu, X.; Peale, F. V.; Lee, C. V.; Meng, Y. G.; Gutierrez, J.; Fu, L.; Malik, A. K.; Gerber, H.-P.; Ferrara, N.; Fuh, G. Cross-Species Vascular Endothelial Growth Factor (VEGF)-Blocking Antibodies Completely Inhibit the Growth of Human Tumor Xenografts and Measure the Contribution of Stromal VEGF. *J. Biol. Chem.* **2006**, *281* (2), 951–961.