Development and applications of a CRISPR activation system for facile genetic overexpression in *Candida albicans*

Nicholas C. Gervais[1], Alyssa A. La Bella[2], Lauren F. Wensing[1], Victoria Acquaviva[1], Madison Best[1], Jehoshua Sharma[1], Deeva Uthayakumar[1][†], Alejandro Chavez[3], Felipe Santiago-Tirado[2], Ana L. Flores-Mireles[2], and Rebecca S. Shapiro[1]*

[1]Department of Molecular and Cellular Biology, University of Guelph, Guelph, Ontario, Canada

[2]Department of Biological Sciences, University of Notre Dame, Notre Dame, Indiana, USA

[3]Department of Pathology and Cell Biology, Columbia University College of Physicians and Surgeons, New York, New York, USA

[†]Current affiliation: Department of Immunology, University of Toronto, Toronto, Ontario, Canada

*Correspondence: shapiror@uoguelph.ca

Abstract

For the fungal pathogen *Candida albicans*, genetic overexpression readily occurs via a diversity of genomic alterations, such as aneuploidy and gain-of-function mutations, with important consequences for host adaptation, virulence, and evolution of antifungal drug resistance. Given the important role of overexpression on *C. albicans* biology, it is critical to develop and harness genetic tools that enable the analysis of genes expressed at high levels in the fungal cell. Here, we describe the development, optimization, and application of a novel, single-plasmid-based CRISPR activation (CRISPRa) platform for targeted genetic overexpression in *C. albicans*. This CRISPRa platform employs a guide RNA to target an activator complex to the promoter region of a gene of interest, thus driving transcriptional expression of that gene. Our CRISPRa system for *C. albicans* is unique, as it is asingle-plasmid system designed for rapid Golden Gate cloning and facile fungal strain generation in diverse strain backgrounds. Our CRISPRa system exploits a nuclease-dead version of Cas9 (dCas9) fused to the tripartite activator complex VP64-p65-Rta (VPR), and a Golden Gate site for efficient guide RNA cloning to accomplish this overexpression. Using this system, we demonstrate the ability of CRISPRa to drive high levels of gene expression in *C. albicans*, and we assess optimal guide RNA targeting for robust and constitutive overexpression. We find that we are able to obtain a range of overexpression from ~2-25-fold over wild-type expression levels, and demonstrate the specificity of the system via RNA sequencing. We further demonstrate the application of CRISPRa to overexpress genes involved in fungal pathogenesis and drug resistance and detect corresponding phenotypic alterations in these traits. Together, this tool will facilitate a broad range of applications for the study of *C. albicans* genetic overexpression.

Keywords

Fungal genetics, CRISPR activation, Candida albicans, Gene regulation, Fungal pathogens

Introduction

Fungal pathogens are an increasingly important cause of human illness, with significant impacts on morbidity and mortality (Fisher *et al.* 2020). *Candida albicans* is an opportunistic fungal pathogen that inhabits the microbiome of most healthy adults as a commensal organism (Gow *et al.* 2011; Pérez 2019). However, it is also responsible for invasive infections, particularly amongst immunocompromised individuals, which are often fatal (Bongomin *et al.* 2017). Instances of *C. albicans* infections are becoming more prevalent globally, in part due to an increasing population of susceptible elderly and immunocompromised individuals (Bongomin *et al.* 2017). Illustratively, systemic *Candida* infections have been identified in up to 15% of patients hospitalized for COVID-19, and patients with invasive candidiasis spent longer in the hospital Intensive Care Unit (ICU) on average, and had more severe symptoms than those COVID-19 patients without a *Candida* infection (Segrelles-Calvo *et al.* 2021). The economic cost of *Candida* infections is estimated to be ~1.4 billion dollars yearly in the United States alone, hospitalizing more than 26,000 people (Benedict *et al.* 2018). With limited development of novel antifungal drugs (Perfect 2017) and increasing prevalence of antifungal drug-resistant strains and species (Sanguinetti *et al.* 2015; Arendrup and Patterson 2017; Geddes-McAlister and Shapiro 2018), *Candida* pathogens remain a serious threat to human health.

Given the importance of *C. albicans* and other fungal pathogens, it is imperative that tools are developed and applied to manipulate these organisms and dissect genetic functions. For *C. albicans*, along with countless other microbial species, CRISPR-based genetic manipulation tools have revolutionized the ability of scientists to modify and study gene function (Shapiro *et al.* 2018a; Uthayakumar *et al.* 2021). Such CRISPR platforms typically rely on a CRISPR-associated endonuclease protein, often Cas9, targeted to a genomic region of interest via a single guide RNA (sgRNA), resulting in targeted DNA double-strand breaks (DSBs), and subsequent repair via non-homologous end-joining or homology-directed repair via a donor

DNA repair template (Adli 2018; Bock *et al.* 2022). Since the initial development of CRISPR tools for genetic editing in *C. albicans* (Vyas *et al.* 2015), numerous systems have been developed to modify this fungal genome with efficiency and specificity (Min *et al.* 2016, 2018; Huang and Mitchell 2017; Ng and Dean 2017; Nguyen *et al.* 2017; Shapiro *et al.* 2018b; Vyas *et al.* 2018; Román *et al.* 2019b; Morio *et al.* 2020; Uthayakumar *et al.* 2021), and to build and screen genetic mutant libraries (Shapiro *et al.* 2018b; Rosiana *et al.* 2021). Together, these gene editing systems have played an important role in providing novel biological insight into this fungal pathogen.

In addition to these canonical CRISPR-based genetic modification systems that introduce mutations, insertions, or deletions into the genome, alternative CRISPR technologies have been developed that modulate the expression of *C. albicans* genes (Román *et al.* 2019a; Wensing *et al.* 2019). These CRISPR systems rely on the inactivation of the Cas9 endonuclease, to generate a nuclease-dead (dCas9) protein, which can be fused to constructs from a transcriptional repressor or activator that generate CRISPR interference (CRISPRi) or CRISPR activation (CRISPRa) systems, respectively (Dominguez *et al.* 2016; Wang *et al.* 2016). These dCas9 fusions, when targeted to the promoter region of a gene via the sgRNA, can repress or activate gene transcription from a gene of interest, and have been widely applied in diverse organisms (Peters *et al.* 2015; Kampmann 2018; Xu and Qi 2019; Schultenkämper *et al.* 2020; Todor *et al.* 2021; Bock *et al.* 2022). In *C. albicans*, these modified CRISPR systems have enabled the modulation and study of important fungal genes, including essential genes (Román *et al.* 2019a; Wensing *et al.* 2019; Wensing and Shapiro 2022).

While a majority of CRISPR systems and other genetic manipulation platforms focus on gene mutation, deletion, or downregulation, it is also critical to develop techniques that enable the upregulation or overexpression of genes. Indeed, genetic overexpression in *C. albicans* occurs readily via gene copy number amplification and aneuploidy, and these copy number amplification events are frequently observed in genetically diverse clinical isolates (Coste *et al.*

2006; Forche *et al.* 2018; Todd *et al.* 2019). The upregulation of gene expression plays a critical role in diverse aspects of fungal biology (Rai *et al.* 2021), including the evolution of antifungal drug resistance (Selmecki *et al.* 2008; Sanglard *et al.* 2009; Schubert *et al.* 2011; Flowers *et al.* 2012; Lohberger *et al.* 2014; Todd and Selmecki 2020), host colonization (Prieto *et al.* 2017; Znaidi *et al.* 2018), and virulence (Fu *et al.* 2008; Lohberger *et al.* 2014). Additionally, in other microbial systems, including bacteria, protozoa, and the model yeast *Saccharomyces cerevisiae*, genetic overexpression libraries have proven to be critical tools for identifying targets of drugs with uncharacterized mechanisms of action, as mutants overexpressing the drug target are typically more resistant to the drug in question (Li *et al.* 2004; Luesch *et al.* 2005; Smith *et al.* 2010; Begolo *et al.* 2014). Overexpression screens can also help dissect molecular pathways (Sopko *et al.* 2006; Sahni *et al.* 2010; Polvi *et al.* 2019). In *C. albicans*, several techniques exist for genetic overexpression (Rai *et al.* 2021), including promoter replacement strategies (Delgado *et al.* 2003; Park and Morschhäuser 2005; Eckert and Mühlschlegel 2009; Sahni *et al.* 2010), the generation of ORFeome collection strains (Chauvel *et al.* 2012; Legrand *et al.* 2018), and CRISPR-based methods (Román *et al.* 2019a)*.* While these these tools have made important contributions to understanding overexpression in *C. albicans*, there remains a need for simple and rapid methods to efficiently target fungal genes of interest for overexpression, and to do so in diverse strain backgrounds.

Here, we introduce a CRISPR-based tool for genetic overexpression in *C. albicans*, to bolster the existing functional genetic toolbox available to manipulate genes in this critical fungal pathogen. Our CRISPRa tool exploits a *C. albicans*-optimized tripartite activator complex fused to dCas9, which was previously developed for highly-efficient transcriptional regulation in *S. cerevisiae* (Chavez *et al.* 2015). This CRISPRa system is distinct from other CRISPR-based gene activation systems for *C. albicans*, as it is an efficient single-plasmid system designed for rapid Golden Gate cloning and facile fungal strain generation, which can be readily applied to diverse strain backgrounds including clinical isolates. We test CRISPRa guide targeting

principles based on predicted transcriptional start sites, confirm on-target guide efficiency, and demonstrate the ability of this CRISPRa system to drive high levels of expression of *C. albicans* genes. Using this optimized CRISPRa technique, we validate its ability to drive overexpression of genes with established roles in antifungal drug resistance and biofilm formation. Together, this work introduces a novel CRISPRa system for genetic overexpression in *C. albicans*, with a wide range of future applications.

Results

Development of a single-plasmid CRISPRa tool in C. albicans

First, we designed a single-plasmid CRISPRa system for targeted gene activation, optimized for *C. albicans*. We exploited an integrating plasmid backbone, which we previously used for CRISPRi in *C. albicans*, containing a codon-optimized dCas9 and Golden Gate assembly-compatible sgRNA cloning site (Wensing *et al.* 2019). This plasmid can be linearized to integrate at the *C. albicans NEUT5L* locus (a large intergenic region whose disruption does not affect fungal fitness (Gerami-Nejad *et al.* 2013)), and contains a dual SapI cloning site for facile and highly-efficient Golden Gate cloning of the sgRNA targeting sequence. This plasmid also contains a dominant nourseothricin resistance (NATr) cassette, to enable strain generation in a diversity of fungal strains, including clinical isolates.

To generate a version of dCas9 that could lead to strong constitutive activation of genes of interest, we generated a plasmid with dCas9 fused to the tripartite activator complex, VP64-p65-Rta (VPR) (**FIGURE 1a,b**). VPR consists of three previously-described transcriptional activators: (1) VP64, a viral transcriptional activator domain composed of four tandem copies of VP16 (Herpes Simplex Viral Protein 16), which has been used extensively in CRISPRa systems (La Russa and Qi 2015); (2) p65, the NF-κB subunit responsible for the strong transcription activation in mammalian cells (Schmitz and Baeuerle 1991); and (3) Rta, the Epstein-Barr virus R transactivator, with strong transcriptional activation properties (Hardwick *et al.* 1992) (**FIGURE**

1a,b). Together the VPR activator complex fused to dCas9 and targetted to the promoter region of genes has previously been demonstrated to generate strong gene activation in *S. cerevisiae* (Chavez *et al.* 2015). Exploiting the ability of dCas9 to bind and be targeted to genomic loci via sgRNAs enables us to design sgRNAs that will target dCas9 along with these strong transcriptional activators to any gene promoter of interest.

This plasmid was generated by homology-based cloning and validated via Sanger sequencing, and has been deposited to Addgene (**TABLE S1**, Addgene plasmid # 182707). To confirm if the expression of the dCas9-VPR construct had an impact on fungal fitness, we compared the growth of a wild-type *C. albicans* strain with a strain transformed with the dCas9-VPR CRISPRa plasmid (expressing a non-targeting sgRNA: a random sequence of nucleotides that have no predicted complementarity to any region in the *C. albicans* genome). We found no difference in growth between these strains (**FIGURE 1c**), suggesting no significant impact of the CRISPRa system on *C. albicans* fitness.

Validation and sgRNA design principles for CRISPRa overexpression in C. albicans

Next, we aimed to validate that the CRISPRa system could successfully overexpress genes in *C. albicans*, and sought to determine the optimal sgRNA design strategy to maximize genetic regulation. Our previous *C. albicans* CRISPRi system exploited sgRNAs ~ 50 - 150bp upstream of the start codon to functionally repress genes of interest (Wensing *et al.* 2019). However, CRISPRa systems are known to have different optimal sgRNA targeting regions (Jensen 2018; Sanson *et al.* 2018), and our previous design did not take into account the transcription start site (TSS), which is known to be an important predictor of regulatory efficiency for CRISPRi/a systems (Smith *et al.* 2016; Jensen 2018), but which had not been robustly characterized across all *C. albicans* genes at the time. More recently, the TSS for many *C. albicans* genes has been predicted (McMillan *et al.* 2019). Thus, we sought to determine the optimal design for CRISPRa sgRNAs based on the start codon and the TSS. We identified three

C. albicans genes with a predicted TSS that was particularly far upstream of the start codon compared to all other genes in the genome (801-952 bp upstream): *KSP1*, *ALS3,* and *FGR28* . We designed six sgRNAs for each gene, targeting upstream of the start codon (as previously designed for CRISPRi (Wensing *et al.* 2019)), as well as upstream of the TSS (based on predicted TSS (McMillan *et al.* 2019)).

We generated CRISPRa plasmids based on these sgRNAs, used these to create *C. albicans* strains, and monitored overexpression of each gene via quantitative reverse-transcription PCR (RT-qPCR). This analysis validated our novel CRISPRa system and its ability to overexpress *C. albicans* genes ~ 2 - 12-fold, based on different sgRNA designs and target genes (**FIGURE 2**). We found that in some instances, such as for *KSP1*, the TSS was in fact a better predictor of CRISPRa overexpression, as the most robust overexpression of this gene was observed with sgRNAs targetted upstream of the TSS, while no overexpression was observed based on targeting upstream of the start codon (**FIGURE 2a**). In other instances, such as for *ALS3*, sgRNA targeting upstream of the start codon led to very high levels of overexpression, while sgRNAs targeted to the TSS region led to moderate overexpression (**FIGURE 2b**). For *FGR28*, modest overexpression (~2-fold) was achieved with sgRNAs targeted to either the start codon or the TSS upstream regions (**FIGURE 2c**), together suggesting that optimal sgRNA design can be predicted based on targeting the TSS and/or start codon, but varies by gene. This data also suggests that our CRISPRa tool may have additional applications in the functional assessment of key transcriptional regulatory regions in *C. albicans*. These differences in sgRNA targeting may be based on differences in overall sgRNA efficiency or may be attributed to inaccurate TSS predictions, or complex promoter dynamics and the preferential use of different TSS regions in different conditions (McMillan *et al.* 2019).

Use of CRISPRa to validate antifungal drug resistance phenotypes

To validate that the CRISPRa system could recapitulate phenotypes associated with *C. albicans* gene overexpression, we targeted genes with known phenotypes associated with high levels of expression. Initially, we focused on *CDR1*, a well-characterized *C. albicans* gene, which encodes a well-characterized drug efflux pump (Prasad *et al.* 2015), and whose overexpression is known to play a role in drug resistance to azole antifungals in laboratory and clinical isolates (Prasad *et al.* 1995; Sanglard *et al.* 1995; White 1997; Marr *et al.* 1998, 2001; Niimi *et al.* 2004; Manoharlal *et al.* 2008). To determine an optimal range in which to target our sgRNA in the *CDR1* promoter, we designed three sgRNAs targeting loci approximately 90 bp, 200 bp, and 290 bp upstream of the *CDR1* TSS. These sgRNAs were each cloned into our CRISPRa dCas9-VPR backbone and transformed into *C. albicans*. Next, we quantified the expression of *CDR1* in the three different fungal strains via RT-qPCR, compared with a control strain containing the CRISPRa plasmid with a non-targeting sgRNA. We found that targeting the dCas9-VPR construct to the *CDR1* promoter enhanced transcription by 2.9 - 5.6 fold (**FIGURE 3a**), indicating that this CRISPRa system is indeed able to drive enhanced levels of gene expression from a target gene of interest. For *CDR1*, we found the highest level of overexpression was obtained by targeting the dCas9-VPR construct ~200 bp upstream of the TSS (**FIGURE 3a**), which helped inform the design of sgRNAs for subsequent target genes.

Next, we sought to determine if *CDR1*-overexpressing strains had measurable changes in antifungal drug resistance phenotypes. We performed minimum inhibitory concentration (MIC) assays with these strains in the presence of the antifungal drug, fluconazole. We found that strains overexpressing *CDR1* exhibited increased resistance to fluconazole compared with the control strain, and that higher levels of overexpression corresponded with higher levels of resistance (**FIGURE 3b**). Together, this confirms that our CRISPRa system can be exploited to functionally overexpress target genes of interest, and recapitulate phenotypes associated with high levels of gene expression. Further, targeting dCas9-VPR to different promoter loci leads to

variable levels of overexpression, which could be used to study correlations between expression levels and associated phenotypes.

Validation of CRISPRa overexpression targeting fungal pathogenesis traits

Next, we sought to validate the CRISPRa system as a means to assess *C. albicans* genes involved in fungal pathogenesis traits. We focused on *ALS1*, a well-characterized *C. albicans* adhesin gene, with important roles in adherence, biofilm formation, and virulence (Fu *et al.* 2002; Kamai *et al.* 2002; Alberti-Segui *et al.* 2004; Zhao *et al.* 2004; Nobile *et al.* 2008; Rosiana *et al.* 2021). Previous work has demonstrated that overexpression of *ALS1* via promoter replacement increases fungal adherence (Fu *et al.* 2002), therefore we chose to overexpress this gene using our CRISPRa system. We designed and cloned two sgRNAs targeting approximately 130bp and 260bp upstream of the *ALS1* TSS, and generated *C. albicans ALS1* CRISPRa strains with each of these constructs. We performed RT-qPCR to confirm the overexpression of these strains, which enhanced the expression of *ALS1* by ~25 and ~6 fold, respectively (**FIGURE 4a**).

To confirm phenotypes associated with overexpression of *ALS1*, we performed two distinct biofilm growth assays that compared the *ALS1* CRISPRa strain to the CRISPRa control strain. In the first assay, we allowed biofilms to establish over 24h in flat-bottom 96-well polystyrene plates, removed planktonic cells, and quantified the metabolic activity of the biofilm via XTT reduction, relative to the planktonic cell population. We found that the *ALS1* CRISPRa strain formed significantly more robust biofilms, relative to the control strain ($P < 0.0005$; **FIGURE 4b**). Interestingly, biofilm growth was not correlated with *ALS1* overexpression levels, as the strain with ~25-fold *ALS1* overexpression formed less robust biofilms compared with the strain expressing ~6-fold more *ALS1*. This suggests that very high levels of *ALS1* may ultimately impair biofilm formation, or that the level of *ALS1* expression varies in the conditions used for

RT-qPCR analysis compared with biofilm growth, perhaps due to condition-specific repositioning in the TSS.

Next, to mimic conditions associated with fungal biofilm formation in catheter-associated urinary tract infections (CAUTIs), we used a urine-based assay and analyzed biofilm formation of a *C. albicans* strain overexpressing *ALS1*, relative to its control strain, using crystal violet staining. For this assay, we followed up on the *ALS1* CRISPRa strain with ~6-fold overexpression of *ALS1*, which had more robust biofilm growth than the control strain (**FIGURE 4a,b**). The *C. albicans* strains were incubated for 48h in human urine supplemented with bovine serum albumin (BSA) in fibrinogen-coated 96-well plates. Fibrinogen has been shown to be a critical protein for pathogen binding in CAUTIs, by providing a scaffold for biofilm formation and a nutritional source for pathogens (Flores-Mireles *et al.* 2016; La Bella *et al.* 2021; Andersen *et al.* 2022). The *ALS1* CRISPRa overexpression strain showed significantly greater biofilm formation compared to the corresponding wild-type strain ($P < 0.0001$; **FIGURE 4c**). This difference in biofilm formation suggests that *ALS1* promotes binding to fibrinogen and plays a vital role in biofilm formation in urine.

To assess the specificity of this CRISPRa system and ensure no off-target effects from the dCas9-sgRNA complex, we monitored global changes in transcription in the *ALS1* CRISPRa strain via RNA-sequencing (RNA-seq). Since *ALS1* is a member of the large agglutinin-like sequence (ALS) gene family (Hoyer *et al.* 2008; Hoyer and Cota 2016), it may be a likely candidate for any potential off-target effects. We also considered *ALS1* as a strong candidate for this RNA-seq off-target profiling since *ALS1* is not known to be a transcriptional regulator, and therefore its overexpression is unlikely to lead to direct downstream transcriptional alterations that would obscure our analysis of CRISPRa off-target effects. Therefore, we performed RNA-seq on the validated CRISPRa strain overexpressing *ALS1* by ~25-fold as well as on a control strain containing the dCas9 cassette and a non-targeting sgRNA when grown in standard laboratory conditions. This analysis confirmed the specific overexpression of *ALS1*

(log2 fold change ~3.3 via RNA-seq analysis, adjusted *P*-value 3.07E-142; **FIGURE 4d, TABLE S2**). Five additional transcripts were identified as significantly modulated (adjusted *P*-value < 0.05, absolute log2 fold change > 1; **TABLE S2**). However, their altered expression was much less significant than *ALS1* (**FIGURE 4d**), none of them having an adjusted P-value of less than 0.03, differing from the adjusted P-value of 3e-142 for *ALS1.* We suspect these additional transcripts' altered expression is likely unrelated to the CRISPRa system based on the fact that 4/5 genes were down-regulated instead of up-regulated, and none of the genes are in proximity to the *ALS1* locus, nor do they appear to share any sgRNA target similarities in their promoter regions. Together this suggests that this CRISPRa system is likely specific with regards to gene targeting.

Discussion

Here, we present the development, validation, and application of a single-plasmid-based CRISPRa platform for efficient genetic overexpression in *C. albicans*. We demonstrate the ability of this system to overexpress genes ~2-20-fold, and recapitulate important biological phenotypes associated with genetic overexpression, including biofilm formation and antifungal drug resistance. In addition, we assess sgRNA design principles based on predicted TSS, and validate the specificity of this system by monitoring off-target effects via RNA-seq. Together, this work demonstrates an effective, efficient, and specific tool for genetic overexpression in *C. albicans* to enhance the available techniques for genetic modulation in this important fungal pathogen.

Studying genetic overexpression is an important strategy for functional genetic analysis, as overexpressing genes can mimic natural biological states where genes are expressed at high levels. In *C. albicans*, there are numerous scenarios in which the overexpression of genes occurs, including mutations that modify the function of transcriptional regulators and promote the expression of downstream genes (Coste *et al.* 2004, 2006), as well as aneuploidies such as

trisomies (Selmecki *et al.* 2006; Forche *et al.* 2009, 2018; Ford *et al.* 2015; Li *et al.* 2015; Anderson *et al.* 2017; Ene *et al.* 2018), isochromosome formation (Selmecki *et al.* 2006, 2008), and copy number variations (Ford *et al.* 2015; Todd and Selmecki 2020). These overexpression events are frequently detected amongst *C. albicans* clinical isolates (Selmecki *et al.* 2006; Ford *et al.* 2015; Anderson *et al.* 2017) and can play an important role in drug resistance (Selmecki *et al.* 2006, 2008; Li *et al.* 2015; Todd and Selmecki 2020), host adaptation (Forche *et al.* 2018, 2019), and other stress tolerance phenotypes (Yang *et al.* 2019b, 2021). For example, amplification of part or all of chromosome 5, via trisomy or isochromosome formation has been tightly linked to the development of resistance to azole antifungals in *C. albicans* via the increased copy number of genes encoding the azole target, Erg11, and the transcriptional regulator of drug efflux, Tac1 (Selmecki *et al.* 2008). The prevalence and importance of these phenomena associated with increased gene expression highlight the relevance of developing new tools to targetedly assess phenotypes associated with genetic overexpression in *C. albicans*.

Harnessing the cell's natural transcriptional machinery to increase gene expression with CRISPRa also leads to a unique set of obstacles. Off-target effects as a result of promiscuous dCas9 binding in the cell presents the possibility of complications if the dCas9 is also targetting other regions in the genome. While our RNA-seq results suggest that our CRISPRa system is specific when paired with one sgRNA targeted to the *ALS1* gene, there is still the possibility of off-target effects occurring in different CRISPRa constructs, especially as binding specificity with CRISPR-Cas9 systems is highly dependent on the exact sgRNA sequence in use (Zhang *et al.* 2015). Using complementary genetic tools to confirm phenotypes observed from CRISPRa strains can help mitigate this. One phenomenon shared among many different CRISPRa activation domains, including VPR, is a reduced capacity to induce overexpression in genes that are already expressed at a high levels (Chavez *et al.* 2016). Incidentally, the gene that we achieved the lowest overexpression with across all of our testing, *FGR28*, had the lowest basal

expression (data not shown) from our RT-qPCR experiments, opposing the idea that genes that are normally expressed at low levels tend to be more easily overexpressed with CRISPRa systems. However, efficiency of the sgRNA molecule and its exact position in the target region of the gene of interest, as well as a host of promoter dynamics, can also influence CRISPR-Cas9 efficacy (Radzisheuskaya *et al.* 2016; Cámara *et al.* 2020; Lu and Lin 2021). This illustrates why multiple sgRNAs need to be designed when targeting a gene with CRISPRa to maximize successful overexpression of the target gene. While the rapid and inexpensive nature of CRISPRa strain construction easily justifies the need to design several sgRNA molecules for any given gene, broader limitations as a result of the specific "NGG" PAM recognition site by traditional *Streptococcus pyogenes* Cas9 proteins, like the one used here, leads to the restraint that on rare occasions it may not be possible to produce any highly-efficient sgRNA molecules in a suitable region of the promoter (Uthayakumar *et al.* 2021). Since this PAM restriction is a feature shared among many CRISPR-Cas9 systems, it may therefore be prudent to further expand the CRISPR toolbox in *C. albicans* by introducing novel Cas9 variants with differing PAM recognition sites into our system, or by constructing novel CRISPR systems exploiting alternative Cas proteins with different and broader PAM recognition capacities, offering the ability to target regions in the genome that cannot be manipulated with current CRISPR-Cas9 tools in *C. albicans* (Hu *et al.* 2018; Ciurkot *et al.* 2021).

While large-scale CRISPRa libraries have not been generated in *C. albicans*, their application in other organisms, namely mammalian cell lines, has enabled important discoveries in cancer biology, stem cell reprogramming, cellular fitness, and drug resistance (Gilbert *et al.* 2014; Konermann *et al.* 2015; Horlbeck *et al.* 2016; Joung *et al.* 2017; Jost *et al.* 2017; Kampmann 2018; Sanson *et al.* 2018; Yang *et al.* 2019a). The development of CRISPRa systems in *C. albicans* could enable the development of larger overexpression libraries for functional genomic screens, to complement existing fungal gene deletion and depletion libraries (Roemer *et al.* 2003; Homann *et al.* 2009; Blankenship *et al.* 2010; Noble *et al.* 2010) that have

been subject to robust screening and analysis under diverse conditions (Gervais *et al.* 2021), to reveal critical new insight into fungal biology (Lohse *et al.* 2016; Hossain *et al.* 2020), pathogenesis (Noble *et al.* 2010; Nobile *et al.* 2012; Ryan *et al.* 2012; O'Meara *et al.* 2015; Lee *et al.* 2016; Hossain *et al.* 2020), and mechanisms of drug and stress tolerance (Motaung *et al.* 2017; Mount *et al.* 2018; Caplan *et al.* 2018; Fu *et al.* 2021).

Currently, non-CRISPR-based overexpression libraries have been constructed and screened in *C. albicans*, revealing novel insights into gene function related to hyphal morphogenesis (Chauvel *et al.* 2012), cell cycle progression (Jaitly *et al.* 2022), drug tolerance (Delarze *et al.* 2020), and biofilm formation (Cabral *et al.* 2014). Previous comparisons done between CRISPRa and ORF-based overexpression libraries such as these in other species have highlighted the potential for these two complementary strategies to both confirm the results from one another as well as unveil significant genetic hits distinct from each other, even when libraries share similar gene targets (Sanson *et al.* 2018). Compared with other overexpression methods, the ability of the CRISPRa system to target any specified region in the genome also confers the unique and powerful ability of allele-specific targeting which could potentially be used to study heterozygosity in *C. albicans*, an aspect of its genomic makeup that is critical to its pathogenicity (Liang and Bennett 2019). CRISPRa also offers the advantage of being easily scalable into a pooled format whereby hundreds or thousands of unique plasmids and resulting mutant strains can be constructed simultaneously, and mutant libraries in different strain backgrounds can therefore be quickly constructed (Sanjana 2017; Momen-Roknabadi *et al.* 2020). Non-CRISPR overexpression libraries that have previously been assembled in *C. albicans* have the potential weakness of missing important phenotypes during screening if the level of overexpression achieved in each mutant strain is too low (Znaidi *et al.* 2018). The ability of our system to cause upwards of 25-fold overexpression in gene targets is thus an important feature. Furthermore, corresponding phenotypes in mutant strains may only be observable if the level of overexpression closely mimics the level of upregulation of the gene induced by the cell

in other relevant stress conditions naturally (Znaidi *et al.* 2018). Future iterations of this CRISPRa system could also exploit an inducible dCas9-VPR construct for tunability of gene expression. It is therefore imperative to employ tools such as CRISPRa that possess the advantage of being inherently flexible and easily tunable in order to study genetic overexpression thoroughly.

The application of similar CRISPRa systems to other fungal organisms would lend further insight into fungal biology across this kingdom. Many CRISPR tools have been developed in a diversity of fungal organisms (Muñoz *et al.* 2019; Cai *et al.* 2019; Román *et al.* 2019b; Schuster and Kahmann 2019; Morio *et al.* 2020; Ouedraogo and Tsang 2020; Shan *et al.* 2021), including numerous non-*albicans Candida* species (Santana and O'Meara; Enkler *et al.* 2016; Lombardi *et al.* 2017, 2019; Zoppo *et al.* 2019; Maroc and Fairhead 2019; Uthayakumar *et al.* 2021; Ennis *et al.* 2021), and other prominent pathogens such as *Cryptococcus neoformans* (Fan and Lin 2018; Wang 2018; Huang *et al.* 2021) and *Aspergillus fumigatus* (Fuller *et al.* 2015; Zhang *et al.* 2016; Al Abdallah *et al.* 2018; van Rhijn *et al.* 2020). These CRISPR platforms could be adapted to similar CRISPRa tools in these organisms, with many possible applications for the analysis of pathogen biology or drug target identification (Li *et al.* 2004; Luesch *et al.* 2005; Smith *et al.* 2010; Arnoldo *et al.* 2014; Begolo *et al.* 2014).

Genetic overexpression systems also have many useful applications for industrially-important fungi, as targeted genetic overexpression can be exploited for metabolic engineering to enhance the production of commercially-important metabolites (Zhao *et al.* 2018; Donohoue *et al.* 2018; Ouedraogo and Tsang 2020; Jiang *et al.* 2021). CRISPR techniques have already been adapted to numerous industrially-relevant fungi (Nødvig *et al.* 2015; Kwon *et al.* 2019; Jiang *et al.* 2021; Wilson and Harrison 2021), and CRISPRa has recently been optimized for use in the filamentous fungi *Aspergillus nidulans* and *Penicillium rubens* (Roux *et al.* 2020; Mózsik *et al.* 2021). These CRISPRa systems were applied to activate transcriptionally-silent biosynthetic gene clusters in these filamentous fungi, with important

applications for the discovery of novel bioactive molecules (Roux *et al.* 2020; Mózsik *et al.* 2021). As many *Candida* species play important industrial roles in manufacturing metabolites for food and pharmaceutical industries, as well as remediating wastewater via hydrocarbon degradation (Joo *et al.* 2008; Gargouri *et al.* 2015; Kieliszek *et al.* 2017; Payen and Thompson 2019; García-Béjar *et al.* 2020), species-specific adaptations to the CRISPRa tool presented here could produce optimized *Candida* strains for valuable biomanufacturing and bioremediation applications.

Materials and Methods

Plasmid design and cloning

The plasmid backbone used in this study was the *C. albicans*-optimized CRISPR-dCas9 plasmid (pRS143, Addgene #122377) used in our previous study (Wensing *et al.* 2019), containing the *NEUT5L* integration site, sgRNA cloning site (*SNR52* promoter, SapI cloning locus, and sgRNA tail), and *dCAS9*. The dCas9-VPR fusion construct was generated via Gibson assembly. The VPR tripartite complex was codon-optimized for *C. albicans* expression and synthesized as gBlocks gene fragments from Integrated DNA Technologies (IDT). This gene fragment was cloned with Gibson assembly into the *dCAS9* plasmid backbone. We have made the CRISPRa (dCas9-VPR) plasmid available via Addgene (reference #182707). All plasmids are listed in **TABLE S1.**

sgRNA design and cloning

The sgRNA CRISPR RNAs (crRNA; 20 nucleotide sequence complementary to the target DNA genomic DNA) were designed based on efficiency and predicted specificity via the sgRNA design tool Eukaryotic Pathogen CRISPR gRNA Design Tool (EuPaGDT; http://grna.ctegd.uga.edu) (Peng and Tarleton 2015). These sgRNA crRNA N20 sequences were cloned into the dCas9-VPR CRISPRa plasmid at the sgRNA cloning locus (SapI cloning

site, flanked by *SNR52* promoter, and sgRNA tail) using Golden Gate cloning (Engler *et al.* 2008), as previously described (Halder *et al.* 2019; Wensing and Shapiro 2022). crRNA N20 sequences were obtained as two oligonucleotides from IDT in forward and reverse complement orientation, each containing a SapI cloning site, and were reconstituted to 100µM in a nuclease-free duplex buffer from IDT. Equal volumes of the two complementary oligonucleotides were combined after being heated separately at 94°C, and then duplexed by heating to 94°C for 2 min and cooling to room temperature. The duplexed fragment was then cloned into the CRISPRa plasmid with the following Golden Gate cloning reaction: 10µl miniprepped CRISPRa plasmid, 1µl duplexed oligonucleotide, 2µl 10X CutSmart buffer, 2µl ATP, 1µl SapI, 1µl T4 DNA ligase, and 3µl nuclease-free water. This mixture was incubated in a thermocycler under the following cycling conditions: (37°C, 2 min; 16°C, 5 min) for 99 cycles; 65°C, 15 min; 80°C, 15 min. After cycling, 1µl of additional SapI was added to each reaction mixture, and the reaction was incubated at 37°C for 1h. All sgRNAs are listed in **TABLE S1.**

Media and growth conditions

Escherichia coli DH5α cells were grown at 37°C in Lysogeny Broth (LB) and LB plates supplemented with 100mg/mL ampicillin (AMP) and 250mg/mL nourseothricin (NAT) for plasmid selection. *C. albicans* cells were grown at 30°C or 37°C in Yeast Peptone Dextrose (YPD) broth and YPD plates supplemented with 250mg/mL nourseothricin (NAT) for plasmid selection.

Bacterial transformation

High Efficiency 5-alpha Competent *E.coli* (NEB) cells were thawed on ice. 1µl of Golden Gate cloning reagents were added to 50µl of competent cells and incubated on ice for 30 min, heat shocked for 30s at 42°C, and incubated on ice for an additional 5 min. 950µl of SOC outgrowth media was added to each cell culture and incubated at 30°C for 1.5h at 250 RPM. Transformed cells were then plated on LB media containing AMP and NAT and grown at 30°C for 1 day.

C. albicans transformation

Plasmids were transformed into *C. albicans* via a chemical transformation strategy, as previously described (Halder *et al.* 2019; Wensing and Shapiro 2022). Briefly, miniprepped CRISPRa plasmids were linearized via a 1.5X restriction digest mix using the PacI enzyme. A transformation master mix was prepared as follows: 800μl of 50% polyethylene glycol (PEG), 100μl of 10X Tris-EDTA (TE) buffer solution, 100μl of 1M lithium acetate (LiAc), 40μl of 10 mg/mL salmon sperm DNA, and 20μl of 1M dithiothreitol (DTT). The transformation mix was added to *C. albicans* cells and linearized CRISPRa plasmid and left to incubate at 30°C for 1h, then heat-shocked at 42°C for 45 min. Cells were washed with fresh YPD, then grown in YPD for 4h to allow for expression of the NAT resistance construct. Transformed cells were then plated on YPD media containing NAT, and grown at 30°C for 2 days. All *C. albicans* strains are listed in **TABLE S1.**

Growth curves

Overnight cultures of *C. albicans* grown in YPD were diluted to an OD600 of 0.05 in fresh YPD and added to a 96-well flat-bottomed plate with a total volume of 200μl per well. Growth was measured at 30°C via optical density at 600nm at 15 min intervals over the course of 24h using an Infinite 200 PRO microplate reader (Tecan), and plates were shaken orbitally for 900s at a 4mm amplitude in between growth measurements.

Minimum inhibitory concentration (MIC) assays

MIC assays were performed in 96-well flat-bottomed plates. A suspension of 40μg/mL fluconazole was prepared in water, of which 100uL was added to the first column of each plate containing 100uL of culture, to obtain a total volume of 200μl per well. The first column was serially diluted 2-fold across the plate in water. The gradient of fluconazole, therefore, ranged

from 20µg/mL to 0µg/mL. Overnight cultures of *C. albicans* grown in YPD were diluted to an OD600 of 0.1 in 2X RPMI 1640 with 40g/L of added D-glucose and mixed into the plates in an equal volume such that the starting OD600 values of each strain were 0.05 and the growth media was diluted to RPMI 1640 with 20g/L of added D-glucose. Plates were incubated at 37°C at 900 RPM, and absorbance values at 600nm were read after 24h using an Infinite 200 PRO microplate reader (Tecan).

Biofilm assays

RPMI-based biofilm assays were performed as previously described (Razzaq *et al.* 2021), with minor modifications. Overnight cultures of *C. albicans* grown in YPD were diluted to an OD600 of 0.001 in 5mL of RPMI 1640 with 20g/L of supplemented D-glucose. 100µl of RPMI 1640 was mixed with 100µl of each diluted overnight culture into a 96-well flat-bottomed plate (12 wells per strain). Plates were grown statically at 37°C for 24h to allow biofilms to form. 120µl of planktonic cell supernatant was removed from each well and moved to a new 96-well flat-bottomed plate. Density of the planktonic cells was measured at 600nm using an Infinite 200 PRO microplate reader (Tecan). Original plates from which planktonic cells were removed were washed by adding 200µl 1X PBS to each well with an electronic multichannel pipette at a moderate dispense speed. 1X PBS was then discarded, and this wash step was repeated. Plates were then left to dry for 1 - 2h. Once biofilms were dry, 90µl of 1mg/mL XTT (prepared in 1X PBS and centrifuged to remove sediment prior to use) and 10µl of 0.32mg/mL PMS (prepared in water) were added to each well. Plates were incubated statically at 30°C for 2h to allow biofilms to reduce XTT, measured at 490nm using an Infinite 200 PRO microplate reader (Tecan), and normalized to the growth of planktonic cells harvested previously.

For urine-based biofilm assays, biofilm formations were performed in 96-well flat-bottomed plates. Plates were coated with 150µg/mL of fibrinogen and incubated overnight at 4°C. The *C. albicans* strains were cultured at 37°C with aeration in 5mL of YPD broth. The

inoculum was normalized to ~1x10^6 CFU/ml and then diluted (1:10) in human urine (female, pH

~6.5). Urine was supplemented with 20mg/mL of BSA for carbon and nitrogen sources

mimicking the catheterized bladder environment. 200µl of the urine with inoculum was incubated

in each well of the 96-well plate at 37°C for 48h while static. Following the 48h incubation, the

cultures were removed and the wells were washed three times with 200µl 1X PBS to remove

unbound fungi. Plates were incubated with 200µl of 0.5% crystal violet for 15 min. Crystal violet

was removed, and plates were washed with water and dried. 200µl of 33% acetic acid was

added to the wells and 100µl was transferred to a new 96-well plate. Absorbance was measured

at 595nm using a SpectraMax ABS Plus microplate reader (Molecular Devices).

RNA extraction and real-time quantitative PCR

To detect differences in gene expression, overnight cultures of *C. albicans* grown in YPD

were diluted to an OD600 of 0.05 in fresh YPD and grown to an OD600 of >0.2 at 30°C.

Cultures were pelleted and frozen at -80°C before RNA was extracted. RNA extractions were

performed using either a Presto Mini RNA Yeast Kit from FroggaBio (cat. RBYD050) or an

RNeasy Mini Kit from Qiagen (cat. 74104). To assess RNA integrity, an RNA ScreenTape assay

was performed on all samples using the TapeStation 4150 system following manufacturer's

instructions (Agilent Technologies). Only samples with RIN values of >5.0 were used for

RT-qPCR. Synthesis of cDNA from 1000ng (10µl) RNA was performed using a High Capacity

cDNA Reverse Transcription Kit from Applied BioSystems (cat. 4368814). Briefly, 20µl reaction

mixtures were prepared with 2µl 10X reverse transcription buffer, 0.8µl of 25X dNTPs at a

concentration of 100mM, 2µl of random primers, 1µl of MultiScribe Reverse Transcriptase at 50

U/µl, 4.2µl of nuclease-free water, and 10µl of RNA (1µg). Reverse transcription run conditions

were as follows: 10 min at 25°C, 120 min at 37°C, 5 min at 85°C. Real-time PCR assays were

conducted using a QuantStudio 7 Pro Real Time PCR system from Thermo Fisher Scientific Inc.

20µl reaction mixtures containing 10µl 2X SsoAdvanced Universal Inhibitor-Tolerant SYBR

supermix (Bio-Rad, cat: 172-5017), 0.8µl of PCR forward and reverse primer mix at 5µM (final concentration of primer at 200 nM), 4.2µl of water and 5µl of diluted cDNA. The run conditions were as follows: 3 min at 98°C polymerase activation step, followed by 40 cycles of a two-step qPCR (10s of 98°C denaturation, 30s of 60°C combined annealing/extension). Primers are listed in **TABLE S1**. Expression profiling calculations were performed according to the comparative C_T method (Schmittgen and Livak 2008). Briefly, expression values for the experimental gene of interest were compared to the housekeeping gene *ACT1* to obtain a ΔC_T value within each strain. The ΔC_T values in the experimental CRISPRa strains were then compared to the non-targeting CRISPRa control strain to obtain a $\Delta\Delta C_T$ value and finally a fold-difference in expression of the experimental gene.

RNA sequencing

RNA preparation and sequencing were performed as previously described, with minor modifications (Razzaq *et al.* 2021). Strains were grown in YPD at 30°C overnight preceding RNA extraction. RNA extraction, sample QC, library preparations, and sequencing reactions were conducted at GENEWIZ, LLC./Azenta US, Inc. as follows: Total RNA was extracted using Qiagen RNeasy Plus Universal kit following manufacturer's instructions (Qiagen). RNA samples were quantified using Qubit 2.0 Fluorometer (ThermoFisher Scientific) and RNA integrity was checked with 4200 TapeStation (Agilent Technologies). ERCC RNA Spike-In Mix kit (cat. 4456740) from ThermoFisher Scientific, was added to normalized total RNA prior to library preparation following the manufacturer's protocol. The RNA sequencing library was prepared using the NEBNext Ultra II RNA Library Prep Kit for Illumina using the manufacturer's instructions (New England Biolabs). Briefly, mRNAs were initially enriched with Oligod(T) beads. Enriched mRNAs were fragmented for 15 min at 94 °C. First-strand and second-strand cDNA were subsequently synthesized. cDNA fragments were end-repaired and adenylated at 3'ends, and universal adapters were ligated to cDNA fragments, followed by index addition and library

enrichment by PCR with limited cycles. The sequencing library was validated on the Agilent

TapeStation (Agilent Technologies), and quantified by using Qubit 2.0 Fluorometer

(ThermoFisher Scientific) as well as by quantitative PCR (KAPA Biosystems). The sequencing

libraries were multiplexed and clustered onto a flowcell. After clustering, the flowcell was loaded

onto the Illumina HiSeq 4000 or equivalent instrument according to the manufacturer's

instructions. The samples were sequenced using a 2x150bp Paired End (PE) configuration.

Image analysis and base calling were conducted by the Illumina Control Software.

Raw sequence data (.bcl files) generated from the Illumina instrument was converted

into fastq files and de-multiplexed using Illumina bcl2fastq 2.20 software. One mis-match was

allowed for index sequence identification. Sequence reads were trimmed to remove possible

adapter sequences and nucleotides with poor quality using Trimmomatic v.0.36, and trimmed

reads were mapped to the candida_albicans reference genome available on ENSEMBL using

the STAR aligner v.2.5.2b. The STAR aligner is a splice aligner that detects splice junctions and

incorporates them to help align the entire read sequences. BAM files were generated as a result

of this step. Unique gene hit counts were then calculated by using featureCounts from the

Subread package v.1.5.2. The hit counts were summarized and reported using the gene_id

feature in the annotation file. Only unique reads that fell within exon regions were counted. After

extraction of gene hit counts, the gene hit counts table was used for downstream differential

expression analysis. Using DESeq2, a comparison of gene expression between the control and

experimental groups of samples was performed. The Wald test was then used to generate

p-values and log2 fold changes. Genes with an adjusted p-value < 0.05 and absolute log2 fold

change > 1 were called as differentially expressed genes for each comparison. The data have

been deposited in NCBI's Gene Expression Omnibus (GEO) (Edgar *et al.* 2002) and are

accessible through GEO Series accession number GSE201904

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE201904).

Data availability

The *C. albicans* CRISPRa plasmid backbone was deposited to Addgene as plasmid number #182707. RNA sequencing data was deposited in NCBI GEO repository (accession number GSE201904; https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE201904).

Acknowledgments

We thank Shapiro lab members, past and present, for helpful discussions and support of this project, as well as Dr. Anna Selmecki and Dr. Petra Vande Zande (University of Minnesota) for helpful feedback. We thank Dr. Zhenguo Lin (Saint Louis University) for helpful discussions on TSS prediction, and Kieran Shah for programming support on this project. We thank Jing Zhang from the University of Guelph Advanced Analysis Centre for RT-qPCR support. This work was supported by an NSERC Discovery Grant (RGPIN-2018-4914), an Ontario Early Research Award, and a CIFAR Azrieli Global Scholar award to RSS. N.C.G and L.W are EvoFunPath Fellows (NSERC CREATE) and L.W. is supported by an Ontario Graduate Scholarship. A.C. is supported by a Career Award for Medical Scientist from the Burroughs Wellcome Fund. Work by A.L.F.M. and F.H.S.T. was supported by institutional funds from the University of Notre Dame as well as National Institute of Health grant R01-DK128805 (to A.L.F.M. and A.A.L.), and the Arthur J. Schmitt Presidential Leadership Foundation (to A.L.L).

Figure Legends

Figure 1. Design of a CRISPRa system for *C. albicans*. (a) The single plasmid system contains a catalytically inactive ("dead") dCas9 protein, paired with the VPR (VP64-P65-Rta) complex of activator domains to cause upregulation at a target locus. The single-guide RNA (sgRNA) region can be subjected to Golden Gate cloning with a custom N20 sequence in order to target the dCas9-VPR complex to a desired region of DNA through complementary base-pairing. The Pac1 enzyme is used to linearize the plasmid at the *NEUT5L* locus, which allows the plasmid to be integrated into the cell's chromosome and remain in the cell following cell division. Panel created with BioRender.com. **(b)** Once the sgRNA recognizes and binds to a target sequence of DNA, the dCas9-VPR complex will cause recruitment of transcription initiation complex factors to the adjacent genetic material, thus increasing gene expression from that region. Created with BioRender.com. **(c)** Wild-type *Candida albicans* cells and cells containing a non-targeting CRISPRa plasmid were grown in YPD at 30°C to compare fitness under standard laboratory conditions. Growth was recorded via optical density at 600nm at 15 min intervals over the course of 24h. Data points show the mean and standard deviation at each time point (n = 32).

Figure 2. Determining optimal guide-design principles for CRISPRa overexpression in *C. albicans*. CRISPRa strains were created using six different sgRNA molecules designed to target different positions in the promoter region of each of **(a)** *KSP1*, **(b)** *ALS3*, and **(c)** *FGR28* upstream of either the start codon or the predicted dominant transcriptional start site (TSS). Fold-change in the expression of the target gene relative to the housekeeping gene *ACT1* in the experimental strains and the non-targeting CRISPRa control strain was measured with RT-qPCR and compared. Data points represent the mean fold-difference and standard deviation in expression of the respective target gene in the CRISPRa strains compared to the non-targeting CRISPRa control (n = 3).

Figure 3. Using CRISPRa to validate antifungal drug resistance phenotypes. (a) Three *C. albicans* CRISPRa strains were created targeting the drug transporter gene *CDR1* for overexpression. Fold-change in the expression of *CDR1* relative to the housekeeping gene *ACT1* in the three experimental strains and the non-targeting CRISPRa control strain was measured with RT-qPCR. All three experimental strains are represented by the position of their sgRNA in the promoter region upstream of the proposed dominant *CDR1* transcriptional start site. represent the mean fold-difference and standard deviation in expression of *CDR1* in the CRISPRa strains compared to the non-targeting CRISPRa control (n = 3). **(b)** The fitness of all three CRISPRa strains targeting *CDR1* for overexpression was measured in the presence of the antifungal drug fluconazole with a minimum inhibitory concentration assay. The OD600 value of each strain in each concentration of fluconazole was measured to represent growth in differing conditions, where OD600 values approaching 1.0 reflect strong fitness.

Figure 4. Validation of CRISPRa overexpression and target-specificity through targeting fungal pathogenesis traits. (a) Two *C. albicans* CRISPRa strains targeting different positions in the promoter region of the adhesin protein gene *ALS1* were created. Fold-change in the expression of *ALS1* relative to the housekeeping gene *ACT1* in both experimental strains and the non-targeting CRISPRa control strain was measured with RT-qPCR. Experimental strains are represented by the position of their sgRNA in the promoter region upstream of the proposed dominant *ALS1* transcriptional start site. Data points represent the mean fold-difference and standard deviation in expression of *ALS1* in the CRISPRa strains compared to the non-targeting CRISPRa control (n = 3). **(b)** Two CRISPRa strains targeting *ALS1* and a non-targeting CRISPRa control strain were grown in the biofilm-inducing media RPMI and left to grow statically at 37°C for 24h to allow biofilms to form. The robustness of biofilms formed by all strains was elucidated through an XTT reduction assay and measuring of the resulting OD490

values before normalizing final values to non-biofilm-forming planktonic cells. Data points represent the mean and the standard error of the mean (n = 72). Differences between groups were tested for significance using Dunnett's multiple comparison test. ****, p-value <0.0001. ***, p-value <0.001. **(c)** A *C. albicans* CRISPRa strain overexpressing *ALS1* ~6-fold and a non-targeting CRISPRa strain were grown in human urine supplemented with fibrinogen and BSA at 37°C for 48h to simulate a catheterized bladder environment. The robustness of biofilms formed by both strains was measured by reading the OD595 value following a crystal violet assay. Data points are combined from three independent experiments where n = 32 for the non-targeting CRISPRa control and n = 48 for the CRISPRa *ALS1* overexpression strain. Differences between groups were tested for significance using the Mann-Whitney U test. ****, p-value <0.0001. **(d)** Global gene expression of a CRISPRa strain overexpressing *ALS1* ~25-fold, as well as a non-targeting CRISPRa control strain, was measured using RNA-Seq. Differential gene expression analysis showed that *ALS1* had a log2-fold change of ~3.3 in the *ALS1* overexpression strain compared to the CRISPRa control strain, with an adjusted p-value of 3.07E-142, and showed no additional transcripts that had differential expression with an adjusted p-value of less than 0.03.

References

Adli, M., 2018 The CRISPR tool kit for genome editing and beyond. Nat. Commun. 9: 1911.

Al Abdallah, Q., A. C. O. Souza, A. Martin-Vicente, W. Ge, and J. R. Fortwendel, 2018 Whole-genome sequencing reveals highly specific gene targeting by *in vitro* assembled Cas9-ribonucleoprotein complexes in *Aspergillus fumigatus*. Fungal Biol Biotechnol 5: 11.

Alberti-Segui, C., A. J. Morales, H. Xing, M. M. Kessler, D. A. Willins *et al.*, 2004 Identification of potential cell-surface proteins in *Candida albicans* and investigation of the role of a putative cell-surface glycosidase in adhesion and virulence. Yeast 21: 285–302.

Andersen, M. J., C. Fong, A. A. La Bella, J. J. Molina, A. Molesan *et al.*, 2022 Inhibiting host-protein deposition on urinary catheters reduces associated urinary tract infections. eLife 11:e75798

Anderson, M. Z., A. Saha, A. Haseeb, and R. J. Bennett, 2017 A chromosome 4 trisomy contributes to increased fluconazole resistance in a clinical isolate of *Candida albicans*. Microbiology 163: 856–865.

Arendrup, M. C., and T. F. Patterson, 2017 Multidrug-resistant *Candida*: epidemiology, Molecular mechanisms, and treatment. J. Infect. Dis. 216: S445–S451.

Arnoldo, A., S. Kittanakom, L. E. Heisler, A. B. Mak, A. I. Shukalyuk *et al.*, 2014 A genome scale overexpression screen to reveal drug activity in human cells. Genome Med. 6: 32.

Begolo, D., E. Erben, and C. Clayton, 2014 Drug target identification using a trypanosome overexpression library. Antimicrob. Agents Chemother. 58: 6260–6264.

Benedict, K., B. R. Jackson, T. Chiller, and K. D. Beer, 2018 Estimation of direct healthcare costs of fungal diseases in the United States. Clin. Infect. Dis.

Blankenship, J. R., S. Fanning, J. J. Hamaker, and A. P. Mitchell, 2010 An Extensive circuitry for cell wall regulation in *Candida albicans*. PLoS Pathogens 6: e1000752.

Bock, C., P. Datlinger, F. Chardon, M. A. Coelho, M. B. Dong *et al.*, 2022 High-content CRISPR

screening. Nature Reviews Methods Primers 2: 1–1.

Bongomin, F., S. Gago, R. Oladele, and D. Denning, 2017 Global and multi-national prevalence of fungal diseases—estimate precision. Journal of Fungi 3: 57.

Cabral, V., S. Znaidi, L. A. Walker, H. Martin-Yken, E. Dague *et al.*, 2014 Targeted changes of the cell wall proteome influence *Candida albicans* ability to form single- and multi-strain biofilms. PLoS Pathog. 10: e1004542.

Cai, P., J. Gao, and Y. Zhou, 2019 CRISPR-mediated genome editing in non-conventional yeasts for biotechnological applications. Microb. Cell Fact. 18: 63.

Cámara, E., I. Lenitz, and Y. Nygård, 2020 A CRISPR activation and interference toolkit for industrial *Saccharomyces cerevisiae* strain KE6-12. Sci. Rep. 10: 14605.

Caplan, T., E. J. Polvi, J. L. Xie, S. Buckhalter, M. D. Leach *et al.*, 2018 Functional genomic screening reveals core modulators of echinocandin stress responses in *Candida albicans*. Cell Rep. 23: 2292–2298.

Chauvel, M., A. Nesseir, V. Cabral, S. Znaidi, S. Goyard *et al.*, 2012 A versatile overexpression strategy in the pathogenic yeast *Candida albicans*: identification of regulators of morphogenesis and fitness. PLoS One 7: e45912.

Chavez, A., J. Scheiman, S. Vora, B. W. Pruitt, M. Tuttle *et al.*, 2015 Highly efficient Cas9-mediated transcriptional programming. Nat. Methods 12: 326–328.

Chavez, A., M. Tuttle, B. W. Pruitt, B. Ewen-Campen, R. Chari *et al.*, 2016 Comparison of Cas9 activators in multiple species. Nature Methods 13: 563–567.

Ciurkot, K., T. E. Gorochowski, J. A. Roubos, and R. Verwaal, 2021 Efficient multiplexed gene regulation in *Saccharomyces cerevisiae* using dCas12a. Nucleic Acids Res. 49: 7775–7790.

Coste, A. T., M. Karababa, F. Ischer, J. Bille, and D. Sanglard, 2004 *TAC1*, transcriptional activator of CDR genes, is a new transcription factor involved in the regulation of *Candida albicans* ABC transporters *CDR1* and *CDR2*. Eukaryot. Cell 3: 1639–1652.

Coste, A., V. Turner, F. Ischer, J. Morschhäuser, A. Forche *et al.*, 2006 A mutation in Tac1p, a transcription factor regulating *CDR1* and *CDR2*, is coupled with loss of heterozygosity at chromosome 5 to mediate antifungal resistance in *Candida albicans*. Genetics 172: 2139–2156.

Delarze, E., L. Brandt, E. Trachsel, M. Patxot, C. Pralong *et al.*, 2020 Identification and characterization of mediators of fluconazole tolerance in *Candida albicans*. Front. Microbiol. 11: 591140.

Delgado, M. L., M. Luisa Delgado, M. Luisa Gil, and D. Gozalbo, 2003 *Candida albicans TDH3* gene promotes secretion of internal invertase when expressed in *Saccharomyces cerevisiae* as a glyceraldehyde-3-phosphate dehydrogenase-invertase fusion protein. Yeast 20: 713–722.

Dominguez, A. A., W. A. Lim, and L. S. Qi, 2016 Beyond editing: repurposing CRISPR-Cas9 for precision genome regulation and interrogation. Nat. Rev. Mol. Cell Biol. 17: 5–15.

Donohoue, P. D., R. Barrangou, and A. P. May, 2018 Advances in industrial biotechnology using CRISPR-Cas systems. Trends Biotechnol. 36: 134–146.

Eckert, S. E., and F. A. Mühlschlegel, 2009 Promoter regulation in *Candida albicans* and related species. FEMS Yeast Res. 9: 2–15.

Ene, I. V., R. A. Farrer, M. P. Hirakawa, K. Agwamba, C. A. Cuomo *et al.*, 2018 Global analysis of mutations driving microevolution of a heterozygous diploid fungal pathogen. Proc. Natl. Acad. Sci. U. S. A. 115: E8688–E8697.

Engler, C., R. Kandzia, and S. Marillonnet, 2008 A one pot, one step, precision cloning method with high throughput capability. PLoS One 3: e3647.

Enkler, L., D. Richer, A. L. Marchand, D. Ferrandon, and F. Jossinet, 2016 Genome engineering in the yeast pathogen *Candida glabrata* using the CRISPR-Cas9 system. Sci. Rep. 6: 35766.

Ennis, C. L., A. D. Hernday, and C. J. Nobile, 2021 A markerless CRISPR-mediated system for

genome editing in *Candida auris* reveals a conserved role for Cas5 in the caspofungin response. Microbiol Spectr 9: e0182021.

Fan, Y., and X. Lin, 2018 Multiple applications of a rransient CRISPR-Cas9 coupled with electroporation (TRACE) system in the *Cryptococcus neoformans* species complex. Genetics 208: 1357–1372.

Fisher, M. C., S. J. Gurr, C. A. Cuomo, D. S. Blehert, H. Jin *et al.*, 2020 Threats posed by the fungal kingdom to humans, wildlife, and agriculture. MBio 11.:

Flores-Mireles, A. L., J. N. Walker, T. M. Bauman, A. M. Potretzke, H. L. Schreiber 4th *et al.*, 2016 Fibrinogen release and deposition on urinary catheters placed during urological procedures. J. Urol. 196: 416–421.

Flowers, S. A., K. S. Barker, E. L. Berkow, G. Toner, S. G. Chadwick *et al.*, 2012 Gain-of-function mutations in *UPC2* are a frequent cause of *ERG11* upregulation in azole-resistant clinical isolates of *Candida albicans*. Eukaryot. Cell 11: 1289–1299.

Forche, A., G. Cromie, A. C. Gerstein, N. V. Solis, T. Pisithkul *et al.*, 2018 Rapid phenotypic and genotypic diversification after exposure to the oral host niche in *Candida albicans*. Genetics 209: 725–741.

Forche, A., P. T. Magee, A. Selmecki, J. Berman, and G. May, 2009 Evolution in *Candida albicans* populations during a single passage through a mouse host. Genetics 182: 799–811.

Forche, A., N. V. Solis, M. Swidergall, R. Thomas, A. Guyer *et al.*, 2019 Selection of *Candida albicans* trisomy during oropharyngeal infection results in a commensal-like phenotype. PLoS Genet. 15: e1008137.

Ford, C. B., J. M. Funt, D. Abbey, L. Issi, C. Guiducci *et al.*, 2015 The evolution of drug resistance in clinical isolates of *Candida albicans*. Elife 4: e00662.

Fu, Y., A. S. Ibrahim, D. C. Sheppard, Y.-C. Chen, S. W. French *et al.*, 2002 *Candida albicans* Als1p: an adhesin that is a downstream effector of the *EFG1* filamentation pathway. Mol.

Microbiol. 44: 61–72.

Fuller, K. K., S. Chen, J. J. Loros, and J. C. Dunlap, 2015 Development of the CRISPR/Cas9
system for targeted gene disruption in *Aspergillus fumigatus*. Eukaryot. Cell 14: 1073–1080.

Fu, Y., G. Luo, B. J. Spellberg, J. E. Edwards Jr, and A. S. Ibrahim, 2008 Gene
overexpression/suppression analysis of candidate virulence factors of *Candida albicans*.
Eukaryot. Cell 7: 483–492.

Fu, C., X. Zhang, A. O. Veri, K. R. Iyer, E. Lash *et al.*, 2021 Leveraging machine learning
essentiality predictions and chemogenomic interactions to identify antifungal targets. Nat.
Commun. 12: 6497.

García-Béjar, B., M. Arévalo-Villena, E. Guisantes-Batan, J. Rodríguez-Flores, and A. Briones,
2020 Study of the bioremediatory capacity of wild yeasts. Sci. Rep. 10: 11265.

Gargouri, B., N. Mhiri, F. Karray, F. Aloui, and S. Sayadi, 2015 Isolation and characterization of
hydrocarbon-degrading yeast strains from petroleum contaminated industrial wastewater.
Biomed Res. Int. 2015: 929424.

Geddes-McAlister, J., and R. S. Shapiro, 2018 New pathogens, new tricks: emerging,
drug-resistant fungal pathogens and future prospects for antifungal therapeutics. Ann. N. Y.
Acad. Sci.

Gerami-Nejad, M., L. F. Zacchi, M. McClellan, K. Matter, and J. Berman, 2013 Shuttle vectors
for facile gap repair cloning and integration into a neutral locus in *Candida albicans*.
Microbiology 159: 565–579.

Gervais, N. C., V. Halder, and R. S. Shapiro, 2021 A data library of *Candida albicans* functional
genomic screens. FEMS Yeast Res. 21.:

Gilbert, L. A., M. A. Horlbeck, B. Adamson, J. E. Villalta, Y. Chen *et al.*, 2014 Genome-scale
CRISPR-mediated control of gene repression and activation. Cell 159: 647–661.

Gow, N. A. R., F. L. van de Veerdonk, A. J. P. Brown, and M. G. Netea, 2011 *Candida albicans*
morphogenesis and host defence: discriminating invasion from colonization. Nat. Rev.

Microbiol. 10: 112–122.

Halder, V., C. B. M. Porter, A. Chavez, and R. S. Shapiro, 2019 Design, execution, and analysis
of CRISPR–Cas9-based deletions and genetic interaction networks in the fungal pathogen
Candida albicans. Nat. Protoc. 14: 955–975.

Hardwick, J. M., L. Tse, N. Applegren, J. Nicholas, and M. A. Veliuona, 1992 The Epstein-Barr
virus R transactivator (Rta) contains a complex, potent activation domain with properties
different from those of VP16. J. Virol. 66: 5500–5508.

Homann, O. R., J. Dea, S. M. Noble, and A. D. Johnson, 2009 A phenotypic profile of the
Candida albicans regulatory network. PLoS Genet. 5: e1000783.

Horlbeck, M. A., L. A. Gilbert, J. E. Villalta, B. Adamson, R. A. Pak *et al.*, 2016 Compact and
highly active next-generation libraries for CRISPR-mediated gene repression and
activation. Elife 5:e19760

Hossain, S., A. O. Veri, and L. E. Cowen, 2020 The proteasome governs fungal morphogenesis
via functional connections with Hsp90 and cAMP-protein kinase A signaling. MBio 11.:

Hoyer, L. L., and E. Cota, 2016 *Candida albicans* agglutinin-like Sequence (Als) family
vignettes: a review of Als protein structure and function. Front. Microbiol. 7: 280.

Hoyer, L. L., C. B. Green, S.-H. Oh, and X. Zhao, 2008 Discovering the secrets of the *Candida
albicans* agglutinin-like sequence (ALS) gene family--a sticky pursuit. Med. Mycol. 46: 1–15.

Huang, M. Y., M. B. Joshi, M. J. Boucher, S. Lee, L. C. Loza *et al.*, 2021 Short
homology-directed repair using optimized Cas9 in the pathogen *Cryptococcus neoformans*
enables rapid gene deletion and tagging. Genetics. 22: iyab180

Huang, M. Y., and A. P. Mitchell, 2017 Marker recycling in *Candida albicans* through
CRISPR-Cas9-induced marker excision. mSphere 2: e00050

Hu, J. H., S. M. Miller, M. H. Geurts, W. Tang, L. Chen *et al.*, 2018 Evolved Cas9 variants with
broad PAM compatibility and high DNA specificity. Nature 556: 57–63.

Jaitly, P., M. Legrand, A. Das, T. Patel, M. Chauvel *et al.*, 2022 A phylogenetically-restricted

essential cell cycle progression factor in the human pathogen *Candida albicans*. Nat. Commun. 13: 4256.

Jensen, M. K., 2018 Design principles for nuclease-deficient CRISPR-based transcriptional regulators. FEMS Yeast Res. 18.:

Jiang, C., G. Lv, Y. Tu, X. Cheng, Y. Duan *et al.*, 2021 Applications of CRISPR/Cas9 in the synthesis of secondary metabolites in filamentous fungi. Front. Microbiol. 12: 638096.

Joo, H.-S., P. M. Ndegwa, M. Shoda, and C.-G. Phae, 2008 Bioremediation of oil-contaminated soil using *Candida catenulata* and food waste. Environmental Pollution 156: 891–896.

Jost, M., Y. Chen, L. A. Gilbert, M. A. Horlbeck, L. Krenning *et al.*, 2017 Combined CRISPRi/a-based chemical genetic screens reveal that Rigosertib is a microtubule-destabilizing Agent. Mol. Cell 68: 210–223.e6.

Joung, J., J. M. Engreitz, S. Konermann, O. O. Abudayyeh, V. K. Verdine *et al.*, 2017 Genome-scale activation screen identifies a lncRNA locus regulating a gene neighbourhood. Nature 548: 343–346.

Kamai, Y., M. Kubota, Y. Kamai, T. Hosokawa, T. Fukuoka *et al.*, 2002 Contribution of *Candida albicans ALS1* to the pathogenesis of experimental oropharyngeal candidiasis. Infect. Immun. 70: 5256–5258.

Kampmann, M., 2018 CRISPRi and CRISPRa screens in mammalian cells for precision biology and medicine. ACS Chem. Biol. 13: 406–416.

Kieliszek, M., A. M. Kot, A. Bzducha-Wróbel, S. BŁażejak, I. Gientka *et al.*, 2017 Biotechnological use of *Candida* yeasts in the food industry: A review. Fungal Biology Reviews 31: 185–198.

Konermann, S., M. D. Brigham, A. E. Trevino, J. Joung, O. O. Abudayyeh *et al.*, 2015 Genome-scale transcriptional activation by an engineered CRISPR-Cas9 complex. Nature 517: 583–588.

Kwon, M. J., T. Schütze, S. Spohner, S. Haefner, and V. Meyer, 2019 Practical guidance for the

implementation of the CRISPR genome editing tool in filamentous fungi. Fungal Biol
Biotechnol 6: 15.

La Bella, A. A., M. J. Andersen, N. C. Gervais, J. J. Molina, A. Molesan *et al.*, 2021 Catheterized
bladder environment induces *Candida albicans* hyphal formation and promotes colonization
and persistence through Als1. bioRxiv.

La Russa, M. F., and L. S. Qi, 2015 The New State of the Art: Cas9 for Gene Activation and
Repression. Mol. Cell. Biol. 35: 3800–3809.

Lee, J. A., N. Robbins, J. L. Xie, T. Ketela, and L. E. Cowen, 2016 Functional genomic analysis
of *Candida albicans* adherence reveals a key role for the Arp2/3 complex in cell wall
remodelling and biofilm formation. PLoS Genet. 12: e1006452.

Legrand, M., S. Bachellier-Bassi, K. K. Lee, Y. Chaudhari, H. Tournu *et al.*, 2018 Generating
genomic platforms to study *Candida albicans* pathogenesis. Nucleic Acids Res. 46:
6935–6949.

Liang, S.-H., and R. J. Bennett, 2019 The impact of gene dosage and heterozygosity on the
diploid pathobiont *Candida albicans*. J Fungi 6:10

Li, X., F. Yang, D. Li, M. Zhou, X. Wang *et al.*, 2015 Trisomy of chromosome R confers
resistance to triazoles in *Candida albicans*. Med. Mycol. 53: 302–309.

Li, X., M. Zolli-Juran, J. D. Cechetto, D. M. Daigle, G. D. Wright *et al.*, 2004 Multicopy
suppressors for novel antibacterial compounds reveal targets and drug efflux susceptibility.
Chem. Biol. 11: 1423–1430.

Lohberger, A., A. T. Coste, and D. Sanglard, 2014 Distinct roles of *Candida albicans* drug
resistance transcription factors *TAC1*, *MRR1*, and *UPC2* in virulence. Eukaryot. Cell 13:
127–142.

Lohse, M. B., I. V. Ene, V. B. Craik, A. D. Hernday, E. Mancera *et al.*, 2016 Systematic genetic
screen for transcriptional regulators of the *Candida albicans* white-opaque switch. Genetics
203: 1679–1692.

Lombardi, L., J. Oliveira-Pacheco, and G. Butler, 2019 Plasmid-based CRISPR-Cas9 gene
    editing in multiple *Candida* species. mSphere 4.:

Lombardi, L., S. A. Turner, F. Zhao, and G. Butler, 2017 Gene editing in clinical isolates of
    Candida parapsilosis using CRISPR/Cas9. Sci. Rep. 7: 8051.

Luesch, H., T. Y. H. Wu, P. Ren, N. S. Gray, P. G. Schultz *et al.*, 2005 A genome-wide
    overexpression screen in yeast for small-molecule target identification. Chem. Biol. 12:
    55–63.

Lu, Z., and Z. Lin, 2021 The origin and evolution of a distinct mechanism of transcription
    initiation in yeasts. Genome Res. 31: 51–63.

Manoharlal, R., N. A. Gaur, S. L. Panwar, J. Morschhäuser, and R. Prasad, 2008 Transcriptional
    activation and increased mRNA stability contribute to overexpression of *CDR1* in
    azole-resistant *Candida albicans*. Antimicrob. Agents Chemother. 52: 1481–1492.

Maroc, L., and C. Fairhead, 2019 A new inducible CRISPR-Cas9 system useful for genome
    editing and study of double-strand break repair in *Candida glabrata*. Yeast 36: 723–731.

Marr, K. A., C. N. Lyons, K. Ha, T. R. Rustad, and T. C. White, 2001 Inducible azole resistance
    associated with a heterogeneous phenotype in *Candida albicans*. Antimicrobial Agents and
    Chemotherapy 45: 52–59.

Marr, K. A., C. N. Lyons, T. R. Rustad, R. A. Bowden, and T. C. White, 1998 Rapid, transient
    fluconazole resistance in *Candida albicans* is associated with increased mRNA levels of
    CDR. Antimicrob. Agents Chemother. 42: 2584–2589.

McMillan, J., Z. Lu, J. S. Rodriguez, T.-H. Ahn, and Z. Lin, 2019 YeasTSS: an integrative web
    database of yeast transcription start sites. Database 2019.:

Min, K., A. Biermann, D. A. Hogan, and J. B. Konopka, 2018 Genetic analysis of *NDT80* family
    transcription factors in *Candida albicans* using new CRISPR-Cas9 approaches. mSphere 3:
    e00545

Min, K., Y. Ichikawa, C. A. Woolford, and A. P. Mitchell, 2016 *Candida albicans* gene deletion

with a transient CRISPR-Cas9 system. mSphere 1: e00130-16.

Momen-Roknabadi, A., P. Oikonomou, M. Zegans, and S. Tavazoie, 2020 An inducible CRISPR
    interference library for genetic interrogation of *Saccharomyces cerevisiae* biology. Commun
    Biol 3: 723.

Morio, F., L. Lombardi, and G. Butler, 2020 The CRISPR toolbox in medical mycology: State of
    the art and perspectives. PLoS Pathog. 16: e1008201.

Motaung, T. E., R. Ells, C. H. Pohl, J. Albertyn, and T. J. Tsilo, 2017 Genome-wide functional
    analysis in *Candida albicans*. Virulence 8: 1563–1579.

Mount, H. O., N. M. Revie, R. T. Todd, K. Anstett, C. Collins *et al.*, 2018 Global analysis of
    genetic circuitry and adaptive mechanisms enabling resistance to the azole antifungal
    drugs. PLoS Genet. 14: e1007319.

Mózsik, L., M. Hoekzema, N. A. W. de Kok, R. A. L. Bovenberg, Y. Nygård *et al.*, 2021
    CRISPR-based transcriptional activation tool for silent genes in filamentous fungi. Sci. Rep.
    11: 1118.

Muñoz, I. V., S. Sarrocco, L. Malfatti, R. Baroncelli, and G. Vannacci, 2019 CRISPR-Cas for
    fungal genome editing: a new tool for the management of plant diseases. Front. Plant Sci.
    10: 135.

Ng, H., and N. Dean, 2017 Dramatic improvement of CRISPR/Cas9 editing in *Candida albicans*
    by increased single guide RNA expression. mSphere 2: e00385

Nguyen, N., M. M. F. Quail, and A. D. Hernday, 2017 An efficient, rapid, and recyclable system
    for CRISPR-mediated genome editing in *Candida albicans*. mSphere 2: e00149

Niimi, M., K. Niimi, Y. Takano, A. R. Holmes, F. J. Fischer *et al.*, 2004 Regulated overexpression
    of *CDR1* in *Candida albicans* confers multidrug resistance. J. Antimicrob. Chemother. 54:
    999–1006.

Nobile, C. J., E. P. Fox, J. E. Nett, T. R. Sorrells, Q. M. Mitrovich *et al.*, 2012 A recently evolved
    transcriptional network controls biofilm development in *Candida albicans*. Cell 148:

126–138.

Nobile, C. J., H. A. Schneider, J. E. Nett, D. C. Sheppard, S. G. Filler *et al.*, 2008
Complementary adhesin function in *C. albicans* biofilm formation. Curr. Biol. 18:
1017–1024.

Noble, S. M., S. French, L. A. Kohn, V. Chen, and A. D. Johnson, 2010 Systematic screens of a
Candida albicans homozygous deletion library decouple morphogenetic switching and
pathogenicity. Nat. Genet. 42: 590–598.

Nødvig, C. S., J. B. Nielsen, M. E. Kogle, and U. H. Mortensen, 2015 A CRISPR-Cas9 system
for genetic engineering of filamentous fungi. PLoS One 10: e0133085.

O'Meara, T. R., A. O. Veri, T. Ketela, B. Jiang, T. Roemer *et al.*, 2015 Global analysis of fungal
morphology exposes mechanisms of host cell escape. Nat. Commun. 6: 6741.

Ouedraogo, J.-P., and A. Tsang, 2020 CRISPR_Cas systems for fungal research. Fungal Biol.
Rev. 34: 189–201.

Park, Y.-N., and J. Morschhäuser, 2005 Tetracycline-inducible gene expression and gene
deletion in *Candida albicans*. Eukaryot. Cell 4: 1328–1342.

Payen, C., and D. Thompson, 2019 The renaissance of yeasts as microbial factories in the
modern age of biomanufacturing. Yeast 36: 685–700.

Peng, D., and R. Tarleton, 2015 EuPaGDT: a web tool tailored to design CRISPR guide RNAs
for eukaryotic pathogens. Microb Genom 1: e000033.

Pérez, J. C., 2019 *Candida albicans* dwelling in the mammalian gut. Curr. Opin. Microbiol. 52:
41–46.

Perfect, J. R., 2017 The antifungal pipeline: a reality check. Nat. Rev. Drug Discov.

Peters, J. M., M. R. Silvis, D. Zhao, J. S. Hawkins, C. A. Gross *et al.*, 2015 Bacterial CRISPR:
accomplishments and prospects. Curr. Opin. Microbiol. 27: 121–126.

Polvi, E. J., A. O. Veri, Z. Liu, S. Hossain, S. Hyde *et al.*, 2019 Functional divergence of a global
regulatory complex governing fungal filamentation. PLOS Genetics 15: e1007901.

Prasad, R., A. Banerjee, N. K. Khandelwal, and S. Dhamgaye, 2015 The ABCs of *Candida albicans* multidrug transporter Cdr1. Eukaryot. Cell 14: 1154–1164.

Prasad, R., P. De Wergifosse, A. Goffeau, and E. Balzi, 1995 Molecular cloning and characterization of a novel gene of *Candida albicans, CDR1*, conferring multiple resistance to drugs and antifungals. Current Genetics 27: 320–329.

Prieto, D., E. Román, R. Alonso-Monge, and J. Pla, 2017 Overexpression of the transcriptional regulator *WOR1* increases susceptibility to bile salts and adhesion to the mouse gut mucosa in *Candida albicans*. Front. Cell. Infect. Microbiol. 7: 389.

Radzisheuskaya, A., D. Shlyueva, I. Müller, and K. Helin, 2016 Optimizing sgRNA position markedly improves the efficiency of CRISPR/dCas9-mediated transcriptional repression. Nucleic Acids Res. 44: e141.

Rai, L. S., L. van Wijlick, M. Chauvel, C. d'Enfert, M. Legrand *et al.*, 2021 Overexpression approaches to advance understanding of *Candida albicans*. Mol. Microbiol.

Razzaq, I., M. D. Berg, Y. Jiang, J. Genereaux, D. Uthayakumar *et al.*, 2021 The SAGA and NuA4 component Tra1 regulates *Candida albicans* drug resistance and pathogenesis. Genetics 219.:

van Rhijn, N., T. Furukawa, C. Zhao, B. L. McCann, E. Bignell *et al.*, 2020 Development of a marker-free mutagenesis system using CRISPR-Cas9 in the pathogenic mould *Aspergillus fumigatus*. Fungal Genet. Biol. 145: 103479.

Roemer, T., B. Jiang, J. Davison, T. Ketela, K. Veillette *et al.*, 2003 Large-scale essential gene identification in *Candida albicans* and applications to antifungal drug discovery. Mol. Microbiol. 50: 167–181.

Román, E., I. Coman, D. Prieto, R. Alonso-Monge, and J. Pla, 2019a Implementation of a CRISPR-based system for gene regulation in *Candida albicans*. mSphere 4.:

Román, E., D. Prieto, R. Alonso-Monge, and J. Pla, 2019b New insights of CRISPR technology in human pathogenic fungi. Future Microbiol. 14: 1243–1255.

Rosiana, S., L. Zhang, G. H. Kim, A. V. Revtovich, D. Uthayakumar *et al.*, 2021 Comprehensive genetic analysis of adhesin proteins and their role in virulence of *Candida albicans*. Genetics.

Roux, I., C. Woodcraft, J. Hu, R. Wolters, C. L. M. Gilchrist *et al.*, 2020 CRISPR-mediated activation of biosynthetic gene clusters for bioactive molecule discovery in filamentous fungi. ACS Synth. Biol. 9: 1843–1854.

Ryan, O., R. S. Shapiro, C. F. Kurat, D. Mayhew, A. Baryshnikova *et al.*, 2012 Global gene deletion analysis exploring yeast filamentous growth. Science 337: 1353–1356.

Sahni, N., S. Yi, K. J. Daniels, G. Huang, T. Srikantha *et al.*, 2010 Tec1 mediates the pheromone response of the white phenotype of *Candida albicans*: insights into the evolution of new signal transduction pathways. PLoS Biology 8: e1000363.

Sanglard, D., A. Coste, and S. Ferrari, 2009 Antifungal drug resistance mechanisms in fungal pathogens from the perspective of transcriptional gene regulation. FEMS Yeast Res. 9: 1029–1050.

Sanglard, D., K. Kuchler, F. Ischer, J. L. Pagani, M. Monod *et al.*, 1995 Mechanisms of resistance to azole antifungal agents in *Candida albicans* isolates from AIDS patients involve specific multidrug transporters. Antimicrobial Agents and Chemotherapy 39: 2378–2386.

Sanguinetti, M., B. Posteraro, and C. Lass-Flörl, 2015 Antifungal drug resistance among *Candida* species: mechanisms and clinical impact. Mycoses 58 Suppl 2: 2–13.

Sanjana, N. E., 2017 Genome-scale CRISPR pooled screens. Anal. Biochem. 532: 95–99.

Sanson, K. R., R. E. Hanna, M. Hegde, K. F. Donovan, C. Strand *et al.*, 2018 Optimized libraries for CRISPR-Cas9 genetic screens with multiple modalities. Nat. Commun. 9: 5416.

Santana, D. J., and T. R. O'Meara Forward and reverse genetic dissection of morphogenesis identifies filament-competent *Candida auris* strains.

Schmittgen, T. D., and K. J. Livak, 2008 Analyzing real-time PCR data by the comparative C(T)

method. Nat. Protoc. 3: 1101–1108.

Schmitz, M. L., and P. A. Baeuerle, 1991 The p65 subunit is responsible for the strong
transcription activating potential of NF-kappa B. The EMBO Journal 10: 3805–3817.

Schubert, S., K. S. Barker, S. Znaidi, S. Schneider, F. Dierolf *et al.*, 2011 Regulation of efflux
pump expression and drug resistance by the transcription factors Mrr1, Upc2, and Cap1 in
Candida albicans. Antimicrob. Agents Chemother. 55: 2212–2223.

Schultenkämper, K., L. F. Brito, and V. F. Wendisch, 2020 Impact of CRISPR interference on
strain development in biotechnology. Biotechnol. Appl. Biochem. 67: 7–21.

Schuster, M., and R. Kahmann, 2019 CRISPR-Cas9 genome editing approaches in filamentous
fungi and oomycetes. Fungal Genet. Biol. 130: 43–53.

Segrelles-Calvo, G., G. R. de S Araújo, E. Llopis-Pastor, J. Carrillo, M. Hernández-Hernández *et
al.*, 2021 *Candida* spp. co-infection in COVID-19 patients with severe pneumonia:
Prevalence study and associated risk factors. Respir. Med. 188: 106619.

Selmecki, A., A. Forche, and J. Berman, 2006 Aneuploidy and isochromosome formation in
drug-resistant *Candida albicans*. Science 313: 367–370.

Selmecki, A., M. Gerami-Nejad, C. Paulson, A. Forche, and J. Berman, 2008 An
isochromosome confers drug resistance in vivo by amplification of two genes, *ERG11* and
TAC1. Mol. Microbiol. 68: 624–641.

Shan, L., Z. Dai, and Q. Wang, 2021 Advances and opportunities of CRISPR/Cas technology in
bioengineering non-conventional yeasts. Front Bioeng Biotechnol 9: 765396.

Shapiro, R. S., A. Chavez, and J. J. Collins, 2018a CRISPR-based genomic tools for the
manipulation of genetically intractable microorganisms. Nat. Rev. Microbiol. 16: 333–339.

Shapiro, R. S., A. Chavez, C. B. M. Porter, M. Hamblin, C. S. Kaas *et al.*, 2018b A
CRISPR-Cas9-based gene drive platform for genetic interaction analysis in *Candida
albicans*. Nat Microbiol 3: 73–82.

Smith, A. M., R. Ammar, C. Nislow, and G. Giaever, 2010 A survey of yeast genomic assays for

drug and target discovery. Pharmacol. Ther. 127: 156–164.

Smith, J. D., S. Suresh, U. Schlecht, M. Wu, O. Wagih *et al.*, 2016 Quantitative CRISPR
interference screens in yeast identify chemical-genetic interactions and new rules for guide
RNA design. Genome Biol. 17: 45.

Sopko, R., D. Huang, N. Preston, G. Chua, B. Papp *et al.*, 2006 Mapping pathways and
phenotypes by systematic gene overexpression. Mol. Cell 21: 319–330.

Todd, R. T., and A. Selmecki, 2020 Expandable and reversible copy number amplification drives
rapid adaptation to antifungal drugs. Elife 9: e58349

Todd, R. T., T. D. Wikoff, A. Forche, and A. Selmecki, 2019 Genome plasticity in *Candida
albicans* is driven by long repeat sequences. Elife 8: e45954

Todor, H., M. R. Silvis, H. Osadnik, and C. A. Gross, 2021 Bacterial CRISPR screens for gene
function. Curr. Opin. Microbiol. 59: 102–109.

Uthayakumar, D., J. Sharma, L. Wensing, and R. S. Shapiro, 2021 CRISPR-based genetic
manipulation of *Candida* species: historical perspectives and current approaches. Frontiers
in Genome Editing 2: 606281

Vyas, V. K., M. I. Barrasa, and G. R. Fink, 2015 A Candida albicans CRISPR system permits
genetic engineering of essential genes and gene families. Sci Adv 1: e1500248.

Vyas, V. K., G. Guy Bushkin, D. A. Bernstein, M. A. Getz, M. Sewastianik *et al.*, 2018 New
CRISPR mutagenesis strategies reveal variation in repair mechanisms among fungi.
mSphere 3: e00154–18.

Wang, P., 2018 Two distinct approaches for CRISPR-Cas9-mediated gene editing in
Cryptococcus neoformans and related species. mSphere 3: e00208

Wang, H., M. La Russa, and L. S. Qi, 2016 CRISPR/Cas9 in genome editing and beyond. Annu.
Rev. Biochem. 85: 227–264.

Wensing, L., and R. S. Shapiro, 2022 Design and generation of a CRISPR interference system
for genetic repression and essential gene analysis in the fungal pathogen *Candida albicans*.

Methods Mol. Biol. 2377: 69–88.

Wensing, L., J. Sharma, D. Uthayakumar, Y. Proteau, A. Chavez *et al.*, 2019 A CRISPR
interference platform for efficient genetic repression in *Candida albicans*. mSphere 4.:

White, T. C., 1997 Increased mRNA levels of *ERG16*, *CDR*, and *MDR1* correlate with increases
in azole resistance in *Candida albicans* isolates from a patient infected with human
immunodeficiency virus. Antimicrob. Agents Chemother. 41: 1482–1487.

Wilson, F. M., and R. J. Harrison, 2021 CRISPR/Cas9 mediated editing of the Quorn fungus
Fusarium venenatum A3/5 by transient expression of Cas9 and sgRNAs targeting
endogenous marker gene *PKS12*. Fungal Biol Biotechnol 8: 15.

Xu, X., and L. S. Qi, 2019 A CRISPR–dCas Toolbox for genetic engineering and synthetic
Biology. J. Mol. Biol. 431: 34–47.

Yang, J., S. S. Rajan, M. J. Friedrich, G. Lan, X. Zou *et al.*, 2019a Genome-scale CRISPRa
screen identifies novel factors for cellular reprogramming. Stem Cell Reports 12: 757–771.

Yang, F., F. Teoh, A. S. M. Tan, Y. Cao, N. Pavelka *et al.*, 2019b Aneuploidy enables
cross-adaptation to unrelated drugs. Mol. Biol. Evol.

Yang, F., R. T. Todd, A. Selmecki, Y.-Y. Jiang, Y.-B. Cao *et al.*, 2021 The fitness costs and
benefits of trisomy of each *Candida albicans* chromosome. Genetics 218.:

Zhang, C., X. Meng, X. Wei, and L. Lu, 2016 Highly efficient CRISPR mutagenesis by
microhomology-mediated end joining in *Aspergillus fumigatus*. Fungal Genet. Biol. 86:
47–57.

Zhang, X.-H., L. Y. Tee, X.-G. Wang, Q.-S. Huang, and S.-H. Yang, 2015 Off-target effects in
CRISPR/Cas9-mediated genome engineering. Mol. Ther. Nucleic Acids 4: e264.

Zhao, X., S.-H. Oh, G. Cheng, C. B. Green, J. A. Nuessen *et al.*, 2004 *ALS3* and *ALS8*
represent a single locus that encodes a *Candida albicans* adhesin; functional comparisons
between Als3p and Als1p. Microbiology 150: 2415–2428.

Zhao, X.-Q., X.-Y. Zhang, F. Zhang, R. Zhang, B.-J. Jiang *et al.*, 2018 Metabolic engineering of

fungal strains for efficient production of cellulolytic enzymes, pp. 27–41 in *Fungal Cellulolytic Enzymes: Microbial Production and Application*, edited by X. Fang and Y. Qu. Springer Singapore, Singapore.

Znaidi, S., L. van Wijlick, A. Hernández-Cervantes, N. Sertour, J.-L. Desseyn *et al.*, 2018 Systematic gene overexpression in *Candida albicans* identifies a regulator of early adaptation to the mammalian gut. Cell. Microbiol. 20: e12890.

Zoppo, M., M. D. Luca, S. N. Villarreal, N. Poma, M. I. Barrasa *et al.*, 2019 A CRISPR/Cas9-based strategy to simultaneously inactivate the entire ALS gene family in *Candida orthopsilosis*. Future Microbiol. 14: 1383–1396.

a

WT

CDR1 CRISPRa strains
- 90bp
- 200bp
- 290bp

Fold Change (*CDR1*)

0   2   4   6   8

b

OD600

0   1.0

[Fluconazole]   10μg/mL

ND

WT

CDR1 CRISPRa strains
- 90bp
- 200bp
- 290bp