1 **Development and applications of a CRISPR activation system for facile genetic**

2 **overexpression in *Candida albicans***

Nicholas C. Gervais[1], Alyssa A. La Bella[2], Lauren F. Wensing[1], Jehoshua Sharma[1], Victoria

Acquaviva[1], Madison Best[1], Ricardo Omar Cadena López[1], Meea Fogal[1], Deeva

Uthayakumar[1†], Alejandro Chavez[3], Felipe Santiago-Tirado[2], Ana L. Flores-Mireles[2], and

Rebecca S. Shapiro[1]*

[1]Department of Molecular and Cellular Biology, University of Guelph, Guelph, Ontario, Canada

[2]Department of Biological Sciences, University of Notre Dame, Notre Dame, Indiana, USA

[3]Department of Pathology and Cell Biology, Columbia University College of Physicians and

Surgeons, New York, New York, USA

†Current affiliation: Department of Immunology, University of Toronto, Toronto, Ontario, Canada

*Correspondence: shapiror@uoguelph.ca

Abstract

For the fungal pathogen *Candida albicans*, genetic overexpression readily occurs via a diversity

of genomic alterations, such as aneuploidy and gain-of-function mutations, with important

consequences for host adaptation, virulence, and evolution of antifungal drug resistance. Given

the important role of overexpression on *C. albicans* biology, it is critical to develop and harness

tools that enable the analysis of genes expressed at high levels in the fungal cell. Here, we

describe the development, optimization, and application of a novel, single-plasmid-based

CRISPR activation (CRISPRa) platform for targeted genetic overexpression in *C. albicans*,

which employs a guide RNA to target an activator complex to the promoter region of a gene of

interest, thus driving transcriptional expression of that gene. Using this system, we demonstrate

the ability of CRISPRa to drive high levels of gene expression in *C. albicans*, and we assess

optimal guide RNA targeting for robust and constitutive overexpression. We further demonstrate

the specificity of the system via RNA sequencing. We highlight the application of CRISPRa to

overexpress genes involved in pathogenesis and drug resistance and contribute towards the

identification of novel phenotypes. Together, this tool will facilitate a broad range of applications

for the study of *C. albicans* genetic overexpression.

Keywords

Fungal genetics, CRISPR activation, Candida albicans, Gene regulation, Fungal pathogens

Introduction

Fungal pathogens are an increasingly important cause of human illness, with significant impacts on morbidity and mortality (1). *Candida albicans* is an opportunistic fungal pathogen that inhabits the microbiome of most healthy adults as a commensal organism (2, 3). However, it is also responsible for invasive infections, particularly amongst immunocompromised individuals, which are often fatal (4). Instances of *C. albicans* infections are becoming more prevalent globally, in part due to an increasing population of susceptible elderly and immunocompromised individuals (4). Illustratively, systemic *Candida* infections have been identified in up to 15% of patients hospitalized for COVID-19, and patients with invasive candidiasis spent longer in the hospital Intensive Care Unit (ICU) on average, and had more severe symptoms than those COVID-19 patients without a *Candida* infection (5). The economic cost of *Candida* infections is estimated to be ~1.4 billion dollars yearly in the United States alone, hospitalizing more than 26,000 people (6). With limited development of novel antifungal drugs (7) and increasing prevalence of antifungal drug-resistant strains and species (8–10), *Candida* pathogens remain a serious threat to human health.

Given the importance of *C. albicans* and other fungal pathogens, it is imperative that tools are developed and applied to manipulate these organisms and dissect genetic functions. For *C. albicans*, along with countless other microbial species, CRISPR-based genetic manipulation tools have revolutionized the ability of scientists to modify and study gene function (11, 12). Such CRISPR platforms typically rely on a CRISPR-associated endonuclease protein, often Cas9, targeted to a genomic region of interest via a single guide RNA (sgRNA), resulting in targeted DNA double-strand breaks (DSBs), and subsequent repair via non-homologous end-joining or homology-directed repair via a donor DNA repair template (13, 14). Since the initial development of CRISPR tools for genetic editing in *C. albicans* (15), numerous systems have been developed to modify this fungal genome with efficiency and specificity (11, 16–24), and to

build and screen genetic mutant libraries (24, 25). Together, these gene editing systems have

played an important role in providing novel biological insight into this fungal pathogen.

In addition to these canonical CRISPR-based genetic modification systems that

introduce mutations, insertions, or deletions into the genome, alternative CRISPR technologies

have been developed that modulate the expression of *C. albicans* genes (26, 27). These

CRISPR systems rely on the inactivation of the Cas9 endonuclease, to generate a nuclease-

dead (dCas9) protein, which can be fused to constructs from a transcriptional repressor or

activator that generate CRISPR interference (CRISPRi) or CRISPR activation (CRISPRa)

systems, respectively (28, 29). These dCas9 fusions, when targeted to the promoter region of a

gene via the sgRNA, can repress or activate gene transcription from a gene of interest, and

have been widely applied in diverse organisms (14, 30–34). In *C. albicans*, these modified

CRISPR systems have enabled the modulation and study of important fungal genes, including

essential genes (26, 27, 35).

While a majority of CRISPR systems and other genetic manipulation platforms focus on

gene mutation, deletion, or downregulation, it is also critical to develop techniques that enable

the upregulation or overexpression of genes. Indeed, genetic overexpression in *C. albicans*

occurs readily via gene copy number amplification and aneuploidy, and these copy number

amplification events are frequently observed in genetically diverse clinical isolates (36–38). The

upregulation of gene expression plays a critical role in diverse aspects of fungal biology (39),

including the evolution of antifungal drug resistance (40–45), host colonization (46, 47), and

virulence (41, 48). Additionally, in other microbial systems, including bacteria, protozoa, and the

model yeast *Saccharomyces cerevisiae*, genetic overexpression libraries have proven to be

critical tools for identifying targets of drugs with uncharacterized mechanisms of action, as

mutants overexpressing the drug target are typically more resistant to the drug in question (49–

52). Overexpression screens can also help dissect molecular pathways (53–55). In *C. albicans*,

several techniques exist for genetic overexpression (39), including promoter replacement

strategies (55–58), the generation of ORFeome collection strains (59, 60), and CRISPR-based

methods (27). While these these tools have made important contributions to understanding

overexpression in *C. albicans*, there remains a need for simple and rapid methods to efficiently

target fungal genes of interest for overexpression, and to do so in diverse strain backgrounds.

Here, we introduce a CRISPR-based tool for genetic overexpression in *C. albicans*, to

bolster the existing functional genetic toolbox available to manipulate genes in this critical fungal

pathogen. Our CRISPRa tool exploits a *C. albicans*-optimized tripartite activator complex fused

to dCas9, which was previously developed for highly-efficient transcriptional regulation in *S.*

cerevisiae (61). This CRISPRa system is distinct from other CRISPR-based gene activation

systems for *C. albicans*, as it is an efficient single-plasmid system designed for rapid Golden

Gate cloning and facile fungal strain generation, which can be readily applied to diverse strain

backgrounds including clinical isolates. We test CRISPRa guide targeting principles based on

predicted transcriptional start sites, confirm on-target guide efficiency, and demonstrate the

ability of this CRISPRa system to drive high levels of expression of *C. albicans* genes. Using

this optimized CRISPRa technique, we validate its ability to drive overexpression of genes with

established roles in antifungal drug resistance and biofilm formation. Together, this work

introduces a novel CRISPRa system for genetic overexpression in *C. albicans*, with a wide

range of future applications.

Results

Development of a single-plasmid CRISPRa tool in C. albicans

First, we designed a single-plasmid CRISPRa system for targeted gene activation,

optimized for *C. albicans*. We exploited an integrating plasmid backbone, which we previously

used for CRISPRi in *C. albicans*, containing a codon-optimized dCas9 and Golden Gate

assembly-compatible sgRNA cloning site (26). This plasmid can be linearized to integrate at the

C. albicans NEUT5L locus (a large intergenic region whose disruption does not affect fungal

fitness (62)), and contains a dual SapI cloning site for facile and highly-efficient Golden Gate

cloning of the sgRNA targeting sequence. This plasmid also contains a dominant nourseothricin

resistance (NATr) cassette, to enable strain generation in a diversity of fungal strains, including

clinical isolates.

To generate a version of dCas9 that could lead to strong constitutive activation of genes

of interest, we generated a plasmid with dCas9 fused to the tripartite activator complex, VP64-

p65-Rta (VPR) (**FIGURE 1a,b**). VPR consists of three previously-described transcriptional

activators: (1) VP64, a viral transcriptional activator domain composed of four tandem copies of

VP16 (Herpes Simplex Viral Protein 16), which has been used extensively in CRISPRa systems

(63); (2) p65, the NF-κB subunit responsible for the strong transcription activation in mammalian

cells (64); and (3) Rta, the Epstein-Barr virus R transactivator, with strong transcriptional

activation properties (65) (**FIGURE 1a,b**). Together the VPR activator complex fused to dCas9

and targetted to the promoter region of genes has previously been demonstrated to generate

strong gene activation in *S. cerevisiae* (61). Exploiting the ability of dCas9 to bind and be

targeted to genomic loci via sgRNAs enables us to design sgRNAs that will target dCas9 along

with these strong transcriptional activators to any gene promoter of interest.

This plasmid was generated by homology-based cloning and validated via Sanger

sequencing, and has been deposited to Addgene (**TABLE S1**, Addgene plasmid #182707). To

confirm if the expression of the dCas9-VPR construct had an impact on fungal fitness, we

compared the growth of a wild-type *C. albicans* strain with a strain transformed with the dCas9-

VPR CRISPRa plasmid (expressing a non-targeting sgRNA: a random sequence of nucleotides

that have no predicted complementarity to any region in the *C. albicans* genome). We found no

difference in growth between these strains (**FIGURE 1c**), suggesting no significant impact of the

CRISPRa system on *C. albicans* fitness.

Validation and sgRNA design principles for CRISPRa overexpression in C. albicans

Next, we aimed to validate that the CRISPRa system could successfully overexpress

genes in *C. albicans*, and sought to determine the optimal sgRNA design strategy to maximize

genetic regulation. Our previous *C. albicans* CRISPRi system exploited sgRNAs ~ 50 - 150bp

upstream of the start codon to functionally repress genes of interest (26). However, CRISPRa

systems are known to have different optimal sgRNA targeting regions (66, 67), and our previous

design did not take into account the transcription start site (TSS), which is known to be an

important predictor of regulatory efficiency for CRISPRi/a systems (66, 68), but which had not

been robustly characterized across all *C. albicans* genes at the time. More recently, the TSS for

many *C. albicans* genes has been predicted (69). Thus, we sought to determine the optimal

design for CRISPRa sgRNAs based on the start codon and the TSS. We identified three *C.*

albicans genes with a predicted TSS that was particularly far upstream of the start codon

compared to all other genes in the genome (801-952 bp upstream): *KSP1*, *ALS3,* and *FGR28* .

We designed six sgRNAs for each gene, targeting upstream of the start codon (as previously

designed for CRISPRi (26)), as well as upstream of the TSS (based on predicted TSS (69)).

We generated CRISPRa plasmids based on these sgRNAs, used these to create *C.*

albicans strains, and monitored overexpression of each gene via quantitative reverse-

transcription PCR (RT-qPCR). This analysis validated our novel CRISPRa system and its ability

to overexpress *C. albicans* genes ~ 2 - 12-fold, based on different sgRNA designs and target

genes (**FIGURE 2**). We found that in some instances, such as for *KSP1*, the TSS was in fact a

better predictor of CRISPRa overexpression, as the most robust overexpression of this gene

was observed with sgRNAs targetted upstream of the TSS, while no overexpression was

observed based on targeting upstream of the start codon (**FIGURE 2a**). In other instances, such

as for *ALS3*, sgRNA targeting upstream of the start codon led to very high levels of

overexpression, while sgRNAs targeted to the TSS region led to moderate overexpression

(**FIGURE 2b**). For *FGR28*, modest overexpression (~2-fold) was achieved with sgRNAs

targeted to either the start codon or the TSS upstream regions (**FIGURE 2c**), together

suggesting that optimal sgRNA design can be predicted based on targeting the TSS and/or start

codon, but varies by gene. This data also suggests that our CRISPRa tool may have additional

applications in the functional assessment of key transcriptional regulatory regions in *C. albicans*.

These differences in sgRNA targeting may be based on differences in overall sgRNA efficiency

or may be attributed to inaccurate TSS predictions, or complex promoter dynamics and the

preferential use of different TSS regions in different conditions (70).

Use of CRISPRa to study antifungal drug resistance phenotypes

To validate that the CRISPRa system could recapitulate phenotypes associated with *C.*

albicans gene overexpression, we targeted genes with known phenotypes associated with high

levels of expression. Initially, we focused on *CDR1*, a well-characterized *C. albicans* gene,

which encodes a well-characterized drug efflux pump (71), and whose overexpression is known

to play a role in drug resistance to azole antifungals in laboratory and clinical isolates (72–78).

To determine an optimal range in which to target our sgRNA in the *CDR1* promoter, we

designed three sgRNAs targeting loci approximately 90 bp, 200 bp, and 290 bp upstream of the

CDR1 TSS. These sgRNAs were each cloned into our CRISPRa dCas9-VPR backbone and

transformed into *C. albicans*. Next, we quantified the expression of *CDR1* in the three different

fungal strains via RT-qPCR, compared with a control strain containing the CRISPRa plasmid

with a non-targeting sgRNA. We found that targeting the dCas9-VPR construct to the *CDR1*

promoter enhanced transcription by 2.9 - 5.6 fold (**FIGURE 3a**), indicating that this CRISPRa

system is indeed able to drive enhanced levels of gene expression from a target gene of

interest. For *CDR1*, we found the highest level of overexpression was obtained by targeting the

dCas9-VPR construct ~200 bp upstream of the TSS (**FIGURE 3a**), which helped inform the

design of sgRNAs for subsequent target genes.

In order to determine if *CDR1*-overexpressing strains had measurable changes in

antifungal drug resistance phenotypes, we performed minimum inhibitory concentration (MIC)

assays with these strains in the presence of the antifungal drug fluconazole. We found that

strains overexpressing *CDR1* exhibited increased resistance to fluconazole compared with the

control strain and that higher levels of overexpression corresponded with higher levels of

resistance (**FIGURE 3b**). Together, this confirms that our CRISPRa system can be exploited to

functionally overexpress target genes of interest, and recapitulate phenotypes associated with

high levels of gene expression. Further, targeting dCas9-VPR to different promoter loci leads to

variable levels of overexpression, which could be used to study correlations between expression

levels and associated phenotypes.

We next sought to determine the ability of our CRISPRa system to be employed to

characterize a mutant strain that, to our knowledge, has never been profiled with regard to

antifungal drug susceptibility. We selected the gene *YAK1* as a target for overexpression with

CRISPRa. *C. albicans YAK1* is a predicted serine-threonine protein kinase, inactivation of which

renders cells more sensitive to the antifungal amphotericin B (79). While *YAK1* overexpression

has been shown to promote filamentation in *C. albicans* (80), and there is emerging evidence

for a direct relationship between the two phenotypes (81), the role of overexpressing this factor

in antifungal drug resistance has not previously been established. We generated a CRISPRa

strain targeting *YAK1* for overexpression (**FIGURE 3c**) and profiled the ability of this strain to

grow in the presence of amphotericin B. We found that the *YAK1* CRISPRa strain was more

resistant to amphotericin B based on MIC testing on this strain compared to the control strain

(**FIGURE 3d**). This highlights our capacity to exploit this CRISPRa system to investigate novel

phenotypes in *C. albicans*.

Validation of CRISPRa overexpression targeting fungal pathogenesis traits

Next, we aimed to validate the CRISPRa system as a means to assess *C. albicans*

genes involved in fungal pathogenesis traits. We focused on *ALS1*, a well-characterized *C.*

albicans adhesin gene, with important roles in adherence, biofilm formation, and virulence (25,

82–86). Previous work has demonstrated that overexpression of *ALS1* via promoter

replacement increases fungal adherence (83), therefore we chose to overexpress this gene

using our CRISPRa system. We designed and cloned two sgRNAs targeting approximately

130bp and 260bp upstream of the *ALS1* TSS, and generated *C. albicans ALS1* CRISPRa

strains with each of these constructs. We performed RT-qPCR to confirm the overexpression of

these strains, which enhanced the expression of *ALS1* by ~25 and ~6 fold, respectively

(**FIGURE 4a**).

To confirm phenotypes associated with overexpression of *ALS1*, we performed two

distinct biofilm growth assays that compared the *ALS1* CRISPRa strain to the CRISPRa control

strain. In the first assay, we allowed biofilms to establish over 24h in flat-bottom 96-well

polystyrene plates, removed planktonic cells, and quantified the metabolic activity of the biofilm

via XTT reduction, relative to the planktonic cell population. We found that the *ALS1* CRISPRa

strain formed significantly more robust biofilms, relative to the control strain ($P < 0.0005$;

FIGURE 4b). Interestingly, biofilm growth was not correlated with *ALS1* overexpression levels,

as the strain with ~25-fold *ALS1* overexpression formed less robust biofilms compared with the

strain expressing ~6-fold more *ALS1*. This suggests that very high levels of *ALS1* may

ultimately impair biofilm formation, or that the level of *ALS1* expression varies in the conditions

used for RT-qPCR analysis compared with biofilm growth, perhaps due to condition-specific

repositioning in the TSS.

Next, to mimic conditions associated with fungal biofilm formation in catheter-associated

urinary tract infections (CAUTIs), we used a urine-based assay and analyzed biofilm formation

of a *C. albicans* strain overexpressing *ALS1*, relative to its control strain, using crystal violet

staining. For this assay, we followed up on the *ALS1* CRISPRa strain with ~6-fold

overexpression of *ALS1*, which had more robust biofilm growth than the control strain (**FIGURE**

4a,b). The *C. albicans* strains were incubated for 48h in human urine supplemented with bovine

serum albumin (BSA) in fibrinogen-coated 96-well plates. Fibrinogen has been shown to be a

critical protein for pathogen binding in CAUTIs, by providing a scaffold for biofilm formation and

a nutritional source for pathogens (87–89). The *ALS1* CRISPRa overexpression strain showed

significantly greater biofilm formation compared to the corresponding wild-type strain ($P <$

0.0001; **FIGURE 4c**). This difference in biofilm formation suggests that *ALS1* promotes binding

to fibrinogen and plays a vital role in biofilm formation in urine.

To assess the specificity of this CRISPRa system and ensure no off-target effects from

the dCas9-sgRNA complex, we monitored global changes in transcription in the *ALS1* CRISPRa

strain via RNA-sequencing (RNA-seq). Since *ALS1* is a member of the large agglutinin-like

sequence (ALS) gene family (90, 91), it may be a likely candidate for any potential off-target

effects. We also considered *ALS1* as a strong candidate for this RNA-seq off-target profiling

since *ALS1* is not known to be a transcriptional regulator, and therefore its overexpression is

unlikely to lead to direct downstream transcriptional alterations that would obscure our analysis

of CRISPRa off-target effects. Therefore, we performed RNA-seq on the validated CRISPRa

strain overexpressing *ALS1* by ~25-fold as well as on a control strain containing the dCas9

cassette and a non-targeting sgRNA when grown in standard laboratory conditions. This

analysis confirmed the specific overexpression of *ALS1* (log2 fold change ~3.3 via RNA-seq

analysis, adjusted *P*-value 3.07E-142; **FIGURE 4d, TABLE S2**). Five additional transcripts were

identified as significantly modulated (adjusted *P*-value < 0.05, absolute log2 fold change > 1;

TABLE S2). However, their altered expression was much less significant than *ALS1* (**FIGURE**

4d), none of them having an adjusted P-value of less than 0.03, differing from the adjusted P-

value of 3e-142 for *ALS1.* We suspect these additional transcripts' altered expression is likely

unrelated to the CRISPRa system based on the fact that 4/5 genes were down-regulated

instead of up-regulated, and none of the genes are in proximity to the *ALS1* locus, nor do they

appear to share any sgRNA target similarities in their promoter regions. Together this suggests

that this CRISPRa system is likely specific with regards to gene targeting.

Discussion

Here, we present the development, validation, and application of a single-plasmid-based CRISPRa platform for efficient genetic overexpression in *C. albicans*. We demonstrate the ability of this system to overexpress genes ~2-20-fold, and recapitulate important biological phenotypes associated with genetic overexpression, including biofilm formation and antifungal drug resistance. In addition, we assess sgRNA design principles based on predicted TSS, and validate the specificity of this system by monitoring off-target effects via RNA-seq. Together, this work demonstrates an effective, efficient, and specific tool for genetic overexpression in *C. albicans* to enhance the available techniques for genetic modulation in this important fungal pathogen.

Studying genetic overexpression is an important strategy for functional genetic analysis, as overexpressing genes can mimic natural biological states where genes are expressed at high levels. In *C. albicans*, there are numerous scenarios in which the overexpression of genes occurs, including mutations that modify the function of transcriptional regulators and promote the expression of downstream genes (36, 92), as well as aneuploidies such as trisomies (38, 93–98), isochromosome formation (40, 94), and copy number variations (45, 93). These overexpression events are frequently detected amongst *C. albicans* clinical isolates (93, 94, 98) and can play an important role in drug resistance (40, 45, 94, 95), host adaptation (38, 99), and other stress tolerance phenotypes (100, 101). For example, amplification of part or all of chromosome 5, via trisomy or isochromosome formation has been tightly linked to the development of resistance to azole antifungals in *C. albicans* via the increased copy number of genes encoding the azole target, Erg11, and the transcriptional regulator of drug efflux, Tac1 (40). The prevalence and importance of these phenomena associated with increased gene expression highlight the relevance of developing new tools to targetedly assess phenotypes associated with genetic overexpression in *C. albicans*.

Harnessing the cell's natural transcriptional machinery to increase gene expression with

CRISPRa also leads to a unique set of obstacles. Off-target effects as a result of promiscuous

dCas9 binding in the cell presents the possibility of complications if the dCas9 is also targetting

other regions in the genome. While our RNA-seq results suggest that our CRISPRa system is

specific when paired with one sgRNA targeted to the *ALS1* gene, there is still the possibility of

off-target effects occurring in different CRISPRa constructs, especially as binding specificity with

CRISPR-Cas9 systems is highly dependent on the exact sgRNA sequence in use (102). Using

complementary genetic tools to confirm phenotypes observed from CRISPRa strains can help

mitigate this. One phenomenon shared among many different CRISPRa activation domains,

including VPR, is a reduced capacity to induce overexpression in genes that are already

expressed at a high levels (103). Incidentally, the gene that we achieved the lowest

overexpression with across all of our testing, *FGR28*, had the lowest basal expression (data not

shown) from our RT-qPCR experiments, opposing the idea that genes that are normally

expressed at low levels tend to be more easily overexpressed with CRISPRa systems.

However, efficiency of the sgRNA molecule and its exact position in the target region of the

gene of interest, as well as a host of promoter dynamics, can also influence CRISPR-Cas9

efficacy (69, 104, 105). This illustrates why multiple sgRNAs need to be designed when

targeting a gene with CRISPRa to maximize successful overexpression of the target gene.

While the rapid and inexpensive nature of CRISPRa strain construction easily justifies the need

to design several sgRNA molecules for any given gene, broader limitations as a result of the

specific "NGG" PAM recognition site by traditional *Streptococcus pyogenes* Cas9 proteins, like

the one used here, leads to the restraint that on rare occasions it may not be possible to

produce any highly-efficient sgRNA molecules in a suitable region of the promoter (11). Since

this PAM restriction is a feature shared among many CRISPR-Cas9 systems, it may therefore

be prudent to further expand the CRISPR toolbox in *C. albicans* by introducing novel Cas9

variants with differing PAM recognition sites into our system, or by constructing novel CRISPR

systems exploiting alternative Cas proteins with different and broader PAM recognition

capacities, offering the ability to target regions in the genome that cannot be manipulated with

current CRISPR-Cas9 tools in *C. albicans* (106, 107).

While large-scale CRISPRa libraries have not been generated in *C. albicans*, their

application in other organisms, namely mammalian cell lines, has enabled important discoveries

in cancer biology, stem cell reprogramming, cellular fitness, and drug resistance (30, 67, 108–

113). The development of CRISPRa systems in *C. albicans* could enable the development of

larger overexpression libraries for functional genomic screens, to complement existing fungal

gene deletion and depletion libraries (114–117) that have been subject to robust screening and

analysis under diverse conditions (118), to reveal critical new insight into fungal biology (119,

120), pathogenesis (115, 120–124), and mechanisms of drug and stress tolerance (125–128).

Currently, non-CRISPR-based overexpression libraries have been constructed and

screened in *C. albicans*, revealing novel insights into gene function related to hyphal

morphogenesis (60), cell cycle progression (129), drug tolerance (130), and biofilm formation

(131). Previous comparisons done between CRISPRa and ORF-based overexpression libraries

such as these in other species have highlighted the potential for these two complementary

strategies to both confirm the results from one another as well as unveil significant genetic hits

distinct from each other, even when libraries share similar gene targets (67). Compared with

other overexpression methods, the ability of the CRISPRa system to target any specified region

in the genome also confers the unique and powerful ability of allele-specific targeting which

could potentially be used to study heterozygosity in *C. albicans*, an aspect of its genomic

makeup that is critical to its pathogenicity (132). CRISPRa also offers the advantage of being

easily scalable into a pooled format whereby hundreds or thousands of unique plasmids and

resulting mutant strains can be constructed simultaneously, and mutant libraries in different

strain backgrounds can therefore be quickly constructed (133, 134). Non-CRISPR

overexpression libraries that have previously been assembled in *C. albicans* have the potential

weakness of missing important phenotypes during screening if the level of overexpression

achieved in each mutant strain is too low (46). The ability of our system to cause upwards of 25-

fold overexpression in gene targets is thus an important feature. Furthermore, corresponding

phenotypes in mutant strains may only be observable if the level of overexpression closely

mimics the level of upregulation of the gene induced by the cell in other relevant stress

conditions naturally (46). Future iterations of this CRISPRa system could also exploit an

inducible dCas9-VPR construct for tunability of gene expression. It is therefore imperative to

employ tools such as CRISPRa that possess the advantage of being inherently flexible and

easily tunable in order to study genetic overexpression thoroughly.

The application of similar CRISPRa systems to other fungal organisms would lend

further insight into fungal biology across this kingdom. Many CRISPR tools have been

developed in a diversity of fungal organisms (17, 22, 135–139), including numerous non-

albicans Candida species (11, 140–146), and other prominent pathogens such as *Cryptococcus*

neoformans (147–149) and *Aspergillus fumigatus* (150–153). These CRISPR platforms could be

adapted to similar CRISPRa tools in these organisms, with many possible applications for the

analysis of pathogen biology or drug target identification (49–52, 154).

Genetic overexpression systems also have many useful applications for industrially-

important fungi, as targeted genetic overexpression can be exploited for metabolic engineering

to enhance the production of commercially-important metabolites (136, 155–157). CRISPR

techniques have already been adapted to numerous industrially-relevant fungi (157–160), and

CRISPRa has recently been optimized for use in the filamentous fungi *Aspergillus nidulans* and

Penicillium rubens (161, 162). These CRISPRa systems were applied to activate

transcriptionally-silent biosynthetic gene clusters in these filamentous fungi, with important

applications for the discovery of novel bioactive molecules (161, 162). As many *Candida*

species play important industrial roles in manufacturing metabolites for food and pharmaceutical

industries, as well as remediating wastewater via hydrocarbon degradation (163–167), species-

specific adaptations to the CRISPRa tool presented here could produce optimized *Candida*

strains for valuable biomanufacturing and bioremediation applications.

Materials and Methods

Plasmid design and cloning

The plasmid backbone used in this study was the *C. albicans*-optimized CRISPR-dCas9

plasmid (pRS143, Addgene #122377) used in our previous study (26), containing the *NEUT5L*

integration site, sgRNA cloning site (*SNR52* promoter, SapI cloning locus, and sgRNA tail), and

dCAS9. The dCas9-VPR fusion construct was generated via Gibson assembly. The VPR

tripartite complex was codon-optimized for *C. albicans* expression and synthesized as gBlocks

gene fragments from Integrated DNA Technologies (IDT). This gene fragment was cloned with

Gibson assembly into the *dCAS9* plasmid backbone. We have made the CRISPRa (dCas9-

VPR) plasmid available via Addgene (reference #182707). All plasmids are listed in **TABLE S1.**

sgRNA design and cloning

The sgRNA CRISPR RNAs (crRNA; 20 nucleotide sequence complementary to the target DNA

genomic DNA) were designed based on efficiency and predicted specificity via the sgRNA

design tool Eukaryotic Pathogen CRISPR gRNA Design Tool (EuPaGDT;

http://grna.ctegd.uga.edu) (168). These sgRNA crRNA N20 sequences were cloned into the

dCas9-VPR CRISPRa plasmid at the sgRNA cloning locus (SapI cloning site, flanked by *SNR52*

promoter, and sgRNA tail) using Golden Gate cloning (169), as previously described (35, 170).

crRNA N20 sequences were obtained as two oligonucleotides from IDT in forward and reverse

complement orientation, each containing a SapI cloning site, and were reconstituted to 100µM

in a nuclease-free duplex buffer from IDT. Equal volumes of the two complementary

oligonucleotides were combined after being heated separately at 94°C, and then duplexed by

heating to 94°C for 2 min and cooling to room temperature. The duplexed fragment was then

cloned into the CRISPRa plasmid with the following Golden Gate cloning reaction: 10µl

miniprepped CRISPRa plasmid, 1µl duplexed oligonucleotide, 2µl 10X CutSmart buffer, 2µl

ATP, 1µl SapI, 1µl T4 DNA ligase, and 3µl nuclease-free water. This mixture was incubated in a

thermocycler under the following cycling conditions: (37°C, 2 min; 16°C, 5 min) for 99 cycles;

65°C, 15 min; 80°C, 15 min. After cycling, 1µl of additional SapI was added to each reaction

mixture, and the reaction was incubated at 37°C for 1h. All sgRNAs are listed in **TABLE S1.**

Media and growth conditions

413   *Escherichia coli* DH5α cells were grown at 37°C in Lysogeny Broth (LB) and LB plates

414   supplemented with 100mg/mL ampicillin (AMP) and 250mg/mL nourseothricin (NAT) for plasmid

415   selection. *C. albicans* cells were grown at 30°C or 37°C in Yeast Peptone Dextrose (YPD) broth

416   and YPD plates supplemented with 250mg/mL nourseothricin (NAT) for plasmid selection.

Bacterial transformation

419   High Efficiency 5-alpha Competent *E. coli* (NEB) cells were thawed on ice. 1µl of Golden Gate

420   cloning reagents were added to 50µl of competent cells and incubated on ice for 30 min, heat

421   shocked for 30s at 42°C, and incubated on ice for an additional 5 min. 950µl of SOC outgrowth

422   media was added to each cell culture and incubated at 30°C for 1.5h at 250 RPM. Transformed

423   cells were then plated on LB media containing AMP and NAT and grown at 30°C for 1 day.

C. albicans transformation

Plasmids were transformed into *C. albicans* via a chemical transformation strategy, as

previously described (35, 170). Briefly, miniprepped CRISPRa plasmids were linearized via a

1.5X restriction digest mix using the PacI enzyme. A transformation master mix was prepared

as follows: 800µl of 50% polyethylene glycol (PEG), 100µl of 10X Tris-EDTA (TE) buffer

solution, 100µl of 1M lithium acetate (LiAc), 40µl of 10 mg/mL salmon sperm DNA, and 20µl of

1M dithiothreitol (DTT). The transformation mix was added to *C. albicans* cells and linearized

CRISPRa plasmid and left to incubate at 30°C for 1h, then heat-shocked at 42°C for 45 min.

Cells were washed with fresh YPD, then grown in YPD for 4h to allow for expression of the NAT

resistance construct. Transformed cells were then plated on YPD media containing NAT, and

grown at 30°C for 2 days. All *C. albicans* strains are listed in **TABLE S1.**

Growth curves

Overnight cultures of *C. albicans* grown in YPD were diluted to an OD600 of 0.05 in fresh YPD

and added to a 96-well flat-bottomed plate with a total volume of 200μl per well. Growth was

measured at 30°C via optical density at 600nm at 15 min intervals over the course of 24h using

an Infinite 200 PRO microplate reader (Tecan), and plates were shaken orbitally for 900s at a

4mm amplitude in between growth measurements.

Minimum inhibitory concentration (MIC) assays

Fluconazole MIC assays were performed in 96-well flat-bottomed plates. A suspension of

40μg/mL fluconazole was prepared in water, of which 100μl was added to the first column of

each plate containing 100μl of culture, to obtain a total volume of 200μl per well. The first

column was serially diluted 2-fold across the plate in water. The gradient of fluconazole,

therefore, ranged from 20μg/mL to 0μg/mL. Overnight cultures of *C. albicans* grown in YPD

were diluted to an OD600 of 0.1 in 2X RPMI 1640 with 40g/L of added D-glucose and mixed into

the plates in an equal volume such that the starting OD600 values of each strain were 0.05 and

the growth media was diluted to RPMI 1640 with 20g/L of added D-glucose. Plates were

incubated at 37°C at 900 RPM, and absorbance values at 600nm were read after 24h using an

Infinite 200 PRO microplate reader (Tecan). The amphotericin B MIC assays were also

performed in 96-well flat-bottomed plates similar to the protocol described above with a few

modifications. A starting concentration of 5μg/mL of amphotericin B was used to create a

gradient ranging from 2.5µg/mL to 0µg/mL. Strains were instead grown in YPD media using a

starting OD600 of 0.001. Plates were incubated at 37°C statically and absorbance values at

600nm were read after 72h using an Infinite 200 PRO microplate reader (Tecan).

Biofilm assays

RPMI-based biofilm assays were performed as previously described (171), with minor

modifications. Overnight cultures of *C. albicans* grown in YPD were diluted to an OD600 of

0.001 in 5mL of RPMI 1640 with 20g/L of supplemented D-glucose. 100µl of RPMI 1640 was

mixed with 100µl of each diluted overnight culture into a 96-well flat-bottomed plate (12 wells

per strain). Plates were grown statically at 37°C for 24h to allow biofilms to form. 120µl of

planktonic cell supernatant was removed from each well and moved to a new 96-well flat-

bottomed plate. Density of the planktonic cells was measured at 600nm using an Infinite 200

PRO microplate reader (Tecan). Original plates from which planktonic cells were removed were

washed by adding 200µl 1X PBS to each well with an electronic multichannel pipette at a

moderate dispense speed. 1X PBS was then discarded, and this wash step was repeated.

Plates were then left to dry for 1 - 2h. Once biofilms were dry, 90µl of 1mg/mL XTT (prepared in

1X PBS and centrifuged to remove sediment prior to use) and 10µl of 0.32mg/mL PMS

(prepared in water) were added to each well. Plates were incubated statically at 30°C for 2h to

allow biofilms to reduce XTT, measured at 490nm using an Infinite 200 PRO microplate reader

(Tecan), and normalized to the growth of planktonic cells harvested previously.

For urine-based biofilm assays, biofilm formations were performed in 96-well flat-

bottomed plates. Plates were coated with 150µg/mL of fibrinogen and incubated overnight at

4°C. The *C. albicans* strains were cultured at 37°C with aeration in 5mL of YPD broth. The

inoculum was normalized to ~1×10^6 CFU/ml and then diluted (1:10) in human urine (female, pH

~6.5). Urine was supplemented with 20mg/mL of BSA for carbon and nitrogen sources

mimicking the catheterized bladder environment. 200µl of the urine with inoculum was incubated

in each well of the 96-well plate at 37°C for 48h while static. Following the 48h incubation, the

cultures were removed and the wells were washed three times with 200µl 1X PBS to remove

unbound fungi. Plates were incubated with 200µl of 0.5% crystal violet for 15 min. Crystal violet

was removed, and plates were washed with water and dried. 200µl of 33% acetic acid was

added to the wells and 100µl was transferred to a new 96-well plate. Absorbance was measured

at 595nm using a SpectraMax ABS Plus microplate reader (Molecular Devices).

RNA extraction and real-time quantitative PCR

To detect differences in gene expression, overnight cultures of *C. albicans* grown in YPD

were diluted to an OD600 of 0.05 in fresh YPD and grown to an OD600 of >0.2 at 30°C.

Cultures were pelleted and frozen at -80°C before RNA was extracted. RNA extractions were

performed using either a Presto Mini RNA Yeast Kit from FroggaBio (cat. RBYD050) or an

RNeasy Mini Kit from Qiagen (cat. 74104). To assess RNA integrity, an RNA ScreenTape assay

was performed on all samples using the TapeStation 4150 system following manufacturer's

instructions (Agilent Technologies). Only samples with RIN values of >5.0 were used for RT-

qPCR. Synthesis of cDNA from 1000ng (10µl) RNA was performed using a High Capacity cDNA

Reverse Transcription Kit from Applied BioSystems (cat. 4368814). Briefly, 20µl reaction

mixtures were prepared with 2µl 10X reverse transcription buffer, 0.8µl of 25X dNTPs at a

concentration of 100mM, 2µl of random primers, 1µl of MultiScribe Reverse Transcriptase at 50

U/µl, 4.2µl of nuclease-free water, and 10µl of RNA (1µg). Reverse transcription run conditions

were as follows: 10 min at 25°C, 120 min at 37°C, 5 min at 85°C. Real-time PCR assays were

conducted using a QuantStudio 7 Pro Real Time PCR system from Thermo Fisher Scientific Inc.

20µl reaction mixtures containing 10µl 2X SsoAdvanced Universal Inhibitor-Tolerant SYBR

supermix (Bio-Rad, cat: 172-5017), 0.8µl of PCR forward and reverse primer mix at 5µM (final

concentration of primer at 200 nM), 4.2µl of water and 5µl of diluted cDNA. The run conditions

were as follows: 3 min at 98°C polymerase activation step, followed by 40 cycles of a two-step

qPCR (10s of 98°C denaturation, 30s of 60°C combined annealing/extension). Primers are

listed in **TABLE S1**. Expression profiling calculations were performed according to the

comparative C_T method (172). Briefly, expression values for the experimental gene of interest

were compared to the housekeeping gene *ACT1* to obtain a ΔC_T value within each strain. The

ΔC_T values in the experimental CRISPRa strains were then compared to the non-targeting

CRISPRa control strain to obtain a $\Delta\Delta C_T$ value and finally a fold-difference in expression of the

experimental gene.

RNA sequencing

RNA preparation and sequencing were performed as previously described, with minor

modifications (171). Strains were grown in YPD at 30°C overnight preceding RNA extraction.

RNA extraction, sample QC, library preparations, and sequencing reactions were conducted at

GENEWIZ, LLC./Azenta US, Inc. as follows: Total RNA was extracted using Qiagen RNeasy

Plus Universal kit following manufacturer's instructions (Qiagen). RNA samples were quantified

using Qubit 2.0 Fluorometer (ThermoFisher Scientific) and RNA integrity was checked with

4200 TapeStation (Agilent Technologies). ERCC RNA Spike-In Mix kit (cat. 4456740) from

ThermoFisher Scientific, was added to normalized total RNA prior to library preparation

following the manufacturer's protocol. The RNA sequencing library was prepared using the

NEBNext Ultra II RNA Library Prep Kit for Illumina using the manufacturer's instructions (New

England Biolabs). Briefly, mRNAs were initially enriched with Oligod(T) beads. Enriched mRNAs

were fragmented for 15 min at 94 °C. First-strand and second-strand cDNA were subsequently

synthesized. cDNA fragments were end-repaired and adenylated at 3'ends, and universal

adapters were ligated to cDNA fragments, followed by index addition and library enrichment by

PCR with limited cycles. The sequencing library was validated on the Agilent TapeStation

(Agilent Technologies), and quantified by using Qubit 2.0 Fluorometer (ThermoFisher Scientific)

as well as by quantitative PCR (KAPA Biosystems). The sequencing libraries were multiplexed

and clustered onto a flowcell. After clustering, the flowcell was loaded onto the Illumina HiSeq

4000 or equivalent instrument according to the manufacturer's instructions. The samples were

sequenced using a 2x150bp Paired End (PE) configuration. Image analysis and base calling

were conducted by the Illumina Control Software.

Raw sequence data (.bcl files) generated from the Illumina instrument was converted

into fastq files and de-multiplexed using Illumina bcl2fastq 2.20 software. One mis-match was

allowed for index sequence identification. Sequence reads were trimmed to remove possible

adapter sequences and nucleotides with poor quality using Trimmomatic v.0.36, and trimmed

reads were mapped to the candida_albicans reference genome available on ENSEMBL using

the STAR aligner v.2.5.2b. The STAR aligner is a splice aligner that detects splice junctions and

incorporates them to help align the entire read sequences. BAM files were generated as a result

of this step. Unique gene hit counts were then calculated by using featureCounts from the

Subread package v.1.5.2. The hit counts were summarized and reported using the gene_id

feature in the annotation file. Only unique reads that fell within exon regions were counted. After

extraction of gene hit counts, the gene hit counts table was used for downstream differential

expression analysis. Using DESeq2, a comparison of gene expression between the control and

experimental groups of samples was performed. The Wald test was then used to generate p-

values and log2 fold changes. Genes with an adjusted p-value < 0.05 and absolute log2 fold

change > 1 were called as differentially expressed genes for each comparison. The data have

been deposited in NCBI's Gene Expression Omnibus (GEO) (Edgar *et al.* 2002) and are

accessible through GEO Series accession number GSE201904

(https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE201904).

Data availability

The *C. albicans* CRISPRa plasmid backbone was deposited to Addgene as plasmid number

#182707. RNA sequencing data was deposited in NCBI GEO repository (accession number

GSE201904; https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE201904).

Acknowledgments

We thank Shapiro lab members, past and present, for helpful discussions and support of this

project, as well as Dr. Anna Selmecki and Dr. Petra Vande Zande (University of Minnesota) for

helpful feedback. We thank Dr. Zhenguo Lin (Saint Louis University) for helpful discussions on

TSS prediction, and Kieran Shah for programming support on this project. We thank Jing Zhang

from the University of Guelph Advanced Analysis Centre for RT-qPCR support. This work was

supported by an NSERC Discovery Grant (RGPIN-2018-4914), an Ontario Early Research

Award, and a CIFAR Azrieli Global Scholar award to RSS. N.C.G, L.W, and M.F. are

EvoFunPath Fellows (NSERC CREATE), L.W. is supported by an Ontario Graduate

Scholarship, M.F. is supported by an NSERC-CGS-M scholarship, and R.O.C.L was supported

by a MITACS Globalink Research Internship. A.C. is supported by a Career Award for Medical

Scientist from the Burroughs Wellcome Fund. Work by A.L.F.M. and F.H.S.T. was supported by

institutional funds from the University of Notre Dame as well as National Institute of Health grant

R01-DK128805 (to A.L.F.M. and A.A.L.), and the Arthur J. Schmitt Presidential Leadership

Foundation (to A.L.L).

Figure Legends

Figure 1. Design of a CRISPRa system for *C. albicans*. (a) The single plasmid system

contains a catalytically inactive ("dead") dCas9 protein, paired with the VPR (VP64-P65-Rta)

complex of activator domains to cause upregulation at a target locus. The single-guide RNA

(sgRNA) region can be subjected to Golden Gate cloning with a custom N20 sequence in order

to target the dCas9-VPR complex to a desired region of DNA through complementary base-

pairing. The PacI enzyme is used to linearize the plasmid at the *NEUT5L* locus, which allows

the plasmid to be integrated into the cell's chromosome and remain in the cell following cell

division. Panel created with BioRender.com. **(b)** Once the sgRNA recognizes and binds to a

target sequence of DNA, the dCas9-VPR complex will cause the recruitment of transcription

initiation complex factors to the adjacent genetic material, thus increasing gene expression from

that region. Created with BioRender.com. **(c)** Wild-type *C. albicans* cells and cells containing a

non-targeting CRISPRa plasmid were grown in YPD at 30°C to compare fitness under standard

laboratory conditions. Growth was recorded via optical density at 600nm at 15 min intervals over

the course of 24h. Data points show the mean and standard deviation at each time point (n =

32).

Figure 2. Determining optimal guide-design principles for CRISPRa overexpression in *C.*

***albicans*.** CRISPRa strains were created using six different sgRNA molecules designed to

target different positions in the promoter region of each of **(a)** *KSP1*, **(b)** *ALS3*, and **(c)** *FGR28*

upstream of either the start codon or the predicted dominant transcriptional start site (TSS).

Fold-change in the expression of the target gene relative to the housekeeping gene *ACT1* in the

experimental strains and the non-targeting CRISPRa control strain was measured with RT-

qPCR and compared. Data points represent the mean fold-difference and standard deviation in

expression of the respective target gene in the CRISPRa strains compared to the non-targeting

CRISPRa control (n = 3).

Figure 3. Using CRISPRa to validate antifungal drug resistance phenotypes. (a) Three *C.*

albicans CRISPRa strains were created targeting the drug transporter gene *CDR1* for

overexpression. Fold-change in the expression of *CDR1* relative to the housekeeping gene

ACT1 in the three experimental strains and the non-targeting CRISPRa control strain was

measured with RT-qPCR. All three experimental strains are represented by the position of their

sgRNA in the promoter region upstream of the proposed dominant *CDR1* transcriptional start

site. Data represent the mean fold-difference and standard deviation in expression of *CDR1* in

the CRISPRa strains compared to the non-targeting CRISPRa control (n = 3). **(b)** The fitness of

all three CRISPRa strains targeting *CDR1* for overexpression was measured in the presence of

the antifungal drug fluconazole with a minimum inhibitory concentration assay. The OD600

value of each strain in each concentration of fluconazole was measured to represent growth in

differing conditions, where OD600 values approaching 1.0 reflect strong fitness. **(c)** Fold-change

in the expression of *YAK1* relative to the housekeeping gene *ACT1* in the experimental strain

and the non-targeting CRISPRa control strain was measured with RT-qPCR. Data represent the

mean fold-difference and standard deviation in expression of *YAK1* in the CRISPRa strains

compared to the non-targeting CRISPRa control (n = 3). **(d)** The fitness of the CRISPRa strain

targeting *YAK1* for overexpression was measured in the presence of the antifungal drug

amphotericin B with a minimum inhibitory concentration assay. The OD600 value of each strain

in each concentration of amphotericin B was measured to represent growth in differing

conditions, where OD600 values approaching 1.0 reflect strong fitness.

Figure 4. Validation of CRISPRa overexpression and target-specificity through targeting

fungal pathogenesis traits. (a) Two *C. albicans* CRISPRa strains targeting different positions

in the promoter region of the adhesin protein gene *ALS1* were created. Fold-change in the

expression of *ALS1* relative to the housekeeping gene *ACT1* in both experimental strains and

the non-targeting CRISPRa control strain was measured with RT-qPCR. Experimental strains

are represented by the position of their sgRNA in the promoter region upstream of the proposed

dominant *ALS1* transcriptional start site. Data points represent the mean fold-difference and

standard deviation in expression of *ALS1* in the CRISPRa strains compared to the non-targeting

CRISPRa control (n = 3). **(b)** Two CRISPRa strains targeting *ALS1* and a non-targeting

CRISPRa control strain were grown in the biofilm-inducing media RPMI and left to grow

statically at 37°C for 24h to allow biofilms to form. The robustness of biofilms formed by all

strains was elucidated through an XTT reduction assay and measuring of the resulting OD490

values before normalizing final values to non-biofilm-forming planktonic cells. Data points

represent the mean and the standard error of the mean (n = 72). Differences between groups

were tested for significance using Dunnett's multiple comparison test. ****, p-value <0.0001. ***,

p-value <0.001. **(c)** A *C. albicans* CRISPRa strain overexpressing *ALS1* ~6-fold and a non-

targeting CRISPRa strain were grown in human urine supplemented with fibrinogen and BSA at

37°C for 48h to simulate a catheterized bladder environment. The robustness of biofilms formed

by both strains was measured by reading the OD595 value following a crystal violet assay. Data

points are combined from three independent experiments where n = 32 for the non-targeting

CRISPRa control and n = 48 for the CRISPRa *ALS1* overexpression strain. Differences

between groups were tested for significance using the Mann-Whitney U test. ****, p-value

<0.0001. **(d)** Global gene expression of a CRISPRa strain overexpressing *ALS1* ~25-fold, as

well as a non-targeting CRISPRa control strain, was measured using RNA-Seq. Differential

gene expression analysis showed that *ALS1* had a log2-fold change of ~3.3 in the *ALS1*

overexpression strain compared to the CRISPRa control strain, with an adjusted p-value of

3.07E-142, and showed no additional transcripts that had differential expression with an

adjusted p-value of less than 0.03.

References

1. Fisher,M.C., Gurr,S.J., Cuomo,C.A., Blehert,D.S., Jin,H., Stukenbrock,E.H., Stajich,J.E.,
Kahmann,R., Boone,C., Denning,D.W., *et al.* (2020) Threats posed by the fungal kingdom
to humans, wildlife, and agriculture. *mBio*, **11**, e00449-20.

2. Pérez,J.C. (2019) *Candida albicans* dwelling in the mammalian gut. *Curr. Opin. Microbiol.*,
52, 41–46.

3. Gow,N.A.R., van de Veerdonk,F.L., Brown,A.J.P. and Netea,M.G. (2011) *Candida albicans*
morphogenesis and host defence: discriminating invasion from colonization. *Nat. Rev.*
Microbiol., **10**, 112–122.

4. Bongomin,F., Gago,S., Oladele,R. and Denning,D. (2017) Global and multi-national
prevalence of fungal diseases—estimate precision. *Journal of Fungi*, **3**, 57.

5. Segrelles-Calvo,G., de S Araújo,G.R., Llopis-Pastor,E., Carrillo,J., Hernández-Hernández,M.,
Rey,L., Melean,N.R., Escribano,I., Antón,E., Zamarro,C., *et al.* (2021) *Candida* spp. co-
infection in COVID-19 patients with severe pneumonia: Prevalence study and associated
risk factors. *Respir. Med.*, **188**, 106619.

6. Benedict,K., Jackson,B.R., Chiller,T. and Beer,K.D. (2018) Estimation of direct healthcare
costs of fungal diseases in the United States. *Clin. Infect. Dis.*, **68**, 1791-1797.

7. Perfect,J.R. (2017) The antifungal pipeline: a reality check. *Nat. Rev. Drug Discov.* **16**, 603-
616.

8. Geddes-McAlister,J. and Shapiro,R.S. (2018) New pathogens, new tricks: emerging, drug-
resistant fungal pathogens and future prospects for antifungal therapeutics. *Ann. N. Y.*
Acad. Sci., **1435**, 57-78.

9. Arendrup,M.C. and Patterson,T.F. (2017) Multidrug-resistant *Candida*: epidemiology,
molecular mechanisms, and treatment. *J. Infect. Dis.*, **216**, S445–S451.

10. Sanguinetti,M., Posteraro,B. and Lass-Flörl,C. (2015) Antifungal drug resistance among
Candida species: mechanisms and clinical impact. *Mycoses*, **58**, 2–13.

11. Uthayakumar,D., Sharma,J., Wensing,L. and Shapiro,R.S. (2021) CRISPR-based genetic
manipulation of *Candida* species: historical perspectives and current approaches. *Frontiers*
in Genome Editing, **2**, 606281.

12. Shapiro,R.S., Chavez,A. and Collins,J.J. (2018) CRISPR-based genomic tools for the
manipulation of genetically intractable microorganisms. *Nat. Rev. Microbiol.*, **16**, 333–339.

13. Adli,M. (2018) The CRISPR tool kit for genome editing and beyond. *Nat. Commun.*, **9**, 1911.

14. Bock,C., Datlinger,P., Chardon,F., Coelho,M.A., Dong,M.B., Lawson,K.A., Lu,T., Maroc,L.,
Norman,T.M., Song,B., *et al.* (2022) High-content CRISPR screening. *Nature Reviews*
Methods Primers, **2**, 1–1.

15. Vyas,V.K., Barrasa,M.I. and Fink,G.R. (2015) A *Candida albicans* CRISPR system permits
genetic engineering of essential genes and gene families. *Sci Adv*, **1**, e1500248.

694 16. Min,K., Biermann,A., Hogan,D.A. and Konopka,J.B. (2018) Genetic analysis of *NDT80*
695   family transcription factors in *Candida albicans* using new CRISPR-Cas9 approaches.
696   *mSphere*, **3**, e00545-18.

697 17. Morio,F., Lombardi,L. and Butler,G. (2020) The CRISPR toolbox in medical mycology: State
698   of the art and perspectives. *PLoS Pathog.*, **16**, e1008201.

699 18. Huang,M.Y. and Mitchell,A.P. (2017) Marker Recycling in *Candida albicans* through
700   CRISPR-Cas9-Induced Marker Excision. *mSphere*, **2**, e00050-17.

701 19. Min,K., Ichikawa,Y., Woolford,C.A. and Mitchell,A.P. (2016) *Candida albicans* gene deletion
702   with a transient CRISPR-Cas9 system. *mSphere* **1**, e00130-16.

703 20. Ng,H. and Dean,N. (2017) Dramatic improvement of CRISPR/Cas9 editing in *Candida*
704   *albicans* by increased single guide RNA expression. *mSphere*, **2**, e00385-16.

705 21. Nguyen,N., Quail,M.M.F. and Hernday,A.D. (2017) An efficient, rapid, and recyclable system
706   for CRISPR-mediated genome editing in *Candida albicans*. *mSphere*, **2**, e00149-17.

707 22. Román,E., Prieto,D., Alonso-Monge,R. and Pla,J. (2019) New insights of CRISPR
708   technology in human pathogenic fungi. *Future Microbiol.*, **14**, 1243–1255.

709 23. Vyas,V.K., Guy Bushkin,G., Bernstein,D.A., Getz,M.A., Sewastianik,M., Inmaculada
710   Barrasa,M., Bartel,D.P. and Fink,G.R. (2018) New CRISPR mutagenesis strategies reveal
711   variation in repair mechanisms among fungi. *mSphere*, **3**, e00154–18.

712 24. Shapiro,R.S., Chavez,A., Porter,C.B.M., Hamblin,M., Kaas,C.S., DiCarlo,J.E., Zeng,G.,
713   Xu,X., Revtovich,A.V., Kirienko,N.V., *et al.* (2018) A CRISPR-Cas9-based gene drive
714   platform for genetic interaction analysis in *Candida albicans*. *Nat Microbiol*, **3**, 73–82.

715 25. Rosiana,S., Zhang,L., Kim,G.H., Revtovich,A.V., Uthayakumar,D., Sukumaran,A., Geddes-
716   McAlister,J., Kirienko,N.V. and Shapiro,R.S. (2021) Comprehensive genetic analysis of
717   adhesin proteins and their role in virulence of *Candida albicans. Genetics*, **217**, iyab003.

718 26. Wensing,L., Sharma,J., Uthayakumar,D., Proteau,Y., Chavez,A. and Shapiro,R.S. (2019) A
719   CRISPR interference platform for efficient genetic repression in *Candida albicans*.
720   *mSphere*, **4**, e00002-19.

721 27. Román,E., Coman,I., Prieto,D., Alonso-Monge,R. and Pla,J. (2019) Implementation of a
722   CRISPR-based system for gene regulation in *Candida albicans. mSphere*, **4**, e00001-19.

723 28. Wang,H., La Russa,M. and Qi,L.S. (2016) CRISPR/Cas9 in genome editing and beyond.
724   *Annu. Rev. Biochem.*, **85**, 227–264.

725 29. Dominguez,A.A., Lim,W.A. and Qi,L.S. (2016) Beyond editing: repurposing CRISPR-Cas9
726   for precision genome regulation and interrogation. *Nat. Rev. Mol. Cell Biol.*, **17**, 5–15.

727 30. Kampmann,M. (2018) CRISPRi and CRISPRa Screens in Mammalian Cells for Precision
728   Biology and Medicine. *ACS Chem. Biol.*, **13**, 406–416.

729 31. Peters,J.M., Silvis,M.R., Zhao,D., Hawkins,J.S., Gross,C.A. and Qi,L.S. (2015) Bacterial
730   CRISPR: accomplishments and prospects. *Curr. Opin. Microbiol.*, **27**, 121–126.

32. Todor,H., Silvis,M.R., Osadnik,H. and Gross,C.A. (2021) Bacterial CRISPR screens for
gene function. *Curr. Opin. Microbiol.*, **59**, 102–109.

33. Schultenkämper,K., Brito,L.F. and Wendisch,V.F. (2020) Impact of CRISPR interference on
strain development in biotechnology. *Biotechnol. Appl. Biochem.*, **67**, 7–21.

34. Xu,X. and Qi,L.S. (2019) A CRISPR–dCas Toolbox for Genetic Engineering and Synthetic
Biology. *J. Mol. Biol.*, **431**, 34–47.

35. Wensing,L. and Shapiro,R.S. (2022) Design and Generation of a CRISPR Interference
System for Genetic Repression and Essential Gene Analysis in the Fungal Pathogen
Candida albicans. *Methods Mol. Biol.*, **2377**, 69–88.

36. Coste,A., Turner,V., Ischer,F., Morschhäuser,J., Forche,A., Selmecki,A., Berman,J., Bille,J.
and Sanglard,D. (2006) A Mutation in Tac1p, a Transcription Factor Regulating *CDR1* and
CDR2, Is Coupled With Loss of Heterozygosity at Chromosome 5 to Mediate Antifungal
Resistance in *Candida albicans. Genetics*, **172**, 2139–2156.

37. Todd,R.T., Wikoff,T.D., Forche,A. and Selmecki,A. (2019) Genome plasticity in *Candida
albicans* is driven by long repeat sequences. *Elife*, **8**, e45954.

38. Forche,A., Cromie,G., Gerstein,A.C., Solis,N.V., Pisithkul,T., Srifa,W., Jeffery,E., Abbey,D.,
Filler,S.G., Dudley,A.M., *et al.* (2018) Rapid Phenotypic and Genotypic Diversification After
Exposure to the Oral Host Niche in *Candida albicans. Genetics*, **209**, 725–741.

39. Rai,L.S., van Wijlick,L., Chauvel,M., d'Enfert,C., Legrand,M. and Bachellier-Bassi,S. (2021)
Overexpression approaches to advance understanding of *Candida albicans*. *Mol.
Microbiol.*, **117**, 589-599.

40. Selmecki,A., Gerami-Nejad,M., Paulson,C., Forche,A. and Berman,J. (2008) An
isochromosome confers drug resistance in vivo by amplification of two genes, *ERG11* and
TAC1. *Mol. Microbiol.*, **68**, 624–641.

41. Lohberger,A., Coste,A.T. and Sanglard,D. (2014) Distinct roles of *Candida albicans* drug
resistance transcription factors *TAC1*, *MRR1*, and *UPC2* in virulence. *Eukaryot. Cell*, **13**,
127–142.

42. Schubert,S., Barker,K.S., Znaidi,S., Schneider,S., Dierolf,F., Dunkel,N., Aïd,M., Boucher,G.,
Rogers,P.D., Raymond,M., *et al.* (2011) Regulation of efflux pump expression and drug
resistance by the transcription factors Mrr1, Upc2, and Cap1 in *Candida albicans*.
Antimicrob. Agents Chemother., **55**, 2212–2223.

43. Flowers,S.A., Barker,K.S., Berkow,E.L., Toner,G., Chadwick,S.G., Gygax,S.E.,
Morschhäuser,J. and Rogers,P.D. (2012) Gain-of-function mutations in *UPC2* are a
frequent cause of *ERG11* upregulation in azole-resistant clinical isolates of *Candida
albicans*. *Eukaryot. Cell*, **11**, 1289–1299.

44. Sanglard,D., Coste,A. and Ferrari,S. (2009) Antifungal drug resistance mechanisms in
fungal pathogens from the perspective of transcriptional gene regulation. *FEMS Yeast
Res.*, **9**, 1029–1050.

45. Todd,R.T. and Selmecki,A. (2020) Expandable and reversible copy number amplification

drives rapid adaptation to antifungal drugs. *eLife*, **9**, e58349.

46. Znaidi,S., van Wijlick,L., Hernández-Cervantes,A., Sertour,N., Desseyn,J.-L., Vincent,F.,
Atanassova,R., Gouyer,V., Munro,C.A., Bachellier-Bassi,S., *et al.* (2018) Systematic gene
overexpression in *Candida albicans* identifies a regulator of early adaptation to the
mammalian gut. *Cell. Microbiol.*, **20**, e12890.

47. Prieto,D., Román,E., Alonso-Monge,R. and Pla,J. (2017) Overexpression of the
Transcriptional Regulator *WOR1* Increases Susceptibility to Bile Salts and Adhesion to the
Mouse Gut Mucosa in *Candida albicans*. *Front. Cell. Infect. Microbiol.*, **7**, 389.

48. Fu,Y., Luo,G., Spellberg,B.J., Edwards,J.E.,Jr and Ibrahim,A.S. (2008) Gene
overexpression/suppression analysis of candidate virulence factors of *Candida albicans*.
Eukaryot. Cell, **7**, 483–492.

49. Luesch,H., Wu,T.Y.H., Ren,P., Gray,N.S., Schultz,P.G. and Supek,F. (2005) A genome-
wide overexpression screen in yeast for small-molecule target identification. *Chem. Biol.*,
12, 55–63.

50. Smith,A.M., Ammar,R., Nislow,C. and Giaever,G. (2010) A survey of yeast genomic assays
for drug and target discovery. *Pharmacol. Ther.*, **127**, 156–164.

51. Li,X., Zolli-Juran,M., Cechetto,J.D., Daigle,D.M., Wright,G.D. and Brown,E.D. (2004)
Multicopy suppressors for novel antibacterial compounds reveal targets and drug efflux
susceptibility. *Chem. Biol.*, **11**, 1423–1430.

52. Begolo,D., Erben,E. and Clayton,C. (2014) Drug target identification using a trypanosome
overexpression library. *Antimicrob. Agents Chemother.*, **58**, 6260–6264.

53. Sopko,R., Huang,D., Preston,N., Chua,G., Papp,B., Kafadar,K., Snyder,M., Oliver,S.G.,
Cyert,M., Hughes,T.R., *et al.* (2006) Mapping pathways and phenotypes by systematic
gene overexpression. *Mol. Cell*, **21**, 319–330.

54. Polvi,E.J., Veri,A.O., Liu,Z., Hossain,S., Hyde,S., Kim,S.H., Tebbji,F., Sellam,A., Todd,R.T.,
Xie,J.L., *et al.* (2019) Functional divergence of a global regulatory complex governing
fungal filamentation. *PLOS Genetics*, **15**, e1007901.

55. Sahni,N., Yi,S., Daniels,K.J., Huang,G., Srikantha,T. and Soll,D.R. (2010) Tec1 Mediates
the Pheromone Response of the White Phenotype of *Candida albicans*: Insights into the
Evolution of New Signal Transduction Pathways. *PLoS Biology*, **8**, e1000363.

56. Delgado,M.L., Luisa Delgado,M., Luisa Gil,M. and Gozalbo,D. (2003) *Candida albicans*
TDH3 gene promotes secretion of internal invertase when expressed in *Saccharomyces*
cerevisiae as a glyceraldehyde-3-phosphate dehydrogenase-invertase fusion protein.
Yeast, **20**, 713–722.

57. Eckert,S.E. and Mühlschlegel,F.A. (2009) Promoter regulation in *Candida albicans* and
related species. *FEMS Yeast Res.*, **9**, 2–15.

58. Park,Y.-N. and Morschhäuser,J. (2005) Tetracycline-inducible gene expression and gene
deletion in *Candida albicans*. *Eukaryot. Cell*, **4**, 1328–1342.

59. Legrand,M., Bachellier-Bassi,S., Lee,K.K., Chaudhari,Y., Tournu,H., Arbogast,L., Boyer,H.,

Chauvel,M., Cabral,V., Maufrais,C., *et al.* (2018) Generating genomic platforms to study
Candida albicans pathogenesis. *Nucleic Acids Res.*, **46**, 6935–6949.

60. Chauvel,M., Nesseir,A., Cabral,V., Znaidi,S., Goyard,S., Bachellier-Bassi,S., Firon,A.,
Legrand,M., Diogo,D., Naulleau,C., *et al.* (2012) A versatile overexpression strategy in the
pathogenic yeast *Candida albicans*: identification of regulators of morphogenesis and
fitness. *PLoS One*, **7**, e45912.

61. Chavez,A., Scheiman,J., Vora,S., Pruitt,B.W., Tuttle,M., P R Iyer,E., Lin,S., Kiani,S.,
Guzman,C.D., Wiegand,D.J., *et al.* (2015) Highly efficient Cas9-mediated transcriptional
programming. *Nat. Methods*, **12**, 326–328.

62. Gerami-Nejad,M., Zacchi,L.F., McClellan,M., Matter,K. and Berman,J. (2013) Shuttle
vectors for facile gap repair cloning and integration into a neutral locus in *Candida albicans*.
Microbiology, **159**, 565–579.

63. La Russa,M.F. and Qi,L.S. (2015) The New State of the Art: Cas9 for Gene Activation and
Repression. *Mol. Cell. Biol.*, **35**, 3800–3809.

64. Schmitz,M.L. and Baeuerle,P.A. (1991) The p65 subunit is responsible for the strong
transcription activating potential of NF-kappa B. *The EMBO Journal*, **10**, 3805–3817.

65. Hardwick,J.M., Tse,L., Applegren,N., Nicholas,J. and Veliuona,M.A. (1992) The Epstein-
Barr virus R transactivator (Rta) contains a complex, potent activation domain with
properties different from those of *VP16*. *J. Virol.*, **66**, 5500–5508.

66. Jensen,M.K. (2018) Design principles for nuclease-deficient CRISPR-based transcriptional
regulators. *FEMS Yeast Res.*, **18**, foy039.

67. Sanson,K.R., Hanna,R.E., Hegde,M., Donovan,K.F., Strand,C., Sullender,M.E.,
Vaimberg,E.W., Goodale,A., Root,D.E., Piccioni,F., *et al.* (2018) Optimized libraries for
CRISPR-Cas9 genetic screens with multiple modalities. *Nat. Commun.*, **9**, 5416.

68. Smith,J.D., Suresh,S., Schlecht,U., Wu,M., Wagih,O., Peltz,G., Davis,R.W., Steinmetz,L.M.,
Parts,L. and St Onge,R.P. (2016) Quantitative CRISPR interference screens in yeast
identify chemical-genetic interactions and new rules for guide RNA design. *Genome Biol.*,
17, 45.

69. Lu,Z. and Lin,Z. (2021) The origin and evolution of a distinct mechanism of transcription
initiation in yeasts. *Genome Res.*, **31**, 51–63.

70. McMillan,J., Lu,Z., Rodriguez,J.S., Ahn,T.-H. and Lin,Z. (2019) YeasTSS: an integrative web
database of yeast transcription start sites. *Database*, **2019**, baz048.

71. Prasad,R., Banerjee,A., Khandelwal,N.K. and Dhamgaye,S. (2015) The ABCs of *Candida*
albicans Multidrug Transporter Cdr1. *Eukaryot. Cell*, **14**, 1154–1164.

72. Prasad,R., De Wergifosse,P., Goffeau,A. and Balzi,E. (1995) Molecular cloning and
characterization of a novel gene of *Candida albicans*, *CDR1*, conferring multiple resistance
to drugs and antifungals. *Current Genetics*, **27**, 320–329.

73. Sanglard,D., Kuchler,K., Ischer,F., Pagani,J.L., Monod,M. and Bille,J. (1995) Mechanisms of
resistance to azole antifungal agents in *Candida albicans* isolates from AIDS patients

involve specific multidrug transporters. *Antimicrobial Agents and Chemotherapy*, **39**, 2378–
2386.

74. White,T.C. (1997) Increased mRNA levels of *ERG16*, *CDR*, and *MDR1* correlate with
increases in azole resistance in *Candida albicans* isolates from a patient infected with
human immunodeficiency virus. *Antimicrob. Agents Chemother.*, **41**, 1482–1487.

75. Marr,K.A., Lyons,C.N., Rustad,T.R., Bowden,R.A. and White,T.C. (1998) Rapid, transient
fluconazole resistance in *Candida albicans* is associated with increased mRNA levels of
CDR. Antimicrob. Agents Chemother., **42**, 2584–2589.

76. Marr,K.A., Lyons,C.N., Ha,K., Rustad,T.R. and White,T.C. (2001) Inducible Azole
Resistance Associated with a Heterogeneous Phenotype in *Candida albicans*. *Antimicrobial*
Agents and Chemotherapy, **45**, 52–59.

77. Manoharlal,R., Gaur,N.A., Panwar,S.L., Morschhäuser,J. and Prasad,R. (2008)
Transcriptional activation and increased mRNA stability contribute to overexpression of
CDR1 in azole-resistant *Candida albicans*. *Antimicrob. Agents Chemother.*, **52**, 1481–1492.

78. Niimi,M., Niimi,K., Takano,Y., Holmes,A.R., Fischer,F.J., Uehara,Y. and Cannon,R.D.
(2004) Regulated overexpression of *CDR1* in *Candida albicans* confers multidrug
resistance. *J. Antimicrob. Chemother.*, **54**, 999–1006.

79. Chen,Y., Mallick,J., Maqnas,A., Sun,Y., Choudhury,B.I., Côte,P., Yan,L., Ni,T.-J.-H., Li,Y.,
Zhang,D., *et al.* (2018) Chemogenomic Profiling of the Fungal Pathogen *Candida albicans*.
Antimicrob. Agents Chemother., **62**, e02365-17.

80. MacAlpine,J., Daniel-Ivad,M., Liu,Z., Yano,J., Revie,N.M., Todd,R.T., Stogios,P.J.,
Sanchez,H., O'Meara,T.R., Tompkins,T.A., *et al.* (2021) A small molecule produced by
Lactobacillus species blocks *Candida albicans* filamentation by inhibiting a DYRK1-family
kinase. *Nat. Commun.*, **12**, 6151.

81. Sharma,J., Rosiana,S., Razzaq,I. and Shapiro,R.S. (2019) Linking Cellular Morphogenesis
with Antifungal Treatment and Susceptibility in *Candida* Pathogens. *J Fungi*, **5**, 17.

82. Kamai,Y., Kubota,M., Kamai,Y., Hosokawa,T., Fukuoka,T. and Filler,S.G. (2002)
Contribution of *Candida albicans ALS1* to the pathogenesis of experimental oropharyngeal
candidiasis. *Infect. Immun.*, **70**, 5256–5258.

83. Fu,Y., Ibrahim,A.S., Sheppard,D.C., Chen,Y.-C., French,S.W., Cutler,J.E., Filler,S.G. and
Edwards,J.E.,Jr (2002) *Candida albicans* Als1p: an adhesin that is a downstream effector
of the *EFG1* filamentation pathway. *Mol. Microbiol.*, **44**, 61–72.

84. Alberti-Segui,C., Morales,A.J., Xing,H., Kessler,M.M., Willins,D.A., Weinstock,K.G.,
Cottarel,G., Fechtel,K. and Rogers,B. (2004) Identification of potential cell-surface proteins
in *Candida albicans* and investigation of the role of a putative cell-surface glycosidase in
adhesion and virulence. *Yeast*, **21**, 285–302.

85. Zhao,X., Oh,S.-H., Cheng,G., Green,C.B., Nuessen,J.A., Yeater,K., Leng,R.P.,
Brown,A.J.P. and Hoyer,L.L. (2004) *ALS3* and *ALS8* represent a single locus that encodes
a *Candida albicans* adhesin; functional comparisons between Als3p and Als1p.
Microbiology, **150**, 2415–2428.

86. Nobile,C.J., Schneider,H.A., Nett,J.E., Sheppard,D.C., Filler,S.G., Andes,D.R. and
Mitchell,A.P. (2008) Complementary adhesin function in *C. albicans* biofilm formation. *Curr.*
Biol., **18**, 1017–1024.

87. Flores-Mireles,A.L., Walker,J.N., Bauman,T.M., Potretzke,A.M., Schreiber,H.L.,4th,
Park,A.M., Pinkner,J.S., Caparon,M.G., Hultgren,S.J. and Desai,A. (2016) Fibrinogen
release and deposition on urinary catheters placed during urological procedures. *J. Urol.*,
196, 416–421.

88. La Bella,A.A., Andersen,M.J., Gervais,N.C., Molina,J.J., Molesan,A., Stuckey,P.V.,
Wensing,L., Shapiro,R.S., Santiago-Tirado,F.H. and Flores-Mireles,A.L. (2021)
Catheterized bladder environment induces *Candida albicans* hyphal formation and
promotes colonization and persistence through Als1. *bioRxiv*, 10.1101/2021.06.01.446547.

89. Andersen,M.J., Fong,C., La Bella,A.A., Molina,J.J., Molesan,A., Champion,M.M., Howell,C.
and Flores-Mireles,A.L. (2022) Inhibiting host-protein deposition on urinary catheters
reduces associated urinary tract infections. *eLife*, **11**, e75798*.

90. Hoyer,L.L., Green,C.B., Oh,S.-H. and Zhao,X. (2008) Discovering the secrets of the
Candida albicans agglutinin-like sequence (*ALS*) gene family--a sticky pursuit. *Med. Mycol.*,
46, 1–15.

91. Hoyer,L.L. and Cota,E. (2016) *Candida albicans* agglutinin-like sequence (Als) family
vignettes: a review of Als protein structure and function. *Front. Microbiol.*, **7**, 280.

92. Coste,A.T., Karababa,M., Ischer,F., Bille,J. and Sanglard,D. (2004) *TAC1*, transcriptional
activator of *CDR* genes, is a new transcription factor involved in the regulation of *Candida*
albicans ABC transporters *CDR1* and *CDR2*. *Eukaryot. Cell*, **3**, 1639–1652.

93. Ford,C.B., Funt,J.M., Abbey,D., Issi,L., Guiducci,C., Martinez,D.A., Delorey,T., Li,B.Y.,
White,T.C., Cuomo,C., *et al.* (2015) The evolution of drug resistance in clinical isolates of
Candida albicans. *eLife*, **4**, e00662.

94. Selmecki,A., Forche,A. and Berman,J. (2006) Aneuploidy and isochromosome formation in
drug-resistant *Candida albicans*. *Science*, **313**, 367–370.

95. Li,X., Yang,F., Li,D., Zhou,M., Wang,X., Xu,Q., Zhang,Y., Yan,L. and Jiang,Y. (2015)
Trisomy of chromosome R confers resistance to triazoles in *Candida albicans*. *Med. Mycol.*,
53, 302–309.

96. Ene,I.V., Farrer,R.A., Hirakawa,M.P., Agwamba,K., Cuomo,C.A. and Bennett,R.J. (2018)
Global analysis of mutations driving microevolution of a heterozygous diploid fungal
pathogen. *Proc. Natl. Acad. Sci. U. S. A.*, **115**, 8688–8697.

97. Forche,A., Magee,P.T., Selmecki,A., Berman,J. and May,G. (2009) Evolution in *Candida*
albicans populations during a single passage through a mouse host. *Genetics*, **182**, 799–
811.

98. Anderson,M.Z., Saha,A., Haseeb,A. and Bennett,R.J. (2017) A chromosome 4 trisomy
contributes to increased fluconazole resistance in a clinical isolate of *Candida albicans*.
Microbiology, **163**, 856–865.

99. Forche,A., Solis,N.V., Swidergall,M., Thomas,R., Guyer,A., Beach,A., Cromie,G.A., Le,G.T.,
Lowell,E., Pavelka,N., *et al.* (2019) Selection of *Candida albicans* trisomy during
oropharyngeal infection results in a commensal-like phenotype. *PLoS Genet.*, **15**,
e1008137.

100. Yang,F., Todd,R.T., Selmecki,A., Jiang,Y.-Y., Cao,Y.-B. and Berman,J. (2021) The fitness
costs and benefits of trisomy of each *Candida albicans* chromosome. *Genetics*, **218**,
iyab056.

101. Yang,F., Teoh,F., Tan,A.S.M., Cao,Y., Pavelka,N. and Berman,J. (2019) Aneuploidy
enables cross-adaptation to unrelated drugs. *Mol. Biol. Evol.*, **36**, 1768-1782.

102. Zhang,X.-H., Tee,L.Y., Wang,X.-G., Huang,Q.-S. and Yang,S.-H. (2015) Off-target Effects
in CRISPR/Cas9-mediated genome engineering. *Mol. Ther. Nucleic Acids*, **4**, e264.

103. Chavez,A., Tuttle,M., Pruitt,B.W., Ewen-Campen,B., Chari,R., Ter-Ovanesyan,D.,
Haque,S.J., Cecchi,R.J., Kowal,E.J.K., Buchthal,J., *et al.* (2016) Comparison of Cas9
activators in multiple species. *Nature Methods*, **13**, 563–567.

104. Cámara,E., Lenitz,I. and Nygård,Y. (2020) A CRISPR activation and interference toolkit for
industrial *Saccharomyces cerevisiae* strain KE6-12. *Sci. Rep.*, **10**, 14605.

105. Radzisheuskaya,A., Shlyueva,D., Müller,I. and Helin,K. (2016) Optimizing sgRNA position
markedly improves the efficiency of CRISPR/dCas9-mediated transcriptional repression.
Nucleic Acids Res., **44**, e141.

106. Hu,J.H., Miller,S.M., Geurts,M.H., Tang,W., Chen,L., Sun,N., Zeina,C.M., Gao,X.,
Rees,H.A., Lin,Z., *et al.* (2018) Evolved Cas9 variants with broad PAM compatibility and
high DNA specificity. *Nature*, **556**, 57–63.

107. Ciurkot,K., Gorochowski,T.E., Roubos,J.A. and Verwaal,R. (2021) Efficient multiplexed
gene regulation in *Saccharomyces cerevisiae* using dCas12a. *Nucleic Acids Res.*, **49**,
7775–7790.

108. Gilbert,L.A., Horlbeck,M.A., Adamson,B., Villalta,J.E., Chen,Y., Whitehead,E.H.,
Guimaraes,C., Panning,B., Ploegh,H.L., Bassik,M.C., *et al.* (2014) Genome-scale CRISPR-
mediated control of gene repression and activation. *Cell*, **159**, 647–661.

109. Horlbeck,M.A., Gilbert,L.A., Villalta,J.E., Adamson,B., Pak,R.A., Chen,Y., Fields,A.P.,
Park,C.Y., Corn,J.E., Kampmann,M., *et al.* (2016) Compact and highly active next-
generation libraries for CRISPR-mediated gene repression and activation. *eLife*, **5**, e19760.

110. Konermann,S., Brigham,M.D., Trevino,A.E., Joung,J., Abudayyeh,O.O., Barcena,C.,
Hsu,P.D., Habib,N., Gootenberg,J.S., Nishimasu,H., *et al.* (2015) Genome-scale
transcriptional activation by an engineered CRISPR-Cas9 complex. *Nature*, **517**, 583–588.

111. Joung,J., Engreitz,J.M., Konermann,S., Abudayyeh,O.O., Verdine,V.K., Aguet,F.,
Gootenberg,J.S., Sanjana,N.E., Wright,J.B., Fulco,C.P., *et al.* (2017) Genome-scale
activation screen identifies a lncRNA locus regulating a gene neighbourhood. *Nature*, **548**,
343–346.

112. Yang,J., Rajan,S.S., Friedrich,M.J., Lan,G., Zou,X., Ponstingl,H., Garyfallos,D.A., Liu,P.,

Bradley,A. and Metzakopian,E. (2019) Genome-scale CRISPRa screen identifies novel
factors for cellular reprogramming. *Stem Cell Reports*, **12**, 757–771.

113. Jost,M., Chen,Y., Gilbert,L.A., Horlbeck,M.A., Krenning,L., Menchon,G., Rai,A., Cho,M.Y.,
Stern,J.J., Prota,A.E., *et al.* (2017) Combined CRISPRi/a-based chemical genetic screens
reveal that Rigosertib is a microtubule-destabilizing agent. *Mol. Cell*, **68**, 210–223.

114. Roemer,T., Jiang,B., Davison,J., Ketela,T., Veillette,K., Breton,A., Tandia,F., Linteau,A.,
Sillaots,S., Marta,C., *et al.* (2003) Large-scale essential gene identification in *Candida
albicans* and applications to antifungal drug discovery. *Mol. Microbiol.*, **50**, 167–181.

115. Noble,S.M., French,S., Kohn,L.A., Chen,V. and Johnson,A.D. (2010) Systematic screens
of a *Candida albicans* homozygous deletion library decouple morphogenetic switching and
pathogenicity. *Nat. Genet.*, **42**, 590–598.

116. Homann,O.R., Dea,J., Noble,S.M. and Johnson,A.D. (2009) A phenotypic profile of the
Candida albicans regulatory network. *PLoS Genet.*, **5**, e1000783.

117. Blankenship,J.R., Fanning,S., Hamaker,J.J. and Mitchell,A.P. (2010) An extensive circuitry
for cell wall regulation in *Candida albicans*. *PLoS Pathogens*, **6**, e1000752.

118. Gervais,N.C., Halder,V. and Shapiro,R.S. (2021) A data library of *Candida albicans*
functional genomic screens. *FEMS Yeast Res.*, **21**, foab060.

119. Lohse,M.B., Ene,I.V., Craik,V.B., Hernday,A.D., Mancera,E., Morschhäuser,J.,
Bennett,R.J. and Johnson,A.D. (2016) Systematic genetic screen for transcriptional
regulators of the *Candida albicans* white-opaque switch. *Genetics*, **203**, 1679–1692.

120. Hossain,S., Veri,A.O. and Cowen,L.E. (2020) The proteasome governs fungal
morphogenesis via functional connections with Hsp90 and cAMP-protein kinase A
signaling. *mBio*, **11**, e00290-20.

121. O'Meara,T.R., Veri,A.O., Ketela,T., Jiang,B., Roemer,T. and Cowen,L.E. (2015) Global
analysis of fungal morphology exposes mechanisms of host cell escape. *Nat. Commun.*, **6**,
6741.

122. Ryan,O., Shapiro,R.S., Kurat,C.F., Mayhew,D., Baryshnikova,A., Chin,B., Lin,Z.-Y.,
Cox,M.J., Vizeacoumar,F., Cheung,D., *et al.* (2012) Global gene deletion analysis exploring
yeast filamentous growth. *Science*, **337**, 1353–1356.

123. Lee,J.A., Robbins,N., Xie,J.L., Ketela,T. and Cowen,L.E. (2016) Functional genomic
analysis of *Candida albicans* adherence reveals a key role for the Arp2/3 complex in cell
wall remodelling and biofilm formation. *PLoS Genet.*, **12**, e1006452.

124. Nobile,C.J., Fox,E.P., Nett,J.E., Sorrells,T.R., Mitrovich,Q.M., Hernday,A.D., Tuch,B.B.,
Andes,D.R. and Johnson,A.D. (2012) A recently evolved transcriptional network controls
biofilm development in *Candida albicans*. *Cell*, **148**, 126–138.

125. Motaung,T.E., Ells,R., Pohl,C.H., Albertyn,J. and Tsilo,T.J. (2017) Genome-wide functional
analysis in *Candida albicans*. *Virulence*, **8**, 1563–1579.

126. Caplan,T., Polvi,E.J., Xie,J.L., Buckhalter,S., Leach,M.D., Robbins,N. and Cowen,L.E.
(2018) Functional genomic screening reveals core modulators of echinocandin stress

responses in *Candida albicans*. *Cell Rep.*, **23**, 2292–2298.

127. Mount,H.O., Revie,N.M., Todd,R.T., Anstett,K., Collins,C., Costanzo,M., Boone,C., Robbins,N., Selmecki,A. and Cowen,L.E. (2018) Global analysis of genetic circuitry and adaptive mechanisms enabling resistance to the azole antifungal drugs. *PLoS Genet.*, **14**, e1007319.

128. Fu,C., Zhang,X., Veri,A.O., Iyer,K.R., Lash,E., Xue,A., Yan,H., Revie,N.M., Wong,C., Lin,Z.-Y., *et al.* (2021) Leveraging machine learning essentiality predictions and chemogenomic interactions to identify antifungal targets. *Nat. Commun.*, **12**, 6497.

129. Jaitly,P., Legrand,M., Das,A., Patel,T., Chauvel,M., Maufrais,C., d'Enfert,C. and Sanyal,K. (2022) A phylogenetically-restricted essential cell cycle progression factor in the human pathogen *Candida albicans*. *Nat. Commun.*, **13**, 4256.

130. Delarze,E., Brandt,L., Trachsel,E., Patxot,M., Pralong,C., Maranzano,F., Chauvel,M., Legrand,M., Znaidi,S., Bougnoux,M.-E., *et al.* (2020) Identification and characterization of mediators of fluconazole tolerance in *Candida albicans*. *Front. Microbiol.*, **11**, 591140.

131. Cabral,V., Znaidi,S., Walker,L.A., Martin-Yken,H., Dague,E., Legrand,M., Lee,K., Chauvel,M., Firon,A., Rossignol,T., *et al.* (2014) Targeted changes of the cell wall proteome influence *Candida albicans* ability to form single- and multi-strain biofilms. *PLoS Pathog.*, **10**, e1004542.

132. Liang,S.-H. and Bennett,R.J. (2019) The Impact of gene dosage and heterozygosity on the diploid pathobiont *Candida albicans*. *J Fungi*, **6**, 10.

133. Momen-Roknabadi,A., Oikonomou,P., Zegans,M. and Tavazoie,S. (2020) An inducible CRISPR interference library for genetic interrogation of *Saccharomyces cerevisiae* biology. *Commun Biol*, **3**, 723.

134. Sanjana,N.E. (2017) Genome-scale CRISPR pooled screens. *Anal. Biochem.*, **532**, 95–99.

135. Muñoz,I.V., Sarrocco,S., Malfatti,L., Baroncelli,R. and Vannacci,G. (2019) CRISPR-Cas for fungal genome editing: a new tool for the management of plant diseases. *Front. Plant Sci.*, **10**, 135.

136. Ouedraogo,J.-P. and Tsang,A. (2020) CRISPR_Cas systems for fungal research. *Fungal Biol. Rev.*, **34**, 189–201.

137. Cai,P., Gao,J. and Zhou,Y. (2019) CRISPR-mediated genome editing in non-conventional yeasts for biotechnological applications. *Microb. Cell Fact.*, **18**, 63.

138. Shan,L., Dai,Z. and Wang,Q. (2021) Advances and opportunities of CRISPR/Cas technology in bioengineering non-conventional yeasts. *Front Bioeng Biotechnol*, **9**, 765396.

139. Schuster,M. and Kahmann,R. (2019) CRISPR-Cas9 genome editing approaches in filamentous fungi and oomycetes. *Fungal Genet. Biol.*, **130**, 43–53.

140. Zoppo,M., Luca,M.D., Villarreal,S.N., Poma,N., Barrasa,M.I., Bottai,D., Vyas,V.K. and Tavanti,A. (2019) A CRISPR/Cas9-based strategy to simultaneously inactivate the entire *ALS* gene family in *Candida orthopsilosis*. *Future Microbiol.*, **14**, 1383–1396.

141. Maroc,L. and Fairhead,C. (2019) A new inducible CRISPR-Cas9 system useful for genome
editing and study of double-strand break repair in *Candida glabrata*. *Yeast*, **36**, 723–731.

142. Enkler,L., Richer,D., Marchand,A.L., Ferrandon,D. and Jossinet,F. (2016) Genome
engineering in the yeast pathogen *Candida glabrata* using the CRISPR-Cas9 system. *Sci.*
Rep., **6**, 35766.

143. Santana,D.J. and O'Meara,T.R. Forward and reverse genetic dissection of morphogenesis
identifies filament-competent *Candida auris* strains. *Nat. Comm.,* **12**, 7197.

144. Ennis,C.L., Hernday,A.D. and Nobile,C.J. (2021) A markerless CRISPR-mediated system
for genome editing in *Candida auris* reveals a conserved role for Cas5 in the caspofungin
response. *Microbiol Spectr*, **9**, e0182021.

145. Lombardi,L., Oliveira-Pacheco,J. and Butler,G. (2019) Plasmid-based CRISPR-Cas9 gene
editing in multiple *Candida* species. *mSphere*, **4**, e00125-19.

146. Lombardi,L., Turner,S.A., Zhao,F. and Butler,G. (2017) Gene editing in clinical isolates of
Candida parapsilosis using CRISPR/Cas9. *Sci. Rep.*, **7**, 8051.

147. Wang,P. (2018) Two distinct approaches for CRISPR-Cas9-mediated gene editing in
Cryptococcus neoformans and related species. *mSphere*, **3**.

148. Huang,M.Y., Joshi,M.B., Boucher,M.J., Lee,S., Loza,L.C., Gaylord,E.A., Doering,T.L. and
Madhani,H.D. (2021) Short homology-directed repair using optimized Cas9 in the pathogen
Cryptococcus neoformans enables rapid gene deletion and tagging. *Genetics*, **220**,
iyab180.

149. Fan,Y. and Lin,X. (2018) Multiple applications of a transient CRISPR-Cas9 coupled with
electroporation (TRACE) system in the *Cryptococcus neoformans* species complex.
Genetics, **208**, 1357–1372.

150. Fuller,K.K., Chen,S., Loros,J.J. and Dunlap,J.C. (2015) Development of the CRISPR/Cas9
system for targeted gene disruption in *Aspergillus fumigatus*. *Eukaryot. Cell*, **14**, 1073–
1080.

151. Al Abdallah,Q., Souza,A.C.O., Martin-Vicente,A., Ge,W. and Fortwendel,J.R. (2018)
Whole-genome sequencing reveals highly specific gene targeting by *in vitro* assembled
Cas9-ribonucleoprotein complexes in *Aspergillus fumigatus*. *Fungal Biol Biotechnol*, **5**, 11.

152. van Rhijn,N., Furukawa,T., Zhao,C., McCann,B.L., Bignell,E. and Bromley,M.J. (2020)
Development of a marker-free mutagenesis system using CRISPR-Cas9 in the pathogenic
mould *Aspergillus fumigatus*. *Fungal Genet. Biol.*, **145**, 103479.

153. Zhang,C., Meng,X., Wei,X. and Lu,L. (2016) Highly efficient CRISPR mutagenesis by
microhomology-mediated end joining in *Aspergillus fumigatus*. *Fungal Genet. Biol.*, **86**, 47–
57.

154. Arnoldo,A., Kittanakom,S., Heisler,L.E., Mak,A.B., Shukalyuk,A.I., Torti,D., Moffat,J.,
Giaever,G. and Nislow,C. (2014) A genome scale overexpression screen to reveal drug
activity in human cells. *Genome Med.*, **6**, 32.

155. Zhao,X.-Q., Zhang,X.-Y., Zhang,F., Zhang,R., Jiang,B.-J. and Bai,F.-W. (2018) Metabolic

Engineering of Fungal Strains for Efficient Production of Cellulolytic Enzymes. In Fang,X., Qu,Y. (eds), *Fungal Cellulolytic Enzymes: Microbial Production and Application*. Springer Singapore, Singapore, pp. 27–41.

156. Donohoue,P.D., Barrangou,R. and May,A.P. (2018) Advances in industrial biotechnology using CRISPR-Cas systems. *Trends Biotechnol.*, **36**, 134–146.

157. Jiang,C., Lv,G., Tu,Y., Cheng,X., Duan,Y., Zeng,B. and He,B. (2021) Applications of CRISPR/Cas9 in the synthesis of secondary metabolites in filamentous fungi. *Front. Microbiol.*, **12**, 638096.

158. Nødvig,C.S., Nielsen,J.B., Kogle,M.E. and Mortensen,U.H. (2015) A CRISPR-Cas9 system for genetic engineering of filamentous fungi. *PLoS One*, **10**, e0133085.

159. Wilson,F.M. and Harrison,R.J. (2021) CRISPR/Cas9 mediated editing of the Quorn fungus *Fusarium venenatum* A3/5 by transient expression of Cas9 and sgRNAs targeting endogenous marker gene *PKS12*. *Fungal Biol Biotechnol*, **8**, 15.

160. Kwon,M.J., Schütze,T., Spohner,S., Haefner,S. and Meyer,V. (2019) Practical guidance for the implementation of the CRISPR genome editing tool in filamentous fungi. *Fungal Biol Biotechnol*, **6**, 15.

161. Roux,I., Woodcraft,C., Hu,J., Wolters,R., Gilchrist,C.L.M. and Chooi,Y.-H. (2020) CRISPR-mediated activation of biosynthetic gene clusters for bioactive molecule discovery in filamentous fungi. *ACS Synth. Biol.*, **9**, 1843–1854.

162. Mózsik,L., Hoekzema,M., de Kok,N.A.W., Bovenberg,R.A.L., Nygård,Y. and Driessen,A.J.M. (2021) CRISPR-based transcriptional activation tool for silent genes in filamentous fungi. *Sci. Rep.*, **11**, 1118.

163. Gargouri,B., Mhiri,N., Karray,F., Aloui,F. and Sayadi,S. (2015) Isolation and characterization of hydrocarbon-degrading yeast strains from petroleum contaminated industrial wastewater. *Biomed Res. Int.*, **2015**, 929424.

164. Kieliszek,M., Kot,A.M., Bzducha-Wróbel,A., BŁażejak,S., Gientka,I. and Kurcz,A. (2017) Biotechnological use of *Candida* yeasts in the food industry: A review. *Fungal Biology Reviews*, **31**, 185–198.

165. Payen,C. and Thompson,D. (2019) The renaissance of yeasts as microbial factories in the modern age of biomanufacturing. *Yeast*, **36**, 685–700.

166. Joo,H.-S., Ndegwa,P.M., Shoda,M. and Phae,C.-G. (2008) Bioremediation of oil-contaminated soil using *Candida catenulata* and food waste. *Environmental Pollution*, **156**, 891–896.

167. García-Béjar,B., Arévalo-Villena,M., Guisantes-Batan,E., Rodríguez-Flores,J. and Briones,A. (2020) Study of the bioremediatory capacity of wild yeasts. *Sci. Rep.*, **10**, 11265.

168. Peng,D. and Tarleton,R. (2015) EuPaGDT: a web tool tailored to design CRISPR guide RNAs for eukaryotic pathogens. *Microb Genom*, **1**, e000033.

169. Engler,C., Kandzia,R. and Marillonnet,S. (2008) A one pot, one step, precision cloning method with high throughput capability. *PLoS One*, **3**, e3647.

170. Halder,V., Porter,C.B.M., Chavez,A. and Shapiro,R.S. (2019) Design, execution, and
analysis of CRISPR–Cas9-based deletions and genetic interaction networks in the fungal
pathogen *Candida albicans*. *Nat. Protoc.*, **14**, 955–975.

171. Razzaq,I., Berg,M.D., Jiang,Y., Genereaux,J., Uthayakumar,D., Kim,G.H., Agyare-
Tabbi,M., Halder,V., Brandl,C.J., Lajoie,P., *et al.* (2021) The SAGA and NuA4 component
Tra1 regulates *Candida albicans* drug resistance and pathogenesis. *Genetics*, **219**,
iyab131.

172. Schmittgen,T.D. and Livak,K.J. (2008) Analyzing real-time PCR data by the comparative
C(T) method. *Nat. Protoc.*, **3**, 1101–1108.

1130

1131

Figure 1. Design of a CRISPRa system for *C. albicans*. (a) The single plasmid system contains a catalytically inactive ("dead") dCas9 protein, paired with the VPR (VP64-P65-Rta) complex of activator domains to cause upregulation at a target locus. The single-guide RNA (sgRNA) region can be subjected to Golden Gate cloning with a custom N20 sequence in order to target the dCas9-VPR complex to a desired region of DNA through complementary base-pairing. The PacI enzyme is used to linearize the plasmid at the *NEUT5L* locus, which allows the plasmid to be integrated into the cell's chromosome and remain in the cell following cell division. Panel created with BioRender.com. **(b)** Once the sgRNA recognizes and binds to a target sequence of DNA, the dCas9-VPR complex will cause the recruitment of transcription initiation complex factors to the adjacent genetic material, thus increasing gene expression from that region. Created with BioRender.com. **(c)** Wild-type *C. albicans* cells and cells containing a non-targeting CRISPRa plasmid were grown in YPD at 30°C to compare fitness under standard laboratory conditions. Growth was recorded via optical density at 600nm at 15 min intervals over the course of 24h. Data points show the mean and standard deviation at each time point (n = 32).

Figure 2. Determining optimal guide-design principles for CRISPRa overexpression in *C. albicans*. CRISPRa strains were created using six different sgRNA molecules designed to target different positions in the promoter region of each of **(a)** *KSP1*, **(b)** *ALS3*, and **(c)** *FGR28* upstream of either the start codon or the predicted dominant transcriptional start site (TSS). Fold-change in the expression of the target gene relative to the housekeeping gene *ACT1* in the experimental strains and the non-targeting CRISPRa control strain was measured with RT-qPCR and compared. Data points represent the mean fold-difference and standard deviation in expression of the respective target gene in the CRISPRa strains compared to the non-targeting CRISPRa control (n = 3).

Figure 3. Using CRISPRa to validate antifungal drug resistance phenotypes. (a) Three *C. albicans* CRISPRa strains were created targeting the drug transporter gene *CDR1* for overexpression. Fold-change in the expression of *CDR1* relative to the housekeeping gene *ACT1* in the three experimental strains and the non-targeting CRISPRa control strain was measured with RT-qPCR. All three experimental strains are represented by the position of their sgRNA in the promoter region upstream of the proposed dominant *CDR1* transcriptional start site. Data represent the mean fold-difference and standard deviation in expression of *CDR1* in the CRISPRa strains compared to the non-targeting CRISPRa control (n = 3). **(b)** The fitness of all three CRISPRa strains targeting *CDR1* for overexpression was measured in the presence of the antifungal drug fluconazole with a minimum inhibitory concentration assay. The OD600 value of each strain in each concentration of fluconazole was measured to represent growth in differing conditions, where OD600 values approaching 1.0 reflect strong fitness. **(c)** Fold-change in the expression of *YAK1* relative to the housekeeping gene *ACT1* in the experimental strain and the non-targeting CRISPRa control strain was measured with RT-qPCR. Data represent the mean fold-difference and standard deviation in expression of *YAK1* in the CRISPRa strains compared to the non-targeting CRISPRa control (n = 3). **(d)** The fitness of the CRISPRa strain targeting *YAK1* for overexpression was measured in the presence of the antifungal drug amphotericin B with a minimum inhibitory concentration assay. The OD600 value of each strain in each concentration of amphotericin B was measured to represent growth in differing conditions, where OD600 values approaching 1.0 reflect strong fitness.

Figure 4. Validation of CRISPRa overexpression and target-specificity through targeting fungal pathogenesis traits. (a) Two *C. albicans* CRISPRa strains targeting different positions in the promoter region of the adhesin protein gene *ALS1* were created. Fold-change in the expression of *ALS1* relative to the housekeeping gene *ACT1* in both experimental strains and the non-targeting CRISPRa control strain was measured with RT-qPCR. Experimental strains are represented by the position of their sgRNA in the promoter region upstream of the proposed dominant *ALS1* transcriptional start site. Data points represent the mean fold-difference and standard deviation in expression of *ALS1* in the CRISPRa strains compared to the non-targeting CRISPRa control (n = 3). **(b)** Two CRISPRa strains targeting *ALS1* and a non-targeting CRISPRa control strain were grown in the biofilm-inducing media RPMI and left to grow statically at 37°C for 24h to allow biofilms to form. The robustness of biofilms formed by all strains was elucidated through an XTT reduction assay and measuring of the resulting OD490 values before normalizing final values to non-biofilm-forming planktonic cells. Data points represent the mean and the standard error of the mean (n = 72). Differences between groups were tested for significance using Dunnett's multiple comparison test. ****, p-value <0.0001. ***, p-value <0.001. **(c)** A *C. albicans* CRISPRa strain overexpressing *ALS1* ~6-fold and a non-targeting CRISPRa strain were grown in human urine supplemented with fibrinogen and BSA at 37°C for 48h to simulate a catheterized bladder environment. The robustness of biofilms formed by both strains was measured by reading the OD595 value following a crystal violet assay. Data points are combined from three independent experiments where n = 32 for the non-targeting CRISPRa control and n = 48 for the CRISPRa *ALS1* overexpression strain. Differences between groups were tested for significance using the Mann-Whitney U test. ****, p-value <0.0001. **(d)** Global gene expression of a CRISPRa strain overexpressing *ALS1* ~25-fold, as well as a non-targeting CRISPRa control strain, was measured using RNA-Seq. Differential gene expression analysis showed that *ALS1* had a log2-fold change of ~3.3 in the *ALS1* overexpression strain compared to the CRISPRa control strain, with an adjusted p-value of 3.07E-142, and showed no additional transcripts that had differential expression with an adjusted p-value of less than 0.03.