1 Genomics of Natural Populations: Gene Conversion Events Reveal Selected Genes within the
2 Inversions of *Drosophila pseudoobscura*
3

4 Stephen W. Schaeffer*, Stephen Richards[1], and Zachary L. Fuller[2]
5
6 *Department of Biology
7 The Pennsylvania State University
8 208 Erwin W. Mueller Laboratory
9 University Park, PA 16802-5301
10
11
12
13 [1]Current address:
14 Human Genome Sequencing Center, Baylor College of Medicine, One Baylor Plaza, Houston,
15 TX 77030
16
17 [2]Current address:
18 23andMe, Inc., Sunnyvale, CA 94086
19

20 Data Availability: Sequence data in FASTA format and aligned NEXUS files are available at
21 https://scholarsphere.psu.edu/resources/720f454e-ceb2-4139-a3a9-c59a5b7fdfaf . Additional
22 data and computer code are available at https://github.com/SWSchaeffer/InversionGC
23
24

30   Corresponding Author: Stephen W. Schaeffer
31   Department of Biology
32   The Pennsylvania State University
33   208 Erwin W. Mueller Laboratory
34   University Park, PA 16802-5301
35   (814) 865-3269
36   sws4@psu.edu
37

Abstract

When adaptive phenotypic variation or QTLs map within an inverted segment of a chromosome, researchers often despair because it is thought that the suppression of crossing over will prevent the discovery of selective target genes through fine scale genetic mapping. If an inversion polymorphism is old enough, then the accumulation of gene conversion tracts offers the promise that QTLs or selected loci within inversions can be identified. This study uses the inversion polymorphism of *Drosophila pseudoobscura* as a model system to determine if gene conversion analysis can be a useful tool for mapping adaptive loci within inversions. *D. pseudoobscura* has over 30 different chromosomal arrangements on the third chromosome (Muller C) in natural populations and their frequencies vary with changes in environmental habitats. Statistical tests of five *D. pseudoobscura* gene arrangements identified outlier genes within inverted regions based on local clusters of fixed SNP differences. These outlier genes also had potentially selectable variation, either fixed amino acid differences or differential expression patterns among arrangements. Here, we use genome sequences of the inverted third chromosome (Muller C) to infer 98,443 gene conversion tracts for a total coverage of 142 Mb. We tested whether the frequency of gene conversion tracts in outlier versus non-outlier genes were equal. Gene conversion tracts occur less frequently in outlier versus non-outlier genes suggesting that selection removes exchanged DNA from the multiple outlier genes. These data support the hypothesis that these inversions captured adaptive alleles and that strong epistatic selection removes recombinants generated through gene conversion. The pattern and organization of genetic diversity in the third chromosome in *D. pseudoobscura* is consistent with the capture of locally adapted combinations of alleles prior to inversion mutation events.

Introduction

Genomic re-sequencing data have provided insights into the mechanisms for the origin, establishment, and maintenance of chromosomal inversions in natural populations. Comparison of sequences between different gene arrangements have allowed the identification of regions of chromosomes where breakage and rejoining events led to the origin of new gene orders (RICHARDS *et al.* 2005; RANZ *et al.* 2007). Ectopic exchange between repeat sequences at breakpoints provide one model for the facilitation of inversion mutations, while staggered cuts that duplicate genes at breakpoints is alternative mechanism observed to generate new gene arrangements.

Once an inversion mutation arises in the population, it is usually lost because its initial frequency (1/2N) is quite low. On the other hand, a new inversion can be established in the population either due to neutral or selective forces. Like point mutations, a selectively neutral inversion can drift to higher frequency through random genetic drift and over time inter-arrangement divergence will be greatest at the breakpoints (NAVARRO *et al.* 2000; GUERRERO *et al.* 2012). There are several possible mechanisms that can explain how an inversion is established by selection. First, direct or position effects (SPERLICH AND PFRIEM 1986) may create positively selectable variation when the breakpoints either disrupt genes, create new hybrid genes, or alter gene expression of boundary genes at or near the lesions. Recent analyses have shown that breakpoints can modify chromatin architecture and disrupt expression of nearby genes in topologically associate domains (WRIGHT AND SCHAEFFER 2022). Second, indirect effects associated with the suppression of recombination can maintain advantageous haplotypes within the inverted region leading to elevated linkage disequilibrium among multiple genes within the inverted segment (KIRKPATRICK AND BARTON 2006). These models assume that the new inversion captures sets of beneficial alleles (CHARLESWORTH AND CHARLESWORTH 1973). Sets of alleles may be beneficial if they are free of deleterious recessive alleles (NEI *et al.* 1967) or are alleles that contribute to local adaptation (KIRKPATRICK AND BARTON 2006). Third, a new gene arrangement may rise to intermediate frequency because one beneficial allele is trapped within the inverted region (MAYNARD SMITH AND HAIGH 1974).

Once established, the new inversion will act as an isolated subpopulation and accumulate genetic differences from its ancestral arrangement (NAVARRO *et al.* 2000). Recombination or genetic flux among different gene arrangements will oppose genetic differentiation keeping sequences homogeneous among inversion types (NAVARRO *et al.* 1997). Genetic flux is initiated by double strand breaks between inverted chromosomes and are either resolved as genetic cross overs or as gene conversion events. Single cross overs within an inverted region of a Drosophila heterokaryotype will generate unbalanced inviable gametes that are selected against leaving only parental gametes (STURTEVANT AND BEADLE 1936). Cross overs are possible if two cross overs occur within the inverted region, but the probability of double cross over events is small unless the size of the inverted segment is large (NAVARRO *et al.* 1997). Gene conversion events, on the other hand, are genetic exchanges at double strand breaks that are resolved as non-cross over events leading to the exchange of small segments of 200 to 300 nucleotides among arrangements without the formation of unbalanced gametes (CHOVNICK 1973; ROZAS AND AGUADÉ 1993; ROZAS AND AGUADÉ 1994; SCHAEFFER AND ANDERSON 2005). Gene conversion events occur uniformly across the inverted segment, but the homogenizing effect is reduced because of the small DNA segment exchanged among chromosomes (NAVARRO *et al.* 1997; MILLER *et al.* 2012; MILLER *et al.* 2016; KORUNES AND NOOR 2019). Overall, crossing over is expected to have a greater homogenizing effect than

gene conversion because larger genomic segments are exchanged, but crossing over is less
likely to occur relative to gene conversion events.
Under a neutral model and an infinite amount of time, maximal nucleotide divergence will
be found at the breakpoints with the central inverted region showing less divergence among
different arrangements (NAVARRO *et al.* 2000; NOOR *et al.* 2007; GUERRERO *et al.* 2012). When
an inversion captures locally adapted alleles, selection opposes the homogenizing effect of
genetic flux leading to local peaks of differentiation (GUERRERO *et al.* 2012). The observation of
peaks of differentiation will become more apparent as time progresses and gene conversion
events homogenize variation among arrangements except at selected loci.
Fuller *et al*. (2017) analyzed complete genome sequences of the third chromosome
inversion polymorphism of *D. pseudoobscura* as a model system to determine mechanisms
responsible for the establishment of inversions in populations. The third chromosome or Muller
C (MULLER 1940) of *D. pseudoobscura* has a wealth of gene arrangements that were generated
through a series of overlapping inversion events (DOBZHANSKY AND STURTEVANT 1938). These
arrangements have been suggested to be targets of selection based on stable clines across
heterogenous environments (DOBZHANSKY 1944; ANDERSON *et al.* 1991), stable altitudinal
gradients (DOBZHANSKY 1948), and seasonal cycling (DOBZHANSKY 1948). Aquadro *et al*.
(1991) used phylogenetic analysis of restriction fragment length polymorphisms to support the
unique origin hypothesis of the *D. pseudoobscura* gene arrangements. Fuller *et al*. (2017)
tested the 2,668 Muller C genes within each of six arrangements for evidence of adaptive
evolution. Loci with significant elevated frequencies of derived nucleotide variants and with
significantly long population specific branch lengths (SHRIVER *et al.* 2004; YI *et al.* 2010) were
identified as outlier genes. In addition, outlier genes that were differentially expressed or had
fixed inversion specific amino acids among arrangements were considered potential targets of
selection.
Five of the six *D. pseudoobscura* arrangements had sufficient sample sizes to detect
multiple outlier loci. These gene arrangements had between 36 and 149 outlier genes that were
distributed within and outside the boundaries of the derived inversion (Figure 1). Outlier genes
had decreased estimates of the recombination parameter rho (ρ) (CHAN *et al.* 2012) compared
to intervening non-outlier loci (see Table 3 in FULLER *et al.* 2017) suggesting that genetic flux is
limiting differentiation at a fine scale. Because genetic flux due to crossing over is expected to
homogenize the central regions of inversions rather than local areas, we hypothesize that gene
conversion is the predominant force homogenizing local regions among arrangements and that
adaptive outlier genes will have fewer detected gene conversion events than non-outlier loci
because natural selection is predicted to remove the gene conversion tracts that bring
maladaptive variants to loci responsible for the increase of inversions.
Direct empirical estimates of gene conversion rates in inverted and colinear regions of *D.*
pseudoobscura have been estimated to be 3.1×10^{-5} and 7.7×10^{-6} converted nucleotides per
base per generation, respectively (KORUNES AND NOOR 2019). In this experiment, Korunes and
Noor (2019) generated genome wide SNP heterozygotes and used computational methods to
infer gene conversion events on four of the five major chromosomal arms. In the cross between
the PP1137 and MSH177 strains, the third chromosome was heterozygous for the ST and PP
arrangements. Ten F_2 offspring were sequenced for this cross and 11 and 23 gene conversion
events were scored within inverted and colinear regions respectively. Identical events were
scored in the cross and that were presumed to occur during mitosis prior to meiosis (KORUNES
AND NOOR 2019). If only unique events are scored, then three and eight gene conversion tracts
were observed in inverted and colinear regions, respectively, which reduces the estimates of

gene conversion rates in inverted and colinear regions to 8.5 x 10^{-6} and 2.4 x 10^{-6},converted

nucleotides per base per generation, respectively. These estimates of gene conversion rates

are two orders of magnitude greater than the mutation rate, which would allow gene conversion

to homogenize genetic diversity among arrangements. Korunes and Noor (2019) show that

gene conversion events can be detected after a single generation, but do not provide insights

about the selective effects of these events. The current study infers gene conversion events

over the accumulated history of the third chromosome inversion polymorphism to determine if

gene conversion events are retained and occur uniformly across inverted regions.

162   A general computational method to detect gene conversion events indirectly was

developed based on the configuration of polymorphic nucleotides (phylogenetic partitions) at

multiple sites across a sample of nucleotide sequences (STEPHENS 1985). This method was

refined to detect gene conversion events in inversion systems when differentiated DNA

segments are exchanged between one arrangement and another (BETRAN *et al.* 1997). Figure

2 shows an example of two detected gene conversion events on Muller C of *D. pseudoobscura*.

The Betran *et al*. (1997) method assumes that the frequency of an informative variant is greater

in the source arrangement than in the recipient. Two probabilities are used to infer gene

conversion tracts from samples of gene arrangement sequence data. The first is the probability

that a SNP is informative of a conversion event based on the frequencies of a variant within the

source and recipient subpopulations or ψ. The second is the probability that a gene conversion

event is extended to an additional nucleotide or φ. This method was applied by Schaeffer and

Anderson (2005) to gene arrangement samples of short aligned sequences (339 to 517 bp)

from seven gene regions of *D. pseudoobscura*. They estimated the mean conversion tract

length to be 205 bp and mean population conversion rate to be 3.4 x 10^{-6}, which is comparable

to the rates inferred by Korunes and Noor (2019). With genome re-sequencing data, we now

use the Bertran *et al*. (1997) method to do a comprehensive analysis of gene conversion tracts

across Muller C of *D. pseudoobscura* to test whether gene conversion events occur uniformly in

genes within the inverted regions. We show here that previously identified outlier genes have

lower coverage from gene conversion events than non-outlier genes providing evidence for

strong epistatic selection maintaining associations of adaptive genes captured by inversion

events.

Materials and Methods

185
Strains. The genome sequencing of 54 strains of *D. pseudoobscura* that carry one of six
different gene arrangements on Muller C have been previously described (FULLER *et al.* 2017).
In brief, isofemale strains were collected from seven localities: Mount St. Helena, CA (MSH),
Santa Cruz Island, CA (SCI); James Reserve, CA (JR); Mather, CA (MA); Kaibab National
Forest, AZ (KB); Davis Mountains, TX (DM), San Pablo Etla, Oaxaca, Mexico (SPE). Crosses
with either the Blade or Lobe third chromosome balancer strains were used to generate
isochromosomal strains (Drosophila Species Stock Center Strain ID 14011-0121.173 is the
Blade balancer for non-AR chromosomes, Genotype: or Bl cv AR/lethal CU; Strain ID 14011-
0121.171 is the Lobe balancer for non-SC chromosomes, Genotype: or L SC/ or + lethal ST;
and Strain ID 14011-0121.172 is an additional strain used in the balancer crosses, Genotype or
pr) (see Figure 1 in DOBZHANSKY AND QUEAL 1938 for the general crossing scheme). The 54
strains used in this study carry one of six Muller C gene arrangements where the names are
based on the locality where the chromosome was first collected (DOBZHANSKY AND STURTEVANT
1938): Arrowhead (AR, n=15); Standard (ST, n=8); Pikes Peak (PP, n=10); Chiricahua (CH,
n=9); Tree Line (TL, n=9); and Cuernavaca (CH,n=3).

Genome Sequences. Each of the 54 strains were sequenced with Illumina HiSeq short
reads (for full details of the sequencing, read mapping, and variant calling, see FULLER *et al.*
2017). Reads were mapped to the *D. pseudoobscura* v3.2 reference sequence. The Arrowhead
MV2-25 reference genome is available at NCBI in BioProject PRJNA18793, the re-sequencing
genomes are available in BioProject PRJNA358242. Two additional Pikes Peak genomes are
available at NCBI as BioSamples SAMN00709017 and SAMN00709004. The set of 54 aligned
sequences for the 19 Mb third chromosome (Muller C) were used in the gene conversion
analysis.

Gene Conversion Analysis. We estimated gene conversion tract coverage in outlier
and non-outlier genes with the following analysis. Coverage refers to how often conversion
tracts occur at a particular nucleotide in a gene region. As gene arrangements age, they
accumulate arrangement specific mutations. These SNPs can be exchanged between
arrangements in heterokaryotypes via crossing over or gene conversion. We used the
approach of Betran *et al*. (1997) to infer gene conversion tracts across Muller C of *D.
pseudoobscura*. This approach compares sets of sequences among different chromosomal
arrangements to identify two or more SNPs in linkage disequilibrium that are found in two or
more arrangements.

The gene conversion events detected by the Betran *et al*. (1997) approach are based on
exchange between gene arrangement heterokaryotypes. The gene conversion analysis requires
that the SNPs use a common gene order and that the distance among SNPs be preserved in
the data set. The seven pairs of *D. pseudoobscura* inversion breakpoints used in this study
divide the third chromosome into 14 regions where genes within the region are in the same
order in all arrangements (Figure 1). While conversion events may occur near breakpoints
(CROWN *et al.* 2018), they are unlikely to straddle the breakpoints.

The Gene Conversion analysis module within DNASP v 6 (ROZAS *et al.* 2017) was used
to infer gene conversion tracts across the 14 cytogenetic regions of Muller C. Each of the 14
regions were subdivided into smaller subregions of 250 kb each. The beginning or end of
subregions were adjusted to accommodate breakpoint positions that resulted in subregions
smaller than 100 kb. The final set of 73 subregions varied in size from 181 to 440 kb with 49
subregions being 250 kb. This was done to accommodate DNASP memory constraints and to
maximize the speed of analysis. All pairs of gene arrangements were compared in each of the

73 subregions with the Gene Conversion analysis in DNASP (15 total comparisons: AR-ST, AR-
PP, AR-CH, AR-TL, AR-CU, ST-PP, ST-CH, ST-TL, ST-CU, PP-CH, PP-TL, PP-CU, CH-TL,
CH-CU, and TL-CU). Each analysis excluded nucleotide sites with alignment gaps and the
output provided lists of conversion tracts within strain, the beginning and end coordinates of the
tract, the tract length excluding gaps, mean value of ψ, number of informatic sites, and a list of
sites with its associated value of ψ. Results of each of the 1,095 pairwise analyses were stored
in separate text files and were also combined into a single file for subsequent summary
analyses.

A gene conversion tract can be detected multiple times in this analysis. There are two
sources of redundancy. First, a gene conversion tract can be inferred from multiple pairwise
comparisons of gene arrangements. For instance, a gene conversion tract in an AR strain might
be inferred from a comparison of AR with CH or with TL. Second, a gene conversion tract can
be inferred in multiple strains of the same gene arrangement. It is assumed that gene
conversion tracts result from unique events and finding the same event in multiple strains is a
result of the tract increasing in frequency within the population (BETRAN *et al.* 1997). Only one
tract from a single strain is used when the conversion event is found in multiple strains. The
effect of removing redundant gene conversion tracts from the data set is a conservative
adjustment that lowers the overall gene conversion tract coverage per site in both outlier and
non-outlier genes.

A histogram of gene conversion coverage relative to outlier and non-outlier genes was
constructed from the set of non-redundant gene conversion tracts (Figure 3). Outlier and Non-
outlier transcripts were mapped to Muller C (Figure 3A). Next, gene conversion tracts were
mapped to Muller C (Figure 3B). Finally, a histogram of gene conversion coverage was
constructed by counting the number of tracts that cover each nucleotide across the
chromosome (Figure 3C).

The mean coverage per nucleotide was estimated for outlier and non-outlier genes from
the gene conversion histogram. The sum of gene conversion tract coverage was estimated
across the exons of each gene and divided by the total number of nucleotides in the gene. The
mean gene conversion coverage was determined separately for outliers and non-outlier genes.

To determine if the donor and recipient conversion events were independent across the
14 syntenic block regions and overall, we permuted the donor and recipient classifications
across the 14 regions. We constrained the permutations so that only non-self-arrangement
events occurred. A total of 10,000 random permutations were used. For each permutation of
the donors and recipients, the number of gene conversion events was re-estimated and
compared with the observed data to determine the probability of obtaining more extreme values
as well as the minimum and maximum values found across all permutations.

Maximum Likelihood Estimates of ϕ and Expected Number of Conversion Tracts.
We used the maximum likelihood approach of Betran *et al*. (1997) to infer ϕ within each of the
14 regions and for each donor-recipient pair using the observed list of conversion tracts for each
pair (See equations 4, 6, and 8). The second derivative of the likelihood equation was used to
estimate the asymptotic variance, which is used to infer a lower and upper bound for tract
length. We did separate analyses for each region and donor-recipient pair because we
assumed that the breakpoints that defined the 14 regions might interfere with the progression of
conversion tracts. This may be a conservative assumption because we define the syntenic
block regions based on the set of 13 breakpoints of gene arrangements used in this study.

Tests for Differences in Gene Conversion Tract Coverage between Outlier and
Non-Outlier Genes. We tested the hypothesis that outlier and non-outlier genes had

equivalent gene conversion coverage using several methods. This assumes that gene
conversion can occur uniformly across the chromosome.

       281   **1.** ***Random Permutation of Gene Class.*** We used a permutation test where we
282   randomly assigned outlier and non-outlier designations to the genes within the
283   arrangement. A total of 10,000 replicates were used to determine the frequency of
284   random outlier assignments that had a mean tract coverage less than the observed
285   value.

286   **2.** ***Random Permutation of Observed Gene Conversion Tracts.*** We randomly
287   assigned the observed gene conversion tracts to Muller C without replacement
288   10,000 times and for each random mapping we re-constructed the conversion tract
289   histogram for the outlier and non-outlier genes. The average coverage in outlier and
290   non-outlier genes was determined for each replicate and the frequency of replicates
291   that had a value less than the observed was estimated.

292   **3.** ***Random Gene Conversion Tracts Based on Maximum Likelihood Parameters.***
293   We used maximum likelihood estimates of region-specific ϕ and the expected
294   number of gene conversion tracts ($E(k_T)$) in the 14 different syntenic block regions to
295   generate a set of conversion tracts. The parameter ϕ is the probability of extending
296   the conversion tract to an additional nucleotide and the process of generating tract
297   lengths is geometric random variable. For each gene conversion tract within a
298   recipient arrangement, we sampled a tract length as a geometric random deviate. If
299   the tract length was greater than the region size, then a new tract length was chosen.
300   Next, a random nucleotide site was chosen from a uniform distribution to begin the
301   gene conversion tract. If the mapped tract spanned two syntenic blocks, a new
302   starting coordinate was chosen. A total of $E(k_T)$ tracts are generated for each region
303   for each of the five donor arrangements, where k_T is the expected number of
304   conversion tracts (See page 95 in BETRAN *et al.* 1997). A total of 10,000 replicates
305   was generated and the mean tract coverage in outlier genes was re-estimated.

Results

  Numbers and Lengths of Conversion Tracts in 14 Syntenic Block Regions. There were 98,443 non-redundant gene conversion tracts observed across Muller C. The mean number of tracts per strain is 1930.2. Sample sizes varied among arrangements, which could bias the number of gene conversion tracts detected. The distribution of gene conversion tracts rejects the null hypothesis that conversion tracts are proportional to the number of strains sampled within each gene arrangement using a Chi-Square Goodness of Fit test (X^2=2281.0, df=4, P<0.0001). The AR, CH, and TL arrangements have more gene conversion tracts than expected while the PP and ST arrangements have fewer tracts than expected (Table 1). Gene conversion tract numbers are not similar in the 14 syntenic block regions in the five different arrangements based on a Chi-Square test for Homogeneity (X^2=366.6, df=52, P<0.0001) (Figure 4). Examination of residuals shows that the AR arrangement has a deficiency of tracts in regions 5 and 9, the CH arrangement has an excess of gene conversion tracts in region 5 and a deficiency of tracts in regions 11 and 14, the PP arrangement has a deficiency of tracts in region 5 and an excess of tracts in region 9, the ST arrangement has deficiencies of tracts in regions 3 and 5 and an excess of tracts in regions 8 and 10, while the TL arrangement has an excess of tracts in region 3 and deficiencies of tracts in regions 5 and 8.

  Median observed gene conversion tract lengths are greatest in the proximal and distal regions of Muller C compared to the central inverted regions (Figure 5). The central areas of the chromosome which are often within inverted regions in heterokaryotypes have an observed median tract length of 112, 160, 97, 127, and 115 bps in the AR, CH. PP, ST, and TL arrangements, respectively. We used a random permutation test to determine if the median gene conversion tract lengths are lower or greater than expected given the distribution of all observed tract lengths. The test was performed by shuffling the list of tracts without replacement, which randomly assigns tracts to the 14 different regions keeping the number of tracts per region the same as the observed. For each of the 10,000 permutation replicates, the regional median was re-estimated and the probability of obtaining a median greater than or less than the observed value within each region was determined. Figure 5 shows the regions that had median gene conversion tract lengths that were significantly less than or greater than expected given the distribution of all tract lengths. Proximal regions 1 and 2 and distal regions 12 through 14 had median tract lengths that were greater than expected in two or more gene arrangements while central regions 4 through 10 had median tract lengths that were less than expected in four or more karyotypes. Regions 3 and 11 had arrangements with median tract lengths less than or greater than expected reflecting boundary regions that the transition between largely collinear or inverted segments of the chromosome.

  We observed gene conversion events between all pairs of arrangements across Muller C ranging between 1,270 conversion events from ST to PP and 9,281 conversion events from CH to AR. Figure 6 shows the polarity of the observed gene conversion events (Donor to Recipient) in the 14 syntenic block regions and across all regions. For all gene conversion events across all regions, two pairs showed a deficiency of events in both directions (CH-PP and ST-TL), one pair showed an excess of events in both directions (PP-TL), and the other seven pairs showed a deficiency in one direction and an excess in the other direction. The test of independence of donor to recipient events for the 14 different regions found subsets of regions with significant deficiencies or excesses of conversion events with a tendency toward what was found for the sum of all events. For instance, if the overall trend was toward a deficiency of events, then most regions with a significant departure tended to also show a deficiency, e.g., the ST to CH transition had a significant overall deficiency (2,195) and nine of 14 regions showed a significant deficiency. For the arrangement pairs likely to form heterozygotes at frequencies of 10% or more in natural populations (CH/ST, CH/PP, AR/ST,

and AR/PP) CH tended to be a donor to AR and ST, ST tended to be a donor to AR, and AR
tended to be a donor to PP.
Maximum Likelihood Estimates of ϕ and Expected Number of Conversion Tracts.
The estimates of ϕ and the expected number of conversion tracts along with other values for all
donor-recipient pairs in each region are in the Supplemental Materials. These values are used
below in tests of gene conversion coverage in outlier genes.
Gene Conversion Tract Coverage in Outlier and Non-Outlier Genes. The 98,443
observed conversion tracts were mapped to Muller C and the mean coverage per nucleotide
was estimated for the exons of each transcript. The expected gene conversion tract coverage
per nucleotide across Muller C is 1.2 per nucleotide for all arrangements except for PP, which
had an expected coverage rate of 0.8 tracts per nucleotide. Thus, at least one conversion tract
should occur in each gene across the chromosome (Table 1).
Figure 7 shows the gene conversion tract coverage for outlier and non-outlier genes
across Muller C in five gene arrangements of *D. pseudoobscura*. Proximal and distal regions of
all arrangements have higher conversion tract coverage than central inverted regions. Outlier
genes tend to have lower levels of coverage than non-outliers. The maximum coverage values
are sample size dependent. Standard, Chiricahua, Pikes Peak, and Tree Line have sample
sizes between 8 and 10 with maximum coverage of 10 tracts/nucleotide. Arrowhead, on the
other hand, has a sample size of 15 with a maximum coverage of 18.
The mean gene conversion tract coverage estimated for the 2,668 genes on Muller C
shows that non-outlier genes have higher mean coverage than outlier genes (Table 2)*.* Gene
conversion tract overage in non-outlier genes matches the expected value for the entire
chromosome, while outlier genes have coverage values < 0.1.
We used three tests to determine if the lower gene conversion coverage is expected by
chance (Table 2). The Random Permutation of Gene Class test found that the observed mean
gene conversion coverage in outlier genes was always less than random permutations in all
gene arrangements. One factor to consider is that we filtered the conversion tracts to remove
redundant tracts. If the redundant tracts were to be included, we would likely see higher
coverage both in outlier and non-outlier genes, but still show a significant difference. The
Random Permutation of Observed Gene Conversion Tracts test showed that the observed
mean coverage in outliers was always less than the replicate permutations. The Random Gene
Conversion Tracts Based on Maximum Likelihood Parameters test found that the observed
mean conversion tract coverage in outlier genes was less than was found with random
conversion tracts. The results of this test should be viewed with caution because the expected
number of true tracts is greater than the observed number of tracts. Thus, it is not surprising
that we observe higher gene conversion tract coverage in outlier genes than in our replicates.
The important result from these three tests is that outlier genes have lower gene conversion
coverage than expected.
394

Discussion

395
The work presented here shows that gene conversion events among arrangements are
readily detectable and serve as a source of genetic flux among the gene arrangements of *D.*
pseudoobscura. The number of observed conversion tracts tends to be uniformly distributed
across Muller C (Figure 4) except for the proximal two syntenic blocks of the chromosome
consistent with theory on genetic flux (NAVARRO *et al.* 1997). Fewer conversion tracts may
result from the initiation of fewer double stranded breaks in the proximal regions. While regions
1 and 2 are closer to the centromere where crossing over is reduced (LINDSLEY AND SANDLER
1977), these regions are in euchromatic DNA, which is unlikely to be affected by the
centromeric effect. More likely is that the Betran *et al*. (1997) method has lower power to detect
gene conversion events in regions 1 and 2. This method relies on informative nucleotide
polymorphisms to detect an exchange between two gene arrangements. Because regions 1
and 2 are outside most inverted regions, genetic exchange is likely to reduce differentiation and
prevent conversion tract detection.
The observed median gene conversion tract lengths are largest in the proximal and
distal regions of Muller C and are reduced in size in the inverted regions (Figure 5). The
detected events in the proximal and distal regions may be the result of cross overs rather than
gene conversion tracts. The Betran *et al*. (1997) method uses informative nucleotide sites to
infer exchanges between arrangements. The accumulation of genetic differences among gene
arrangements can extend beyond the boundaries of the inversion breakpoints providing
information about gene exchanges. In this case, the exchange events are longer than what is
observed within the inverted regions. The most proximal region (1) has the largest median
conversion track length compared to the most distal region (14). This may reflect that region 1
is further from the influence of interference caused by inversion breakpoints (GONG *et al.* 2005).
Most common arrangement pairs are homosequential across regions 1 to 3 with the exception
of the PP arrangement, which has its proximal breakpoint at the boundary of regions 2 and 3.
As a result, there is likely to be less interference of inversion breakpoints in the more proximal
regions leading to resolution of exchanges as cross overs and longer sequences being
exchanged.
We observed exchanges among all pairs of gene arrangements, although the number of
events from donor to recipients are either lower or greater than expected in a permutation test
(Figure 6). For most pairs of arrangements, there is a bias in which arrangement is the donor
versus the recipient. For instance, AR is the recipient of more tracts from CH than expected
while there is a deficiency of events in the reverse direction. There is no clear pattern that might
suggest the nature of the bias. The expected frequency of heterozygotes, common versus rare,
does not explain the biases (pink versus black arrows, respectively in Figure 6). No arrangement
tends to be biased for being a donor or recipient. For instance, four arrangements (AR, PP, ST,
and TL) have excess conversions with two arrangements and a deficient of exchanges with two
arrangements. CH is the only arrangement that shows a bias against being a recipient of
conversion tracts with deficiencies with three arrangements and one case of showing an excess
of being a donor with AR. Thus, we failed to find a plausible explanation for the deficiencies and
excesses of being donors or recipients.
The rate of gene conversion is sufficient to homogenize the sequences among the gene
arrangements, yet one observes differentiation of some genes (outliers) and not others across
the inverted segments. Our analysis of gene conversion coverage tracts shows that gene
conversion occurs less frequently in outlier versus non-outlier genes. Tests that random shuffle
the gene conversion tracts shows that there are enough events to homogenize the sequences.
The reason we can detect these outlier loci is that we observe differentiation in the face of the
homogenizing effect of gene flux, which works in a similar manner as gene flow. This requires
that the various gene arrangements accumulate genetic differences and that there is sufficient

time for gene conversion events to accumulate. The Muller C inversion polymorphism in *D.*
pseudoobscura is estimated to be at least 1 million years old (AQUADRO *et al.* 1991; WALLACE *et*
al. 2011) with the AR arrangement being the newest arrangement of the set originating 350,000
years ago. Despite being a young arrangement, our sample of AR chromosomes has over 30
thousand gene conversion events with a mean coverage of 2.33 events per nucleotide and 36
detected outlier genes with a mean coverage of 0.11. A similar pattern of reduced coverage in
outlier genes is observed in the other four older arrangements.
Essentially what is observed here is an example of survivorship bias. Gene conversion
events survive over time in less consequential genes within the inverted regions, while gene
conversion events in outlier genes from different arrangements do not survive because they
damage the important features of the inversion. We assume that these outlier genes are in fact
adaptive genes essential for the gene arrangements survival. Thus, gene conversion allows
one to detect a signal of genes involved in adaptive evolution within gene arrangements despite
cross over suppression in heterokaryotypes.
Each of the major gene arrangements examined in *D. pseudoobscura* have evidence for
multiple outlier genes that show arrangement specific variation. We show here that the
arrangement specific variation in outlier genes is observed despite the homogenizing effect of
extensive genetic flux among arrangements driven by gene conversion events through the
history of these chromosomes. These data are consistent with a model where the inversions
captured sets of genes involved in local adaptation (KIRKPATRICK AND BARTON 2006). We
propose that the local populations of *D. pseudoobscura* in different physiographic provinces
(DOBZHANSKY 1944; LOBECK 1948) are selected for different locally adapted phenotypes
encoded by multiple genes. These polygenic traits were generated through shuffling of allelic
variation to generate adaptive combinations in different environments. Adaptive combinations in
one environment may not be beneficial in another habitat and are likely to generate maladaptive
types through recombination. This creates a setting for chromosomal inversions to establish if
they capture adaptive combinations of genes (CHARLESWORTH AND CHARLESWORTH 1973). The
set of adaptive genes is not likely to be recognized as selective targets initially because there
will be extensive linkage disequilibrium of SNPs across the chromosome from the ancestral
arrangement. As the ancestral and derived arrangements diverge in sequence, this sets up
conditions to recognize adaptive loci. Gene conversion begins to homogenize the less
important regions within the inversion and with sufficient time differentiated adaptive genes can
be recognized.
If genes within the inverted region are neutral, then we would expect to see the highest
levels of genetic differentiation at or near the inversion breakpoints and not within the central
regions of the inversion (NAVARRO *et al.* 2000; MUNTE *et al.* 2005; NOOR *et al.* 2007; GUERRERO
AND KIRKPATRICK 2014). We observe outlier genes distributed across the inverted regions and
are not restricted to the breakpoint regions even in the largest inversion event considered in this
study, Pikes Peak. The inversion that gave rise to the PP arrangement accounts for 60% of
Muller C. Double cross overs would be expected to homogenize the central region of the
inverted region, yet 129 outlier loci are distributed across the rearranged segment. This
differentiation pattern is the so-called suspension bridge pattern where the towers of the bridge
are the regions of high differentiation or outlier genes and the cables are the regions of reduced
differentiation caused by the accumulation of gene conversion events.
An alternative model to the capture model is one where new inversions gain beneficial
alleles within the arrangement over time. This model does not seem plausible because in this
scenario, one would predict that the repeated selected sweeps that add multiple adaptive alleles
would continue to remove segregating variation from the inverted region making it impossible to
identify outlier genes. A majority of nucleotide sites within the arrangement would be in near
absolute linkage disequilibrium at all times due to the repeated sweeps (FULLER *et al.* 2019).

In summary, we show here that gene conversion is a powerful force for genetic flux
among different gene arrangements that can allow population geneticists to identify adaptive
loci within the inversion. Adaptive loci are discovered as differentiated loci surrounded by
regions of relative homogeneity among arrangements. This approach will not work for recent
inversions because sufficient time must pass for differentiation among arrangements and for
gene conversion to homogenize non-consequential genes. The inversion polymorphism of *D.*
pseudoobscura is sufficiently old that we have identified multiple loci within each arrangement
are identified as selective targets. We think this supports the model that each arrangement
captured sets of locally adapted alleles. Locally adapted inversions were favored because they
prevented the formation of maladaptive combinations from other locally adapted migrant
chromosomes (KIRKPATRICK AND BARTON 2006). When an adaptive QTL is discovered to map
within an inversion, researchers often have a look of despair on their faces. This study
suggests there is hope to identify an adaptive QTL within the confines of an inversion.
508

Acknowledgements

The data presented in this work was generated through support from the National Institutes of Health R01 GM 098478 to (S.W.S.). We thank the sequencing team at the Baylor College of Medicine Human Genome Sequencing Center for their help in sequencing data generation. The funders had no role in study design, data collection, data analysis, the decision to publish, or preparation of the manuscript.

515   Table 1. Gene conversion tracts detected in five recipient gene arrangements in *D. pseudoobscura*.

Gene Arrangement	Strains	Observed Tracts	Expected Tracts	Tracts/Strain	Σ Tract Length	GC Coverage
Arrowhead (AR)	15	30,129	44,342	2,008.6	46,173,391	2.33
Standard (ST)	8	14,982	26,185	1,872.8	25,122,088	1.27
Pikes Peak PP)	10	14,043	20,971	1,404.3	15,948,639	0.81
Chiricahua (CH)	9	20,935	33,259	2,326.1	31,529,250	1.59
Tree Line (TL)	9	18,354	34,861	2,039.3	23,830,343	1.20
Total	51	98,443	159,618	1,930.2	142,603,711	7.20

Strains, number of strains within arrangement; Observed Tracts, observed number of gene conversion tracts detected; Expected
Tracts, expected number of gene conversion tracts = Observed Tracts/(1-Pr(unobserved tracts)), where Pr(unobserved tracts) is
from equation 9 in Betran *et al*. (1997); Tracts/Strain, mean number of gene conversion tracts per strain; Tracts, Σ Tract Length, sum
of nucleotides in all gene conversion tracts; GC Coverage, Σ Tract Length divided by nucleotides in Muller C (19,787,792).

Table 2. Gene Conversion Tract Coverage of Outlier and Non-Outlier Genes on Muller C of
Drosophila pseudoobscura.

Gene Arrangement	Outlier Obs Coverage (SD)	Non-Outlier Obs Coverage (SD)	Min – Max
Arrowhead (AR)			
Transcripts	36	2,632	
Nucleotides	78,468 bp	4,099,525 bp	
Permute Out	0.112 (0.503)*	2.585 (3.064)	1.406 – 4.037
Shuffle CT			0.679 – 2.652
ML Parameter			2.018 – 4.066
Standard (ST)			
Transcripts	82	2,596	
Nucleotides	179,724 bp	3,998,269 bp	
Permute Out	0.067 (0.356)*	1.415 (1.662)	0.974 – 1.926
Shuffle CT			0.563 – 1.758
ML Parameter			1.415 – 2.565
Pikes Peak (PP)			
Transcripts	149	2,519	
Nucleotides	365,249 bp	3,812,744 bp	
Permute Out	0.048 (0.295)*	0.922 (1.488)	0.609 – 1.156
Shuffle CT			0.233 – 0.922
ML Parameter			0.922 – 2.242
Chiricahua (CH)			
Transcripts	40	2,628	
Nucleotides	95,262 bp	4,082,731 bp	
Permute Out	0.055 (0.322)*	1.735 (1.873)	1.087 – 2.709
Shuffle CT			0.483 – 1.735
ML Parameter			1.595 – 3.199
Tree Line (TL)			
Transcripts	54	2,614	
Nucleotides	113,586 bp	4,064,407 bp	
Permute Out	0.101 (0.444)*	1.290 (1.656)	0.832 – 1.917
Shuffle CT			0.216 – 1.290
ML Parameter			0.658 – 1.660

*, P<0.0001 the observed coverage value for the outlier genes is less than that observed in the
randomly permuted data. Min – Max, minimum and maximum values from 10,000 replicates;
Permute Out, permutation test where outlier label was permuted; Shuffle CT, permutation test
where the observed conversion tracts were shuffled; ML Parameter, test where the expected
number of conversion tracts with random tract length with maximum likelihood parameter ϕ is
used.

Figure Legends

Figure 1. Network of *Drosophila pseudoobscura* gene arrangements. The network of gene arrangements (HY, ST, AR, PP, SC, CU, TL, and CH) and the inversion events (colored arrows) are shown to the left where the HY arrangement is the ancestral arrangement. One the right side are the cytogenetic maps for the gene arrangements along with the inverted segments indicated by the colored brackets. The color-coded lines below each arrangement show the locations of the different inversion breakpoints. Outlier genes in each of the gene arrangements are indicated by a black dot. An outlier gene had a significant elevation of derived alleles or population specific branch lengths and had either fixed amino acid differences or differential expression. Below the AR arrangement, 14 regions where genes are in the same order among all chromosomes in the set. The dotted line below the CH, ST, PP, and AR arrangements indicates the regions that would be inverted in ST/AR, ST/CH, AR/CH, or AR/PP heterozygotes whose frequency is at least 10% at least one locality in the Southwestern United States (DOBZHANSKY 1944).

Figure 2. Detection of gene conversion. A map of SNP polymorphism is shown for a short region of the third chromosome of *D. pseudoobscura*. Each column represents a SNP with gray boxes matching the reference genome in the first row. A nucleotide that differs from the reference genome is represented in one of four colors (see legend on the right-hand side). Multi-site genotypes are shown for six gene arrangements along with their sample sizes: Arrowhead (AR), Standard (ST), Pikes Peak (PP), Chiricahua (CH), Tree Line (TL), Cuernavaca (CU), and the outgroup *D. miranda*. Two inferred gene conversion events are shown with the black outline box indicating the recipient sequence and the light blue outline boxes shown in light blue representing the source sequence. The length of the conversion tract is determined from the first and last nucleotide coordinate of the detected tract.

Figure 3. Example of a gene conversion map. A. The locations of non-outlier (black) and outlier genes (blue) are shown in this section of the Muller C nucleotide sequence. B. The positions of mapped gene conversion tracts are shown on the Muller C nucleotide sequence. C. The gene conversion tracts are converted to a histogram based on the number of tracts that cover each nucleotide of the Muller C sequence.

Figure 4. Distribution of the number of conversion tracts within recipient gene arrangement for 14 regions along Muller C of *D. pseudoobscura*. Dots underneath the x-axis indicate regions within gene arrangement with a deficiency (blue) or excess of gene conversion tracts based on residuals from a Chi Square test of homogeneity.

Figure 5. Distribution of conversion tract lengths within recipient gene arrangement for 14 regions along Muller C of *D. pseudoobscura*. Dots underneath the x-axis indicate regions within gene arrangement with a deficiency (blue) or excess of gene conversion tract lengths based on a random permutation test.

Figure 6. Number of gene conversion events per strain between pairs of gene arrangements in *D. pseudoobscura* in 14 different syntenic blocks. The five gene arrangements are abbreviated with their two letter codes: AR, Arrowhead; CH, Chiricahua; PP, Pikes Peak; ST, Standard; TL, Tree Line. Histograms show the number of gene conversion events in the 14 syntenic blocks with the maximum value on all scales being 2,500. Blue and pink boxes below each bar indicate regions with a significant deficiency (blue) or excess (pink) of events based on a random permutation test (see the text). The number above each histogram indicate the total number of events across all 14 regions with a blue number indicating a deficiency and a pink number indicating an excess of events. The

double headed arrows connect pairs of arrangements with the arrows pointing to the
number of events in the recipient arrangement from the donor arrangement. For
instance, 1802 conversion events were from ST to TL and 1973 conversion events were
from TL to ST. The pink arrows indicate pairs of arrangements that are expected to
have frequencies greater than 10% in populations in the Southwestern United States.
Gray shading is provided to delineate recipient data for each arrangement.
Figure 7. Mean gene conversion tract coverage in 2,668 transcripts across Muller C in *D.*
pseudoobscura in five gene arrangements. Coverage is the number of times that a
nucleotide is covered by a gene conversion tract. Coverage was estimated for each
exon within a transcript and the mean coverage for the entire transcript was averaged
over all exons. Outlier genes are shown with an orange marker while non-outliers are
shown with a gray marker. The location of the derived inversion giving rise to each gene
arrangement are shown with a black bar.

Figure 1

Figure 2.

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Arrowhead

Standard

Pikes Peak

Chiricahua

Tree Line

Literature Cited

Anderson, W. W., J. Arnold, D. G. Baldwin, A. T. Beckenbach, C. J. Brown *et al.*, 1991 Four decades of inversion polymorphism in *Drosophila pseudoobscura*. Proceedings of the National Academy of Sciences USA 88**:** 10367-10371.

Aquadro, C. F., A. L. Weaver, S. W. Schaeffer and W. W. Anderson, 1991 Molecular evolution of inversions in *Drosophila pseudoobscura* : The amylase gene region. Proceedings of the National Academy of Sciences USA 88**:** 305-309.

Betran, E., J. Rozas, A. Navarro and A. Barbadilla, 1997 The estimation of the number and the length distribution of gene conversion tracts from population DNA sequence data. Genetics 146**:** 89-99.

Chan, A. H., P. A. Jenkins and Y. S. Song, 2012 Genome-Wide Fine-Scale Recombination Rate Variation in Drosophila melanogaster. PLOS Genetics 8**:** e1003090.

Charlesworth, B., and D. Charlesworth, 1973 Selection of new inversion in multi-locus genetic systems. Genetical Research 21**:** 167-183.

Chovnick, A., 1973 Gene conversion and transfer of genetic information within the inverted region of inversion heterozygotes. Genetics 75**:** 123-131.

Crown, K. N., D. E. Miller, J. Sekelsky and R. S. Hawley, 2018 Local Inversion Heterozygosity Alters Recombination throughout the Genome. Current Biology 28**:** 2984-2990.

Dobzhansky, T., 1944 Chromosomal races in *Drosophila pseudoobscura* and *Drosophila persimilis*. Carnegie Inst. Washington Publ. 554**:** 47-144.

Dobzhansky, T., 1948 Genetics of natural populations. XVI. Altitudinal and seasonal changes in certain populations of *Drosophila pseudoobscura* and *Drosophila persimilis*. Genetics 33**:** 158-176.

Dobzhansky, T., and M. L. Queal, 1938 Genetics of natural populations. II. Genic variation in populations of *Drosophila pseudoobscura* inhabiting isolated mountain ranges. Genetics 23**:** 463-484.

Dobzhansky, T., and A. H. Sturtevant, 1938 Inversions in the chromosomes of *Drosophila pseudoobscura*. Genetics 23**:** 28-64.

Fuller, Z. L., G. D. Haynes, S. Richards and S. W. Schaeffer, 2017 Genomics of natural populations: Evolutionary forces that establish and maintain gene arrangements in *Drosophila pseudoobscura*. Molecular Ecology 26**:** 6539-6562.

Fuller, Z. L., S. A. Koury, N. Phadnis and S. W. Schaeffer, 2019 How chromosomal rearrangements shape adaptation and speciation: Case studies in *Drosophila pseudoobscura* and its sibling species *Drosophila persimilis*. Molecular Ecology 28**:** 1283-1301.

Gong, W. J., K. S. McKim and R. S. Hawley, 2005 All Paired Up with No Place to Go: Pairing, Synapsis, and DSB Formation in a Balancer Heterozygote. PLOS Genetics 1**:** e67.

Guerrero, R. F., and M. Kirkpatrick, 2014 Local adaptation and the evolution of chromosome fusions. Evolution 68**:** 2747-2756.

Guerrero, R. F., F. Rousset and M. Kirkpatrick, 2012 Coalescent patterns for chromosomal inversions in divergent populations. Philosophical Transactions of the Royal Society B: Biological Sciences 367**:** 430-438.

Kirkpatrick, M., and N. Barton, 2006 Chromosome inversions, local adaptation and speciation. Genetics 173**:** 419-434.

Korunes, K. L., and M. A. F. Noor, 2019 Pervasive gene conversion in chromosomal inversion heterozygotes. Mol Ecol 28**:** 1302-1315.

Lindsley, D. L., and L. Sandler, 1977 The genetic analysis of meiosis in female *Drosophila melanogaster*. Philosophical Transactions of the Royal Society of London Biology 277B**:** 295-312.

Lobeck, A. K., 1948 Physiographic diagram of North America, pp. C. S. Hammond & Co., Maplewood, NJ.

Maynard Smith, J., and J. Haigh, 1974 The hitch-hiking effect of a favorable gene. Genetical Research 23: 23-35.

Miller, D. E., C. B. Smith, N. Y. Kazemi, A. J. Cockrell, A. V. Arvanitakas *et al.*, 2016 Whole-genome analysis of individual meiotic events in *Drosophila melanogaster* reveals That noncrossover gene conversions are insensitive to interference and the centromere effect. Genetics 203: 159-171.

Miller, D. E., S. Takeo, K. Nandanan, A. Paulson, M. M. Gogol *et al.*, 2012 A Whole-Chromosome Analysis of Meiotic Recombination in Drosophila melanogaster. G3 Genes|Genomes|Genetics 2: 249-260.

Muller, H. J., 1940 Bearings of the 'Drosophila' work on systematics., pp. 185-268 in *The New Systematics*, edited by J. Huxley. Clarendon Press, Oxford.

Munte, A., J. Rozas, M. Aguade and C. Segarra, 2005 Chromosomal inversion polymorphism leads to extensive genetic structure: a multilocus survey in *Drosophila subobscura*. Genetics 169: 1573-1581.

Navarro, A., A. Barbadilla and A. Ruiz, 2000 Effect of inversion polymorphism on the neutral nucleotide variability of linked chromosomal regions in *Drosophila*. Genetics 155: 685-698.

Navarro, A., E. Betrán, A. Barbadilla and A. Ruiz, 1997 Recombination and gene flux caused by gene conversion and crossing over in inversion heterokaryotypes. Genetics 146: 695-709.

Nei, M., K. I. Kojima and H. E. Schaffer, 1967 Frequency changes of new inversions in populations under mutation-selection equilibria. Genetics 57: 741-750.

Noor, M. A., D. A. Garfield, S. W. Schaeffer and C. A. Machado, 2007 Divergence between the *Drosophila pseudoobscura* and *D. persimilis* genome sequences in relation to chromosomal inversions. Genetics 177: 1417–1428.

Ranz, J. M., D. Maurin, Y. S. Chan, M. v. Grotthuss, L. W. Hillier *et al.*, 2007 Principles of genome evolution in the *Drosophila melanogaster* species group Public Library of Science Biology 5: 1366-1381.

Richards, S., Y. Liu, B. R. Bettencourt, P. Hradecky, S. Letovsky *et al.*, 2005 Comparative genome sequencing of *Drosophila pseudoobscura*: Chromosomal, gene and *cis*-element evolution. Genome Research 15: 1-18.

Rozas, J., and M. Aguadé, 1993 Transfer of genetic information in the *rp49* region of *Drosophila subobscura* between different gene arrangements. Proceedings of the National Academy of Sciences USA 90: 8083-8087.

Rozas, J., and M. Aguadé, 1994 Gene conversion is involved in the transfer of genetic information between naturally occurring inversions of *Drosophila*. Proceedings of the National Academy of Sciences USA 91: 11517-11521.

Rozas, J., A. Ferrer-Mata, J. C. Sánchez-DelBarrio, S. Guirao-Rico, P. Librado *et al.*, 2017 DnaSP 6: DNA Sequence Polymorphism Analysis of Large Data Sets. Molecular Biology and Evolution 34: 3299-3302.

Schaeffer, S. W., and W. W. Anderson, 2005 Mechanisms of genetic exchange within the chromosomal inversions of *Drosophila pseudoobscura*. Genetics 171: 1729-1739.

Shriver, M., G. Kennedy, E. Parra, H. Lawson, V. Sonpar *et al.*, 2004 The genomic distribution of population substructure in four populations using 8,525 autosomal SNPs. Human Genomics 1: 274 - 286.

Sperlich, D., and P. Pfriem, 1986 Chromosomal polymorphism in natural and experimental populations, pp. 257-309 in *The Genetics and Biology of Drosophila 3e*, edited by M. Ashburner, H. L. Carson and J. N. Thomson. Academic Press, New York, NY.

Stephens, J. C., 1985 Statistical methods of DNA sequence analysis: Detection of intragenic recombination or gene conversion. Molecular Biology and Evolution 2: 539-556.

Sturtevant, A. H., and G. W. Beadle, 1936 The relations of inversions in the X chromosome of
Drosophila melanogaster to crossing over and disjunction. Genetics 21**:** 544-604.
Wallace, A. G., D. Detweiler and S. W. Schaeffer, 2011 Evolutionary history of the third
chromosome gene arrangements of *Drosophila pseudoobscura* inferred from inversion
breakpoints. Molecular Biology and Evolution 28**:** 2219-2229.
Wright, D., and S. W. Schaeffer, 2022 The relevance of chromatin architecture to genome
rearrangements in *Drosophila*. Philosophical Transactions of the Royal Society B:
Biological Sciences 377**:** 20210206.
Yi, X., Y. Liang, E. Huerta-Sanchez, X. Jin, Z. X. P. Cuo *et al.*, 2010 Sequencing of 50 human
exomes reveals adaptation to high altitude. Science 329**:** 75-78.