Three R2R3-MYB transcription factors from banana (*Musa* spp.) activate structural

anthocyanin biosynthesis genes as part of an MBW complex

Mareike Busche[1], Boas Pucker[2], Bernd Weisshaar[1], Ralf Stracke[1]*

[1] Genetics and Genomics of Plants, Faculty of Biology, Bielefeld University, 33615 Bielefeld,

Germany

[2] Institute of Plant Biology & Braunschweig Integrated Centre of Systems Biology (BRICS), TU

Braunschweig, 38106 Braunschweig, Germany

*Corresponding author:

ralf.stracke@uni-bielefeld.de

Abstract

Bananas are among the most popular fruits in the world and provide food security and

employment opportunities in several developing countries. An increased anthocyanin content

could enhance the health promoting properties of banana fruits. The biosynthesis of

anthocyanins is largely regulated at the transcriptional level. However, little is known about

transcriptional activation of anthocyanin biosynthesis in banana. We analysed the regulatory

activity of three *Musa*MYBs predicted by bioinformatic analysis to transcriptionally regulate

anthocyanin biosynthesis in banana. *MusaMYBA1, MusaMYBA2* and *MusaMYBPA2* did not

complement the anthocyanin deficiency phenotype of the *A. thaliana pap1/pap2* mutant.

However, co-transfection experiments in *A. thaliana* protoplasts showed that *Musa*MYBA1,

*Musa*MYBA2 and *Musa*MYBPA2 function as components of a transcription factor complex with

a bHLH and WD40 protein, called MBW complex, resulting in the activation of the *anthocyanin*

synthase and *dihydroflavonol 4-reductase* promoters from *A. thaliana*. The activation potential

of *Musa*MYBA1, *Musa*MYBA2 and *Musa*MYBPA2 increased when combined with the monocot

bHLH *Zm*R instead of the dicot *At*EGL3. This work paves the path towards decoding the MBW

complex-mediated transcriptional activation of anthocyanin biosynthesis in banana. Moreover,

it facilitates research towards an elevated anthocyanin content in banana and other monocot

crops.

Keywords: *Musa acuminata*, banana, specialised metabolites, transcription factor, flavonoid

biosynthesis, R2R3-MYB

Introduction

Bananas (*Musa*) are monocotyledonous, perennial plants which are grown in many tropical and subtropical countries. They are one of the most important food crops, particularly in the developing world (1). While the sweet fruits of dessert bananas are popular in Europe and Northern America, plantains or cooking bananas are commonly eaten as a staple food in Africa and Latin America where they provide food security, as well as employment opportunities (2). Furthermore, banana fruits are rich in several health promoting minerals and beneficial phytochemicals such as vitamins and flavonoids (3).

Flavonoids are a major group of plant specialised metabolites which share a basic structure of two aromatic C6-rings connected by a heterocyclic ring (4). Reorganisations and modifications e.g. oxidation, glycosylation, acylation, and methylation of the carbon skeleton create a versatile group comprising more than 9,000 different flavonoid derivatives (5). Consequently, flavonoids do not only contribute to the nutritional value of fruits, but also play important roles in manifold processes. While the group of red coloured anthocyanin pigments attracts animals for pollination and dispersal of seeds by colouring flowers and fruits, other flavonoids protect plants against UV-B irradiation or increase plant fertility (6-10). Flavonoids from many species have been reported to have anti-pathogenic properties, this includes flavonoids from *Dianthus caryophyllus* which exhibit antifungal activity against the plant's major pest *Fusarium oxysporum* f.sp. *dianthi* (11, 12). The tropical race 4 (TR4) of the banana Fusarium wilt (commonly known as 'panama disease') is caused by another *Fusarium* subspecies called *Fusarium oxysporum* f. sp. *cubense* (Foc) and threatens the global banana production (13). Transcriptome analysis of susceptible and resistant banana cultivars infected by Foc TR4 revealed an increased transcription of flavonoid biosynthesis related genes in the resistant cultivar, suggesting an involvement of flavonoids in the defence against Foc TR4 (14).

Flavonoid biosynthesis is one of the best characterised pathways of the secondary metabolism and has been extensively studied in many plant species (15). In banana, several flavonoid biosynthesis related enzymes have been identified and characterised (16, 17). Regulation of structural genes on a transcriptional level allows a specific response to environmental

influences as well as development and organ specific expression (18-20). MYB transcription

factors are common transcriptional regulators of flavonoid biosynthesis. While some MYBs act

independently, others interact with basic helix-loop-helix (bHLH) and WD40 proteins to form a

protein complex called MBW complex (21). MYB proteins are present in all eukaryotes and

characterised by highly conserved DNA-binding domains (22). These MYB domains consist of

up to three imperfect amino acid repeat sequences, based on which they are classified. R2R3-

MYBs are the most abundant class of plant MYBs and reveal versatile functions in plant-

specific processes (23). Besides core- and specialised metabolism they are also involved in

cell fate and -identity definition, developmental processes and the response to biotic and

abiotic stresses (23). Well-known R2R3-MYBs which act as activators of anthocyanin

biosynthesis include C1 (COLOURED ALEURONE1) from *Zea mays* (*Z. mays*), as well as

PAP1 (PRODUCTION OF ANTHOCYANIN PIGMENT1/MYB75) and PAP2 (MYB90) from

Arabidopsis thaliana (*A. thaliana*) (24, 25). They act as part of an MBW complex and control

the promoters of the anthocyanin biosynthesis related structural genes as for example

ANTHOCYANIDIN SYNTHASE (*ANS*) and *DIHYDROFLAVONOL 4-REDUCTASE* (*DFR*) (26-

29).

In banana (*Musa acuminata, M. acuminata*), 285 R2R3-MYB proteins, including several

putative regulators of flavonoid biosynthesis, have been identified in a genome-wide study

(30). Furthermore, MYB31, MYB4 and MYBPR1 - MYBPR4 have been identified as negative

regulators of flavonoid biosynthesis in banana (31, 32). Despite the recent identification of two

proanthocyanidin biosynthesis activating R2R3-MYBs (33), little functional data is available on

positive regulators (activators) of flavonoid and in particular anthocyanin biosynthesis in

M. acuminata.

Here, we describe the regulatory properties of three *Musa*MYBs, named *Musa*MYBA1,

*Musa*MYBA2 and *Musa*MYBPA2, with a possible role in the regulation of anthocyanin

biosynthesis. As one of these *Musa*MYBs was very recently published under the term

*Musa*MYBPA2 (33), we used this name to avoid confusion due to multiple protein naming.

Regulatory activity was investigated by *in planta* complementation experiments of the

anthocyanin deficient *A. thaliana* regulatory *pap1/pap2* mutant and co-transfection

experiments in *A. thaliana* protoplasts (for detailed methods refer to Supplementary File 2).

Our results show that *Musa*MYBA1, *Musa*MYBA2 and *Musa*MYBPA2 are able to activate the

promoters of *AtANS* and *AtDFR* as part of an MBW complex. Moreover, we show that the

activation potential of *Musa*MYBA1, *Musa*MYBA2 and *Musa*MYBPA2 is increased when

combined with the monocot bHLH *Zm*R instead of the dicot bHLH protein ENHANCER OF

GLABRA 3 (*At*EGL3).

Main text

We aimed to analyse the regulatory properties of three *Musa*MYBs which have been previously

assigned with a possible role in positive regulation of anthocyanin biosynthesis (Ma06_g05960

or *MusaMYBA1*, Ma09_g27990 or *MusaMYBA2*, Ma10_g17650 or *MusaMYBPA2*). Since all

three *MusaMYB* genes were detected in the haploid *M. acuminata* reference genome

sequence DH (doubled-haploid) Phang v2 (34, 35), these *MYBs* appear to be present in the

same sub-genome, suggesting that they are different genes and not haplo-copies. We tried to

amplify the corresponding coding sequences (CDSs) on a template collection containing cDNA

from different banana samples. The CDSs of all three *MusaMYB*s were successfully amplified

on cDNA derived from peel tissue.

In a first approach, we performed a complementation assay with the regulatory *A. thaliana*

pap1/pap2 double mutant which cannot produce anthocyanins in the seedling (Figure 1).

Seedlings were grown on anthocyanin synthesis-inducing media to analyse the ability of

2x35S-driven *MusaMYB*s to complement the *pap1/pap2* anthocyanin deficiency. While wild-

type seedlings accumulated high amounts of red anthocyanin pigments, *pap1/pap2* seedlings

did not. Even though *MusaMYBA1*, *MusaMYBA2* or *MusaMYBPA2* were successfully

expressed in the transgenic seedlings (Supplementary Figure S1), the anthocyanin level in

pap1/pap2 plants expressing *MusaMYBA1*, *MusaMYBA2* or *MusaMYBPA2* did not differ from

the double mutant. Accordingly, *MusaMYBA1*, *MusaMYBA2* and *MusaMYBPA2* do not seem

to be able to complement the mutant phenotype and thus to regulate anthocyanin biosynthesis

in *A. thaliana* in combination with the *bHLH* and *WD40* genes expressed in *A. thaliana*

seedlings.

Figure 1: *MusaMYBA1, MusaMYBA2* and *MusaMYBPA2* cannot complement the
anthocyanin deficient phenotype of *A. thaliana pap1/pap2* mutant seedlings. (A)
Representative pictures of anthocyanin accumulation in 6-day-old *MusaMYB*-expressing
pap1/pap2 seedlings. A representative plant per construct is displayed. (B) Photometrical
measurement of the sucrose induced anthocyanin content in *pap1/pap2* seedlings expressing
2x35S-driven *MusaMYB*s. Col-0, Nö-0 (wildtypes) and *pap1/pap2* were used as controls. Error
bars indicate the standard deviation of three biological replicates. The different numbers
represent individual, independent, transgenic lines.

Lloyd *et al.* (28) analysed the *Z. mays* anthocyanin regulator C1 by generation of transgenic

A. thaliana overexpression lines. In their study, Lloyd *et al.* generated three independent

ZmC1-expressing *A. thaliana* lines that did not show an increased anthocyanin content

compared to wildtype. Similar results were obtained in transgenic tobacco. However, further

experiments suggested that *Zm*C1 needs to interact with the maize bHLH *Zm*R to activate

anthocyanin biosynthesis in the heterologous *A. thaliana* system (28). Phylogenetic analysis

(36) showed that the anthocyanin biosynthesis regulating R2R3-MYBs from several monocots

such as *Z. mays* and *Oryza sativa* (*O. sativa*) are part of a different phylogenetic clade than

the anthocyanin regulators from several dicots, such as *A. thaliana* or *Vitis vinifera* and further

monocots, including *Allium cepa* and *Lilium hybrida*. Experiments in snapdragon also showed

that the monocot *Ac*MYB1 can activate anthocyanin production in dicots (36). The phylogenetic

differences between anthocyanin biosynthesis activating R2R3-MYBs and the dependency of

*Zm*C1 on *Zm*R in dicots hint at an explanation for our observations. For example, the

*Musa*MYBs could depend on their endogenous or at least monocot bHLH for effective

activation of structural anthocyanin biosynthesis genes. To investigate the phylogenetic

differences between anthocyanin biosynthesis regulating R2R3-MYBs from different plant

species, an approximately maximum-likelihood tree was built (Figure 2A). The resulting tree,

which also contained R2R3-MYBs which activate other branches of flavonoid biosynthesis,

revealed two distinct clades of anthocyanin-related R2R3-MYBs (highlighted in red).

*Musa*MYBA1, *Musa*MYBA2 and *Musa*MYBPA2 form a clade with MYB10 from *Triticum*

aestivum and exhibit a close evolutionary relationship with *Zm*C1, *Os*C1 and anthocyanin

regulating MYBs from further monocots. In contrast, *At*PAP1 and *At*PAP2 fall into a second

clade of anthocyanin-related MYBs. Furthermore, differences between anthocyanin

biosynthesis regulating bHLHs were analysed in a second approximately maximum-likelihood

tree (Figure 2B). The tree showed that monocot bHLH proteins involved in the regulation of

anthocyanin biosynthesis form a separate clade (highlighted in red). Both, *Zm*R and several

putative anthocyanin biosynthesis regulating bHLHs from banana fall into this clade. These

phylogenetic analyses show that anthocyanin biosynthesis regulating R2R3-MYB and bHLH

proteins from several monocot species seem to be distinct from other R2R3-MYB and bHLH

proteins regulating anthocyanin biosynthesis, for example from *A. thaliana*. These differences

could imply a dependency of *Musa*MYBA1, *Musa*MYBA2 and *Musa*MYBPA2 on a banana or

other monocot bHLH.

To follow up this idea, we further investigated the regulatory properties of the three *Musa*MYBs

performing co-transfection assays (Figure 3) in At7 protoplasts with different bHLH proteins

from *A. thaliana* (*At*EGL3) or *Z. mays* (*Zm*R) and the *A. thaliana* WD40 protein

TRANSPARENT TESTA GLABRA 1 (*At*TTG1). Thereby, their potential to activate the

promoters of

169
Figure 2: Rooted approximately maximum-likelihood trees of MYB (A) and bHLH (B)
transcription factors. Circle sizes represent bootstrap values. Banana proteins are
highlighted in blue.

AtDFR and *AtANS*, which are important structural genes of anthocyanin biosynthesis (37), was

analysed. While none of the *Musa*MYBs was able to independently activate *proAtANS* or

proAtDFR, which both contain conserved *cis*-regulatory elements, a slight activation of

proAtDFR was detectable when *Musa*MYBA1 or *Musa*MYBPA2 were combined with *At*EGL3

and *At*TTG1. In combination with *At*EGL3 and *At*TTG1, *Musa*MYBPA2 showed the strongest

activation of *proAtDFR* and was also able to activate *proAtANS*. When combined with *Zm*R

and *At*TTG1, the three MYBs *Musa*MYBA1, *Musa*MYBA2, and *Musa*MYBPA2 exhibited a

significant activation potential on *proAtDFR* and *proAtANS*. All three *Musa*MYBs showed a

higher activation potential in an MBW complex with *Zm*R and *At*TTG1 than in combination with

*At*EGL3 and *At*TTG1.

These results display that *Musa*MYBA1, *Musa*MYBA2 and *Musa*MYBPA2 are able to activate

proAtANS and *proAtDFR* as part of an MBW complex and *Musa*MYBPA2 shows the strongest

activation potential. Furthermore, *Musa*MYBA1, *Musa*MYBA2 and *Musa*MYBPA2 show a

higher activation potential when combined with the monocot bHLH *Zm*R instead of the dicot

bHLH *At*EGL3.

The previously mentioned phylogenetic differences could be an explanation for the observed

higher activation potentials of the analysed *Musa*MYBs in combination with *Zm*R and *At*TTG1

instead of *At*EGL3 and *At*TTG1. It is conceivable that these differences impede the interaction

between the tested *Musa*MYBs and the dicot bHLH *At*EGL3. As *Musa*MYBPA2 shows the

highest activation potential and is also able to activate *proAtANS* in combination with *At*EGL3

or *Zm*R, differences between the three *Musa*MYBs that affect the interaction with the *At*EGL3

are likely and should be further studied. Interestingly, *Musa*MYBPA2 did not complement the

anthocyanin deficient phenotype of *A. thaliana pap1/pap2* mutant seedlings, but was able to

activate *proAtANS* and *proAtDFR* when combined with *At*EGL3 in co-transfection experiments

in *A. thaliana* protoplasts. Even though previous studies showed *AtEGL3* expression in

A. thaliana seedlings (21), it is possible that the level was too low to activate anthocyanin

biosynthesis in combination with *Musa*MYBPA2. Moreover, the At7 cell line used for co-

transfection experiments was established more than 25 years ago. This long time of

propagation in suspension cell culture caused a variety of genomic and transcriptomic changes

(38). These changes could also explain differences between the analysis in seedlings and cell

culture.

Figure 3: *Musa*MYBA1, *Musa*MYBA2 and *Musa*MYBPA2 can activate *proAtDFR* and *proAtANS* as part of an MBW complex. The ability of *Musa*MYBs to activate *proAtDFR-GUS* and *proAtANS-GUS* reporter constructs in combination with different bHLH proteins (*At*EGL3, *Zm*R) and a WDR (*At*TTG1) was analysed by co-transfection in *A. thaliana* At7 protoplasts. The relative promoter activity refers to the measured GUS reporter enzyme activity. Promoter activity is given relative to the values obtained for the *A. thaliana* MBW complex (*At*PAP1, *At*EGL3, *At*TTG1). Error bars indicate the standard deviation of five independent biological replicates. Statistical significance is indicated by asterisks which mark p-values < 0.05.

The expression profiles of *MusaMYBA1*, *MusaMYBA2* and *MusaMYBPA2* have previously

been analysed in embryogenic cell suspension, seedling, root and different developmental

stages of leaf, pulp and peel (30). The expression of *MusaMYBA1* was strongest in pulp

(developmental stage S2-S3), *MusaMYBA2* expression was relatively low in all samples and

MusaMYBPA2 showed highest expression in seedlings and early developmental stages of

pulp (S1). This gene expression pattern could give a hint at organ specific *MusaMYBA1*,

MusaMYBA2 and *MusaMYBPA2* activity in banana. Since flavonoid biosynthesis is largely

regulated at the transcriptional level, it would be particularly interesting to analyse the

expression levels of *MusaMYBA1*, *MusaMYBA2,* and *MusaMYBPA2* in anthocyanin rich

organs such as bract or pseudostem to relate anthocyanin content to *MusaMYB* transcript

abundance.

Deng *et al*. (32) performed an expression analysis of flavonoid biosynthesis related genes

using leaves of banana plants overexpressing the anthocyanin repressor *MusaMYB4*. They

found that the expression of *MusaMYBA1* and *MusaMYBPA2* is decreased along with

MusaDFR and *MusaANS* in plants overexpressing *Musa*MYB4, compared to wildtype. This

data supports a proposed functionality of *Musa*MYBA1 and *Musa*MYBPA2 as transcriptional

activators of anthocyanin biosynthesis in banana, as *MusaMYB4* could cause a feedback

regulation of the positive regulators of anthocyanin biosynthesis, of *MusaMYBA1* and

MusaMYBPA2.

Very recently, Rajput *et al*. (33) showed that *Musa*MYBPA2 can activate the banana *ANS,*

ANR and LAR promoters. Moreover, they showed that *Musa*MYBPA2 can partially rescue the

proanthocyanin deficiency of the *A. thaliana tt2-1* mutant. The reported ability of *Musa*MYBPA2

to activate *proMusaANS* supports the proposed role of *Musa*MYBPA2 in the regulation of

anthocyanin biosynthesis. However, the partial complementation of the proanthocyanin

deficient phenotype of the *A. thaliana tt2-1* R2R3-MYB mutant, as well as the ability to activate

the banana *ANR* and *LAR* promoters, shows an additional role in the regulation of

proanthocyanidin biosynthesis. Such dual roles in the regulation of flavonoid biosynthesis have

previously been suggested for R2R3-MYB transcription factors from grape, blueberry and

apple (39-42). In several species, including *A. thaliana, Z. mays* and *Petunia hybrida*,

anthocyanin biosynthesis is regulated by an MBW complex (21, 28, 43-45). Our co-transfection

assays showed that *Musa*MYBA1, *Musa*MYBA2 and *Musa*MYBPA2 are able to activate

proAtANS and *proAtDFR* as part of an MBW complex with *Zm*R and *At*TTG1 *in planta*. As

presented in a previous study by Pucker *et al.* (30), it is known that *Musa*MYBA1, *Musa*MYBA2,

and *Musa*MYBPA2 all contain a bHLH-binding consensus motif (27). Thus, the regulation of

anthocyanin biosynthesis by *Musa*MYBA1, *Musa*MYBA2, and *Musa*MYBPA2 likely depends

on bHLH and WD40 proteins. Based on our tree (Figure 2B), the bHLH encoding genes

Ma05_g23010, Ma11_g19640, Ma11_g16060, Ma08_g11190 and Ma11_g03740 could be

suitable candidates for further studies on the transcriptional activation of anthocyanin

biosynthesis by MBW complexes. Moreover, future analyses should additionally include

*Musa*TTG1 to identify a whole functional MBW complex in banana and to find out if the use of

endogenous WD40 protein further improves the activation potential of the MBW complex.

The presented results show that *Musa*MYBA1, *Musa*MYBA2, and *Musa*MYBPA2 have the

capacity to transcriptionally activate structural anthocyanin biosynthesis genes in an MBW

complex with a suitable bHLH partner. This is a step towards decoding the MBW complex-

mediated transcriptional activation of flavonoid biosynthesis in banana. Moreover, it is a basis

for further research towards increased anthocyanin production in banana which could promote

fruit quality and disease resistance.

Limitations

The co-transfection assays revealed that *Musa*MYBA1, *Musa*MYBA2, and *Musa*MYBPA2 can

activate two structural genes of anthocyanin biosynthesis from *A. thaliana* as part of an MBW

complex. In the future this analysis could be expanded to further genes. Moreover, co-

transfection experiments with the promoters of structural anthocyanin biosynthesis genes and

bHLH and WDR candidates from banana should be performed. In order to elucidate the

regulatory role in banana and confirm possible target genes, future studies should include an

overexpression of the three *Musa*MYBs in banana.

Declarations

Competing interests

273 The authors declare that they have no competing interests.

Funding

276 This work was supported by the basic funding of the chair of Genetics and Genomics of Plants

277 provided by the Bielefeld University/Faculty of Biology and the Open Access Publication Fund

278 of Bielefeld University.

Authors' Contributions

281 MB, BP and RS planned the experiments. MB performed the experiments and analysed the

282 data. RS and BW supervised the project. MB wrote the initial draft. RS and BP revised the

283 manuscript.

Acknowledgements

286 We are grateful to Melanie Kuhlmann and Janina Grabowski for their excellent technical

287 assistance and to Andrea Voigt for her competent help in the greenhouse. We thank Martin

288 Hülskamp for providing the *Zm*R entry clone plasmid and Ashutosh Pandey for providing the

289 banana cDNAs.

Abbreviations

292 ANS   anthocyanidin synthase

293 *A. thaliana*   *Arabidopsis thaliana*

294 bHLH   basic helix-loop-helix

295 C1   COLOURED ALEURONE1

296 DFR   dihydroflavonol 4-reductase

297	EGL3	ENHANCER OF GLABRA 3
298	Foc	*Fusarium oxysporum* f. sp. *cubense*
299	*M. acuminata*	*Musa acuminata*
300	*O. sativa*	*Oryza sativa*
301	PAP	production of anthocyanin pigment
302	TR4	tropical race 4
303	TTG1	TRANSPARENT TESTA GLABRA 1
304	*Z. mays*	*Zea mays*

Supplementary information

Figure S1: Expression analysis of *MusaMYBA1, MusaMYBA2 and MusaMYBPA2* in *A.*

thaliana pap1/pap2 seedlings.

Figure S2: Full-length gels from Figure S1.

Table S1: Oligonucleotide primers used in this work.

Table S2: IDs of protein sequences used for the construction of the phylogenetic tree.

File S2: Methods

References

1. Scott GJ. A review of root, tuber and banana crops in developing countries: past,
present and future. International Journal of Food Science and Technology. 2021;56(3):1093-
114.
2. Arias P, Dankers C, Liu P, Pilkauskas P. The World Banana Economy, 1985-2002 2003
[http://www.fao.org/3/y5102e/ye00.htm].   Available   from:
http://www.fao.org/3/y5102e/y5102e00.htm.
3. Singh B, Singh JP, Kaur A, Singh N. Bioactive compounds in banana and their
associated health benefits - A review. Food Chemistry. 2016;206:1-11.
4. Winkel-Shirley B. Flavonoid biosynthesis. A colorful model for genetics, biochemistry,
cell biology, and biotechnology. Plant Physiology. 2001;126:485-93.
5. Wen W, Alseekh S, Fernie AR. Conservation and diversification of flavonoid
metabolism in the plant kingdom. Current Opinion in Plant Biology. 2020;55:100-8.
6. Ishikura N, Yoshitama K. Anthocyanin-Flavonol Co-pigmentation in Blue Seed Coats
of Ophiopogon jaburan. Journal of Plant Physiology. 1984;115(2):171-5.
7. Mo Y, Nagel C, Taylor LP. Biochemical complementation of chalcone synthase mutants
defines a role for flavonols in functional pollen. Proceedings of the National Academy of
Sciences of the United States of America. 1992;89:7213-7.
8. Li J, Ou-Lee TM, Raba R, Amundson RG, Last RL. Arabidopsis Flavonoid Mutants Are
Hypersensitive to UV-B Irradiation. The Plant Cell. 1993;5(2):171-9.

9.   Gronquist M, Bezzerides A, Attygalle A, Meinwald J, Eisner M, Eisner T. Attractive and
defensive functions of the ultraviolet pigments of a flower *(Hypericum calycinum)*. Proceedings
of the National Academie of Sciences of the United States of America. 2001;98(24):13745-50.
10.   Grotewold E. The Science of Flavonoids. Grotewold E, editor. Columbus: The Ohio
State University; 2006. 273 p.
11.   Galeotti F, Barile E, Curir P, Dolci M, Lanzotti V. Flavonoids from carnation (Dianthus
caryophyllus) and their antifungal activity. Phytochemistry Letters. 2008;1(1):44-8.
12.   Mierziak J, Kostyn K, Kulma A. Flavonoids as Important Molecules of Plant Interactions
with the Environment. Molecules. 2014;19(10):16240-65.
13.   Dita M, Barquero M, Heck D, Mizubuti ESG, Staver CP. Fusarium Wilt of Banana:
Current Knowledge on Epidemiology and Research Needs Toward Sustainable Disease
Management. Frontiers in Plant Science. 2018;9.
14.   Sun J, Zhang J, Fang H, Peng L, Wei S, Li C, et al. Comparative transcriptome analysis
reveals resistance-related genes and pathways in *Musa acuminata* banana 'Guijiao 9' in
response to Fusarium wilt. Plant Physiology and Biochemistry. 2019;141:83-94.
15.   Liu WX, Feng Y, Yu SH, Fan ZQ, Li XL, Li JY, et al. The Flavonoid Biosynthesis Network
in Plants. International Journal of Molecular Sciences. 2021;22(23).
16.   Pandey A, Alok A, Lakhwani D, Singh J, Asif MH, Trivedi PK. Genome-wide Expression
Analysis and Metabolite Profiling Elucidate Transcriptional Regulation of Flavonoid
Biosynthesis and Modulation under Abiotic Stresses in Banana. Scientific Reports. 2016;6.
17.   Busche M, Acatay C, Martens S, Weisshaar B, Stracke R. Functional Characterisation
of Banana (Musa spp.) 2-Oxoglutarate-Dependent Dioxygenases Involved in Flavonoid
Biosynthesis. Frontiers in Plant Science. 2021;12:701780.
18.   Procissi A, Dolfini S, Ronchi A, Tonelli C. Light-dependent spatial and temporal
expression of pigment regulatory genes in developing maize seeds. The Plant Cell.
1997;9(9):1547-57.
19.   Nesi N, Jond C, Debeaujon I, Caboche M, Lepiniec L. The Arabidopsis *TT2* gene
encodes an R2R3 MYB domain protein that acts as a key determinant for proanthocyanidin
accumulation in developing seed. The Plant Cell. 2001;13(9):2099-114.
20.   Stracke R, Jahns O, Keck M, Tohge T, Niehaus K, Fernie AR, et al. Analysis of
PRODUCTION OF FLAVONOL GLYCOSIDES-dependent flavonol glycoside accumulation in
Arabidopsis thaliana plants reveals MYB11-, MYB12- and MYB111-independent flavonol
glycoside accumulation. New Phytologist. 2010;188(4):985-1000.
21.   Gonzalez A, Zhao M, Leavitt JM, Lloyd AM. Regulation of the anthocyanin biosynthetic
pathway by the TTG1/bHLH/Myb transcriptional complex in Arabidopsis seedlings. The Plant
Journal. 2008;53(5):814-27.
22.   Ramsay NA, Glover BJ. MYB-bHLH-WD40 protein complex and the evolution of
cellular diversity. Trends in Plant Science. 2005;10(2):63-70.
23.   Dubos C, Stracke R, Grotewold E, Weisshaar B, Martin C, Lepiniec L. MYB
transcription factors in Arabidopsis. Trends in Plant Science. 2010;15(10):573-81.
24.   Borevitz JO, Xia YJ, Blount J, Dixon RA, Lamb C. Activation tagging identifies a
conserved MYB regulator of phenylpropanoid biosynthesis. The Plant Cell. 2000;12(12):2383-
93.
25.   Paz-Ares J, Ghosal D, Wienand U, Peterson PA, Saedler H. The regulatory c1 locus of
Zea mays encodes a protein with homology to myb proto-oncogene products and with
structural similarities to transcriptional activators. The EMBO Journal. 1987;6(12):3553-8.
26.   Appelhagen I, Jahns O, Bartelniewoehner L, Sagasser M, Weisshaar B, Stracke R.
Leucoanthocyanidin Dioxygenase in Arabidopsis thaliana: characterization of mutant alleles
and regulation by MYB-BHLH-TTG1 transcription factor complexes. Gene. 2011;484(1-2):61-
8.
27.   Zimmermann IM, Heim MA, Weisshaar B, Uhrig JF. Comprehensive identification of
Arabidopsis thaliana MYB transcription factors interacting with R/B-like BHLH proteins. The
Plant Journal. 2004;40:22-34.
28.   Lloyd AM, Walbot V, Davis RW. *Arabidopsis* and *Nicotiana* anthocyanin production
activated by maize regulators R and C1. Science. 1992;258:1773-5.

29.   Carey CC, Strahle JT, Selinger DA, Chandler VL. Mutations in the pale aleurone color1
Regulatory Gene of the Zea mays Anthocyanin Pathway Have Distinct Phenotypes Relative
to the Functionally Similar TRANSPARENT TESTA GLABRA1 Gene in Arabidopsis thaliana.
The Plant Cell. 2004;16:450-64.
30.   Pucker B, Pandey A, Weisshaar B, Stracke R. The R2R3-MYB gene family in banana
(*Musa acuminata*): Genome-wide identification, classification and expression patterns. PLoS
One. 2020;15(10):e0239275.
31.   Tak H, Negi S, Ganapathi TR. Overexpression of MusaMYB31, a R2R3 type MYB
transcription factor gene indicate its role as a negative regulator of lignin biosynthesis in
banana. PLoS ONE. 2017;12(2):e0172695.
32.   Deng GM, Zhang S, Yang QS, Gao HJ, Sheng O, Bi FC, et al. MaMYB4, an R2R3-
MYB Repressor Transcription Factor, Negatively Regulates the Biosynthesis of Anthocyanin
in Banana. Frontiers in Plant Science. 2021;11:600704.
33.   Rajput R, Naik J, Stracke R, Pandey A. Interplay between R2R3 MYB-type activators
and repressors regulates proanthocyanidin biosynthesis in banana (Musa acuminata). New
Phytologist. 2022.
34.   D´Hont A, Denoeud F, Aury JM, Baurens FC, Carreel F, Garsmeur O, et al. The banana
(Musa acuminata) genome and the evolution of monocotyledonous plants. Nature.
2012;488(7410):213-7.
35.   Martin G, Baurens FC, Droc G, Rouard M, Cenci A, Kilian A, et al. Improvement of the
banana "Musa acuminata" reference sequence using NGS data and semi-automated
bioinformatics methods. BMC Genetics. 2016;17(243).
36.   Schwinn KE, Ngo H, Kenel F, Brummell DA, Albert NW, McCallum JA, et al. The Onion
(Allium cepa L.) R2R3-MYB Gene MYB1 Regulates Anthocyanin Biosynthesis. Frontiers in
Plant Science. 2016;7.
37.   Shirley BW, Hanley S, Goodman HM. Effects of ionizing radiation on a plant genome:
analysis of two Arabidopsis transparent testa mutations. The Plant Cell. 1992;4:333-47.
38.   Pucker B, Ruckert C, Stracke R, Viehover P, Kalinowski J, Weisshaar B. Twenty-Five
Years of Propagation in Suspension Cell Culture Results in Substantial Alterations of the
Arabidopsis Thaliana Genome. Genes. 2019;10(9):671.
39.   Deluc L, Bogs J, Walker AR, Ferrier T, Decendit A, Merillon JM, et al. The transcription
factor VvMYB5b contributes to the regulation of anthocyanin and proanthocyanidin
biosynthesis in developing grape berries. Plant Physiology. 2008;147(4):2041-53.
40.   Wang N, Qu C, Jiang S, Chen Z, Xu H, Fang H, et al. The proanthocyanidin-specific
transcription factor MdMYBPA1 initiates anthocyanin synthesis under low temperature
conditions in red-fleshed apple. The Plant Journal. 2018;96(1):tpj.14013.
41.   Gunther CS, Dare AP, McGhie TK, Deng C, Lafferty DJ, Plunkett BJ, et al.
Spatiotemporal Modulation of Flavonoid Metabolism in Blueberries. Frontiers in Plant Science.
2020;11:545.
42.   Lafferty DJ, Espley RV, Deng CH, Gunther CS, Plunkett B, Turner JL, et al. Hierarchical
regulation of MYBPA1 by anthocyanin- and proanthocyanidin-related MYB proteins is
conserved in *Vaccinium* species. Journal of Experimental Botany. 2022;73(5):1344-56.
43.   Goff SA, Cone KC, Chandler VL. Functional analysis of the transcriptional activator
encoded by the maize *B* gene: evidence for a direct functional interaction between two classes
of regulatory proteins. Genes & Development. 1992;6:864-675.
44.   Quattrocchio F, Verweij W, Kroon A, Spelt C, Mol J, Koes R. PH4 of Petunia is an R2R3
MYB protein that activates vacuolar acidification through interactions with basic-helix-loop-
helix transcription factors of the anthocyanin pathway. The Plant Cell. 2006;18(5):1274-91.
45.   Petroni K, Tonelli C. Recent advances on the regulation of anthocyanin synthesis in
reproductive organs. Plant Science. 2011;181(3):219-29.
439