Three R2R3-MYB transcription factors from banana (*Musa* spp.) activate structural

anthocyanin biosynthesis genes as part of an MBW complex

Mareike Busche[1], Boas Pucker[2], Bernd Weisshaar[1], Ralf Stracke[1]*

[1] Genetics and Genomics of Plants, Faculty of Biology, Bielefeld University, 33615 Bielefeld,

Germany

[2] Institute of Plant Biology & Braunschweig Integrated Centre of Systems Biology (BRICS), TU

Braunschweig, 38106 Braunschweig, Germany

*Corresponding author:

ralf.stracke@uni-bielefeld.de

Abstract

Bananas are among the most popular fruits in the world and provide food security and

employment opportunities in several developing countries. An increased anthocyanin content

could enhance the health promoting properties of banana fruits. The biosynthesis of

anthocyanins is largely regulated at the transcriptional level. However, little is known about

transcriptional activation of anthocyanin biosynthesis in banana. We analysed the regulatory

activity of three *Musa*MYBs predicted to transcriptionally regulate anthocyanin biosynthesis in

banana. Co-transfection experiments in *A. thaliana* protoplasts showed that *Musa*MYB-α,

*Musa*MYB-β and *Musa*MYB-γ function as components of a transcription factor complex with a

bHLH and WD40 protein, called MBW complex, resulting in the activation of the *anthocyanin*

synthase and *dihydroflavonol 4-reductase* promoters from *A. thaliana*. The activation potential

of *Musa*MYB-α, *Musa*MYB-β and *Musa*MYB-γ increased when combined with the monocot

bHLH *Zm*R instead of the dicot *At*EGL3. This work paves the path towards decoding the MBW

complex-mediated transcriptional activation of anthocyanin biosynthesis in banana. Moreover,

it facilitates research towards an elevated anthocyanin content in banana fruits.

Keywords: *Musa acuminata*, banana, specialised metabolites, transcription factor, flavonoid

biosynthesis, R2R3-MYB

Introduction

Bananas (*Musa*) are monocotyledonous, perennial plants which are grown in many tropical and subtropical countries. They are one of the most important food crops, particularly in the developing world (1). While the sweet fruits of dessert bananas are popular in Europe and Northern America, plantains or cooking bananas are commonly eaten as a staple food in Africa and Latin America where they provide food security, as well as employment opportunities (2). Furthermore, banana fruits are rich in several health promoting minerals and beneficial phytochemicals such as vitamins and flavonoids (3).

Flavonoids are a major group of plant specialised metabolites which share a basic structure of two aromatic C6-rings connected by a heterocyclic ring (4). Reorganisations and modifications e.g. oxidation, glycosylation, acylation, and methylation of the carbon skeleton create a versatile group comprising more than 9,000 different flavonoid derivatives (5). Consequently, flavonoids do not only contribute to the nutritional value of fruits, but also play important roles in manifold processes. While the group of red coloured anthocyanin pigments attracts animals for pollination and dispersal of seeds by colouring flowers and fruits, other flavonoids protect plants against UV-B irradiation or increase plant fertility (6-10). Flavonoids from many species have been reported to have anti-pathogenic properties, this includes flavonoids from *Dianthus caryophyllus* which exhibit antifungal activity against the plant's major pest *Fusarium oxysporum* f.sp. *dianthi* (11, 12). The tropical race 4 (TR4) of the banana Fusarium wilt (commonly known as 'panama disease') is caused by another *Fusarium* subspecies called *Fusarium oxysporum* f. sp. *cubense* (Foc) and threatens the global banana production (13). Transcriptome analysis of susceptible and resistant banana cultivars infected by Foc TR4 revealed an increased transcription of flavonoid biosynthesis related genes in the resistant cultivar, suggesting an involvement of flavonoids in the defence against Foc TR4 (14).

Flavonoid biosynthesis is one of the best characterised pathways of the secondary metabolism and has been extensively studied in many plant species (15). In banana, several flavonoid biosynthesis related enzymes have been identified and characterised (16, 17). Regulation of structural genes on a transcriptional level allows a specific response to environmental

influences as well as development and organ specific expression (summarised in18). MYB

transcription factors are common transcriptional regulators of flavonoid biosynthesis. While

some MYBs act independently, others interact with basic helix-loop-helix (bHLH) and WD40

proteins to form a protein complex called MBW complex (19). MYB proteins are present in all

eukaryotes and characterised by highly conserved DNA-binding domains (20). These MYB

domains consist of up to three imperfect amino acid repeat sequences, based on which they

are classified. R2R3-MYBs are the most abundant class of plant MYBs and reveal versatile

functions in plant-specific processes (21). Besides core- and specialised metabolism they are

also involved in cell fate and -identity definition, developmental processes and the response to

biotic and abiotic stresses (21). Well-known R2R3-MYBs which act as activators of

anthocyanin biosynthesis include C1 (COLOURED ALEURONE1) from *Zea mays* (*Z. mays*),

as well as PAP1 (PRODUCTION OF ANTHOCYANIN PIGMENT1/MYB75) and PAP2

(MYB90) from *Arabidopsis thaliana* (*A. thaliana*) (22, 23). They act as part of an MBW complex

and control the promoters of the anthocyanin biosynthesis related structural genes as for

example *ANTHOCYANIDIN SYNTHASE* (*ANS*) and *DIHYDROFLAVONOL 4-REDUCTASE*

(*DFR*) (24-27).

In banana (*Musa acuminata, M. acuminata*), 285 R2R3-MYB proteins, including several

putative regulators of flavonoid biosynthesis, have been identified in a genome-wide study

(28). Furthermore, MYB31, MYB4 and MYBPR1 - MYBPR4 have been identified as negative

regulators of flavonoid biosynthesis in banana (29, 30). Despite the recent identification of two

proanthocyanidin biosynthesis activating R2R3-MYBs (31), little functional data is available on

positive regulators (activators) of flavonoid and in particular anthocyanin biosynthesis in

M. acuminata.

Here, we describe the regulatory properties of three *Musa*MYBs, named *Musa*MYB-α,

*Musa*MYB-β and *Musa*MYB-γ, with a possible role in the regulation of anthocyanin

biosynthesis. Regulatory activity was investigated by *in planta* complementation experiments

of the anthocyanin deficient *A. thaliana* regulatory *pap1/pap2* mutant and co-transfection

experiments in *A. thaliana* protoplasts. Our results show that *Musa*MYB-α, *Musa*MYB-β and

*Musa*MYB-γ are able to activate the promoters of *AtANS* and *AtDFR* as part of an MBW

complex. Moreover, we show that the activation potential of *Musa*MYB-α, *Musa*MYB-β and

*Musa*MYB-γ is increased when combined with the monocot bHLH *Zm*R instead of the dicot

bHLH protein ENHANCER OF GLABRA 3 (*At*EGL3).

Methods

Cladistic analysis

The approximately maximum-likelihood trees of plant MYB domain sequences and bHLH

protein sequences were constructed as described previously (32): Multiple protein sequence

alignments were created using MAFFT v7 (33) with default settings and cleaned with pxclsq

(34). The tree was constructed with FastTree v2.1.10 using the WAG+CAT model (35) and

visualised with interactive tree of life (36). GenBank IDs of protein sequences used for the

construction of the tree are given in Supplementary Table S2.

Plant material

Banana plants (*M. acuminata* (AAA Group) 'Grand Naine') for RNA extraction were grown in

the field in Lucknow, India. *Musa* gene annotation identifiers refer to the study of Martin *et al.*

(37). *A. thaliana* Columbia-0 (Col-0, NASC ID N1092) and Nössen-0 (Nö-0, NASC ID N3081)

accessions were used as wildtype controls corresponding to the *pap1/pap2* (*pap1*: Nö-0

background, PST16228; *pap2*: Col-0 background, SALK093731; (24)) double mutant, used for

complementation assays. Hypocotyl-derived, cultured *A. thaliana* At7 cells (38) for co-

transfection experiments were grown in MS medium (39) and maintained at 26 °C in the dark

on a rotary shaker (105 rpm). Cells were subcultured weekly and cultures for protoplast

isolation were inoculated 5 days prior to harvest.

cDNA synthesis and molecular cloning

RNA from Grand Naine peel was isolated according to a protocol from Asif *et al*. (40). cDNA

synthesis was performed from 1 µg total RNA using the ProtoScript First Strand cDNA

Synthesis Kit (New England Biolabs, NEB). Full length coding sequences (CDSs) were

amplified using Q5® High-Fidelity DNA polymerase (NEB) and gene-specific primers

(Supplementary Table S1). To generate Gateway Entry plasmids, PCR products were

recombined into pDONR™/Zeo (Invitrogen) with BP clonase (Invitrogen) according to the

manufacturer's protocol. Full length CDSs were introduced into the binary 2x35S

overexpression vector pLEELA (41) and the 2x35S overexpression vector pBTdest (42) using

Gateway LR reaction (Invitrogen). All plasmids were verified by Sanger sequencing.

Agrobacterium tumefaciens*-mediated transformation of *A. thaliana

Agrobacterium tumefaciens (GV3101::pMP90RK (43)) mediated gene transfer was carried out

using the floral dip method (44). T-DNA integration from pLEELA-based plasmids into the

A. thaliana genome was screened by BASTA selection and PCR-based genotyping (refer to

Supplementary Figure 1 and Supplementary Table 1 for details).

Determination of anthocyanin content

A. thaliana seedlings were grown with 16 h of light illumination per day on 0.5x MS agar plates

with 4 % sucrose to induce anthocyanin production. Anthocyanin content in a bulk of six-day-

old seedlings was quantified as described by Mehrtens *et al.* (45). All samples were measured

in three independent biological replicates. Error bars indicate standard deviation of the average

(mean) anthocyanin content.

Co-transfection experiments in *A. thaliana* protoplasts

Cell culture, protoplast isolation, PEG-mediated DNA transfer into protoplasts and

determination of standardised GUS activity were performed as described by Hartmann *et al*.

(46). As an adaption we used 1.3 % (w/v) Cellulase Onozuka RS (Duchefa) and 0.8 % (w/v)

Macerozyme R-10 (Duchefa) in 240 mM $CaCl_2$ as the enzyme solution for protoplast

preparation and 30 µg of plasmid DNA for the PEG-mediated DNA transfer into protoplasts.

This comprised 10 µg of the promoter-GUS reporter construct, 0.5 µg of each

2x35S:transcription factor effector construct, 5 µg of the standardisation plasmid (pBT8 UBI-

LUCm3), and an inactive luciferase expression vector (pBTΔLUC) to fill up to the amount

of 30 µg of plasmid DNA. GUS-fused promoters, 520 bp *proAtDFR* and 1.95 kb *proAtANS,* as

well as expression constructs for *At*PAP1 (*At*MYB75), *At*EGL3 (*At*bHLH2), *At*TTG1 and *Zm*C1

were available from previous studies (24, 25, 42). A *Zm*R entry clone (47) was kindly provided

by Martin Hülskamp. Statistical analysis was performed using the Mann-Whitney *U* test (48).

Results and Discussion

To analyse the regulatory properties of three *Musa*MYBs which have been previously assigned

with a possible role in positive regulation of anthocyanin biosynthesis (Ma06_g05960 or

MusaMYB-α, Ma09_g27990 or *MusaMYB-β*, Ma10_g17650 or *MusaMYB-γ*), we tried to

amplify the corresponding coding sequences (CDSs) on a template collection containing cDNA

from different banana samples. The CDSs of all three *MusaMYB*s were successfully amplified

on cDNA derived from peel tissue.

In a first approach, we performed a complementation assay with the regulatory *A. thaliana*

pap1/pap2 double mutant which cannot produce anthocyanins in the seedling (Figure 1).

Seedlings were grown on anthocyanin synthesis-inducing media to analyse the ability of

2x35S-driven *MusaMYB*s to complement the *pap1/pap2* anthocyanin deficiency. While wild-

type seedlings accumulated high amounts of red anthocyanin pigments, *pap1/pap2* seedlings

did not. Even though *MusaMYB-α*, *MusaMYB-β* or *MusaMYB-γ* were successfully expressed

in the transgenic seedlings (Supplementary Figure S1), the anthocyanin level in *pap1/pap2*

plants expressing *MusaMYB-α*, *MusaMYB-β* or *MusaMYB-γ* did not differ from the double

mutant. Accordingly, *MusaMYB-α*, *MusaMYB-β* and *MusaMYB-γ* do not seem to be able to

complement the mutant phenotype and thus to regulate anthocyanin biosynthesis in *A. thaliana*

in combination with the *bHLH* and *WD40* genes expressed in *A. thaliana* seedlings.

Figure 1: *MusaMYB-α, MusaMYB-β* and *MusaMYB-γ* cannot complement the anthocyanin deficient phenotype of *A. thaliana pap1/pap2* mutant seedlings. (A) Representative pictures of anthocyanin accumulation in 6-day-old *MusaMYB*-expressing *pap1/pap2* seedlings. A representative plant per construct is displayed. (B) Photometrical measurement of the sucrose induced anthocyanin content in *pap1/pap2* seedlings expressing 2x35S-driven *MusaMYBs*. Col-0, Nö-0 (wildtypes) and *pap1/pap2* were used as controls. Error bars indicate the standard deviation of three biological replicates. The different numbers represent individual, independent, transgenic lines.

Lloyd *et al.* (26) analysed the *Z. mays* anthocyanin regulator C1 by generation of transgenic

A. thaliana overexpression lines. In their study, Lloyd *et al.* generated three independent

ZmC1-expressing *A. thaliana* lines that did not show an increased anthocyanin content

compared to wildtype. Similar results were obtained in transgenic tobacco. However, further

experiments suggested that *Zm*C1 needs to interact with the maize bHLH *Zm*R to activate

anthocyanin biosynthesis in the heterologous *A. thaliana* system (26). Phylogenetic analysis

(49) showed that the anthocyanin biosynthesis regulating R2R3-MYBs from several monocots

such as *Z. mays* and *Oryza sativa* (*O. sativa*) are part of a different phylogenetic clade than

the anthocyanin regulators from several dicots, such as *A. thaliana* or *Vitis vinifera* and further

monocots, including *Allium cepa* and *Lilium hybrida*. Experiments in snapdragon also showed

that the monocot *Ac*MYB1 can activate anthocyanin production in dicots (49). The phylogenetic

differences between anthocyanin biosynthesis activating R2R3-MYBs and the dependency of

*Zm*C1 on *Zm*R in dicots hint at an explanation for our observations. For example, the

*Musa*MYBs could depend on their endogenous or at least monocot bHLH for effective

activation of structural anthocyanin biosynthesis genes. To investigate the phylogenetic

differences between anthocyanin biosynthesis regulating R2R3-MYBs from different plant

species, an approximately maximum-likelihood tree was built (Figure 2A). The resulting tree,

which also contained R2R3-MYBs which activate other branches of flavonoid biosynthesis,

revealed two distinct clades of anthocyanin-related R2R3-MYBs (highlighted in red).

*Musa*MYB-α, *Musa*MYB-β and *Musa*MYB-γ form a clade with MYB10 from *Triticum aestivum*

and exhibit a close evolutionary relationship with *Zm*C1, *Os*C1 and anthocyanin regulating

MYBs from further monocots. In contrast, *At*PAP1 and *At*PAP2 fall into a second clade of

anthocyanin-related MYBs. Furthermore, differences between anthocyanin biosynthesis

regulating bHLHs were analysed in a second approximately maximum-likelihood tree (Figure

2B). The tree showed that monocot bHLH proteins involved in the regulation of anthocyanin

biosynthesis form a separate clade (highlighted in red). Both, *Zm*R and several putative

anthocyanin biosynthesis regulating bHLHs from banana fall into this clade. These

phylogenetic analyses show that anthocyanin biosynthesis regulating R2R3-MYB and bHLH

proteins from several monocot species seem to be distinct from other R2R3-MYB and bHLH

proteins regulating anthocyanin biosynthesis, for example from *A. thaliana*. These differences

could imply a dependency of *Musa*MYB-α, *Musa*MYB-β and *Musa*MYB-γ on a banana or other

monocot bHLH.

To follow up this idea, we further investigated the regulatory properties of the three *Musa*MYBs

performing co-transfection assays in At7 protoplasts with different bHLH proteins from

A. thaliana (*At*EGL3) or *Z. mays* (*Zm*R) and the *A. thaliana* WD40 protein TRANSPARENT

TESTA GLABRA 1 (*At*TTG1, Figure 3). Thereby, their potential to activate the promoters of

Figure 2: Rooted approximately maximum-likelihood trees of MYB (A) and bHLH (B) transcription factors. Circle sizes represent bootstrap values. Banana proteins are highlighted in blue.

AtANS and *AtDFR*, which are important structural genes of anthocyanin biosynthesis (summarised in 50), was analysed. While none of the *Musa*MYBs was able to independently

activate *proAtANS* or *proAtDFR*, which both contain conserved *cis*-regulatory elements, a

slight activation of *proAtDFR* was detectable when *Musa*MYB-α or *Musa*MYB-γ were

combined with *At*EGL3 and *At*TTG1. In combination with *At*EGL3 and *At*TTG1, *Musa*MYB-γ

showed the strongest activation of *proAtDFR* and was also able to activate *proAtANS*. When

combined with *Zm*R and *At*TTG1, the three MYBs *Musa*MYB-α, *Musa*MYB-β, and *Musa*MYB-γ

exhibited a significant activation potential on *proAtDFR* and *proAtANS*. All three *Musa*MYBs

showed a higher activation potential in an MBW complex with *Zm*R and *At*TTG1 than in

combination with *At*EGL3 and *At*TTG1.

These results display that *Musa*MYB-α, *Musa*MYB-β and *Musa*MYB-γ are able to activate

proAtANS and *proAtDFR* as part of an MBW complex and *Musa*MYB-γ shows the strongest

activation potential. Furthermore, *Musa*MYB-α, *Musa*MYB-β and *Musa*MYB-γ show a higher

activation potential when combined with the monocot bHLH *Zm*R instead of the dicot bHLH

*At*EGL3.

The previously mentioned phylogenetic differences could be an explanation for the observed

higher activation potentials of the analysed *Musa*MYBs in combination with *Zm*R and *At*TTG1

instead of *At*EGL3 and *At*TTG1. It is conceivable that these differences impede the interaction

between the tested *Musa*MYBs and the dicot bHLH *At*EGL3. As *Musa*MYB-γ shows the

highest activation potential and is also able to activate *proAtANS* in combination with *At*EGL3

or *Zm*R, differences between the three *Musa*MYBs that affect the interaction with the *At*EGL3

are likely and should be further studied. Interestingly, *Musa*MYB-γ did not complement the

anthocyanin deficient phenotype of *A. thaliana pap1/pap2* mutant seedlings, but was able to

activate *proAtANS* and *proAtDFR* when combined with *At*EGL3 in co-transfection experiments

in *A. thaliana* protoplasts. Even though previous studies showed *AtEGL3* expression in

A. thaliana seedlings (19), it is possible that the level was too low to activate anthocyanin

biosynthesis in combination with *Musa*MYB-γ. Moreover, the At7 cell line used for co-

transfection experiments was established more than 25 years ago. This long time of

propagation in suspension cell culture caused a variety of genomic and transcriptomic changes

(51). These changes could also explain differences between the analysis in seedlings and cell

culture.

Figure 3: *Musa*MYB-α, *Musa*MYB-β and *Musa*MYB-γ can activate *proAtDFR* and
***proAtANS* as part of an MBW complex**. The ability of *Musa*MYBs to activate *proAtDFR-GUS*
and *proAtANS-GUS* reporter constructs in combination with different bHLH proteins (*At*EGL3,
*Zm*R) and a WDR (*At*TTG1) was analysed by co-transfection in *A. thaliana* At7 protoplasts.
The relative promoter activity refers to the measured GUS reporter enzyme activity. Promoter
activity is given relative to the values obtained for the *A. thaliana* MBW complex (*At*PAP1,
*At*EGL3, *At*TTG1). Error bars indicate the standard deviation of five independent biological
replicates. Statistical significance is indicated by asterisks which mark p-values < 0.05.

The expression profiles of *MusaMYB-α*, *MusaMYB-β* and *MusaMYB-γ* have previously been

analysed in embryogenic cell suspension, seedling, root and different developmental stages

of leaf, pulp and peel (28). The expression of *MusaMYB-α* was strongest in pulp

(developmental stage S2-S3), *MusaMYB-β* expression was relatively low in all samples and

MusaMYB-γ showed highest expression in seedlings and early developmental stages of pulp

(S1). This gene expression pattern could give a hint at organ specific *MusaMYB-α*,

MusaMYB-β and *MusaMYB-γ* activity in banana. Since flavonoid biosynthesis is largely

regulated at the transcriptional level, it would be particularly interesting to analyse the

expression levels of *MusaMYB-α*, *MusaMYB-β*, and *MusaMYB-γ* in anthocyanin rich organs

such as bract or pseudostem to relate anthocyanin content to *MusaMYB* transcript abundance.

Deng *et al*. (30) performed an expression analysis of flavonoid biosynthesis related genes

using leaves of banana plants overexpressing the anthocyanin repressor *MusaMYB4*. They

found that the expression of *MusaMYB-α* and *MusaMYB-γ* is decreased along with *MusaDFR*

and *MusaANS* in plants overexpressing *Musa*MYB4, compared to wildtype. This data supports

a proposed functionality of *Musa*MYB-α and *Musa*MYB-γ as transcriptional activators of

anthocyanin biosynthesis in banana, as *MusaMYB4* could cause a feedback regulation of the

positive regulators of anthocyanin biosynthesis, of *MusaMYB-α* and *MusaMYB-γ*.

Very recently, Rajput *et al*. (31) showed that *Musa*MYB-γ, which they refer to as MaMYBPA2,

can activate the banana *ANS, ANR* and LAR promoters. Moreover, they showed that

*Musa*MYB-γ/MaMYBPA2 can partially rescue the proanthocyanin deficiency of the *A. thaliana*

tt2-1 mutant. The reported ability of *Musa*MYB-γ/MaMYBPA2 to activate *proMusaANS*

supports the proposed role of *Musa*MYB-γ/MaMYBPA2 in the regulation of anthocyanin

biosynthesis. However, the partial complementation of the proanthocyanin deficient phenotype

of the *A. thaliana tt2-1* R2R3-MYB mutant, as well as the ability to activate the banana *ANR*

and *LAR* promoters, shows an additional role in the regulation of proanthocyanidin

biosynthesis. Such dual roles in the regulation of flavonoid biosynthesis have previously been

suggested for R2R3-MYB transcription factors from grape and apple (52, 53). In several

species, including *A. thaliana, Z. mays* and *Petunia hybrida*, anthocyanin biosynthesis is

regulated by an MBW complex (summarised in 54). Our co-transfection assays showed that

*Musa*MYB-α, *Musa*MYB-β and *Musa*MYB-γ are able to activate *proAtANS* and *proAtDFR* as

part of an MBW complex with *Zm*R and *At*TTG1 *in planta*. As presented in a previous study by

Pucker *et al.* (28), it is known that *Musa*MYB-α, *Musa*MYB-β, and *Musa*MYB-γ all contain a

bHLH-binding consensus motif (25). Thus, the regulation of anthocyanin biosynthesis by

*Musa*MYB-α, *Musa*MYB-β, and *Musa*MYB-γ likely depends on bHLH and WD40 proteins.

Based on our tree (Figure 2B), the bHLH encoding genes Ma05_g23010, Ma11_g19640,

Ma11_g16060, Ma08_g11190 and Ma11_g03740 could be suitable candidates for further

studies on the transcriptional activation of anthocyanin biosynthesis by MBW complexes.

Moreover, future analyses should additionally include *Musa*TTG1 to identify a whole functional

MBW complex in banana and to find out if the use of endogenous WD40 protein further

improves the activation potential of the MBW complex.

Resumed, the presented results show that *Musa*MYB-α, *Musa*MYB-β, and *Musa*MYB-γ have

the capacity to transcriptionally activate structural anthocyanin biosynthesis genes in an MBW

complex with a suitable bHLH partner. This is a step towards decoding the MBW complex-

mediated transcriptional activation of flavonoid biosynthesis in banana. Moreover, it is a basis

for further research towards increased anthocyanin production in banana which could promote

fruit quality and disease resistance.

Acknowledgements

We are grateful to Melanie Kuhlmann and Janina Grabowski for their excellent technical

assistance and to Andrea Voigt for her competent help in the greenhouse. We thank Martin

Hülskamp for providing the *Zm*R entry clone plasmid and Ashutosh Pandey for providing the

banana cDNAs.

Author Contributions

MB, BP and RS planned the experiments. MB performed the experiments and analysed the

data. RS and BW supervised the project. MB wrote the initial draft. RS and BP revised the

manuscript.

Abbreviations

ANS   anthocyanidin synthase

A. thaliana   *Arabidopsis thaliana*

bHLH   basic helix-loop-helix

C1   COLOURED ALEURONE1

DFR   dihydroflavonol 4-reductase

EGL3   ENHANCER OF GLABRA 3

Foc   *Fusarium oxysporum* f. sp. *cubense*

M. acuminata *Musa acuminata*

O. sativa   *Oryza sativa*

PAP   production of anthocyanin pigment

TR4   tropical race 4

TTG1   TRANSPARENT TESTA GLABRA 1

Z. mays   *Zea mays*

Supplementary information

Figure S1: Expression analysis of *MusaMYB-α, MusaMYB-β and MusaMYB-γ* in *A. thaliana*

pap1/pap2 seedlings.

Table S1: Oligonucleotide primers used in this work.

Table S2: IDs of protein sequences used for the construction of the phylogenetic tree

References

1.   Scott GJ. A review of root, tuber and banana crops in developing countries: past,
present and future. International Journal of Food Science and Technology. 2021;56(3):1093-
114.

2. Arias P, Dankers C, Liu P, Pilkauskas P. The World Banana Economy, 1985-2002 2003
[http://www.fao.org/3/y5102e/ye00.htm]. Available from:
http://www.fao.org/3/y5102e/y5102e00.htm.
3. Singh B, Singh JP, Kaur A, Singh N. Bioactive compounds in banana and their
associated health benefits - A review. Food Chemistry. 2016;206:1-11.
4. Winkel-Shirley B. Flavonoid biosynthesis. A colorful model for genetics, biochemistry,
cell biology, and biotechnology. Plant Physiology. 2001;126:485-93.
5. Wen W, Alseekh S, Fernie AR. Conservation and diversification of flavonoid
metabolism in the plant kingdom. Current Opinion in Plant Biology. 2020;55:100-8.
6. Ishikura N, Yoshitama K. Anthocyanin-Flavonol Co-pigmentation in Blue Seed Coats
of Ophiopogon jaburan. Journal of Plant Physiology. 1984;115(2):171-5.
7. Mo Y, Nagel C, Taylor LP. Biochemical complementation of chalcone synthase mutants
defines a role for flavonols in functional pollen. Proceedings of the National Academy of
Sciences of the United States of America. 1992;89:7213-7.
8. Li J, Ou-Lee TM, Raba R, Amundson RG, Last RL. Arabidopsis Flavonoid Mutants Are
Hypersensitive to UV-B Irradiation. The Plant Cell. 1993;5(2):171-9.
9. Gronquist M, Bezzerides A, Attygalle A, Meinwald J, Eisner M, Eisner T. Attractive and
defensive functions of the ultraviolet pigments of a flower *(Hypericum calycinum)*. Proceedings
of the National Academie of Sciences of the United States of America. 2001;98(24):13745-50.
10. Grotewold E. The Science of Flavonoids. Grotewold E, editor. Columbus: The Ohio
State University; 2006. 273 p.
11. Galeotti F, Barile E, Curir P, Dolci M, Lanzotti V. Flavonoids from carnation (Dianthus
caryophyllus) and their antifungal activity. Phytochemistry Letters. 2008;1(1):44-8.
12. Mierziak J, Kostyn K, Kulma A. Flavonoids as Important Molecules of Plant Interactions
with the Environment. Molecules. 2014;19(10):16240-65.
13. Dita M, Barquero M, Heck D, Mizubuti ESG, Staver CP. Fusarium Wilt of Banana:
Current Knowledge on Epidemiology and Research Needs Toward Sustainable Disease
Management. Frontiers in Plant Science. 2018;9.
14. Sun J, Zhang J, Fang H, Peng L, Wei S, Li C, et al. Comparative transcriptome analysis
reveals resistance-related genes and pathways in *Musa acuminata* banana 'Guijiao 9' in
response to Fusarium wilt. Plant Physiology and Biochemistry. 2019;141:83-94.
15. Liu WX, Feng Y, Yu SH, Fan ZQ, Li XL, Li JY, et al. The Flavonoid Biosynthesis Network
in Plants. International Journal of Molecular Sciences. 2021;22(23).
16. Pandey A, Alok A, Lakhwani D, Singh J, Asif MH, Trivedi PK. Genome-wide Expression
Analysis and Metabolite Profiling Elucidate Transcriptional Regulation of Flavonoid
Biosynthesis and Modulation under Abiotic Stresses in Banana. Scientific Reports. 2016;6.
17. Busche M, Acatay C, Martens S, Weisshaar B, Stracke R. Functional Characterisation
of Banana (Musa spp.) 2-Oxoglutarate-Dependent Dioxygenases Involved in Flavonoid
Biosynthesis. Frontiers in Plant Science. 2021;12:701780.
18. Koes R, Verweij W, Quattrocchio F. Flavonoids: a colorful model for the regulation and
evolution of biochemical pathways. Trends in Plant Science. 2005;10(5):236-42.
19. Gonzalez A, Zhao M, Leavitt JM, Lloyd AM. Regulation of the anthocyanin biosynthetic
pathway by the TTG1/bHLH/Myb transcriptional complex in Arabidopsis seedlings. The Plant
Journal. 2008;53(5):814-27.
20. Ramsay NA, Glover BJ. MYB-bHLH-WD40 protein complex and the evolution of
cellular diversity. Trends in Plant Science. 2005;10(2):63-70.
21. Dubos C, Stracke R, Grotewold E, Weisshaar B, Martin C, Lepiniec L. MYB
transcription factors in Arabidopsis. Trends in Plant Science. 2010;15(10):573-81.
22. Borevitz JO, Xia YJ, Blount J, Dixon RA, Lamb C. Activation tagging identifies a
conserved MYB regulator of phenylpropanoid biosynthesis. The Plant Cell. 2000;12(12):2383-
93.
23. Paz-Ares J, Ghosal D, Wienand U, Peterson PA, Saedler H. The regulatory c1 locus of
Zea mays encodes a protein with homology to myb proto-oncogene products and with
structural similarities to transcriptional activators. The EMBO Journal. 1987;6(12):3553-8.

24. Appelhagen I, Jahns O, Bartelniewoehner L, Sagasser M, Weisshaar B, Stracke R.
Leucoanthocyanidin Dioxygenase in Arabidopsis thaliana: characterization of mutant alleles
and regulation by MYB-BHLH-TTG1 transcription factor complexes. Gene. 2011;484(1-2):61-
8.
25. Zimmermann IM, Heim MA, Weisshaar B, Uhrig JF. Comprehensive identification of
Arabidopsis thaliana MYB transcription factors interacting with R/B-like BHLH proteins. The
Plant Journal. 2004;40:22-34.
26. Lloyd AM, Walbot V, Davis RW. *Arabidopsis* and *Nicotiana* anthocyanin production
activated by maize regulators R and C1. Science. 1992;258:1773-5.
27. Carey CC, Strahle JT, Selinger DA, Chandler VL. Mutations in the pale aleurone color1
Regulatory Gene of the Zea mays Anthocyanin Pathway Have Distinct Phenotypes Relative
to the Functionally Similar TRANSPARENT TESTA GLABRA1 Gene in Arabidopsis thaliana.
The Plant Cell. 2004;16:450-64.
28. Pucker B, Pandey A, Weisshaar B, Stracke R. The R2R3-MYB gene family in banana
(*Musa acuminata*): Genome-wide identification, classification and expression patterns. PLoS
One. 2020;15(10):e0239275.
29. Tak H, Negi S, Ganapathi TR. Overexpression of MusaMYB31, a R2R3 type MYB
transcription factor gene indicate its role as a negative regulator of lignin biosynthesis in
banana. PLoS ONE. 2017;12(2):e0172695.
30. Deng GM, Zhang S, Yang QS, Gao HJ, Sheng O, Bi FC, et al. MaMYB4, an R2R3-
MYB Repressor Transcription Factor, Negatively Regulates the Biosynthesis of Anthocyanin
in Banana. Frontiers in Plant Science. 2021;11:600704.
31. Rajput R, Naik J, Stracke R, Pandey A. Interplay between R2R3-MYB-type activators
and repressors regulates proanthocyanidin biosynthesis in banana (*Musa acuminata*). New
Phytologist. 2022.
32. Pucker B, Reiher F, Schilbert HM. Automatic Identification of Players in the Flavonoid
Biosynthesis with Application on the Biomedicinal Plant *Croton tiglium*. Plants. 2020;9(9).
33. Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Molecular Biology and Evolution. 2013;30(4):772-
80.
34. Brown JW, Walker JF, Smith SA. Phyx: phylogenetic tools for unix. Bioinformatics.
2017;33(12):1886-8.
35. Price MN, Dehal PS, Arkin AP. FastTree 2--approximately maximum-likelihood trees
for large alignments. PLoS One. 2010;5(3):e9490.
36. Letunic I, Bork P. Interactive Tree Of Life (iTOL) v4: recent updates and new
developments. Nucleic Acids Research. 2019;47(W1):W256-W9.
37. Martin G, Baurens FC, Droc G, Rouard M, Cenci A, Kilian A, et al. Improvement of the
banana "Musa acuminata" reference sequence using NGS data and semi-automated
bioinformatics methods. BMC Genetics. 2016;17(243).
38. Trezzini GF, Horrichs A, Somssich IE. Isolation of putative defense-related genes from
Arabidopsis thaliana and expression in fungal elicitor-treated cells. Plant Molecular Biology.
1993;21:385-9.
39. Murashige T, Skoog F. A revised Medium for rapid growth and bioassays with tobacco
tissue cultures. Plant Physiology. 1962;15:473-97.
40. Asif MH, Dhawan P, Nath P. A simple procedure for the isolation of high quality RNA
from ripening banana fruit. Plant Molecular Biology Reporter. 2000;18:109-15.
41. Jakoby M, Wang HY, Reidt W, Weisshaar B, Bauer P. FRU (BHLH029) is required for
induction of iron mobilization genes in Arabidopsis thaliana. FEBS Letters. 2004;577(3):528-
34.
42. Baudry A, Heim MA, Dubreucq B, Caboche M, Weisshaar B, Lepiniec L. TT2, TT8, and
TTG1 synergistically specify the expression of *BANYULS* and proanthocyanidin biosynthesis
in *Arabidopsis thaliana*. The Plant Journal. 2004;39(3):366-80.
43. Koncz C, Schell J. The promoter of TL-DNA gene 5 controls the tissue-specific
expression of chimaeric genes carried by a novel type of Agrobacterium binary vector.
Molecular Genetics and Genomics. 1986;204:383-96.

44.   Clough SJ, Bent AF. Floral dip: a simplified method for Agrobacterium-mediated
transformation of *Arabidopsis thaliana*. The Plant Journal. 1998;16(6):735-43.
45.   Mehrtens F, Kranz H, Bednarek P, Weisshaar B. The Arabidopsis transcription factor
MYB12 is a flavonol-specific regulator of phenylpropanoid biosynthesis. Plant Physiology.
2005;138(2):1083-96.
46.   Hartmann U, Valentine WJ, Christie JM, Hays J, Jenkins GI, Weisshaar B. Identification
of UV/blue light-response elements in the *Arabidopsis thaliana* chalcone synthase promoter
using a homologous protoplast transient expression system. Plant Molecular Biology.
1998;36:741-54.
47.   Zhang B, Chopra D, Schrader A, Hulskamp M. Evolutionary comparison of competitive
protein-complex formation of MYB, bHLH, and WDR proteins in plants. Journal of Experimental
Botany. 2019;70(12):3197-209.
48.   Mann HB, Whitney DR. On a Test of Whether One of Two Random Variables Is
Stochastically Larger than the Other. The Annals of Mathematical Statistics. 1947;18(1):50-60.
49.   Schwinn KE, Ngo H, Kenel F, Brummell DA, Albert NW, McCallum JA, et al. The Onion
(Allium cepa L.) R2R3-MYB Gene MYB1 Regulates Anthocyanin Biosynthesis. Frontiers in
Plant Science. 2016;7.
50.   Martens S, Preuß A, Matern U. Multifunctional flavonoid dioxygenases: Flavonol and
anthocyanin biosynthesis in Arabidopsis thaliana L. Phytochemistry. 2010;71(10):1040-9.
51.   Pucker B, Ruckert C, Stracke R, Viehover P, Kalinowski J, Weisshaar B. Twenty-Five
Years of Propagation in Suspension Cell Culture Results in Substantial Alterations of the
Arabidopsis Thaliana Genome. Genes. 2019;10(9):671.
52.   Deluc L, Bogs J, Walker AR, Ferrier T, Decendit A, Merillon JM, et al. The transcription
factor VvMYB5b contributes to the regulation of anthocyanin and proanthocyanidin
biosynthesis in developing grape berries. Plant Physiology. 2008;147(4):2041-53.
53.   Wang N, Qu C, Jiang S, Chen Z, Xu H, Fang H, et al. The proanthocyanidin-specific
transcription factor MdMYBPA1 initiates anthocyanin synthesis under low temperature
conditions in red-fleshed apple. The Plant Journal. 2018;96(1):tpj.14013.
54.   Petroni K, Tonelli C. Recent advances on the regulation of anthocyanin synthesis in
reproductive organs. Plant Science. 2011;181(3):219-29.
482