The role of calcium, Akt and ERK signaling in cadmium-induced hair cell death

Jennifer Galdieri[a], Chloe Adams[a], María Padilla[b], Tamara M. Stawicki[a,*]

[a] Neuroscience Program, Lafayette College, Easton, Pennsylvania, USA 18042

[b] Biology Department, Lafayette College, Easton, Pennsylvania, USA 18042

*Corresponding author:
Tamara Stawicki
stawickt@lafayette.edu
350 Hamilton St
Easton, PA 18042

ABSTRACT

Exposure to heavy metals has been shown to cause damage to a variety of different tissues and cell types including hair cells, the sensory cells of our inner ears responsible for hearing and balance. Elevated levels of one such metal, cadmium, have been associated with hearing loss and shown to cause hair cell death in multiple experimental models. While the mechanisms of cadmium-induced cell death have been extensively studied in other cell types they remain relatively unknown in hair cells. We have found that calcium signaling, which is known to play a role in cadmium-induced cell death in other cell types through calmodulin and CaMKII activation as well as IP3 receptor and mitochondrial calcium uniporter mediated calcium flow, does not appear to play a significant role in cadmium-induced hair cell death. While calmodulin inhibition can partially protect hair cells this may be due to impacts on mechanotransduction activity. Removal of extracellular calcium, and inhibiting CaMKII, the IP3 receptor and

the mitochondrial calcium uniporter all failed to protect against cadmium-induced hair

cell death. We also found cadmium treatment increased pAkt levels in hair cells and

pERK levels in supporting cells. This activation may be protective as inhibiting these

pathways enhances cadmium-induced hair cell death rather than protecting cells. Thus

cadmium-induced hair cell death appears distinct from cadmium-induced cell death in

other cell types where calcium, Akt and ERK signaling all promote cell death.

Keywords: Akt, ERK, cadmium, calcium signaling, hair cells, ototoxicity

INTRODUCTION

Hair cells, the sensory cells we use for hearing and balance, are extremely sensitive to insult, and as such disorders of these sensory systems are common. Recently environmental toxins such as heavy metals have gained attention as a potential cause of hearing loss (Choi and Kim, 2014; Schaal et al., 2017). One of these metals is cadmium. Cadmium can be found in some industrial run-off and individuals can be exposed to it in certain occupations, via smoking or via contaminated food (Faroon et al., 2012; IARC, 2012). Multiple studies have found a link between blood or urinary cadmium levels and hearing or balance impairment (Choi et al., 2012; Choi and Park, 2017; Dalton et al., 2020; Huh et al., 2018; Jung, 2019; Kang et al., 2018; Min et al., 2012; Shargorodsky et al., 2011; Wang et al., 2020). Studies have also found hearing loss and/or hair cell death in mammals and fish exposed to cadmium (Faucher et al., 2006; Kim et al., 2008; Liu et al., 2014; Low and Higgs, 2015; Montalbano et al., 2018; Ozcaglar et al., 2001; Schmid et al., 2020; Sonnack et al., 2015). There is evidence that this toxicity may stem from reactive oxygen species generation (ROS). Increased ROS production is seen following treatment with cadmium in the hair cell-like cell line HEI-OC1, and antioxidant treatment protects against cadmium-induced cell death in both HEI-OC1 cells and cochlear explants. Antioxidant treatment also protects against cadmium-induced increases in auditory-brain stem response thresholds in mice (Kim et al., 2008). However, the role of other potential cell-signaling pathways involved in cadmium-induced cell death has not been extensively studied in hair cells to date.

In addition to its association with hearing loss, cadmium has also been linked to a

number of other health conditions including cancer, osteoporosis, and dysfunction of the

cardiovascular system, kidneys, liver, nervous system and reproductive system.

(Bhardwaj et al., 2021; Ferraro et al., 2010; Gallagher et al., 2008; García-Esquinas et

al., 2014; Hyder et al., 2013; Mezynska and Brzóska, 2018; Ruczaj and Brzóska, 2022;

Tellez-Plaza et al., 2012). While mechanisms of cadmium toxicity in hair cells are still

relatively unknown, they have been more extensively studied in other cell types. Multiple

cell types show an increase in calcium in response to cadmium (Shih et al., 2005; Smith

et al., 1989; Xu et al., 2011; Yamagami et al., 1998; Yang et al., 2007; Yeh et al., 2009).

In some cases this increase appears due to the release of calcium from endoplasmic

reticulum (ER) stores, as it can be blocked by an inositol 1,4,5-triphosphate (IP3)

receptor antagonist but not by the removal of extracellular calcium (Liu et al., 2016;

Smith et al., 1989; Yamagami et al., 1998). Cadmium has also been shown to increase

IP3 levels in cells (Benters et al., 1997; Hague et al., 2000; Smith et al., 1989;

Yamagami et al., 1998). However, in other cases removing extracellular calcium or

inhibiting calcium channels can block the cadmium-induced calcium increase

suggesting extracellular calcium is contributing to this increase (Liu et al., 2018; Xu et

al., 2011; Yeh et al., 2009).

This increase in calcium appears to be important for cadmium-induced cell death

as calcium chelation and IP3 receptor inhibition have both been shown to protect

against cadmium-induced toxicity (Liu et al., 2016; L. Wang et al., 2009; Wang et al.,

2008; Yang et al., 2008, 2007; Yeh et al., 2009). Calcium may ultimately be inducing

cell death through disruptions in the mitochondria and the production of ROS as calcium

chelation can block cadmium-induced mitochondrial depolarization and increases in

ROS production (Xu et al., 2011; Yang et al., 2008, 2007; Yuan et al., 2013). Cadmium

is unable to increase ROS production when directly applied to isolated mitochondria

further supporting the role of calcium as a potential intermediary signal in this process

(Shih et al., 2005). Mitochondrial calcium also appears important in cadmium-induced

cell death as inhibition of the mitochondria calcium uniporter blocks cadmium-induced

cytochrome c release, mitochondrial depolarization and cell toxicity (Lee et al., 2005; Li

et al., 2003; Shih et al., 2005).

Downstream of calcium both calmodulin and calcium/calmodulin-dependent

protein kinase II (CaMKII) appear to be playing a role in cadmium-induced cell death.

Cadmium is able to directly activate calmodulin as well as increase calcium's activation

of calmodulin, and it is believed this may result in a toxic overactivation of this molecule

(Chao et al., 1984; Sutoo et al., 1990). Inhibition of both calmodulin and CaMKII can

protect cells against cadmium-induced cell death (Liu et al., 2018; Liu and Templeton,

2007).

Cadmium has also been shown to increase phosphorylated levels of Akt (pAkt),

and Extracellular Receptor Kinase (pERK) in vitro in a calcium-dependent manner

(Jiang et al., 2015; Xu et al., 2011). Increased pAkt levels have also been seen in the

brain in vivo in response to cadmium (Chen et al., 2014). The calcium-dependent

activation of these pathways appears to be through the activation of calmodulin and

CaMKII as inhabiting either of these molecules can block the cadmium-dependent

increase in pAkt and pERK (Chen et al., 2011; Fujiki et al., 2013; Liu et al., 2018; Xiao

et al., 2009; Xu et al., 2011). Akt and ERK activation appears to be playing a role in the

cell death process as inhibitors of both signaling pathways have been shown to protect

against cadmium-induced cell toxicity (Chen et al., 2008; Iryo et al., 2000; Jiang et al.,

2014, 2015; Y. K. Kim et al., 2005; Wang et al., 2014; Zhang et al., 2017; Zhao et al.,

2015).

Calcium, Akt and ERK signaling have also all been previously implicated in hair

cell damage from either aminoglycoside antibiotic toxicity or noise. Inhibiting calcium

movement from the ER or mitochondria has been shown to protect against hair cell

damage (Esterberg et al., 2014; Wang et al., 2019). There is evidence that a transfer of

calcium from the ER to the mitochondria in response to aminoglycoside treatment is

triggering ROS generation leading to hair cell death (Esterberg et al., 2016). The Akt

and ERK signaling pathways on the other hand are believed to play a protective role in

hair cell toxicity as inhibition of these pathways is most commonly shown to increase

hair cell damage (Battaglia et al., 2003; Brand et al., 2015; Chung et al., 2006; Haake et

al., 2009; Hayashi et al., 2013; Kurioka et al., 2015). Therefore, we felt these three

signaling molecules were strong candidates to investigate as potential signaling

pathways involved in cadmium-induced hair cell death. Activation of ERK has also

previously been seen in the cochlea in response to cadmium and inhibition of ERK can

protect HE-OC1 cells, a hair cell like cell line, from cadmium-induced cell death (Kim et

al., 2008).

We investigated the role of these signaling molecules in hair cell death using the

zebrafish lateral line. The lateral line is a sensory system in fish used to detect water

motion that consists of clusters of hair cells and support cells, collectively known as

neuromasts, on the surface of the animal. Support cells are glia-like cells surrounding

hair cells that have been shown to play key roles in the development, maintenance and

regeneration of hair cells (reviewed in Monzack and Cunningham, 2013). Hair cells of

the zebrafish lateral line have been shown to be functionally, molecularly and

morphologically similar to mammalian hair cells (reviewed in Nicolson, 2005; Whitfield,

2002).

We have previously shown that cadmium causes reliable hair cell death in the

zebrafish lateral line within three hours in a dose-dependent manner, and therefore

used that treatment paradigm in this study (Schmid et al., 2020). We first found that

removal of extracellular calcium following cadmium treatment caused an increase in hair

cell death both in control conditions and following cadmium treatment. We also found

that co-treatment of a calmodulin inhibitor with cadmium could significantly protect

against cadmium-induced hair cell death. However, when the calmodulin inhibitor was

applied after fish were washed out of cadmium, a condition we refer to as

posttreatment, protection was no longer seen. Additionally, other calcium signaling

pathway inhibitors including a CaMKII inhibitor, inositol IP3 receptor inhibitor, and

mitochondria calcium uniporter inhibitor failed to protect against cadmium-induced hair

cell death. We also saw an increase in levels of pAkt in hair cells and pERK in

supporting cells in response to cadmium, however, inhibition of these pathways

enhanced hair cell death rather than protecting against it. Therefore, we conclude that

the signaling pathways responsible for cadmium-induced hair cell death appear distinct

from those commonly seen in cadmium-induced toxicity in other cell types and partially

distinct from those seen in other forms of hair cell toxicity.

MATERIALS AND METHODS

Animals

All experiments were carried out in 5 days post-fertilization (dpf) *AB strain wild-type zebrafish larvae. Fish larvae were housed in an incubator maintained at 28.5°C with a 14/10 hour light/dark cycle and raised in embryo media (EM) consisting of 1 mM $MgSO_4$, 150 µM KH_2PO_4, 42 µM Na_2HPO_4, 1 mM $CaCl_2$, 500 µM KCl, 15 mM NaCl, and 714 µM $NaHCO_3$. The Lafayette College Institution Animal Care and Use Committee approved all experiments.

Drug Treatments

The following drugs were used: cadmium chloride hemipentahydrate (Fisher Scientific), water-soluble KN-93 (Abcam), MK-2206 (Millipore Sigma), PD184352 (Millipore Sigma), RU360 (Millipore Sigma), W-13 (Enzo) and Xestospongin (Abcam). Stock solutions for cadmium chloride, KN-93, and Ru360 were made in water. Stock solutions of all other drugs were made in dimethyl sulfoxide (DMSO, Millipore Sigma). All drugs were then diluted to their treatment concentrations in EM. For drugs diluted in DMSO the control groups contained DMSO at the same volume as the drug. For the water diluted drugs EM was used for the control group.

For the drug treatments, fish were initially put into netwell inserts (Corning) in 6 well plates containing EM. The netwell inserts with the larvae were then moved to new plates containing the solutions the fish were to be treated in. At the end of treatment, fish were washed three times in EM, again by moving the netwell inserts to new plates, between treatments or before fixation. For experiments where fish were moved to EM

with reduced calcium levels after cadmium treatment, they were washed three times in

zero calcium EM instead of standard EM before being moved to the reduced or zero

calcium EM and then returned to standard EM immediately prior to fixation.

Immunostaining

Fish were anesthetized with MS-222 and then fixed for 2 hours at room

temperature in 4% paraformaldehyde. Antibody labeling was carried out as previously

described (Stawicki et al., 2014). The following primary antibodies were used: rabbit

anti-parvalbumin (ThermoFisher, PA1-933) diluted at 1:1,000, mouse anti-otoferlin

(Developmental Studies Hybridoma Bank, HCS-1) diluted at 1:100, rabbit anti-phospho-

Akt (Ser473) (Cell Signaling, 9271) diluted at 1:100, and rabbit anti-phospho-p44/42

MAPK (ERK1/2) (Thr202/Tyr204) (Cell Signaling, 4370) diluted at 1:200.

Hair Cell Counts

For hair cell counts fish were either stained with just parvalbumin or costained

with otoferlin and parvalbumin. In the majority of experiments the parvalbumin stain was

used for counts, however, in fish that were costained with otoferlin the otoferlin stain

was used if the parvalbumin stain was too dim to reliably count hair cells. Within a single

experiment the same antibody was always used for all counts. To perform hair cell

counts fish were observed on an Accu-Scope EXC-350 microscope using the 40X

objective. The number of hair cells in the following neuromasts OP1, M2, IO4, O2, MI2,

and MI1 (Raible and Kruse, 2000) were counted for each larvae. The total number of

hair cells counted for that fish was then divided by 6 to get an average number of hair

cells per neuromast. To normalize data for the zero calcium and ERK inhibitor

experiment, the average hair cells/neuromast number for each treatment group was

divided by the average hair cells/neuromast number for their respective control groups

with no cadmium and then multiplied by 100 to be converted to a percentage.

Confocal Imaging and Analysis of pAkt and pERK levels

To calculate pAkt and pERK levels fish stained with either the pAkt or pERK

antibodies and the otoferlin antibody were imaged on a Zeiss LSM800 confocal

microscope using the Zen Blue software. For pAkt, the approximate center of the

neuromast was selected and 10 planes separated by 1 μm were imaged for each

neuromast. For pERK, there was significant neuronal labeling in the control condition

that we did not want to include in our analysis. Therefore, the top and bottom of the

neuromast were selected and planes separated by 1μm were imaged throughout the

neuromast regardless of number with images ranging from 11-29 planes. Then 10

planes with minimal neuronal labeling were selected for subsequent analysis. One

neuromast of the anterior lateral line was imaged per fish with the exact neuromast to

be imaged selected based on which neuromasts were best positioned to image. Image

analysis was conducted in Fiji. Maximum projection images were made for each 10-

plane stack. The neuromast was outlined using the otoferlin label and the average

fluorescent intensity of the pAkt or pERK signal was measured.

Statistics and Graphical Representation of Data

All experiments were carried out with n=10 animals per group. Statistics were

calculated in GraphPad Prism 6 which was also used to generate graphs. For

experiments where two or more different groups were tested against a range of calcium

or cadmium doses, a two-way ANOVA was used with the either a Tukey's or Šídák's

multiple comparisons test depending on which was recommended by Prism. The

multiple comparisons test used for each experiment is mentioned in the figure legend.

Graphs of this data were represented by the average of each data set with error bars

showing the standard deviation. For all other experiments, a one-way ANOVA was used

again with a Tukey's multiple comparison test as all groups were compared. Graphs of

this data show the individual data points as well as the average and standard

deviations. For experiments where the ANOVA showed a significant difference between

relevant groups the significance found by the multiple comparisons tests are shown on

the graphs as asterisks.

233　　　　　　　　　　　　　　　　**RESULTS**

Alterations in extracellular calcium levels following cadmium treatment do not

cause dramatic changes in cadmium-induced hair cell death

Previous research has shown that calcium plays an important role in cadmium-

induced cell death in many cell types. In some cell types, it appears that extracellular

calcium is important as chelating it or treating cells with calcium channel blockers can

protect against cadmium-induced toxicity (Liu et al., 2018; Yang et al., 2008; Yeh et al.,

2009). Unfortunately, determining the role of extracellular calcium in cadmium-induced

hair cell toxicity is complicated by the fact that we have previously shown that cadmium-

induced hair cell toxicity is mechanotransduction-activity dependent (Schmid et al.,

2020). Extracellular calcium levels are known to impact hair cell mechanotransduction

activity (reviewed in Fettiplace, 2017; Ó Maoiléidigh and Ricci, 2019) with calcium

chelation completely blocking it (Assad et al., 1991). It has previously been shown that

the entry and toxicity of other mechanotransduction-dependent ototoxins can be altered

by alterations in extracellular calcium levels for this reason (Coffin et al., 2009).

Therefore, to investigate whether calcium signaling was impacting cadmium-induced

hair cell death we first wanted to determine if there was continued hair cell death

following cadmium removal. If there was, we could potentially separate cadmium entry,

which is the phase of cadmium-induced hair cell death most likely to be

mechanotransduction dependent, from the subsequent cell death process. To do this

we treated fish with 30 μM of cadmium chloride for either 1 or 2 hours, fish were then

washed out of cadmium, and either fixed immediately or fixed three hours after the start

of the experiment. We found that following both the 1 and 2 hour cadmium chloride

treatment continued hair cell death was seen after cadmium removal (Figure 1).

Figure 1: Cadmium causes continued cell death after removal. Hair cells remaining after various

treatment paradigms with 30μM of cadmium. Fish were either treated with cadmium chloride for 1 hour

and immediately fixed (squares), treated with cadmium for 1 hour then left in EM for two hours before

fixation (triangles), treated with cadmium for 2 hours and then immediately fixed (upside down triangles),

or treated with cadmium for 2 hours and then left in EM for 1 hour before fixation (diamonds). Control fish

were kept in EM for three hours (circles). There was a significant amount of cell death after cadmium

removal for both the 1 hour and 2 hours cadmium treatment paradigms (*** = p<0.001 by ANOVA and

Tukey post-hoc test).

As we saw significant continued cadmium-induced hair cell death after cadmium

removal, we next wanted to test if this continued cell death could be altered by changing

extracellular calcium levels. To do this we used the treatment paradigm where fish were

treated with cadmium chloride for one hour, washed out of cadmium, and then fixed two

hours later. We first tested a range of calcium concentrations by treating fish with either

EM or 30 μM cadmium chloride for 1 hour and then washing them out of that solution

and moving them into EM with calcium concentrations ranging from 0-2 mM for 2 hours

prior to fixation. We found the only calcium concentration where a significant difference

was seen in the amount of cadmium-induced hair cell death as compared to the

standard EM calcium concentration of 1mM was 0mM (Figure 2A). However, rather than

protecting hair cells the removal of extracellular calcium appeared to increase cadmium-

induced hair cell death. We then treated fish with a range of cadmium doses from 0-120

μM for 1 hour before washing them out of cadmium and moving them to either standard

EM with 1 mM calcium or EM with 0 mM calcium for 2 hours prior to fixation. Again, a

significant increase in the amount of cadmium-induced hair cell death was seen in the

fish that were moved to 0 mM calcium following cadmium treatment (Figure 2B).

However, there was also a significant reduction in the number of control hair cells in this

condition similar to what has previously been seen when zebrafish are put in calcium-

free EM (Coffin et al., 2009). Therefore, to determine whether the incubation in 0 mM

calcium EM was sensitizing fish to cadmium-induced hair cell death or simply killing hair

cells on its own we normalized these data for both the 1 mM and 0 mM calcium EM

groups to the 0 cadmium control for each group. We found in this case while there is still

a trend toward a decrease in hair cell number at all cadmium concentrations tested, the

only concentration that showed a significant decrease was 120 μM cadmium (Figure

2C). Taken together this data suggests alterations in calcium concentration following

cadmium removal do not have dramatic impacts on cadmium-induced hair cell death.

However, due to the complications with calcium's effect on mechanotransduction

activity, we cannot rule out an impact from extracellular calcium at earlier time points in

the cell death process.

300
301
Figure 2: Removal of extracellular calcium following cadmium treatment kills hair cells on its own

but does not dramatically impact cadmium-induced hair cell death. (A) Hair cells remaining after a 1

hour treatment with either EM, control (solid line), or 30 μM cadmium chloride (dotted line) followed by a 2

hour incubation in EM containing varying concentrations of calcium. The only calcium concentration

where there was a significant difference in the number of hair cells following cadmium treatment

compared to the standard EM 1 mM calcium group was the 0 mM calcium group (** = $p<0.01$ by Two-

Way ANOVA and Tukey's post-hoc test). (B) Hair cells remaining after a 1 hour treatment with varying

doses of cadmium chloride followed by a 2 hour incubation in either EM containing 1mM calcium (solid

line) or EM containing 0 mM calcium (dotted line). A significant decrease in hair cells in the 0 mM calcium

group was seen for all concentrations of cadmium tested except 60 μM (* = $p<0.05$, ** = $p<0.01$, *** =

$p<0.001$, by Two-Way ANOVA and Šídák's post-hoc test). (C) The same data as in B, but normalized to

the zero cadmium control for each group. Following normalization, a significant decrease in hair cells in

the 0 mM calcium group was only seen at 120 μM cadmium. (** = p<0.01, by Two-Way ANOVA and

Šídák's post-hoc test). N = 10 fish for each group. Error bars = Standard deviation.

Calmodulin and CaMKII inhibitors do not consistently protect against cadmium-

induced hair cell death

Calmodulin is a major intracellular receptor for calcium and calmodulin inhibition

has been shown to protect against cadmium-induced cell death (Niewenhuis and

Prozialeck, 1987; Xu et al., 2011; Yang and Yang, 1997). Calmodulin inhibition has also

previously been shown to protect against aminoglycoside-induced hair cell death

(Esterberg et al., 2013). To see if calmodulin inhibition could protect against cadmium-

induced hair cell death we first cotreated fish with 30 μM of the calmodulin inhibitor W-

13 hydrochloride and varying doses of calmodulin chloride ranging from 0-120 μM for

three hours. We found that calmodulin inhibition significantly reduced cadmium-induced

hair cell death at lower doses of cadmium chloride, 15 and 30 μM, but not at higher

doses, 60 and 120 μM (Figure 3A).

However, calmodulin inhibition has also previously been shown to inhibit rapid

apical endocytosis in hair cells as measured by rapid FM1-43 uptake, a process

dependent on mechanotransduction activity (Seiler and Nicolson, 1999). As mentioned

above inhibiting mechanotransduction can protect against cadmium-induced hair cell

death (Schmid et al., 2020), therefore, we tested whether inhibiting calmodulin only after

cadmium removal would still significantly protect against cadmium-induced hair cell

death. To do this fish were treated with varying doses of cadmium chloride from 0-30

μM for one hour, washed out of cadmium, and then immediately fixed or placed into

DMSO or 30 μM W-13 for two hours before fixation. Using this treatment paradigm we

no longer saw protection from calmodulin inhibition (Figure 3B). This suggests the

protection seen with cotreatment with the calmodulin inhibitor may be due to impacts on

mechanotransduction-dependent processes, though we can also not rule out a distinct

early role for calmodulin in the cell death process.

Calmodulin is known to activate CaMKII and inhibiting CaMKII has also been

shown to protect against cadmium-induced cell death in multiple cell types (Chen et al.,

2011; Liu et al., 2018; Liu and Templeton, 2007). Therefore, as our calmodulin results

did now show a clear protective impact we, subsequently tested if CaMKII inhibition

would impact cadmium-induced hair cell death. To do this we cotreated fish with 10 μM

of the CaMKII inhibitor KN-93, a dose previously shown to be effective in zebrafish

(Rothschild et al., 2013), and varying doses of cadmium chloride for three hours. We

found CaMKII inhibition did not significantly impact cadmium-induced hair cell death

(Figure 3C). This further supports the idea that calmodulin and CaMKII are not playing a

role in cadmium-induced hair cell death outside of calmodulin's impact on

mechanotransduction activity.

Figure 3: Calmodulin inhibitor cotreatment protects against cadmium-induced hair cell death whereas calmodulin inhibitor posttreatment and CaMKII inhibitor cotreatment fail to protect. (A) Hair cells remaining after a 3 hour cotreatment of either DMSO, a vehicle control, (solid line), or 30 μM of the calmodulin inhibitor W-13 (dotted line) and varying doses of cadmium chloride. W-13 cotreatment caused a significant reduction in hair cell death seen at lower cadmium doses (** = $p<0.01$, **** = $p<0.0001$ by Two-Way ANOVA and Šídák's post-hoc test). (B) Hair cells remaining after a 1 hour treatment with varying doses of cadmium chloride and either immediate fixation (dashed line), or a two

hour post-treatment with DMSO (solid line) or W-13 (dotted line). No significant protection was seen

comparing the DMSO and W-13 groups by Two-Way ANOVA and Tukey's post-hoc test. (C) Hair cells

remaining after a 3 hour cotreatment of either plain embryo media or 10 μM of the CaMKII inhibitor KN-93

and varying doses of cadmium chloride. No significant difference was seen between the two groups by

Two-Way ANOVA. n = 10 fish for each group. Error bars = Standard deviation.

Inhibition of the IP3 Receptor and Mitochondrial Uniporter fail to protect against

cadmium-induced hair cell death

We next wanted to test whether calcium flow from the ER or mitochondria would

impact cadmium-induced hair cell death as blocking the IP3 receptor and the

mitochondria calcium uniporter has been shown to protect other cell types against

cadmium-induced toxicity (Lee et al., 2005; Liu et al., 2016, 2018; Shih et al., 2005; S.

H. Wang et al., 2009; Wang et al., 2008, 2022). These inhibitors have also been shown

to protect hair cells from other forms of toxicity. (Esterberg et al., 2014; Wang et al.,

2019). To test this in hair cells we cotreated fish for three hours with either 1 μM of the

IP3 receptor inhibitor Xestospongin C or 500 nM of the mitochondrial uniporter inhibitor

RU360, doses that have previously been shown to protect against aminoglycoside-

induced hair cell death in the zebrafish lateral line (Esterberg et al., 2014), and varying

doses of cadmium chloride for three hours. In both cases, the inhibitor failed to

significantly protect against cadmium-induced hair cell death with RU360 causing a

slight enhancement of cadmium-induced hair cell death at the lowest cadmium dose

(Figure 4). Thus, our results indicate calcium flow from the ER and mitochondria is not

playing a significant role in cadmium-induced hair cell death.

Figure 4: Inhibiting ER and mitochondria calcium transfer does not protect against cadmium-induced hair cell death. Hair cells remaining after a three hour cotreatment of either EM or DMSO (solid lines), or (A) 1 μM of the IP3 receptor inhibitor Xestospongin C, or (B) 500 nM of the mitochondrial uniporter inhibitor RU360 (dotted lines) and varying doses of cadmium chloride. Neither of the inhibitors significantly protected against cadmium-induced hair cell death with RU360 showing a significant increase in cadmium-induced hair cell death at the lowest dose (** = $p < 0.01$ = by Two-Way ANOVA and Šídák's post-hoc test). N = 10 fish/group. Error bars = standard deviation.

Cadmium increases Akt and ERK signaling in neuromasts

In many cell types cadmium increases the level of phosphorylated Akt and ERK (Chen et al., 2008, 2011; Fujiki et al., 2013; Iryo et al., 2000; J. Kim et al., 2005; Kim et al., 2008; Liu and Templeton, 2007; Misra et al., 2003; Xiao et al., 2009; Xu et al., 2011). Activation of these pathways has also been seen in hair cells or supporting cells in response to other forms of hair cell damage (Chung et al., 2006; Herranen et al., 2018; Ingersoll et al., 2020; Lahne and Gale, 2008; Maeda et al., 2013). Based on this evidence we tested whether the Akt or ERK pathways would be activated in either hair cells or supporting cells of the zebrafish lateral line in response to cadmium. To do this we treated fish with 30 μM of cadmium chloride for one hour and then fixed and stained

with an antibody for either pAkt or pERK. We found that treatment with cadmium

significantly increased the levels of pAkt seen in hair cells (Figure 5A & B). This

increase was lost when fish were cotreated with cadmium and 5 μM of the Akt inhibitor

MK-2206. We also saw a significant increase in pERK levels following cadmium

treatment that was lost when fish were cotreated with cadmium and 3 μM of the MEK

inhibitor PD184352 which suppresses ERK phosphorylation (Figure 5C & D). However,

unlike pAkt labeling which appeared localized to hair cells, pERK labeling appears

localized to supporting cells (Figure 5B & D). This is in agreement with past research in

mammals showing increased pERK in support cells but not hair cells in response to

other forms of hair cell damage (Herranen et al., 2018; Ingersoll et al., 2020; Lahne and

Gale, 2008; Maeda et al., 2013).

Figure 5: Cadmium treatment increases pAkt levels in hair cells and pERK levels in supporting cells. (A) Quantification of the fluorescence intensity of pAkt in fish treated for 1 hour with either DMSO alone, 30μM cadmium chloride (Cd) + DMSO, or 30μM Cd and 5 μM of MK-2206. (**** = p<0.0001 by ANOVA and Tukey post-hoc test). (B) Images of a neuromast from the DMSO alone group (top), the Cd + DMSO group (middle), and Cd + MK-2206 group (bottom). Hair cells are labeled with the otoferlin (HCS-1) antibody in green (left) and pAkt in magenta (center). (C) Quantification of the fluorescence intensity of pERK in fish treated for 1 hour with either DMSO alone, 30μM Cd + DMSO, or 30μM Cd and 3 μM of PD184352. (*** = p<0.001 by ANOVA and Tukey post-hoc test). (D) Images of a neuromast from the DMSO alone group (top) and the Cd + DMSO group (middle), and Cd + MK-2206 group (bottom). Hair cells are labeled with the HCS-1 antibody in green (left) and pERK in magenta (center). Scale bar = 10 μm.

Inhibiting Akt and ERK signaling pathways increases cadmium-induced hair cell death

430
431

Having seen an increase in pAkt and pERK levels following cadmium treatment,
we wanted to determine whether the Akt or ERK signaling pathways impacted
cadmium-induced hair cell death. Akt and/or ERK inhibitors have been shown to protect
against cadmium-induced toxicity in multiple cell types (Chen et al., 2008; Iryo et al.,
2000; Jiang et al., 2015; J. Kim et al., 2005; Kim et al., 2008; Wang et al., 2014; Zhang
et al., 2017), suggesting these pathways normally function to promote cadmium-induced
cell death. In hair cells blocking Akt or ERK signaling commonly increases hair cell
death, suggesting these pathways may be activated to protect hair cells (Battaglia et al.,
2003; Brand et al., 2015; Chung et al., 2006; Haake et al., 2009; Hayashi et al., 2013;
Kurioka et al., 2015).

To determine if the Akt or ERK pathways were impacting cadmium-induced hair
cell death, we cotreated fish with either 5 μM of the Akt inhibitor MK-2206, or 1.5 μM of
the MEK inhibitor PD184352, which inhibits ERK activation, and varying doses of
cadmium chloride for three hours. A smaller dose of PD184352 was used in this
experiment than the pERK levels experiment due to increased hair cell toxicity from
PD184352 alone with the longer treatment time. We found both drugs caused a
decrease in hair cell number following cadmium treatment particularly, at the 30 and 60
μM cadmium chloride doses (Figure 6A & B). However, as noted above PD184352
caused a loss of hair cells independent of cadmium treatment, a similar loss of hair cells
from simple ERK inhibition has also previously been seen in cochlear explants
(Battaglia et al., 2003). Therefore, to determine if PD184352 was enhancing cadmium-

induced hair cell death or simply killing hair cells on its own, we normalized the data for

both the DMSO and PD184352 groups to the zero cadmium control (Figure 6C). In

doing so we still saw a significant increase in cadmium-induced hair cell death at the 30

and 60 μM cadmium chloride doses. Thus, we conclude that unlike in other cell types

Akt and ERK signaling do not appear to be facilitating cadmium-induced hair cell death

and may be playing a protective role similar to what is seen in other forms of hair cell

toxicity.

Figure 6: Inhibiting Akt and ERK signaling increases cadmium-induced cell death. (A) Hair cells

remaining after a 3 hour cotreatment of either DMSO (solid line) or 5 μM of the Akt inhibitor MK-2206

(dotted line) and varying doses of cadmium chloride. MK-2206 cotreatment caused a significant increase

in cadmium-induced hair cell death. (B) Hair cells remaining after a 3 hour cotreatment of either DMSO, a

vehicle control, (solid line), or 1.5 μM of the ERK pathway inhibitor PD184352 (dotted line) and varying

doses of cadmium chloride. PD184352 caused a significant decrease in the number of hair cells seen. (C)

The same data as in B, but normalized to the zero cadmium control for each group. PD184352 caused a

significant increase in the amount of cadmium-induced hair cell death seen. (* = $p<0.05$, *** = $p<0.001$,

**** = $p<0.0001$, by Two-Way ANOVA and Šídák's post-hoc test). N = 10 fish for each group. Error bars =

Standard deviation.

473   **DISCUSSION**
474

Overall we found that cadmium-induced hair cell death showed both some

distinctions and similarities with cadmium-induced cell death in other cell types and

other forms of hair cell toxicity. One of the distinctions was that we failed to find a

significant role for calcium signaling. This is in contrast to both cadmium-induced cell

death in other cell types and other forms of hair cell death where calcium appears to be

a key mediator in the cell death process (Esterberg et al., 2014, 2013; Liu et al., 2016;

L. Wang et al., 2009; Wang et al., 2008, 2019; Xu et al., 2011; Yang et al., 2008, 2007;

Yeh et al., 2009). We did however, see activation of both the Akt and ERK signaling

pathways. These two pathways are known to be activated both in response to cadmium

in other cell types (Chen et al., 2008, 2011; Fujiki et al., 2013; Iryo et al., 2000; J. Kim et

al., 2005; Kim et al., 2008; Liu and Templeton, 2007; Misra et al., 2003; Xiao et al.,

2009; Xu et al., 2011) and in other forms of hair cell toxicity (Chung et al., 2006;

Herranen et al., 2018; Ingersoll et al., 2020; Lahne and Gale, 2008; Maeda et al., 2013).

Though unlike in other cell types, inhibiting these signaling pathways enhanced

cadmium-induced hair cell death rather than protecting against it (Chen et al., 2008; Iryo

et al., 2000; Jiang et al., 2014, 2015; Y. K. Kim et al., 2005; Wang et al., 2014; Zhang et

al., 2017; Zhao et al., 2015) similar to what has been shown in other forms of hair cell

toxicity (Battaglia et al., 2003; Brand et al., 2015; Chung et al., 2006; Haake et al., 2009;

Hayashi et al., 2013; Kurioka et al., 2015).

While most of our manipulations of calcium signaling failed to protect against

cadmium-induced hair cell death, one manipulation that did cause significant protection

was the cotreatment of cadmium and a calmodulin inhibitor. It has been shown that

cadmium can activate calmodulin in lieu of calcium and it has been hypothesized this

may be one potential mechanism of cadmium-induced toxicity (Chao et al., 1984; Sutoo

et al., 1990). Treatment with inhibitors blocking calmodulin-dependent signaling

pathways has been shown to protect against cadmium in multiple cell types and tissues

including the kidneys, osteoblasts, and testes (Ha et al., 2016; Niewenhuis and

Prozialeck, 1987; Yang and Yang, 1997). However, these experiments are complicated

in hair cells by the fact that calmodulin inhibition also inhibits rapid apical endocytosis in

hair cells, a process dependent on mechanotransduction activity (Seiler and Nicolson,

1999). As inhibiting mechanotransduction activity is known to inhibit cadmium-induced

hair cell death (Schmid et al., 2020) we cannot definitively say that the protection we

see from calmodulin inhibition is due to a role for calmodulin in the cell death process.

The fact that protection is no longer seen when the calmodulin inhibitor is added after

cadmium treatment, rather than as a cotreatment, and the fact that CaMKII inhibition

was unable to protect hair cells from cadmium-induced hair cell death further supports

the idea that calmodulin inhibition is mainly impacting hair cell mechanotransduction

activity and cadmium entry rather than the cell death process.

We also failed to see protection against cadmium-induced hair cell death using

other calcium signaling manipulations including removing extracellular calcium and

inhibiting the IP3 receptor and mitochondrial calcium uniporter. These manipulations

have been shown to protect against cadmium-induced cell death in other cell types (Lee

et al., 2005; Liu et al., 2016, 2018; Shih et al., 2005; Wang et al., 2008; Xu et al., 2011).

However, these previous studies were carried out in mitotic cultured cells, including

mesangial cells, PC-12 cells, proximal tubule cells, osteoblasts, and osteosarcoma

cells, whereas our study was carried out *in vivo* in postmitotic cells, so this may be the

reason for the discrepancy. Indeed another study looking at neurons, another

postmitotic cell, failed to see protection from cadmium-induced cell death when

inhibiting the mitochondrial calcium uniporter (Tang et al., 2021). Calcium signaling in

general and specifically calmodulin, calcium release via the IP3 receptor, and activation

of the mitochondrial uniporter have all been shown to be important for the progression

of mitosis (Lagos-Cabré et al., 2020; Nugues et al., 2022; Rasmussen and Means,

1989; Takuwa et al., 1995; Zhao et al., 2019). Therefore, mitotic cells may be more

sensitive to cadmium-induced alterations in these calcium signaling pathways. Also,

even in mitotic cells, there have been discrepancies over the role of calcium signaling in

cadmium-induced cell death. For example, when looking at CaMKII inhibition a study in

an osteoblast cell line showed no effect (Ha et al., 2016), and a study in mesangial cells

showed an increase in cell death (Xiao et al., 2009).

The fact that blocking the IP3 receptor and mitochondrial calcium uniporter failed

to protect against cadmium-induced hair cell death also contrasted with what has

previously been shown in other forms of hair cell death (Esterberg et al., 2014; Wang et

al., 2019). However, while there are clearly some overlapping mechanisms between

different forms of hair cell death, drug screens frequently find drugs that can protect

against only a subset of ototoxic insults (Coffin et al., 2013; Owens et al., 2009; Vlasits

et al., 2012) showing there are distinct mechanisms by which ototoxic compounds

induce cell death in hair cells.

While we did not find a significant impact of calcium signaling in cadmium-

induced hair cell death we did see an increase in Akt and ERK signaling similar to what

has been seen in other cell types. This was somewhat unexpected as in other cell types

this increase is calcium signaling dependent (Chen et al., 2011; Fujiki et al., 2013; Jiang

et al., 2015; Liu et al., 2018; Xiao et al., 2009; Xu et al., 2011). However, it has

previously been shown that ERK signaling can be activated in supporting cells even

following calcium chelation (Lahne and Gale, 2008). One potential alternative

mechanism for the activation of these pathways includes Ras and Raf signaling which

has previously been implicated in activating ERK signaling following hair cell damage

(Battaglia et al., 2003; Lahne and Gale, 2008). Another alternative activation route is

growth factor signaling. Increases in growth factor receptors have been seen after hair

cell damage (Lee and Cotanche, 1996; Zine and de Ribaupierre, 1999), and inhibiting

the IGF1R receptor has been shown to also increase susceptibility to aminoglycoside-

induced hair cell damage (Alassaf and Halloran, 2021), suggesting a role for growth

factor signaling in protecting hair cells from damage.

While the activation of ERK and Akt signaling was similar to what was seen in

other forms of cadmium-induced cell death, the fact that inhibiting these pathways

increased rather than protected against cell death was unexpected. This, however, is

not unprecedented. While many studies have found blocking Akt or ERK activation

protects cells against cadmium-induced cell death (Chen et al., 2008; Iryo et al., 2000;

Jiang et al., 2015; J. Kim et al., 2005; Zhang et al., 2017) others have found that it has

no effect (Låg et al., 2005; Liu and Templeton, 2007; Rigon et al., 2008; Wang et al.,

2008), and still others that death is increased similar to what we observed (Chao and

Yang, 2001; Chuang et al., 2000; Xiao et al., 2009). Some of this disparity may come

from differences in treatment paradigms. For example, in astrocytes ERK inhibition

protects against chronic but not acute cadmium treatment (Jiang et al., 2015). The type

of cell death also appears important as in macrophages ERK inhibition can protect

against necrosis but not apoptosis induced by cadmium (J. Kim et al., 2005).

Unlike what we found in calcium signaling, the role of the Akt and ERK signaling

pathways in cadmium-induced hair cell death appears similar to what has previously

been shown for other forms of hair cell damage. These pathways are activated in

response to both noise and other ototoxic drugs (Chung et al., 2006; Herranen et al.,

2018; Ingersoll et al., 2020; Lahne and Gale, 2008; Maeda et al., 2013), and inhibiting

them has been shown to increase hair cell death (Battaglia et al., 2003; Brand et al.,

2015; Chung et al., 2006; Haake et al., 2009; Hayashi et al., 2013; Kurioka et al., 2015).

Particularly striking is the fact that ERK activation is seen not in the hair cells

themselves but rather in the support cells. This has also been seen in the mammalian

inner ear in response to aminoglycosides, cisplatin, and noise (Herranen et al., 2018;

Ingersoll et al., 2020; Lahne and Gale, 2008; Maeda et al., 2013).

Cadmium is known to damage support cells (Ding et al., 2020), and thus this

ERK activation could be coming from a direct effect of cadmium on the support cells.

Alternatively, the damaged hair cells could be sending a signal to the support cells to

induce ERK activation. It has been shown that the damage-induced release of ATP and

gap junction signaling are important for support cell ERK activation following mechanical

injury of hair cells (Lahne and Gale, 2008). Once ERK is activated in support cells it

may mediate its protective effect by modulating support cell released HSP70. ERK

activation is important for HSP70 expression and folding (Lim et al., 2019; Xu et al.,

2018; Zhai et al., 2019). Support cells are known to release HSP70 in exosomes

following heat shock, with that HSP70 subsequently protecting hair cells from

aminoglycoside-induced hair cell death (Breglio et al., 2020; May et al., 2013). The

mechanism by which Akt activation is working in hair cell death is likely more

straightforward as it is activated in the hair cells themselves. The activation of Akt

signaling is well known to promote cell survival and inhibit cell death and multiple

apoptotic regulators are Akt substrates (reviewed in Downward, 2004; Franke et al.,

2003), providing a potential mechanism by which Akt activation could protect hair cells

from cadmium-induced cell death.

The highly conserved activation and potential protective nature of Akt and ERK

pathways suggest enhancing their activation could be a potential target for protecting

hair cells from damage, however, as with other forms of cadmium-induced cell death,

there are other examples where the activation of ERK appears to promote hair cell

death (Lahne and Gale, 2008; So et al., 2007). The reasons behind these dual roles of

ERK would need to be further investigated before ERK activation could be used as a

potential therapeutic target.

ACKNOWLEDGEMENTS
606

607   This work was supported by a Lafayette College Academic Research Committee

608 faculty research grant to TMS and the Lafayette College Neuroscience program. We

609 thank Amy Badillo for assistance with zebrafish care.

610
611 **AUTHOR CONTRIBUTIONS**
612
613 **Jennifer Galdieri:** investigation, writing – original draft. **Chloe Adams:** investigation,

614 writing – review & editing. **María Padilla:** investigation, writing – review & editing.

615 **Tamara Stawicki:** conceptualization, methodology, formal analysis, investigation,

616 writing – original draft and revision, and review and editing.

618 **COMPETING INTERESTS**
619
620 The authors have no competing interests to declare

621 **REFERENCES**
622

Alassaf, M., Halloran, M.C., 2021. Pregnancy-associated plasma protein-aa regulates

endoplasmic reticulum–mitochondria associations. Elife 10, 259687.

https://doi.org/10.7554/eLife.59687

Assad, J.A., Shepherd, G.M.G., Corey, D.P., 1991. Tip-link integrity and mechanical

transduction in vertebrate hair cells. Neuron 7, 985–994.

https://doi.org/10.1016/0896-6273(91)90343-X

Battaglia, A., Pak, K., Brors, D., Bodmer, D., Frangos, J.A., Ryan, A.F., 2003.

Involvement of ras activation in toxic hair cell damage of the mammalian cochlea.

Neuroscience 122, 1025–1035. https://doi.org/10.1016/j.neuroscience.2003.08.041

Benters, J., Flögel, U., Schäfer, T., Leibfritz, D., Hechtenberg, S., Beyersmann, D.,

1997. Study of the interactions of cadmium and zinc ions with cellular calcium

homoeostasis using 19F-NMR spectroscopy. Biochem J 322 (Pt 3, 793–799.

https://doi.org/10.1042/BJ3220793

Bhardwaj, J.K., Panchal, H., Saraf, P., 2021. Cadmium as a testicular toxicant: A

Review. J Appl Toxicol 41, 105–117. https://doi.org/10.1002/JAT.4055

Brand, Y., Levano, S., Radojevic, V., Naldi, A.M., Setz, C., Ryan, A.F., Pak, K.,

Hemmings, B.A., Bodmer, D., 2015. All Akt isoforms (Akt1, Akt2, Akt3) are involved

in normal hearing, but only Akt2 and Akt3 are involved in auditory hair cell survival

in the mammalian inner ear. PLoS One 10, e0121599.

https://doi.org/10.1371/journal.pone.0121599

Breglio, A.M., May, L.A., Barzik, M., Welsh, N.C., Francis, S.P., Costain, T.Q., Wang, L.,

Anderson, D.E., Petralia, R.S., Wang, Y.X., Friedman, T.B., Wood, M.J.A.,

Cunningham, L.L., 2020. Exosomes mediate sensory hair cell protection in the

inner ear. J Clin Invest 130, 2657–2672. https://doi.org/10.1172/JCI128867

Chao, J.I., Yang, J., 2001. Opposite roles of ERK and p38 mitogen-activated protein

kinases in cadmium-induced genotoxicity and mitotic arrest. Chem Res Toxicol 14,

1193–1202. https://doi.org/10.1021/TX010041O

Chao, S., Suzuki, Y., Zysk, J., Cheung, W., 1984. Activation of calmodulin by various

metal cations as a function of ionic radius. Mol Pharmacol 26, 75–82.

Chen, L., Liu, L., Luo, Y., Huang, S., 2008. MAPK and mTOR pathways are involved in

cadmium-induced neuronal apoptosis. J Neurochem 105, 251–261.

https://doi.org/10.1111/J.1471-4159.2007.05133.X

Chen, S., Ren, Q., Zhang, J., Ye, Y., Zhang, Z., Xu, Y., Guo, M., Ji, H., Xu, C., Gu, C.,

Gao, W., Huang, S., Chen, L., 2014. N-acetyl-L-cysteine protects against cadmium-

induced neuronal apoptosis by inhibiting ROS-dependent activation of Akt/mTOR

pathway in mouse brain. Neuropathol Appl Neurobiol 40, 759–777.

https://doi.org/10.1111/nan.12103

Chen, S., Xu, Y., Xu, B., Guo, M., Zhang, Z., Liu, L., Ma, H., Chen, Z., Luo, Y., Huang,

S., Chen, L., 2011. CaMKII is involved in cadmium activation of MAPK and mTOR

pathways leading to neuronal cell death. J Neurochem 119, 1108–1118.

https://doi.org/10.1111/j.1471-4159.2011.07493.x

Choi, Y.-H., Hu, H., Mukherjee, B., Miller, J., Park, S.K., 2012. Environmental cadmium

and lead exposures and hearing loss in U.S. adults: the National Health and

Nutrition Examination Survey, 1999 to 2004. Environ Health Perspect 120, 1544–

50. https://doi.org/10.1289/ehp.1104863

Choi, Y.-H., Kim, K., 2014. Noise-Induced Hearing Loss in Korean Workers: Co-

Exposure to Organic Solvents and Heavy Metals in Nationwide Industries. PLoS

One 9, e97538. https://doi.org/10.1371/journal.pone.0097538

Choi, Y.-H., Park, S.K., 2017. Environmental Exposures to Lead, Mercury, and

Cadmium and Hearing Loss in Adults and Adolescents: KNHANES 2010-2012.

Environ Health Perspect 125, 067003. https://doi.org/10.1289/EHP565

Chuang, S.M., Wang, I.C., Yang, J.L., 2000. Roles of JNK, p38 and ERK mitogen-

activated protein kinases in the growth inhibition and apoptosis induced by

cadmium. Carcinogenesis 21, 1423–1432. https://doi.org/10.1093/carcin/21.5.423

Chung, W.H., Pak, K., Lin, B., Webster, N., Ryan, A.F., 2006. A PI3K pathway mediates

hair cell survival and opposes gentamicin toxicity in neonatal rat organ of corti.

JARO - Journal of the Association for Research in Otolaryngology 7, 373–382.

https://doi.org/10.1007/s10162-006-0050-y

Coffin, A.B., Reinhart, K.E., Owens, K.N., Raible, D.W., Rubel, E.W., 2009. Extracellular

divalent cations modulate aminoglycoside-induced hair cell death in the zebrafish

lateral line. Hear Res 253, 42–51. https://doi.org/10.1016/J.HEARES.2009.03.004

Coffin, A.B., Williamson, K.L., Mamiya, A., Raible, D.W., Rubel, E.W., 2013. Profiling

drug-induced cell death pathways in the zebrafish lateral line. Apoptosis 18, 393–

408. https://doi.org/10.1007/s10495-013-0816-8

Dalton, D.S., Schubert, C.R., Pinto, A., Fischer, M.E., Huang, G.-H., Klein, B.E.K., Klein,

R., Pankow, J.S., Paulsen, A.J., Tsai, M.Y., Tweed, T.S., Cruickshanks, K.J., 2020.

Cadmium, obesity, and education, and the 10-year incidence of hearing

impairment: The beaver dam offspring study. Laryngoscope 130, 1396–1401.

https://doi.org/10.1002/LARY.28244

Ding, D., Zhang, J., Jiang, H., Xuan, W., Qi, W., Salvi, R., 2020. Some Ototoxic Drugs

Destroy Cochlear Support Cells Before Damaging Sensory Hair Cells. Neurotoxicity

Research 2020 37:3 37, 743–752. https://doi.org/10.1007/S12640-020-00170-8

Downward, J., 2004. PI 3-kinase, Akt and cell survival. Semin Cell Dev Biol 15, 177–

182. https://doi.org/10.1016/J.SEMCDB.2004.01.002

Esterberg, R., Hailey, D.W., Coffin, A.B., Raible, D.W., Rubel, E.W., 2013. Disruption of

intracellular calcium regulation is integral to aminoglycoside-induced hair cell death.

Journal of Neuroscience 33, 7513–7525.

https://doi.org/10.1523/JNEUROSCI.4559-12.2013

Esterberg, R., Hailey, D.W., Rubel, E.W., Raible, D.W., 2014. ER-mitochondrial calcium

flow underlies vulnerability of mechanosensory hair cells to damage. The Journal of

Neuroscience 34, 9703–19. https://doi.org/10.1523/JNEUROSCI.0281-14.2014

Esterberg, R., Linbo, T., Pickett, S.B., Wu, P., Ou, H.C., Rubel, E.W., Raible, D.W.,

2016. Mitochondrial calcium uptake underlies ROS generation during

aminoglycoside-induced hair cell death. Journal of Clinical Investigation 126, 3556–

3566. https://doi.org/10.1172/JCI84939

Faroon, O., Ashizawa, A., Wright, S., Tucker, P., Jenkins, K., Ingerman, L., Rudisill, C.,

2012. Toxicological Profile for Cadmium, Toxicological Profile for Cadmium.

Agency for Toxic Substances and Disease Registry (US), Atlanta (GA).

Faucher, K., Fichet, D., Miramand, P., Lagardère, J.P., 2006. Impact of acute cadmium

exposure on the trunk lateral line neuromasts and consequences on the "C-start"

response behaviour of the sea bass (Dicentrarchus labrax L.; Teleostei,

Moronidae). Aquatic Toxicology 76, 278–294.

https://doi.org/10.1016/J.AQUATOX.2005.10.004

Ferraro, P.M., Costanzi, S., Naticchia, A., Sturniolo, A., Gambaro, G., 2010. Low level

exposure to cadmium increases the risk of chronic kidney disease: analysis of the

NHANES 1999-2006. BMC Public Health 10, 304. https://doi.org/10.1186/1471-

2458-10-304

Fettiplace, R., 2017. Hair Cell Transduction, Tuning, and Synaptic Transmission in the

Mammalian Cochlea. Compr Physiol 7, 1197–1227.

https://doi.org/10.1002/CPHY.C160049

Franke, T.F., Hornik, C.P., Segev, L., Shostak, G.A., Sugimoto, C., 2003. PI3K/Akt and

apoptosis: size matters. Oncogene 2003 22:56 22, 8983–8998.

https://doi.org/10.1038/sj.onc.1207115

Fujiki, K., Inamura, H., Matsuoka, M., 2013. Phosphorylation of FOXO3a by PI3K/Akt

pathway in HK-2 renal proximal tubular epithelial cells exposed to cadmium. Arch

Toxicol 87, 2119–2127. https://doi.org/10.1007/S00204-013-1077-6/FIGURES/6

Gallagher, C.M., Kovach, J.S., Meliker, J.R., 2008. Urinary Cadmium and Osteoporosis

in U.S. Women ≥50 Years of Age: NHANES 1988–1994 and 1999–2004. Environ

Health Perspect 116, 1338–1343. https://doi.org/10.1289/ehp.11452

García-Esquinas, E., Pollan, M., Tellez-Plaza, M., Francesconi, K.A., Goessler, W.,

Guallar, E., Umans, J.G., Yeh, J., Best, L.G., Navas-Acien, A., 2014. Cadmium

Exposure and Cancer Mortality in a Prospective Cohort: The Strong Heart Study.

Environ Health Perspect 122, 363–370. https://doi.org/10.1289/ehp.1306587

Ha, T., Burwell, S., Goodwin, M., Noeker, J., Heggland, S., 2016. Pleiotropic roles of Ca

+2/calmodulin-dependent pathways in regulating cadmium-induced toxicity in

human osteoblast-like cell lines. Toxicol Lett 260, 18–27.

https://doi.org/10.1016/J.TOXLET.2016.08.020

Haake, S.M., Dinh, C.T., Chen, S., Eshraghi, A.A., van de Water, T.R., 2009.

Dexamethasone protects auditory hair cells against TNFα-initiated apoptosis via

activation of PI3K/Akt and NFκB signaling. Hear Res 255, 22–32.

https://doi.org/10.1016/j.heares.2009.05.003

Hague, F., Matifat, F., Louvet, L., Brûle, G., Collin, T., 2000. The carcinogen Cd2+

activates InsP3-mediated Ca2+ release through a specific metal ions receptor in

Xenopus oocyte. Cell Signal 12, 419–424. https://doi.org/10.1016/S0898-

6568(00)00079-6

Hayashi, Y., Yamamoto, N., Nakagawa, T., Ito, J., 2013. Insulin-like growth factor 1

inhibits hair cell apoptosis and promotes the cell cycle of supporting cells by

activating different downstream cascades after pharmacological hair cell injury in

neonatal mice. Molecular and Cellular Neuroscience 56, 29–38.

https://doi.org/10.1016/J.MCN.2013.03.003

Herranen, A., Ikäheimo, K., Virkkala, J., Pirvola, U., 2018. The Stress Response in the

Non-sensory Cells of the Cochlea Under Pathological Conditions—Possible Role in

Mediating Noise Vulnerability. JARO - Journal of the Association for Research in

Otolaryngology 19, 637–652. https://doi.org/10.1007/s10162-018-00691-2

Huh, D.A., Choi, Y.H., Ji, M.S., Moon, K.W., Yoon, S.J., Sohn, J.R., 2018. Comparison

of Pure-Tone Average Methods for Estimation of Hearing Loss Caused by

Environmental Exposure to Lead and Cadmium: Does the Pure-Tone Average

Method Which Uses Low-Frequency Ranges Underestimate the Actual Hearing

Loss Caused by Environmental Le. Audiol Neurootol 23, 259–269.

https://doi.org/10.1159/000494049

Hyder, O., Chung, M., Cosgrove, D., Herman, J.M., Li, Z., Firoozmand, A., Gurakar, A.,

Koteish, A., Pawlik, T.M., 2013. Cadmium Exposure and Liver Disease among US

Adults. Journal of Gastrointestinal Surgery 17, 1265–1273.

https://doi.org/10.1007/s11605-013-2210-9

IARC, 2012. Cadmium and cadmium compounds, in: Arsenic, Metals, Fibres, and Dusts

IARC Monographs on the Evaluation of Carcinogenic Risks to Humans.

International Agency for Research on Cancer, Lyon, France, pp. 121–145.

Ingersoll, M.A., Malloy, E.A., Caster, L.E., Holland, E.M., Xu, Z., Zallocchi, M., Currier,

D., Liu, H., He, D.Z.Z., Min, J., Chen, T., Zuo, J., Teitz, T., 2020. BRAF inhibition

protects against hearing loss in mice. Sci Adv 6, eabd0561.

https://doi.org/10.1126/sciadv.abd0561

Iryo, Y., Matsuoka, M., Wispriyono, B., Sugiura, T., Igisu, H., 2000. Involvement of the

extracellular signal-regulated protein kinase (ERK) pathway in the induction of

apoptosis by cadmium chloride in CCRF-CEM cells. Biochem Pharmacol 60, 1875–

1882. https://doi.org/10.1016/S0006-2952(00)00510-4

Jiang, C., Yuan, Y., Hu, F., Wang, Q., Zhang, K., Wang, Y., Gu, J., Liu, X., Bian, J., Liu,

Z., 2014. Cadmium induces PC12 cells apoptosis via an extracellular signal-

regulated kinase and c-Jun N-terminal kinase-mediated mitochondrial apoptotic

pathway. Biol Trace Elem Res 158, 249–258. https://doi.org/10.1007/S12011-014-

9918-6

Jiang, J.H., Ge, G., Gao, K., Pang, Y., Chai, R.C., Jia, X.H., Kong, J.G., Yu, A.C.H.,

2015. Calcium Signaling Involvement in Cadmium-Induced Astrocyte Cytotoxicity

and Cell Death Through Activation of MAPK and PI3K/Akt Signaling Pathways.

Neurochem Res 40, 1929–1944. https://doi.org/10.1007/s11064-015-1686-y

Jung, D.J., 2019. Association between cadmium exposure and hearing impairment: a

population-based study in Korean adults. Yeungnam Univ J Med 36, 141–147.

https://doi.org/10.12701/YUJM.2019.00178

Kang, G.H., Uhm, J.Y., Choi, Y.G., Kang, E.K., Kim, S.Y., Choo, W.O., Chang, S.S.,

2018. Environmental exposure of heavy metal (lead and cadmium) and hearing

loss: data from the Korea National Health and Nutrition Examination Survey

(KNHANES 2010–2013). Ann Occup Environ Med 30, 22.

https://doi.org/10.1186/s40557-018-0237-9

Kim, J., Kim, S.H., Johnson, V.J., Sharma, R.P., 2005. Extracellular signal-regulated

kinase-signaling-dependent G2/M arrest and cell death in murine macrophages by

cadmium. Environ Toxicol Chem 24, 3069–3077. https://doi.org/10.1897/04-

503R3.1

Kim, S.-J., Jeong, H.-J., Myung, N.-Y., Kim, M., Lee, J.-H., So, H., Park, R.-K., Kim, H.-

M., Um, J.-Y., Hong, S.-H., 2008. The protective mechanism of antioxidants in

cadmium-induced ototoxicity in vitro and in vivo. Environ Health Perspect 116, 854–

62. https://doi.org/10.1289/ehp.10467

Kim, Y.K., Kim, H.J., Kwon, C.H., Kim, J.H., Woo, J.S., Jung, J.S., Kim, J.M., 2005.

Role of ERK activation in cisplatin-induced apoptosis in OK renal epithelial cells.

Journal of Applied Toxicology 25, 374–382. https://doi.org/10.1002/jat.1081

Kurioka, T., Matsunobu, T., Satoh, Y., Niwa, K., Endo, S., Fujioka, M., Shiotani, A.,

2015. ERK2 mediates inner hair cell survival and decreases susceptibility to noise-

induced hearing loss. Sci Rep 5, 16839. https://doi.org/10.1038/srep16839

Låg, M., Refsnes, M., Lilleaas, E.M., Holme, J.A., Becher, R., Schwarze, P.E., 2005.

Role of mitogen activated protein kinases and protein kinase C in cadmium-induced

apoptosis of primary epithelial lung cells. Toxicology 211, 253–264.

https://doi.org/10.1016/J.TOX.2005.03.012

Lagos-Cabré, R., Ivanova, A., Taylor, C.W., 2020. Ca2+ Release by IP3 Receptors Is

Required to Orient the Mitotic Spindle. Cell Rep 33.

https://doi.org/10.1016/J.CELREP.2020.108483

Lahne, M., Gale, J.E., 2008. Damage-induced activation of ERK1/2 in cochlear

supporting cells is a hair cell death-promoting signal that depends on extracellular

ATP and calcium. Journal of Neuroscience 28, 4918–4928.

https://doi.org/10.1523/JNEUROSCI.4914-07.2008

Lee, K.H., Cotanche, D.A., 1996. Potential role of bFGF and retinoic acid in the

regeneration of chicken cochlear hair cells. Hear Res 94, 1–13.

Lee, W.K., Bork, U., Gholamrezaei, F., Thévenod, F., 2005. Cd(2+)-induced cytochrome

c release in apoptotic proximal tubule cells: role of mitochondrial permeability

transition pore and Ca(2+) uniporter. Am J Physiol Renal Physiol 288, F27–F39.

https://doi.org/10.1152/AJPRENAL.00224.2004

Li, M., Xia, T., Jiang, C., Li, L., Fu, J., Zhou, Z., 2003. Cadmium directly induced the

opening of membrane permeability pore of mitochondria which possibly involved in

cadmium-triggered apoptosis. Toxicology 194, 19–33.

https://doi.org/10.1016/S0300-483X(03)00327-5

Lim, S., Kim, D.G., Kim, S., 2019. ERK-dependent phosphorylation of the linker and

substrate-binding domain of HSP70 increases folding activity and cell proliferation.

Exp Mol Med 51. https://doi.org/10.1038/S12276-019-0317-0

Liu, F., Li, Z.-F., Wang, Z.-Y., Wang, L., 2016. Role of subcellular calcium redistribution

in regulating apoptosis and autophagy in cadmium-exposed primary rat proximal

tubular cells. J Inorg Biochem 164, 99–109.

https://doi.org/10.1016/j.jinorgbio.2016.09.005

Liu, H., Ding, D., Sun, H., Jiang, H., Wu, X., Roth, J.A., Salvi, R., 2014. Cadmium-

induced ototoxicity in rat cochlear organotypic cultures. Neurotox Res 26, 179–89.

https://doi.org/10.1007/s12640-014-9461-4

Liu, W., Xu, C., Ran, D., Wang, Y., Zhao, H., Gu, J., Liu, X., Bian, J., Yuan, Y., Liu, Z.,

2018. CaMKⅡ mediates cadmium induced apoptosis in rat primary osteoblasts

through MAPK activation and endoplasmic reticulum stress. Toxicology 406–407,

70–80. https://doi.org/10.1016/J.TOX.2018.06.002

Liu, Y., Templeton, D.M., 2007. Cadmium activates CaMK-II and initiates CaMK-II-

dependent apoptosis in mesangial cells. FEBS Lett 581, 1481–1486.

https://doi.org/10.1016/j.febslet.2007.03.003

Low, J., Higgs, D.M., 2015. Sublethal effects of cadmium on auditory structure and

function in fathead minnows (Pimephales promelas). Fish Physiol Biochem 41,

357–369. https://doi.org/10.1007/s10695-014-9988-6

Maeda, Y., Fukushima, K., Omichi, R., Kariya, S., Nishizaki, K., 2013. Time Courses of

Changes in Phospho- and Total- MAP Kinases in the Cochlea after Intense Noise

Exposure. PLoS One 8, e58775. https://doi.org/10.1371/journal.pone.0058775

May, L.A., Kramarenko, I.I., Brandon, C.S., Voelkel-Johnson, C., Roy, S., Truong, K.,

Francis, S.P., Monzack, E.L., Lee, F.S., Cunningham, L.L., 2013. Inner ear

supporting cells protect hair cells by secreting HSP70. J Clin Invest 123, 3577–

3587. https://doi.org/10.1172/JCI68480

Mezynska, M., Brzóska, M.M., 2018. Environmental exposure to cadmium-a risk for

health of the general population in industrialized countries and preventive

strategies. Environ Sci Pollut Res Int 25, 3211–3232.

https://doi.org/10.1007/S11356-017-0827-Z

Min, K.-B., Lee, K.-J., Park, J.-B., Min, J.-Y., 2012. Lead and Cadmium Levels and

Balance and Vestibular Dysfunction among Adult Participants in the National

Health and Nutrition Examination Survey (NHANES) 1999–2004. Environ Health

Perspect 120, 413–417. https://doi.org/10.1289/ehp.1103643

Misra, U.K., Gawdi, G., Pizzo, S.V., 2003. Induction of mitogenic signalling in the 1LN

prostate cell line on exposure to submicromolar concentrations of cadmium+. Cell

Signal 15, 1059–1070. https://doi.org/10.1016/S0898-6568(03)00117-7

Montalbano, G., Capillo, G., Laurà, R., Abbate, F., Levanti, M., Guerrera, M.C., Ciriaco,

E., Germanà, A., 2018. Neuromast hair cells retain the capacity of regeneration

870  during heavy metal exposure. Annals of Anatomy - Anatomischer Anzeiger 218,

871   183–189. https://doi.org/10.1016/j.aanat.2018.03.007

872 Monzack, E.L., Cunningham, L.L., 2013. Lead roles for supporting actors: Critical

873   functions of inner ear supporting cells. Hear Res 303, 20–29.

874   https://doi.org/10.1016/J.HEARES.2013.01.008

875 Nicolson, T., 2005. The genetics of hearing and balance in zebrafish. Annu Rev Genet.

876   https://doi.org/10.1146/annurev.genet.39.073003.105049

877 Niewenhuis, R., Prozialeck, W., 1987. Calmodulin inhibitors protect against cadmium-

878   induced testicular damage in mice. Biol Reprod 37, 127–133.

879   https://doi.org/10.1095/BIOLREPROD37.1.127

880 Nugues, C., Helassa, N., Haynes, L.P., 2022. Mitosis, Focus on Calcium. Front Physiol

881   13, 1215. https://doi.org/10.3389/FPHYS.2022.951979/BIBTEX

882 Ó Maoiléidigh, D., Ricci, A.J., 2019. A bundle of mechanisms: Inner-ear hair-cell

883   mechanotransduction. Trends Neurosci 42, 221.

884   https://doi.org/10.1016/J.TINS.2018.12.006

885 Owens, K.N., Coffin, A.B., Hong, L.S., Bennett, K.O., Rubel, E.W., Raible, D.W., 2009.

886   Response of mechanosensory hair cells of the zebrafish lateral line to

887   aminoglycosides reveals distinct cell death pathways. Hear Res 253, 32–41.

888   https://doi.org/10.1016/j.heares.2009.03.001

889 Ozcaglar, H.U., Agirdir, B., Dinc, O., Turhan, M., Kilinçarslan, S., Oner, G., 2001.

890   Effects of cadmium on the hearing system. Acta Otolaryngol 121, 393–7.

891 Raible, D.W., Kruse, G.J., 2000. Organization of the lateral line system in embryonic

892   zebrafish. J Comp Neurol 421, 189–198.

Rasmussen, C.D., Means, A.R., 1989. Calmodulin is required for cell-cycle progression

during G1 and mitosis. EMBO J 8, 73. https://doi.org/10.1002/J.1460-

2075.1989.TB03350.X

Rigon, A., Cordova, F., Oliveira, C., Posser, T., Costa, A., Silva, I., Santos, D., Rossi,

F., Rocha, J., Leal, R., 2008. Neurotoxicity of cadmium on immature hippocampus

and a neuroprotective role for p38 MAPK. Neurotoxicology 29, 727–734.

https://doi.org/10.1016/J.NEURO.2008.04.017

Rothschild, S.C., Lahvic, J., Francescatto, L., McLeod, J.J.A., Burgess, S.M., Tombes,

R.M., 2013. CaMK-II activation is essential for zebrafish inner ear development and

acts through Delta-Notch signaling. Dev Biol 381, 179–188.

https://doi.org/10.1016/j.ydbio.2013.05.028

Ruczaj, A., Brzóska, M.M., 2022. Environmental exposure of the general population to

cadmium as a risk factor of the damage to the nervous system: A critical review of

current data. J Appl Toxicol. https://doi.org/10.1002/JAT.4322

Schaal, N., Slagley, J., Zreiqat, M., Paschold, H., 2017. Effects of combined exposure to

metals, solvents, and noise on permanent threshold shifts. Am J Ind Med 60, 227–

238. https://doi.org/10.1002/ajim.22690

Schmid, C., Alampi, I., Briggs, J., Tarcza, K., Stawicki, T.M., 2020.

Mechanotransduction Activity Facilitates Hair Cell Toxicity Caused by the Heavy

Metal Cadmium. Front Cell Neurosci 14.

https://doi.org/10.3389/FNCEL.2020.00037

Seiler, C., Nicolson, T., 1999. Defective calmodulin-dependent rapid apical endocytosis

in zebrafish sensory hair cell mutants. J Neurobiol 41, 424–434.

Shargorodsky, J., Curhan, S.G., Henderson, E., Eavey, R., Curhan, G.C., 2011. Heavy

Metals Exposure and Hearing Loss in US Adolescents. Arch Otolaryngol Head

Neck Surg 137, 1183–1189. https://doi.org/10.1001/archoto.2011.202

Shih, Y.L., Lin, C.J., Hsu, S.W., Wang, S.H., Chen, W.L., Lee, M.T., Wei, Y.H., Shih,

C.M., 2005. Cadmium toxicity toward caspase-independent apoptosis through the

mitochondria-calcium pathway in mtDNA-depleted cells, in: Annals of the New York

Academy of Sciences. New York Academy of Sciences, pp. 497–505.

https://doi.org/10.1196/annals.1338.043

Smith, J.B., Dwyer, S.D., Smith, L., 1989. Cadmium evokes inositol polyphosphate

formation and calcium mobilization: Evidence for a cell surface receptor that

cadmium stimulates and zinc antagonizes. Journal of Biological Chemistry 264,

7115–7118. https://doi.org/10.1016/S0021-9258(18)83208-2

So, H., Kim, H.J., Lee, J.H., Park, C., Kim, Y., Kim, E., Kim, J.K., Yun, K.J., Lee, K.M.,

Lee, H.Y., Moon, S.K., Lim, D.J., Park, R., 2007. Cisplatin cytotoxicity of auditory

cells requires secretions of proinflammatory cytokines via activation of ERK and

NF-κB. JARO - Journal of the Association for Research in Otolaryngology 8, 338–

355. https://doi.org/10.1007/s10162-007-0084-9

Sonnack, L., Kampe, S., Muth-Köhne, E., Erdinger, L., Henny, N., Hollert, H., Schäfers,

C., Fenske, M., 2015. Effects of metal exposure on motor neuron development,

neuromasts and the escape response of zebrafish embryos. Neurotoxicol Teratol

50, 33–42. https://doi.org/10.1016/j.ntt.2015.05.006

Stawicki, T.M., Owens, K.N., Linbo, T., Reinhart, K.E., Rubel, E.W., Raible, D.W., 2014.

The zebrafish merovingian mutant reveals a role for pH regulation in hair cell

toxicity and function. Dis Model Mech 7, 847–856.

https://doi.org/10.1242/dmm.016576

Sutoo, D., Akiyama, K., Imamiya, S., 1990. A mechanism of cadmium poisoning: the

cross effect of calcium and cadmium in the calmodulin-dependent system. Arch

Toxicol 64, 161–164. https://doi.org/10.1007/BF01974404

Takuwa, N., Zhou, W., Takuwa, Y., 1995. Calcium, calmodulin and cell cycle

progression. Cell Signal 7, 93–104. https://doi.org/10.1016/0898-6568(94)00074-L

Tang, J., Duan, W., Deng, P., Li, H., Liu, C., Duan, Y., Feng, M., Xu, S., 2021. Cadmium

disrupts mitochondrial distribution and activates excessive mitochondrial fission by

elevating cytosolic calcium independent of MCU-mediated mitochondrial calcium

uptake in its neurotoxicity. Toxicology 453, 152726.

https://doi.org/10.1016/j.tox.2021.152726

Tellez-Plaza, M., Navas-Acien, A., Menke, A., Crainiceanu, C.M., Pastor-Barriuso, R.,

Guallar, E., 2012. Cadmium Exposure and All-Cause and Cardiovascular Mortality

in the U.S. General Population. Environ Health Perspect 120, 1017–1022.

https://doi.org/10.1289/ehp.1104352

Vlasits, A.L., Simon, J.A., Raible, D.W., Rubel, E.W., Owens, K.N., 2012. Screen of

FDA-approved drug library reveals compounds that protect hair cells from

aminoglycosides and cisplatin. Hear Res 294, 153–165.

Wang, B., Xiao, J.L., Ling, Y.H., Meng, X.J., Wu, B., Yang, X.Y., Zou, F., 2014. BNIP3

upregulation by ERK and JNK mediates cadmium-induced necrosis in neuronal

cells. Toxicol Sci 140, 393–402. https://doi.org/10.1093/TOXSCI/KFU091

Wang, D.-H., Xu, H., Zheng, Y.-H., Gu, D.-S., Zhu, Y.-J., Ren, Y., Wang, S.-C., Yang,

L., Xu, L.-W., 2020. Environmental exposure to lead and cadmium and hearing loss

in Chinese adults: A case-control study. PLoS One 15, e0233165.

https://doi.org/10.1371/JOURNAL.PONE.0233165

Wang, L., Cao, J., Chen, D., Liu, X., Lu, H., Liu, Z., 2009. Role of oxidative stress,

apoptosis, and intracellular homeostasis in primary cultures of rat proximal tubular

cells exposed to cadmium. Biol Trace Elem Res 127, 53–68.

https://doi.org/10.1007/s12011-008-8223-7

Wang, S.H., Shih, Y.L., Ko, W.C., Wei, Y.H., Shih, C.M., 2008. Cadmium-induced

autophagy and apoptosis are mediated by a calcium signaling pathway. Cellular

and Molecular Life Sciences 65, 3640–3652. https://doi.org/10.1007/s00018-008-

8383-9

Wang, S.H., Shih, Y.L., Lee, C.C., Chen, W.L., Lin, C.J., Lin, Y.S., Wu, K.H., Shih, C.M.,

2009. The role of endoplasmic reticulum in cadmium-induced mesangial cell

apoptosis. Chem Biol Interact 181, 45–51. https://doi.org/10.1016/j.cbi.2009.05.004

Wang, T., Zhu, Q., Cao, B., Cai, Y., Wen, S., Bian, J., Zou, H., Song, R., Gu, J., Liu, X.,

Liu, Z., Yuan, Y., 2022. Ca 2+ transfer via the ER-mitochondria tethering complex

in neuronal cells contribute to cadmium-induced autophagy. Cell Biol Toxicol 38.

https://doi.org/10.1007/S10565-021-09623-Y

Wang, X., Zhu, Y., Long, H., Pan, S., Xiong, H., Fang, Q., Hill, K., Lai, R., Yuan, H.,

Sha, S.H., 2019. Mitochondrial Calcium Transporters Mediate Sensitivity to Noise-

Induced Losses of Hair Cells and Cochlear Synapses. Front Mol Neurosci 11, 469.

https://doi.org/10.3389/FNMOL.2018.00469

Whitfield, T.T., 2002. Zebrafish as a model for hearing and deafness. J Neurobiol.

https://doi.org/10.1002/neu.10123

Xiao, W., Liu, Y., Templeton, D.M., 2009. Pleiotropic effects of cadmium in mesangial

cells. Toxicol Appl Pharmacol 238, 315–326.

https://doi.org/10.1016/J.TAAP.2009.02.005

Xu, B., Chen, S., Luo, Y., Chen, Z., Liu, L., Zhou, H., Chen, W., Shen, T., Han, X.,

Chen, L., Huang, S., 2011. Calcium Signaling Is Involved in Cadmium-Induced

Neuronal Apoptosis via Induction of Reactive Oxygen Species and Activation of

MAPK/mTOR Network. PLoS One 6, e19052.

https://doi.org/10.1371/journal.pone.0019052

Xu, J., Tang, S., Yin, B., Sun, J., Bao, E., 2018. Co-enzyme Q10 upregulates Hsp70

and protects chicken primary myocardial cells under in vitro heat stress via

PKC/MAPK. Mol Cell Biochem 449, 195–206. https://doi.org/10.1007/S11010-018-

3356-2/FIGURES/8

Yamagami, K., Nishimura, S., Sorimachi, M., 1998. Cd2+ and Co2+ at micromolar

concentrations mobilize intracellular Ca2+ via the generation of inositol 1,4,5-

triphosphate in bovine chromaffin cells. Brain Res 798, 316–319.

https://doi.org/10.1016/S0006-8993(98)00445-4

Yang, C.S., Tzou, B.C., Liu, Y.P., Tsai, M.J., Shyue, S.K., Tzeng, S.F., 2008. Inhibition

of cadmium-induced oxidative injury in rat primary astrocytes by the addition of

antioxidants and the reduction of intracellular calcium. J Cell Biochem 103, 825–

834. https://doi.org/10.1002/jcb.21452

Yang, P.M., Chen, H.C., Tsai, J.S., Lin, L.Y., 2007. Cadmium induces Ca2+-dependent necrotic cell death through calpain-triggered mitochondrial depolarization and reactive oxygen species-mediated inhibition of nuclear factor-κB activity. Chem Res Toxicol 20, 406–415. https://doi.org/10.1021/tx060144c

Yang, X.F., Yang, Y.N., 1997. Protective effects of calcium antagonists on cadmium-induced toxicity in rats. Biomedical and Environmental Sciences 10, 402–407.

Yeh, J.H., Huang, C.C., Yeh, M.Y., Wang, J.S., Lee, J.K., Jan, C.R., 2009. Cadmium-induced cytosolic Ca2+ elevation and subsequent apoptosis in renal tubular cells. Basic Clin Pharmacol Toxicol 104, 345–351. https://doi.org/10.1111/j.1742-7843.2009.00391.x

Yuan, Y., Jiang, C., Xu, H., Sun, Y., Hu, F., Bian, J., Liu, X., Gu, J., Liu, Z., 2013. Cadmium-Induced Apoptosis in Primary Rat Cerebral Cortical Neurons Culture Is Mediated by a Calcium Signaling Pathway. PLoS One 8, e64330. https://doi.org/10.1371/journal.pone.0064330

Zhai, C., Lv, J., Wang, K., Li, Q., Qu, Y., 2019. HSP70 silencing aggravates apoptosis induced by hypoxia/reoxygenation in vitro. Exp Ther Med 18, 1013–1020. https://doi.org/10.3892/ETM.2019.7697

Zhang, R., Zhu, Y., Dong, X., Liu, B., Zhang, N., Wang, X., Liu, L., Xu, C., Huang, S., Chen, L., 2017. Celastrol Attenuates Cadmium-Induced Neuronal Apoptosis via Inhibiting Ca2+-CaMKII-Dependent Akt/mTOR Pathway. J Cell Physiol 232, 2145–2157. https://doi.org/10.1002/jcp.25703

Zhao, H., Li, Teng, Wang, K., Zhao, F., Chen, J., Xu, G., Zhao, J., Li, Ting, Chen, L., Li, L., Xia, Q., Zhou, T., Li, H.Y., Li, A.L., Finkel, T., Zhang, X.M., Pan, X., 2019.

AMPK-mediated activation of MCU stimulates mitochondrial Ca2+ entry to promote

mitotic progression. Nature Cell Biology 2019 21:4 21, 476–486.

https://doi.org/10.1038/s41556-019-0296-3

Zhao, H., Liu, W., Wang, Y., Dai, N., Gu, J., Yuan, Y., Liu, X., Bian, J., Liu, Z., 2015.

Cadmium induces apoptosis in primary rat osteoblasts through caspase and

mitogen-activated protein kinase pathways. J Vet Sci 16, 297–306.

https://doi.org/10.4142/JVS.2015.16.3.297

Zine, A., de Ribaupierre, F., 1999. Tissue-specific levels and cellular distribution of

epidermal growth factor receptors within control and neomycin-damaged neonatal

rat organ of corti. J Neurobiol 38, 313–322. https://doi.org/10.1002/(SICI)1097-

4695(19990215)38:3<313::AID-NEU2>3.0.CO;2-O

1040