Abiotic conditions alter effectiveness of a key seed defensive trait against granivores similarly

across many plant species

Eric F. LoPresti[1,2], Madison E. Stessman[2], Sara E. Warren[2], Katherine Toll[3]

1: Department of Biology, University of South Carolina, Columbia, SC, USA.

2: Department of Plant Biology, Oklahoma State University, Stillwater, OK, USA.

3: Dept. Plant Biology, Michigan State University, East Lansing, MI, USA.

Abstract

Environmental conditions may alter the function of many plant traits that drive species

interactions, producing context-dependency in the outcomes of those interactions. Seed

mucilage is a common, convergently-evolved trait found in thousands of plant species. When

wetted, the seed coat swells into a viscid mass; when dried, the mucilage strands strongly

cement the seed to whatever it is in contact with. This binding to the ground has been previously

shown to protect seeds from granivory. Previous research found both that mucilage volume –

and the correlated attachment strength – are higher in species from hot, dry, areas suggesting

an environmental component of this trait's function. Here we (1) quantified the effect of

temperature on attachment across many species in a lab setting, (2) tested the potential

mechanism behind this correlation by accelerating desiccation speed without changing

temperature, and (3) tested whether these relationships introduce context dependency of the

defensive function of mucilage in the field, using field trials with harvester ants. We found that

(1) increasing temperature during mucilage drying strongly reduced the force needed to

dislodge seeds for most species, (2) drying time was likely the driving mechanism behind the

loss of attachment strength at higher temperatures, not temperature *per se*, (3) seeds attached

to substrate during higher temperatures or under accelerated drying conditions were far more

susceptible to granivory. These results show not only the mechanism behind an abiotic

modification of a functional trait of seeds, but this change majorly alters a key interaction

contributing to seed survival. These results add to a small, but growing, literature on the

importance of seed mucilage in seed survival and demonstrate strong and largely predictable

context-dependency in this trait's defensive function.

Introduction

The abiotic environment may alter the function of plant traits important in species interactions,

leading to context-dependency in the outcome of these interactions (Maron et al 2014). Under

one set of conditions, the effect of the trait on the interaction may be very pronounced and

under others, far less so. While it is easy to conclude that "community ecology is a mess, with

so much contingency that useful generalisations are hard to find" (Lawton 1999), it is indeed

likely that species with similar traits will have parallel responses to changes in the abiotic

environment, leading to similar alterations of interactions. Having a mechanistic knowledge of

how trait functions are modified by environmental conditions can make this contingency

predictable and even lead to broad ecological and evolutionary insight about trait function and

trait evolution. Seed traits, often quantified in a dry state from herbarium specimens or seed

envelopes, are usually considered fixed once the seed has matured and because of the slow

rates of cellular processes, far less likely to be affected by the environment than vegetative

characters (though shaped by environmental selection pressures over evolutionary time). Here,

we investigate how the temperature experienced by a moistened seed alters a key functional

seed trait and affects seed survival in the presence of granivores.

Variation in expression of a defensive trait across some abiotic gradient may be genetic (Hahn

and Maron 2016), plastic, or some combination of the two. Here we are mostly concerned with

the plastic component, how local variation in abiotic conditions changes traits and alters the

outcomes of these interactions. Wagner and Mitchell-Olds (2018) showed that local conditions

across common gardens had large effects on defensive glucosinolate concentrations (up to a

doubling!) for many *Boechera* genotypes. In a warming experiment in natural plant communities,

including 132 plants, Descombes et al (2020) demonstrated this treatment changed putatively

defensive traits across most species, but that the outcomes of these changes on interactions

with herbivores were idiosyncratic with respect to the treatment. While these, and many other,

studies nicely demonstrate environmental effects on defensive traits and interactions, making

sense of this context-dependency requires some amount of mechanistic knowledge: how

exactly does the abiotic environment alter the trait's role in the interaction across many

species?

A powerful method for identifying the underlying mechanism of context dependence is to

experimentally manipulate the environment and quantify how it effects traits and species

interactions under different environmental conditions (e.g., McGuire and Agrawal 2005, Mody et

al 2009, Pezzola et al 2017). However, interactions may differ between abiotic conditions in the

absence of trait changes in one partner in the interaction. The classic study done by Lincoln and

Mooney (1984) found that herbivory on a monkeyflower was far greater in sun than shade.

However, none of the hypothesized defensive traits varied concurrently, instead they concluded

that adult butterfly oviposition preferences for sunny areas drove the very strong context-

dependency in this system. In contrast, Louda and Rodman (1996) found that reduced

glucosinolate concentrations in experimentally sun-exposed mustards, coupled with higher

insect abundance in sun, explained the same markedly higher herbivory in higher light areas.

Despite both studies finding higher herbivory in the sun, the former study found the context-

dependency was due to arthropod behavior, while concentrations of a defensive chemical

contributed in the latter, demonstrating the importance of understanding the mechanisms behind

altered interactions.

Though often treated simply as a unit of reproduction, each seed is an individual plant. Most

seeds never survive to reproduce, with a major source of mortality in many systems being

varied granivores (e.g. Tevis 1958). Therefore, defensive traits of the seed may be extremely

important in determining survival. A very common, multifunctional trait in plants - diaspore

mucilage - occurs in thousands of plant species, including many ruderal, cosmopolitan weeds

found on eroded slopes, roadsides, sidewalk cracks, and other disturbed areas. Diaspore

mucilage (referred here as seed mucilage for simplicity), whether on the seed, fruit coat (i.e.

mint nutlets), or accessory fruit (i.e. anthocarps of *Mirabilis*) is a common, convergently evolved

trait across thousands of plant species. These seeds are unassuming and entirely nonsticky

when dry, yet when wetted, the polysaccharide coating swells into a viscid mass (Figure 1, top

row); when dried again, the mucilage strands strongly cement the seed to whatever it is in

contact with (Figure 1, middle row). The composition (whether pectins or cellulose strands),

volume, structure (strands or envelope), and the physical properties of the mucilage layer vary

greatly across species (Western 2012, Kreitschitz et al 2021). These differences have functional

ecological significance – species with larger mucilage volumes generally had greater

attachment force and greater protection from granivorous ants (Pan et al 2021) and erosive

dislodgement (Pan et al 2022). While previous studies have demonstrated mucilage properties

across species drive defensive differences (LoPresti et al 2019, Pan et al 2021), whether the

effects of environmental modification or intraspecific differences in these properties affect

interactions with granivores has not been investigated, nor whether this context-dependency is

idiosyncratic or predictable.

Global, and local, patterns in the distribution of plants with seed mucilage and traits of that

mucilage suggest an environmental influence on the evolution of the trait. Seed mucilage is

found in plants in virtually every terrestrial ecosystem worldwide, yet it is thought to be most

common and most pronounced in species and populations in hotter, dryer areas and in ruderal

plants (e.g. Ellner and Schmida 1981; Al-Shehbaz 1986; Kreitschitz and Valles 2007,

Nordenstram et al 2009; Yang et al 2012). While often-stated, only four quantitative

comparisons have examined this relationship. Murbeck (1919) compared the flora of

Scandinavia (mesic) to North Africa (xeric) and found the latter had a higher proportion of

mucilaginous seeds. Ryding (2001) found a similar result, mucilage was more common, and

more pronounced in mint species from xeric regions compared to mesic regions. Comparing

data from 432 species with mucilage, Pan et al (2021) found that species from areas of higher

temperature, solar radiation, humidity, and fewer wet days per year had higher mucilage

attachment potential. Finally, Villellas and Garcia (2012) examined it at the population scale and

found that populations of *Plantago coronopus* from areas with lower summer precipitation had

higher quantities of mucilage. While these patterns conclusively demonstrate macroevolutionary

and population-level changes in mucilage, the ecological effects of either environmentally-driven

or heritable intraspecific variation is unknown.

Environmental factors clearly affect the functional significance of seed mucilage. In an

environment entirely devoid of moisture, mucilage will never become imbibed and any functions

of imbibition, including defense during the viscid phase (Yang et al 2013; Pan et al 2021),

defense due to mucilage-bound substrate (Fuller and Hay 1983; LoPresti et al 2019), defense

due to binding to the ground (Engelbrecht and Garcia-Fayos 2012; Pan et al 2021), or

resistance to dislodgement (Engelbrecht et al 2014; Pan et al 2022). However, a complete lack

of water at all times would prevent any plant growth, therefore, any seed produced is in an area

that, at least occasionally, has water and occasional mucilage imbibition. The effect of variation

in environmental factors on ecological context-dependency in mucilage is entirely unstudied (but

see Cowley et al 2022 for a related agricultural study), though preliminary data referenced in

Pan et al (2021), state that temperature during drying may affect functional properties of the

mucilage. Though convergently evolved a great many times, it is likely that seed mucilage of

diverse species will respond similarly to environmental factors and that the context-dependency

imparted by this response may affect interactions and may drive evolutionary changes in the

trait.

Here, we first investigate the effect of temperature on seed anchorage across many species

with mucilaginous seeds. Secondly, we investigate physical mechanisms behind that

relationship. Lastly, we test whether the temperature modification of anchorage strength affects

the defensive capabilities of seeds against harvester ants in the field.

Materials and Methods

Seed selection

In order to make these results generalizable, we chose 38 species across 10 plant families

(Acanthaceae, Brassicaceae, Cistaceae, Euphorbiaceae, Lamiaceae, Linaceae, Nyctaginaceae,

Plantaginaceae, Polemoniaceae & Solanaceae). Seeds were obtained commercially or through

local collections. The sources of each seed, as well as which species were used in each of the

following experiments is detailed in Table S1.

Drying temperature effects on anchorage strength

To determine if the environmental conditions experienced by the seed during a drying cycle

affected anchorage of the seed, we subjected seeds of twenty-two species (Table S1) to a

series of temperatures during mucilage drying and then measured the force needed to dislodge

them. These protocols are based on those in Pan et al (2021), the measurements from which

were found to be ecologically-informative. Seeds of each species were wetted for an hour in

order for mucilage to fully imbibe. Ten seeds were then placed onto each of 55 new glass slides

(for a total of up to 550 seeds/species tested); washed slides have a lower dislodgement force

(supplementary data), therefore new slides were used for each trial. The slides were placed in

dehydrators (Ivation, 1000W, 10 tray) and set for nine different temperatures, 5 slides per

temperature/species. They were dried for an hour, sufficient for full drying. The other two

treatments were a room temperature control (5 slides/species), and in the refrigerator (5

slides/species). Even though the dehydrator slides all took <1 hour to dry, all seeds were left at

least overnight to dry before measuring dislodgement force. Dislodgement force was measured

exactly as in Pan et al (2022; a modification of Pan et al [2021)); briefly, the seeds were

dislodged with a shear force applied perpendicularly with a Pesola pressure meter (300, 600,

1000 grams, depending on species) and the force at which dislodgement occurred was

recorded. In these trials, we used 10,000 seeds of the 22 species, though one measurement

was recorded in error, leaving a final sample of 9,999 dislodgement force measurements.

To analyze these data, we used a mixed-model with a gaussian error distribution, with

temperature as a fixed effect and random effect of slide ID nested within species. Given that the

response to temperature may be nonlinear, we used a linear term (temperature), a quadratic

term (temperature2), and an asymptotic term (1/temperature). We tested models with each

combination of these predictors and a no-predictor null model. We also fit a model with each

nonlinear predictor for each species independently; the output from these models were plotted

to show the individual species relationships (for Supplementary Figure 1), but only the overall

model with species as a random effect (as described earlier) is reported.

Accelerated drying via vacuum desiccator, glass slides:

The previous experiment strongly demonstrated that the drying environment temperature

strongly influences how tightly the seed attaches during mucilage drying. However,

mechanistically, while changing temperature could change the attachment strength directly, it

also affects drying speed - at higher temperatures, seeds often appeared fully dry in <10

minutes, whereas the room temperature trials often took 5+ hours for the seeds to appear dry

(and they were kept overnight to guarantee that by the time of testing). In order to parse out

whether the broad reductions in dislodgement force at higher temperatures were due to this

accelerated drying time or due to temperature effects directly, perhaps on mucilage structure,

we manipulated these two variables independently. To do this, we dried seeds onto glass slides

in two treatments. For each species, we prepared 5 slides, each with 12 seeds, in each of the

following ways. Treatment 1 was a control, with seeds left to dry in the lab overnight (as done in

the prior experiment, for the 20C treatment). Treatment 2 was a faster desiccation, without

raising the temperature (for a comparison with the temperature results). For this treatment,

slides of seeds were placed into a vacuum desiccator that contained drierrite, a desiccant. For

these trials, we tested 1181 seeds of 10 species.

Accelerated drying on sandpaper, via heating and vacuum desiccator:

Additionally, we manipulated the substrate, and ran an additional test with those two treatments,

plus a third (a 75C heat treatment, as in the prior section) on glass slides covered in 80-grit

sandpaper (3M - 80 grit), as used in the ant trials. As in the other treatments, these were dried

overnight, though actual drying time for the latter two treatments, visually, was less than an

hour. During drying, humidity in the vacuum chambers was 18%-21%, oven 10-18%, and lab

45%-65%. The force needed to dislodge was measured for these seeds in the same manner as

previously. All species used in ant trials were planned to be tested in this way in order to

quantitatively link this reduction to field performance, but the vacuum went out in the building

before all could be tested. In these trials, we test 1725 seed of 10 species.

For both the glass slide test and the sandpaper tests, we analyzed the overall treatment effect

on dislodgement force, using a mixed-model with a guassian error distribution and treatment as

a fixed effect, with species as a random effect (adding date as a random effect did not improve

the fit). In order to calculate post-hoc comparisons within species, we did a linear regression of

dislodgement force as a function of the interaction between treatment and species, calculating

posthoc comparisons of treatment within species using a Tukey test in the R package

emmeans.

Seed defense trials

To test whether the relationships established in the first two parts of this study translated into a

context-dependent interaction with granivores, we did granivory trials using the common

harvester ant, *Pogonomyrmex barbatus*. These experiments occurred along gravel paths and

mown lawns at the Oklahoma State University Botanical Garden, in Stillwater, OK, where *P.*

barbatus is common. Seven nearby nests were selected for use in this experiment. Seed

treatments were prepared as in the previous experiment (control, vacuum desiccated, heat

dried) with one important difference. Instead of being attached directly to glass slides, the glass

slides again had a layer of sandpaper (3M - 80 grit) glued on, which is a more appropriate

substrate proxy than glass for ant foraging. A trial consisted of three slides (containing 5-12

seeds, depending on seed size), one of each treatment, that were placed in a semicircle ~0.5m

from the entrance of the nest. As in LoPresti et al (2019), these were monitored until one

treatment was entirely consumed, or for two or four hours, depending on the ant activity (i.e.

some days, probably driven by colder weather, very few seeds would be consumed in two hours

and those trials were lengthened to four hours). Multiple trials of different species were done at

the same nests simultaneously, but a species was never repeated at the same nest on the

same day. Two species (*Nicandra physalodes* & *Salvia sclarea*) had no vacuum treatment, only

the heat and control treatments. The sample size was 7699 seeds in total from 34 species.

We analyzed the ant trial results using seed survival as a binomial response variable (survived

or was removed) with treatment as a fixed effect and random effects of date, ant nest, species,

as well as a composite ID for each slide nested within one for each species trial (i.e., *Camelina*

sativa at ant nest #3 on July 7[th]). We compared treatment means using a Tukey's posthoc

comparison of means.

Data and analyses

All data and code will be available on FigShare. All analyses were conducted and figures made

in R vers. 4.1.0. Specific model parameters are described in the sections above.

Results

Higher drying temperatures reduce anchorage strength

Generally, higher drying temperatures greatly reduced the amount of force necessary to

dislodge the seeds (Figure 2; Supplementary Figure 1). A linear mixed-model with drying

temperature as linear, exponential, and asymptotic fixed effects and species as a random

intercept (i.e. Force ~ Temp + Temp2 + 1/Temp + [1|Species]) fit better than any simpler model

and each of these coefficients was highly significant ($p < 0.01$ for each). However, individual

species responses varied greatly (Supplementary Figure 1). The effect of temperature was

especially pronounced in the plantains (*Plantago* spp.) and flaxes (*Linum* spp.). The highest

temperature treatment for *P. aristata* required only 1.5% the force of the lowest temperature

treatment for dislodgement! *P. rhodosperma* (5.5%), and *P. virginica* (8.7%) were similar, with

P. patagonica having a slightly smaller effect (26.2%). The highest temperature for *Linum lewisii*

had a mean dislodgement force of just 2.8% of that at the lowest temperature, with *L.*

grandiflorum at 13.3%, *L. usitassimum* at 19.1%, and *L. perenne* at 30.0%. In contrast, other

groups were far less affected; certain species showed no significant decrease in dislodgement

force at higher temperatures (i.e. *Helianthemum nummularium*). Nevertheless, on average

seeds dried at the highest temperatures required ~33% of the force to dislodge compared to

those dried in the refrigerator or at room temperature, demonstrating that attachment is strongly

dependent on temperature.

Humidity manipulations to drying speed, independent of temperature

Accelerated drying via vacuum desiccator on glass slides:

Using a vacuum desiccator greatly accelerated drying speed while remaining at room

temperature, isolating the effect of drying speed from temperature (Figure 3). Seeds in the

vacuum desiccated treatment required significantly less force to dislodge than those at room

temperature overall (Mixed-effect linear model, with species as a random effect, posthoc

comparison of treatment means, $p < 0.001$), mirroring the results of higher temperatures. Post-

hoc comparisons of treatment means were significant for six of the ten species tested and the

means of the control groups were higher than the vacuum for nine of the ten species tested

(Figure 3).

Accelerated drying via vacuum desiccator and heating on sandpaper:

Accelerated drying, via heat and vacuum dehydration, greatly reduced the force needed to

dislodge the seed (Mixed-effect linear model, with species as a random effect, posthoc

comparison of treatment means, $p < 0.001$). The control group had a higher mean dislodgement

force than the heat and vacuum treatments in all of the species tested; post-hoc comparisons

the control group showed it differed significantly from the heat treatment in seven of ten species,

the same number (and same species) differing in control versus vacuum treatments.

Functional role of attachment drying speed in defense against granivory

Accelerated drying due to heat or vacuum increased granivory on seeds (Figure 5)

compared to the control group (Tukey's post-hoc comparison of means, all pairwise

comparisons $p < 0.01$). Seeds dried in the vacuum desiccator were also removed at a

significantly higher rate than those in the heat-dried treatment. This effect, however,

was not uniform – some were removed nearly completely, others barely removed, and

the treatment effects differed depending on the species. The overall effect, with species

as one of the random effects, was strongly significant, yet it may be useful to examine

the species separately to determine the generality of this result (supplementary figure

2). While only 56% of the species (19/34) showed a significant reduction in survival in

the heat or vacuum compared to the control (post-hoc comparisons in an interactive

model), 68% (23/34) had a higher mean survival in control than heat, with only 29%

lower survival (10/34, 1 had no difference). This was a significant result using a sign-test

($p=0.035$). Mirroring the overall results, the difference was even stronger between the

control and vacuum; 91% (29/32) of species had higher survival of controls than

vacuum-dried, with the remaining 9% exactly equal (3/32), a highly significant difference

with a sign test ($p<0.001$).

Discussion

Environmental conditions modified the ecological function of an important seed trait –

higher drying temperatures led to lower seed survival in the presence of harvester ants,

due to easier removal of seeds. As hypothesized, this context dependency was largely

predictable across species, though the mucilage traits differ greatly across these many

evolutionary origins (Grubert 1974, Kreitschitz et al, Pan et al 2021). While not every

species showed a significant reduction in dislodgement force at higher temperatures or

faster drying speeds, most species had a measurable reduction. Similarly, the reduction

in defensive efficiency was rather uniform for those species palatable to the ants, with

few species having either treatment with lower attachment strength (i.e. high temp or

vacuum dried) surviving at higher levels than the room control treatment. Unlike a

previous study that found an effect of attachment strength in interspecific comparisons

(Pan et al 2021), in which other traits surely played a large role, the intraspecific

comparisons here are much stronger support for a protective benefit of strength of

attachment as we manipulated it within each species. The interplay between this

widespread trait - found in 1000's of plant species (Grubert 1983) - the environment,

and interacting organisms is an exciting topic that we hope these results inspire future

studies of.

The environment affects mucilage attachment strength

High temperatures greatly reduced the force required to dislodge seeds. We

experimentally demonstrated the mechanisms underlying the modification of the

interaction - higher temperatures led to weaker attachment force, via a decrease in

drying time. While these two factors are mechanistic at some level, this is not a full

physical explanation for how this reduction occurs - that would require further study.

However, we have a strong mechanistic hypothesis integrating these results and

previous research. In interspecific comparisons of over fifty species, attachment

strength was strongly correlated with mucilage volume (Pan et al 2021, 2022). Mucilage

volume probably affects attachment strength by altering the area of contact

(hypothesized in Grubert 1974); therefore, it is likely that the rapid decline in volume at

higher temperatures or at low humidity simply led to a smaller area of substrate that the

mucilage strands were in contact with. For this reason, we can also hypothesize that

other environmental factors that affect mucilage volume (such as pH, ion

concentrations, number of wetting-drying cycles: Grubert 1974, Muñoz et al 2012) or

area that the mucilage contacts (i.e., seed size or substrate grit, Stessman unpublished

data) would similarly alter interactions and subsequent seed success.

While most plants cannot survive very high temperatures, substrate surfaces, where

many seeds are, reach the highest treatments we applied (Nobel et al 1986, pers. obs.).

While it is unlikely that a seed would be imbibed at that temperature (as any substrate

reaching that temperature requires abundant sunshine), it is likely that many of the

higher temperatures we used are regularly encountered by seeds and harvester ants

(Whitford and Ettershank 1975) and that this fundamental interaction is therefore likely

to be greatly modified by temperature in realistic systems. Of course, temperature itself

was not the ultimate driver, and while seeds are rather unlikely to encounter vacuum

chambers of desiccant in the real world, realistic differences in humidity are also

certainly likely to modify these interactions as they would alter drying speed.

Temperature may affect other aspects of the functional ecology of seed mucilage

besides granivory. Few studies have examined attachment strength of mucilage (but

see Grubert 1974, Kreitschitz et al 2021, Pan et al 2021) and none have examined the

effect of temperature, but it is well-known that other seed mucilage properties change

with temperature in its food science, pharmaceutical and industrial applications.

Temperature-dependent changes in physical properties of mucilage have been

characterized for economically-important species used in this study (i.e. flax, *Linum*

usitatisimum: Mazza and Biliaderis 1989, chia, *Salvia hispanica*: Muñoz et al 2012, and

arugula, *Eruca sativa*, Koocheki et al 2012). However, few studies have examined the

effects of temperature on ecologically-important processes involving seed mucilage.

The two studies that have both concern germination, a process that mucilage may

either promote or retard (Western 2012). Gorai et al (2014) examined germination of

Henophyton deserti under seven temperature treatments and found germination

differences between mucilage intact and demucilaged seeds under only one

temperature. Similarly, Bhatt et al (2016) found a significant effect of the interaction

between temperature and mucilage presence in two of five species tested (*Lavandula*

subnuda and *Lepidium aucheri*). Given the diverse roles of mucilage (Grubert 1974;

Western 2012), it is likely that much further complexity exists and when thinking about

mucilage's roles in the field, the effect of simultaneous functions, including defense and

germination, ought to be examined concurrently.

Reduced strength of attachment affects seed survival

Unsurprisingly, the effect of this reduced attachment strength due to higher

temperatures or more rapid drying made the seeds far easier to exploit by ants. This

result makes good sense; a seed bound to the ground requires a dislodgement force

between 10,000-30,000x its own weight to dislodge (Grubert 1974), and interspecific

differences in attachment strength explain protection from ants well (Pan et al 2021). An

ant cannot, therefore, simply pick up even the most weakly-bound seeds (i.e. <10 g

required to dislodge). Instead, they clip the dried mucilage strands (Figure 1), increasing

the time needed to collect and probably greatly increasing the cost of collecting the

seed (see supplementary information in Pan et al 2021 for a video of this behavior). It is

worth stressing that we were not explicitly testing the defensive benefit of mucilage *per*

se, which would have had a non-bound treatment. Previous experiments have found

very strong effects of binding on survival across many species (e.g., Gutterman and

Shem-Tov 1997, Engelbrecht and Garcia-Fayos 2012, Pan et al 2021), and ants

certainly struggled and gave up on tightly bound seeds in this experiment (LoPresti,

Stessman, pers. obs.) and therefore the defensive function was assumed, and here we

investigated only the context-dependency of the benefit.

Attachment strength, however, is far from the only defensive trait that seeds have. The

drying environment had little effect on survival of a few species in the study

(Supplementary Figure 2), most notably those species that were not collected at all

(*Ocimum americanum* & *Plantago ovata*), or collected in extremely small numbers

(*Cobaea scandens*, *Lepidium densiflorum, Gilia tricolor, Plantago aristata*, & *Salvia*

coccinia). For *C. scandens*, the seeds were simply too large to be easily exploited, but

for others, the ants simply rejected them after antennating or palpating them with their

mandibles (pers. obs.). It is possible - even likely - that the mucilage layer plays a role in

this advertisement of unpalatability as well. It is both the outer layer of the seed, which

is first contacted by granivores and therefore, one that may deter them (see LoPresti

2017). In fact, the diversity of secondary chemistry in mucilage (e.g., Wardle et al 1991),

could be an advertisement of unpalatability. Mucilage also protects seeds when imbibed

(Yang et al 2013, LoPresti et al 2019), and when it binds substrate particles to the seed

(Fuller and Hay 1983, LoPresti et al 2019). Therefore, mucilage may play a greater role

in defense than simply attachment strength and the chemistry and other physical

aspects of this layer deserve greater attention as a multifunctional defensive trait.

What is the environment like when seeds become imbibed?

Many mucilaginous seeds simply drop off plants upon dehiscion (e.g. Fuller and Hay

1983), though there may be some plant control of when this occurs. *Cardamine hirsuta*

seeds are ballistically dispersed, and while Vaughn et al (2011) suggest that herbivore

touch stimulates them in natural populations, our observations following rain events in

Oklahoma, US, suggest that many siliques rupture and the seeds disperse during rain

storms. These highly mucilaginous seeds are thus imbibed in the wet environment and

often cemented to surrounding vegetation or objects (tree trunks, fallen wood, the

ground, e.g. Figure 6). Rain may be a common mechanism for mucilaginous seeds to

disperse from a plant, other examples include the dispersal of many Aizoaceae, *Thlaspi*,

many Lamiaceae, *Collomia*, and others (Parolin 2005, LoPresti, pers. obs.). As these

seeds leave the plants during wet periods, the mucilage begins imbibing quickly (e.g.

Yang et al 2013) and the seeds get firmly stuck to a substrate during this initial dispersal

period.

Despite repeated wetting-drying periods due to precipitation and even morning dew

(Grubert 1974, Gutterman and Shem-Tov 1996, Huang et al 2008), those seeds that

attach at that point are unlikely to move far. While mucilage is rehydratable, seeds

attached to a substrate are not easily dislodged. The exact mechanism for this

phenomenon deserves further study; Pan et al (2022) found that even after a week of

being under constant water flow, most previously dried seeds remained firmly stuck to

tiles. This unexpected result, coupled with observations of mucilaginous seeds firmly

attached to substrates and objects naturally, demonstrates that the attachment of these

seeds is both ecologically realistic and that the conditions that they experience during

that initial drying period probably determine their subsequent success.

Do plants that regularly experience high temperatures evolve to have more mucilage

and stronger attachment?

Previous research found that seed mucilage of species from warmer areas had stronger

attachment and greater mucilage volume than that of species from cooler areas (Pan et

al 2021, also see Villelas and Garcia 2012 for population level differences). Coupled

with the findings in this study of stronger attachment at lower drying temperatures, we

think it is reasonable to assume that species or populations regularly experiencing

higher temperatures may need greater attachment strength to obtain the same

protection that may come with the rapid drying in warmer areas. For example, [an

example calculation here]. Therefore, we believe that in addition to the environment

modifying the function of mucilage, it also likely is a potent selective force that may, in

part, drive the diversity of mucilage properties we see across plants (Grubert 1974,

Western 2012, LoPresti et al 2019, Pan et al 2021, 2022).

While differences in attachment strength are likely to drive differential survival between

seeds, it is likely that mucilage volume, as the underlying trait strongly correlating with

force needed to dislodge the seed (Pan et al 2021, 2022) is the heritable component

that selection would act on. Mucilage volume differs greatly among species, populations

and between individuals, likely indicating some degree of heritability (e.g. Villelas and

Garcia 2012; Pan et al 2021,2022). Population examinations, especially of widespread

species spanning a large climatic range, (e.g. Villelas and Garcia 2012), or

examinations of a complete clade would be the strongest tests of this hypothesis.

Fortunately, many mucilaginous-seeded species are nearly cosmopolitan as

introductions (e.g., *Capsella bursa-pastoris*, *Euphorbia prostrata*, *Glechoma heracea,*

Arabidopsis thaliana, Plantago coronopus, lanceolata, & major, Prunella vulgaris) or

extremely widespread naturally (e.g., *Plantago maritima*) and present rich opportunities

to study.

Future directions

The results of these experiments demonstrate strong context-dependency in the

defensive function of seed mucilage, driven by altered drying speed. This totally new

line of research brings up more questions than it answers. We suggest that future

research investigate:

1) How these temperature-mediated physical changes are affected by different

chemical compositions of mucilage and mucilage location (e.g., Phan and Burton

2019, Phan et al 2021).

2) The interplay between attachment strength and area of attachment, with

consideration of both mucilage volume and seed shape (i.e., the flat seeds of flax

versus the nearly spherical seeds of chia).

3) How much attachment strength affects foraging decisions of ants (i.e., by

changing the costs of foraging).

4) Functional consequences of other factors that alter attachment strength (i.e., pH,

substrate, wetting-drying cycles) for granivory and other functions of mucilage.

5) How attachment strength and seed survival vary across a season for seeds in real populations, due to dial and seasonal differences in temperature, precipitation, and humidity.

Acknowledgements

Vincent Pan discovered the temperature and dislodgment force relationship and encouraged us to pursue this study. We thank Gabby Barber, Addison Darby, Michael Foisy, Sierra Jaeger, and Lizz Waring for feedback on the study and manuscript. Michelle Dextraze collected the *Plantago rugelii* seeds. This study was funded by the OSU College of Arts and Sciences AURCA program, with special thanks to Rachel Eaton, and with start-up funds to EFL from Oklahoma State University.

References

Bhatt A, Santo A, Gallacher D. 2016. Seed mucilage effect on water uptake and germination in five species from the hyper-arid Arabian desert. Journal of Arid Environments 128: 73–79.

Engelbrecht M, Bochet E, García-Fayos P. 2014. Mucilage secretion: an adaptive mechanism to reduce seed removal by soil erosion?: mucilage secretion and seed removal. Biological Journal of the Linnean Society 111: 241–251.

Engelbrecht M, García-Fayos P. 2012. Mucilage secretion by seeds doubles the chance to escape removal by ants. Plant Ecology 213: 1167–1175.

García-Fayos P, Engelbrecht M, Bochet E. 2013. Post-dispersal seed anchorage to

soil in semiarid plant communities, a test of the hypothesis of Ellner and Shmida.

Plant Ecology 214: 941–952.

Gorai M, El Aloui W, Yang X, Neffati M. 2014. Toward understanding the ecological role

of mucilage in seed germination of a desert shrub Henophyton deserti:

Interactive effects of temperature, salinity and osmotic stress. Plant and Soil 374:

727–738.

Grubert M. 1974. Studies on the distribution of myxospermy among seeds and fruits of

Angiospermae and its ecological importance. Acta Biologica Venezuelica 8:

315–551.

Grubert M. 1981. Mucilage or gum in seeds and fruits of angiosperms: a review.

Munich, Germany: Minerva-Publikation.

Gutterman Y, Shem-Tov S. 1997. The efficiency of the strategy of mucilaginous seeds

of some common annuals of the negev adhering to the soil crust to delay

collection by ants. Israel Journal of Plant Sciences 45: 317–327.

Huang Z, Boubriak I, Osborne DJ, Dong M, Gutterman Y. 2008. Possible role of

pectin-containing mucilage and dew in repairing embryo DNA of seeds adapted

to desert conditions. Annals of Botany 101: 277–283. doi:10.1093/aob/mcm089.

Kreitschitz A, Kovalev A, Gorb SN. 2015. Slipping vs sticking: Water-dependent

adhesive and frictional properties of Linum usitatissimum L. seed mucilaginous

envelope and its biological significance. Acta Biomaterialia 17: 152–159.

Kreitschitz A, Haase E, Gorb SN. 2021a. The role of mucilage envelope in the

endozoochory of selected plant taxa. The Science of Nature 108: 2.

Kreitschitz A, Kovalev A, Gorb SN. 2021b. Plant seed mucilage as a glue: Adhesive

properties of hydrated and dried-in-contact seed mucilage of five plant species.

International Journal of Molecular Sciences 22: 1443.

LoPresti EF. 2016. Chemicals on plant surfaces as a heretofore unrecognized, but

ecologically informative, class for investigations into plant defence. Biological

Reviews 91: 1102–1117.

LoPresti EF, Pan V, Goidell J, Weber MG, Karban R. 2019. Mucilage-bound sand

reduces seed predation by ants but not by reducing apparency: a field test of 53

plant species. Ecology 100: e02809.

Pan VS, McMunn M, Karban R, Goidell J, Weber MG, LoPresti EF. 2021.

Mucilage binding to ground protects seeds of many plants from harvester ants: A

functional investigation. Functional Ecology 35: 2448–2460.

Phan, J.L., Cowley, J.M., Neumann, K.A., Herliana, L., O'Donovan, L.A. and Burton,

R.A., 2020. The novel features of Plantago ovata seed mucilage accumulation,

storage and release. Scientific Reports, 10:1-14.

Phan, J.L. and Burton, R.A., 2018. New insights into the composition and structure of

seed mucilage. Annual Plant Reviews Online, pp.63-104.

Ryding O. 2001. Myxocarpy in the Nepetoideae (Lamiaceae) with notes on

myxodiaspory in general. Systematics and Geography of Plants 71: 503–514.

Villellas J, García MB. 2013. The role of the tolerance-fecundity trade-off in maintaining

intraspecific seed trait variation in a widespread dimorphic herb. Plant Biology 15:

899–909.

Wardle DA, Ahmed M, Nicholson KS. 1991. Allelopathic influence of nodding thistle

(Carduus nutans l) seeds on germination and radicle growth of pasture plants.

New Zealand Journal of Agricultural Research 34: 185–191.

Western TL. 2012. The sticky tale of seed coat mucilages: production, genetics, and

role in seed germination and dispersal. Seed Science Research 22: 1–25.

Yang X, Baskin CC, Baskin JM, et al. 2013. Hydrated mucilage reduces post-dispersal

seed removal of a sand desert shrub by ants in a semiarid ecosystem. Oecologia

173: 1451–1458.

Yang X, Zhang W, Dong M, Boubriak I, Huang Z. 2011. The achene mucilage hydrated

in desert dew assists seed cells in maintaining DNA integrity: Adaptive strategy

of desert plant Artemisia sphaerocephala. PLoS ONE 6.

Zhou Z, Xing J, Zhao J, Liu L, Gu L, Lan H. 2021. The ecological roles of seed

mucilage on germination of Lepidium perfoliatum, a desert herb with typical

myxospermy in Xinjiang. Plant Growth Regulation in press. doi:10.1007/s10725-

021-00702-y.

Figures

Figure 1: Top row– left to right: mucilage imbibed *Croton captitatus* seed, *Plantago*

ovata seed, *Salvia lyrata* nutlet & *Mirabilis nyctaginea* anthocarp. Middle row—left to

right: mucilage-bound *Ocimum americanum* nutlet, *Plantago aristata* seed, *Mirabilis*

nyctaginea anthocarp, and *Salvia azurea* nutlet. Bottom row—Left: harvester ant

clipping mucilage strands of *Plantago rhodosperma*. Right: with visible dried mucilage

hanging from a dislodged *Salvia lyrata* nutlet.

Figure 2: Overall relationship, between drying temperature and force necessary to

dislodge a seed from a new glass slide for the 16 species tested. The overall model

prediction is plotted, species was treated as a random effect. For plots of individual

species responses, see supplementary figure 1.

Figure 3: Force necessary to dislodge seeds dried to glass slides at control, room temperature

(room) and in vacuum desiccators with Drierite (vac). Letters indicate post-hoc comparison of

means for each species, from a single interactive model.

Figure 4: Force necessary to dislodge seeds dried to sandpaper at room temperature (Control),

in a drying oven at 70C (HighTemp) and in vacuum desiccators with Drierite (Vac). Letters

indicate post-hoc comparison of means for each species, from a single interactive model.

Figure 5: Survival of seeds exposed to harvester ants, when dried to dried to sandpaper at room

temperature (Control), in a drying oven at 70C (Heat) and in vacuum desiccators with Drierite

(Vacuum). Each treatment differed significantly from each other in a post-hoc comparison of

means. For plots of individual species responses, see supplementary figure 2.

Figure 6: *Plantago rhodosperma* seeds naturally mucilage-bound to a rock, 6-July-2022. The

previous precipitation was ~1 cm, 10-June, therefore, these must have remaining attached for at

least that long, if not longer. Boomer Lake Park, Stillwater, Oklahoma, USA. Photo: KT

Supplement

Supplementary Table 1: Species used in each study, as well as the source of the seed.

Supplementary Figure 1: Temperature dependence of 16 of the 22 species tested (all data is

archived). The other species were either too weakly attached to measure reliably, we had too

little seed to do a full temperature spread, or all treatments were not completed before SW

graduated.

Supplementary Figure 2: Survival of seeds exposed to harvester ants, when dried to dried to

sandpaper at room temperature (Control), in a drying oven at 70C (Heat) and in vacuum

desiccators with Drierite (Vacuum). Letters indicated treatments which differed significantly from

each other in a post-hoc comparison of means.