Four decades of phenology in an alpine amphibian: trends, stasis and climatic

drivers

Omar Lenzi[1], Kurt Grossenbacher[2], Silvia Zumbach[3], Beatrice Lüscher[4], Sarah Althaus[4],

Daniela Schmocker[5], Helmut Recher[6], Marco Thoma[7], Arpat Ozgul[1], Benedikt R. Schmidt[1,3]

[1]Department of Evolutionary Biology and Environmental Studies, University of Zurich,

Winterthurerstrasse 190, 8057 Zurich, Switzerland

[2]Eichholzstrasse 18F, 3027 Bern, Switzerland

[3]Info Fauna Karch, Bellevaux 51, 2000 Neuchâtel, Switzerland

[4]Schwand 3, 3110 Münsingen, Switzerland

[5]Impuls AG, Seestrasse 2, 3600 Thun, Switzerland

[6]Abteilung Biodiversität und Landschaft, Bundesamt für Umwelt, Worblentalstrasse 68, 3063

Ittigen, Switzerland

[7]Gesellschaftsstrasse 89, 3012 Bern, Switzerland

Corresponding author's email address: omar.lenzi@ieu.uzh.ch

ORCID: OL, 0000-0003-1938-6786
MT, 0000-0003-2729-2376
AO, 0000-0001-7477-2642
BRS, 0000-0002-4023-1001
21

22 Word count (abstract through acknowledgements): 7'426

23 Number of citations: 93

24 Number of tables in main document: 3

25 Number of figures in main document: 3

Abstract

1. Strong phenological shifts in response to changes in climatic conditions have

been reported for many species, including amphibians, which are expected to

breed earlier. Phenological shifts in breeding are observed in a wide number

of amphibian populations, but less is known about populations living at high

elevations, which are predicted to be more sensitive to climate change than

lowland populations.

2. The goal of this study is to assess the main factors determining the timing of

breeding in an alpine population of the common toad (*Bufo bufo*) and to

describe the observed shifts in its breeding phenology.

3. We modelled the effect of environmental variables on the start and peak dates

of the breeding season using 39 years of individual-based data. In addition,

we investigated the effect of the lunar cycle, as well as the individual variation

in breeding phenology. Finally, to understand the role of the genetic

component of the breeding phenology we calculated the repeatability of the

timing of arrival at the breeding site.

4. Breeding advanced to earlier dates in the first years of the study but the trend

continued only until the mid 1990s, and stabilised afterwards. Overall, toads

are now breeding on average around 30 days earlier than at the start of the

study period. High temperatures and low snow cover in winter and spring, as

well as reduced spring precipitation were all associated with earlier breeding.

Additionally, we found evidence of males arriving on average before females

at the breeding site but no clear and strong effect of the lunar cycle. We only

found weak evidence of among-individual variation in shifts in the breeding

phenology, as well as a low repeatability of arrival timing.

5. Our findings show that the observed changes in breeding phenology are

strongly associated with the environmental conditions. These results

contribute to filling a knowledge gap on the effects of climate change on

alpine amphibian populations. Moreover, we show that changes in phenology,

especially in the mountains, can be hard to predict as local microclimatic

conditions do not necessarily reflect the observed global climatic trends.

Keywords: Phenology, climate change, *Bufo bufo*, mountains, principal component

analysis, repeatability

Introduction

Phenology refers to the timing of periodical events (e.g., seasonal migration, spring flowering) in relation to biotic and abiotic factors, and is a key element of the life cycle in a multitude of organisms. Phenology is normally determined by a combination of a genetic and an environmental component (Quinn & Wetherington, 2002; Tang et al., 2016). Thus, climate change can shift the phenology of many species, potentially leading to mismatches between demand and availability of resources (Parmesan & Yohe, 2003; Visser & Gienapp, 2019; Iler et al., 2021). These shifts can have large effects on the demography of populations, as individuals cannot benefit from the optimal conditions at the right time, with consequences on their fitness (Visser & Gienapp, 2019; Iler et al., 2021). Therefore, it is important to describe and quantify phenological shifts and their causes.

Phenology has a key role in amphibians as well, especially in species living in temperate regions, where various aspects of the annual cycle are determined by seasonality (Duellman & Trueb, 1986; Gotthard, 2001; Hartel et al., 2007). The environmental component is more important than the genetic component in explosive breeders (*sensu* Wells, 1977). In fact, explosive breeders reproduce once a year around springtime and the timing is linked to specific environmental signals such as increasing day length, temperature, and rainfall, which can trigger the migration of amphibians from the hibernation sites to the breeding ponds (Semlitsch, 1985; Oseen & Wassersug, 2002; While & Uller, 2014; Ficetola & Maiorano, 2016). Other important environmental factors affecting the timing of breeding in explosive breeders can be the lunar cycle (Grant et al., 2009; Green et al., 2016; Arnfield et al., 2012; Jarvis et al., 2021) or the hydrological cycle of breeding ponds (Semlitsch et

al., 1993). Previous studies have also identified a possible genetic component in

triggering the migration to the breeding site and thus the start of the breeding season

(Heusser & Ott, 1968; Semlitsch et al., 1993; Phillimore et al., 2010). Breeding

phenology also shows individual variation, as the animals will arrive at different times

at the breeding site. The causes of individual-level variation are multifold and can

include genetics (Heusser & Ott, 1968; Semlitsch et al., 1993), sex and size (Loman

& Madsen, 1986), body condition (Kokko, 1999) as well as features of the

hibernation site, such as distance from the breeding site, which in *Bufo bufo* can be

up to more than 1000 m (Sztatecsny & Schabetsberger, 2005; Kovar et al., 2009).

While most studies on amphibians across species and locations have found

earlier spring breeding in response to climate change (Beebee, 1995; Blaustein et

al., 2001; Parmesan, 2007; While & Uller, 2014), phenological delays have also been

observed (e.g., Arnfield et al., 2012; Arietta et al., 2020). The direction and

magnitude of these phenological shifts are therefore variable among and within

species, as they can depend on the specific environmental conditions that the

populations are experiencing at the local scale, or on the genetic structure of said

populations (Phillimore et al., 2010; Bison et al., 2021).

Shifts in phenology can have adverse effects on amphibians, as phenological

mismatches can affect predator-prey dynamics and food availability (Todd et al.,

2011; Reinhardt et al., 2015; Jara et al., 2019; Visser & Gienapp, 2019). Early

breeding can expose eggs and hatched tadpoles more frequently to late frost events,

thus increasing mortality (Muir et al., 2014; Bison et al., 2021). On the other hand, in

the absence of frost or drying events, earlier breeding might be beneficial as it allows

post-metamorphic toadlets more time to fully develop in summer before hibernation

(Reading & Clarke, 1999; Reading, 2010). Delayed breeding can also have a

negative outcome on the population, as it can result in increased mortality in

juveniles that could not fully grow before their first hibernation (Morin et al., 1990;

Garner et al., 2011; Sinsch & Schäfer, 2016). Even though this phenomenon can be

compensated in some cases with an accelerated growth rate, this can come at the

cost of reduced defences against predation (Orizaola et al., 2016). Thus,

phenological shifts and their causes should be identified and better understood, as

they can help design and prioritise conservation and management actions.

The consequences of phenological shifts could be exacerbated in ecosystems

less resilient to climate change. Mountains are among the most threatened

ecosystems (Thompson, 2000; Diaz et al., 2003, but see Körner & Hiltbrunner, 2021)

and are predicted to warm more rapidly in the northern hemisphere (Nogués-Bravo

et al., 2007; Keiler et al., 2010; Vitasse et al., 2021). The phenology of plant and

animal populations at high elevations is shifting on average towards earlier dates

(Vitasse et al., 2021). Long-term studies on amphibian populations living at high

elevations are scarce, and not much is known about how their breeding phenology is

changing. These populations experience different environmental conditions (e.g.,

increased amount of snow and colder temperatures) compared to their lowland

counterparts. Thus, different environmental variables potentially play a bigger role in

determining breeding phenology compared to what is observed at lower elevations

(Nufio et al., 2010; Bison et al., 2020).

Using 39 years of data on an explosive-breeding amphibian population living

at a high elevation (*B. bufo*), we study the relationship between breeding phenology

and both the environment and genetics (i.e., heritability of the breeding timing trait).

More specifically, our goal is to (i) identify the environmental variables (e.g.,

temperature, snow cover, moon cycle) that could be driving the observed breeding

phenology of this population (both the start and the peak of the breeding season), (ii)

analyse if there is significant variation in the phenological shifts among individuals,

(iii) obtain a measure of the importance of the genetic component, by calculating

individual-level repeatability (i.e., upper limit of heritability; Falconer, 1981; Lessells &

Boag, 1987; Semlitsch et al., 1993) of the timing of arrival at the breeding site for

both males and females.

Material and Methods

Life-history data

The study site is a pond located above Grindelwald, below the Grosse Scheidegg mountain pass (canton of Bern, Switzerland, 46.65240 N, 8.09683 E), at an elevation of 1841 m a.s.l. The pond measures approximately 10 m x 30 m, with a maximum depth of about 1 m. Since 1982, we have captured annually all the toads that come to breed at the study pond. We then marked (first by toe-clipping, then starting in 1993 by implanting PIT tags), measured, and released them in the same place (Hemelaar, 1988; Grossenbacher, 2002). To make sure we captured both early and late arrivers, we repeated this procedure for on average 5–6 nights, with breaks in-between of about 2-4 days (i.e, the data conform to Pollock's (1982) robust design). The length of the fieldwork period usually covers the breeding season duration, which typically lasts about two weeks at our study pond. This design also had the advantage of not overly stressing the toads. In total, for the period 1982–2020, 3053 uniquely recognizable individuals have been caught, of which 1852 were males and 1201 females. For each individual we have a record of presence for each capture night over the study period. Given the reduced size of the pond and the repeated capture rounds within a capture night, we assumed high capture probabilities (capture probability $p \approx 0.85$ per year based on a preliminary analysis of the mark-recapture data). At the population level we determined for each year a start, a peak, and an end date of breeding (i.e., first capture night, the capture night when most toads were captured, and last capture night, respectively). These calendar dates were all transformed into days of the year (where January 1st = 1), to facilitate

modelling of long-term trends. These dates come with a degree of uncertainty, given

the sampling done every 2–4 days and not daily. The date of start of the breeding

comes with additional uncertainty as the first capture night is not always reflective of

the same toad activity at the pond over the study period. We accounted for these

sources of uncertainty in all following analyses, using simulated data on start and

peak breeding dates.

Climatic data

We obtained climatic data for the period 1980–2020 from the DaymetCH dataset

(data obtained from *Bioclimatic maps of Switzerland © WSL, based on station data*

from the Federal Office of Meteorology and Climatology MeteoSwiss, and elaborated

by the *Land Change Science group, WSL*). This dataset consists of a 100-metre

resolution grid of interpolated estimates of weather variables, using meteorological

data from ground stations and the Daymet software (Thornton et al., 1997). We

obtained data for the cell containing the breeding pond for the following variables:

daily minimum, maximum, and mean temperature, daily total precipitation, and daily

snow water equivalent (SWE; the equivalent amount of water stored in the

snowpack). We then calculated average seasonal minimum, maximum, and mean

daily temperatures, and cumulative seasonal precipitation and SWE.

 Data analysis

 Population trend

A visual inspection of the data suggests that the trends in the breeding phenology

across the study period are non-linear, both for start and peak breeding (Figure 1).

Therefore, to better describe the observed trends, we conducted a piecewise

regression on both start and peak breeding using the R package *segmented*

(Muggeo, 2008). This analysis enables the identification of possible breakpoints in a

trend, in our case a year (or several years) when a significant change occurs in the

temporal trends of the breeding phenology. To account for possible imperfect

assignment of start and peak dates, we simulated 1000 datasets of breeding start

dates over the study period, allowing the date of the start of the breeding to be as

early as seven days before the initially assigned first capture night. We also

simulated 1000 datasets for peak breeding dates, allowing the dates to deviate from

the originally assigned date by letting it vary between the previous and the following

capture night. Using these simulated datasets, we ran 1000 piecewise regressions

for both start and peak breeding dates, and calculated 95% confidence intervals (CI)

for each model parameter, including p-values testing for the significance of the

breakpoint.

Moreover, to check how the standard deviation (SD) of the start or the peak

breeding dates changes over time, we calculated for both start and peak breeding

the SD of the residuals of each of the 1000 piecewise regressions, using a rolling

window approach (with a 10-year window) with the function *rollapply* of the package

zoo (Zeileis & Grothendieck, 2005).

Determinants of variation in the breeding phenology in the population

To understand the climatic causes of the observed shifts in the breeding phenology

of this population, we investigated the effects of several climatic variables on the

timing of breeding at the population level. We identified *a priori* the climatic

covariates that most reasonably could influence the breeding phenology in spring

based on previous literature and expert knowledge (Oseen & Wassersug, 2002;

Reading, 2003; While & Uller, 2014; Ficetola & Maiorano, 2016; Green, 2017).

These climatic covariates are: average minimum daily temperature in spring (T_{Sp})

and winter (T_W), total precipitation in spring ($Prec_{Sp}$, which includes both rainfall and

snowfall), total snow water equivalent in spring (SWE_{Sp}), and winter (SWE_W). We

then performed a piecewise regression on the time series of these five climatic

covariates (Figure 2, Table S1). We used minimum temperatures because toads are

nocturnal animals and are therefore more exposed to colder temperatures and less

to average or warmer temperatures. Moreover, minimum temperatures will

determine if the ground stays above freezing conditions. Changing the temperature

variable (mean vs minimum vs maximum) in the subsequent analyses did not

change the results as they were highly correlated ($r > 0.93$).

With warmer winters and springs, toads should emerge sooner from their

hibernation burrows as the snow will melt and the ground unfreeze earlier (Corn,

2003). The higher the snow water equivalent, the later the toads will emerge, as the

snow cover will keep them blocked underground (Corn, 2003). Finally, precipitation

can either favour or delay the breeding season. Snowfall should delay breeding as

the snow cover will increase (Corn, 2003), but rainfall could potentially lead to an

earlier start of the breeding season, as toads need high humidity levels to be active

(Todd et al., 2011; Green, 2017). After standardising these climatic variables by

subtracting the mean value and dividing by the standard deviation, we performed a

principal component analysis (PCA, function *prcomp*, R package *stats* (R Core

Team, 2020)), to reduce dimensionality and obtain uncorrelated variables (Figure

S1).

In addition to these five climatic variables, the lunar cycle has also been

identified to be an important factor for the timing of breeding in amphibians, with in

general peak migration to the breeding site under waxing or full moon phases (Grant

et al., 2009; Arnfield et al., 2012; Green et al., 2016; Jarvis et al., 2021). To assess

the effect of the lunar cycle on the breeding phenology in our population, we first

obtained the moon phase for each date of breeding start and peak breeding over the

study period using the package *lunar* (Lazaridis, 2014). Following Arnfield et al.

(2012) and Jarvis et al. (2021), we transformed the lunar phases in lunar angles (in

radians, where 0 = new moon and π = full moon).

To quantify both the effects of climate and of the moon cycle on the breeding

phenology, we modelled two separate linear regressions on the day of the breeding

start and the day of peak breeding over the period 1982–2020. As explanatory

variables we used the scores of the first two principal components (PC), as they

explained an important amount of the variance in the data (>70%), while also

checking for the effect of their interaction. As an additional explanatory variable, to

better understand the role of the moon cycle, we included the cosine of the lunar

angles of the start and peak breeding dates respectively. As we did for the piecewise

regression, we let the dates of start and peak breeding vary to account for their

uncertainty. For both start and peak breeding, we ran 1000 models with 1000

different datasets of dates and calculated 95% CI for all model parameters.

To further study the association between the moon cycle and breeding

phenology we tested if start and peak breeding tended to happen more frequently

under certain moon phases. To do this, we used the *rayleigh.test* function of the

circular R package (Agostinelli & Lund, 2017) to perform the Rayleigh test, a circular

goodness-of-fit test that is particularly suited for checking if the values of a circular

variable show a unimodal departure from a uniform distribution (Landler et al., 2018).

To check for significant multimodal departures we performed the Hermans-Rasson

test instead, using the *HR_test* from the *CircMLE* package (Fitak & Johnsen, 2017;

Landler et al., 2018)**.** Both tests were performed on the values in radians of the lunar

angles. Also for these analyses we simulated 1000 datasets of start and peak

breeding dates and obtained 95% CI for the p-values of the two tests.

Determinants of individual variation in breeding phenology

In addition to considering phenology at the population level, we also wanted to

understand whether individuals can show different patterns of changes in their

reproductive phenology over time through different responses to climatic variables,

possibly indicating a genetic component that mediates the effect of the changing

environment. We therefore modelled the effect of the previously used principal

components PC1 and PC2, as well as of the cosine of the lunar angle on each

individual first capture occasion in any given year (6735 occurrences for 3053

uniquely marked individuals, as many individuals were breeding in multiple years

(mean = 2.21 years, SE = 0.02)), using a linear mixed model (package *lmerTest*;

Kuznetsova et al., 2017). Also in this case, we accounted for uncertainty in the

assignment of the date of arrival to the pond by simulating 1000 new datasets where

every individual arrival date is newly sampled and can be as early as the capture

night preceding the original arrival date, or if it was the first capture night of the

season, up to seven days before. As a random effect, applied on both the intercept

and the slope of both PC1 and PC2, we included individual identity (ID). This was

done not only to observe if individuals react differently to changing environmental

conditions, but also to account for the non-independence of the data. Moreover, we

also included *year* as a random effect on the intercept, to account for unexplained

year-specific variation in the data. Finally, we included the effect of sex to account for

differences between males and female. To properly be able to compare the effects of

continuous variables (i.e., the two PCs and the cosine of the lunar angles) with the

effect of a categorical variable (i.e., sex), we standardised the three continuous

variables by subtracting the mean and dividing by two times the standard deviation

(Gelman, 2008). Finally, as a measure of model fit, we calculated the conditional R^2

value for each model using the *r.squaredGLMM* function from the package *MuMIn*

(Barton, 2019). We ran 1000 models and obtained mean and 95% CI values for each

parameter.

Repeatability of arrival date

Finally, we also estimated repeatability (i.e., the upper limit of heritability) of arrival

dates at the breeding site. High values of repeatability (*r*) mean that individuals are

consistent in their relative arrival timing (e.g., always among the first ones), and vice

versa. To calculate *r*, we used for each individual the date of first capture for each

year that it was captured. This date is a relatively good proxy for the date of arrival at

the breeding site, as the data collection usually starts every year approximately when

the first toads arrive at the pond. As for the individual-level analysis of the breeding

phenology, to account for the uncertainty in the assignment of the dates, we

repeated the calculation of r 1000 times, sampling different arrival dates every time.

The date was converted to the day of the year (where January 1st = 1), and then

standardised by subtracting the year-specific mean and dividing by the yyear-specific

standard deviation. We then used the function rpt from the package $rptR$ to calculate

r using individual ID as the group variable (Stoffel et al., 2017), and bootstrapping

1000 times to obtain the 95% CI. Given the different reproductive strategies that

males and females toads have, with females on average coming to the breeding site

later than males and for a shorter period of time (Reading & Clarke, 1983; Loman &

Madsen, 1986), we performed sex-specific calculations of r.

We conducted all the analyses in R (R version 4.1.1; R Core Team, 2020) with

RStudio (version 2022.7.1.554; R Studio Team, 2022).

Results

Population trend

Both the breeding start dates and the dates of peak breeding show very similar trends (Pearson's correlation coefficient = 0.91), with both also showing marked between-year variation over the study period. Nonetheless, a shift towards earlier breeding dates is observable, with breeding happening now on average around 30 days earlier compared to the start of the study period (Figure 1). In 910 cases out of 1000, the piecewise regression on breeding start dates identified a single breakpoint in the temporal trend in the year 1993, with a pre-1993 steep advancement of breeding dates followed by a post-1993 almost flat trend (Figure 1A; Table 1). The breakpoint years identified the most in the 1000 piecewise regressions on peak breeding dates were 1993 or 1996 (274 and 283 out of 1000 respectively) (Figure 1B; Table 1). Moreover, we found the standard deviation (SD) of the residuals of the piecewise regressions on both start and peak breeding dates to vary considerably, with higher SDs at the start and the end of the study period (Figure S2).

Figure 1. *Trends of breeding phenology over the study period (1982–2021). **(A)** Mean first*
day of the breeding season (day of the year, where January 1st = 1) out of the 1000 simulated.
The segmented green line is the result of a piecewise regression, where the year 1993 (± 5;
*95% CI) was identified as a breakpoint, thus creating two distinct trends. **(B)** Mean day of peak*
breeding (i.e., date where most toads were captured in a given breeding season) out of the
1000 simulated. The segmented green line is the result of a piecewise regression, where the
year 1996 (± 7, 95% CI) was identified as a breakpoint. The green band in both plots
represents the 95% CI for the piecewise regression.

Table 1. Summary of the piecewise regressions on the start and peak of the breeding season. For both intercept and slopes we show the estimate, its standard error, and the t-value and p-value associated with it. Moreover, in square brackets we show the 95% CI values obtained by running 1000 models. Slope1 refers to the segment before the breakpoint and Slope2 refers to the segment after the breakpoint. Asterisks next to the p-values show significance at the 0.05 level. The p-value for Slope2 is NA since standard asymptotics do not apply (Muggeo, 2008).

		Estimates	Std. Error	t-value	p-value
Start	Intercept	6342.69 [6317.19 ; 6368.20]	1874.42 [1864.24 ; 1884.59]	3.40 [3.38 ; 3.42]	0.0021* [0.0020 ; 0.0022]
	Slope1	-3.116 [-3.130 ; -3.103]	0.943 [0.938 ; 0.948]	-3.319 [-3.334 ; -3.303]	0.0027* [0.0026 ; 0.0028]
	Slope2	0.063 [0.059 ; 0.066]	0.229 [0.229 ; 0.230]	0.272 [0.258 ; 0.285]	NA
	Breakpoint	1993	2.294 [2.282 ; 2.305]	-	< 0.001*
Peak	Intercept	4491.64 [4412.97 ; 4570.31]	1379.40 [1342.63 ; 1416.16]	3.55 [3.50 ; 3.59]	0.0037* [0.0034 ; 0.0040]
	Slope1	-2.18 [-2.22 ; -2.14]	0.69 [0.68 ; 0.71]	-3.40 [-3.44 ; -3.36]	0.0046* [0.042 ; 0.049]
	Slope2	0.32 [0.29 ; 0.35]	0.44 [0.42 ; 0.45]	0.41 [0.37 ; 0.44]	NA
	Breakpoint	1993 / 1996	3.13 [3.11 ; 3.15]	-	0.0021* [0.0020 ; 0.0021]

355
356
Figure 2. *Trends over the study period of the five focal environmental variables. **(A)** Average*
minimum daily temperature in spring. The piecewise regression identified the year 2007 (± 9,
*95% CI) as a breakpoint. **(B)** Average minimum daily temperature in winter. The year 1990 (±*
*7, 95% CI) was a breakpoint. **(C)** Cumulative precipitation in spring. The year 1986 (± 8, 95%*
*CI) was a breakpoint. **(D)** Cumulative snow water equivalent (SWE) in winter. The year 1985*
*(± 5, 95% CI) was a breakpoint. **(E)** Cumulative SWE in Spring. The year 1990 (± 7, 95% CI)*
was a breakpoint. In all plots green ribbons represent the 95% CI for the linear regressions.
Table S1 in the Appendix shows the summary of these five piecewise regressions.

Determinants of variation in the breeding phenology in the population

The first two principal components (PC) of the principal component analysis (PCA) described together more than 70% of the variation in the data, and both had a standard deviation above one (Figure S1; Table S2). Therefore, we kept the scores of these two PCs (PC1 and PC2) as explanatory variables in the following linear regressions on the start of the breeding season and on peak breeding (also including the scaled cosine of lunar angle). PC1 was mostly determined by winter temperature (+0.45 loading) and winter and spring SWE (-0.61 and -0.64, respectively). PC2 was mostly determined by spring weather conditions. Spring temperature had a negative loading (-0.68), while precipitation had a positive loading (+0.68) (Figure S1; Table S3).

The 1000 linear regressions on the start of the breeding season and the 1000 on the date of peak breeding performed similarly. Regarding the start of the breeding season, the model (mean adjusted R^2 = 0.384 [0.382 ; 0.385]) indicated a significant negative relationship with PC1 and a significant positive relationship with PC2 (Table 2). The interaction of PC1 and PC2 had a non-significant and small effect, and the cosine of the lunar angle also had a non-significant effect. Similarly, for the regression on peak night, we found a significant negative relationship with PC1 and a significant positive relationship with PC2, while the interaction of PC1 and PC2 and the cosine of the lunar angle were non-significant (Table 2). The mean adjusted R^2 was 0.561 [0.558 ; 0.564]. In both cases the outcome is that warmer temperatures in winter and spring, less snow cover, and weaker precipitations are all associated with an earlier start and peak of the breeding season.

Table 2. Summary of the linear regression on the start and peak of the breeding season. For each variable we show the estimate, its standard error, and the t-value and p-value associated with it. In square brackets we show the 95% CI values, obtained by bootstrapping 1000 times. Asterisks next to the p-value show significance at the 0.05 level.

		Estimates	Std. Error	t-value	p-value
Start	Intercept	138.22 [138.18 ; 138.24]	1.713 [1.709 ; 1.716]	80.82 [80.64 ; 81.00]	< 0.0001*
	PC1	-5.53 [-5.56 ; -5.51]	1.754 [1.750 ; 1.759]	-3.16 [-3.17 ; -3.14]	0.0042* [0.0040 ; 0.0044]
	PC2	6.96 [6.93 ; 6.99]	1.838 [1.833 ; 1.843]	3.80 [3.78 ; 3.82]	0.00087* [0.00079 ; 0.00094]
	cos(moon)	0.98 [0.91 ; 1.04]	1.814 [1.808 ; 1.819]	0.54 [0.50 ; 0.58]	0.56 [0.54 ; 0.57]
	PC1:PC2	-0.058 [-0.095 ; -0.020]	2.281 [2.275 ; 2.286]	-0.026 [-0.043 ; -0.010]	0.84 [0.83 ; 0.85]
Peak	Intercept	150.36 [150.34 ; 150.39]	1.571 [1.566 ; 1.574]	95.94 [95.67 ; 96.20]	< 0.0001*
	PC1	-5.43 [-5.39 ; -5.46]	1.604 [1.600 ; 1.609]	-3.39 [-3.42 ; -3.37]	0.0030* [0.0028 ; 0.0032]
	PC2	8.91 [8.88 ; 8.95]	1.675 [1.670 ; 1.680]	5.34 [5.31 ; 5.36]	< 0.0001*
	cos(moon)	-2.50 [-2.56 ; -2.44]	1.646 [1.641 ; 1.651]	-1.53 [-1.57 ; -1.49]	0.20 [0.19 ; 0.21]
	PC1:PC2	-2.39 [-2.44 ; -2.35]	2.102 [2.096 ; 2.107]	-1.14 [-1.16 ; -1.12]	0.29 [0.28 ; 0.30]

Effect of the moon cycle on breeding phenology

To further understand if the lunar cycle is associated with the breeding phenology,
we performed two statistical tests. To check for unimodal deviation we ran a
Rayleigh's test on 1000 datasets of moon phases on breeding season start and on
peak dates. In both cases we obtained a non-significant p-value (0.37 [0.36 ; 0.39]
and 0.15 [0.14 ; 0.16] respectively), indicating that we could not confidently reject the
null-hypothesis of the data being uniformly distributed in the circular space. Also the

398 outcome of the Hermans-Rasson test for multivariate deviations indicated that the

399 null hypothesis could not be rejected for both start and peak breeding (p-value = 0.45

400 [0.44 ; 0.47] and 0.26 [0.25 ; 0.27] respectively). This means that there was no clear

401 pattern between lunar phases and the start of the breeding season or the peak

402 breeding (Figure 3).

403
404 *Figure 3.* Circular histogram showing counts of *(A)* the original breeding start dates and *(B)*
405 *the original peak breeding dates under different lunar phases for the period 1982–2021 (e.g.,*
406 *the breeding season started eight times under a moon phase with a lunar angle between 0°*
407 *and 60°). Lunar angles, initially in radians, were back-transformed to degrees, so that the new*
408 *moon is at 0° and full moon is at 180°.*

Determinants of individual variation in breeding phenology

To better understand if there are among-individual differences in the phenological response to changing climatic variables, we used a linear mixed model to test for the effect of climatic variables on the individual breeding start dates (i.e., the date on which an individual was first captured). We ran 1000 models, always changing the arrival date of each individual by letting it vary between the assigned date and up to seven days before. We then obtained mean and 95% CI values for each model parameter. We found only a small difference in the response of breeding phenology to climatic variables among individuals (i.e., low values for the random effect ID, both on intercept and slopes, Table 3). We found a strong significant positive effect of PC2 on the breeding dates (16.32 ± 3.38 SE), meaning that stronger precipitation and lower minimum spring temperatures are associated with a delay in the breeding. We also found a significant and strong negative effect of PC1 (-9.58 ± 2.97 SE), indicating that colder winter temperatures and higher SWE are associated with a delay in the breeding. The interaction between PC1 and PC2 was not significant. We also found a significant but weak effect of the cosine of the lunar angle (1.07 ± 0.15 SE), suggesting a possible small role of the lunar cycle. Finally, we observed an effect of sex indicating that males arrived on average earlier than females (-1.76 ± 0.16 SE) (Table 3).

Table 3. *Detailed description of the model used to check for the effect of environmental variables on the phenology at the individual level. Sex is included to observe differences between males and females. The response variable ArrivalDate is a vector of dates of arrival at the breeding site for each individual over the study period. PC1 and PC2 are the first two components of the PCA performed on the climatic data. Cos(moon) is the cosine of the lunar angle for the arrival date. ID refers to the identity of each individual, and it is used as a random effect on both intercept and the slopes of PC1 and PC2. Finally, Year is included as a random effect to account for additional unexplained variation that might be caused by sampling variation. The second and third part of the table provide details on the estimates for the fixed and random effects respectively. In square brackets we show the 95% CI obtained by bootstrapping 1000 times the model.*

Model name	Variables			Conditional R^2
Full_model	ArrivalDate ~ PC1 + PC2 + PC1:PC2 + Sex + cos(moon) + (1|Year) + (1|ID) + (0 + PC1|ID) + (0 + PC2|ID)			0.8822 [0.8821 ; 0.8823]

	Effect size	Std. Error	t-value	P-value
Intercept	145.902 [145.899 ; 145.905]	1.6910 [1.6907 ; 1.6913]	86.28 [86.26 ; 86.30]	< 0.0001*
Sex (male)	-1.757 [-1.760 ; -1.754]	0.1640 [0.1639 ; 0.1641]	-10.71 [-10.73 ; -10.70]	< 0.0001*
PC1	-9.581 [-9.585 ; -9.577]	2.974 [2.973 ; 2.975]	-3.221 [-3.223 ; -3.220]	0.00276* [0.00275 ; 0.00277]
PC2	16.315 [16.309 ; 16.320]	3.3844 [3.3837 ; 3.3851]	4.820 [4.819 ; 4.822]	< 0.0001*
cos(moon)	1.07 [1.06 ; 1.08]	0.1465 [0.1464 ; 0.1466]	7.31 [7.26 ; 7.36]	< 0.0001*
PC1:PC2	-5.63 [-5.64 ; -5.62]	7.542 [7.540 ; 7.543]	-0.747 [-0.748 ; -0.745]	0.460 [0.459 ; 0.461]

	Variance
ID (intercept)	3.07 [3.06 ; 3.08]
ID on PC1	1.91 [1.89 ; 1.94]
ID on PC2	0.34 [0.31 ; 0.36]
Year (intercept)	105.65 [105.61 ; 105.70]
Residuals	26.07 [26.05 ; 26.09]

429   ***Repeatability of arrival date***

431   In total, 453 females and 1092 males visited the pond over multiple years. We

432   calculated 1000 repeatability values for females and 1000 for males, sampling every

433   time new arrival dates. We found a mean repeatability value *r* of 0.141 [CI 95%

434   0.140; 0.142] for females and 0.1017 [0.1014; 0.1021] for males.

Discussion

Our results show that variation in the breeding phenology is strongly associated with climatic conditions, which vary substantially among years but also show trends across times. We also found low repeatability values and low variability in individual responses, suggesting the absence of a strong genetic component contributing to the observed variation of individuals in the breeding phenology. Finally, we found indications of a possibly significant, but weak, effect of the lunar cycle. A signal might indeed exist, but the climatic variables probably have a stronger effect.

The hypothesised strong link between the breeding phenology of high-elevation amphibian populations and climatic conditions appears to be confirmed. Increasing temperatures are a key driver of snow melt and ground defrosting, which in turn act as important environmental cues for toads to initiate migration to their breeding grounds (Corn & Muths, 2002; Green, 2017). During particularly warm springs, the snow melts and the ground defrosts earlier, leading to a shift of the onset of breeding to earlier dates. Our findings on the importance of temperature are in line with previous studies on *B. bufo* (Reading & Clarke, 1983; Reading, 2003; Tryjanowski et al., 2003; Arnfield et al., 2012). On the other hand, where past studies have identified rainfall to be an important trigger for migration in lowland populations (Reading & Clarke, 1983; Sinsch, 1988; Jarvis et al., 2021), we did not clearly observe this in our data, as our measure of precipitation included both snow- and rainfall. We found that a higher amount of precipitation in spring (combined with a decrease of spring temperature) was associated with a later breeding date. In fact, at low temperatures, snowfall or freezing rain can delay the melting of the snow cover, therefore leading to

a delay in the breeding. Finally, the negative association between snow water

equivalent (SWE) and breeding timing is in line with the rest of the findings. In fact,

SWE depends considerably on temperatures and precipitation, as well as other

aspects such as exposition, and it is a key factor that influences phenology (Corn,

2003). The very similar trend observed for peak activity in breeding indicates that

both start and peak breeding are influenced mostly in the same way by the same

climatic variables.

When looking at the individual timing of arrival we still found an important effect on

the breeding phenology of PC1 (T_W and $SWE_{Sp/W}$) and PC2 (T_{Sp} and $Prec_{Sp}$) (Table

3). However, we found only non-significant and small among-individual variation in

phenological response to changing climatic conditions (Table 3). As reproduction

happens only once a year in temperate-living explosive breeders, synchronisation in

breeding could be key to maximise reproductive output (Ims, 1990). Such an

accurate synchronisation can be achieved more easily when all individuals

hibernating close to each other have similar responses to external cues triggering

their migration to the breeding pond, instead of responding individually in different

ways, highlighting once more that the breeding phenology is mainly driven by

climatic conditions.

Moreover, the low values of r (i.e., the upper limit of heritability) that we found

for the timing of arrival further indicate that there is only a small contribution of the

genetic component to variation in the breeding phenology. This conclusion is in line

with what most studies on amphibian phenology found (Semlitsch et al., 1993;

Blaustein et al., 2001; Parmesan, 2007; While & Uller, 2014; but see Heusser & Ott,

1968; Phillimore et al., 2010). In other species, for instance birds, higher values of

repeatability have been found for migration phenology, a trait linked to breeding.

Franklin and colleagues (2022) found in their meta-analysis an average value of

repeatability of 0.414, while Kürten and colleagues (2022) found repeatability values

above 0.60 for various traits (but see Clermont et al., 2018; Vaillant et al., 2021 for

examples of low repeatability in birds), but in amphibians that follow an explosive

breeding strategy, the genetic component does not appear to be the main

determinant of variation in breeding phenology. This is due to either populations

being truly able to respond plastically to changing climatic conditions, and therefore

there is no strong selection on genetic variation in the trait, or there is little genetic

variation in the population to begin with. Low values of repeatability might also

indicate a non-consistent choice of the hibernation site (and therefore distance to the

pond). Not much is known about hibernation site fidelity in anurans, and future

studies should address this question.

Finally, we found that on average males tend to arrive earlier than females (Table 3),

similarly to what has been found in lowland populations of *B. bufo* (Loman &

Madsen, 1986; Höglund & Robertson, 1987, 1988; but see Gittins et al., 1980). In

these studies, males, especially bigger ones, were observed to arrive on average

earlier to the breeding pond. Smaller males, on the other hand, were observed

intercepting females on their way to the pond, betting on the fact that the females

would lay the eggs as soon as they arrived at the pond, avoiding competition from

the other bigger males. A more detailed future analysis of body size and its effects

on the timing of migration to the breeding site could confirm this theory also for this

population.

Climate change is leading to on-average increasing temperatures both globally but

also at smaller scales such as the European Alps (Vitasse et al., 2021) and

Switzerland (Rebetez & Reinhard, 2008). The start of the study period (early 1980s)

coincides with an important increase of temperatures in Switzerland (Bundesamt für

Umwelt (BAFU), 2020). In fact, each year since the mid-80s, the deviation from the

mean yearly temperature (average calculated over the period 1864–2019) has

always been positive (Begert & Frei, 2018). In the Swiss Alps, mean temperature

increased by about 1.7 °C from 1975 to 2004, nearly twice the global average

(Rebetez & Reinhard, 2008). Despite these general trends, we observe at our study

site stable or even decreasing trends in temperatures during the study period,

especially in the second part of the study period (Figure 2). Initially, the shift towards

earlier breeding (pre-1993/1996) can be explained by warming temperatures and

decreasing SWE (Figure 2). On the other hand, the absence of a trend in the

breeding dates observed after the mid-1990s (Figure 1) could be explained by a

change in trajectories of winter temperature, which started decreasing around 1990

(Figure 2), as well as of winter and spring SWE, which started increasing around the

same time. These combined changes are acting against the increasing spring

temperature (which has increased until around 2007; Figure 2), therefore slowing

down and ultimately halting the shift towards earlier breeding dates of the toads.

While we could expect climate change to act linearly on the shift towards

earlier breeding dates, it is possible that other site-specific conditions prevail at

smaller temporal and geographical scales, creating an heterogenous mosaic of

climate conditions, especially at high elevations (Scherrer & Körner, 2011; Feldmeier

et al., 2020). Populations therefore will not necessarily respond the same way, as

they are experiencing different microclimates (Miller et al., 2018; Arietta et al., 2020;

Turner & Maclean, 2022). Our data on climatic variables was limited to the 100 m x

100 m cell which includes the pond, and since we do not exactly know where the

toads hibernate in the surrounding landscape, we cannot exclude that they are

experiencing different microclimates compared to the pond. Toads have been found

hibernating more than one kilometre away from the breeding site horizontally, and up

to almost 400 metres away vertically (Sztatecsny & Schabetsberger, 2005). Since

the breeding pond and surrounding area are often still partially covered by snow

during peak night, the hibernation grounds are probably warmer than the pond itself.

This difference in microclimates between hibernation sites and breeding site could

further explain individual variation in breeding timing (e.g., arrival at the pond).

Further studies on the hibernation grounds of the toads in this population and how

the choice of the hibernation site affects breeding phenology should be conducted.

Despite the observed stabilisation of the trend of the breeding dates (Figure 1), the

study population is experiencing increased variation in the dates of the start of the

breeding season (Figure S2). This increased variation could be explained by

extreme weather events whose occurrence is expected to increase under climate

change (Rahmstorf & Coumou, 2011; National Academies of Sciences, Engineering,

and Medicine, 2016). These unpredictability and extremeness of environmental

conditions can threaten the population if they lead to either excessively early or late

breeding. In fact, extreme early breeding is associated with reduced hibernation

periods which can decrease the body condition in spring (Reading, 2007).

Additionally, early breeding can expose eggs and tadpoles to late frost events (Muir

et al., 2014; Bison et al., 2021; Turner & Maclean, 2022). Delayed breeding can

potentially pose a problem as well if the pond dries out during late spring warm

conditions or if juveniles cannot accomplish full growth before hibernation. Indeed,

smaller and younger juveniles are more at risk of death before and during the first

hibernation period (Morin et al., 1990; Sinsch & Schäfer, 2016). This seems to be

compensated in some cases by an accelerated growth at the larval stage in case of

late breeding, but with a cost of reduced defences against predation (Orizaola et al.,

2016). Such riskful situations can have strong negative effects on individual survival

and reproductive output, ultimately leading to population declines (Reading, 2007;

Iler et al., 2021). On the other hand, at least initially, climate change could lead to

longer growing seasons during which individuals would have the opportunity to

gather more energy before the onset of hibernation (Zani, 2008; Iler et al., 2021),

with potentially positive effects at the population level. Climate change can as well

lead to species expanding upward (Vitasse et al., 2021), with *Bufo bufo* populations

observed locally extending their upper range limit to higher elevations (Lüscher et al.,

2016). If moving upwards is not possible, high-elevation populations adapted to their

environments could face local extirpation (Urban, 2018).

Conclusion

In this study we showed the important association between climatic variables such as temperature, snow cover, and precipitation with the breeding phenology of a *Bufo bufo* population living at high elevations. Breeding happens on average around 30 days earlier now compared to four decades ago, and interestingly the shift towards earlier breeding dates has not been constant, but is better described by two different trends. After an initial steep advancement until the mid-90s, the trend stabilised. This is reflected in the trends of the time series of the focal climatic variables, which explain the observed temporal variation in breeding phenology. The stabilisation in the trend suggests that there might be spatial heterogeneity in climate change and its effects, therefore different populations might show different trends in their breeding phenology. This stabilisation is accompanied by an increased variation in the dates of the start of the breeding season, with potential consequences for the population that should be further investigated in the future. To conclude, this 40-year study is one of the first and most detailed studies on the breeding phenology of alpine populations of *B. bufo*, and it highlights the influence of changing environmental conditions on the timing of reproduction.

Acknowledgements

We warmly thank all the people involved in data collection and management since 1982 and everybody who facilitated fieldwork during the past 40 years. Capture, handling and marking of toads were done under an animal welfare permit issued by the Veterinäramt des Kantons Bern. We also thank the Land Change Science of the Swiss Federal Research Institute WSL for kindly providing us with the climatic data.

Authors contribution

O.L. and B.S. conceived the study. K.G., S.Z., S.A., B.L., D.S., M.T., and H.R. collected data. O.L. prepared and analysed the data. B.S and A.O. provided feedback on the analyses. O.L. wrote the paper with input from all authors.

Data and script accessibility

Data and scripts for this publication will be online on the Zenodo Repository.

Supplementary material

Extra tables and figures are available in the Appendix.

Conflict of interest and disclosure

The authors of this preprint declare that they have no financial conflict of interest with the content of this article.

Funding

This project was funded by the Federal Office for the Environment (contract numbers: 20.0001.PJ/46DBED0F1 and 06.0126-PZ Artenförderung / N494-1379) and the Stiftung Temperatio.

References

618 **References**

619 Agostinelli, C., & Lund, U. (2017). *R package 'circular': Circular statistics*. https://r-

620   forge.r-project.org/projects/circular/

621 Arietta, A. Z. A., Freidenburg, L. K., Urban, M. C., Rodrigues, S. B., Rubinstein, A., &

622   Skelly, D. K. (2020). Phenological delay despite warming in wood frog *Rana*

623   *sylvatica* reproductive timing: A 20-year study. *Ecography*, *43*(12), 1791–

624   1800. https://doi.org/10.1111/ecog.05297

625 Arnfield, H., Grant, R. A., Monk, C., & Uller, T. (2012). Factors influencing the timing

626   of spring migration in common toads (*Bufo bufo*): Timing of spring migration in

627   toads. *Journal of Zoology*, *288*(2), 112–118. https://doi.org/10.1111/j.1469-

628   7998.2012.00933.x

629 Barton, K. (2019). *Multi-Model Inference*.

630 Beebee, T. J. C. (1995). Amphibian breeding and climate. *Nature*, *374*(6519), 219–

631   220. https://doi.org/10.1038/374219a0

632 Begert, M., & Frei, C. (2018). Long-term area-mean temperature series for

633   Switzerland—Combining homogenized station data and high resolution grid

634   data. *International Journal of Climatology*, *38*(6), 2792–2807.

635   https://doi.org/10.1002/joc.5460

636 Bison, M., Yoccoz, N. G., Carlson, B. Z., Klein, G., Laigle, I., Van Reeth, C., Asse,

637   D., & Delestrade, A. (2020). Best environmental predictors of breeding

638   phenology differ with elevation in a common woodland bird species. *Ecology*

639   *and Evolution*, *10*(18), 10219–10229. https://doi.org/10.1002/ece3.6684

640 Bison, M., Yoccoz, N. G., Carlson, B. Z., Klein, G., Laigle, I., Van Reeth, C., &

641   Delestrade, A. (2021). Earlier snowmelt advances breeding phenology of the

642   common frog (*Rana temporaria*) but increases the risk of frost exposure and

643   wetland drying. *Frontiers in Ecology and Evolution*, *9*, 645585.

644   https://doi.org/10.3389/fevo.2021.645585

645 Blaustein, A. R., Belden, L. K., Olson, D. H., Green, D. M., Root, T. L., & Kiesecker,

646   J. M. (2001). Amphibian breeding and climate change. *Conservation Biology*,

647   *15*(6), 1804–1809. https://doi.org/10.1046/j.1523-1739.2001.00307.x

648 Bundesamt für Umwelt (BAFU). (2020). *Klimawandel in der Schweiz* (p. 105).

649 Clermont, J., Réale, D., & Giroux, J.-F. (2018). Plasticity in laying dates of Canada

650   geese in response to spring phenology. *Ibis*, *160*(3), 597–607.

651   https://doi.org/10.1111/ibi.12560

652 Corn, P. S. (2003). Amphibian breeding and climate change: Importance of snow in

653   the mountains. *Conservation Biology*, *17*(2), 622–625.

654   https://doi.org/10.1046/j.1523-1739.2003.02111.x

655 Corn, P. S., & Muths, E. (2002). Variable breeding phenology affects the exposure of

656   amphibian embryos to ultraviolet radiation. *Ecology*, *83*(11), 2958–2963.

657   https://doi.org/10.1890/0012-9658(2002)083[2958:VBPATE]2.0.CO;2

658 Diaz, H. F., Grosjean, M., & Graumlich, L. (2003). Climate variability and change in

659   high elevation regions: Past, present and future. *Climatic Change*, *59*, 1–4.

660   https://doi.org/10.1007/978-94-015-1252-7_1

661 Duellman, W. E., & Trueb, L. (1986). *Biology of Amphibians*. McGraw-Hill Book

662   Company.

663 Falconer, D. S. (1981). *Introduction to quantitative genetics* (2d ed.). Longman.

664 Feldmeier, S., Schmidt, B. R., Zimmermann, N. E., Veith, M., Ficetola, G. F., &

665   Lötters, S. (2020). Shifting aspect or elevation? The climate change response

666   of ectotherms in a complex mountain topography. *Diversity and Distributions*,

667   *26*(11), 1483–1495. https://doi.org/10.1111/ddi.13146

Ficetola, G. F., & Maiorano, L. (2016). Contrasting effects of temperature and

precipitation change on amphibian phenology, abundance and performance.

Oecologia, *181*(3), 683–693. https://doi.org/10.1007/s00442-016-3610-9

Fitak, R. R., & Johnsen, S. (2017). Bringing the analysis of animal orientation data

full circle: Model-based approaches with maximum likelihood. *Journal of*

Experimental Biology, *220*(21), 3878–3882.

https://doi.org/10.1242/jeb.167056

Franklin, K. A., Nicoll, M. A. C., Butler, S. J., Norris, K., Ratcliffe, N., Nakagawa, S.,

& Gill, J. A. (2022). Individual repeatability of avian migration phenology: A

systematic review and meta-analysis. *Journal of Animal Ecology*, *91*(7),

1416–1430. https://doi.org/10.1111/1365-2656.13697

Garner, T. W. J., Rowcliffe, J. M., & Fisher, M. C. (2011). Climate change,

chytridiomycosis or condition: An experimental test of amphibian survival.

Global Change Biology, *17*(2), 667–675. https://doi.org/10.1111/j.1365-

2486.2010.02272.x

Gelman, A. (2008). Scaling regression inputs by dividing by two standard deviations.

Statistics in Medicine, *27*(15), 2865–2873. https://doi.org/10.1002/sim.3107

Gittins, S. P., Parker, A. G., & Slater, F. M. (1980). Population characteristics of the

common toad (*Bufo bufo*) visiting a breeding site in Mid-Wales. *Journal of*

Animal Ecology, *49*(1), 161–173. https://doi.org/10.2307/4281

Gotthard, K. (2001). Growth strategies of ectothermic animals in temperate

environments. In D. Atkinson & M. Thorndyke (Eds.), *Environment and animal*

development: Genes, life histories and plasticity (pp. 287–304). BIOS

Scientific.

Grant, R. A., Chadwick, E. A., & Halliday, T. R. (2009). The lunar cycle: A cue for

amphibian reproductive phenology? *Animal Behaviour*, *78*(2), 349–357.

https://doi.org/10.1016/j.anbehav.2009.05.007

Green, D. M. (2017). Amphibian breeding phenology trends under climate change:
Predicting the past to forecast the future. *Global Change Biology*, *23*(2), 646–656. https://doi.org/10.1111/gcb.13390

Green, T., Das, E., & Green, D. M. (2016). Springtime emergence of overwintering toads, *Anaxyrus fowleri* , in relation to environmental factors. *Copeia*, *104*(2), 393–401. https://doi.org/10.1643/CE-15-323

Grossenbacher, K. (2002). First results of a 20-year-study on common toad *Bufo bufo* in the Swiss Alps. *Biota*, *3*(1–2), 43–48.

Hartel, T., Sas-Kovacs, I., Pernetta, A., & Geltsch, I. (2007). The reproductive dynamics of temperate amphibians: A review. *North-Western Journal of Zoology*, *3*.

Hemelaar, A. (1988). Age, growth and other population characteristics of *Bufo bufo* from different latitudes and altitudes. *Journal of Herpetology*, *22*(4), 369.

https://doi.org/10.2307/1564332

Heusser, H., & Ott, J. (1968). Sollzeit der Laichwanderung bei der Erdkröte, *Bufo bufo* (L.). *Revue Suisse de Zoologie*, *75*, 1005–1022.

Höglund, J., & Robertson, J. G. M. (1987). Random mating by size in a population of common toads (*Bufo bufo*). *Amphibia-Reptilia*, *8*(4), 321–330.

https://doi.org/10.1163/156853887X00108

Höglund, J., & Robertson, J. G. M. (1988). Chorusing behaviour, a density-dependent alternative mating strategy in male common toads (*Bufo bufo*).

Ethology, *79*(4), 324–332. https://doi.org/10.1111/j.1439-0310.1988.tb00721.x

Iler, A. M., CaraDonna, P. J., Forrest, J. R. K., & Post, E. (2021). Demographic

consequences of phenological shifts in response to climate change. *Annual*

Review of Ecology, Evolution, and Systematics, *52*(1), annurev-ecolsys-

011921-032939. https://doi.org/10.1146/annurev-ecolsys-011921-032939

Ims, R. A. (1990). The ecology and evolution of reproductive synchrony. *Trends in*

Ecology & Evolution, *5*(5), 135–140. https://doi.org/10.1016/0169-

5347(90)90218-3

Jara, F. G., Thurman, L. L., Montiglio, P.-O., Sih, A., & Garcia, T. S. (2019).

Warming-induced shifts in amphibian phenology and behavior lead to altered

predator–prey dynamics. *Oecologia*, *189*(3), 803–813.

https://doi.org/10.1007/s00442-019-04360-w

Jarvis, L. E., Grant, R. A., & SenGupta, A. (2021). Lunar phase as a cue for

migrations to two species of explosive breeding amphibians—Implications for

conservation. *European Journal of Wildlife Research*, *67*(1), 11.

https://doi.org/10.1007/s10344-020-01453-3

Keiler, M., Knight, J., & Harrison, S. (2010). Climate change and geomorphological

hazards in the eastern European Alps. *Philosophical Transactions of the*

Royal Society A: Mathematical, Physical and Engineering Sciences,

368(1919), 2461–2479. https://doi.org/10.1098/rsta.2010.0047

Kokko, H. (1999). Competition for early arrival in migratory birds. *Journal of Animal*

Ecology, *68*(5), 940–950. https://doi.org/10.1046/j.1365-2656.1999.00343.x

Körner, C., & Hiltbrunner, E. (2021). Why is the alpine flora comparatively robust

against climatic warming? *Diversity*, *13*(8), 383.

https://doi.org/10.3390/d13080383

Kovar, R., Brabec, M., Bocek, R., & Vita, R. (2009). Spring migration distances of

some Central European amphibian species. *Amphibia-Reptilia*, *30*(3), 367–

378. https://doi.org/10.1163/156853809788795236

Kürten, N., Schmaljohann, H., Bichet, C., Haest, B., Vedder, O., González-Solís, J.,
& Bouwhuis, S. (2022). High individual repeatability of the migratory behaviour
of a long-distance migratory seabird. *Movement Ecology*, *10*(1), 5.
https://doi.org/10.1186/s40462-022-00303-y

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package:
Tests in linear mixed effects models. *Journal of Statistical Software*, *82*(13),
1–26.

Landler, L., Ruxton, G. D., & Malkemper, E. P. (2018). Circular data in biology:
Advice for effectively implementing statistical procedures. *Behavioral Ecology
and Sociobiology*, *72*(8), 128. https://doi.org/10.1007/s00265-018-2538-y

Lazaridis, E. (2014). *lunar: Lunar phase and distance, seasons and other
environmental factors* (Version 0.1-04).

Lessells, C. M., & Boag, P. T. (1987). Unrepeatable repeatabilities: A common
mistake. *The Auk*, *104*(1), 116–121. https://doi.org/10.2307/4087240

Loman, J., & Madsen, T. (1986). Reproductive tactics of large and small male toads
Bufo bufo. *Oikos*, *46*(1), 57. https://doi.org/10.2307/3565380

Lüscher, B., Beer, S., & Grossenbacher, K. (2016). Die Höhenverbreitung der
Erdkröte (*Bufo bufo*) im Berner Oberland (Schweiz) unter sich verändernden
Klimabedingungen. *Zeitschrift für Feldherpetologie*, *23*, 47–58.

Miller, D. A. W., Grant, E. H. C., Muths, E., Amburgey, S. M., Adams, M. J., Joseph,
M. B., Waddle, J. H., Johnson, P. T. J., Ryan, M. E., Schmidt, B. R., Calhoun,
D. L., Davis, C. L., Fisher, R. N., Green, D. M., Hossack, B. R., Rittenhouse,
T. A. G., Walls, S. C., Bailey, L. L., Cruickshank, S. S., … Sigafus, B. H.
(2018). Quantifying climate sensitivity and climate-driven change in North

American amphibian communities. *Nature Communications*, *9*(1), 3926. https://doi.org/10.1038/s41467-018-06157-6

Morin, P. J., Lawler, S. P., & Johnson, E. A. (1990). Ecology and breeding phenology of larval *Hyla andersonii*: The disadvantages of breeding late. *Ecology*, *71*(4), 1590–1598. https://doi.org/10.2307/1938294

Muggeo, V. M. R. (2008). Segmented: An R package to fit regression models with broken-line. *R News*, *8*(1), 20–25.

Muir, A. P., Biek, R., Thomas, R., & Mable, B. K. (2014). Local adaptation with high gene flow: Temperature parameters drive adaptation to altitude in the common frog (*Rana temporaria*). *Molecular Ecology*, *23*(3), 561–574. https://doi.org/10.1111/mec.12624

National Academies of Sciences, Engineering, and Medicine. (2016). *Attribution of extreme weather events in the context of climate change*. National Academies Press.

Nogués-Bravo, D., Araújo, M. B., Errea, M. P., & Martínez-Rica, J. P. (2007). Exposure of global mountain systems to climate warming during the 21st Century. *Global Environmental Change*, *17*(3–4), 420–428. https://doi.org/10.1016/j.gloenvcha.2006.11.007

Nufio, C. R., McGuire, C. R., Bowers, M. D., & Guralnick, R. P. (2010). Grasshopper community response to climatic change: Variation along an elevational gradient. *PLoS ONE*, *5*(9), 1–11. https://doi.org/10.1371/journal.pone.0012977

Orizaola, G., Richter-Boix, A., & Laurila, A. (2016). Transgenerational effects and impact of compensatory responses to changes in breeding phenology on antipredator defenses. *Ecology*, *97*(9), 2470–2478.

793 https://doi.org/10.1002/ecy.1464

794 Oseen, K. L., & Wassersug, R. J. (2002). Environmental factors influencing calling in

795 sympatric anurans. *Oecologia*, *133*(4), 616–625.

796 https://doi.org/10.1007/s00442-002-1067-5

797 Parmesan, C. (2007). Influences of species, latitudes and methodologies on

798 estimates of phenological response to global warming. *Global Change*

799 *Biology*, *13*(9), 1860–1872. https://doi.org/10.1111/j.1365-2486.2007.01404.x

800 Parmesan, C., & Yohe, G. (2003). A globally coherent fingerprint of climate change

801 impacts across natural systems. *Nature*, *421*(6918), 37–42.

802 https://doi.org/10.1038/nature01286

803 Phillimore, A. B., Hadfield, J. D., Jones, O. R., & Smithers, R. J. (2010). Differences

804 in spawning date between populations of common frog reveal local

805 adaptation. *Proceedings of the National Academy of Sciences*, *107*(18),

806 8292–8297. https://doi.org/10.1073/pnas.0913792107

807 Pollock, K. H. (1982). A capture-recapture design robust to unequal probability of

808 capture. *The Journal of Wildlife Management*, *46*(3), 752–757.

809 https://doi.org/10.2307/3808568

810 Quinn, J. A., & Wetherington, J. D. (2002). Genetic variability and phenotypic

811 plasticity in flowering phenology in populations of two grasses. *The Journal of*

812 *the Torrey Botanical Society*, *129*(2), 96–106. https://doi.org/10.2307/3088723

813 R Core Team. (2020). *R: A language and environment for statistical computing. R*

814 *foundation for statistical computing, Vienna, Austria. URL https://www.R-*

815 *project.org/*.

816 R Studio Team. (2022). *RStudio: Integrated Development for R*. RStudio, PBC.,

817 Boston, MA.

Rahmstorf, S., & Coumou, D. (2011). Increase of extreme events in a warming

world. *Proceedings of the National Academy of Sciences*, *108*(44), 17905–

17909. https://doi.org/10.1073/pnas.1101766108

Reading, C. J. (2003). The effects of variation in climatic temperature (1980–2001)

on breeding activity and tadpole stage duration in the common toad, *Bufo*

bufo. *Science of The Total Environment*, *310*(1–3), 231–236.

https://doi.org/10.1016/S0048-9697(02)00643-5

Reading, C. J. (2007). Linking global warming to amphibian declines through its

effects on female body condition and survivorship. *Oecologia*, *151*(1), 125–

131. https://doi.org/10.1007/s00442-006-0558-1

Reading, C. J. (2010). The impact of environmental temperature on larval

development and metamorph body condition in the common toad, *Bufo bufo*.

Amphibia-Reptilia, *31*(4), 483–488.

https://doi.org/10.1163/017353710X521537

Reading, C. J., & Clarke, R. T. (1983). Male breeding behaviour and mate

acquisition in the common toad, *Bufo bufo*. *Journal of Zoology*, *201*(2), 237–

246. https://doi.org/10.1111/j.1469-7998.1983.tb04273.x

Reading, C. J., & Clarke, R. T. (1999). Impacts of climate and density on the duration

of the tadpole stage of the common toad *Bufo bufo*. *Oecologia*, *121*(3), 310–

315. https://doi.org/10.1007/s004420050933

Rebetez, M., & Reinhard, M. (2008). Monthly air temperature trends in Switzerland

1901–2000 and 1975–2004. *Theoretical and Applied Climatology*, *91*(1), 27–

34. https://doi.org/10.1007/s00704-007-0296-2

Reinhardt, T., Steinfartz, S., & Weitere, M. (2015). Inter-annual weather variability

can drive the outcome of predator prey match in ponds. *Amphibia-Reptilia*,

36(2), 97–109. https://doi.org/10.1163/15685381-00002982

Scherrer, D., & Körner, C. (2011). Topographically controlled thermal-habitat

differentiation buffers alpine plant diversity against climate warming. *Journal*

of Biogeography, *38*(2), 406–416. https://doi.org/10.1111/j.1365-

2699.2010.02407.x

Semlitsch, R. D. (1985). Analysis of climatic factors influencing migrations of the

salamander *Ambystoma talpoideum. Copeia*, *1985*(2), 477–489.

https://doi.org/10.2307/1444862

Semlitsch, R. D., Scott, D. E., Pechmann, J. H. K., & Gibbons, J. W. (1993).

Phenotypic variation in the arrival time of breeding salamanders: Individual

repeatability and environmental influences. *The Journal of Animal Ecology*,

62(2), 334. https://doi.org/10.2307/5364

Sinsch, U. (1988). Seasonal changes in the migratory behaviour of the toad *Bufo*

bufo: Direction and magnitude of movements. *Oecologia*, *76*, 390–398.

Sinsch, U., & Schäfer, A. M. (2016). Density regulation in toad populations (*Epidalea*

calamita, *Bufotes viridis*) by differential winter survival of juveniles. *Journal of*

Thermal Biology, *55*, 20–29. https://doi.org/10.1016/j.jtherbio.2015.11.007

Stoffel, M. A., Nakagawa, S., & Schielzeth, H. (2017). RptR: repeatability estimation

and variance decomposition by generalized linear mixed-effects models.

Methods in Ecology and Evolution, *8*, 1639–1644.

https://doi.org/doi:10.1111/2041-210X.12797

Sztatecsny, M., & Schabetsberger, R. (2005). Into thin air: Vertical migration, body

condition, and quality of terrestrial habitats of alpine common toads, *Bufo*

bufo. Canadian Journal of Zoology, *83*, 788–796.

Tang, J., Körner, C., Muraoka, H., Piao, S., Shen, M., Thackeray, S. J., & Yang, X.

(2016). Emerging opportunities and challenges in phenology: A review.

Ecosphere, *7*(8), e01436. https://doi.org/10.1002/ecs2.1436

Thompson, L. (2000). Ice core evidence for climate change in the Tropics:

Implications for our future. *Quaternary Science Reviews*, *19*, 19–35.

https://doi.org/10.1016/S0277-3791(99)00052-9

Thornton, P. E., Running, S. W., & White, M. A. (1997). Generating surfaces of daily

meteorological variables over large regions of complex terrain. *Journal of*

Hydrology, *190*(3), 214–251. https://doi.org/10.1016/S0022-1694(96)03128-9

Todd, B., Scott, D., Pechmann, J., & Gibbons, J. (2011). Climate change correlates

with rapid delays and advancements in reproductive timing in an amphibian

community. *Proceedings. Biological Sciences / The Royal Society*, *278*,

2191–2197. https://doi.org/10.1098/rspb.2010.1768

Tryjanowski, P., Rybacki, M., & Sparks, T. (2003). Changes in the first spawning

dates of common frogs and common toads in western Poland in 1978–2002.

Annales Zoologici Fennici, *40*, 459–464.

Turner, R. K., & Maclean, I. M. D. (2022). Microclimate-driven trends in spring-

emergence phenology in a temperate reptile (*Vipera berus*): Evidence for a

potential "climate trap"? *Ecology and Evolution*, *12*(2), e8623.

https://doi.org/10.1002/ece3.8623

Urban, M. C. (2018). Escalator to extinction. *Proceedings of the National Academy of*

Sciences, *115*(47), 11871–11873. https://doi.org/10.1073/pnas.1817416115

Vaillant, J. L., Potti, J., Camacho, C., Canal, D., & Martínez-Padilla, J. (2021). Low

repeatability of breeding events reflects flexibility in reproductive timing in the

pied flycatcher *Ficedula hypoleuca* in Spain. *Ardeola*, *69*(1), 21–39.

https://doi.org/10.13157/arla.69.1.2022.ra2

Visser, M. E., & Gienapp, P. (2019). Evolutionary and demographic consequences of

phenological mismatches. *Nature Ecology & Evolution*, *3*(6), 879–885.

https://doi.org/10.1038/s41559-019-0880-8

Vitasse, Y., Ursenbacher, S., Klein, G., Bohnenstengel, T., Chittaro, Y., Delestrade,

A., Monnerat, C., Rebetez, M., Rixen, C., Strebel, N., Schmidt, B. R., Wipf, S.,

Wohlgemuth, T., Yoccoz, N. G., & Lenoir, J. (2021). Phenological and

elevational shifts of plants, animals and fungi under climate change in the

European Alps. *Biological Reviews*, *96*(5), 1816–1835.

https://doi.org/10.1111/brv.12727

Wells, K. D. (1977). The social behaviour of anuran amphibians. *Animal Behaviour*,

25, 666–693. https://doi.org/10.1016/0003-3472(77)90118-X

While, G. M., & Uller, T. (2014). Quo vadis amphibia? Global warming and breeding

phenology in frogs, toads and salamanders. *Ecography*, *37*(10), 921–929.

https://doi.org/10.1111/ecog.00521

Zani, P. A. (2008). Climate change trade-offs in the side-blotched lizard (*Uta*

stansburiana): Effects of growing-season length and mild temperatures on

winter survival. *Physiological and Biochemical Zoology*, *81*(6), 797–809.

https://doi.org/10.1086/588305

Zeileis, A., & Grothendieck, G. (2005). zoo: S3 infrastructure for regular and irregular

time series. *Journal of Statistical Software*, *14*, 1–27.

https://doi.org/10.18637/jss.v014.i06

Appendix

Figure S1. *Graphical visualisation of the first two principal components of our principal component analysis. The points represent the scores over the two axes, while the arrows represent the loadings of the five environmental variables. The first principal component explained 43.2% of the variance in the data, while the second principal component explained 27.7%. T_{Sp} is the average minimum daily spring temperature, T_W is the average minimum daily winter temperature, $Prec_{Sp}$ is the total precipitation in spring, SWE_{Sp} is the total spring snow water equivalent and SWE_W is the total winter snow water equivalent*

924
925 ***Figure S2.*** *Mean standard deviation of the residuals of the 1000 piecewise regressions on*
926 *start of the breeding season **(A)** and 1000 piecewise regression on peak of the breeding*
927 *season **(B)**, calculated with a rolling window approach. Window size was 25% of the length of*
928 *the timeseries (i.e., 10 years out of 40 years, therefore we had 31 windows).*

Table S1. *Summary of the piecewise regressions on the five focal environmental variables. MinT_Spring is the average minimum daily spring temperature, MinT_Winter is the average minimum daily winter temperature, Prec_Spring is the total precipitation in spring, SWE_Spring is the total spring snow water equivalent and SWE_Winter is the total winter snow water equivalent. For both intercept and slopes we show the estimate, its standard error, and the t-value and p-value associated with it. Slope1 refers to the segment before the breakpoint and Slope2 refers to the segment after the breakpoint. We also show the statistics associated with the identified breakpoint. Asterisks next to the p-values show significance at the 0.05 level. The p-value for Slope2 is NA since standard asymptotics do not apply (Muggeo, 2008).*

		Estimates	Std. Error	t-value	p-value
MinT_Spring	Intercept	-130.32	39.20	-3.33	0.002*
	Slope1	0.07	0.02	3.33	0.002*
	Slope2	-0.05	0.05	-0.87	NA
	Breakpoint	2007	4.60	-1.92	0.06
MinT_Winter	Intercept	-479.94	268.55	-1.79	0.08
	Slope1	0.24	0.14	1.76	0.09
	Slope2	-0.03	0.03	-1.07	NA
	Breakpoint	1990	3.56	-1.05	0.30
Prec_Spring	Intercept	-4441.03	4795.75	-0.93	0.36
	Slope1	2.26	2.42	0.94	0.36
	Slope2	-0.30	0.17	-1.75	NA
	Breakpoint	1986	3.91	-1.51	0.14
SWE_Spring	Intercept	437670.0	262926.3	1.67	0.10
	Slope1	-219.2	132.5	-1.66	0.11
	Slope2	37.68	24.17	1.56	NA
	Breakpoint	1989.87	3.55	2.23	0.03*
SWE_Winter	Intercept	630172.0	526017.4	1.20	0.24
	Slope1	-316.8	265.4	-1.19	0.24
	Slope2	19.40	13.47	1.44	NA
	Breakpoint	1984.96	2.72	1.86	0.07

Table S2. *Details of the five principal components. For each principal component we report its standard deviation, the proportion of variance explained and the cumulative proportion of this variance. In our linear regression we kept the first two principal components as their standard deviation is >1 and combined they explain >70 % of the variance.*

	PC1	PC2	PC3	PC4	PC5
Standard deviation	1.47	1.18	0.83	0.82	0.33
Proportion of variance	0.43	0.28	0.14	0.13	0.02
Cumulative proportion	0.43	0.71	0.85	0.98	1.00

Table S3. *Loadings of the five original environmental variables from which the five principal components are constructed. MinT_Spring is the average minimum daily spring temperature, MinT_Winter is the average minimum daily winter temperature, Prec_Spring is the total precipitation in spring, SWE_Spring is the total spring snow water equivalent and SWE_Winter is the total winter snow water equivalent. PC1 is mainly driven by MinT_Winter, SWE_Spring and SWE_Winter, while PC2 mostly by MinT_Spring and Prec_Spring.*

	PC1	PC2	PC3	PC4	PC5
MinT_Spring	0.08	-0.68	0.33	-0.65	0.09
MinT_Winter	0.45	-0.19	0.64	0.58	0.07
Prec_Spring	-0.04	0.68	0.60	-0.41	-0.06
SWE_Spring	-0.64	-0.05	0.20	0.17	0.72
SWE_Winter	-0.61	-0.22	0.27	0.19	-0.69