1

Genomic health is dependent on long-term population demographic history

Eric Wootton[1], Claude Robert[2], Joëlle Taillon[3], Steeve Côté[4], Aaron B.A. Shafer[5]

[1]Biochemistry & Molecular Biology, Trent University, K9J 7B8 Peterborough, Ontario, Canada

[2]Département des Sciences Animales, Université Laval, G1V 0A6, Québec, QC, Canada

[3]Direction de l'Expertise sur la Faune Terrestre, l'Herpétofaune et l'Avifaune, Ministère des
Forêts, de la Faune et des Parcs, Gouvernement du Québec, Québec, QC, Canada

[4]Département de Biologie, Caribou Ungava and Centre d'Études Nordiques, Université Laval,
G1V 0A6, Québec, QC, Canada

[5]Environmental and Life Sciences Graduate Program, Trent University, 7K9J 7B8 Peterborough,
Ontario, Canada

Corresponding authors:

Eric Wootton: ericwootton@trentu.ca

Aaron Shafer: aaronshafer@trentu.ca

Abstract

Current genetic methods of population assessment in conservation biology have been challenged by genome-scale analyses due to their quantitatively novel insights. These analyses include assessments of runs-of-homozygosity (ROH), genomic evolutionary rate profiling (GERP), and mutational load. Here, we aim to elucidate the relationships between these measures using three divergent ungulates: the white-tailed deer, caribou, and mountain goat. The white-tailed deer is currently expanding, while caribou are in the midst of a significant decline. Mountain goats remain stable, having suffered a large historical bottleneck. We assessed genome-wide signatures of inbreeding using the inbreeding coefficient F and %ROH (F_{ROH}) and identified evolutionarily constrained regions with GERP. Mutational load was estimated by identifying mutations in highly constrained elements (CEs) and sorting intolerant from tolerant (SIFT) mutations. Our results show that F and F_{ROH} are higher in mountain goats than in caribou and white-tailed deer. Given the extended bottleneck and low N_e of the mountain goat, this supports the idea that the genome-wide effects of demographic change take time to accrue. Similarly, we found that mountain goats possess more highly constrained CEs which are indicative of greater purifying selection. This is reflected by fewer mutations in CEs and deleterious mutations identified by SIFT. In contrast, white-tailed deer presented the highest mutational load with both metrics, while caribou were intermediate. Our results demonstrate that extended bottlenecks may lead to reduced diversity and increased F_{ROH} in ungulates, but not necessarily the accumulation of deleterious alleles, likely due to the purging of deleterious alleles in small populations.

Introduction

Organizations like the International Union for Conservation of Nature (IUCN) rely primarily on population sizes and trends when assigning wildlife conservation statuses (IUCN Standards and Petitions Committee, 2019). The predictive relationship between population size (N_C) and genetic diversity is challenging, as N_C, for example, tends to be substantially more variable than diversity metrics like nucleotide diversity (π), referred to as Lewontin's paradox (Ellegren & Galtier, 2016; Lewontin, 1974), and therefore rarely accounts for recent demographic history and the genetic health of populations. While genetic diversity estimates generally correlate with IUCN status (Peart et al., 2020; Petit-Marty et al., 2021), the relationship (i.e., direction) is varied and predictive association is low (Schmidt et al., 2022). The unprecedented loss of global biodiversity (Pascual et al., 2021) and ambiguous associations (Schmidt et al., 2022) necessitates the need for conservation-informative genetic metrics.

Both novel genome metrics and functional versus presumed neutral variation have garnered attention in the conservation genetics community. The importance of neutral genetic diversity for example, despite a long history (e.g. DeWoody et al., 2021; Kohn et al., 2006), was recently called into question (Teixeira & Huber, 2021). Citing Lewontin (1974), Teixera & Huber (2021) argued for a renewed focus on functional variation and gene-environment interactions, which prompted two high-profile responses (Garcia-Dorado & Caballero, 2021; Kardos et al., 2021). Kardos et al. (2021) in particular echoed earlier concerns of [functional] gene-targeted conservation (i.e. Kardos & Shafer, 2018), and highlighted some potential problems in Teixeira & Huber's simulations. Kardos et al. (2021) argued against abandoning genome-wide neutral markers; supporting this view would be empirical work from Peart et al. (2020) that showed genome wide estimates of Tajima's *D*, a composite statistic derived from

Watterson's θ and π (Tajima, 1989), appears predictive of conservation status, further reiterating

the value of classic neutral summary statistics derived from genome-wide sequences.

The high correlation between nucleotide diversity and microsatellite heterozygosity

suggests that genome-scale data reflect smaller targeted panels of loci (Väli et al., 2008). This

observation is ultimately what formed the premise of Shafer et al. (2015) that argued for the use

of quantitatively novel genomic metrics for conservation. Runs-of-homozygosity (ROH),

genomic evolutionary rate profiling (GERP), and mutational load are metrics dependent on

genome-scale data and have only recently been applied to species and populations of concern

(Grossen et al., 2020; van der Valk et al., 2019; von Seth et al., 2021). ROHs are long stretches

of homozygosity that arise as a result of inbreeding (Ceballos et al., 2018), and reflect the true

inbreeding coefficient and can identify genes underlying inbreeding depression (Kardos et al.,

2016). GERP profiling can be used to identify evolutionarily constrained genomic elements,

defined as regions with fewer substitutions than expected under neutral evolution (Davydov et

al., 2010), which are often the subject of selective pressures (Cooper et al., 2005; Davydov et al.,

2010). Any mutations that appear at highly conserved sites are generally assumed to be

deleterious (Cooper et al., 2005; Davydov et al., 2010). Likewise, tolerant and intolerant

mutations – also referred to as SIFT scores (Vaser et al., 2015) – that are considered of human

clinical interest (P. Kumar et al., 2009) – have now been applied to natural populations

(Robinson et al., 2016). ROH, GERP and SIFT scores collectively allow for inferring the

mutational load and genomic health of a population or species.

While quantitatively useful, the relationships between these metrics and conservation

status and diversity metrics are generally not well defined and at-times counter to expectation.

van der Valk et al. (2019) showed a weak relationship between conservation status and

mutational load across mammalian taxa; the highest mutational loads were in fact seen in the

most abundant and diverse species. Similarly, species with the highest inbreeding (ROH) also

had the lowest mutational load and GERP scores (van der Valk et al., 2019). This pattern could

be due the purging of deleterious alleles in small populations, which is consistent with

simulations by Kyriazis et al. (2021). van der Valk et al. (2019) argued that this process has

implications for genetic rescue and suggested that individuals be selected for translocation based

on low mutational load and not diversity. Work on the Sumatran rhinoceros (*Dicerorhinus*

sumatrensis) added another layer of nuance, with populations showing low to moderate levels of

inbreeding (ROH), but a relatively high mutational load (von Seth et al., 2021). In contrast, the

Alpine ibex (*Capra ibex*) has experienced successive demographic bottlenecks that led to

reduced diversity and increased ROH, but not the accumulation of highly deleterious alleles

(Grossen et al., 2020). Thus, there seems to be a link between historical demography, ROH, and

mutational load. Understanding such processes has important implications for both predicting

long-term viability, but also selecting animals for genetic rescue.

Here, we analysed signatures of genomic health across populations of three divergent

ungulates to elucidate these relationships. The caribou (*Rangifer tarandus*) is currently

experiencing significant population declines (N_C) prompting international concern (COSEWIC,

2015). In Quebec, five of the major populations have seen drastic declines from their highest

estimates ranging between 10% and 99%, depending on the herd (COSEWIC, 2015, 2017). The

white-tailed deer (*Odocoileus virginianus*), in contrast, is expanding northward and has

accordingly high genomic diversity (Kessler et al., 2022). While white-tailed deer currently

number in the 10s of millions (Kauffman et al., 2018), they were historically extirpated across

much their range due to colonial harvest and forestry practices (Kauffman et al., 2018); meaning

this demographic expansion is very recent. The mountain goat (*Oreamnos americanus*)

underwent a massive bottleneck during the last glacial maximum (Martchenko et al., 2020) but

has since remained relatively stable with a range-wide population of ~100,000 individuals

(Festa-Bianchet & Côté, 2008). These three species, contrasted in their ecology but also

conservation status, historical and demographic trajectories, present useful comparisons to

explore patterns of novel genomic metrics as they pertain to conservation status. In line with

recent empirical work, we predicted that populations with low diversity and high inbreeding do

not necessarily have higher mutational load and that genome-wide effects of demographic

change like ROH take time to accrue. Similarly, we expected to see greater purging of

deleterious alleles in populations with long-term small N_e. Collectively, this means that poor

genomic metrics of health would not correlate with current negative population trends.

Methods

Caribou samples were obtained from 67 individuals across Quebec and Labrador (Dedato et al.

2022); 53 white-tailed deer were sampled across their North American range (Kessler et al.

2022). Additionally, 35 mountain goats were sampled across their native range from Alaska to as

far south as Idaho (Martchenko et al. 2022; Figure 1). All samples were sequenced with Illumina

HiSeqX technology to approximately 5X coverage. Non-target sequences were trimmed from

raw FASTQ files with Trimmomatic v.0.36 (Bolger et al., 2014), and reads were mapped to their

reference genome using BWA-MEM v.0.7.17 and SAMtools v.1.15 (Li et al., 2009; Li &

Durbin, 2009). Duplicate mapped reads were marked and removed using Picard v.2.23.0 (Broad

Institute, 2019). Uniquely mapped reads were subset by Sambamba v.0.8.2 (Streit et al., 2012),

and indels were realigned with GATK v.3.8 (Van der Auwera et al., 2013). We estimated

genotype likelihoods using ANGSD v.0.934 using the GATK model: -doVcf 1 -doGeno -4 -

doPost 1 -gl 2 -SNP_pval 1e-6 -minMapQ 20 -minQ 20 -doCounts 1 -skipTriallelic 1 -doGlf 2

(Korneliussen et al., 2014). Hard calls required for analyses were performed by Beagle v.5.1

(Browning et al., 2018) with the following parameters: impute=false overlap=3000

window=100000 gprobs=TRUE parameters. Posterior genotype probabilities were estimated

using allele frequencies, and only genotypes with a posterior probability above 0.95 were called.

Tajima's D and nucleotide diversity (π) were calculated using VCFtools v.0.1.13 with windows

of 10 Kb and using all positions, regardless of allele frequency (Danecek et al., 2011).

Inbreeding and Homozygosity

A subset of SNPs in approximate linkage equilibrium was obtained using PLINK v.1.9 with a

window size of 50 SNPs, a window shift size of 10 SNPs, and a pairwise (R^2) linkage

disequilibrium threshold of 0.1 (Chang et al., 2015; Purcell et al., 2007). We then identified runs

of homozygosity (ROH) on scaffolds greater than 10 Mb in length and with a minor allele

frequency cut-off of 0.05 (Foote et al., 2021). Sliding windows of 50 SNPs were used, and each

ROH was defined as a sequence at least 100 Kb long with at least 50 homozygous SNPs (Foote

et al., 2021). Each ROH required a minimum density of 1 SNP per 50 Kb and no more than 5

missing or heterozygous calls in a window. Any SNPs more than 1 Mb apart within a segment

were used to split that segment in two. Once identified, each ROH was classified by size, and the

percent ROH (F_{ROH}) for each scaffold and individual were calculated. PLINK parameters were

taken from Foote et al. (2021) with output validated using the program ROHan (Renaud et al.,

2019). ROHan was run on the same data with the same ROH size definition of 100 Kb and a

transition to transversion ratio (Ts/Tv) of 2.25. This ratio was calculated using BCFtools v.1.14

and the genotypes provided by ANGSD (Li et al., 2009). The inbreeding coefficient (F) for each

individual was also estimated using PLINK and the –het flag.

Evolutionary Constraint and Mutational Load

Twenty ungulate reference genomes from the NCBI GenBank genome database (Table S1) were selected to represent most extant families with genomes available in both the orders of Perissodactyla (odd-toed ungulates) and Artiodactyla (even-toed ungulates). Constrained regions for each individual of our three focal species were identified using the Genomic Evolutionary Rate Profiling (GERP++); this approach requires a phylogenetic tree which we constructed for all 23 species using TimeTree (Figure S1) (Kumar et al., 2017). The individual fastq files of all individuals of the three species were then aligned to each of the 20 reference genomes in 50 bp segments using BWA-MEM. All duplicate alignments were marked and removed with Picard. Then, the alignments of each individual were concatenated into one multiple alignment file (MAF), resulting in 155 total MAFs. The gerpcol function included in GERP++ was run on each MAF to generate rejected substitution (RS) scores for each alignment column. RS scores are used as a metric for constraint intensity and are calculated by subtracting the estimated evolutionary rate at each position from the neutral rate (Cooper et al., 2005; Davydov et al., 2010). Both rates were estimated by gerpcol using the aforementioned phylogenetic tree (Figure 2). RS scores quantify the number of differences between the expected and observed number of SNPs at a site, and a positive score is indicative of constraint (Cooper et al., 2005; Davydov et al., 2010). An RS score greater than 2 represents a highly constrained region whereas a score lower than 1 is putatively neutral and discarded (Cooper et al., 2005; Davydov et al., 2010). gerpcol was run with the previously calculated Ts/Tv ratio of 2.25. We also used GERP++ to identify constrained elements with the gerpelem function.

To estimate mutational load, we determined the number of SNPs within these constrained

elements using BEDtools v.2.30.0 (Quinlan & Hall, 2010). These SNPs are putatively

deleterious due to the common functional importance of constrained regions (Cooper et al.,

2005; Davydov et al., 2010). For each species, the functional consequence of each SNP was

determined using Ensembl's Variant Effect Predictor (VEP) (McLaren et al., 2016). The

deleteriousness of each SNP across the genome was then estimated using functional tolerance

scores provided by SIFT and SIFT4G (Vaser et al., 2015). Using multiple alignment information,

SIFT and SIFT4G calculate normalised probabilities where substitutions with scores lower than

0.05 are predicted to be deleterious to protein function (Vaser et al., 2015).

Caribou did not have an available annotation at the time of analysis which is required for

this analysis; thus, we used a mixture of *de novo* and homology-based predictions. We aligned

protein sequences from *Bos taurus, Equus caballus, Homo sapiens,* and *Ovis aries* to the caribou

genome using blastx v.2.7.1 (Altschul et al., 1990). These sequences were all obtained from

Ensembl release 89 (Hunt et al., 2018). We also aligned transcripts from the NCBI mammalian

RefSeq transcript database v.1.0 using BLAT v.1.0 to help identify exons. Caribou RNAseq data

(SRX12486288) from the NCBI Sequence Read Archive were aligned to the genome with

HISAT v.2.10 for *de novo* prediction (Kim et al., 2015). Using these data, genes were predicted

by Augustus v.3.1.1 as well as GeneMarkES v.4.69 (Lomsadze et al., 2005; Stanke et al., 2006).

We generated a set of consensus genes using EVidenceModeler v.1.1.1 with the following

weighting scheme (Haas et al., 2008): transcript alignment via BLAT (10x weight); protein

alignment via blastx (5x weight); gene prediction via Augustus (1x weight); gene prediction via

GeneMarkES (1x weight). PASA v.2.3.3 was used last to refine gene identifications (Haas et al.,

2008).

For all individual metrics above we quantified the correlation to sample location (latitude,

longitude). Because white-tailed deer were extirpated across much of their range, we used

translocation data from McDonald & Miller, (2004) to determine if these metrics were correlated

to restocking efforts in the United States. We examined correlations with stock start and end

dates, estimated population lows, number of individuals stocked, and the number of states

involved in stocking.

Results

Diversity and Inbreeding

Nucleotide diversity (π) was found to be highest in deer (mean = 0.00771) and lowest in

mountain goats (mean = 0.00101) (Figure 1). Tajima's D was highest in mountain goats (mean =

0.625) and lowest in white-tailed deer (mean = -0.645) (Figure 1). For caribou, Tajima's D was

positive (mean = 0.124), but lower than in mountain goats (t-test p-value $< 2.2 \times 10^{-16}$). Mountain

goats were found to have significantly higher F_{ROH} (mean = 0.038) than both caribou (none

detected) and white-tailed deer (mean = 0.0020) (Figure 2a). This pattern is reflected in the

relative values of F where mountain goats had the highest value (mean = -0.018) in comparison

to caribou (mean = -0.44) and deer (mean = -0.27) (Figure 2b). For the mountain goat, most

ROHs were found to be in the 500-1000 Kb range whereas deer possessed ROHs almost

exclusively below 300 Kb (Figure 3). Despite a much higher number of ROHs overall, mountain

goats possessed significantly fewer in the 100-300 Kb range than deer (Figure 3). Both measures

of inbreeding were found to be correlated in mountain goats (Pearson's $r = 0.840$, $p < 0.05$) and

deer (Pearson's $r = 0.395$, $p < 0.05$) (Figure S2), noting again there were no F_{ROH} in caribou

preventing a similar assessment.

Evolutionary Constraint and Mutational Load

Mountain goats had the highest number of constrained elements (CEs) (mean = 2926) while caribou have the fewest (mean = 935; Figure S3). Highly constrained elements followed a nearly identical distribution between species with all three having approximately 50% as many CEs with RS > 2 (Figure S3). In addition to having the most CEs, mountain goats showed slightly longer CEs on average (Figure S4). The three species showed little variability in average CE RS-score (Figure S4, S5). Despite having significantly fewer CEs, white-tailed deer had a larger number of SNPs in CEs (mean = 411) followed by mountain goats (mean = 212) and caribou (mean = 74) (Table 1). The relative differences between species remained the same for highly constrained CEs (Table 1). We observed that a much higher fraction of constrained positions was deleterious in deer (mean = 0.625%) than in caribou (mean = 0.326%) and mountain goats (mean = 0.121%; Figure 4).

We found 42.1% of coding sequence consequences identified by VEP to be missense variants in white-tailed deer, 49.3% in caribou, and 57.4% in mountain goats (Figure 5a). Despite this distribution, white-tailed deer had the highest number of intolerant mutations (mean = 6852) identified by SIFT followed by caribou (mean = 4944) and mountain goats (mean = 3190) (Table 1). Similarly, we observed white-tailed deer to have the highest fraction of genes with deleterious mutations (mean = 0.330%) while caribou (mean = 0.149%) and mountain goats (mean = 0.143%) had the lowest (Figure 5b). None of these metrics were correlated to the number of genes in the annotation (Table 1). Lastly, we found F and latitude to be moderately correlated in caribou (Pearson's r = -0.466, $p < 0.05$; Figure S6) and deer (Pearson's r = -0.513, $p < 0.05$; Figure S6), but not in mountain goats (Pearson's r = 0.003, $p < 0.05$). There were no

significant correlations between the individual genome health metrics and restocking data in

white-tailed deer (Pearson's $r < 0.4$, $p > 0.05$; Table S2).

Discussion

The interplay between nucleotide diversity (π), inbreeding, mutational load, and census size (N_C),

is nuanced and full of contrasting examples. Here, we demonstrate that mutational load may be

lower than expected in populations with relatively high inbreeding and low diversity. Our

observed patterns suggest that the purging of deleterious alleles in these cases is stronger in small

populations (Huber et al., 2020), and therefore would reduce risks from inbreeding depression.

This supports our hypotheses that low diversity and high inbreeding do not on their own

necessitate special conservation status, and that purging is likely occurring in populations with

low N_e. We also predicted that genomic signatures of demographic events take time to accrue,

which is supported by evidence of F_{ROH} and CEs requiring extended bottlenecks to surface and

increase. Caribou have seen recent decreases in N_C yet have no detectable ROH; in contrast,

mountain goats show elevated ROH likely due to the ancestral nature of their bottleneck

(Martchenko et al., 2020). The varied relationships between different measures of genomic

health and the lack of correlation to the population trends that are typically used for status

assignment highlights the value of this comparative framework (see also Huber et al., 2020).

This information is especially valuable for small populations of conservation concern, where

mutational load combined with ROH and CE assessments is probably the most informative way

forward, in addition to metrics like Tajima's *D* (Peart et al., 2020).

Inbreeding Effects

Genomic consequences stemming from demographic take time to accumulate (e.g. Grossen et

al., 2020; von Seth et al., 2021). Several caribou populations are currently in the midst of a

demographic bottleneck resulting in an extremely low contemporary effective population size

(N_e) and nucleotide diversity (Dedato et al., 2022). In contrast, mountain goats underwent a large

historical bottleneck during the last glacial maximum, and while they have not recovered, they

have remained relatively stable, resulting in a moderately low N_e (Martchenko & Shafer, 2022).

Low diversity stemming from bottleneck is supported by positive values of Tajima's D in these

two species, which indicate a lack of rare alleles and population contraction (Tajima, 1989). The

lower π in mountain goats is clearly due to their extended bottleneck that would have

exacerbated any effects of inbreeding, but also their historically lower census sizes. These effects

are reflected by F_{ROH}, where mountain goats were found to have 3.8% of the genome in ROH

whereas caribou had virtually no detectable ROH segments over 100 Kb long. Here, despite a

less drastic decline in N_C, mountain goats have accumulated a significantly greater F_{ROH} than

caribou. These differences are also unlikely to be the result of genome assembly quality, as the

mean number and length of analysed scaffolds were comparable (Table 1). Instead, it is likely

that any stretches of homozygosity in caribou were too short to be classified as ROHs by our

methods (<100 Kb). This is a common finding in mammalian genomes, where most species have

no ROHs above this threshold and possess a low ROH burden (Brüniche-Olsen et al., 2018).

Compared to white-tailed deer, mountain goats have developed significantly longer and

more numerous ROHs, which further outline the effects of a prolonged bottleneck. Caribou

simply have not been subject to a low N_C for long enough to approach mountain goats or even

differentiate themselves from deer in their levels of inbreeding. Of note, F_{ROH} was more

correlated to F after an extended bottleneck (i.e., mountain goats vs. white-tailed deer; Figure

S2), which further implies that ROHs require both time and demographic change (Gillespie,

2004) to accumulate. Overall, these results support the idea that genome-wide effects of

demographic change will not accrue unless given enough time; F and F_{ROH} are therefore unlikely

to align with IUCN conservation decisions, similarly noted by Schmidt et al. (2022).

Mutational Load and Purging

The relationship between N_e, inbreeding, π, and mutational load are variable (Grossen et al.,

2020; van der Valk et al., 2019; von Seth et al., 2021). Empirical studies on mammals have led

us to predict that low diversity and high inbreeding do not necessarily result in the accumulation

of deleterious alleles (Grossen et al., 2020; Robinson et al., 2018, 2018; van der Valk et al.,

2019). Here, we found a much greater number of putatively deleterious mutations in white-tailed

deer than in mountain goats and caribou by identifying CE-bound SNPs with GERP. These

results were confirmed by the deleterious alleles identified by SIFT that showed a similar gap

between white-tailed deer and the other species, regardless of the annotation quality (Table 1).

This means that despite lower diversity and small N_e, caribou and mountain goats have relatively

low mutational loads; further, despite a relatively low number of CEs, deer still possessed

significantly more CE-bound SNPs indicative of deleterious mutations. Mountain goats, in

contrast, were found to have the highest amounts of inbreeding but still not accumulated many

deleterious alleles. This reiterates the finding that inbreeding does not necessarily result in high

mutational load; in the case of caribou, if N_C can return (e.g., Peart et al. 2020) or if purging of

deleterious alleles can occur (e.g., Huber et al., 2020), the genetic cost to the current decline in

N_C could be minimized. This, however, remains an open question and wolves in Europe provide

an informative example (Smeds & Ellegren, 2022).

The purging of deleterious alleles in small populations is rarely observed (Hedrick &

Garcia-Dorado, 2016; Robinson et al., 2018; Xue et al., 2015), but there is evidence that it may

occur in the presence of severe bottlenecks (Grossen et al., 2020). Mountain goats were shown to

have low mutational load in conjunction with generally low diversity, while caribou present

intermediate profiles. This would be consistent with N_C numbers pre-20[th] century for caribou,

that were presumably much higher (Festa-Bianchet et al., 2011). Reductions in N_e from severe

bottlenecks can increase the fitness effects of recessive deleterious alleles by increasing

homozygosity (Garcia-Dorado, 2012, 2015); this strengthens the pressure of negative selection

and can ultimately lower the mutational load of a population (Garcia-Dorado, 2012, 2015;

Glémin, 2003). However, this effect can be insufficient for overcoming cases of extensive

homozygosity. Sumatran rhinos for example have a very similar historical N_e to that of mountain

goats (von Seth et al., 2021), but considerably more ROH. While purging might have played

some role in some populations (von Seth et al., 2021), the overall negative effect of autozygosity

and continued presence of high frequency deleterious alleles (e.g., Khan et al., 2021) was likely

insurmountable from a population viability standpoint.

Purifying selection should remove SNPs in functionally important regions and constrain

these areas against change, which is why we observed less SNPs than expected in CEs (Cooper

et al., 2005; Davydov et al., 2010). The mountain goat's longer and more numerous CEs relative

to the other species are possibly indicative of stronger purifying selection, the driving force

behind constraint. In contrast, caribou have a smaller N_e than white-tailed deer; thus purging has

either yet to take place or selection is less efficient, or a combination of both. The trajectory of

caribou, would be predicted to be towards the patterns of mountain goat based on mammalian

wide patterns and profiles (Cooper et al., 2005; Huber et al., 2020). It is also important to note

that purifying selection reduces nucleotide diversity by eliminating sites under direct selection

and those linked through background selection (Charlesworth, 1996; Cvijović et al., 2018),

which partially explains patterns of diversity seen here (Figure 1; Buffalo, 2021).

This work has implications for conservation decisions and the expected genomic

consequences of numerical population declines. While deleterious alleles may be purged in

small, inbred populations, high levels of autozygosity can render this benefit insignificant. Thus,

we posit that high mutational load must be accompanied by high ROH to be of conservation

concern, similar to what was observed in woolly mammoths (Rogers & Slatkin, 2017). Further,

genetic rescue decisions should not rely solely on inbreeding and diversity metrics, especially

when their relationships to fitness are not straightforward (Schmidt et al., 2022). This is also true

for assessing restocking efforts, as we demonstrate that translocations are not necessarily related

to population health metrics. Collectively, this provides further evidence of the time needed for

genome-scale changes to accrue after significant changes in demography. Therefore, we

recommend that genomic health metrics like F_{ROH}, mutational load, and evolutionary constraint

be examined alongside classic metrics of population viability for a holistic view that can provide

more accurate assessments of genome health.

Acknowledgements

We wish to acknowledge that the work for this study was carried out on the traditional territory of the Mississauga (Michi Saagiig) Anishinaabeg people. This work was supported by a Natural Sciences and Engineering Research Council Discovery Grant, Canada Foundation for Innovation: John R. Evans Leaders Fund, and Compute Canada Awards to ABAS; EW was supported by three NSERC USRAs. We thank Daria Martchenko, Morgan Dedato, and Camille Kessler for help with data acquisition. Sample providers are listed in Dedato et al. (2022), Martchenko & Shafer (2022), and Kessler et al. (2022).

References

Altschul, S. F., Gish, W., Miller, W., Myers, E. W., & Lipman, D. J. (1990). Basic local alignment search tool. *Journal of Molecular Biology*, *215*(3), 403–410. https://doi.org/10.1016/S0022-2836(05)80360-2

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: A flexible trimmer for Illumina sequence data. *Bioinformatics*, *30*(15), 2114–2120. https://doi.org/10.1093/bioinformatics/btu170

Broad Institute. (2019). *Picard Toolkit*.

Brüniche-Olsen, A., Kellner, K. F., Anderson, C. J., & DeWoody, J. A. (2018). Runs of homozygosity have utility in mammalian conservation and evolutionary studies. *Conservation Genetics*, *19*(6), 1295–1307. https://doi.org/10.1007/s10592-018-1099-y

Buffalo, V. (2021). Quantifying the relationship between genetic diversity and population size suggests natural selection cannot explain Lewontin's Paradox. *ELife*, *10*, e67509. https://doi.org/10.7554/eLife.67509

Ceballos, F. C., Joshi, P. K., Clark, D. W., Ramsay, M., & Wilson, J. F. (2018). Runs of homozygosity:

Windows into population history and trait architecture. *Nature Reviews Genetics 2018 19:4*,

19(4), 220–234. https://doi.org/10.1038/nrg.2017.109

Chang, C. C., Chow, C. C., Tellier, L. C. A. M., Vattikuti, S., Purcell, S. M., & Lee, J. J. (2015). Second-

generation PLINK: rising to the challenge of larger and richer datasets. *GigaScience*, *4*(1), 7.

https://doi.org/10.1186/S13742-015-0047-8

Charlesworth, B. (1996). Background selection and patterns of genetic diversity in Drosophila

melanogaster. *Genetical Research*, *68*(2), 131–149. https://doi.org/10.1017/s0016672300034029

Cooper, G. M., Stone, E. A., Asimenos, G., Comparative, N., Program, S., Green, E. D., Batzoglou, S., &

Sidow, A. (2005). Distribution and intensity of constraint in mammalian genomic sequence.

Genome Research, *15*(7), 901–913. https://doi.org/10.1101/gr.3577405

COSEWIC. (2015). *COSEWIC Assessment and Status Report on the Caribou Rangifer tarandus,*

Newfoundland population, Atlantic-Gaspésie population, Boreal population in Canada.

https://species-registry.canada.ca/index-en.html#/documents/2769

COSEWIC. (2017). *COSEWIC Assessment and Status Report on the Caribou Rangifer tarandus, Eastern*

Migratory population and Torngat Mountains population, in Canada.

http://www.registrelepsararegistry.gc.ca/default.asp?lang=en&n=24F7211B-1

Cvijović, I., Good, B. H., & Desai, M. M. (2018). The Effect of Strong Purifying Selection on Genetic

Diversity. *Genetics*, *209*(4), 1235–1278. https://doi.org/10.1534/genetics.118.301058

Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Handsaker, R. E.,

Lunter, G., Marth, G. T., Sherry, S. T., McVean, G., Durbin, R., & 1000 Genomes Project

Analysis Group. (2011). The variant call format and VCFtools. *Bioinformatics*, *27*(15), 2156–

2158. https://doi.org/10.1093/bioinformatics/btr330

Davydov, E. V., Goode, D. L., Sirota, M., Cooper, G. M., Sidow, A., & Batzoglou, S. (2010). Identifying

a High Fraction of the Human Genome to be under Selective Constraint Using GERP++. *PLOS*

Computational Biology, *6*(12), e1001025. https://doi.org/10.1371/JOURNAL.PCBI.1001025

Dedato, M., Robert, C., Taillon, J., Shafer, A., & Cote, S. (2022). *Demographic history and conservation*

genomics of caribou (Rangifer tarandus) in Québec [Preprint]. Authorea.

https://doi.org/10.22541/au.164165840.02956352/v1

DeWoody, J. A., Harder, A. M., Mathur, S., & Willoughby, J. R. (2021). The long-standing significance

of genetic diversity in conservation. *Molecular Ecology*, *30*(17), 4147–4154.

https://doi.org/10.1111/mec.16051

Ellegren, H., & Galtier, N. (2016). Determinants of genetic diversity. *Nature Reviews Genetics*, *17*(7),

422–433. https://doi.org/10.1038/nrg.2016.58

Festa-Bianchet, M., & Côté, S. D. (2008). *Mountain Goats: Ecology, Behavior, and Conservation of an*

Alpine Ungulate. Island Press.

Festa-Bianchet, M., Ray, J. C., Boutin, S., Côté, S. D., & Gunn, A. (2011). Conservation of caribou

(Rangifer tarandus) in Canada: An uncertain future1This review is part of the virtual symposium

"Flagship Species☐–☐Flagship Problems" that deals with ecology, biodiversity and management

issues, and climate impacts on species at risk and of Canadian importance, including the polar

bear (Ursus maritimus), Atlantic cod (Gadus morhua), Piping Plover (Charadrius melodus), and

caribou (Rangifer tarandus). *Canadian Journal of Zoology*, *89*(5), 419–434.

https://doi.org/10.1139/z11-025

Foote, A. D., Hooper, R., Alexander, A., Baird, R. W., Baker, C. S., Ballance, L., Barlow, J., Brownlow,

A., Collins, T., Constantine, R., Dalla Rosa, L., Davison, N. J., Durban, J. W., Esteban, R.,

Excoffier, L., Martin, S. L. F., Forney, K. A., Gerrodette, T., Gilbert, M. T. P., … Morin, P. A.

(2021). Runs of homozygosity in killer whale genomes provide a global record of demographic

histories. *Molecular Ecology*, *30*(23), 6162–6177. https://doi.org/10.1111/mec.16137

Garcia-Dorado, A. (2012). Understanding and Predicting the Fitness Decline of Shrunk Populations:

Inbreeding, Purging, Mutation, and Standard Selection. *Genetics*, *190*(4), 1461–1476.

https://doi.org/10.1534/genetics.111.135541

Garcia-Dorado, A. (2015). On the consequences of ignoring purging on genetic recommendations for

minimum viable population rules. *Heredity*, *115*(3), 185–187.

https://doi.org/10.1038/hdy.2015.28

Garcia-Dorado, A., & Caballero, A. (2021). Neutral genetic diversity as a useful tool for conservation

biology. *Conservation Genetics*, *22*(4), 541–545. https://doi.org/10.1007/s10592-021-01384-9

Gillespie, J. H. (2004). *Population Genetics: A Concise Guide*. JHU Press.

Glémin, S. (2003). How Are Deleterious Mutations Purged? Drift Versus Nonrandom Mating. *Evolution*,

57(12), 2678–2687. https://doi.org/10.1111/j.0014-3820.2003.tb01512.x

Grossen, C., Guillaume, F., Keller, L. F., & Croll, D. (2020). Purging of highly deleterious mutations

through severe bottlenecks in Alpine ibex. *Nature Communications*, *11*(1), 1001.

https://doi.org/10.1038/s41467-020-14803-1

Haas, B. J., Salzberg, S. L., Zhu, W., Pertea, M., Allen, J. E., Orvis, J., White, O., Buell, C. R., &

Wortman, J. R. (2008). Automated eukaryotic gene structure annotation using EVidenceModeler

and the Program to Assemble Spliced Alignments. *Genome Biology*, *9*(1), R7.

https://doi.org/10.1186/gb-2008-9-1-r7

Hedrick, P. W., & Garcia-Dorado, A. (2016). Understanding Inbreeding Depression, Purging, and

Genetic Rescue. *Trends in Ecology & Evolution*, *31*(12), 940–952.

https://doi.org/10.1016/j.tree.2016.09.005

Huber, C. D., Kim, B. Y., & Lohmueller, K. E. (2020). Population genetic models of GERP scores

suggest pervasive turnover of constrained sites across mammalian evolution. *PLoS Genetics*,

16(5), e1008827. https://doi.org/10.1371/journal.pgen.1008827

Hunt, S. E., McLaren, W., Gil, L., Thormann, A., Schuilenburg, H., Sheppard, D., Parton, A., Armean, I.

M., Trevanion, S. J., Flicek, P., & Cunningham, F. (2018). Ensembl variation resources.

Database: The Journal of Biological Databases and Curation, *2018*.

https://doi.org/10.1093/database/bay119

IUCN Standards and Petitions Committee. (2019). *Guidelines for Using the IUCN Red List Categories*

and Criteria (No. 14; pp. 1–60). IUCN Red List.

Kardos, M., Armstrong, E. E., Fitzpatrick, S. W., Hauser, S., Hedrick, P. W., Miller, J. M., Tallmon, D.

A., & Funk, W. C. (2021). The crucial role of genome-wide genetic variation in conservation.

Proceedings of the National Academy of Sciences, *118*(48), e2104642118.

https://doi.org/10.1073/pnas.2104642118

Kardos, M., & Shafer, A. B. A. (2018). The Peril of Gene-Targeted Conservation. *Trends in Ecology &*

Evolution, *33*(11), 827–839. https://doi.org/10.1016/j.tree.2018.08.011

Kardos, M., Taylor, H. R., Ellegren, H., Luikart, G., & Allendorf, F. W. (2016). Genomics advances the

study of inbreeding depression in the wild. *Evolutionary Applications*, *9*(10), 1205–1218.

https://doi.org/10.1111/eva.12414

Kauffman, M. J., Meacham, J. E., Hall, S., Steingisser, A. Y., Rudd, W. J., & Ostlind, E. (2018). *Wild*

Migrations: Atlas of Wyoming's Ungulates (Vol. 31). University of Nebraska Press.

https://muse.jhu.edu/article/789690

Kessler, C., Wootton, E., & Shafer, A. B. A. (2022). *Speciation without gene-flow in hybridizing deer*.

https://doi.org/10.1101/2022.04.20.488928

Khan, A., Patel, K., Shukla, H., Viswanathan, A., van der Valk, T., Borthakur, U., Nigam, P., Zachariah,

A., Jhala, Y. V., Kardos, M., & Ramakrishnan, U. (2021). Genomic evidence for inbreeding

depression and purging of deleterious genetic variation in Indian tigers. *Proceedings of the*

National Academy of Sciences, *118*(49), e2023018118. https://doi.org/10.1073/pnas.2023018118

Kim, D., Langmead, B., & Salzberg, S. L. (2015). HISAT: A fast spliced aligner with low memory

requirements. *Nature Methods*, *12*(4), 357–360. https://doi.org/10.1038/nmeth.3317

Kohn, M. H., Murphy, W. J., Ostrander, E. A., & Wayne, R. K. (2006). Genomics and conservation

genetics. *Trends in Ecology & Evolution*, *21*(11), 629–637.

https://doi.org/10.1016/J.TREE.2006.08.001

Korneliussen, T. S., Albrechtsen, A., & Nielsen, R. (2014). ANGSD: Analysis of Next Generation

Sequencing Data. *BMC Bioinformatics 2014 15:1*, *15*(1), 1–13. https://doi.org/10.1186/S12859-

014-0356-4

Kumar, P., Henikoff, S., & Ng, P. C. (2009). Predicting the effects of coding non-synonymous variants on

protein function using the SIFT algorithm. *Nature Protocols*, *4*(7), 1073–1081.

https://doi.org/10.1038/nprot.2009.86

Kumar, S., Stecher, G., Suleski, M., & Hedges, S. B. (2017). TimeTree: A Resource for Timelines,

Timetrees, and Divergence Times. *Molecular Biology and Evolution*, *34*(7), 1812–1819.

https://doi.org/10.1093/MOLBEV/MSX116

Kyriazis, C. C., Wayne, R. K., & Lohmueller, K. E. (2021). Strongly deleterious mutations are a primary

determinant of extinction risk due to inbreeding depression. *Evolution Letters*, *5*(1), 33–47.

https://doi.org/10.1002/evl3.209

Lewontin, R. C. (1974). *The Genetic Basis of Evolutionary Change*. Columbia University Press.

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler transform.

Bioinformatics, *25*(14), 1754–1760. https://doi.org/10.1093/BIOINFORMATICS/BTP324

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., & Durbin,

R. (2009). The Sequence Alignment/Map format and SAMtools. *Bioinformatics (Oxford,*

England), *25*(16), 2078–2079. https://doi.org/10.1093/BIOINFORMATICS/BTP352

Lomsadze, A., Ter-Hovhannisyan, V., Chernoff, Y. O., & Borodovsky, M. (2005). Gene identification in

novel eukaryotic genomes by self-training algorithm. *Nucleic Acids Research*, *33*(20), 6494–

6506. https://doi.org/10.1093/nar/gki937

Martchenko, D., Chikhi, R., & Shafer, A. B. A. (2020). Genome Assembly and Analysis of the North

American Mountain Goat (Oreamnos americanus) Reveals Species-Level Responses to Extreme

Environments. *G3: Genes, Genomes, Genetics*, *10*(2), 437–442.

https://doi.org/10.1534/G3.119.400747

Martchenko, D., & Shafer, A. (2022). *Contrasting whole-genome and reduced representation sequencing*

for population demographic inference: An alpine mammal exampl.

https://doi.org/10.22541/au.164190604.45542461/v1

McDonald, J. S., & Miller, K. V. (2004). *A History of White-tailed Deer Restocking in the United States,*

1878 to 2004. Quality Deer Management Assoc.

McLaren, W., Gil, L., Hunt, S. E., Riat, H. S., Ritchie, G. R. S., Thormann, A., Flicek, P., &

Cunningham, F. (2016). The Ensembl Variant Effect Predictor. *Genome Biology 2016 17:1*,

17(1), 1–14. https://doi.org/10.1186/S13059-016-0974-4

Pascual, U., Adams, W. M., Díaz, S., Lele, S., Mace, G. M., & Turnhout, E. (2021). Biodiversity and the

challenge of pluralism. *Nature Sustainability*, *4*(7), 567–572. https://doi.org/10.1038/s41893-021-

00694-7

Peart, C. R., Tusso, S., Pophaly, S. D., Botero-Castro, F., Wu, C.-C., Aurioles-Gamboa, D., Baird, A. B.,

Bickham, J. W., Forcada, J., Galimberti, F., Gemmell, N. J., Hoffman, J. I., Kovacs, K. M.,

Kunnasranta, M., Lydersen, C., Nyman, T., de Oliveira, L. R., Orr, A. J., Sanvito, S., … Wolf, J.

B. W. (2020). Determinants of genetic variation across eco-evolutionary scales in pinnipeds.

Nature Ecology & Evolution, *4*(8), 1095–1104. https://doi.org/10.1038/s41559-020-1215-5

Petit-Marty, N., Vázquez-Luis, M., & Hendriks, I. E. (2021). Use of the nucleotide diversity in COI

mitochondrial gene as an early diagnostic of conservation status of animal species. *Conservation*

Letters, *14*(1), e12756. https://doi.org/10.1111/conl.12756

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A. R., Bender, D., Maller, J., Sklar, P.,

De Bakker, P. I. W., Daly, M. J., & Sham, P. C. (2007). PLINK: A Tool Set for Whole-Genome

Association and Population-Based Linkage Analyses. *The American Journal of Human Genetics*,

81(3), 559–575. https://doi.org/10.1086/519795

Quinlan, A. R., & Hall, I. M. (2010). BEDTools: A flexible suite of utilities for comparing genomic

features. *Bioinformatics*, *26*(6), 841–842. https://doi.org/10.1093/BIOINFORMATICS/BTQ033

Renaud, G., Hanghøj, K., Korneliussen, T. S., Willerslev, E., & Orlando, L. (2019). Joint Estimates of

Heterozygosity and Runs of Homozygosity for Modern and Ancient Samples. *Genetics*, *212*(3),

587–614. https://doi.org/10.1534/GENETICS.119.302057

Robinson, J. A., Brown, C., Kim, B. Y., Lohmueller, K. E., & Wayne, R. K. (2018). Purging of Strongly

Deleterious Mutations Explains Long-Term Persistence and Absence of Inbreeding Depression in

Island Foxes. *Current Biology*, *28*(21), 3487-3494.e4. https://doi.org/10.1016/j.cub.2018.08.066

Robinson, J. A., Ortega-Del Vecchyo, D., Fan, Z., Kim, B. Y., vonHoldt, B. M., Marsden, C. D.,

Lohmueller, K. E., & Wayne, R. K. (2016). Genomic Flatlining in the Endangered Island Fox.

Current Biology, *26*(9), 1183–1189. https://doi.org/10.1016/j.cub.2016.02.062

Rogers, R. L., & Slatkin, M. (2017). Excess of genomic defects in a woolly mammoth on Wrangel island.

PLOS Genetics, *13*(3), e1006601. https://doi.org/10.1371/journal.pgen.1006601

Schmidt, C., Hoban, S., Hunter, M., Paz-Vinas, I., & Garroway, C. J. (2022). *The IUCN Red List is not*

sufficient to protect genetic diversity. EcoEvoRxiv. https://doi.org/10.32942/osf.io/hn4by

Shafer, A. B. A., Wolf, J. B. W., Alves, P. C., Bergström, L., Bruford, M. W., Brännström, I., Colling, G.,

Dalén, L., Meester, L. D., Ekblom, R., Fawcett, K. D., Fior, S., Hajibabaei, M., Hill, J. A.,

Hoezel, A. R., Höglund, J., Jensen, E. L., Krause, J., Kristensen, T. N., … Zieliński, P. (2015).

Genomics and the challenging translation into conservation practice. *Trends in Ecology &*

Evolution, *30*(2), 78–87. https://doi.org/10.1016/j.tree.2014.11.009

Smeds, L., & Ellegren, H. (2022). *From high masked to high realized genetic load in inbred*

Scandinavian wolves. Authorea. https://doi.org/10.22541/au.165907442.23809998/v1

Stanke, M., Tzvetkova, A., & Morgenstern, B. (2006). AUGUSTUS at EGASP: Using EST, protein and

genomic alignments for improved gene prediction in the human genome. *Genome Biology*, *7*

Suppl 1, S11.1-8. https://doi.org/10.1186/gb-2006-7-s1-s11

Streit, K., Hammacher, C., Zeller, A., & Hack, S. (2012). Sambamba: A Runtime System for Online

Adaptive Parallelization. In M. O'Boyle (Ed.), *Compiler Construction* (pp. 240–243). Springer.

https://doi.org/10.1007/978-3-642-28652-0_13

Tajima, F. (1989). Statistical Method for Testing the Neutral Mutation Hypothesis by DNA

Polymorphism. *Genetics*, *123*(3), 585–595.

Teixeira, J. C., & Huber, C. D. (2021). The inflated significance of neutral genetic diversity in

conservation genetics. *Proceedings of the National Academy of Sciences*, *118*(10), e2015096118.

https://doi.org/10.1073/pnas.2015096118

Väli, Ü., Einarsson, A., Waits, L., & Ellegren, H. (2008). To what extent do microsatellite markers reflect

genome-wide genetic diversity in natural populations? *Molecular Ecology*, *17*(17), 3808–3817.

https://doi.org/10.1111/j.1365-294X.2008.03876.x

Van der Auwera, G. A., Carneiro, M. O., Hartl, C., Poplin, R., del Angel, G., Levy-Moonshine, A.,

Jordan, T., Shakir, K., Roazen, D., Thibault, J., Banks, E., Garimella, K. V., Altshuler, D.,

Gabriel, S., & DePristo, M. A. (2013). From FastQ Data to High-Confidence Variant Calls: The

Genome Analysis Toolkit Best Practices Pipeline. *Current Protocols in Bioinformatics*, *43*(1),

11.10.1-11.10.33. https://doi.org/10.1002/0471250953.bi1110s43

van der Valk, T., Díez-del-Molino, D., Marques-Bonet, T., Guschanski, K., & Dalén, L. (2019).

Historical Genomes Reveal the Genomic Consequences of Recent Population Decline in Eastern

Gorillas. *Current Biology : CB*, *29*(1), 165-170.e6. https://doi.org/10.1016/J.CUB.2018.11.055

Vaser, R., Adusumalli, S., Ngak Leng, S., Sikic, M., & Ng, P. C. (2015). SIFT missense predictions for

genomes. *Nature Protocols*, *11*. https://doi.org/10.1038/nprot.2015.123

von Seth, J., Dussex, N., Díez-del-Molino, D., van der Valk, T., Kutschera, V. E., Kierczak, M., Steiner,

C. C., Liu, S., Gilbert, M. T. P., Sinding, M. H. S., Prost, S., Guschanski, K., Nathan, S. K. S. S.,

Brace, S., Chan, Y. L., Wheat, C. W., Skoglund, P., Ryder, O. A., Goossens, B., … Dalén, L.

(2021). Genomic insights into the conservation status of the world's last remaining Sumatran

rhinoceros populations. *Nature Communications 2021 12:1*, *12*(1), 1–11.

https://doi.org/10.1038/s41467-021-22386-8

Xue, Y., Prado-Martinez, J., Sudmant, P. H., Narasimhan, V., Ayub, Q., Szpak, M., Frandsen, P., Chen,

Y., Yngvadottir, B., Cooper, D. N., de Manuel, M., Hernandez-Rodriguez, J., Lobon, I.,

Siegismund, H. R., Pagani, L., Quail, M. A., Hvilsom, C., Mudakikwa, A., Eichler, E. E., …

Scally, A. (2015). Mountain gorilla genomes reveal the impact of long-term population decline

and inbreeding. *Science*, *348*(6231), 242–245. https://doi.org/10.1126/science.aaa3952

Data Availability and Benefit Sharing

All genome assemblies can be found on GenBank (details in Appendix Table S1). Re-sequencing

data has been submitted to SRA and accession numbers are pending. Bash and R scripts for all

analyses are available on GitLab: https://gitlab.com/WiDGeT_TrentU/undergrad-theses/-

/tree/master/Wootton_2022

Benefits Generated: This research addresses important questions about the genomic health of

populations of priority concern and illustrates relationships between genomic health and long-

term demographic history. Collaborations were formed with researchers across North America,

and all major collaborators are included as co-authors. All contributors are listed in the

Acknowledgements. We are committed to institutional capacity-building and have shared the

results of our research with our collaborators and the broader scientific community.

Author Contributions

ABAS and EW designed the research project. EW conducted all analyses. ABAS and EW wrote

the manuscript using comments and contributions from all coauthors.

639   **Tables and Figures**

640   **Table 1:** The number of scaffolds > 10 Mb, mean scaffold length > 10 Mb, number of genes, and average
641   number of deleterious mutations per species identified by GERP and SIFT. The number of genes was
642   taken from the annotation of each species.

Species	Scaffolds > 10 Mb (n)	Mean Scaffold Length > 10 Mb	Genes (n)	SNPs in CEs (GERP)	SNPs in CEs with RS > 2 (GERP)	Deleterious mutations (SIFT)
Caribou	68	19,723,800	33,242	74	32	4,681
Mountain Goat	49	49,667,287	22,292	212	74	3,317
White-Tailed Deer	86	21,282,945	20,750	411	187	6,502

Figure 1: Sample locations and images of each species. (Red) *Rangifer tarandus* (N=67), (Yellow) *Oreamnos americanus* (N=35), (Blue) *Odocoileus virginianus* (N=53). Current demographic trends are outlined in the top left corner of each image. Boxplots showing Tajima's D and nucleotide diversity (π) per species are also pictured.

Figure 2: (A) Boxplot showing F_{ROH} as calculated by PLINK for all three species. Only ROHs greater than 100 Kb are included in each estimate. No ROHs were detected in caribou. (B) Boxplot showing the method-of-moments inbreeding coefficient (*F*) for each species as calculated by PLINK.

Figure 3: (A) Bar plot showing the total number of ROHs in each species stratified by ROH size category in Kb. No ROHs were detected in caribou. (B) Bar plot showing the total length (Mb) of all ROHs in each size category (Kb). No ROHs were detected in caribou.

Figure 4: Boxplot showing the fraction of constrained positions with SNPs by species. Such SNPs are putatively deleterious, making this the fraction of positions that are deleterious in constrained elements (CEs).

676 **Figure 5:** (A) Bar plot showing the fractions of each coding consequence identified by VEP in each
677 species. (B) Boxplot showing the fraction of genes that possess deleterious mutations identified by SIFT.

684 **Supplemental Information**

685 **Table S1:** GenBank assembly accession numbers for each species used by GERP.

Species	GenBank Assembly Accession
Equus caballus	GCA_002863925.1
Equus asinus	GCA_016077325.2
Tapirus terrestris	GCA_004025025.1
Diceros bicornis	GCA_004027315.2
Ceratotherium simum	GCA_000283155.1
Camelus dromedarius	GCA_000803125.3
Camelus bactrianus	GCA_000767855.1
Hippopotamus amphibius	GCA_004027065.2
Tragulus javanicus	GCA_004024965.2
Moschus moschiferus	GCA_004024705.2
Moschus chryogaster	GCA_006461725.1
Bos taurus	GCA_002263795.2

Connochaetes taurinus	GCA_006408615.1
Ovis canadensis	GCA_004026945.1
Giraffa camelopardalis	GCA_017591445.1
Antilocapra americana	GCA_007570785.1
Alces alces	GCA_007570765.1
Cervus elaphus	GCA_910594005.1
Phacochoerus africanus	GCA_016906955.1
Sus scrofa	GCA_000003025.6
Rangifer tarandus	GCA_019903745.1
Oreamnos americanus	GCA_009758055.1
Odocoileus virginianus	GCA_014726795.1

687 **Table S2:** Correlation matrix between genomic health metrics and translocation data for white-tailed
688 deer.

	F_{ROH}	F	Fraction of deleterious CEs	Fraction of deleterious genes
Stock start	0.032	0.127	0.076	-0.225
Stock end	0.372	0.392	0.205	0.347
Population low	-0.182	-0.059	-0.177	0.308
Number stocked	-0.102	-0.062	0.162	-0.103
Number of states	-0.161	-0.164	-0.093	0.096

Figure S1: Phylogenetic tree generated by TimeTree showing all 23 ungulate species analysed by GERP++ and estimated times of divergence in MYA (Kumar et al., 2017). Study species with sample data are starred in red (*Odocoileus virginianus*, *Rangifer tarandus*, and *Oreamnos americanus*).

Figure S2: (A) Scatterplot and linear regression line showing the correlation between F and F_{ROH} in
mountain goats (Pearson's $r = 0.840$, $p < 0.05$). (B) Scatterplot and linear regression line showing the
correlation between F and F_{ROH} in white-tailed deer (Pearson's $r = 0.395$, $p < 0.05$).

Figure S3: (A) Boxplot showing the number of constrained elements (CEs) identified per species by
GERP++'s gerpelem function. (B) Boxplot showing the number of highly constrained CEs (RS > 2)
identified in each species by gerpelem.

Figure S4: (A) Boxplot showing the average length of constrained elements (CEs) identified per species.
(B) Boxplot showing the average intensity of constraint (RS-score) of CEs identified in each species.

Figure S5: Histogram showing the fractional distribution of constrained element (CE) RS-scores.

734   **Figure S6:** (A) Scatterplot and linear regression line showing the correlation between *F* and latitude in
735   caribou (Pearson's $r = -0.466$, $p < 0.05$). (B) Scatterplot and linear regression line showing the correlation
736   between *F* and latitude in white-tailed deer (Pearson's $r = -0.513$, $p < 0.05$).