Adverse effects of hunting with hounds on participants and bystanders

Adrian Treves[a¥*], Laura Menefee[b†*]

[a] Nelson Institute for Environmental Studies, 30A Science Hall, 550 North Park Street, University

of Wisconsin, Madison, WI, 53706, USA, +1-608-890-1450, atreves@wisc.edu,

[b] Department of Literature, Leeward College, University of Hawai'i, Pearl City, HI and Sierra Club

Wisconsin Chapter, mail c/o A. Treves, 30A Science Hall, 550 North Park Street, University of

Wisconsin, Madison, WI, 53706, USA, menefee.laura@gmail.com

[†] These authors have contributed equally and share first authorship.

Correspondence: atreves@wisc.edu, menefee.laura@gmail.com

Abstract

In the face of biodiversity crises and concerns raised about animal ethics, some societies are re-

examining many human uses of animals. One activity that may once have been unobjectionable

has undergone little scrutiny in most countries: hunting mammals with free-running hounds.

We present two novel datasets about this under-studied `behavioral interaction and hunting

method. In the state of Wisconsin, USA, hounds and gray wolves occasionally interacted

aggressively and human by-standers reported adverse interactions with hounds and their

handlers. Such self-selected samples cannot be used to extrapolate rates in space or time but

do provide new information. Between 1999 and 2012, 176 hounds were reported to have

suffered injury during encounters with wolves. No government data were collected on how

many wolves or other non-target animals were injured by hounds. We describe the grouping

`behavior, patterns of injury and vulnerability to attack, hound phenotypes, and other

ethological data associated with aggressive wolf-hound interactions. We analyze wolf life

history, seasonal life history patterns, bite locations on hound bodies, consumption of hound

carcasses, and the outcomes of interactions to test two hypotheses for wolf-hound

interactions. We find weak or no support for either and propose new hypotheses focusing on

the behavior of hounds and handlers primarily. We also report handler and hound interactions

with human bystanders. Between 2015-2021, bystanders (n=105) reported various adverse

incidents with hounds and handlers, interactions with law enforcement, and other events

associated with hunting with hounds. We propose reforms to policy for wildlife, hunting, and

law enforcement, and reforms to husbandry by hound handlers.

Keywords: human-carnivore conflict and coexistence, dog, illegal take, poaching, wolf

Introduction

In the face of biodiversity crises and concerns raised about animal ethics partly caused

by climate change and partly by human-induced mortality, some societies are re-examining

many human uses of animals that once seemed unobjectionable. For example, uses of poison,

off-road vehicles, fire suppression, etc. have undergone scrutiny for their societal benefit-cost

estimates and their effects on domestic animals, wildlife, and ecosystem health. One human

activity that may once have been unobjectionable has undergone little scrutiny: hunting with

free-running hounds loosed far from their owners.

In our search for work on hunting with dogs or hounds in Google Scholar, the search

phrase 'hunt with (hound or dog)'' yielded 38,300 results declining by half when '-bird' was

added to the search string to exclude bird-hunting with dogs. By contrast, 'hunt -bird 'yielded

3.16 million results. Therefore, hunting with hounds is under-studied. Judging from one court

case in Wisconsin, the practice of hound-hunting is protected from outside scrutiny (Brown v

Kemp2021). Yet, the practice of hunting mammals with hounds has been recorded since 8,000

years at least (Guagnin et al., 2018), praised by President Theodore Roosevelt in 1902

(Roosevelt, 1902), and is legal in numerous countries and several U.S. states (Heberlein, 2000;

Hristienko and McDonald, 2007). Despite its long history, loosing mammal-hunting hounds to

pursue prey, some as large as brown (grizzly) bears, *Ursus arctos*, may have harmful effects on

people in their path, on the hounds themselves, and on target and non-target wildlife they

encounter or pursue (Grignolio et al., 2011; Gompper, 2013). Therefore, we present data on

reports by self-selected owners alleging harm to their hounds, following interactions with

wolves. We also present self-reported perceptions of human by-standers who experienced the

`behavior of hounds during training or hunting. We offer these two data sets to describe the

side-effects of this poorly studied hunting method.

Non-target animals and their interactions with hounds

When pets kill wildlife, biodiversity may diminish or ecosystem health may deteriorate

(Grignolio et al., 2011; Gompper, 2013). Dogs are potential predators of domestic animals and

wild ones (Bowers, 1953; Ciucci and Boitani, 1998). `behavioral changes in wildlife are often

reported. For example, black bears *U. americanus* were reported to avoid encounters with

hounds and in so doing approach people and major roads more frequently (Stillfried et al.,

2015). Some animals will stand their ground when hounds encounter them (Treves et al., 2002;

Wydeven et al., 2004; Backeryd, 2007). Larger size and greater competitive ability of the wild

animals may alter the risk posed by hounds and vice versa. Hounds are often used for pursuit of

mammals larger than individual hounds. When hounds encounter animals that can defend

themselves effectively, the hounds may be injured. Researchers have examined aggressive

encounters between wolves and dogs in many regions (Fritts and Paul, 1989; Backeryd, 2007;

Kaartinen et al., 2009).

The States of Wisconsin and Michigan, USA, have a relatively longer history of such

research. Spatial patterns of wolf *Canis lupus* attacks on hounds are somewhat predictable

(Treves et al., 2002; Wydeven et al., 2004; Bump et al., 2013; Olson et al., 2014). The risk of an

attack appeared to be higher in areas with more public land, larger wolf packs, closer to a wolf

pack, and when baits were left out longer. Here, we examined self-reports by owners on the

characteristics of both canids involved, and self-reported correlates of the outcomes of wolf-

hound interactions (WHI).

At the time of our study, it was illegal for hounds to attack wild animals. Nevertheless,

such attacks might have occurred anyway. We do not have evidence of which animal in a WHI

initiated aggression or escalated it to the point of injury or death. We only present data on the

outcomes for hounds because outcomes for wolves were not documented. Federal law under

the Endangered Species Act and state law against animal fighting would both require

authorities to collect data on the latter. We cannot explain why the database contained no such

information. Therefore, we cannot rule out the possibility that wolves responded defensively to

hound attacks. The evidence for wolf attacks on hounds came from handlers seeking

compensation or other forms of redress (Treves et al., 2002; Ruid et al., 2009; Treves et al.,

2009). In a prior study, a number of wolf deaths caused by other canids were invariably

attributed to other wolves (Treves et al., 2017b). Yet, veterinary pathologists might not be able

to reliably distinguish large dogs, such as hunting hounds, from wolves by simple scrutiny of

bite marks without DNA analysis (Plumer et al., 2018). Therefore, our sample is necessarily

biased toward handler concerns and outcomes for hounds. This does not rule out that wolves

were the aggressors or that wolves remained unharmed by hounds.

We focus on the `behavior of hounds, wolves, and handlers as reported by by-standers.

We test two ethological hypotheses about WHI. First, the predation hypothesis predicts WHI

occur when wolves attack hounds for food, and second, the territoriality hypothesis predicts

that WHI occur when wolves defend territory or pups (Treves et al., 2002; Wydeven et al.,

2004).

Hounds and humans

Much research has measured attitudes of hound-handlers towards wolves (Supplementary

Material 1). In brief, as a group, hound-handlers have the lowest tolerance for wild wolves in

any group measured thus far. There are no published data on hound-handler' encounters

between by-standers and either handlers or their free-running hounds. The Sierra Club

Wisconsin Chapter (SCWC) began to fill the gap with the survey we report here. As part of a

National Sierra Club initiative, the SCWC subcommittee discussed the first-hand reports

members had received of by-stander experience. Because such first-hand reports are likely to

be remarkable, most reports were adverse. The SCWC subcommittee requested information

from local law enforcement in the counties from which reports and also the Wisconsin

Department of Natural Resources (WDNR) for incident report data. They were informed by

both agencies that "no such data were recorded." Thus, SCWC members led an effort at

gathering information more broadly and systematically.

Material and methods

Wolf-hound interactions (WHI) `behavior, grouping, and phenotype

During our study period, hounds were legally used to hunt many mammals, including smaller

carnivores and black bears. Bear hunting in Wisconsin occurred legally from the last week of

August into October, and hound training was legal in July and August (Treves et al., 2009;

Treves et al., 2010). Hound hunters accounted for approximately 40% of the annual take by

bear hunters (Treves et al., 2010; Olson et al., 2014). Typically hounds were loosed from

vehicles and allowed to run far from owners, without control (Wydeven et al., 2004). Hounds

were often fitted with global positioning systems (GPS) or VHF radio-collars, allowing the owner

to follow remotely the movements of hounds and determine when and where a bear had been

treed. Hunters used groups of up to 6 hounds to track and trail prey during training or hunting

(Wydeven et al., 2004).

We examined WHI from 7 August 1999 through 19 January 2012. Case files documented

145 killed and 31 injured hounds deemed by state or federal verifiers as "confirmed" (85%) or

"probable" (15%) WHI, see methods in (Treves et al., 2002). We believe most WHI incidents

were reported because of a compensation program characterized as more generous than other

jurisdictions (Naughton-Treves et al., 2003; Treves et al., 2009). Bump et al. (Bump et al., 2013)

suggest fewer WHI are reported in Michigan's wolf range because hound owners receive no

compensation. Owners with confirmed losses in Wisconsin were eligible to receive up to $2,500

per hound based on the estimated value of the hound. Since the compensation program began

in 1985, nearly $350,000 dollars were paid to hunters to compensate for hounds injured or

killed by wolves. Between 1985 and 2006, payments for hunting hounds comprised 37% of all

compensation (Treves et al., 2002; Naughton-Treves et al., 2003; Ruid et al., 2009; Treves et al.,

2009). WDNR provided compensation for domestic animals injured or killed by wolves,

including hounds (Treves et al., 2002; Ruid et al., 2009; Treves et al., 2009). United States

Department of Agriculture agents assumed responsibility for verifying WHI in 1990, and

conducted most of the investigations used in our analysis of WHI (Willging and Wydeven, 1997).

WHI case files included written reports and forms documenting field investigations,

including necropsy data, photos, veterinary reports, and anecdotal reports handlers. During the

early years of record-keeping, documentation and reporting of depredations lacked uniformity;

thus, some portions of the data were missing, resulting in lower sample sizes for various

analyses. We limited our analysis of WHI to hounds used to hunt bears, bobcats, or coyotes. A

total of 91% of WHI occurred while pursuing these species, which involves different breeds and

use of hounds than for other quarry, such as waterfowl, upland birds, or rabbits. Hounds that

hunt large prey such as bears or coyotes are typically breeds of large size and build, frequently

Plott (18-27 kg), or larger hound breeds (Bluetick, Redbone, or Walker coon-hounds (18-36 kg).

Occasionally, WHI files did not specify the type of prey being pursued. In these cases, if the

breed of hound was one of the five listed above, we assumed the WHI occurred while pursuing

the above three wildlife species. In total, we report on 176 case files. We quantified the

frequency of WHI among breeds of hounds. If the hound was reported as a mix of multiple

breeds, we used the first breed listed. We pooled breeds in an "other" category when a single

breed had too few WHI to meet the assumptions of the *chi*-squared test. No data are available

on breed frequencies or preference by hunters in Wisconsin, with which we could estimate

relative risk by breed.

Our analysis of the body site bitten was limited because a number of hound carcasses

were partially or wholly consumed before retrieval by an owner arriving late at the scene. We

pooled head, neck, and throat into one category and all other sites in another category, to test

if outcomes of WHI differed by bite site.

We compared hound group size, number of hounds involved in the WHI, and wolf pack

size using (1) number of wolves seen and reported by hunters (observed), and (2) WDNR-

reported wolf pack sizes from the winter preceding the WHI (censused). Because wolf packs

exhibit fission-fusion sociality and packs disaggregate often in the summer (Mech, 1970), when

many WHI occurred. Wolf packs in Wisconsin average 4 adults in late winter (range 2-12),

usually composed of offspring of the two (usually) unrelated alpha breeders. By summer, pups

are exiting the dens but range near them or near other pack rendezvous sites until the pack

regularly fuses and travels long distance in mid- to late Fall. During this time, most wolf pack

members return periodically to rendezvous or den sites to assist with pup-rearing, and

consequently have higher food demands, perhaps requiring wolves to forage more frequently

(Ruprecht et al., 2012). From birth until the end of August, wolf pups experience the highest

growth rates, with September representing a critical month for weight gain (Kreeger, 2003). In

some cases, wolf pups have been observed gaining as much as 3.6 pounds per week (Van

Ballenberghe and Mech, 1975). Pup growth, critical to survival, is limited by food quality and

availability. By late August, growth begins to taper (Pulliainen, 1965), as does rendezvous site

use (Van Ballenberghe and Mech, 1975; Kreeger, 2003; Ruprecht et al., 2012). We acknowledge

R. P. Thiel for the suggestion that seasonality of wolf life-history on an annual cycle would

predict more risk of WHI in late summer and autumn because of all pack members and

especially pups' caloric needs. Finally, we analyzed the temporal occurrence of WHI as it relates

to public hunting seasons, and we compared the frequency of WHI during the hound-training

period (July-August) to that during the bear-hunting season (treated as September-October).

Handler `behavior and husbandry

The state wildlife agency implemented several methods for mitigating or preventing

WHI, including compensation for handlers' self-reported losses, encouraging the use of warning

devices on collars to deter wolves, and statewide communication to hunters on recent hound

injuries and their locations, and designation of Wolf Caution Areas (WCAs). The state

communicated the location of higher-risk WCAs online, posted in the field, and in other ways to

handlers (Wydeven et al., 2004; Olson et al., 2014). Within WCAs, the WDNR recommended

that bear hunters release hounds >2 miles from known rendezvous sites. Prior work

documented handlers' willingness to risk hounds in posted WCAs, even within the same season

and even within hours of previous WHI or WCA posting (Wydeven et al., 2004; Olson et al.,

2014). Compensation records also document multiple payments to the same owner or handler

within a single season (Treves et al., 2009). These data suggest not all hunters heeded the

state's warnings. Compensation programs did not clearly reduce WHI rates. Much scholarly

discussion has focused on the moral hazard (in short, negligent owners have no incentive to

protect their animals if they will be paid for losses). Prior research on Wisconsin's

compensation program addressed moral hazards (Naughton-Treves et al., 2003; Treves et al.,

2009). The state only changed the source of the funds, not the conditions for payment since

that work was published. Therefore, cooperation of the state legislature may be essential to

changing an incentive for WHI to occur into a disincentive for hounds and non-target animals to

be put at risk. Handlers may be able to prevent WHI by using protective vests or stronger collars

(Backeryd, 2007; Khorozyan et al., 2020), keeping hounds leashed until the targeted game

species is located, or bringing first aid kits on the hunt, although the possible effect of these

interventions has not been studied in Wisconsin.

Survey of by-standers

The Sierra Club Wildlife Committee (SCWC) subcommittee on Protecting Native Forests and

Wildlife collected systematic information from by-standers' experiences of hound hunting. The

SCWC led and posted the survey on the Sierra Club Wisconsin Chapter website through 2015-

2021, and administered the survey. The survey appeared at

https://www.sierraclub.org/wisconsin/protecting-native-forests-wildlife accessed 20 August

2022. SCWC members also printed hard copies of the instrument and distributed these at wolf

and wildlife related meetings and conferences in Wisconsin in 2015 and 2016. SCWC also

invited any members of the public who described adverse hounding encounters to fill the

online report. Therefore, respondents were self-selected with attendant biases.

We analyzed anonymized data stripped of identifying information by the SCWC

administrators. The 25-question survey (Supplementary Material Appendix 1) is organized in

four sections: Observations; Trespass; Property Damage, Personal Injury or Threats; and

Interactions with Law Enforcement, totaling 22 yes/no questions and 4 items that allowed

unstructured responses to elaborate on answers. Respondents could identify county of

residence and interaction, if different. Respondents were asked how many hounds they saw

during each interaction they reported. When two respondents mentioned the same interaction

but different numbers of hounds, LM averaged and rounded up for the number of hounds.

About 80% of respondents used the online form to report anonymously, and 20% sent

their responses directly to SCWC via mail, phone, email, or in person while being assured of

anonymity. LM screened the sample to eliminate responses which identified no adverse

incident involving hunting hounds, as these respondents generally used the report format to

express an opinion about the practice of hunting with hounds, which we did not analyze

because it was outside the purview of this survey (Supplementary Material 2). We screened for

multiple reports of the same incident with identifying factors such as location and date. After

the screening, the sample presented here appears to come from independent incidents

although we had no way to verify location or date.

Statistical analyses

We performed statistical analyses (SAS Institute, 2013) using Student's paired t tests to

compare the differences in average estimated ages of hounds and numbers of wolves during

the attack, in relation to the outcome of the WHI (i.e., killed or injured) after evaluating if

variances were equivalent (F test). All statements of statistical significance are based on $P \leq$

0.05. We used Spearman rank correlations to detect associations between multiple continuous

variables.

Ethical note

The data presented are exempt from approvals for research on human subjects because

(a) the data were collected by another organization (WDNR and SCWC) and distributed to the

authors without individually identifiable information; (b) all SCWC survey responses were

completely anonymous; and (c) the identity of complainant handlers was likewise unknown to

us and redacted from records by the state agency that received the complaints. Use of hounds

for hunting is a state-regulated activity. No research was conducted on the hounds.

Results and Discussion (as allowed by instructions to authors, p.9 para 3)

243   Wolf-hound interactions (WHI) `behavior, grouping, and phenotype

244   In 176 case files, we found 140 independent WHI cases during our study period, where a

case reported on the same day and location by different owners were pooled into one WHI.

Records included 145 killed (83%) and 31 injured (17%) hounds, similar to 71% and 82%

reported in Nordic countries (Kojola and Kuittinen, 2002; Backeryd, 2007). The high percentage

of fatalities in both cases might reflect that handlers sometimes took hours to find a distant

hound. Therefore, sub-lethal injuries might not be attributed to a WHI if handlers and hounds

reunited long after it ended; this effect would bias WHI records upward to more fatalities.

Similarly, the cases involving distant handlers might not have escalated to lethal injuries if the

handlers had arrived earlier.

253   Wolf injuries and deaths in WHI were not documented nor reported in case files.

Another possible source of missing data might come from misidentifying wolves as the wild

interactant, given the availability of compensation for such WHI and not for casualties from

other wildlife (Treves et al., 2009). In total, 89% of WHI occurred while hunters reported

pursuing black bears, bobcats *Lynx rufus* 6%, coyotes *C. latrans* 4%, raccoons *Procyon lotor* 1%).

However, we lack independent data on the animal being pursued by those hounds at the time

of WHI. We also lack the relative frequencies of targeting each species with hounds statewide

and over time. There was no association between the outcome of WHI categorized as either

injury or death and the prey being pursued by hunters (X^2 = 1.9, P = 0.75, df = 4, n = 140). The

bear-hound-training period accounted for 62% of WHI, whereas the bear-hunting season

accounted for 28% despite being the same length approximately. Outcomes were not

associated with month annually (X^2 = 8.5, P = 0.38, df = 8, n = 176, Table 1). These two results

suggest that seasonality and Wolf reproductive timing did not predict injury or death of hounds,

which undermine both the predation and territoriality hypotheses. We lack information on the

frequency with which WHI was initiated by hounds, which could result in very different causal

hypotheses focused on the motivations of hounds and handlers.

Neither sex nor age of the hounds was associated with the outcome of WHI (sex X^2 =

1.32, P = 0.25, df = 1, n = 151; age t = -0.71, P = 0.49; variances were equal F = 0.49). The

Treeing Walker Coonhound was the most common breed in WHI (33.3%, n = 51), followed by

the Plott (27.5%, n = 42). There was a significant association between breed and outcome (X^2 =

10.7, P = 0.03, df = 4, n = 176). Notably, the Plott fatality frequency of 95.4% was higher than

the average 81.2% (Table 1). Plott hounds are the smallest and most vocal hound breed

commonly used in Wisconsin for mammals. Small body size has been implicated in the risks and

fatalities associated with WHI (Fritts and Paul, 1989; Peterson, 1995; Backeryd, 2007; Kaartinen

et al., 2009). The Plott breed is also known for its baying vocalizations, which might alert wolves

from a long distance. Similar elevated risks of dog vocalizations were reported in Nordic

countries (Backeryd, 2007; Kaartinen et al., 2009). Small size may make a hound more

vulnerable to head and neck bites. Bites to the neck were associated with higher fatality rates in

a Scandinavian study (Backeryd, 2007).

Our analysis on hound body site bitten was limited to 109 WHI. We cannot be certain

that wolves inflicted every bite because of the above-mentioned delays in discovering hounds

and the missing center created by compensation only for attacks by wolves. Of the 109

carcasses with bite information, 50 provided one bite location (46%), 37 provided two locations

(34%), and 22 provided 3 or more locations (20%). Considering all bite locations (n = 193), the

single most frequent bite site was the neck (33%), followed by back (17%), upper thigh (12%),

and chest (10%). We considered bites to the head, shoulders, neck (as opposed to throat), back,

and upper thighs as indicative the hound had been lower than its attacker. Those upper body

parts were represented in 72% of the 193 bites whereas under-parts (throat, groin, sternum,

ribs, lower legs, abdomen) were represented in 28% of bite locations. We found no relationship

between body site bitten and outcome, when we separated neck and head bites from others

(X^2 = 1.5, P = 0.22, df = 1, n = 66).

Another factor affecting the vulnerability of canids to attack by other canids is numerical

superiority. An average of 1.3 hounds were injured or killed per WHI (maximum 5 in a single

WHI). The size of the hound group averaged 3.8 SD 1.4, n = 57; only 3 (5%) of those WHI involve

a single hound in the handler's control; and also the number involved in the WHI (2.6 SD 1.3, n

= 47 with 9 (19%) of those WHI involving only 1 hound, were both similar to the number of

wolves observed by handlers (2.9 SD 1.2, n = 15), the census pack size for the pack blamed by

the state or federal agent tasked with verifying the report (2.4 SD 1.0, n = 19; n = 4 included

information for both observed and censused). The outcomes were not associated with the

number of hounds, number of wolves, or difference between the two in a given WHI by any of

the measures of group size or pack size above (Welch test assumes unequal variance, F < 0.72,

P > 0.41 in every test). Wolves injured or killed hounds in groups with superior numbers in 44%

of WHI with such data (n = 16). Aggression between wolves and coyotes in Yellowstone

National Park had fatal consequences when wolves outnumbered the smaller coyotes, but not

when coyotes outnumbered wolves, suggesting that group size exerted less influence than

individual body size differences in determining outcomes between canids (Ballard et al., 2003;

Merkle et al., 2009). In our study, if wolves were motivated by predation or territorial defense

as suggested by the two hypotheses, we might expect pack size to predict the outcomes of

WHI. Again, this suggests the need for new hypotheses as above.

Of 80 deaths with data on consumption of a carcass, 49% of hound carcasses were

partially consumed. Of those 80 hounds consumed by wolves, 71% occurred July–August and

27% in September–October. When we combine the former timing of WHI, site location of bites,

and the consumption data above, we can test two hypotheses about the cause of WHI. The

timing of WHI presents equivocal evidence for both hypotheses. Higher frequencies of WHI

occurred during the hound training period in July and August than during the autumn black

bear hunt in September and October (Table 1). Elevated risk in July and August might have been

associated with the practice of baiting, as wolves visit bear bait sites in search of food (Bump et

al., 2013). In Wisconsin, bear bait sites could be legally established as early as April and could

last the entire wolf pup-rearing season. Bump et al. (2013) documented that the risk of WHI

was three to seven times greater in Wisconsin than in adjacent Michigan, citing the extended

bear-baiting period as a probable cause for the much higher risk of WHI in Wisconsin. That

might support the predation hypothesis. However, bear baiting was confounded with wolf pup

defense predicted by the territoriality hypothesis. The hound training period coincided during

the study with wolf use of rendezvous sites or den sites. See the introduction for the life-history

on an annual cycle of wolves, which the majority of hound packs would not experience in the

same way because of human provisioning, aseasonal reproduction, and more human

intervention in breeding `behavior. The consumption of hound carcasses might corroborate the

predation hypothesis, but that is not persuasive because consumption was recorded in only

approximately half of the WHI and we do not know if the wolves that attacked were the

consumers. Nor can we rule out that consumption followed after the primary motivation for

aggression in either the hounds or the wolves. The hound carcasses and bite locations provided

limited insight. Bites to head, neck, and throat represented 41% of bite locations on hound

carcasses. The predation hypothesis might find support from this result because cranio-cervical

killing bites are associated with predation by many mammals (Steklis and King, 1978; Van

Valkenburgh and Ruff., 1987). But the greater height of wolves off the ground then off most

towns would predict such bite locations in any case. In sum, we find equivocal support for both

hypotheses. This could imply both are correct or we are missing information, such as whether

the hounds were pursuing Wolves. We also do not have information on the body condition of

any of the participants, which seems essential data to test the prediction hypothesis.

Handler `behavior and husbandry

Husbandry, such as avoidance of rendezvous sites and use of bells on collars were

difficult to evaluate, because of a lack of data on prevalence, use, and effectiveness of these

potentially preventive methods. Overall, 55 (31%) records reported whether hounds in WHI

wore bells on their collars, with 20 (36%) wearing them and 35 did not (64%). But we have no

data on the use of bells among hounds that did not enter the WHI database and the majority of

records did not contain any information on devices. Outcomes were not associated with hunter

self-reports of affixing bells to collars. Another step handlers might take to protect hounds and

wolves would be to release hounds only in low-risk areas. The state did not systematically

collect data on whether air warnings about high-risk WCAs were heeded. The WHI records did

not contain such information.

Survey of by-standers

In all, 105 respondents reported adverse incidents with hunting hounds from 51

Wisconsin counties, 4 Michigan counties, and 5 counties from other states. The average

number of incidents per county was 2. Seven respondents declined to specify location, but

timing distinguished the reports from other respondents' reports. The 105 respondents

reported 119 separate incidents (Table 2). Of the 105, 42% reported the hounds observed were

not accompanied by a handler; similarly, 41% reported finding abandoned or lost hounds on

their property. In those cases, some respondents reported contacting local animal shelters, law

enforcement or handlers via phone numbers on collars. Overall, 63% of 105 respondents

described incidents of trespass including hounds running on their property without permission,

handlers found on property without permission, or running hounds on property after being

denied permission. Beyond trespass, 18% of respondents described illegal or nuisance

`behavior: damage to property caused by hounds, including downed fencing, damaged

landscaping and gardens, injury to self and livestock, dead wildlife left on property, vandalism

or litter. Most seriously, 11% reported injury to pets or livestock by hounds, although 24%

reported knowledge of hounds attacking others 'pets or livestock. Of the 105, 8% describe

direct encounters with hounds resulting in personal injury or being chased.

Moreover, 31% reported threatening altercations with hound handlers, including being

unwillingly detained by hound handlers 'trucks on public roads, or their own private driveways.

Of 105 respondents, 51% of respondents reported they "feel intimidated by hound handlers,"

and 44% feared retaliation from handlers for reporting confrontations to law enforcement.

Finally, respondents describe being given false information by law enforcement officers,

including local sheriffs and game wardens, and also reported filing official complaints upon

which no discernible action was taken. Overall, 36% of 105 respondents believed a conflict of

interest existed for law enforcement officers, including game wardens, because of relationships

between officers and handlers, or because the officers were believed to hunt with hounds

themselves. Prior research studies report that hound handlers as a group contain a substantial

contingent willing to break the law and flout regulations intended to protect animals and more

so than other groups active in wolf range (summarized in SM 1). Actual poaching evidence is

consistent with the above survey reports (see Conclusions)

Comparing numbers of hounds from WHI and survey data

Survey respondents reported 2–8 hounds per interaction (average 3.7, mode 2). That

average is identical to the average number of hounds that handlers reported in their pack in

WHI above. This seems to be corroborating evidence of accuracy in reports of hound pack sizes

in both datasets, as neither set of complainants was aware of the other. Given the rarity of

single hounds (5%) in WHI, the bystander reports of >1 hound seem unsurprising. Similarly,

bystanders reported >6 hounds in 3 events (8% of reports that include these data) but handlers

never reported >6 in their pack after a WHI. The legal limit per handler was multiple handlers

may release more than 6 hounds. The topic of comparing self-reports by handlers and self-

reports by bystanders raises an issue of expertise and deference to it. A common assumption in

peer-reviewed commentary on hunting is that hunters have lay knowledge gained through local

experience and expertise in their practices. This often plays out among academics and

manager-authors as favoring the views and preferences of local communities and of hunters

over others, especially outside experts (von Essen et al., 2015; Redpath et al., 2017; Richardson,

2022). But that assumption is unjustified when two local, lay types of expertise are pitted

against each other as in an unknown proportion of our data when hound handlers and the

bystanders complaining about them were equally local and both held lay expertise. The second

flaw is when the lay bystanders complained to law enforcement experts about hounds or

handlers, then the power asymmetry is reversed from the usual academic debate. In short, the

local, lay expert complainant deserves the attention and compassion typically reserved for local

hunters. Therefore, we call for more just and fair consideration of the under-represented and

marginalized in wildlife management.

Conclusion

Hunting mammals with hounds has rarely been investigated. Here we fill the gap partly

with self-reported data on the `behavior of hounds and wolves associated with aggressive wolf-

hound interactions (WHI). Also, we present self-reported data on the husbandry of handlers

and the experiences of bystanders exposed to hound and handler `behavior. We report a

shortage of data on wolf casualties. Our inability to find strong support for either the predation

hypothesis or the territoriality hypothesis, which focus on wolf motivation for WHI, suggests

either a multi-causal phenomenon or we lack hypotheses for the motivation of hounds to

initiate WHI. The lack of a seasonal pattern in our data argues against the notion that wolves

are the primary aggressors; WHI may be a mutual affair or may be largely initiated by hounds.

Treeing Walker coonhound was the breed most commonly involved in WHI. Most WHI

were fatal for hounds. Plott hounds were disproportionally represented among fatalities. We

could not confirm or reject a protective role of special collars with bells or defensive features.

Nevertheless, we recommend the state obligate veterinary clinics that treat hounds for wildlife

injuries to report each such incident so the welfare of hounds and preventive actions taken by

handlers can be evaluated by professional veterinarians and wildlife scientists. In light of recent

findings that studded leather collars protect cattle against leopards (Khorozyan et al., 2020),

recommending more precautions by handlers seems prudent. A majority of WHI involved

hounds pursuing black bears compared to other prey but none could confirm what animal the

hounds were pursuing when WHI began. Outcomes of WHI (injury or death of hounds) were

not associated with the number of wolves observed or censused near the site, or the numerical

differences between wolves and hounds, hound age, hound sex, the species of prey targeted by

hunters, or the month in which WHI occurred. Regarding numerical superiority, perhaps both

the large group sizes of hounds (up to 6 per owner) in Wisconsin and uncertainty about the

number of wolves involved, has obscured associations with the fatality rates during WHI.

We report bystanders perceptions of unlawful actions involving trespass, property

damage, attacks, threats, intimidation and harassment of by-standers or their domestic

animals, perpetrated by both hounds and handlers. The survey data and WHI self-reports by

handlers are self-selected samples, which cannot be verify directly or used to extrapolate rates,

frequencies, or representativeness in space, time, or demography. Nevertheless, the data are

the first of their kind to our knowledge and deserve consideration. If even a small proportion

are accurate, the data support improved husbandry of hounds, more transparent law

enforcement, and wildlife protections against hounds. We found little financial or non-financial

incentive for the self-reports other than bystander dissatisfaction with law enforcement

response or value-based disapproval of hounding. By contrast, our other self-selected data set

(handler complaints of hound losses in WHI) were motivated by a compensation program that

paid for injured or dead hounds. That program also seems to need reform given that no

information on harm to wolves was collected despite legal protections for wolves. It might be

impossible to verify that the handlers or hounds were acting lawfully at the time of the WHI.

Furthermore, the size of packs of hounds (up to 6) and the possibility that multiple handlers can

run multiple packs in the same area simultaneously creates a clear danger for wolves and other

non-target animals.

These results highlight a need for improved regulation, greater oversight and more

energetic enforcement of activities involving the use of hounds during hunting. Our results are

also consistent with earlier findings of more serious wildlife crimes (SM 1). Wolf-poaching is the

major cause of mortality in US wolf populations (Treves et al., 2017a). Much of the poaching

involves concealment or destruction of evidence, which reflects intent to break the law, has

repeatedly risen in incidence along with policies that permit some legal wolf-killing in several US

wolf populations (SM 1). These findings indicate that would-be poachers profit from

governmental laxity to act unlawfully or that would-be poachers use the cover of legal hunting

to act unlawfully (Chapron and Treves, 2016a; b; Santiago-Ávila et al., 2020; Louchouarn et al.,

2021; Santiago-Ávila and Treves, 2022). Most recently, data reveal that cryptic poaching rose

during hound training, bear-hunting seasons, and deer-hunting seasons (Santiago-Ávila and

Treves, 2022). The 2021 Wisconsin wolf-hunt saw the most rapid season closure and over-kill in

Wisconsin wolf management history with 218 wolves killed in <72 hours with >80% being killed

by hunters using hounds (Treves et al., 2021). Unpublished necropsy data collected by the

Great Lakes Indian Fish & Wildlife Commission indicate hounds delivered potentially lethal bites

during those hunting incidents, which represent unlawful take by hunters (Treves et al., 2021).

Taken altogether, these data suggest law enforcement is needed.

In USA wildlife law, the Endangered Species Act (ESA) and federal court cases

surrounding it make clear that some hound handlers are vulnerable to prosecution. First, any

"take" of wolves (including harassment, pursuit, injury, killing, etc.) is prohibited under the ESA

regardless of whether the perpetrator knew the wild animal harmed was listed (Newcomer et

al., 2011). The absence of data revealed by this study indicates that hounding is not adequately

regulated. Because wolves were often a federal- or state-listed species during our study, the

practice of hound-hunting in wolf pack territories should be prohibited when wolves are a listed

species unless it can be regulated to reduce WHI near zero. Prohibitions on non-selective killing

methods in the range of endangered species and prohibitions on hunting non-listed species of

similar appearance such as coyotes (Newsome et al., 2015), might resolve some WHI. Potential

legal jeopardy does not stop at the hound handlers. Given the state wildlife agency and law

enforcement refused the SCWC request for information or had no such documents (see

Introduction), the alleged infractions by handlers may still be within the statute of limitations.

Systematic and repeated failure to investigate or prosecute complaints of illegal activity can

make a department vulnerable to lawsuits and the imposition of oversight by higher authorities

(e.g., federal consent decrees). Legal jeopardy arises for the agency because the doctrine of

prosecutorial discretion may not protect a law enforcement agency from charges of systematic

neglect of unlawful activities (Newcomer et al., 2011) and see allegations of such systematic

neglect in a case involving wolves (Wild Earth Guardians v USA2015).

Acknowledgments: R. P. Thiel and C. Williamson shared data and analyses, E. Huntley helped

design the survey of by-standers, Heathcliff Wright designed graphics, and Sierra Club

Wisconsin Chapter administered the online survey and shared data with LM.

Declaration of competing interests: AT and LM perceive no potential competing interests. All

funding awarded to AT as of 13 April 2022 are listed here

http://faculty.nelson.wisc.edu/treves/archive_BAS/funding.pdf accessed 21 August 2022, which

we provide for transparency and a CV for disclosure of potential competing interests:

http://faculty.nelson.wisc.edu/treves/archive_BAS/Treves_vita_latest.pdf accessed 22 August

2022. LM is a retired Associate Professor from the Department of Literature, Leeward College,

University of Hawaii and currently a member of: Endangered Species Coalition, Society for the

Preservation of Poultry Antiquities, National Wolfwatchers Coalition, Wisconsin Wolf Trackers,

Wolf Patrol, Executive Committee, Fox Valley Group, Sierra Club.

495   References and Citations

2015. WildEarth Guardians, et al. v U.S. Department of Justice, U.S. District Court Arizona.

2021. Brown, J. et al. v Kemp, J. et al., U.S. Court of Appeals, 7th circuit.

Backeryd, J., 2007. Wolf attacks on dogs in Scandinavia 1995-2005, Ecology Institute, Swedish

University of Agricultural Sciences, Grimso.

Ballard, W.B., Carbyn, L.N., Smith, D.W., 2003. Wolf interactions with non-prey, in: Mech, L.,

Boitani, L. (Eds.), Wolves: behavior, ecology, and conservation, University of Chicago Press,

Chicago, pp. 259-271.

Bowers, R.R., 1953. The free-running dog menace. Virginia Wildlife 14, 5-7.

Bump, J.K., Murawski, C.M., Kartano, L.M., Beyer, D.E., Roell, B.J., 2013. Bear-Baiting May

Exacerbate Wolf-Hunting Dog Conflict. PLoS ONE 10.1371/journal.pone.0061708.

Chapron, G., Treves, A., 2016a. Blood does not buy goodwill: allowing culling increases poaching

of a large carnivore. Proceedings of the Royal Society B 283, 20152939.

Chapron, G., Treves, A., 2016b. Correction to 'Blood does not buy goodwill: allowing culling

increases poaching of a large carnivore'. Proceedings of the Royal Society B Volume 283,

20162577.

Ciucci, P., Boitani, L., 1998. Wolf and dog depredation on livestock in central Italy. Wildl. Soc.

Bull. 26, 504-514.

Fritts, S.H., Paul, W.J., 1989. Interactions of wolves and dogs in Minnesota. Wildl. Soc. Bull. 17,

121-123.

Gompper, M.E., 2013. Free-ranging dogs and wildlife conservation. Oxford University Press.,

Oxford.

Grignolio, S., Merli, E., Bongi, P., Ciuti, S., Apollonio, M., 2011. Effects of hunting with hounds on

a non-target species living on the edge of a protected area. Biol. Conserv. 144, 641-649.

10.1016/j.biocon.2010.10.022G

Guagnin, M., Perri, A.R., Petraglia, M.D., 2018. Pre-Neolithic evidence for dog-assisted hunting

strategies in Arabia. Journal of Anthropological Archaeology 49, 225-236.

Heberlein, T.A., 2000. The gun, the dog and the thermos: Culture and hunting in Sweden and

the United States. Sweden & America Fall 2000, 24-29.

Hristienko, H., McDonald, J.E.J., 2007. Going into the 21st century: a perspective on trends and

controversies in the management of the American black bear Ursus 18, 72-88.

Kaartinen, S., Luoto, M., Kojola, I., 2009. Carnivore-livestock conflicts: determinants of wolf

(Canis lupus) depredation on sheep farms in Finland. Biodivers. Conserv. 18, 3503-3517.

Khorozyan, I., Siavash, G., Mobin, S., Soofi, M., Waltert, M., 2020. Studded leather collars are

very effective in protecting cattle from leopard (Panthera pardus) attacks. Ecological S;lutions

and Evidence 1, e12013. 10.1002/2688-8319.12013

Kojola, I., Kuittinen, J., 2002. Wolf Attacks on Dogs in Finland. Wildl. Soc. Bull. 30, 498-501.

Kreeger, T., 2003. The internal wolf: physiology, pathology, and pharmacology., in: Mech LD,

B.L. (Ed.), Wolves: behavior, ecology, and conservation, University of Chicago Press, Chicago,

pp. 192-217.

Louchouarn, N.X., Santiago-Ávila, F.J., Parsons, D.R., Treves, A., 2021. Evaluating how lethal

management affects poaching of Mexican wolves Open Science 8 (registered report), 200330.

Mech, L.D., 1970. The Wolf: The Ecology and Behavior of an Endangered Species. University of

Minnesota Press, Minneapolis.

Merkle, J., Stahler, D., DW, S., 2009. Interference competition between gray wolves and coyotes

in Yellowstone National Park. Canadian Journal of Zoology 87, 56-63.

Naughton-Treves, L., Grossberg, R., Treves, A., 2003. Paying for tolerance: The impact of

livestock depredation and compensation payments on rural citizens' attitudes toward wolves.

Conserv. Biol. 17, 1500-1511.

Newcomer, E., Palladini, M., Jones, L., 2011. The Endangered Species Act v. the United States

Department of Justice: How the Department of Justice derailed criminal prosecutions under the

Endangered Species Act. Animal Law 17, 241-271.

Newsome, T., Bruskotter, J.T., Ripple, W.J., 2015. When shooting a coyote kills a wolf: Mistaken

identity or misguided management? Biodivers. Conserv. 24, 3145-3149.

Olson, E.R., Treves, A., Wydeven, A.P., Ventura, S., 2014. Landscape predictors of wolf attacks

on bear-hunting dogs in Wisconsin, USA. Wildl. Res. 41, 584–597.

Peterson, R., 1995. Wolves as interspecific competitors in canid ecology, in: Carbyn, L., Fritts, S.,

Seip, D. (Eds.), Ecology and conservation of wolves in a changing world., Canadian Circumpolar

Institute., Edmonton, Alberta, Canada, pp. 315-323.

Plumer, L., Talvi, T.n., Männil, P., Saarma, U., 2018. Assessing the roles of wolves and dogs in

livestock predation and suggestions for mitigating human-wildlife conflict and conservation of

wolves. Conserv. Genet. https://doi.org/10.1007/s10592-017-1045-4.

Pulliainen, E., 1965. Studies on the wolf (Canis lupus) in Finland. Ann. Zool. Fenn. 2, 215-259.

Redpath, S., Linnell, J.D.C., Festa-Bianchet, M., Boitani, L., Bunnefeld, N., Dickman, A., Gutiérrez,

R., Irvine, J., Johansson, M., Majić, A., Mcmanus, B., Pooley, S., Sandström, C., Sjölander-

Lindqvist, A., Skogen, K., Swenson, J.E., Trouwhborst, A., Young, J., Milner-Gulland, E.J., 2017.

Don't forget to look down – collaborative approaches to predator conservation. Biological

Reviews 92, 2157–2163.

Richardson, J.E., 2022. The Cows May Safely Graze: Placing Expert-Lay Relationships at the

Center of Overcoming the Expert-Lay Knowledge Divide. Rural Sociology in press.

10.1111/ruso.12426

Roosevelt, T.D., 1902. Hunting the Grisly and Other Sketches. G. P. Putnam's Sons, New York.

Ruid, D.B., Paul, W.J., Roell, B.J., Wideven, A.P., Willging, R.C., Jurewicz, R.L., Lonsway, D.H.,

2009. Wolf–human conflicts and management in Minnesota, Wisconsin, and Michigan, in:

Wydeven, A.P., Van Deelen, T.R., Heske, E.J. (Eds.), Recovery of Gray Wolves in the Great Lakes

Region of the United States: An Endangered Species Success Story, Springer, New York, pp. 279-

295.

Ruprecht, J., Ausband, D., Mitchell, M., Garton, M., Zager, P., 2012. Homesite attendance based

on sex, breeding status, and number of helpers in gray wolf packs. J. Mammal. 93, 1001-1005.

Santiago-Ávila, F.J., Chappell, R.J., Treves, A., 2020. Liberalizing the killing of endangered wolves

was associated with more disappearances of collared individuals in Wisconsin, USA. Scientific

Reports 10, 13881. /10.1038

Santiago-Ávila, F.J., Treves, A., 2022. Poaching of protected wolves fluctuated seasonally and

with non-wolf hunting. Scientific Reports 12, e1738 10.1038/s41598-022-05679-w

SAS Institute, I., 2013. JMP 11.0, Cary, North Carolina. .

Steklis, H.D., King, G.E., 1978. The craniocervical killing bite: Toward an ethology of primate

predatory behavior. J. Hum. Evol. 7, 567-581.

Stillfried, M., Belant, J., Svoboda, N., Beyer, D., Kramer-Schadt, S., 2015. When top predators

become prey: Black bears alter movement behaviour in response to hunting pressure. Behav.

Processes 120, 30-39. 0.1016/j.beproc.2015.08.003

Treves, A., Artelle, K.A., Darimont, C.T., Parsons, D.R., 2017a. Mismeasured mortality: correcting

estimates of wolf poaching in the United States. J. Mammal. 98, 1256–1264.

10.1093/jmammal/gyx052

Treves, A., Jurewicz, R.L., Naughton-Treves, L., Rose, R.A., Willging, R.C., Wydeven, A.P., 2002.

Wolf depredation on domestic animals: control and compensation in Wisconsin, 1976-2000.

Wildl. Soc. Bull. 30, 231-241.

Treves, A., Jurewicz, R.L., Naughton-Treves, L., Wilcove, D., 2009. The price of tolerance: Wolf

damage payments after recovery. Biodivers. Conserv. 18, 4003–4021.

Treves, A., Kapp, K.J., Macfarland, D.M., 2010. American black bear nuisance complaints and

hunter take. Ursus 21, 30–42.

Treves, A., Langenberg, J.A., López-Bao, J.V., Rabenhorst, M.F., 2017b. Gray wolf mortality

patterns in Wisconsin from 1979 to 2012. J. Mammal. 98, 17-32. 10.1093/jmammal/gyw145

Treves, A., Santiago-Ávila, F.J., Putrevu, K., 2021. Quantifying the effects of delisting wolves

after the first state began lethal management. PeerJ 9, e11666. 10.7717/peerj.11666

Van Ballenberghe, V., Mech, L., 1975. Weights, growth, and survival of timber wolf pups in

Minnesota. J. Mammal. 56, 44-63.

Van Valkenburgh, B., Ruff., C.B., 1987. Canine tooth strength and killing behaviour in large

carnivores. J. Zool. 212, 379-397.

von Essen, E., Hansen, H.P., Kallstrom, H.N., Peterson, M.N., Peterson, T.R., 2015. The

radicalisation of rural resistance: How hunting counterpublics in the Nordic countries contribute

to illegal hunting. Journal of Rural Studies 39, 199-209.

Willging, R., Wydeven, A.P., 1997. Cooperative wolf depredation management in Wisconsin, in:

Lee, C.D., Hygnstrom, S.E. (Eds.), Thirteenth Great Plains Wildlife Damage Control Workshop

Proceedings., Kansas State University Agricultural Experiment Station and Cooperative

Extension Service, Lincoln, NE, pp. 46-51.

Wydeven, A.P., Treves, A., Brost, B., Wiedenhoeft, J.E., 2004. Characteristics of wolf packs in

Wisconsin: Identification of traits influencing depredation, in: Fascione, N., Delach, A., Smith,

M.E. (Eds.), People and Predators: From Conflict to Coexistence, Island Press, Washington, D. C.,

pp. 28-50.

Table 1. The number of hounds reported in wolf-hound interactions by breed and month

Breed	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Bluetick	0	1	0	0	0	0	3	6	8	0	0	1
Plott	0	0	0	0	0	0	11	20	9	3	0	0
Redbone	0	1	0	0	0	0	4	4	1	0	0	1
Walker	0	0	0	1	0	0	16	19	16	3	1	5
Other	1	2	0	0	0	0	12	15	8	0	0	4
Totals	**1**	**4**	**0**	**1**	**0**	**0**	**46**	**64**	**42**	**6**	**1**	**11**

Table 2. Bystander reports (n=105) of 119 adverse interactions with hounds or their handlers

across counties of Wisconsin, USA. Because a respondent might report more than one

interaction, we present the counties from most reports 6) to fewest (1), the names of the 51

counties mentioned in reports, the sum of interactions per row, and the maximum number of

hounds reported in a single interaction. When multiple reports were filed about the same

interaction and the number of hounds differed, we counted only one interaction and averaged

the number of hounds, rounding to the higher integer.

Reports per county	Sum of the interactions	Maximum number of hounds in a single interaction	Counties with interactions reported
6	15	8	Bayfield, Iron, Sawyer
5	5	5	Forest
4	4	6	Langlade
3	13	6	Chippewa, Dane, Marathon, Polk, Washburn
2	18	6	Dodge, Douglas, Dunn, Florence, Lincoln, Oconto, Price, Shawano

| 1 | 62 | 8 | Ashland, Barron, Brown, Burnett, Calumet, Cheboygan MI, Columbia, Cuyahoga OH, Door, Eau Claire, Fond du Lac, Gogebic, MI, Green, Houghton, MI, Jackson, Kenosha, Kewaunee, Macon, GA, Manitowoc, Marin, CA, Marinette, Milwaukee, Nash, NC, Oneida, Ontonagon, MI, Outagamie, Ozaukee, Rock, Rusk, St. Croix, Sheboygan, Taylor, Trempealeau, Vernon, Vilas, Walworth, Washington, Waukesha, Winnebago, Wood |

628