Applied Animal Behaviour Science

Adverse effects of hunting with hounds on participants and bystanders
--Manuscript Draft--

Manuscript Number:	
Article Type:	Research Paper
Section/Category:	Non-Farmyard
Keywords:	human-carnivore conflict and coexistence; dog; illegal take; poaching; wolf
Corresponding Author:	Adrian Treves, PhD UNITED STATES
First Author:	Adrian Treves, PhD
Order of Authors:	Adrian Treves, PhD
	Laura Menefee
Abstract:	In the face of biodiversity crises and concerns raised about animal ethics, some societies are re-examining many human uses of animals. One activity that may once have been unobjectionable has undergone little scrutiny in most countries: hunting mammals with free-running hounds. We present two novel datasets about this under-studied behavioural interaction and hunting method. In the state of Wisconsin, USA, hounds and gray wolves occasionally interacted aggressively and human by-standers reported adverse interactions with hounds and their handlers. Such self-selected samples cannot be used to extrapolate in space or time but do provide new information. Between 1999 and 2012, 176 hounds were reported to have suffered injury during encounters with wolves. No government data were collected on how many wolves or other non-target animals were injured by hounds despite legal requirements. We describe the grouping behaviour, patterns of injury and vulnerability to attack, hound phenotypes, and other ethological data associated with aggressive wolf-hound interactions. We analyze wolf life history, seasonal life history patterns, bite locations on hound bodies, consumption of hound carcasses, and the outcomes of interactions to test two hypotheses for wolf-hound interactions. We find weak or no support for either and propose new hypotheses focusing on the behavior of hounds and handlers primarily. We also report handler and hound interactions with human bystanders. Between 2015-2021, bystanders (n=105) reported various incidents of illegal behaviour by hounds and handlers, adverse interactions with law enforcement, and other events associated with hunting with hounds. We propose reforms to policy for wildlife, hunting, and law enforcement, and reforms to husbandry by hound handlers.
Suggested Reviewers:	Paul Paquet University of Victoria ppaquet@baudoux.ca wolf, dog, human conflict expert
	Patricia Mcconnell, PhD retired, University of Wisconsin-Madison trisha@patriciamcconnell.com Dog, sheep and applied animal behaviour expert
	Catherine Hill, PhD Professor, Oxford Brookes University cmhill@brookes.ac.uk human-wildlife interactions and the social scientific dimensions

Highlights

- We describe self-reports of hound and handler behavior during wolf-hound interactions (WHI) and correlates of outcomes (injury or fatality).
- We describe self-reports of by-standers' experiences of hound and handler behaviors, including allegations of unlawful hunting practices and interpersonal interactions.
- We find equivocal support for two hypotheses centered on wolf behaviour to explain wolf-hound interactions (WHI).
- We propose a new hypothesis focused on hound and handler behaviour to explain WHI frequency and severity.

Adverse effects of hunting with hounds on participants and bystanders

Adrian Treves[a¥*], Laura Menefee[b†*]

[a] Nelson Institute for Environmental Studies, 30A Science Hall, 550 North Park Street, University of

Wisconsin, Madison, WI, 53706, USA, +1-608-890-1450, atreves@wisc.edu,

[b] Department of Literature, Leeward College, University of Hawai'i, Pearl City, HI and Sierra Club

Wisconsin Chapter, mail c/o A. Treves, 30A Science Hall, 550 North Park Street, University of Wisconsin,

Madison, WI, 53706, USA, menefee.laura@gmail.com

[†] These authors have contributed equally and share first authorship.

* Correspondence: atreves@wisc.edu, menefee.laura@gmail.com

Abstract

In the face of biodiversity crises and concerns raised about animal ethics, some societies are re-

examining many human uses of animals. One activity that may once have been unobjectionable has

undergone little scrutiny in most countries: hunting mammals with free-running hounds. We present

two novel datasets about this under-studied behavioural interaction and hunting method. In the state of

Wisconsin, USA, hounds and gray wolves occasionally interacted aggressively and human by-standers

reported adverse interactions with hounds and their handlers. Such self-selected samples cannot be

used to extrapolate in space or time but do provide new information. Between 1999 and 2012, 176

hounds were reported to have suffered injury during encounters with wolves. No government data were

collected on how many wolves or other non-target animals were injured by hounds despite legal

requirements. We describe the grouping behaviour, patterns of injury and vulnerability to attack, hound

phenotypes, and other ethological data associated with aggressive wolf-hound interactions. We analyze

wolf life history, seasonal life history patterns, bite locations on hound bodies, consumption of hound

carcasses, and the outcomes of interactions to test two hypotheses for wolf-hound interactions. We find

weak or no support for either and propose new hypotheses focusing on the behavior of hounds and

handlers primarily. We also report handler and hound interactions with human bystanders. Between

2015-2021, bystanders (n=105) reported various incidents of illegal behaviour by hounds and handlers,

adverse interactions with law enforcement, and other events associated with hunting with hounds. We

propose reforms to policy for wildlife, hunting, and law enforcement, and reforms to husbandry by

hound handlers.

Keywords: human-carnivore conflict and coexistence, dog, illegal take, poaching, wolf

1.   Introduction

In the face of biodiversity crises and concerns raised about animal ethics partly caused by climate

change and partly by human-induced mortality, some societies are re-examining many human uses

of animals that once seemed unobjectionable. For example, uses of poison, off-road vehicles, fire

suppression, etc. have undergone scrutiny for their societal benefit-cost estimates and their effects

on domestic animals, wildlife, and ecosystem health. One human activity that may once have been

unobjectionable has undergone little scrutiny: hunting with free-running hounds loosed far from

their owners.

In our search for work on hunting with dogs or hounds in Google Scholar, the search phrase

'hunt with (hound or dog)'' yielded 38,300 results declining by half when '-bird' was added to the search

string to exclude bird-hunting with dogs. By contrast, 'hunt -bird 'yielded 3.16 million results. Therefore,

hunting with hounds is under-studied. Current practitioners are also secretive, judging from their

introduction of a bill in the state of Wisconsin, USA legislature to prohibit video-recording of the activity

which is currently under appeal in federal court (2021). Yet, the practice of hunting mammals with

hounds has been recorded since 8,000 years at least (Guagnin et al., 2018), praised by President

Theodore Roosevelt in 1902 (Roosevelt, 1902), and is legal in numerous countries and several U.S. states

(Heberlein, 2000; Hristienko and McDonald, 2007). Despite its long history, loosing mammal-hunting

hounds to pursue prey, some as large as brown (grizzly) bears *Ursus arctos*, may have harmful effects on

people in their path, on the hounds themselves, and on target and non-target wildlife they encounter or

pursue (Grignolio et al., 2011; Gompper, 2013). Therefore, we present data on reports by self-selected

owners alleging harm to their hounds, following interactions with wolves. We also present self-reported

perceptions of human by-standers who experienced the behaviour of hounds during training or hunting.

We offer these two data sets to describe this poorly studied hunting method.

1.1 Non-target animals and their interactions with hounds

When pets kill wildlife, biodiversity may diminish or ecosystem health may deteriorate (Grignolio et al.,

2011; Gompper, 2013). Dogs are potential predators of domestic animals and wild ones (Bowers, 1953;

Ciucci and Boitani, 1998). Larger size and greater competitive ability of the wild animals may alter the

risk posed by hounds, just as the danger to non-target animals may increase if hounds outnumber or

outweigh them. Hounds are often used for pursuit of mammals larger than individual hounds;

behavioural changes in wildlife are often reported. For example, black bears *U. americanus* were

reported to avoid encounters with hounds and in so doing approach people and major roads more

frequently (Stillfried et al., 2015). Some animals will stand their ground when hounds encounter them

(Treves et al., 2002; Wydeven et al., 2004; Backeryd, 2007). When hounds encounter animals that can

defend themselves effectively, the hounds may be injured.

Researchers have examined aggressive encounters between wolves and dogs in many regions

(Fritts and Paul, 1989; Backeryd, 2007; Kaartinen et al., 2009). The States of Wisconsin and Michigan,

USA, have a relatively longer history of such research. Spatial patterns of wolf *Canis lupus* attacks on

hounds are somewhat predictable (Treves et al., 2002; Wydeven et al., 2004; Bump et al., 2013; Olson et

al., 2014). The risk of an attack appeared to be higher in areas with more public land, larger wolf packs,

closer to a wolf pack, and when baits were left out longer. Here, we examined self-reports by owners on

the characteristics of both canids involved, and self-reported correlates of the outcomes of wolf-hound

interactions (WHI).

At the time of our study, it was illegal for hounds to attack wild animals, but such attacks might

have occurred anyway. We do not have evidence of which animal in a WHI initiated aggression or

escalated it to the point of injury or death. We only present data on the outcomes for hounds because

outcomes for wolves were not documented. Federal law under the Endangered Species Act and state

law against animal fighting would both require authorities to collect data on the latter. We cannot

explain why the database containing of such information. Therefore, we cannot rule out the possibility

that wolves responded defensively to hound attacks. The evidence for wolf attacks on hounds came

from handlers seeking compensation or other forms of redress (Treves et al., 2002; Ruid et al., 2009;

Treves et al., 2009). In a prior study, a number of wolf deaths caused by other canids were invariably

attributed to other wolves (Treves et al., 2017b). Yet, veterinary pathologists might not be able to

reliably distinguish large dogs, such as hunting hounds, from wolves by simple scrutiny of bite marks

without DNA analysis (Plumer et al., 2018). Therefore, our sample is necessarily biased toward handler

concerns and outcomes for hounds. This should not be construed as evidence that wolves were the

aggressors or that wolves remained unharmed by hounds.

1.2 Hounds and humans

Much research has measured attitudes of hound-handlers towards wolves (Supplementary Material 1).

In brief, as a group, hound-handlers have the lowest tolerance for wild wolves in any group measured

thus far. There are no published data on hound-handler' encounters between by-standers and either

handlers or their free-running hounds. The Sierra Club Wisconsin Chapter (SCWC) began to fill the gap

with the survey we report here. As part of a National Sierra Club initiative, the SCWC subcommittee

discussed the first-hand reports members had received of by-stander experience. Because such first-

hand reports are likely to be remarkable, most reports were adverse. The SCWC subcommittee

requested information from local law enforcement in the counties from which reports and also the

Wisconsin Department of Natural Resources (WDNR) for incident report data. They were informed by

both agencies that "no such data were recorded." Thus, SCWC members led an effort at gathering

information more broadly and systematically. We focus on the behaviour of hounds, wolves, and

handlers as reported by by-standers. We test two ethological hypotheses about WHI. First, the

predation hypothesis predicts WHI occur when wolves attack hounds for food, and second, the

territoriality hypothesis predicts that WHI occur when wolves defend territory or pups (Treves et al.,

2002; Wydeven et al., 2004).

2.0 Material and methods

2.1 Wolf-hound interactions (WHI) behaviour, grouping, and phenotype

During our study period, hounds were legally used to hunt many mammals, including smaller carnivores

and black bears. Bear hunting in Wisconsin occurred legally from the last week of August into October,

and hound training was legal in July and August (Treves et al., 2009; Treves et al., 2010). Hound hunters

accounted for approximately 40% of the annual take by bear hunters (Treves et al., 2010; Olson et al.,

2014). Typically hounds were loosed from vehicles and allowed to run far from owners, without control

(Wydeven et al., 2004). Hounds were often fitted with global positioning systems (GPS) or VHF radio-

collars, allowing the owner to follow remotely the movements of hounds and determine when and

where a bear had been treed. Hunters used groups of up to 6 hounds to track and trail prey during

training or hunting (Wydeven et al., 2004).

We examined WHI from 7 August 1999 through 19 January 2012. Case files documented 145

killed and 31 injured hounds deemed by state or federal verifiers as "confirmed" (85%) or "probable"

(15%) WHI, see methods in (Treves et al., 2002). We believe most WHI incidents were reported because

of a compensation program characterized as more generous than other jurisdictions (Naughton-Treves

et al., 2003; Treves et al., 2009). Bump et al. (Bump et al., 2013) suggest fewer WHI are reported in

Michigan's wolf range because hound owners receive no compensation. Owners with confirmed losses

in Wisconsin were eligible to receive up to $2,500 per hound based on the estimated value of the

hound. Since the compensation program began in 1985, nearly $350,000 dollars were paid to hunters to

compensate for hounds injured or killed by wolves. Between 1985 and 2006, payments for hunting

hounds comprised 37% of all compensation (Treves et al., 2002; Naughton-Treves et al., 2003; Ruid et

al., 2009; Treves et al., 2009). WDNR provided compensation for domestic animals injured or killed by

wolves, including hounds (Treves et al., 2002; Ruid et al., 2009; Treves et al., 2009). United States

Department of Agriculture agents assumed responsibility for verifying WHI in 1990, and conducted most

of the investigations used in our analysis of WHI (Willging and Wydeven, 1997).

WHI case files included written reports and forms documenting field investigations, including

necropsy data, photos, veterinary reports, and anecdotal reports handlers. During the early years of

record-keeping, documentation and reporting of depredations lacked uniformity; thus, some portions of

the data were missing, resulting in lower sample sizes for various analyses. We limited our analysis of

WHI to hounds used to hunt bears, bobcats, or coyotes. A total of 91% of WHI occurred while pursuing

these species, which involves different breeds and use of hounds than for other quarry, such as

waterfowl, upland birds, or rabbits. Hounds that hunt large prey such as bears or coyotes are typically

breeds of large size and build, frequently Plott (18-27 kg), or larger hound breeds (Bluetick, Redbone, or

Walker coon-hounds (18-36 kg). Occasionally, WHI files did not specify the type of prey being pursued.

In these cases, if the breed of hound was one of the five listed above, we assumed the WHI occurred

while pursuing the above three wildlife species. In total, we report on 176 case files. We quantified the

frequency of WHI among breeds of hounds. If the hound was reported as a mix of multiple breeds, we

used the first breed listed. We pooled breeds in an "other" category when a single breed had too few

WHI to meet the assumptions of the *chi*-squared test. No data are available on breed frequencies or

preference by hunters in Wisconsin, with which we could estimate relative risk by breed.

Our analysis of the body site bitten was limited because a number of hound carcasses were

partially or wholly consumed before retrieval by an owner arriving late at the scene. We pooled head,

neck, and throat into one category and all other sites in another category, to test if outcomes of WHI

differed by bite site.

We compared hound group size, number of hounds involved in the WHI, and wolf pack size

using (1) number of wolves seen and reported by hunters (observed), and (2) WDNR-reported wolf pack

sizes from the winter preceding the WHI (censused). Because wolf packs exhibit fission-fusion sociality

and packs disaggregate often in the summer (Mech, 1970), when many WHI occurred. Wolf packs in

Wisconsin average 4 adults in late winter (range 2-12), usually composed of offspring of the two

(usually) unrelated alpha breeders. By summer, pups are exiting the dens but range near them or near

other pack rendezvous sites until the pack regularly fuses and travels long distance in mid- to late Fall.

During this time, most wolf pack members return periodically to rendezvous or den sites to assist with

pup-rearing, and consequently have higher food demands, perhaps requiring wolves to forage more

frequently (Ruprecht et al., 2012). From birth until the end of August, wolf pups experience the highest

growth rates, with September representing a critical month for weight gain (Kreeger, 2003). In some

cases, wolf pups have been observed gaining as much as 3.6 pounds per week (Van Ballenberghe and

Mech, 1975). Pup growth, critical to survival, is limited by food quality and availability. By late August,

growth begins to taper (Pulliainen, 1965), as does rendezvous site use (Van Ballenberghe and Mech,

1975; Kreeger, 2003; Ruprecht et al., 2012). We acknowledge R. P. Thiel for the suggestion that

seasonality of wolf life-history on an annual cycle would predict more risk of WHI in late summer and

autumn because of all pack members and especially pups' caloric needs. Finally, we analyzed the

temporal occurrence of WHI as it relates to public hunting seasons, and we compared the frequency of

WHI during the hound-training period (July-August) to that during the bear-hunting season (treated as

September-October).

2.2 Handler behaviour and husbandry

The state wildlife agency implemented several methods for mitigating or preventing WHI,

including compensation for handlers' self-reported losses, encouraging the use of warning devices on

collars to deter wolves, and statewide communication to hunters on recent hound injuries and their

locations, and designation of Wolf Caution Areas (WCAs). The state communicated the location of

higher-risk WCAs online, posted in the field, and in other ways to handlers (Wydeven et al., 2004; Olson

et al., 2014). Within WCAs, the WDNR recommended that bear hunters release hounds >2 miles from

known rendezvous sites. Prior work documented handlers' willingness to risk hounds in posted WCAs,

even within the same season and even within hours of previous WHI or WCA posting (Wydeven et al.,

2004; Olson et al., 2014). Compensation records also document multiple payments to the same owner

or handler within a single season (Treves et al., 2009). These data suggest not all hunters heeded the

state's warnings. Compensation programs did not clearly reduce WHI rates. Much scholarly discussion

has focused on the moral hazard (in short, negligent owners have no incentive to protect their animals if

they will be paid for losses). Prior research on Wisconsin's compensation program addressed moral

hazards (Naughton-Treves et al., 2003; Treves et al., 2009). The state only changed the source of the

funds, not the conditions for payment since that work was published. Therefore, cooperation of the

state legislature may be essential to changing an incentive for WHI to occur into a disincentive for

hounds and non-target animals to be put at risk. Handlers may be able to prevent WHI by using

protective vests or stronger collars (Backeryd, 2007; Khorozyan et al., 2020), keeping hounds leashed

until the targeted game species is located, or bringing first aid kits on the hunt, although the possible

effect of these interventions has not been studied in Wisconsin.

2.3 Survey of by-standers

The Sierra Club Wildlife Committee (SCWC) subcommittee on Protecting Native Forests and Wildlife

collected systematic information from by-standers' experiences of hound hunting. The SCWC led and

posted the survey on the Sierra Club Wisconsin Chapter website through 2015-2021, and administered

the survey. The survey appeared at https://www.sierraclub.org/wisconsin/protecting-native-forests-

wildlife accessed 20 August 2022. SCWC members also printed hard copies of the instrument and

distributed these at wolf and wildlife related meetings and conferences in Wisconsin in 2015 and 2016.

SCWC also invited any members of the public who described adverse hounding encounters to fill the

online report. Therefore, respondents were self-selected with attendant biases.

We analyzed anonymized data stripped of identifying information by the SCWC administrators.

The 25-question survey (Supplementary Material Appendix 1) is organized in four sections:

Observations; Trespass; Property Damage, Personal Injury or Threats; and Interactions with Law

Enforcement, totaling 22 yes/no questions and 4 items that allowed unstructured responses to

elaborate on answers. Respondents could identify county of residence and interaction, if different.

Respondents were asked how many hounds they saw during each interaction they reported. When two

respondents mentioned the same interaction but different numbers of hounds, LM averaged and

rounded up for the number of hounds.

About 80% of respondents used the online form to report anonymously, and 20% sent their

responses directly to SCWC via mail, phone, email, or in person while being assured of anonymity. LM

screened the sample to eliminate responses which identified no adverse incident involving hunting

hounds, as these respondents generally used the report format to express an opinion about the practice

of hunting with hounds, which we did not analyze because it was outside the purview of this survey

(Supplementary Material 2). We screened for multiple reports of the same incident with identifying

factors such as location and date. After the screening, the sample presented here appears to come from

independent incidents although we had no way to verify location or date.

2.4 Statistical analyses

We performed statistical analyses (SAS Institute, 2013) using Student's paired *t* tests to compare the

differences in average estimated ages of hounds and numbers of wolves during the attack, in relation to

the outcome of the WHI (i.e., killed or injured) after evaluating if variances were equivalent (*F* test). All

statements of statistical significance are based on *P* ≤ 0.05. We used Spearman rank correlations to

detect associations between multiple continuous variables.

2.5 Ethical note

The data presented are exempt from approvals for research on human subjects because (a) the

data were collected by another organization (WDNR and SCWC) and distributed to the authors without

individually identifiable information; (b) all SCWC survey responses were completely anonymous; and (c)

the identity of complainant handlers was likewise unknown to us and redacted from records by the state

agency that received the complaints. Use of hounds for hunting is a state-regulated activity. No research

was conducted on the hounds.

3.0 Results and Discussion (as allowed by instructions to authors, p.9 para 3)

3.1 Wolf-hound interactions (WHI) behaviour, grouping, and phenotype

In 176 case files, we found 140 independent WHI cases during our study period, where a case

reported on the same day and location by different owners were pooled into one WHI. Records included

145 killed (83%) and 31 injured (17%) hounds, similar to 71% and 82% reported in Nordic countries

(Kojola and Kuittinen, 2002; Backeryd, 2007). The high percentage of fatalities in both cases might

reflect that handlers sometimes took hours to find a distant hound. Therefore, sub-lethal injuries might

not be attributed to a WHI if handlers and hounds reunited long after it ended; this effect would bias

WHI records upward to more fatalities. Similarly, the cases involving distant handlers might not have

escalated to lethal injuries if the handlers had arrived earlier.

Wolf injuries and deaths in WHI were not documented nor reported in case files. Another

possible source of missing data might come from misidentifying wolves as the wild interactant, given the

availability of compensation for such WHI and not for casualties from other wildlife (Treves et al., 2009).

In total, 89% of WHI occurred while hunters reported pursuing black bears, bobcats *Lynx rufus* 6%,

coyotes *C. latrans* 4%, raccoons *Procyon lotor* 1%). However, we lack independent data on the animal

being pursued by those hounds at the time of WHI. We also lack the relative frequencies of targeting

each species with hounds statewide and over time. There was no association between the outcome of

WHI categorized as either injury or death and the prey being pursued by hunters (X^2 = 1.9, *P* = 0.75, *df* =

4, *n* = 140). The bear-hound-training period accounted for 62% of WHI, whereas the bear-hunting

season accounted for 28% despite being the same length approximately. Outcomes were not associated

with month annually (X^2 = 8.5, *P* = 0.38, *df* = 8, *n* = 176, Table 1). These two results suggest that

seasonality and Wolf reproductive timing did not predict injury or death of hounds, which undermine

both the predation and territoriality hypotheses. We lack information on the frequency with which WHI

was initiated by hounds, which could result in very different causal hypotheses focused on the

motivations of hounds and handlers.

251   Neither sex nor age of the hounds was associated with the outcome of WHI (sex X^2 = 1.32, *P* =

0.25, *df* = 1, *n* = 151; age *t* = -0.71, *P* = 0.49; variances were equal *F* = 0.49). The Treeing Walker

Coonhound was the most common breed in WHI (33.3%, *n* = 51), followed by the Plott (27.5%, *n* = 42).

There was a significant association between breed and outcome (X^2 = 10.7, *P* = 0.03, *df* = 4, *n* = 176).

Notably, the Plott fatality frequency of 95.4% was higher than the average 81.2% (Table 1). Plott hounds

are the smallest and most vocal hound breed commonly used in Wisconsin for mammals. Small body

size has been implicated in the risks and fatalities associated with WHI (Fritts and Paul, 1989; Peterson,

1995; Backeryd, 2007; Kaartinen et al., 2009). The Plott breed is also known for its baying vocalizations,

which might alert wolves from a long distance. Similar elevated risks of dog vocalizations were reported

in Nordic countries (Backeryd, 2007; Kaartinen et al., 2009). Small size may make a hound more

vulnerable to head and neck bites. Bites to the neck were associated with higher fatality rates in a

Scandinavian study (Backeryd, 2007).

Our analysis on hound body site bitten was limited to 109 WHI. We cannot be certain that

wolves inflicted every bite because of the above-mentioned delays in discovering hounds and the

missing center created by compensation only for attacks by wolves. Of the 109 carcasses with bite

information, 50 provided one bite location (46%), 37 provided two locations (34%), and 22 provided 3 or

more locations (20%). Considering all bite locations (n = 193), the single most frequent bite site was the

neck (33%), followed by back (17%), upper thigh (12%), and chest (10%). We considered bites to the

head, shoulders, neck (as opposed to throat), back, and upper thighs as indicative the hound had been

lower than its attacker. Those upper body parts were represented in 72% of the 193 bites whereas

under-parts (throat, groin, sternum, ribs, lower legs, abdomen) were represented in 28% of bite

locations. We found no relationship between body site bitten and outcome, when we separated neck

and head bites from others (X^2 = 1.5, P = 0.22, df = 1, n = 66).

Another factor affecting the vulnerability of canids to attack by other canids is numerical

superiority. An average of 1.3 hounds were injured or killed per WHI (maximum 5 in a single WHI). The

size of the hound group averaged 3.8 SD 1.4, n = 57; only 3 (5%) of those WHI involve a single hound in

the handler's control; and also the number involved in the WHI (2.6 SD 1.3, n = 47 with 9 (19%) of those

WHI involving only 1 hound, were both similar to the number of wolves observed by handlers (2.9 SD

1.2, n = 15), the census pack size for the pack blamed by the state or federal agent tasked with verifying

the report (2.4 SD 1.0, n = 19; n = 4 included information for both observed and censused). The

outcomes were not associated with the number of hounds, number of wolves, or difference between

the two in a given WHI by any of the measures of group size or pack size above (Welch test assumes

unequal variance, F < 0.72, P > 0.41 in every test). Wolves injured or killed hounds in groups with

superior numbers in 44% of WHI with such data (n = 16). Aggression between wolves and coyotes in

Yellowstone National Park had fatal consequences when wolves outnumbered the smaller coyotes, but

not when coyotes outnumbered wolves, suggesting that group size exerted less influence than

individual body size differences in determining outcomes between canids (Ballard et al., 2003; Merkle et

al., 2009). In our study, if wolves were motivated by predation or territorial defense as suggested by the

two hypotheses, we might expect pack size to predict the outcomes of WHI. Again, this suggests the

need for new hypotheses as above.

Of 80 deaths with data on consumption of a carcass, 49% of hound carcasses were partially

consumed. Of those 80 hounds consumed by wolves, 71% occurred July–August and 27% in September–

October. When we combine the former timing of WHI, site location of bites, and the consumption data

above, we can test two hypotheses about the cause of WHI. The timing of WHI presents equivocal

evidence for both hypotheses. Higher frequencies of WHI occurred during the hound training period in

July and August than during the autumn black bear hunt in September and October (Table 1). Elevated

risk in July and August might have been associated with the practice of baiting, as wolves visit bear bait

sites in search of food (Bump et al., 2013). In Wisconsin, bear bait sites could be legally established as

early as April and could last the entire wolf pup-rearing season. Bump et al. (2013) documented that the

risk of WHI was three to seven times greater in Wisconsin than in adjacent Michigan, citing the extended

bear-baiting period as a probable cause for the much higher risk of WHI in Wisconsin. That might

support the predation hypothesis. However, bear baiting was confounded with wolf pup defense

predicted by the territoriality hypothesis. The hound training period coincided during the study with

wolf use of rendezvous sites or den sites. See the introduction for the life-history on an annual cycle of

wolves, which the majority of hound packs would not experience in the same way because of human

provisioning, aseasonal reproduction, and more human intervention in breeding behaviour. The

consumption of hound carcasses might corroborate the predation hypothesis, but that is not persuasive

because consumption was recorded in only approximately half of the WHI and we do not know if the

wolves that attacked were the consumers. Nor can we rule out that consumption followed after the

primary motivation for aggression in either the hounds or the wolves. The hound carcasses and bite

locations provided limited insight. Bites to head, neck, and throat represented 41% of bite locations on

hound carcasses. The predation hypothesis might find support from this result because cranio-cervical

killing bites are associated with predation by many mammals (Steklis and King, 1978; Van Valkenburgh

and Ruff., 1987). But the greater height of wolves off the ground then off most towns would predict

such bite locations in any case. In sum, we find equivocal support for both hypotheses. This could imply

both are correct or we are missing information, such as whether the hounds were pursuing Wolves. We

also do not have information on the body condition of any of the participants, which seems essential

data to test the prediction hypothesis.

319   3.2 Handler behaviour and husbandry

Husbandry, such as avoidance of rendezvous sites and use of bells on collars were difficult to evaluate,

because of a lack of data on prevalence, use, and effectiveness of these potentially preventive methods.

Overall, 55 (31%) records reported whether hounds in WHI wore bells on their collars, with 20 (36%)

wearing them and 35 did not (64%). But we have no data on the use of bells among hounds that did not

enter the WHI database and the majority of records did not contain any information on devices.

Outcomes were not associated with hunter self-reports of affixing bells to collars. Another step handlers

might take to protect hounds and wolves would be to release hounds only in low-risk areas. The state

did not systematically collect data on whether air warnings about high-risk WCAs were heeded. The WHI

records did not contain such information.

329   3.3 Survey of by-standers

In all, 105 respondents reported adverse incidents with hunting hounds from 51 Wisconsin counties, 4

Michigan counties, and 5 counties from other states. The average number of incidents per county was 2.

Seven respondents declined to specify location, but timing distinguished the reports from other

respondents' reports. The 105 respondents reported 119 separate incidents (Table 2). Of the 105, 42%

reported the hounds observed were not accompanied by a handler; similarly, 41% reported finding

abandoned or lost hounds on their property. In those cases, some respondents reported contacting local

animal shelters, law enforcement or handlers via phone numbers on collars. Overall, 63% of 105

respondents described incidents of trespass including hounds running on their property without

permission, handlers found on property without permission, or running hounds on property after being

denied permission. Beyond trespass, 18% of respondents described illegal or nuisance behaviour:

damage to property caused by hounds, including downed fencing, damaged landscaping and gardens,

injury to self and livestock, dead wildlife left on property, vandalism or litter. Most seriously, 11%

reported injury to pets or livestock by hounds, although 24% reported knowledge of hounds attacking

others 'pets or livestock. Of the 105, 8% describe direct encounters with hounds resulting in personal

injury or being chased.

345   Moreover, 31% reported threatening altercations with hound handlers, including being

346   unwillingly detained by hound handlers 'trucks on public roads, or their own private driveways. Of 105

347   respondents, 51% of respondents reported they "feel intimidated by hound handlers," and 44% feared

348   retaliation from handlers for reporting confrontations to law enforcement.

349   Finally, respondents describe being given false information by law enforcement officers,

including local sheriffs and game wardens, and also reported filing official complaints upon which no

discernible action was taken. Overall, 36% of 105 respondents believed a conflict of interest existed for

law enforcement officers, including game wardens, because of relationships between officers and

handlers, or because the officers were believed to hunt with hounds themselves. Prior research studies

report that hound handlers as a group contain a substantial contingent willing to break the law and flout

regulations intended to protect animals and more so than other groups active in wolf range

(summarized in SM 1). Actual poaching evidence is consistent with the above survey reports (see

Conclusions)

358   3.4 Comparing numbers of hounds from WHI and survey data

Survey respondents reported 2–8 hounds per interaction (average 3.7, mode 2). That average is

identical to the average number of hounds that handlers reported in their pack in WHI above. This

seems to be corroborating evidence of accuracy in reports of hound pack sizes in both datasets, as

neither set of complainants was aware of the other. Given the rarity of single hounds (5%) in WHI, the

bystander reports of >1 hound seem unsurprising. Similarly, bystanders reported >6 hounds in 3 events

(8% of reports that include these data) but handlers never reported >6 in their pack after a WHI. The

legal limit per handler was multiple handlers may release more than 6 hounds. The topic of comparing

self-reports by handlers and self-reports by bystanders raises an issue of expertise and deference to it. A

common assumption in peer-reviewed commentary on hunting is that hunters have lay knowledge

gained through local experience and expertise in their practices. This often plays out among academics

and manager-authors as favoring the views and preferences of local communities and of hunters over

others, especially outside experts (von Essen et al., 2015; Redpath et al., 2017; Richardson, 2022). But

that assumption is unjustified when two local, lay types of expertise are pitted against each other as in

an unknown proportion of our data when hound handlers and the bystanders complaining about them

were equally local and both held lay expertise. The second flaw is when the lay bystanders complained

to law enforcement experts about hounds or handlers, then the power asymmetry is reversed from the

usual academic debate. In short, the local, lay expert complainant deserves the attention and

compassion typically reserved for local hunters. Therefore, we call for more just and fair consideration

of the under-represented and marginalized in wildlife management.

4.0 Conclusion

Hunting with hounds is rarely studied. Here we fill the gap partly with data on the behaviour of

hounds and wolves associated with aggressive wolf-hound interactions (WHI). Also, we present self-

reported data on the husbandry of handlers and the experiences of bystanders exposed to hound and

handler behaviour. The shortage of data on wolf casualties makes the lack of research a concern for

wildlife law enforcers and wolf interests. Our inability to find strong support for either the predation

hypothesis or the territoriality hypothesis, which focus on wolf motivation for WHI, suggests either a

multi-causal phenomenon or we lack hypotheses for the motivation of hounds to initiate WHI. The lack

of a seasonal pattern in our data argues against the notion that wolves are the primary aggressors; WHI

may be a mutual affair or may be largely initiated by hounds. Treeing Walker coonhound was the breed

most commonly involved in WHI. Most WHI were fatal for hounds. Plott hounds were disproportionally

represented among fatalities. We could not confirm or reject a protective role of special collars with

bells or defensive features. Nevertheless, we recommend the state obligate veterinary clinics that treat

hounds for wildlife injuries to report each such incident so the welfare of hounds and preventive actions

taken by handlers can be evaluated by professional veterinarians and wildlife scientists. In light of recent

findings that studded leather collars protect cattle against leopards (Khorozyan et al., 2020), more

precautions by handlers seems a prudent recommendation. A majority of WHI involved hounds

pursuing black bears compared to other prey but no one can say with confidence what animal the

hounds were pursuing when WHI began. Outcomes of WHI (injury or death of hounds) were not

associated with the number of wolves observed or censused near the site, or the numerical differences

between wolves and hounds, hound age, hound sex, the species of prey targeted by hunters, or the

month in which WHI occurred. Regarding numerical superiority, perhaps the large group sizes of hounds

in Wisconsin and uncertainty about the number of wolves involved both obscured associations with the

fatality rates during WHI. We found a critical support for the production and territoriality hypotheses,

which may reflect the lack of data on which canid initiated a WHI or a mixture of motivations for WHI.

We report unlawful actions involving trespass, property damage, attacks, threats, intimidation and

harassment of by-standers or their domestic animals, perpetrated by both hounds and handlers. The

survey data and WHI self-reports by handlers are self-selected samples, which cannot be verify directly

or used to extrapolate rates, frequencies, or representativeness in space, time, or demography.

Nevertheless, the data support improved husbandry of hounds, more transparent law enforcement, and

wildlife protections. By contrast, our other self-selected data set (handler complaints of hound losses in

WHI) were motivated by a compensation program that paid for injured or dead hounds. That program

also seems to need reform given that no information on harm to wolves was collected and it might be

impossible to verify that the handlers or hounds were acting lawfully at the time of the WHI.

Furthermore, the size of packs of hounds (up to 6) and the possibility that multiple handlers can run

multiple packs in the same area simultaneously creates a clear danger for wolves, given our results

about numerical superiority. These results highlight a need for improved regulation, greater oversight

and more energetic enforcement of activities involving the use of hounds during hunting. Our results are

also consistent with findings of more serious wildlife crimes. Wolf-poaching is the major cause of

mortality in US wolf populations (Treves et al., 2017a). Wisconsin is no exception. Much of the poaching

involves concealment or destruction of evidence, which reflects intent to break the law, has repeatedly

risen in incidence along with policies that permit some legal wolf-killing in several US wolf populations.

These findings indicate that would-be poachers profit from governmental laxity to act unlawfully or that

would-be poachers use the cover of legal hunting to act unlawfully (Chapron and Treves, 2016a; b;

Santiago-Ávila et al., 2020; Louchouarn et al., 2021). Most recently, researchers found that cryptic

poaching rose during hound training, bear-hunting seasons, and deer-hunting seasons (Santiago-Ávila

and Treves, 2022). The 2021 Wisconsin wolf-hunt saw the most rapid season closure and over-kill in

Wisconsin wolf management history with 218 wolves killed in <72 hours with >80% being killed by

hunters using hounds (Treves et al., 2021). Unpublished necropsy data collected by the Great Lakes

Indian Fish & Wildlife Commission indicate hounds delivered potentially lethal bites during those

hunting incidents, which represent unlawful take by hunters (Treves et al., 2021). Finally, the allegations

of unlawful hound and handler behaviour suggest law enforcement is needed.

In USA wildlife law, the Endangered Species Act (ESA) and federal court cases surrounding it

make clear that some hound handlers are vulnerable to prosecution. First, any "take" of wolves

(including harassment, pursuit, injury, killing, etc.) is prohibited under the ESA regardless of whether the

perpetrator knew the wild animal harmed was listed (Newcomer et al., 2011). The absence of data

revealed by this study indicates that hounding is not adequately regulated. Because wolves were often a

federal- or state-listed species during our study, the practice of hound-hunting in wolf pack territories

should be prohibited when wolves are a listed species unless it can be regulated to reduce WHI near

zero. Prohibitions on non-selective killing methods in the range of endangered species and prohibitions

on hunting non-listed species of similar appearance such as coyotes (Newsome et al., 2015), might

resolve some WHI. Potential legal jeopardy does not stop at the hound handlers. Given the state wildlife

agency and law enforcement refused the SCWC request for information or had no such documents (see

Introduction), the alleged infractions by handlers may still be within the statute of limitations.

Systematic and repeated failure to investigate or prosecute complaints of illegal activity can make a

department vulnerable to lawsuits and the imposition of oversight by higher authorities (e.g., federal

consent decrees). Legal jeopardy arises for the agency because the doctrine of prosecutorial discretion

may not protect a law enforcement agency from charges of systematic neglect of unlawful activities

(Newcomer et al., 2011) and see allegations of such systematic neglect in a case involving wolves (2015).

5.0 Acknowledgments: R. P. Thiel and C. Williamson shared data and analyses, E. Huntley helped design

the survey of by-standers, Heathcliff Wright designed graphics, and Sierra Club Wisconsin Chapter

administered the online survey and shared data with LM.

6.0 Declaration of competing interests: AT and LM perceive no potential competing interests. All

funding awarded to AT as of 13 April 2022 are listed here

http://faculty.nelson.wisc.edu/treves/archive_BAS/funding.pdf accessed 21 August 2022, which we

provide for transparency and a CV for disclosure of potential competing interests:

http://faculty.nelson.wisc.edu/treves/archive_BAS/Treves_vita_latest.pdf accessed 22 August 2022. LM

is a retired Associate Professor from the Department of Literature, Leeward College, University of

Hawaii and currently a member of: Endangered Species Coalition, Society for the Preservation of Poultry

Antiquities, National Wolfwatchers Coalition, Wisconsin Wolf Trackers, Wolf Patrol, Executive

Committee, Fox Valley Group, Sierra Club.

459   References and Citations

2015. WildEarth Guardians, et al. v U.S. Department of Justice, U.S. District Court Arizona.

2021. Brown, J. et al. v Kemp, J. et al., U.S. Court of Appeals, 7th circuit.

Backeryd, J., 2007. Wolf attacks on dogs in Scandinavia 1995-2005, Ecology Institute, Swedish University

of Agricultural Sciences, Grimso.

Ballard, W.B., Carbyn, L.N., Smith, D.W., 2003. Wolf interactions with non-prey, in: Mech, L., Boitani, L.

(Eds.), Wolves: behaviour, ecology, and conservation, University of Chicago Press, Chicago, pp. 259-271.

Bowers, R.R., 1953. The free-running dog menace. Virginia Wildlife 14, 5-7.

Bump, J.K., Murawski, C.M., Kartano, L.M., Beyer, D.E., Roell, B.J., 2013. Bear-Baiting May Exacerbate

Wolf-Hunting Dog Conflict. PLoS ONE 10.1371/journal.pone.0061708.

Chapron, G., Treves, A., 2016a. Blood does not buy goodwill: allowing culling increases poaching of a

large carnivore. Proceedings of the Royal Society B 283, 20152939.

Chapron, G., Treves, A., 2016b. Correction to 'Blood does not buy goodwill: allowing culling increases

poaching of a large carnivore'. Proceedings of the Royal Society B Volume 283, 20162577.

Ciucci, P., Boitani, L., 1998. Wolf and dog depredation on livestock in central Italy. Wildl. Soc. Bull. 26,

504-514.

Fritts, S.H., Paul, W.J., 1989. Interactions of wolves and dogs in Minnesota. Wildl. Soc. Bull. 17, 121-123.

Gompper, M.E., 2013. Free-ranging dogs and wildlife conservation. Oxford University Press., Oxford.

Grignolio, S., Merli, E., Bongi, P., Ciuti, S., Apollonio, M., 2011. Effects of hunting with hounds on a non-

target species living on the edge of a protected area. Biol. Conserv. 144, 641-649.

10.1016/j.biocon.2010.10.022G

Guagnin, M., Perri, A.R., Petraglia, M.D., 2018. Pre-Neolithic evidence for dog-assisted hunting strategies

in Arabia. Journal of Anthropological Archaeology 49, 225-236.

Heberlein, T.A., 2000. The gun, the dog and the thermos: Culture and hunting in Sweden and the United

States. Sweden & America Fall 2000, 24-29.

Hristienko, H., McDonald, J.E.J., 2007. Going into the 21st century: a perspective on trends and

controversies in the management of the American black bear Ursus 18, 72-88.

Kaartinen, S., Luoto, M., Kojola, I., 2009. Carnivore-livestock conflicts: determinants of wolf (Canis lupus)

depredation on sheep farms in Finland. Biodivers. Conserv. 18, 3503-3517.

Khorozyan, I., Siavash, G., Mobin, S., Soofi, M., Waltert, M., 2020. Studded leather collars are very

effective in protecting cattle from leopard (Panthera pardus) attacks. 10.1002/2688-8319.12013

Kojola, I., Kuittinen, J., 2002. Wolf Attacks on Dogs in Finland. Wildl. Soc. Bull. 30, 498-501.

Kreeger, T., 2003. The internal wolf: physiology, pathology, and pharmacology., in: Mech LD, B.L. (Ed.),

Wolves: behaviour, ecology, and conservation, University of Chicago Press, Chicago, pp. 192-217.

Louchouarn, N.X., Santiago-Ávila, F.J., Parsons, D.R., Treves, A., 2021. Evaluating how lethal

management affects poaching of Mexican wolves Open Science 8 (registered report), 200330.

Mech, L.D., 1970. The Wolf: The Ecology and Behaviour of an Endangered Species. University of

Minnesota Press, Minneapolis.

Merkle, J., Stahler, D., DW, S., 2009. Interference competition between gray wolves and coyotes in

Yellowstone National Park. Canadian Journal of Zoology 87, 56-63.

Naughton-Treves, L., Grossberg, R., Treves, A., 2003. Paying for tolerance: The impact of livestock

depredation and compensation payments on rural citizens' attitudes toward wolves. Conserv. Biol. 17,

1500-1511.

Newcomer, E., Palladini, M., Jones, L., 2011. The Endangered Species Act v. the United States

Department of Justice: How the Department of Justice derailed criminal prosecutions under the

Endangered Species Act. Animal Law 17, 241-271.

Newsome, T., Bruskotter, J.T., Ripple, W.J., 2015. When shooting a coyote kills a wolf: Mistaken identity

or misguided management? Biodivers. Conserv. 24, 3145-3149.

Olson, E.R., Treves, A., Wydeven, A.P., Ventura, S., 2014. Landscape predictors of wolf attacks on bear-

hunting dogs in Wisconsin, USA. Wildl. Res. 41, 584–597.

Peterson, R., 1995. Wolves as interspecific competitors in canid ecology, in: Carbyn, L., Fritts, S., Seip, D.

(Eds.), Ecology and conservation of wolves in a changing world., Canadian Circumpolar Institute.,

Edmonton, Alberta, Canada, pp. 315-323.

Plumer, L., Talvi, T.n., Männil, P., Saarma, U., 2018. Assessing the roles of wolves and dogs in livestock

predation and suggestions for mitigating human-wildlife conflict and conservation of wolves. Conserv.

Genet. https://doi.org/10.1007/s10592-017-1045-4.

Pulliainen, E., 1965. Studies on the wolf (Canis lupus) in Finland. Ann. Zool. Fenn. 2, 215-259.

Redpath, S., Linnell, J.D.C., Festa-Bianchet, M., Boitani, L., Bunnefeld, N., Dickman, A., Gutiérrez, R.,

Irvine, J., Johansson, M., Majić, A., Mcmanus, B., Pooley, S., Sandström, C., Sjölander-Lindqvist, A.,

Skogen, K., Swenson, J.E., Trouwhborst, A., Young, J., Milner-Gulland, E.J., 2017. Don't forget to look

down – collaborative approaches to predator conservation. Biological Reviews 92, 2157–2163.

Richardson, J.E., 2022. The Cows May Safely Graze: Placing Expert-Lay Relationships at the Center of

Overcoming the Expert-Lay Knowledge Divide. Rural Sociology in press. 10.1111/ruso.12426

Roosevelt, T.D., 1902. Hunting the Grisly and Other Sketches. G. P. Putnam's Sons, New York.

Ruid, D.B., Paul, W.J., Roell, B.J., Wideven, A.P., Willging, R.C., Jurewicz, R.L., Lonsway, D.H., 2009. Wolf–

human conflicts and management in Minnesota, Wisconsin, and Michigan, in: Wydeven, A.P., Van

Deelen, T.R., Heske, E.J. (Eds.), Recovery of Gray Wolves in the Great Lakes Region of the United States:

An Endangered Species Success Story, Springer, New York, pp. 279-295.

Ruprecht, J., Ausband, D., Mitchell, M., Garton, M., Zager, P., 2012. Homesite attendance based on sex,

breeding status, and number of helpers in gray wolf packs. J. Mammal. 93, 1001-1005.

Santiago-Ávila, F.J., Chappell, R.J., Treves, A., 2020. Liberalizing the killing of endangered wolves was

associated with more disappearances of collared individuals in Wisconsin, USA. Scientific Reports 10,

13881. /10.1038

Santiago-Ávila, F.J., Treves, A., 2022. Poaching of protected wolves fluctuated seasonally and with non-

wolf hunting. Scientific Reports 12, e1738 10.1038/s41598-022-05679-w

SAS Institute, I., 2013. JMP 11.0, Cary, North Carolina. .

Steklis, H.D., King, G.E., 1978. The craniocervical killing bite: Toward an ethology of primate predatory

behaviour. J. Hum. Evol. 7, 567-581.

Stillfried, M., Belant, J., Svoboda, N., Beyer, D., Kramer-Schadt, S., 2015. When top predators become

prey: Black bears alter movement behaviour in response to hunting pressure. Behav. Processes 120, 30-

39. 0.1016/j.beproc.2015.08.003

Treves, A., Artelle, K.A., Darimont, C.T., Parsons, D.R., 2017a. Mismeasured mortality: correcting

estimates of wolf poaching in the United States. J. Mammal. 98, 1256–1264. 10.1093/jmammal/gyx052

Treves, A., Jurewicz, R.L., Naughton-Treves, L., Rose, R.A., Willging, R.C., Wydeven, A.P., 2002. Wolf

depredation on domestic animals: control and compensation in Wisconsin, 1976-2000. Wildl. Soc. Bull.

30, 231-241.

Treves, A., Jurewicz, R.L., Naughton-Treves, L., Wilcove, D., 2009. The price of tolerance: Wolf damage

payments after recovery. Biodivers. Conserv. 18, 4003–4021.

Treves, A., Kapp, K.J., Macfarland, D.M., 2010. American black bear nuisance complaints and hunter

take. Ursus 21, 30–42.

Treves, A., Langenberg, J.A., López-Bao, J.V., Rabenhorst, M.F., 2017b. Gray wolf mortality patterns in

Wisconsin from 1979 to 2012. J. Mammal. 98, 17-32. 10.1093/jmammal/gyw145

Treves, A., Santiago-Ávila, F.J., Putrevu, K., 2021. Quantifying the effects of delisting wolves after the

first state began lethal management. PeerJ 9, e11666. 10.7717/peerj.11666

Van Ballenberghe, V., Mech, L., 1975. Weights, growth, and survival of timber wolf pups in Minnesota. J.

Mammal. 56, 44-63.

Van Valkenburgh, B., Ruff., C.B., 1987. Canine tooth strength and killing behaviour in large carnivores. J.

Zool. 212, 379-397.

von Essen, E., Hansen, H.P., Kallstrom, H.N., Peterson, M.N., Peterson, T.R., 2015. The radicalisation of

rural resistance: How hunting counterpublics in the Nordic countries contribute to illegal hunting.

Journal of Rural Studies 39, 199-209.

Willging, R., Wydeven, A.P., 1997. Cooperative wolf depredation management in Wisconsin, in: Lee,

C.D., Hygnstrom, S.E. (Eds.), Thirteenth Great Plains Wildlife Damage Control Workshop Proceedings.,

Kansas State University Agricultural Experiment Station and Cooperative Extension Service, Lincoln, NE,

pp. 46-51.

Wydeven, A.P., Treves, A., Brost, B., Wiedenhoeft, J.E., 2004. Characteristics of wolf packs in Wisconsin:

Identification of traits influencing depredation, in: Fascione, N., Delach, A., Smith, M.E. (Eds.), People

and Predators: From Conflict to Coexistence, Island Press, Washington, D. C., pp. 28-50.

Table 1. The number of hounds reported in wolf-hound interactions by breed and month

Breed	Jan	Feb	Mar	Apr	May	Jun	Jul	Aug	Sep	Oct	Nov	Dec
Bluetick	0	1	0	0	0	0	3	6	8	0	0	1
Plott	0	0	0	0	0	0	11	20	9	3	0	0
Redbone	0	1	0	0	0	0	4	4	1	0	0	1
Walker	0	0	0	1	0	0	16	19	16	3	1	5
Other	1	2	0	0	0	0	12	15	8	0	0	4
Totals	**1**	**4**	**0**	**1**	**0**	**0**	**46**	**64**	**42**	**6**	**1**	**11**

572
573

Table 2. Bystander reports (n=105) of 119 adverse interactions with hounds or their handlers across

counties of Wisconsin, USA. Because a respondent might report more than one interaction, we present

the counties from most reports 6) to fewest (1), the names of the 51 counties mentioned in reports, the

sum of interactions per row, and the maximum number of hounds reported in a single interaction.

When multiple reports were filed about the same interaction and the number of hounds differed, we

counted only one interaction and averaged the number of hounds, rounding to the higher integer.

Reports per county	Sum of the interactions	Maximum number of hounds in a single interaction	Counties with interactions reported
6	15	8	Bayfield, Iron, Sawyer
5	5	5	Forest
4	4	6	Langlade
3	13	6	Chippewa, Dane, Marathon, Polk, Washburn
2	18	6	Dodge, Douglas, Dunn, Florence, Lincoln, Oconto, Price, Shawano
1	62	8	Ashland, Barron, Brown, Burnett, Calumet, Cheboygan MI, Columbia, Cuyahoga OH, Door, Eau Claire, Fond du Lac, Gogebic, MI, Green, Houghton, MI, Jackson, Kenosha, Kewaunee, Macon, GA, Manitowoc, Marin, CA, Marinette, Milwaukee, Nash, NC, Oneida, Ontonagon, MI, Outagamie, Ozaukee, Rock, Rusk, St. Croix, Sheboygan, Taylor, Trempealeau, Vernon, Vilas, Walworth, Washington, Waukesha, Winnebago, Wood

SM 2 & 3 Survey & qualitative responses

Click here to access/download
Supplementary Material
SM 2 & 3 Survey & qualitative responses.docx

Click here to access/download
Supplementary Material
SM1 Hound-handlers attitudes to wolves.docx

Declaration of interests

1. Declaration of interests

☐ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☒The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

> AT and LM perceive no potential competing interests. All funding awards to AT as of 13 April 2022 are listed here http://faculty.nelson.wisc.edu/treves/archive_BAS/funding.pdf accessed 21 August 2022, which we provide for transparency and a CV for disclosure of potential competing interests: http://faculty.nelson.wisc.edu/treves/archive_BAS/Treves_vita_latest.pdf accessed 22 August 2022. LM is a retired Associate Professor from the Department of Literature, Leeward College, University of Hawaii and currently a member of: Endangered Species Coalition, Society for the Preservation of Poultry Antiquities, National Wolfwatchers Coalition, Wisconsin Wolf Trackers, Wolf Patrol, Executive Committee, Fox Valley Group, Sierra Club.

2. Contributions

☒ Each author declares substantial contributions through the following:
(1) the conception and design of the study, or acquisition of data, or analysis and interpretation of data,
(2) drafting the article or revising it critically for important intellectual content,

Please indicate for each author the author contributions in the text field below. Signatures are not required.

> AT and LM contributed data equally, AT wrote the manuscript, LM helped with proof-reading.

3. Approval of the submitted version of the manuscript

☒ Please check this box to confirm that all co-authors have read and approved the version of the manuscript that is submitted. Signatures are not required.