Proteomic analysis revealed that the oomyceticide phosphite exhibits multi-modal
action in an oomycete pathosystem.

Christina E. Andronis*[1,2], Silke Jacques[1], Francisco J. Lopez-Ruiz[1], Richard Lipscombe[2] and
Kar-Chun Tan[1*]

*Corresponding authors

Email: Christina.Andronis@postgrad.curtin.edu.au (ORCID: 0000-0003-4173-1232) and

Kar-Chun.Tan@curtin.edu.au (ORCID: 0000-0001-6094-823X).

Author affiliations

[1]Centre for Crop and Disease Management, Curtin University, Bentley, WA, Australia

[2]Proteomics International, Nedlands, WA, Australia

ABSTRACT

BACKGROUND: Phytopathogenic oomycetes constitute some of the most devastating plant
pathogens that cause significant crop and horticultural loss. *Phytophthora cinnamomi* is a
phytopathogenic oomycete that causes dieback disease in native vegetation and a variety of
crops. This pathogen can survive through harsh environmental conditions which gives it an
advantage over its susceptible hosts. The only implemented chemical used to control *P.*
cinnamomi is the oomyceticide phosphite. Despite its widespread use, the mode of action of
phosphite is not well understood and it is unclear whether it works directly on the pathogen
or through the host. Additionally, resistance to phosphite is emerging in *P. cinnamomi*
isolates and other oomycete phytopathogens.

RESULTS: The mode of action of phosphite on the pathogen and through a model host was
investigated using label-free quantitative proteomics. *In vitro* treatment of *P. cinnamomi* with
phosphite hinders growth by interfering with metabolism, signalling and gene expression,
traits that are not observed in the tolerant isolate. When the model host *L. angustifolius* was
treated with phosphite, enrichment of proteins that are associated with photosynthesis,
carbon fixation and lipid metabolism in the host was observed. An increase in the production
of a range of defence-related proteins was observed.

CONCLUSION: We hypothesise direct and indirect models of the multi-modal action of
phosphite that directly targets the pathogen as well as alters plant metabolism and immune
response.

Keywords

Phytophthora cinnamomi, oomycete, phosphite, resistance, mode of action, oomyceticide

1 INTRODUCTION

Phytopathogenic oomycetes are significant plant pathogens in natural ecosystems and

agriculture. These pathogens cause substantial environmental and economic losses from

plant death and management costs. *Phytophthora cinnamomi* is an oomycete that causes

dieback disease in native vegetation and several crops including avocado, macadamia,

pineapple, and a variety of stone fruits. This necrotic pathogen attacks the roots of

susceptible hosts and causes plant death via saprophytic growth [1]. The dynamic lifecycle of

this organism drives its resilience and success as a plant pathogen [2]. Similarly to other

Phytophthora spp, its spores enable it to survive through harsh environmental conditions and

thrive once conditions become more favourable. This gives the pathogen an advantage over

its susceptible hosts. Characteristically, oomycetes produce an assortment of virulence

molecules throughout their life cycles which facilitate the infection process [3,4].

Phosphite is the only commercially available chemical used to control *P. cinnamomi* [5].

Inorganic phosphites are reduced forms of phosphate that are commonly used as

oomyceticides to manage diseases caused by the *Phytophthora spp.* This includes *P.*

cinnamomi, P. nicotianae, P. palmivora, P. capsici and *P. infestans,* along with other

oomycetes such as downy mildews such as *Pseudoperonospora humuli* and *Bremia*

lactucae [6–10]. It is applied on native trees and horticultural crops by foliar spray or direct

injection into the trunk where it circulates throughout the plant system [11]. Phosphite cannot

eradicate *P. cinnamomi* in the field but it can be used to reduce the severity and spread of

the disease. It is also used as a preventative measure in uninfected areas [11–13]. The only

other strategies implemented to combat dieback disease are the use of tolerant root stocks if

they are available (such as commercially available avocado root stocks) and hygiene

measures applied to vehicles and personnel, which aim to minimise spread [14–17]. *In vitro*

fungicide screening of *P. cinnamomi* has shown growth inhibitory effects of chemical

alternatives such as metalaxyl, fosetyl-A1, benzethonium chloride and copper salts however,

these remain to be tested in the field [18,19]. As such, phosphite remains the only option for the

chemical control of *P. cinnamomi.*

Despite its widespread use in dieback management, the mode of action of phosphite is not

well understood. It is suggested that it acts both directly on the pathogen and indirectly

through the plant host by priming the plant immune system [7,8,20–22]. It has been reported that

mycelial growth and sporulation are inhibited by phosphite *in vitro* [20,21]. *In planta*, the

increase in transcription of defence-related genes associated with the salicylic and jasmonic

acid pathways in *Arabidopsis thaliana* and potato crops have been demonstrated post-

treatment with phosphite [7,8,22]. The effect of phosphite on potato leaves indicates an increase

in defence responses and altered metabolism, glycolysis, and carbon fixation [22]. These

findings have led to the hypothesis that phosphite primes the immune system of plants for

potential infection. Despite these observations, the mode of action of phosphite on the

pathogen has not been defined, nor have the mechanisms of phosphite primed plant

defence.

Tolerance of *P. cinnamomi* to phosphite is widespread with most reports originating from

horticultural plantations and variability in phosphite sensitivity has been demonstrated *in vitro*

and *in planta* [20,23]. Tolerance to phosphite has also been reported in *P. nicotianae, P.*

capsici, Bremia lactucae and *Pseudoperonospora humuli* [9,10,24–26]. This is a suspected result

of prolonged use of phosphite in agriculture and poses a significant threat to natural

ecosystems and the agricultural industry. These tolerant isolates are driving the pressure for

improved management strategies.

To elucidate the mode of action of phosphite on the pathogen and the host, we used a label-

free quantitative proteomic approach to conduct a study encompassing the effects of

phosphite on a sensitive and a tolerant isolate of *P. cinnamomi*, as well as investigating the

effects of phosphite on the physiology in a model plant system that is susceptible to dieback.

P. cinnamomi is used as model to understand phosphite tolerance as it is treated with

phosphite in the field and tolerant isolates have emerged. By using a shotgun approach, we

can obtain a snapshot of the biochemical processes that are altered as a result of treatment

with phosphite and determine whether phosphite exhibits a direct and/or indirect mode of

action.

2 MATERIALS AND METHODS

2.1 *In vitro* treatment of *P. cinnamomi* with phosphite

Stocks of two *P. cinnamomi* isolates, MU94-48 and CPSM366 were obtained from the

Centre of Phytophthora Science and Management, Perth, Australia. MU94-48 was collected

from susceptible *Eucalyptus marginata* in Willowdale, Western Australia with no history of

phosphite use and CPSM366 was collected from a Western Australian avocado orchard with

a history of extensive phosphite use and reduced efficacy of protection on dieback was

observed [23,27]. To determine the level of phosphite sensitivity of both isolates, they were

grown on Ribeiro's media amended with phosphite at concentrations ranging from 0.0 µg

mL^{-1} to 1000 µg mL^{-1} [28]. Mycelial cultures were incubated at room temperature in the dark.

After 14 days of growth, the mycelial radial growth was measured to determine the growth

inhibition. The EC_{50} and minimum inhibitory concentration (MIC) for both isolates were

calculated as previously described [29]. For proteomics analysis, growth inhibition of 0% and

20% were used as untreated and sub-lethal doses of phosphite to cause a physiological
effect and provide enough biomass for experimentation. Mycelia were harvested for protein
extraction by scraping from the surface of the plate, snap-frozen in liquid nitrogen and
freeze-dried. Lyophilised mycelia were ground using metal beads and a tissue mill (Retsch,
Haan, Germany) at a frequency of 3 Hz/s for three minutes.

2.2 *In planta* treatment of *Lupinus angustifolius* and inoculation with *P. cinnamomi*

The narrow leaf lupin *Lupinus angustifolius* (cv. 'Tanjil') seeds were surface sterilised with
5% sodium hypochlorite, washed twice with 70% ethanol and washed three times with water.
Seeds were placed in clear containers lined with moist Whatman paper (Cytiva,
Massachusetts, USA) and left to germinate for 3 days at room temperature under natural
light [30].

Three-day-old germinated seedlings were sprayed with a 0.5% pH 7 solution of phosphite
(Sigma, St Louis, USA) using a hand-triggered spray bottle and untreated seedlings were
sprayed with water [31–33]. Seedlings were incubated on a light shelf at room temperature with
a 12-hour photoperiod and seedlings were sprayed daily with water [30]. Lupin root tips were
harvested one day post first treatment. 1.5 cm of root tips were excised and immediately
snap frozen, freeze dry, and ground to a fine powder using metal beads and a tissue mill
(Retsch, Haan, Germany) at a frequency of 3 Hz/s for three minutes. Three biological
replicates (of which three root tips were collected) were used for each sample.

To test the effect of phosphite on the colonisation ability of *P. cinnamomi,* 24 hours post-
spraying, the infected seedlings were inoculated with 5 mm mycelial discs of each isolate.
Containers were placed back on the light shelf for three days and lesion scores from zero to
three were recorded.

2.3 Protein extraction and digestion

300 µl of extraction buffer (25 mM Tris-HCl pH 7.5, 0.25% SDS, 50 mM Na_2PO_4, 1 mM
Na_2F, 50 µM Na_3VO_4, 1 mM phenylmethylsulfonyl fluoride and a protease inhibitor cocktail)
was added to the ground mycelia and the samples were kept on ice for 30 minutes with
regular gentle mixing. The crude extract was centrifuged at 20,000 g at 4 °C for 30 minutes,
the solubilised proteins were decanted, and proteins were precipitated using six volumes of
ice-cold acetone and incubated at -20 °C overnight [34].

300 µl of extraction buffer (125 mM Tris-HCl pH 7.0, 7 % SDS, 0.5 % PVP-40) was added to
the whole ground plant material and the samples were kept on ice for 30 minutes with
regular gentle mixing. The crude extract was centrifuged at 20,000 g at 4°C for 30 minutes.
Proteins were purified by adding 800 µL of ice-cold methanol and 200 µL of ice-cold

chloroform to 200 μL of the solubilised protein samples. Samples were vortexed, and 500 μL

of water was added and centrifuged for 5 minutes at 15000g a 4 °C. The aqueous phase

was removed and 500 μL of methanol was added. Samples were inverted and the

supernatant was discarded. 1 mL of ice-cold acetone was added and the samples were

incubated at -20 °C overnight [35].

The mycelial and root protein pellets were washed twice with ice-cold acetone and

reconstituted in 200 μl of 0.5 M triethylammonium bicarbonate (pH 8.5) before reduction and

alkylation with 20 μL of 50 mM tris(2-carboxyethyl)phosphine (Thermo Scientific,

Massachusetts, USA) and 10 μL 200 mM methyl methanethiosulfonate. Samples were

tryptically digested overnight at 37 °C at a ratio of 1:10, subsequently desalted on a Strata-X

33 um polymeric reverse phase column (Phenomenex, Torrance, CA, USA) and dried in a

vacuum centrifuge [36].

2.4 Mass spectrometry and data analysis

1 μg of each sample was loaded on columns and peptides were resolved with a gradient of

10-40% acetonitrile (0.1% formic acid) at 300 nL/min over 90 minutes and eluted through a

nanospray interface into a Q-Exactive Orbitrap mass spectrometer (Thermofisher Scientific).

Qualitative and label-free quantification was performed using Proteome Discoverer 2.3 [37].

Mass spectra from the *in vitro* assay were matched to the *P. cinnamomi* MU94-48 genome

consisting of 26,151 protein-coding sequences [38]. The proteomes between phosphite-treated

MU94-48 and CPSM366 were compared to simultaneously gain insight into the effects of

phosphite on sensitive *P. cinnamomi* and understand the differences between phosphite

sensitivity and tolerance. The mass spectra obtained from the *in planta* assay were matched

to the *L. angustifolius* genome consisting of 39,339 protein-coding sequences [39]. For protein

identification, 1 or more 95% confidence peptides were used. For label-free quantification,

proteins with 2 or more 95% confidence peptides were used and for significant differential

abundance a p-value threshold of <0.05 was used. Ratios and p values for quantitative

analysis were generated using the default t-test hypothesis testing over biological replicates

[37].

To elucidate the functions of the detected proteins and understand the biochemical

differentiation between samples, Gene Ontology (GO), KEGG and Interpro were used [40–42].

For qualitative analysis, a Fisher's exact test was performed on assigned GOs to indicate

GO enrichment within samples compared to assigned ontologies of the whole genome

annotation in each respective organismp.

3 RESULTS

3.1 Growth inhibition of *P. cinnamomi* isolates by phosphite

To determine the sensitivity of *P. cinnamomi* isolates MU94-48 and CPSM366 to phosphite, both isolates were grown on phosphite-supplemented media, and sensitivity was determined based on the radial growth of hyphal colonies. Both isolates exhibited a similar radial growth rate in the absence of phosphite (Figure 1). The MIC for the sensitive and tolerant isolates were 500 µg/ mL and >1000 µg/ mL respectively. At concentrations of 1000 µg/ mL the growth of CPSM366 was reduced but not completely inhibited. The EC_{50} for the MU94-48 and CPSM366 were 10.8 µg/ mL and 415.6 µg/ mL respectively.

For proteomic analyses, 0.0 µg/ mL and 2.5 µg/ mL were used as these represent an untreated and sub-lethal dose of phosphite resulting in reduced growth of 20% and 4% for MU94-48 and CPSM366, respectively.

Figure 1. *In vitro* growth inhibition of MU94-48 and a CPSM366 in response to phosphite treatment indicating the MIC and EC_{50} for each isolate. Images of mycelial growth on the phosphite supplemented media are also displayed.

3.2 Phosphite induced significant alterations in the proteome of MU94-48 and CPSM366.

To determine the biochemical differences between MU94-48 and CPSM366 treated with
phosphite, the soluble intracellular proteome of each isolate was compared by qualitative
and quantitative proteomic analyses. 1393 proteins were unique to MU94-48, 280 were
unique to the CPSM366 and 1572 were common between the two isolates (Figure 2). Of the
common proteins, 171 were higher in abundance in MU94-48 compared to CPSM699 and
90 were lower in abundance. 1311 were not differentially abundant. Overall, there was a
significantly higher number of proteins identified in MU94-48, with 42.9% only observed in
this isolate.

Figure 2. Protein identification in phosphite treated MU94-48 and CPSM366. **a)** Indicating
the total number of unique and common proteins between the sensitive and tolerant isolates.
b) Differential abundance of the common proteins by label-free quantification shows the
increased and decreased abundance of proteins in phosphite-treated MU94-48 compared to
phosphite-treated CPSM366. The number of proteins identified in all samples obtained for
the *in vitro* assay are shown in Supplementary material 1.

3.3 Phosphite induces a stress response in MU94-48.

GO enrichment analysis and differential protein abundance between the MU94-48 and
CPSM366 were used to elucidate biochemical changes from phosphite treatment (Figure 3).
In MU94-48, an enrichment of putative stress response proteins was observed including
glutathione S- transferases (GST), thioredoxins, peptidases, proteasomes and proteolytic
enzymes [43]. These may be indicators of programmed cell death or attempts of the pathogen
to stay alive. Putative stress proteins were not observed in CPSM366.

3.4 An increase in signalling in response to phosphite treatment

Enrichment of GOs that are associated with protein signalling including phosphorylation (eg.
ATPases, GTPases, PKs), and ubiquitination were observed [44] (Figure 3). GOs that are
associated with positive regulation of DNA damage were enriched in addition to tRNA
binding, DNA binding, tRNA methylation and mRNA splicing and processing. Inositol
signalling and phosphorylation were significantly lower in abundance in MU94-48 coinciding
with the enrichment of phosphatidylinositol binding in CPSM366. Positive regulation of
TORC1 signalling and Wnt transmembrane signalling GOs were enriched. This indicates
stress response signalling, increase in gene expression to compensate for the loss of
biological material, and signalling to coordinate cellular processes during stress conditions
[45,46].

3.5 More transporters in MU94-48 than CPSM366 when treated with phopshite.

A higher diversity of transporters was found in MU94-48, including transmembrane
transporters, and intracellular transporters, along with facilitators of transport such as COPI
vesicle coatomer proteins, COP9 signalosome, armadillo-like proteins and clathrin proteins
(Figure 3). MU94-48 may be using transporters to pump out phosphite or toxic biproducts of
phopshite metabolism as a means to facilitate the high level of signalling.

3.6 Phosphite alters mitochondrial respiration in MU94-48.

A significantly lower abundance of mitochondrial-associated ontologies were observed in
MU94-48 as indicated by mitochondrial respiratory chain, cytochrome C oxidase,
mitochondrial electron transport and mitochondrial ribosomal subunit GOs. This could be a
result of oxidative stress in MU94-48 or may suggest that similarly to other fungicides,
phosphite targets mitochondrial respiration [47,48]. The KEGG ontologies between phosphite-
treated MU94-48 and CPSM366 did not show changes in disctinct metabolic pathway
clusters but rather showed general metabolic disorder (Supplementary material 2). However,
the disordered KEGG orthologues in MU94-48 suggest that phosphite is exerting a cytotoxic
effect on the sensitive isolate.

We then examined the proteome of MU94-48 and CPSM66 compared to their respective
untreated controls to ensure that these observations were not artifacts of the isolates. It was
observed that the proteome profile of untreated and phosphite-treated MU94-48 and
CPSM366 are comparable to those found between MU94-48 (Supplementary material 3).
This includes the enrichment in putative stress proteins such as oxidoreductase activity,
intracellular signalling and protein activation, and gene expression-related ontologies.
Additionally, a significant reduction in inositol biosynthesis and signalling, mitochondrial
electron transport-related GOs were observed, reflecting the observations between the two
phosphite-treated isolates. Similarly, we compared the proteome of untreated and phosphite-
treated CPSM366 and found less biochemical responses than when MU94-48 was
compared to CPSM366. The full GOs, KEGG orthologues and gene functions for all *in vitro*
qualitative and quantitative comparisons are shown in Supplementary Material 4.

Figure 3. GO enrichment and differential abundance between MU94-48 and CPSM366

treated with phosphite. **a)** and **b)** show GO enrichment in the phosphite treated tolerant and

sensitive isolates respectively, where the P value is generated by GO enrichment and each

point is separated on a log10 scale generated by each GO count to the total GO count for

the sample set. **d)** The enriched gene ontologies in the sensitive isolate at P < 0.001. **d)**

Differential abundance of proteins between the sensitive and tolerant isolate. Ratios are

generated by the peptide signal of each protein in MU94-48 compared to CPSM366. P

values   depict   the   significance   of   the   differential   abundance.

3.7 Differential abundance of proteins between untreated and phosphite treated
Lupinus angustifolius.

We hypothesise that phosphite in the plant physiology by facilitating plant immunity to
prevent infection by phytopathogens such as *P. cinnamomi.* To investigate this, the
proteome of *L. angustifolius* treated with and without phosphite were compared. 498 proteins
were unique in the untreated Lupin sample and 1725 were unique in the phosphite-treated
Lupin (Figure 4). 245 proteins were common between the untreated and treated lupin
samples of which 46 were significantly higher in abundance when phosphite was applied
and 4 were significantly lower.

Figure 4. Protein identification between untreated and phosphite treated lupins. **a)** Indicating
the total number of unique and common proteins between the untreated and phosphite-
treated lupin. **b)** Differential abundance of the common proteins by label-free quantification
shows the increased and decreased abundance of proteins in phosphite-treated lupin
compared to the untreated control. The number of proteins identified in all samples obtained
for the *in planta* assay are shown in Supplementary material 1.

3.8 Phosphite increased the abundance of photosynthetic and carbon fixation proteins in *L. angustifolius*.

GO enrichment and KEGG pathway enrichment analyses were used to examine biochemical effects in lupin treated with phosphite, (Figure 5 and 6). It was observed that GOs containing proteins that are associated with photosynthesis and starch metabolism were significantly enriched in phosphite-treated lupins. Photosynthesis, photosystem units, geranylgeranyl reductase, chlorophyll biosynthesis, carbon fixation and TCA cycle were all enriched in phosphite-treated lupins. Clustering of KEGG ontologies only found in the phosphite-treated lupin mapped to photorespiration, photosynthesis, reductive pentose phosphate cycle, isoprenoid biosynthesis and carbon fixation (Figure 6). As lupin seedlings were grown under artificial light with roots exposed leading to the development of photosynthetic-capable tissues, these were excised as part of the excised root tip for protein extraction (Figure 7).

Proteins associated with glucose, starch, and lipid metabolism GOs were enriched in the phosphite-treated lupin along with related KEGGs such as starch and sucrose metabolism, beta-oxidation and fatty elongation and lactosylceramide. Gluconeogenesis was reduced in abundance which supports the utilisation of carbohydrates and sugars as an energy source to fuel the heightened metabolism. Similarly, KEGG orthologues that are associated with lipid metabolism were also observed as indicated by the clustering of identified KEGG pathways only found in the phosphite-treated lupin (Figure 6). Phosphite treatment results in increased abundance of proteins that are associated with photosynthesis, carbon metabolism and energy production in lupin.

Figure 5. GO enrichment and differential abundance between untreated and phosphite-treated lupin. **a)** and **b)** show GO enrichment in the untreated and phosphite-treated lupin respectively, where the P value is generated by GO enrichment and each point is separated on a log10 scale generated by each GO count to the total GO count for the sample set. **c)** The enriched gene ontologies in the phosphite-treated lupin at P < 0.001. **d)** differential abundance of proteins between the untreated and treated lupin. Ratios are generated by the peptide signal of treated lupin compared to the untreated control. P values depict the significance of the differential abundance.

3.9 Transcriptional activities in phosphite-treated *L. angustifolius* are enhanced.

As a complement to the abundance of photosynthetic and metabolic gene ontologies, gene expression was also overrepresented. Translation, ribosomal proteins, DNA topological change, and RNA processing were all enriched in the phosphite treatment (Figure 5). This is also shown by the clustering of KEGG pathways in the phosphite-treated lupin including pyrimidine biosynthesis and keratan sulfate degradation (Figure 6). Hence, an increase in gene expression and biosynthesis accompanies massive physiological and metabolic changes.

3.10 Phosphite triggers the accumulation of defence-related proteins in lupin

Phosphite has previously been reported to increase the production of defence-related molecules in plants, particularly salicylic and jasmonic acid [7,8,49]. An enrichment of defence-related GOs was observed in the phosphite-treated lupin. These included proteolysis, oxidoreductase activity, hydrolases, cellular oxidant detoxification and response to oxidative stress (Figure 5). Secretory peroxidases, superoxide metabolism, abscisic acid signalling, and defence response GOs encompassing genes such as steroid chaperones, programmed cell death and apoptosis were also enriched. Several secondary metabolites were enriched including nicotinamidase and isochorismatase were also found only in the phosphite-treated lupin.

Precursors to the salicylic acid (SA) pathway including isochorismatase and actin depolymerisation GOs were enriched in the phosphite-treated lupin. KEGG analysis revealed components associated with the jasmonic acid (JA) pathway were only found in the phosphite-treated lupin. This shows that more defence-related proteins including secondary metabolites were enriched and higher in abundance in the phosphite-treated lupin.

Legend: KEGG Module | Significantly enriched in phosphite treated lupin

Pathways labelled in figure: Lactosylceramide biosynthesis, Keratan sulfate degradation, Jasmonic acid biosynthesis, Starch and sucrose metabolism, Pyrimidine biosynthesis, Beta-oxidation/fatty acid elongation, Photorespiration, Isoprenoid biosynthesis, Carbon fixation, Reductive pentose phosphate cycle, Photosynthesis

Figure 6. A representative set of the KEGG pathways enriched in the phosphite-treated lupin. Each dot represents a KEGG entry, and each line represents individual KEGG ontology identifiers. Pathway sets from overrepresented modules are labelled.

3.11 The effect of phosphite on MU94-48 and CPSM366 isolates during host infection.

To test the effect of phosphite during host infection*, L. angustifolius* was treated with phosphite and subsequently inoculated with the sensitive or tolerant isolate of *P. cinnamomi* (Figure 7). Untreated lupin infected with MU94-48 and CPSM366 developed comparable lesions. When phosphite-treated *L. angustifolius* was infected with MU94-38 no lesion was observed showing sensitivity to phosphite *in planta.* When inoculated with CPSM366, there was no reduction in lesion score compared to the untreated lupin.

Figure 7. Lesion scores of lupin roots when treated with phosphite and infected with the two

isolates of *P. cinnamomi*. Lesion scores were taken on day 3.

4 DISCUSSION

The mode of action of phosphite in *Pythophthora* pathosystems is a central question in the

development of future management strategies. As resistance to the only effective chemical

for the control of *P. cinnamomi* and other ooomycetes emerges, the pressure to find

resistance genes or alternative management strategies mounts. An understanding of the

mode of action of phosphite could aid in the development of better oomyceticides, which in

turn would contribute to improving existing management strategies. Previous literature has

suggested both direct and indirect mechanisms of phosphite on the pathogen and its host

plants however the biochemical mechanisms in which these occur have not been defined.

Our study aims to deconvolute the biochemistry of this system and gain a clearer insight into

the pathways altered by phosphite. *L. angustifolius* (lupin) is used as a model as it is

susceptible to *P. cinnamomi* infection, it has a published genome sequence required for

proteomics work, and it grows rapidly compared to native and horticultural *P. cinnamomi*

hosts [39,50].

The *in vitro* growth assay demonstrates that phosphite has a direct inhibitory effect on

mycelial growth in *P. cinnamomi*. The EC_{50} of previously reported phosphite-sensitive *P.*

cinnamomi isolates ranges between 4 µg mL^{-1} and 25 µg mL^{-1} [20,51]. The EC_{50} MU94-48 falls

within this range and can be considered highly sensitive to phosphite. Reported tolerant

isolates of *P. cinnamomi* have EC_{50} concentrations up to 150 µg mL^{-1} and other species of

Phytophthora have EC_{50} values up to 350 µg mL^{-1} , of which CMSP366 exceeds [20,51–53]. *P.*

citrophthor and *P. syringae* isolates screened for sensitivity to phosphite were considered

sensitive when their EC_{50} values were below 25 µg mL^{-1}), whereas *P. nicotianae* isolates

with EC_{50} values above 75 µg mL^{-1} were considered moderately resistant or resistant [24].

CPSM366 is therefore considered as highly resistant to phosphite, highlighting the need for

understanding its biochemistry and the development of alternative management strategies.

A proteomic approach was taken to understand how phosphite interacts with the two *P.*

cinnamomi isolates MU94-48 and CPSM366 and causes growth reduction. The data

obtained from this work was used to build a model suggesting the possible mechanisms in

which phosphite directly affects the pathogen (Figure 8). In the presence of phosphite, an

abundance of putative stress proteins is observed in the sensitive isolate (Figure 8a). The

pathogen is trying to find ways to cope with the influx of phosphite by expressing GSTs and

thioredoxins [54]. Hydrolase, proteolysis and peptidase proteins could also be produced by the

pathogen in this case for nutrient recycling, detoxification of xenobiotics, or could be

products of cell death caused by phosphite [43,55]. Putative stress response proteins were only

observed in MU94-48 indicating that CPSM366 does not undergo abiotic stress when

treated with phosphite.

Signalling molecules related to stress response were also induced along with the disruption

of the regulation of cellular processes caused by the exposure of MU94-48 to phosphite

(Figure 8b). Several core regulators of cell growth, metabolism and signalling were altered

as a result of phosphite treatment. For example, TORCI signalling was enriched in MU94-

48, which is involved in aspects of cell growth and metabolism [46]. This indicates that the

pathogen attempts to cope with disrupted cell growth and metabolism caused by phosphite.

Positive response to regulation of DNA damage and increase in gene expression-related

ontologies shows that phosphite could cause DNA damage and subsequent increase in

gene expression to compensate for the loss of DNA, proteins and cell material through

biosynthesis (Figure 8d) [56]. Canonical Wnt proteins are core regulators of development,

another biochemical pathway that is altered by phosphite. Inositol signalling and

phosphorylation, which play important roles in aspects of cell growth, structure and

signalling, were significantly lower in abundance in MU94-48 (Figure 8e). These

mechanisms of coping with extensive damage do not seem to be effective enough to protect

the sensitive isolate from damage by phosphite.

Greater diversity of membrane transporters were observed in the proteome of MU94-48 than

in CPSM366 (Figure 8c). These transporters include ABC and MSF-type, which have been

described to participate in the removal of xenobiotics from the cell [57,58]. Additionally,

facilitators of intracellular transport were also enriched in MU94-48 suggesting that, unlike
CPSM366, MU94-48 attempts to remove the toxic xenobiotic.

Compared to the tolerant isolate CPSM366, several constituents of the mitochondrial
respiratory pathway were reduced in abundance in the sensitive isolate (Figure 8f). In fungal
pathogens, programmed cell death can be triggered by mitochondrial-initiated signalling that
can be activated by cell damage, exposure to toxic xenobiotics and oxidative stress [59].
During these stress conditions, reactive oxygen species can disrupt components of the
mitochondrial respiration chain causing subsequent cell death [60–62]. As this is a core pathway
for cellular function, mitochondrial respiration is commonly used as a target for chemical
control of fungal phytopathogens by single-site fungicides [48,63]. For example, strobilurins are
a broad-spectrum class of fungicide applied to control fungal crop diseases. Strobilurin binds
cytochrome b complex III, inhibiting mitochondrial respiration [64]. Similarly, azoxystrobin
blocks electron transport in mitochondrial respiration by blocking electron transport [64,65,66].
Evidence presented here might suggest a broader inhibitory activity in the case of phosphite
as it seems to be affecting the pathogen's metabolism at multiple levels. Many fungicides
have multi-site modes of action, where several biochemical processes are disrupted such as
chlorothalonil, folpet, thiram, sulphur and copper [67,68].

The metabolic, signalling, regulatory and stress responses were not observed in CPSM366.
If phosphite acts as a multi-site oomyceticide, CPSM366 has likely developed physiological
adaptation to tolerate phosphite or alteration in target sites [69,70].

Figure 8- A proposed model on the direct effect of phosphite on *P. cinnamomi* MU94-48
from the combination of qualitative and quantitative comparisons between MU94-48 and

421 CPSM366. Black arrows indicate a possible cascading effect of core biochemical pathways
422 that are disrupted as a result of phosphite treatment. The size of bubbles represents
423 enrichment relative to each other for visualisation.

424 It has been suggested that phosphite alters the plant system to better cope with potential
425 attacks from oomycete pathogens by priming the plant immune system [7,8]. In addition,
426 phosphite has been reported to act as a biostimulant in plants [71]. A proteomic approach was
427 used to obtain a detailed biochemical snapshot of the proteins differentially abundant in
428 phosphite-treated *L. angustifolius.* The proteomic data obtained from the *in planta* assay was
429 used to generate a model that describes how phosphite alters plant metabolism and induces
430 an increase in defence-related proteins (Figure 9).

431 The biochemical differentiation between phosphite treated and untreated *L. angustifolius*
432 suggests an increase in metabolic activity. Constituents of the photosynthetic process,
433 carbon fixation and citric acid cycle suggest that phosphite is driving metabolism in plants
434 (Figure 9a). In addition, an increase in lipid and carbohydrate metabolism was observed,
435 indicating that stored carbohydrates can be metabolised for energy generation (Figure 9b).
436 The reductive pentose phosphate cycle was observed in the KEGG ontologies of the
437 phosphite-treated lupin and generates NADPH that feeds back into glycolysis [72].
438 Gluconeogenesis is significantly reduced in the phosphite-treated lupin highlighting that
439 sugars are utilised for energy production in this state, not stored. This is reinforced by
440 enrichment in gene expression to keep up with the metabolism and to cope with the increase
441 in photosynthesis through the process of biosynthesis (Figure 9c).

442 An increased metabolism has been observed in other phosphite-treated plants compared to
443 untreated controls along with its biostimulant effect [22,71,73,74]. The application of phosphite
444 improves yield, biomass, fruiting and growth, with a field trial applying phosphite in avocado
445 plantations showing a significant increase in fruit production [75,76]. In potato, phosphite
446 application resulted in reduced seedling emergence time, increased leaf size and biomass
447 [77]. This 'greening effect' has been demonstrated amongst other fungicides in agriculture,
448 where yield, biomass, leaf surface area, and protein content are increased as a result of
449 fungicide application [78–80]. The response observed in the phosphite-treated lupin seedlings is
450 similar to the greening effect, where phosphite boosts the metabolism of the plant. Non-leaf
451 plant structures such as roots, stems, flowers and seeds have photosynthetic potential when
452 exposed to light and carbon fixation has been reported to occur in many plant roots along
453 with green roots [81–83].

454 Defence-related gene expression has been demonstrated in several phosphite-treated
455 crops. In potato, phosphite treatment caused a significant increase in transcription of salicylic

and jasmonic acid [8]. Proteomic analysis of a similar system showed an increase in the
abundance of peroxidases, glutathione S-transferase and proteinase inhibitors [84]. In the
present study, enrichment of defence and stress-related proteins in *L. angustifolius* were
detected as a result of phosphite treatment, which may act in favour of the defence response
of the plant.

Secretory peroxidases were also enriched in the phosphite-treated lupin which can oxidise
toxic compounds and have functional roles in defence and biosynthesis [85]. Superoxide
metabolism and phospholipid binding composed of annexin genes were also enriched.
Superoxide dismutase is used in plant defence against reactive oxygen species [86,87].
Isoprenoids identified in the phosphite-treated lupin are not only carriers in photosynthetic
and respiratory electron transport and have additional functions as antioxidants [88]. Defence-
related ontologies with gene functions related to steroid chaperoning indicating programmed
cell death and apoptosis were also abundant [89]. Signalling and binding of abscisic acid, a
key hormone involved in signalling during stress and defence response to abiotic and
pathogens was significantly higher in abundance in the phosphite-treated lupin.
Nicotinamidase and isochorismatase are involved in plant growth, hydrolase activity and
synthesis of salicylic acid [90–93] JA pathway is associated with defence response against
necrotrophic microbial pathogens and abiotic stresses and the SA pathway is involved with
biotic stressors, cell death and hypersensitive responses in plants [94,95].

Figure 9- A proposed model on the effects of phosphite on *L. angustifolius* from the
combination of qualitative and quantitative proteome data between the untreated and
treated lupin.

In this system, phosphite is driving both metabolism and defence by directly promoting their
biochemical pathways [8]. The production of stress molecules such as abscisic acid can at the
same time be used as signalling molecules to trigger plant immunity [96]. The production of
defence and stress-related molecules in the plant could also be acting in response to
phosphite as a xenobiotic substance [97,98]. In the field, the ideal use of phosphite part of a
preventative strategy for the management of *P. cinnamomi* infection [11–13]. The *in planta*
assay demonstrates that phosphite application does not reduce the observed lesion when
lupin plants are infected with the tolerant isolate. If the elevated defence response impacts
the colonisation ability of *P. cinnamomi,* it is not observed in this system. Potentially the
induction of defense molecules by phosphite in this system may not have reached a
sufficient amplitude for host resistance.

5 CONCLUSION

Our data present a new perspective on the mode of action of phosphite. We have provided
evidence to propose models of the direct and indirect mode of action of phosphite. This data
demonstrates a comprehensive snapshot of the metabolic dysregulation in the sensitive
isolate when treated with phosphite, suggesting it is a substrate for one or more core
biochemical pathways. The tolerant isolate is likely to have adapted to block phosphite from
entering the cell, or if phosphite targets constituents of mitochondrial respiration, adapted to
block this interaction. Our model proposes an alternative avenue of plant responses to
phosphite, where phosphite drives multiple biochemical pathways and as a bi-product, more
defence-related proteins are produced. We proposed probable mechanisms based on
proteomic data that indicates a bi-modal mode of action of phosphite on both the pathogen
and the host plant. Further studies are required to determine systemic changes in phosphite-
treated *L. angustifolius* such as changes in photosynthesis and respiration in leaves to
confirm the current observations. Additionally, the accumulation of phosphite in the plant
tissue and subsequent uptake by the pathogen cannot be excluded. The outcome of this
study presented opportunities for functional validation to determine the phosphite mode of
action in crop protection.

Availability of data and materials

Spectral   data   used   for   this   study   are   available   at   Figshare
(DOI:10.6084/m9.figshare.20026214).

Acknowledgements

Proteomics International provided funding for the project. Curtin University provided funding
for sample preparation through the postgraduate maintenance fund. KCT, FLR and SJ are

supported by the Centre for Crop and Disease Management, a joint initiative of Curtin
University and the Grains Research and Development Corporation (CUR00023). We thank
Dr Lars Kamphuis from the Centre for Crop and Disease Management and CSIRO for
providing the seed stocks. We also thank Johannes Debler for his technical input for
functional annotation. We thank Prof. Giles Hardy and Dr Bill Dunstan from the Centre of
Phytophthora and Science Management at Murdoch University for providing the *P.*
cinnamomi isolates. We also thank Mr Leon Lenzo for his critical feedback on the manuscript
and Dr Paula Moolhuijzen for her input into bioinformatic visualisation.

Supplementary material 1- Protein identification of *in vitro phosphite* treated MU94-48 and
CPSM366 treated with phosphite and of the phosphite treated lupin roots. FDR was set to
<1%.

Supplementary material 2- Th KEGG pathways identified from the protenis only found in
MU94-48. Green lines indicate the KEGG pathways assigned to the unique proteins in
phopshite treated MU94-48 compared to the untreated control. There are no clear clusters of
pathways in this sample indicating that phosphite affects multiple processes in the sensitive
isolate.

Supplementary material 3- Representative GOs obtained from phosphite treated MU94-48
and CPSM366 compared to their respective untreated controls.. **a)** Enriched GOs in the
proteins uniquely identidied in untreated MU94-48 compared to treated MU94-48. **b)** GOs
associates with the differentially abundant proteins between untreated and phopshite treated
MU94-48. **c)** Enriched GOs in the proteins uniquely identidied in treated MU94-48 compared
to tnreated MU94-48. **d)** Enriched GOs in the proteins uniquely identidied in untreated
CPSM366 compared to treated CPSM366. **e)** GOs associates with the differentially
abundant proteins between untreated and phopshite treated CPSM366. **f)** Enriched GOs in
the proteins uniquely identidied in treated MU94-48 compared to tnreated MU94-48. Green
bubbles in **a), c), d)** and **f)** indicate significantly enriched GOs, green bubbles in **b)** and **e)**
indicated significantly higher in abundance in the treated samples and red bubbles indicate
significantly lower abundance in the treated samples compared to untreated samples.

Supplementary material 4- The full list of GOs, Interpro and KEGGs for the qualitative and
quantitative *in vitro* data set.

Supplementary material 5- The full list of GOs, Interpro and KEGGs for the qualitative and
quantitative *in planta* data set.

References

1   Hardham AR, Blackman LM. Phytophthora cinnamomi. Mol Plant Pathol **19**:260–285
(2018).

2   Hardham AR. Pathogen Profile: Phytophthora cinnamomi. Mol Plant Pathol **6**:589–
604 (2005).

3   Sonah H, Deshmukh RK, Bélanger RR. Computational prediction of effector proteins
in fungi: Opportunities and challenges. Front Plant Sci **7**:126 (2016).

4   Girard V, Dieryckx C, Job C, Job D. Secretomes: The fungal strike force. Proteomics
13:597–608 (2013).

5   Hardy GESJ, Barrett S, Shearer BL. The future of phosphite as a fungicide to control
the soilborne plant pathogen Phytophthora cinnamomi in natural ecosystems.
Australas Plant Pathol **30**:133–139 (2001).

6   Smillie R. The Mode of Action of Phosphite: Evidence for Both Direct and Indirect
Modes of Action on Three Phytophthora spp. in Plants. Phytopathology **79**:921–926
(1989).

7   Eshraghi L, Anderson J, Aryamanesh N, Shearer B, Mccomb J, … O'Brien PA.
Phosphite primed defence responses and enhanced expression of defence genes in
Arabidopsis thaliana infected with Phytophthora cinnamomi. Plant Pathol **60**:1086–
1095 (2011).

8   Machinandiarena MF, Lobato MC, Feldman ML, Daleo GR, Andreu AB. Potassium
phosphite primes defense responses in potato against Phytophthora infestans. J Plant
Physiol **169**:1417–1424 (2012).

9   Brown S, Koike ST, Ochoa OE, Laemmlen F, Michelmore RW. Insensitivity to the
Fungicide Fosetyl-Aluminum in California Isolates of the Lettuce Downy Mildew
Pathogen, Bremia lactucae. Plant Dis **88**:502–508 (2004).

10   Nelson ME, Eastwell KC, Grove GG, Barbour JD, Ocamb CM, Alldredge JR.
Sensitivity of Pseudoperonospora humuli (the Causal Agent of Hop Downy Mildew)
from Oregon, Idaho, and Washington to Fosetyl-Al (Aliette). doi:10.1094/PHP-2004-
0811-01-RShttps://doi.org/101094/PHP-2004-0811-01-RS **5** (2018).

11   Masikane SL, Novela P, Mohale P, McLeod A. Effect of phosphonate application
timing and -strategy on phosphite fruit and root residues of avocado. Crop Prot
128:105008 (2020).

12   Grobler M, Meyer J, Liebenberg J. Phosphite treatments as part of early preventative
control of Phytophthora root rot, pre- and post-planting. South African avocado Grow
Assoc Yearb **39**:30–35 (2016).

13   Ramallo AC, Cerioni L, Olmedo GM, Volentini SI, Ramallo J, Rapisarda VA. Control of
Phytophthora brown rot of lemons by pre- and postharvest applications of potassium
phosphite. Eur J Plant Pathol **154**:975–982 (2019).

14   Shearer BL, Crane CE, Barrett S, Cochrane A. Phytophthora cinnamomi invasion, a
major threatening process to conservation of flora diversity in the South-west
Botanical Province of Western Australia. Aust J Bot. **55**:225–238 (2007).

15   Cahill D, Rookes J, Wilson B, Gibson L, Mcdougall K. Phytophthora cinnamomi and
Australia ' s biodiversity : impacts , predictions and progress towards control. Aust J
Bot **56**:279–310 (2008).

16   Smith LA, Dann EK, Pegg KG, Whiley AW, Giblin FR, … Kopittke R. Field
assessment of avocado rootstock selections for resistance to phytophthora root rot.
Australas Plant Pathol **40**:39–47 (2011).

17   Kellam MK, Coffey MD. Quantitative Comparison of the Resistance to Phytophthora
Root Rot in Three Avocado Rootstocks. Phytopathology **75**:230–234 (1985).

18   González M, Caetano P, Sánchez ME. Testing systemic fungicides for control of
Phytophthora oak root disease. For Pathol **47**:e12343 (2017).

19   Lawrence SA, Armstrong CB, Patrick WM, Gerth ML. High-throughput chemical
screening identifies compounds that inhibit different stages of the Phytophthora
agathidicida and Phytophthora cinnamomi life cycles. Front Microbiol **8**:1340 (2017).

20   Wilkinson CJ, Shearer BL, Jackson TJ, Hardy GESJ. Variation in sensitivity of
Western Australian isolates of Phytophthora cinnamomi to phosphite in vitro. Plant
Pathol **50**:83–89 (2001).

21   Wong MH, McComb J, Hardy GESJ, Brien PAO. Phosphite induces expression of a
putative proteophosphoglycan gene in Phytophthora cinnamomi. Australas Plant
Pathol **38**:235–241 (2009).

22   Lim S, Borza T, Peters RD, Coffin RH, Al-Mughrabi KI, … Wang-Pruski G. Proteomics
analysis suggests broad functional changes in potato leaves triggered by phosphites
and a complex indirect mode of action against Phytophthora infestans. J Proteomics
93:207–223 (2013).

23   Dobrowolski MP, Shearer BL, Colquhoun IJ, O'Brien PA, Hardy GESJ. Selection for
decreased sensitivity to phosphite in Phytophthora cinnamomi with prolonged use of
fungicide. Plant Pathol **57**:928–936 (2008).

24   Hao W, Forster H, Adaskaveg JE. Resistance to potassium phosphite in phytophthora
species causing citrus brown rot and integrated practices for management of resistant
isolates. Plant Dis **105**:972–977 (2021).

25   Veena SS, Anandaraj M, Sarma Y. Variability in the sensitivity of Phytophthora capsici
isolates to potassium phosphonate. | Semantic Scholar. Indian Phytopathol **63**:71–75
(2010).

26   Gent DH, Block M, Claassen BJ. High levels of insensitivity to phosphonate fungicides
in Pseudoperonospora humuli. Plant Dis **104**:1400–1406 (2020).

27   Simamora A V., Stukely MJC, Barber PA, Hardy GESJ, Burgess TI. Age-related
susceptibility of Eucalyptus species to Phytophthora boodjera. Plant Pathol **66**:501–
512 (2017).

28   Ribeiro OK. A Source Book of the Genus Phytophthora. Mycologia **71**:74–77 (1979).

29   Lóopez-Ruiz FJ, Pérez-García A, Fernández-Ortuño D, Romero D, García E, …
Torés JA. Sensitivities to DMI fungicides in populations of Podosphaera fusca in south
central Spain. Pest Manag Sci **66**:801–808 (2010).

30   Smillie R, Grant BR, Guest D. The Mode of Action of Phosphite: Evidence for Both
Direct and Indirect Modes of Action on Three Phytophthora spp. in Plants.
Phytopathology **79**:921–926 (1989).

31   Pilbeam R a, Colquhoun IJ, Shearer B, Hardy GES. Phosphite concentration: its
effect on phytotoxicity symptoms and colonisation by Phytophthora cinnamomi in
three understorey species of Eucalyptus marginata forest. Australas Plant Pathol
29:86–95 (2000).

32   Cerioni L, Rapisarda VA. Use of Phosphite Salts in Laboratory and Semicommercial
Tests to Control Citrus Postharvest Decay. Plant Dis **97**:201–212 (2013).

33   Barrett SR, Shearer BL, Hardy GESJ. The efficacy of phosphite a pp lied after
inoculation on the colonisation of Banksia brownii sterns by Phytophthora cinnamomi.
Australas Plant Pathol 2003 321 **32**:1–7 (2003).

34   Bianco L, Perrotta G. Methodologies and perspectives of proteomics applied to
filamentous fungi: From sample preparation to secretome analysis. Int J Mol Sci
16:5803–5829 (2015).

35   Wessel D, Flügge UI. A method for the quantitative recovery of protein in dilute
solution in the presence of detergents and lipids. Anal Biochem **138**:141–143 (1984).

36   Casey TM, Khan JM, Bringans SD, Koudelka T, Takle PS, … Lipscombe RJ. Analysis
of Reproducibility of Proteome Coverage and Quantitation Using Isobaric Mass Tags
(iTRAQ and TMT). J Proteome Res **16**:384–392 (2017).

37   Orsburn BC. Proteome Discoverer—A Community Enhanced Data Processing Suite
for Protein Informatics. Proteomes **9**:15 (2021).

38   Andronis CE, Hane JK, Bringans S, S Hardy GE, Jacques S, Tan K-C. Gene
validation and remodelling using proteogenomics of Phytophthora cinnamomi, the
causal agent of Dieback. Front Microbiol **12**:1519 (2021).

39   Garg G, Kamphuis LG, Bayer PE, Kaur P, Dudchenko O, … Singh KB. A pan-genome
and chromosome-length reference genome of narrow-leafed lupin (Lupinus
angustifolius) reveals genomic diversity and insights into key industry and biological
traits. doi:10.1111/TPJ.15885Plant J (2022).

40   Zdobnov EM, Apweiler R. InterProScan – an integration platform for the signature-
recognition methods in InterPro. Bioinformatics **17**:847–848 (2001).

41   Harris MA, Clark J, Ireland A, Lomax J, Ashburner M, … White R. The Gene Ontology
(GO) database and informatics resource. Nucleic Acids Res **32**:D258–D261 (2004).

42   Kanehisa M, Goto S. KEGG: Kyoto Encyclopedia of Genes and Genomes. Nucleic
Acids Res **28**:27–30 (2000).

43   Meister AMEA. Glutathione. Annu Rev Biochem **52**:711–760 (1983).

44   Lengeler KB, Davidson RC, D'souza C, Harashima T, Shen W-C, … Heitman J.
Signal Transduction Cascades Regulating Fungal Development and Virulence.
Microbiol Mol Biol Rev **64**:746 (2000).

45   Niehrs C. The complex world of WNT receptor signalling. Nat Rev Mol Cell Biol 2012
1312 **13**:767–779 (2012).

46   Yan G, Lai Y, Jiang Y. TOR under stress: Targeting TORC1 by Rho1 GTPase. Cell
Cycle **11**:3384 (2012).

47   Li Q, Harvey LM, McNeil B. Oxidative stress in industrial fungi. Crit Rev Biotechnol
29:199–213 (2009).

48   Li D, Calderone R. Exploiting mitochondria as targets for the development of new
antifungals. Virulence **8**:159 (2017).

49   Eshraghi L, Anderson JP, Aryamanesh N, McComb JA, Shearer B, Giles GE.

Defence Signalling Pathways Involved in Plant Resistance and Phosphite-Mediated
Control of Phytophthora Cinnamomi. Plant Mol Biol Report **32**:342–356 (2014).

50   Qiu R, Qu D, Hardy GESJ, Trengove R, Agarwal M, Ren Y. Optimization of
headspace solid-phase microextraction conditions for the identification of
Phytophthora cinnamomi rands. Plant Dis **98**:1088–1098 (2014).

51   Coffey MD, Bower LA. In vitro variability among isolates of eight Phytophthora species
in response to phosphorous acid. Phytopathology **6**:738–742 (1984).

52   Hunter S, McDougal R, Williams N, Scott P. Variability in phosphite sensitivity
observed within and between seven Phytophthora species. Australas Plant Pathol
51:273–279 (2022).

53   Hunter S, Williams N, McDougal R, Scott P, Garbelotto M. Evidence for rapid adaptive
evolution of tolerance to chemical treatments in Phytophthora species and its practical
implications. PLoS One **13**:e0208961 (2018).

54   Lenoir I, Fontaine J, Lounès-Hadj Sahraoui A. Arbuscular mycorrhizal fungal
responses to abiotic stresses: A review. Phytochemistry **123**:4–15 (2016).

55   Barata-Antunes C, Alves R, Talaia G, Casal M, Gerós H, … Paiva S. Endocytosis of
nutrient transporters in fungi: The ART of connecting signaling and trafficking. Comput
Struct Biotechnol J **19**:1713–1737 (2021).

56   Moye-Rowley WS. Regulation of the Transcriptional Response to Oxidative Stress in
Fungi: Similarities and Differences. Eukaryot Cell **2**:381 (2003).

57   Víglaš J, Olejníková P. An update on ABC transporters of filamentous fungi – from
physiological substrates to xenobiotics. Microbiol Res **246**:126684 (2021).

58   Sang H, Hulvey JP, Green R, Xu H, Im J, … Jung G. A xenobiotic detoxification
pathway through transcriptional regulation in filamentous fungi. MBio **9**:e00457-18
(2018).

59   Sharon A, Finkelstein A, Shlezinger N, Hatam I. Fungal apoptosis: function, genes
and gene function. FEMS Microbiol Rev **33**:833–854 (2009).

60   Calderone R, Li D, Traven A. System-level impact of mitochondria on fungal
virulence: to metabolism and beyond. FEMS Yeast Res **15**:27 (2015).

61   Breitenbach M, Weber M, Rinnerthaler M, Karl T, Breitenbach-Koller L. Oxidative
Stress in Fungi: Its Function in Signal Transduction, Interaction with Plant Hosts, and
Lignocellulose Degradation. Biomolecules **5**:318 (2015).

62   Oliver RP, Hewitt GH. Fungicides in Crop Protection. 2nd ed. Cabi, 2014.

63   Kim YS, Dixon EW, Vincelli P, Farman ML. Field Resistance to Strobilurin (QoI)
Fungicides in Pyricularia grisea Caused by Mutations in the Mitochondrial
Cytochrome b Gene. http://dx.doi.org/101094/PHYTO2003937891 **93**:891–900
(2007).

64   Feng Y, Huang Y, Zhan H, Bhatt P, Chen S. An Overview of Strobilurin Fungicide
Degradation:Current Status and Future Perspective. Front Microbiol **11**:389 (2020).

65   Affourtit C, Heaney SP, Moore AL. Mitochondrial electron transfer in the wheat
pathogenic fungus Septoria tritici: on the role of alternative respiratory enzymes in
fungicide resistance. Biochim Biophys Acta - Bioenerg **1459**:291–298 (2000).

66   Fernández-Ortuño D, Tores JA, Pérez-García A. Mechanisms of resistance to QoI
fungicides in phytopathogenic fungi. Int Microbiol **11**:1–9 (2008).

67   Tillman RW, Siegel MR, Long JW. Mechanism of action and fate of the fungicide
chlorothalonil (2,4,5,6-tetrachloroisophthalonitrile) in biological systems: I. Reactions
with cells and subcellular components of Saccharomyces pastorianus. Pestic
Biochem Physiol **3**:160–167 (1973).

68   McCallan SEA. The nature of the fungicidal action of copper and sulfur. Bot Rev
15:629–643 (1949).

69   Corkley I, Fraaije B, Hawkins N. Fungicide resistance management: Maximizing the
effective life of plant protection products. Plant Pathol **71**:150–169 (2022).

70   Grimmer MK, van den Bosch F, Powers SJ, Paveley ND. Evaluation of a matrix to
calculate fungicide resistance risk. Pest Manag Sci **70**:1008–1016 (2014).

71   Achary VMM, Ram B, Manna M, Datta D, Bhatt A, … Agrawal PK. Phosphite: a novel
P fertilizer for weed management and pathogen control. Plant Biotechnol J **15**:1493–
1508 (2017).

72   Ramos-Martinez JI. The regulation of the pentose phosphate pathway: Remember
Krebs. Arch Biochem Biophys **614**:50–52 (2017).

73   Feldman ML, Guzzo MC, Machinandiarena MF, Rey-Burusco MF, Beligni MV, …
Andreu AB. New insights into the molecular basis of induced resistance triggered by
potassium phosphite in potato. Physiol Mol Plant Pathol **109**:101452 (2020).

74   Wu L, Gao X, Xia F, Joshi J, Borza T, Wang-Pruski G. Biostimulant and fungicidal
effects of phosphite assessed by GC-TOF-MS analysis of potato leaf metabolome.

Physiol Mol Plant Pathol **106**:49–56 (2019).

75   Gómez-Merino FC, Trejo-Téllez LI. Biostimulant activity of phosphite in horticulture.
Sci Hortic (Amsterdam) **196**:82–90 (2015).

76   Lovatt CJ. Properly Timing Foliar-applied Fertilizers Increases Efficacy: A Review and
Update on Timing Foliar Nutrient Applications to Citrus and Avocado. Horttechnology
23:536–541 (2013).

77   Tambascio C, Covacevich F, Lobato MC, Lasa C de, Caldiz D, … Andreu A. The
Application of K Phosphites to Seed Tubers Enhanced Emergence, Early Growth and
Mycorrhizal Colonization in Potato (Solanum tuberosum). Am J Plant Sci **5**:132–137
(2014).

78   Eiiner FM. Results of long-term field studies into the effect of strobilurin containing
fungicides on the production of mycotoxins in several winter wheat varieties.
Mycotoxin Res **21**:112–115 (2005).

79   Kyveryga PM, Blackmer TM, Mueller DS. When Do Foliar Pyraclostrobin Fungicide
Applications Produce Profitable Soybean Yield Responses? doi:10.1094/PHP-2013-
0928-01-RSPlant Heal Prog **14** (2013).

80   Spitzer T, Matušinsky P, Klemová Z, Kazda J. Effect of Fungicide Application Date
against Sclerotinia sclerotiorum on Yield and Greening of Winter Rape. Plant Prot Sci
48:105–109 (2012).

81   Aschan G, Pfanz H. Non-foliar photosynthesis – a strategy of additional carbon
acquisition. Flora - Morphol Distrib Funct Ecol Plants **198**:81–97 (2003).

82   Flores HE, Dai YR, Cuello JL, Maldonado-Mendoza IE, Loyola-Vargas VM. Green
Roots: Photosynthesis and Photoautotrophy in an Underground Plant Organ. Plant
Physiol **101**:363–371 (1993).

83   Henry RJ, Furtado A, Rangan P. Pathways of Photosynthesis in Non-Leaf Tissues.
Biol 2020, Vol 9, Page 438 **9**:438 (2020).

84   Lim S, Borza T, Peters RD, Coffin RH, Al-Mughrabi KI, … Wang-Pruski G. Proteomics
analysis suggests broad functional changes in potato leaves triggered by phosphites
and a complex indirect mode of action against Phytophthora infestans. J Proteomics
93:207–223 (2013).

85   Yoshida K, Kaothien P, Matsui T, Kawaoka A, Shinmyo A. Molecular biology and
application of plant peroxidase genes. Appl Microbiol Biotechnol **60**:665–670 (2003).

86　Yadav D, Boyidi P, Ahmed I, Kirti PB. Plant annexins and their involvement in stress
responses. Environ Exp Bot **155**:293–306 (2018).

87　Alscher RG, Erturk N, Heath LS. Role of superoxide dismutases (SODs) in controlling
oxidative stress in plants. J Exp Bot **53**:1331–1341 (2002).

88　Chappell J. Biochemistry and Molecular Biology of the Isoprenoid Biosynthetic
Pathway in Plants. http://dx.doi.org/101146/annurev.pp46060195002513 **46**:521–547
(2003).

89　Tang D, Ade J, Frye CA, Innes RW. Regulation of plant defense responses in
Arabidopsis by EDR2, a PH and START domain-containing protein. Plant J **44**:245–
257 (2005).

90　Hunt L, Holdsworth MJ, Gray JE. Nicotinamidase activity is important for germination.
Plant J **51**:341–351 (2007).

91　Wang G, Pichersky E. Nicotinamidase participates in the salvage pathway of NAD
biosynthesis in Arabidopsis. Plant J **49**:1020–1029 (2007).

92　Wildermuth MC, Dewdney J, Wu G, Ausubel FM. Isochorismate synthase is required
to synthesize salicylic acid for plant defence. Nature **414**:562–565 (2001).

93　Leontovyčová H, Kalachova T, Trdá L, Pospíchalová R, Lamparová L, … Janda M.
Actin depolymerization is able to increase plant resistance against pathogens via
activation of salicylic acid signalling pathway. Sci Reports 2019 91 **9**:1–10 (2019).

94　Ruan J, Zhou Y, Zhou M, Yan J, Khurshid M, … Zhang K. Jasmonic Acid Signaling
Pathway in Plants. Int J Mol Sci 2019, Vol 20, Page 2479 **20**:2479 (2019).

95　Ding P, Ding Y. Stories of Salicylic Acid: A Plant Defense Hormone. Trends Plant Sci
25:549–565 (2020).

96　Morgutti S, Negrini N, Pucciariello C, Sacchi GA. Role of trehalose and regulation of
its levels as a signal molecule to abiotic stresses in plants. doi:10.1016/B978-0-12-
816451-8.00014-9 Elsevier, 2019.

97　Ramel F, Sulmon C, Serra AA, Gouesbet G, Couée I. Xenobiotic sensing and
signalling in higher plants. J Exp Bot **63**:3999–4014 (2012).

98　Singh A, Sheo ·, Prasad M, Singh RP. Plant Responses to Xenobiotics. Springer,
2016.