Proteomic analysis revealed that the oomyceticide phosphite exhibits multi-modal action in an

oomycete pathosystem.

Christina E. Andronis*[12], Silke Jacques[1], Francisco J. Lopez-Ruiz[1], Richard Lipscombe[2] and Kar-Chun

Tan[1*]

*Corresponding authors

Email: Christina.Andronis@postgrad.curtin.edu.au (ORCID: 0000-0003-4173-1232) and

Kar-Chun.Tan@curtin.edu.au (ORCID: 0000-0001-6094-823X).

Author affiliations

[1]Centre for Crop and Disease Management, Curtin University, Bentley, WA, Australia

[2]Proteomics International, Nedlands, WA, Australia

ABSTRACT

BACKGROUND: Phytopathogenic oomycetes constitute some of the most devastating plant pathogens

that cause significant crop and horticultural loss. *Phytophthora cinnamomi* is a phytopathogenic

oomycete that causes dieback disease in native vegetation and a variety of crops. This pathogen can

survive through harsh environmental conditions which gives it an advantage over its susceptible hosts.

The only implemented chemical used to control *P. cinnamomi* is the oomyceticide phosphite. Despite

its widespread use, the mode of action of phosphite is not well understood and it is unclear whether it

works directly on the pathogen or through the host. Additionally, resistance to phosphite is emerging

in *P. cinnamomi* isolates and other oomycete phytopathogens.

RESULTS: The mode of action of phosphite on the pathogen and through a model host was investigated

using label-free quantitative proteomics. *In vitro* treatment of *P. cinnamomi* with phosphite hinders

growth by interfering with metabolism, signalling and gene expression, traits that are not observed in

the tolerant isolate. When the model host *L. angustifolius* was treated with phosphite, enrichment of

proteins that are associated with photosynthesis, carbon fixation and lipid metabolism in the host was

observed. An increase in the production of a range of defence-related proteins was observed.

CONCLUSION: We hypothesise direct and indirect models of the multi-modal action of phosphite that

directly targets the pathogen as well as alters plant metabolism and immune response.

Keywords

Phytophthora cinnamomi, oomycete, phosphite, resistance, mode of action, oomyceticide

1 INTRODUCTION

Phytopathogenic oomycetes are significant plant pathogens in natural ecosystems and agriculture. These pathogens cause substantial environmental and economic losses from plant death and management costs. *Phytophthora cinnamomi* is an oomycete that causes dieback disease in native vegetation and several crops including avocado, macadamia, pineapple, and a variety of stone fruits. This necrotic pathogen attacks the roots of susceptible hosts and causes plant death via saprophytic growth [1]. The dynamic lifecycle of this organism drives its resilience and success as a plant pathogen [2]. Similarly to other *Phytophthora spp,* its spores enable it to survive through harsh environmental conditions and thrive once conditions become more favourable. This gives the pathogen an advantage over its susceptible hosts. Characteristically, oomycetes produce an assortment of virulence molecules throughout their life cycles which facilitate the infection process [3,4].

Phosphite is the only commercially available chemical used to control *P. cinnamomi* [5]. Inorganic phosphites are reduced forms of phosphate that are commonly used as oomyceticides to manage diseases caused by the *Phytophthora spp.* This includes *P. cinnamomi, P. nicotianae, P. palmivora, P. capsici* and *P. infestans,* along with other oomycetes such as downy mildews such as *Pseudoperonospora humuli* and *Bremia lactucae* [6–10]. It is applied on native trees and horticultural crops by foliar spray or direct injection into the trunk where it circulates throughout the plant system [11]. Phosphite cannot eradicate *P. cinnamomi* in the field but it can be used to reduce the severity and spread of the disease. It is also used as a preventative measure in uninfected areas [11–13]. The only other strategies implemented to combat dieback disease are the use of tolerant root stocks if they are available (such as commercially available avocado root stocks) and hygiene measures applied to vehicles and personnel, which aim to minimise spread [14–17]. *In vitro* fungicide screening of *P. cinnamomi* has shown growth inhibitory effects of chemical alternatives such as metalaxyl, fosetyl-A1, benzethonium chloride and copper salts however, these remain to be tested in the field [18,19]. As such, phosphite remains the only option for the chemical control of *P. cinnamomi.*

Despite its widespread use in dieback management, the mode of action of phosphite is not well understood. It is suggested that it acts both directly on the pathogen and indirectly through the plant host by priming the plant immune system [7,8,20–22]. It has been reported that mycelial growth and sporulation are inhibited by phosphite *in vitro* [20,21]. *In planta,* the increase in transcription of defence-related genes associated with the salicylic and jasmonic acid pathways in *Arabidopsis thaliana* and potato crops have been demonstrated post-treatment with phosphite [7,8,22]. The effect of phosphite on potato leaves indicates an increase in defence responses and altered metabolism, glycolysis, and carbon fixation [22]. These findings have led to the hypothesis that phosphite primes the immune system

of plants for potential infection. Despite these observations, the mode of action of phosphite on the

pathogen has not been defined, nor have the mechanisms of phosphite primed plant defence.

Tolerance of *P. cinnamomi* to phosphite is widespread with most reports originating from horticultural

plantations and variability in phosphite sensitivity has been demonstrated *in vitro* and *in planta* [20,23].

Tolerance to phosphite has also been reported in *P. nicotianae, P. capsici, Bremia lactucae* and

Pseudoperonospora humuli [9,10,24–26]. This is a suspected result of prolonged use of phosphite in

agriculture and poses a significant threat to natural ecosystems and the agricultural industry. These

tolerant isolates are driving the pressure for improved management strategies.

To elucidate the mode of action of phosphite on the pathogen and the host, we used a label-free

quantitative proteomic approach to conduct a study encompassing the effects of phosphite on a

sensitive and a tolerant isolate of *P. cinnamomi*, as well as investigating the effects of phosphite on the

physiology in a model plant system that is susceptible to dieback. *P. cinnamomi* is used as model to

understand phosphite tolerance as it is treated with phosphite in the field and tolerant isolates have

emerged. By using a shotgun approach, we can obtain a snapshot of the biochemical processes that are

altered as a result of treatment with phosphite and determine whether phosphite exhibits a direct

and/or indirect mode of action.

2 MATERIALS AND METHODS

2.1 *In vitro* treatment of *P. cinnamomi* with phosphite

Stocks of two *P. cinnamomi* isolates, MU94-48 and CPSM366 were obtained from the Centre of

Phytophthora Science and Management, Perth, Australia. MU94-48 was collected from susceptible

Eucalyptus marginata in Willowdale, Western Australia with no history of phosphite use and CPSM366

was collected from a Western Australian avocado orchard with a history of extensive phosphite use and

reduced efficacy of protection on dieback was observed [23,27]. To determine the level of phosphite

sensitivity of both isolates, they were grown on Ribeiro's media amended with phosphite at

concentrations ranging from 0.0 µg mL^{-1} to 1000 µg mL^{-1} [28]. Mycelial cultures were incubated at room

temperature in the dark. After 14 days of growth, the mycelial radial growth was measured to

determine the growth inhibition. The EC$_{50}$ and minimum inhibitory concentration (MIC) for both isolates

were calculated as previously described [29]. For proteomics analysis, growth inhibition of 0% and 20%

were used as untreated and sub-lethal doses of phosphite to cause a physiological effect and provide

enough biomass for experimentation. Mycelia were harvested for protein extraction by scraping from

the surface of the plate, snap-frozen in liquid nitrogen and freeze-dried. Lyophilised mycelia were

ground using metal beads and a tissue mill (Retsch, Haan, Germany) at a frequency of 3 Hz/s for three

minutes.

2.2 *In planta* treatment of *Lupinus angustifolius* and inoculation with *P. cinnamomi*

The narrow leaf lupin *Lupinus angustifolius* (cv. 'Tanjil') seeds were surface sterilised with 5% sodium

hypochlorite, washed twice with 70% ethanol and washed three times with water. Seeds were placed

in clear containers lined with moist Whatman paper (Cytiva, Massachusetts, USA) and left to germinate

for 3 days at room temperature under natural light [30].

Three-day-old germinated seedlings were sprayed with a 0.5% pH 7 solution of phosphite (Sigma, St

Louis, USA) using a hand-triggered spray bottle and untreated seedlings were sprayed with water [31–33].

Seedlings were incubated on a light shelf at room temperature with a 12-hour photoperiod and

seedlings were sprayed daily with water [30]. Lupin root tips were harvested one day post first treatment.

1.5 cm of root tips were excised and immediately snap frozen, freeze dry, and ground to a fine powder

using metal beads and a tissue mill (Retsch, Haan, Germany) at a frequency of 3 Hz/s for three minutes.

Three biological replicates (of which three root tips were collected) were used for each sample.

To test the effect of phosphite on the colonisation ability of *P. cinnamomi,* 24 hours post- spraying, the

infected seedlings were inoculated with 5 mm mycelial discs of each isolate. Containers were placed

back on the light shelf for three days and lesion scores from zero to three were recorded.

2.3 Protein extraction and digestion

300 µl of extraction buffer (25 mM Tris-HCl pH 7.5, 0.25% SDS, 50 mM Na_2PO_4, 1 mM Na_2F, 50 µM

Na_3VO_4, 1 mM phenylmethylsulfonyl fluoride and a protease inhibitor cocktail) was added to the ground

mycelia and the samples were kept on ice for 30 minutes with regular gentle mixing. The crude extract

was centrifuged at 20,000 g at 4 °C for 30 minutes, the solubilised proteins were decanted, and proteins

were precipitated using six volumes of ice-cold acetone and incubated at -20 °C overnight [34].

300 µl of extraction buffer (125 mM Tris-HCl pH 7.0, 7 % SDS, 0.5 % PVP-40) was added to the whole

ground plant material and the samples were kept on ice for 30 minutes with regular gentle mixing. The

crude extract was centrifuged at 20,000 g at 4°C for 30 minutes. Proteins were purified by adding 800

µL of ice-cold methanol and 200 µL of ice-cold chloroform to 200 µL of the solubilised protein samples.

Samples were vortexed, and 500 µL of water was added and centrifuged for 5 minutes at 15000g a 4

°C. The aqueous phase was removed and 500 µL of methanol was added. Samples were inverted and

the supernatant was discarded. 1 mL of ice-cold acetone was added and the samples were incubated

at -20 °C overnight [35].

The mycelial and root protein pellets were washed twice with ice-cold acetone and reconstituted in
200 µl of 0.5 M triethylammonium bicarbonate (pH 8.5) before reduction and alkylation with 20 µL of
50 mM tris(2-carboxyethyl)phosphine (Thermo Scientific, Massachusetts, USA) and 10 µL 200 mM
methyl methanethiosulfonate. Samples were tryptically digested overnight at 37 °C at a ratio of 1:10,
subsequently desalted on a Strata-X 33 um polymeric reverse phase column (Phenomenex, Torrance,
CA, USA) and dried in a vacuum centrifuge [36].

2.4 Mass spectrometry and data analysis

1 µg of each sample was loaded on columns and peptides were resolved with a gradient of 10-40%
acetonitrile (0.1% formic acid) at 300 nL/min over 90 minutes and eluted through a nanospray interface
into a Q-Exactive Orbitrap mass spectrometer (Thermofisher Scientific). Qualitative and label-free
quantification was performed using Proteome Discoverer 2.3 [37]. Mass spectra from the *in vitro* assay
were matched to the *P. cinnamomi* MU94-48 genome consisting of 26,151 protein-coding sequences
[38]. The proteomes between phosphite-treated MU94-48 and CPSM366 were compared to
simultaneously gain insight into the effects of phosphite on sensitive *P. cinnamomi* and understand the
differences between phosphite sensitivity and tolerance. The mass spectra obtained from the *in planta*
assay were matched to the *L. angustifolius* genome consisting of 39,339 protein-coding sequences [39].
For protein identification, 1 or more 95% confidence peptides were used. For label-free quantification,
proteins with 2 or more 95% confidence peptides were used and for significant differential abundance
a p-value threshold of <0.05 was used. Ratios and p values for quantitative analysis were generated
using the default t-test hypothesis testing over biological replicates [37].

To elucidate the functions of the detected proteins and understand the biochemical differentiation
between samples, Gene Ontology (GO), KEGG and Interpro were used [40–42]. For qualitative analysis, a
Fisher's exact test was performed on assigned GOs to indicate GO enrichment within samples compared
to assigned ontologies of the whole genome annotation in each respective organismp.

3 RESULTS

3.1 Growth inhibition of *P. cinnamomi* isolates by phosphite

To determine the sensitivity of *P. cinnamomi* isolates MU94-48 and CPSM366 to phosphite, both
isolates were grown on phosphite-supplemented media, and sensitivity was determined based on the
radial growth of hyphal colonies. Both isolates exhibited a similar radial growth rate in the absence of
phosphite (Figure 1). The MIC for the sensitive and tolerant isolates were 500 µg/ mL and >1000 µg/
mL respectively. At concentrations of 1000 µg/ mL the growth of CPSM366 was reduced but not

completely inhibited. The EC_{50} for the MU94-48 and CPSM366 were 10.8 μg/ mL and 415.6 μg/ mL

respectively.

For proteomic analyses, 0.0 μg/ mL and 2.5 μg/ mL were used as these represent an untreated and sub-

lethal dose of phosphite resulting in reduced growth of 20% and 4% for MU94-48 and CPSM366,

respectively.

Figure 1. *In vitro* growth inhibition of MU94-48 and a CPSM366 in response to phosphite treatment

indicating the MIC and EC_{50} for each isolate. Images of mycelial growth on the phosphite supplemented

media are also displayed.

3.2 Phosphite induced significant alterations in the proteome of MU94-48 and CPSM366.

To determine the biochemical differences between MU94-48 and CPSM366 treated with phosphite,

the soluble intracellular proteome of each isolate was compared by qualitative and quantitative

proteomic analyses. 1393 proteins were unique to MU94-48, 280 were unique to the CPSM366 and

1572 were common between the two isolates (Figure 2). Of the common proteins, 171 were higher in

abundance in MU94-48 compared to CPSM699 and 90 were lower in abundance. 1311 were not

differentially abundant. Overall, there was a significantly higher number of proteins identified in MU94-

48, with 42.9% only observed in this isolate.

Figure 2. Protein identification in phosphite treated MU94-48 and CPSM366. **a)** Indicating the total

number of unique and common proteins between the sensitive and tolerant isolates. **b)** Differential

abundance of the common proteins by label-free quantification shows the increased and decreased

abundance of proteins in phosphite-treated MU94-48 compared to phosphite-treated CPSM366. The

number of proteins identified in all samples obtained for the *in vitro* assay are shown in Supplementary

material 1.

3.3 Phosphite induces a stress response in MU94-48.

GO enrichment analysis and differential protein abundance between the MU94-48 and CPSM366 were

used to elucidate biochemical changes from phosphite treatment (Figure 3). In MU94-48, an

enrichment of putative stress response proteins was observed including glutathione S- transferases

(GST), thioredoxins, peptidases, proteasomes and proteolytic enzymes [43]. These may be indicators of

programmed cell death or attempts of the pathogen to stay alive. Putative stress proteins were not

observed in CPSM366.

3.4 An increase in signalling in response to phosphite treatment

Enrichment of GOs that are associated with protein signalling including phosphorylation (eg. ATPases,

GTPases, PKs), and ubiquitination were observed [44] (Figure 3). GOs that are associated with positive

regulation of DNA damage were enriched in addition to tRNA binding, DNA binding, tRNA methylation

and mRNA splicing and processing. Inositol signalling and phosphorylation were significantly lower in

abundance in MU94-48 coinciding with the enrichment of phosphatidylinositol binding in CPSM366.

Positive regulation of TORC1 signalling and Wnt transmembrane signalling GOs were enriched. This

indicates stress response signalling, increase in gene expression to compensate for the loss of biological

material, and signalling to coordinate cellular processes during stress conditions [45,46].

3.5 More transporters in MU94-48 than CPSM366 when treated with phopshite.

A higher diversity of transporters was found in MU94-48, including transmembrane transporters, and

intracellular transporters, along with facilitators of transport such as COPI vesicle coatomer proteins,

COP9 signalosome, armadillo-like proteins and clathrin proteins (Figure 3). MU94-48 may be using

transporters to pump out phosphite or toxic biproducts of phopshite metabolism as a means to

facilitate the high level of signalling.

3.6 Phosphite alters mitochondrial respiration in MU94-48.

A significantly lower abundance of mitochondrial-associated ontologies were observed in MU94-48 as

indicated by mitochondrial respiratory chain, cytochrome C oxidase, mitochondrial electron transport

and mitochondrial ribosomal subunit GOs. This could be a result of oxidative stress in MU94-48 or may

suggest that similarly to other fungicides, phosphite targets mitochondrial respiration [47,48]. The KEGG

ontologies between phosphite-treated MU94-48 and CPSM366 did not show changes in disctinct

metabolic pathway clusters but rather showed general metabolic disorder (Supplementary material 2).

However, the disordered KEGG orthologues in MU94-48 suggest that phosphite is exerting a cytotoxic

effect on the sensitive isolate.

We then examined the proteome of MU94-48 and CPSM66 compared to their respective untreated

controls to ensure that these observations were not artifacts of the isolates. It was observed that the

proteome profile of untreated and phosphite-treated MU94-48 and CPSM366 are comparable to those

found between MU94-48 (Supplementary material 3). This includes the enrichment in putative stress

proteins such as oxidoreductase activity, intracellular signalling and protein activation, and gene

expression-related ontologies. Additionally, a significant reduction in inositol biosynthesis and

signalling, mitochondrial electron transport-related GOs were observed, reflecting the observations

between the two phosphite-treated isolates. Similarly, we compared the proteome of untreated and

phosphite-treated CPSM366 and found less biochemical responses than when MU94-48 was compared

to CPSM366. The full GOs, KEGG orthologues and gene functions for all *in vitro* qualitative and

quantitative comparisons are shown in Supplementary Material 4.

Figure 3. GO enrichment and differential abundance between MU94-48 and CPSM366 treated with phosphite. a) and b) show GO enrichment in the phosphite treated tolerant and sensitive isolates respectively, where the P value is generated by GO enrichment and each point is separated on a log10 scale generated by each GO count to the total GO count for the sample set. d) The enriched gene ontologies in the sensitive isolate at P < 0.001. d) Differential abundance of proteins between the sensitive and tolerant isolate. Ratios are generated by the peptide signal of each protein in MU94-48 compared to CPSM366. P values depict the significance of the differential abundance.

3.7 Differential abundance of proteins between untreated and phosphite treated *Lupinus angustifolius*.

We hypothesise that phosphite in the plant physiology by facilitating plant immunity to prevent

infection by phytopathogens such as *P. cinnamomi.* To investigate this, the proteome of *L. angustifolius*

treated with and without phosphite were compared. 498 proteins were unique in the untreated Lupin

sample and 1725 were unique in the phosphite-treated Lupin (Figure 4). 245 proteins were common

between the untreated and treated lupin samples of which 46 were significantly higher in abundance

when phosphite was applied and 4 were significantly lower.

Figure 4. Protein identification between untreated and phosphite treated lupins. **a)** Indicating the total

number of unique and common proteins between the untreated and phosphite-treated lupin. **b)**

Differential abundance of the common proteins by label-free quantification shows the increased and

decreased abundance of proteins in phosphite-treated lupin compared to the untreated control. The

number of proteins identified in all samples obtained for the *in planta* assay are shown in

Supplementary material 1.

3.8 Phosphite increased the abundance of photosynthetic and carbon fixation proteins in *L. angustifolius.*

GO enrichment and KEGG pathway enrichment analyses were used to examine biochemical effects in lupin treated with phosphite, (Figure 5 and 6). It was observed that GOs containing proteins that are associated with photosynthesis and starch metabolism were significantly enriched in phosphite-treated lupins. Photosynthesis, photosystem units, geranylgeranyl reductase, chlorophyll biosynthesis, carbon fixation and TCA cycle were all enriched in phosphite-treated lupins. Clustering of KEGG ontologies only found in the phosphite-treated lupin mapped to photorespiration, photosynthesis, reductive pentose phosphate cycle, isoprenoid biosynthesis and carbon fixation (Figure 6). As lupin seedlings were grown under artificial light with roots exposed leading to the development of photosynthetic-capable tissues, these were excised as part of the excised root tip for protein extraction (Figure 7).

Proteins associated with glucose, starch, and lipid metabolism GOs were enriched in the phosphite-treated lupin along with related KEGGs such as starch and sucrose metabolism, beta-oxidation and fatty elongation and lactosylceramide. Gluconeogenesis was reduced in abundance which supports the utilisation of carbohydrates and sugars as an energy source to fuel the heightened metabolism. Similarly, KEGG orthologues that are associated with lipid metabolism were also observed as indicated by the clustering of identified KEGG pathways only found in the phosphite-treated lupin (Figure 6). Phosphite treatment results in increased abundance of proteins that are associated with photosynthesis, carbon metabolism and energy production in lupin.

Figure 5. GO enrichment and differential abundance between untreated and phosphite-treated lupin. **a)** and **b)** show GO enrichment in the untreated and phosphite-treated lupin respectively, where the P value is generated by GO enrichment and each point is separated on a log10 scale generated by each GO count to the total GO count for the sample set. **c)** The enriched gene ontologies in the phosphite-treated lupin at P < 0.001. **d)** differential abundance of proteins between the untreated and treated lupin. Ratios are generated by the peptide signal of treated lupin compared to the untreated control. P values depict the significance of the differential abundance.

3.9 Transcriptional activities in phosphite-treated *L. angustifolius* are enhanced.

As a complement to the abundance of photosynthetic and metabolic gene ontologies, gene expression was also overrepresented. Translation, ribosomal proteins, DNA topological change, and RNA processing were all enriched in the phosphite treatment (Figure 5). This is also shown by the clustering of KEGG pathways in the phosphite-treated lupin including pyrimidine biosynthesis and keratan sulfate degradation (Figure 6). Hence, an increase in gene expression and biosynthesis accompanies massive physiological and metabolic changes.

3.10 Phosphite triggers the accumulation of defence-related proteins in lupin

Phosphite has previously been reported to increase the production of defence-related molecules in plants, particularly salicylic and jasmonic acid [7,8,49]. An enrichment of defence-related GOs was observed in the phosphite-treated lupin. These included proteolysis, oxidoreductase activity, hydrolases, cellular oxidant detoxification and response to oxidative stress (Figure 5). Secretory peroxidases, superoxide metabolism, abscisic acid signalling, and defence response GOs encompassing genes such as steroid chaperones, programmed cell death and apoptosis were also enriched. Several secondary metabolites were enriched including nicotinamidase and isochorismatase were also found only in the phosphite-treated lupin.

Precursors to the salicylic acid (SA) pathway including isochorismatase and actin depolymerisation GOs were enriched in the phosphite-treated lupin. KEGG analysis revealed components associated with the jasmonic acid (JA) pathway were only found in the phosphite-treated lupin. This shows that more defence-related proteins including secondary metabolites were enriched and higher in abundance in the phosphite-treated lupin.

Figure 6. A representative set of the KEGG pathways enriched in the phosphite-treated lupin. Each dot represents a KEGG entry, and each line represents individual KEGG ontology identifiers. Pathway sets from overrepresented modules are labelled.

3.11 The effect of phosphite on MU94-48 and CPSM366 isolates during host infection.

To test the effect of phosphite during host infection, *L. angustifolius* was treated with phosphite and subsequently inoculated with the sensitive or tolerant isolate of *P. cinnamomi* (Figure 7). Untreated lupin infected with MU94-48 and CPSM366 developed comparable lesions. When phosphite-treated *L. angustifolius* was infected with MU94-38 no lesion was observed showing sensitivity to phosphite *in planta*. When inoculated with CPSM366, there was no reduction in lesion score compared to the untreated lupin.

Figure 7. Lesion scores of lupin roots when treated with phosphite and infected with the two isolates

of *P. cinnamomi*. Lesion scores were taken on day 3.

4 DISCUSSION

The mode of action of phosphite in *Pythophthora* pathosystems is a central question in the development of future management strategies. As resistance to the only effective chemical for the control of *P. cinnamomi* and other ooomycetes emerges, the pressure to find resistance genes or alternative management strategies mounts. An understanding of the mode of action of phosphite could aid in the development of better oomyceticides, which in turn would contribute to improving existing management strategies. Previous literature has suggested both direct and indirect mechanisms of phosphite on the pathogen and its host plants however the biochemical mechanisms in which these occur have not been defined. Our study aims to deconvolute the biochemistry of this system and gain a clearer insight into the pathways altered by phosphite. *L. angustifolius* (lupin) is used as a model as it is susceptible to *P. cinnamomi* infection, it has a published genome sequence required for proteomics work, and it grows rapidly compared to native and horticultural *P. cinnamomi* hosts [39,50].

The *in vitro* growth assay demonstrates that phosphite has a direct inhibitory effect on mycelial growth in *P. cinnamomi*. The EC_{50} of previously reported phosphite-sensitive *P. cinnamomi* isolates ranges between 4 µg mL^{-1} and 25 µg mL^{-1} [20,51]. The EC_{50} MU94-48 falls within this range and can be considered highly sensitive to phosphite. Reported tolerant isolates of *P. cinnamomi* have EC_{50} concentrations up to 150 µg mL^{-1} and other species of *Phytophthora* have EC_{50} values up to 350 µg mL^{-1}, of which CMSP366 exceeds [20,51–53]. *P. citrophthor* and *P. syringae* isolates screened for sensitivity to phosphite

were considered sensitive when their EC_{50} values were below 25 µg mL^{-1}), whereas *P. nicotianae*

isolates with EC_{50} values above 75 µg mL^{-1} were considered moderately resistant or resistant [24].

CPSM366 is therefore considered as highly resistant to phosphite, highlighting the need for

understanding its biochemistry and the development of alternative management strategies.

A proteomic approach was taken to understand how phosphite interacts with the two *P. cinnamomi*

isolates MU94-48 and CPSM366 and causes growth reduction. The data obtained from this work was

used to build a model suggesting the possible mechanisms in which phosphite directly affects the

pathogen (Figure 8). In the presence of phosphite, an abundance of putative stress proteins is observed

in the sensitive isolate (Figure 8a). The pathogen is trying to find ways to cope with the influx of

phosphite by expressing GSTs and thioredoxins [54]. Hydrolase, proteolysis and peptidase proteins could

also be produced by the pathogen in this case for nutrient recycling, detoxification of xenobiotics, or

could be products of cell death caused by phosphite [43,55]. Putative stress response proteins were only

observed in MU94-48 indicating that CPSM366 does not undergo abiotic stress when treated with

phosphite.

Signalling molecules related to stress response were also induced along with the disruption of the

regulation of cellular processes caused by the exposure of MU94-48 to phosphite (Figure 8b). Several

core regulators of cell growth, metabolism and signalling were altered as a result of phosphite

treatment. For example, TORCI signalling was enriched in MU94-48, which is involved in aspects of cell

growth and metabolism [46]. This indicates that the pathogen attempts to cope with disrupted cell growth

and metabolism caused by phosphite. Positive response to regulation of DNA damage and increase in

gene expression-related ontologies shows that phosphite could cause DNA damage and subsequent

increase in gene expression to compensate for the loss of DNA, proteins and cell material through

biosynthesis (Figure 8d) [56]. Canonical Wnt proteins are core regulators of development, another

biochemical pathway that is altered by phosphite. Inositol signalling and phosphorylation, which play

important roles in aspects of cell growth, structure and signalling, were significantly lower in abundance

in MU94-48 (Figure 8e). These mechanisms of coping with extensive damage do not seem to be

effective enough to protect the sensitive isolate from damage by phosphite.

Greater diversity of membrane transporters were observed in the proteome of MU94-48 than in

CPSM366 (Figure 8c). These transporters include ABC and MSF-type, which have been described to

participate in the removal of xenobiotics from the cell [57,58]. Additionally, facilitators of intracellular

transport were also enriched in MU94-48 suggesting that, unlike CPSM366, MU94-48 attempts to

remove the toxic xenobiotic.

Compared to the tolerant isolate CPSM366, several constituents of the mitochondrial respiratory

pathway were reduced in abundance in the sensitive isolate (Figure 8f). In fungal pathogens,

programmed cell death can be triggered by mitochondrial-initiated signalling that can be activated by

cell damage, exposure to toxic xenobiotics and oxidative stress [59]. During these stress conditions,

reactive oxygen species can disrupt components of the mitochondrial respiration chain causing

subsequent cell death [60–62]. As this is a core pathway for cellular function, mitochondrial respiration is

commonly used as a target for chemical control of fungal phytopathogens by single-site fungicides [48,63].

For example, strobilurins are a broad-spectrum class of fungicide applied to control fungal crop

diseases. Strobilurin binds cytochrome b complex III, inhibiting mitochondrial respiration [64]. Similarly,

azoxystrobin blocks electron transport in mitochondrial respiration by blocking electron transport

[64,65,66]. Evidence presented here might suggest a broader inhibitory activity in the case of phosphite as

it seems to be affecting the pathogen's metabolism at multiple levels. Many fungicides have multi-

site modes of action, where several biochemical processes are disrupted such as chlorothalonil, folpet,

thiram, sulphur and copper [67,68].

The metabolic, signalling, regulatory and stress responses were not observed in CPSM366. If phosphite

acts as a multi-site oomyceticide, CPSM366 has likely developed physiological adaptation to tolerate

phosphite or alteration in target sites [69,70].

Figure 8- A proposed model on the direct effect of phosphite on *P. cinnamomi* MU94-48 from the

combination of qualitative and quantitative comparisons between MU94-48 and CPSM366. Black

arrows indicate a possible cascading effect of core biochemical pathways that are disrupted as a result

of phosphite treatment. The size of bubbles represents enrichment relative to each other for
visualisation.

It has been suggested that phosphite alters the plant system to better cope with potential attacks from
oomycete pathogens by priming the plant immune system [7,8]. In addition, phosphite has been reported
to act as a biostimulant in plants [71]. A proteomic approach was used to obtain a detailed biochemical
snapshot of the proteins differentially abundant in phosphite-treated *L. angustifolius.* The proteomic
data obtained from the *in planta* assay was used to generate a model that describes how phosphite
alters plant metabolism and induces an increase in defence-related proteins (Figure 9).

The biochemical differentiation between phosphite treated and untreated *L. angustifolius* suggests an
increase in metabolic activity. Constituents of the photosynthetic process, carbon fixation and citric
acid cycle suggest that phosphite is driving metabolism in plants (Figure 9a). In addition, an increase in
lipid and carbohydrate metabolism was observed, indicating that stored carbohydrates can be
metabolised for energy generation (Figure 9b). The reductive pentose phosphate cycle was observed
in the KEGG ontologies of the phosphite-treated lupin and generates NADPH that feeds back into
glycolysis [72]. Gluconeogenesis is significantly reduced in the phosphite-treated lupin highlighting that
sugars are utilised for energy production in this state, not stored. This is reinforced by enrichment in
gene expression to keep up with the metabolism and to cope with the increase in photosynthesis
through the process of biosynthesis (Figure 9c).

An increased metabolism has been observed in other phosphite-treated plants compared to untreated
controls along with its biostimulant effect [22,71,73,74]. The application of phosphite improves yield,
biomass, fruiting and growth, with a field trial applying phosphite in avocado plantations showing a
significant increase in fruit production [75,76]. In potato, phosphite application resulted in reduced
seedling emergence time, increased leaf size and biomass [77]. This 'greening effect' has been
demonstrated amongst other fungicides in agriculture, where yield, biomass, leaf surface area, and
protein content are increased as a result of fungicide application [78–80]. The response observed in the
phosphite-treated lupin seedlings is similar to the greening effect, where phosphite boosts the
metabolism of the plant. Non-leaf plant structures such as roots, stems, flowers and seeds have
photosynthetic potential when exposed to light and carbon fixation has been reported to occur in many
plant roots along with green roots [81–83].

Defence-related gene expression has been demonstrated in several phosphite-treated crops. In potato,
phosphite treatment caused a significant increase in transcription of salicylic and jasmonic acid [8].
Proteomic analysis of a similar system showed an increase in the abundance of peroxidases, glutathione
S-transferase and proteinase inhibitors [84]. In the present study, enrichment of defence and stress-

related proteins in *L. angustifolius* were detected as a result of phosphite treatment, which may act in
favour of the defence response of the plant.

Secretory peroxidases were also enriched in the phosphite-treated lupin which can oxidise toxic
compounds and have functional roles in defence and biosynthesis [85]. Superoxide metabolism and
phospholipid binding composed of annexin genes were also enriched. Superoxide dismutase is used in
plant defence against reactive oxygen species [86,87]. Isoprenoids identified in the phosphite-treated lupin
are not only carriers in photosynthetic and respiratory electron transport and have additional functions
as antioxidants [88]. Defence-related ontologies with gene functions related to steroid chaperoning
indicating programmed cell death and apoptosis were also abundant [89]. Signalling and binding of
abscisic acid, a key hormone involved in signalling during stress and defence response to abiotic and
pathogens was significantly higher in abundance in the phosphite-treated lupin. Nicotinamidase and
isochorismatase are involved in plant growth, hydrolase activity and synthesis of salicylic acid [90–93] JA
pathway is associated with defence response against necrotrophic microbial pathogens and abiotic
stresses and the SA pathway is involved with biotic stressors, cell death and hypersensitive responses
in plants [94,95].

Figure 9- A proposed model on the effects of phosphite on *L. angustifolius* from the combination of
qualitative and quantitative proteome data between the untreated and treated lupin.

In this system, phosphite is driving both metabolism and defence by directly promoting their
biochemical pathways [8]. The production of stress molecules such as abscisic acid can at the same time
be used as signalling molecules to trigger plant immunity [96]. The production of defence and stress-

related molecules in the plant could also be acting in response to phosphite as a xenobiotic substance

[97,98]. In the field, the ideal use of phosphite part of a preventative strategy for the management of *P.*

cinnamomi infection [11–13]. The *in planta* assay demonstrates that phosphite application does not reduce

the observed lesion when lupin plants are infected with the tolerant isolate. If the elevated defence

response impacts the colonisation ability of *P. cinnamomi,* it is not observed in this system. Potentially

the induction of defense molecules by phosphite in this system may not have reached a sufficient

amplitude for host resistance.

5 CONCLUSION

Our data present a new perspective on the mode of action of phosphite. We have provided evidence

to propose models of the direct and indirect mode of action of phosphite. This data demonstrates a

comprehensive snapshot of the metabolic dysregulation in the sensitive isolate when treated with

phosphite, suggesting it is a substrate for one or more core biochemical pathways. The tolerant isolate

is likely to have adapted to block phosphite from entering the cell, or if phosphite targets constituents

of mitochondrial respiration, adapted to block this interaction. Our model proposes an alternative

avenue of plant responses to phosphite, where phosphite drives multiple biochemical pathways and as

a bi-product, more defence-related proteins are produced. We proposed probable mechanisms based

on proteomic data that indicates a bi-modal mode of action of phosphite on both the pathogen and the

host plant. Further studies are required to determine systemic changes in phosphite-treated *L.*

angustifolius such as changes in photosynthesis and respiration in leaves to confirm the current

observations. Additionally, the accumulation of phosphite in the plant tissue and subsequent uptake by

the pathogen cannot be excluded. The outcome of this study presented opportunities for functional

validation to determine the phosphite mode of action in crop protection.

Availability of data and materials

Acknowledgements

Proteomics International provided funding for the project. Curtin University provided funding for

sample preparation through the postgraduate maintenance fund. KCT, FLR and SJ are supported by the

Centre for Crop and Disease Management, a joint initiative of Curtin University and the Grains Research

and Development Corporation (CUR00023). We thank Dr Lars Kamphuis from the Centre for Crop and

Disease Management and CSIRO for providing the seed stocks. We also thank Johannes Debler for his

technical input for functional annotation. We thank Prof. Giles Hardy and Dr Bill Dunstan from the

Centre of Phytophthora and Science Management at Murdoch University for providing the *P.*

cinnamomi isolates. We also thank Mr Leon Lenzo for his critical feedback on the manuscript and Dr

Paula Moolhuijzen for her input into bioinformatic visualisation.

Conflict of interest

The authors declare that the manuscript 'The oomyceticide phosphite exhibits potential multi-modal

action in an oomycete pathosystem' was undertaken with no conflicts of interest. No additional

financial support, affiliations, patents or copyrights are associated with this manuscript.

Supplementary material 1- Protein identification of *in vitro phosphite* treated MU94-48 and CPSM366
treated with phosphite and of the phosphite treated lupin roots. FDR was set to <1%.

Supplementary material 2- Th KEGG pathways identified from the protenis only found in MU94-48.

Green lines indicate the KEGG pathways assigned to the unique proteins in phopshite treated MU94-

48 compared to the untreated control. There are no clear clusters of pathways in this sample indicating

that phosphite affects multiple processes in the sensitive isolate.

Supplementary material 3- Representative GOs obtained from phosphite treated MU94-48 and

CPSM366 compared to their respective untreated controls.. **a)** Enriched GOs in the proteins uniquely

identidied in untreated MU94-48 compared to treated MU94-48. **b)** GOs associates with the

differentially abundant proteins between untreated and phopshite treated MU94-48. **c)** Enriched GOs

in the proteins uniquely identidied in treated MU94-48 compared to tnreated MU94-48. **d)** Enriched

GOs in the proteins uniquely identidied in untreated CPSM366 compared to treated CPSM366. **e)** GOs

associates with the differentially abundant proteins between untreated and phopshite treated

CPSM366. **f)** Enriched GOs in the proteins uniquely identidied in treated MU94-48 compared to

tnreated MU94-48. Green bubbles in **a), c), d)** and **f)** indicate significantly enriched GOs, green bubbles

in **b)** and **e)** indicated significantly higher in abundance in the treated samples and red bubbles indicate

significantly lower abundance in the treated samples compared to untreated samples.

Supplementary material 4- The full list of GOs, Interpro and KEGGs for the qualitative and quantitative
in vitro data set.

Supplementary material 5- The full list of GOs, Interpro and KEGGs for the qualitative and quantitative
in planta data set.

References

1   Hardham AR, Blackman LM. Phytophthora cinnamomi. Mol Plant Pathol **19**:260–285 (2018).

2   Hardham AR. Pathogen Profile: Phytophthora cinnamomi. Mol Plant Pathol **6**:589–604 (2005).

3   Sonah H, Deshmukh RK, Bélanger RR. Computational prediction of effector proteins in fungi:

Opportunities and challenges. Front Plant Sci **7**:126 (2016).

4   Girard V, Dieryckx C, Job C, Job D. Secretomes: The fungal strike force. Proteomics **13**:597–608
(2013).

5   Hardy GESJ, Barrett S, Shearer BL. The future of phosphite as a fungicide to control the
soilborne plant pathogen Phytophthora cinnamomi in natural ecosystems. Australas Plant
Pathol **30**:133–139 (2001).

6   Smillie R. The Mode of Action of Phosphite: Evidence for Both Direct and Indirect Modes of
Action on Three Phytophthora spp. in Plants. Phytopathology **79**:921–926 (1989).

7   Eshraghi L, Anderson J, Aryamanesh N, Shearer B, Mccomb J, … O'Brien PA. Phosphite primed
defence responses and enhanced expression of defence genes in Arabidopsis thaliana infected
with Phytophthora cinnamomi. Plant Pathol **60**:1086–1095 (2011).

8   Machinandiarena MF, Lobato MC, Feldman ML, Daleo GR, Andreu AB. Potassium phosphite
primes defense responses in potato against Phytophthora infestans. J Plant Physiol **169**:1417–
1424 (2012).

9   Brown S, Koike ST, Ochoa OE, Laemmlen F, Michelmore RW. Insensitivity to the Fungicide
Fosetyl-Aluminum in California Isolates of the Lettuce Downy Mildew Pathogen, Bremia
lactucae. Plant Dis **88**:502–508 (2004).

10 Nelson ME, Eastwell KC, Grove GG, Barbour JD, Ocamb CM, Alldredge JR. Sensitivity of
Pseudoperonospora humuli (the Causal Agent of Hop Downy Mildew) from Oregon, Idaho, and
Washington to Fosetyl-Al (Aliette). doi:10.1094/PHP-2004-0811-01-
RShttps://doi.org/101094/PHP-2004-0811-01-RS **5** (2018).

11 Masikane SL, Novela P, Mohale P, McLeod A. Effect of phosphonate application timing and -
strategy on phosphite fruit and root residues of avocado. Crop Prot **128**:105008 (2020).

12 Grobler M, Meyer J, Liebenberg J. Phosphite treatments as part of early preventative control
of Phytophthora root rot, pre- and post-planting. South African avocado Grow Assoc Yearb
39:30–35 (2016).

13 Ramallo AC, Cerioni L, Olmedo GM, Volentini SI, Ramallo J, Rapisarda VA. Control of
Phytophthora brown rot of lemons by pre- and postharvest applications of potassium
phosphite. Eur J Plant Pathol **154**:975–982 (2019).

14 Shearer BL, Crane CE, Barrett S, Cochrane A. Phytophthora cinnamomi invasion, a major

threatening process to conservation of flora diversity in the South-west Botanical Province of
Western Australia. Aust J Bot. **55**:225–238 (2007).

15 Cahill D, Rookes J, Wilson B, Gibson L, Mcdougall K. Phytophthora cinnamomi and Australia ' s
biodiversity : impacts , predictions and progress towards control. Aust J Bot **56**:279–310
(2008).

16 Smith LA, Dann EK, Pegg KG, Whiley AW, Giblin FR, … Kopittke R. Field assessment of avocado
rootstock selections for resistance to phytophthora root rot. Australas Plant Pathol **40**:39–47
(2011).

17 Kellam MK, Coffey MD. Quantitative Comparison of the Resistance to Phytophthora Root Rot
in Three Avocado Rootstocks. Phytopathology **75**:230–234 (1985).

18 González M, Caetano P, Sánchez ME. Testing systemic fungicides for control of Phytophthora
oak root disease. For Pathol **47**:e12343 (2017).

19 Lawrence SA, Armstrong CB, Patrick WM, Gerth ML. High-throughput chemical screening
identifies compounds that inhibit different stages of the Phytophthora agathidicida and
Phytophthora cinnamomi life cycles. Front Microbiol **8**:1340 (2017).

20 Wilkinson CJ, Shearer BL, Jackson TJ, Hardy GESJ. Variation in sensitivity of Western Australian
isolates of Phytophthora cinnamomi to phosphite in vitro. Plant Pathol **50**:83–89 (2001).

21 Wong MH, McComb J, Hardy GESJ, Brien PAO. Phosphite induces expression of a putative
proteophosphoglycan gene in Phytophthora cinnamomi. Australas Plant Pathol **38**:235–241
(2009).

22 Lim S, Borza T, Peters RD, Coffin RH, Al-Mughrabi KI, … Wang-Pruski G. Proteomics analysis
suggests broad functional changes in potato leaves triggered by phosphites and a complex
indirect mode of action against Phytophthora infestans. J Proteomics **93**:207–223 (2013).

23 Dobrowolski MP, Shearer BL, Colquhoun IJ, O'Brien PA, Hardy GESJ. Selection for decreased
sensitivity to phosphite in Phytophthora cinnamomi with prolonged use of fungicide. Plant
Pathol **57**:928–936 (2008).

24 Hao W, Forster H, Adaskaveg JE. Resistance to potassium phosphite in phytophthora species
causing citrus brown rot and integrated practices for management of resistant isolates. Plant
Dis **105**:972–977 (2021).

25 Veena SS, Anandaraj M, Sarma Y. Variability in the sensitivity of Phytophthora capsici isolates

to potassium phosphonate. | Semantic Scholar. Indian Phytopathol **63**:71–75 (2010).

26   Gent DH, Block M, Claassen BJ. High levels of insensitivity to phosphonate fungicides in
Pseudoperonospora humuli. Plant Dis **104**:1400–1406 (2020).

27   Simamora A V., Stukely MJC, Barber PA, Hardy GESJ, Burgess TI. Age-related susceptibility of
Eucalyptus species to Phytophthora boodjera. Plant Pathol **66**:501–512 (2017).

28   Ribeiro OK. A Source Book of the Genus Phytophthora. Mycologia **71**:74–77 (1979).

29   Lóopez-Ruiz FJ, Pérez-García A, Fernández-Ortuño D, Romero D, García E, … Torés JA.
Sensitivities to DMI fungicides in populations of Podosphaera fusca in south central Spain. Pest
Manag Sci **66**:801–808 (2010).

30   Smillie R, Grant BR, Guest D. The Mode of Action of Phosphite: Evidence for Both Direct and
Indirect Modes of Action on Three Phytophthora spp. in Plants. Phytopathology **79**:921–926
(1989).

31   Pilbeam R a, Colquhoun IJ, Shearer B, Hardy GES. Phosphite concentration: its effect on
phytotoxicity symptoms and colonisation by Phytophthora cinnamomi in three understorey
species of Eucalyptus marginata forest. Australas Plant Pathol **29**:86–95 (2000).

32   Cerioni L, Rapisarda VA. Use of Phosphite Salts in Laboratory and Semicommercial Tests to
Control Citrus Postharvest Decay. Plant Dis **97**:201–212 (2013).

33   Barrett SR, Shearer BL, Hardy GESJ. The efficacy of phosphite a pp lied after inoculation on the
colonisation of Banksia brownii sterns by Phytophthora cinnamomi. Australas Plant Pathol
2003 321 **32**:1–7 (2003).

34   Bianco L, Perrotta G. Methodologies and perspectives of proteomics applied to filamentous
fungi: From sample preparation to secretome analysis. Int J Mol Sci **16**:5803–5829 (2015).

35   Wessel D, Flügge UI. A method for the quantitative recovery of protein in dilute solution in the
presence of detergents and lipids. Anal Biochem **138**:141–143 (1984).

36   Casey TM, Khan JM, Bringans SD, Koudelka T, Takle PS, … Lipscombe RJ. Analysis of
Reproducibility of Proteome Coverage and Quantitation Using Isobaric Mass Tags (iTRAQ and
TMT). J Proteome Res **16**:384–392 (2017).

37   Orsburn BC. Proteome Discoverer—A Community Enhanced Data Processing Suite for Protein
Informatics. Proteomes **9**:15 (2021).

38   Andronis CE, Hane JK, Bringans S, S Hardy GE, Jacques S, Tan K-C. Gene validation and
remodelling using proteogenomics of Phytophthora cinnamomi, the causal agent of Dieback.
Front Microbiol **12**:1519 (2021).

39   Garg G, Kamphuis LG, Bayer PE, Kaur P, Dudchenko O, … Singh KB. A pan-genome and
chromosome-length reference genome of narrow-leafed lupin (Lupinus angustifolius) reveals
genomic diversity and insights into key industry and biological traits.
doi:10.1111/TPJ.15885Plant J (2022).

40   Zdobnov EM, Apweiler R. InterProScan – an integration platform for the signature-recognition
methods in InterPro. Bioinformatics **17**:847–848 (2001).

41   Harris MA, Clark J, Ireland A, Lomax J, Ashburner M, … White R. The Gene Ontology (GO)
database and informatics resource. Nucleic Acids Res **32**:D258–D261 (2004).

42   Kanehisa M, Goto S. KEGG: Kyoto Encyclopedia of Genes and Genomes. Nucleic Acids Res
28:27–30 (2000).

43   Meister AMEA. Glutathione. Annu Rev Biochem **52**:711–760 (1983).

44   Lengeler KB, Davidson RC, D'souza C, Harashima T, Shen W-C, … Heitman J. Signal Transduction
Cascades Regulating Fungal Development and Virulence. Microbiol Mol Biol Rev **64**:746
(2000).

45   Niehrs C. The complex world of WNT receptor signalling. Nat Rev Mol Cell Biol 2012 1312
13:767–779 (2012).

46   Yan G, Lai Y, Jiang Y. TOR under stress: Targeting TORC1 by Rho1 GTPase. Cell Cycle **11**:3384
(2012).

47   Li Q, Harvey LM, McNeil B. Oxidative stress in industrial fungi. Crit Rev Biotechnol **29**:199–213
(2009).

48   Li D, Calderone R. Exploiting mitochondria as targets for the development of new antifungals.
Virulence **8**:159 (2017).

49   Eshraghi L, Anderson JP, Aryamanesh N, McComb JA, Shearer B, Giles GE. Defence Signalling
Pathways Involved in Plant Resistance and Phosphite-Mediated Control of Phytophthora
Cinnamomi. Plant Mol Biol Report **32**:342–356 (2014).

50   Qiu R, Qu D, Hardy GESJ, Trengove R, Agarwal M, Ren Y. Optimization of headspace solid-
phase microextraction conditions for the identification of Phytophthora cinnamomi rands.

Plant Dis **98**:1088–1098 (2014).

51   Coffey MD, Bower LA. In vitro variability among isolates of eight Phytophthora species in
response to phosphorous acid. Phytopathology **6**:738–742 (1984).

52   Hunter S, McDougal R, Williams N, Scott P. Variability in phosphite sensitivity observed within
and between seven Phytophthora species. Australas Plant Pathol **51**:273–279 (2022).

53   Hunter S, Williams N, McDougal R, Scott P, Garbelotto M. Evidence for rapid adaptive
evolution of tolerance to chemical treatments in Phytophthora species and its practical
implications. PLoS One **13**:e0208961 (2018).

54   Lenoir I, Fontaine J, Lounès-Hadj Sahraoui A. Arbuscular mycorrhizal fungal responses to
abiotic stresses: A review. Phytochemistry **123**:4–15 (2016).

55   Barata-Antunes C, Alves R, Talaia G, Casal M, Gerós H, … Paiva S. Endocytosis of nutrient
transporters in fungi: The ART of connecting signaling and trafficking. Comput Struct
Biotechnol J **19**:1713–1737 (2021).

56   Moye-Rowley WS. Regulation of the Transcriptional Response to Oxidative Stress in Fungi:
Similarities and Differences. Eukaryot Cell **2**:381 (2003).

57   Víglaš J, Olejníková P. An update on ABC transporters of filamentous fungi – from physiological
substrates to xenobiotics. Microbiol Res **246**:126684 (2021).

58   Sang H, Hulvey JP, Green R, Xu H, Im J, … Jung G. A xenobiotic detoxification pathway through
transcriptional regulation in filamentous fungi. MBio **9**:e00457-18 (2018).

59   Sharon A, Finkelstein A, Shlezinger N, Hatam I. Fungal apoptosis: function, genes and gene
function. FEMS Microbiol Rev **33**:833–854 (2009).

60   Calderone R, Li D, Traven A. System-level impact of mitochondria on fungal virulence: to
metabolism and beyond. FEMS Yeast Res **15**:27 (2015).

61   Breitenbach M, Weber M, Rinnerthaler M, Karl T, Breitenbach-Koller L. Oxidative Stress in
Fungi: Its Function in Signal Transduction, Interaction with Plant Hosts, and Lignocellulose
Degradation. Biomolecules **5**:318 (2015).

62   Oliver RP, Hewitt GH. Fungicides in Crop Protection. 2nd ed. Cabi, 2014.

63   Kim YS, Dixon EW, Vincelli P, Farman ML. Field Resistance to Strobilurin (QoI) Fungicides in
Pyricularia grisea Caused by Mutations in the Mitochondrial Cytochrome b Gene.

http://dx.doi.org/101094/PHYTO2003937891 **93**:891–900 (2007).

64 Feng Y, Huang Y, Zhan H, Bhatt P, Chen S. An Overview of Strobilurin Fungicide
Degradation:Current Status and Future Perspective. Front Microbiol **11**:389 (2020).

65 Affourtit C, Heaney SP, Moore AL. Mitochondrial electron transfer in the wheat pathogenic
fungus Septoria tritici: on the role of alternative respiratory enzymes in fungicide resistance.
Biochim Biophys Acta - Bioenerg **1459**:291–298 (2000).

66 Fernández-Ortuño D, Tores JA, Pérez-García A. Mechanisms of resistance to QoI fungicides in
phytopathogenic fungi. Int Microbiol **11**:1–9 (2008).

67 Tillman RW, Siegel MR, Long JW. Mechanism of action and fate of the fungicide chlorothalonil
(2,4,5,6-tetrachloroisophthalonitrile) in biological systems: I. Reactions with cells and
subcellular components of Saccharomyces pastorianus. Pestic Biochem Physiol **3**:160–167
(1973).

68 McCallan SEA. The nature of the fungicidal action of copper and sulfur. Bot Rev **15**:629–643
(1949).

69 Corkley I, Fraaije B, Hawkins N. Fungicide resistance management: Maximizing the effective life
of plant protection products. Plant Pathol **71**:150–169 (2022).

70 Grimmer MK, van den Bosch F, Powers SJ, Paveley ND. Evaluation of a matrix to calculate
fungicide resistance risk. Pest Manag Sci **70**:1008–1016 (2014).

71 Achary VMM, Ram B, Manna M, Datta D, Bhatt A, … Agrawal PK. Phosphite: a novel P fertilizer
for weed management and pathogen control. Plant Biotechnol J **15**:1493–1508 (2017).

72 Ramos-Martinez JI. The regulation of the pentose phosphate pathway: Remember Krebs. Arch
Biochem Biophys **614**:50–52 (2017).

73 Feldman ML, Guzzo MC, Machinandiarena MF, Rey-Burusco MF, Beligni MV, … Andreu AB.
New insights into the molecular basis of induced resistance triggered by potassium phosphite
in potato. Physiol Mol Plant Pathol **109**:101452 (2020).

74 Wu L, Gao X, Xia F, Joshi J, Borza T, Wang-Pruski G. Biostimulant and fungicidal effects of
phosphite assessed by GC-TOF-MS analysis of potato leaf metabolome. Physiol Mol Plant
Pathol **106**:49–56 (2019).

75 Gómez-Merino FC, Trejo-Téllez LI. Biostimulant activity of phosphite in horticulture. Sci Hortic
(Amsterdam) **196**:82–90 (2015).

76   Lovatt CJ. Properly Timing Foliar-applied Fertilizers Increases Efficacy: A Review and Update on
Timing Foliar Nutrient Applications to Citrus and Avocado. Horttechnology **23**:536–541 (2013).

77   Tambascio C, Covacevich F, Lobato MC, Lasa C de, Caldiz D, … Andreu A. The Application of K
Phosphites to Seed Tubers Enhanced Emergence, Early Growth and Mycorrhizal Colonization
in Potato (Solanum tuberosum). Am J Plant Sci **5**:132–137 (2014).

78   Eiiner FM. Results of long-term field studies into the effect of strobilurin containing fungicides
on the production of mycotoxins in several winter wheat varieties. Mycotoxin Res **21**:112–115
(2005).

79   Kyveryga PM, Blackmer TM, Mueller DS. When Do Foliar Pyraclostrobin Fungicide Applications
Produce Profitable Soybean Yield Responses? doi:10.1094/PHP-2013-0928-01-RSPlant Heal
Prog **14** (2013).

80   Spitzer T, Matušinsky P, Klemová Z, Kazda J. Effect of Fungicide Application Date against
Sclerotinia sclerotiorum on Yield and Greening of Winter Rape. Plant Prot Sci **48**:105–109
(2012).

81   Aschan G, Pfanz H. Non-foliar photosynthesis – a strategy of additional carbon acquisition.
Flora - Morphol Distrib Funct Ecol Plants **198**:81–97 (2003).

82   Flores HE, Dai YR, Cuello JL, Maldonado-Mendoza IE, Loyola-Vargas VM. Green Roots:
Photosynthesis and Photoautotrophy in an Underground Plant Organ. Plant Physiol **101**:363–
371 (1993).

83   Henry RJ, Furtado A, Rangan P. Pathways of Photosynthesis in Non-Leaf Tissues. Biol 2020, Vol
9, Page 438 **9**:438 (2020).

84   Lim S, Borza T, Peters RD, Coffin RH, Al-Mughrabi KI, … Wang-Pruski G. Proteomics analysis
suggests broad functional changes in potato leaves triggered by phosphites and a complex
indirect mode of action against Phytophthora infestans. J Proteomics **93**:207–223 (2013).

85   Yoshida K, Kaothien P, Matsui T, Kawaoka A, Shinmyo A. Molecular biology and application of
plant peroxidase genes. Appl Microbiol Biotechnol **60**:665–670 (2003).

86   Yadav D, Boyidi P, Ahmed I, Kirti PB. Plant annexins and their involvement in stress responses.
Environ Exp Bot **155**:293–306 (2018).

87   Alscher RG, Erturk N, Heath LS. Role of superoxide dismutases (SODs) in controlling oxidative
stress in plants. J Exp Bot **53**:1331–1341 (2002).

88   Chappell J. Biochemistry and Molecular Biology of the Isoprenoid Biosynthetic Pathway in
Plants. http://dx.doi.org/101146/annurev.pp46060195002513 **46**:521–547 (2003).

89   Tang D, Ade J, Frye CA, Innes RW. Regulation of plant defense responses in Arabidopsis by
EDR2, a PH and START domain-containing protein. Plant J **44**:245–257 (2005).

90   Hunt L, Holdsworth MJ, Gray JE. Nicotinamidase activity is important for germination. Plant J
51:341–351 (2007).

91   Wang G, Pichersky E. Nicotinamidase participates in the salvage pathway of NAD biosynthesis
in Arabidopsis. Plant J **49**:1020–1029 (2007).

92   Wildermuth MC, Dewdney J, Wu G, Ausubel FM. Isochorismate synthase is required to
synthesize salicylic acid for plant defence. Nature **414**:562–565 (2001).

93   Leontovyčová H, Kalachova T, Trdá L, Pospíchalová R, Lamparová L, … Janda M. Actin
depolymerization is able to increase plant resistance against pathogens via activation of
salicylic acid signalling pathway. Sci Reports 2019 91 **9**:1–10 (2019).

94   Ruan J, Zhou Y, Zhou M, Yan J, Khurshid M, … Zhang K. Jasmonic Acid Signaling Pathway in
Plants. Int J Mol Sci 2019, Vol 20, Page 2479 **20**:2479 (2019).

95   Ding P, Ding Y. Stories of Salicylic Acid: A Plant Defense Hormone. Trends Plant Sci **25**:549–565
(2020).

96   Morgutti S, Negrini N, Pucciariello C, Sacchi GA. Role of trehalose and regulation of its levels as
a signal molecule to abiotic stresses in plants. doi:10.1016/B978-0-12-816451-8.00014-9
Elsevier, 2019.

97   Ramel F, Sulmon C, Serra AA, Gouesbet G, Couée I. Xenobiotic sensing and signalling in higher
plants. J Exp Bot **63**:3999–4014 (2012).

98   Singh A, Sheo ·, Prasad M, Singh RP. Plant Responses to Xenobiotics. Springer, 2016.