ENGRAILED-1 transcription factor exerts a paracrine neurotrophic activity on adult spinal cord α-motoneurons

Abbreviated title: **ENGRAILED-1 adult trophic activity**

Mélanie Lebœuf[1,2]*, Stephanie E. Vargas-Abonce[1,2]*, Eugénie Pezé-Hedsieck[1], Edmond Dupont[1], Lucia Jimenez-Alonso[2], Kenneth L. Moya[1] and Alain Prochiantz[1,2]

[1] Center for Interdisciplinary Research in Biology, Collège de France, CNRS, INSERM, Université PSL, Paris, France

[2] BrainEver SAS, Paris France

* Equal contribution to the study

Correspondence to Kenneth Moya (ken.moya@college-de-france.fr) and Alain Prochiantz (alain.prochiantz@college-de-france.fr).

Author contributions: ML and SEVA contributed to experiment design and execution; E.P-H developed the single-chain antibodies; L.J-A participated in all histochemistry experiments; ED did the immunoprecipitation and western blot experiments; KLM and AP designed the research and wrote the manuscript.

Acknowledgements: We thank the College de France imaging and animal facilities and Katia Avila Gutierrez for her help with some of the experiments. Drs. Anne Bousseau, Elizabeth Di Lullo, Ariel Di Nardo and Julien Spatazza provided very helpful comments on the manuscript.

Conflict of interest statement: AP and KLM are co-founders and hold shares in BrainEver, a company developing HPs for therapeutic use.

Funding sources: Memolife Labex PhD fellowship for SEVA, Association Nationale de la Recherche et de la Technologie (Cifre/ANRT n°2017/0488) to ML, BrainEver, Homeosign ERC-AdG n°339379, Fondation Schueller-Bettencourt.

ABSTRACT (204 words)

In addition to their cell-autonomous activities, a number of homeoprotein transcription factors transfer between cells and regulate gene expression, protein translation and chromatin organization in a non-cell autonomous way. ENGRAILED-1 is one such homeoprotein and is expressed in spinal V1 interneurons that synapse on α-motoneurons. Neutralizing extracellular ENGRAILED-1 by expressing a secreted single chain antibody blocks its capture by spinal motoneurons and induces the loss of neuromuscular strength and of some α-motoneurons. A similar but stronger phenotype is observed in the *Engrailed-1* heterozygote mouse, confirming that ENGRAILED-1 exerts a non-cell autonomous neurotrophic activity on spinal cord α-motoneurons. Intrathecal injection of ENGRAILED-1 leads to its specific internalization by spinal motoneurons and has long-lasting protective effects against neurodegeneration and neuromuscular weakness. Midbrain dopaminergic neurons express *Engrailed-1* and, similarly to spinal cord α-motoneurons, degenerate in the heterozygote. The identification of genes with modified expression in *Engrailed-1* heterozygote versus wild-type midbrain neurons that are also expressed in spinal cord motoneurons and with modified expression in the main familial ALS forms, led us to identify SQTSM1/p62 expression as an age marker in spinal cord motoneurons. The increase of SQTSM1/p62 expression is accelerated in the *Engrailed-1* heterozygote and upregulated upon extracellular ENGRAILED-1 neutralization. We propose that ENGRAILED-1 is a regulator of motoneuron ageing with non-cell autonomous neurotrophic activity.

INTRODUCTION

Homeoprotein (HP) transcription factors are key transcriptional regulators with well-established developmental functions, including control of morphogenesis, lineage decisions and cell differentiation[1,2]. In addition to their classical cell autonomous activities, non-cell autonomous developmental functions based on HP intercellular transfer were identified for ENGRAILED-1 (EN1) and EN2, VAX1, OTX2 and PAX6. Early in development, paracrine ENGRAILED induces anterior cross vein formation in the fly wing disk and shapes the zebrafish optic tectum[3–5]. ENGRAILED-2 and VAX1 respectively regulate retinal ganglion cell (RGC) axon guidance in the tectum and decussation at the optic chiasma[6–9], while PAX6 regulates the size of the zebrafish eye anlagen, enhances oligodendrocyte progenitor cell migration in the chick

neural tube and guides Cajal-Retzius cell migration in the mouse neuroepithelium[10–12]. At late developmental stages and in the adult, OTX2 non-cell autonomous activity regulates cerebral cortex plasticity[13,14], inner retinal physiological functions and RGC survival[15–17]. This novel signaling pathway is likely common to a larger number of HPs since the transfer domains are highly conserved between most HPs[18]. The finding that more than 150 HPs can transfer between cells, *in vitro* and in the embryonic brain, even though without identified functions for most of them, lends weight to this hypothesis[19].

Several vertebrate HPs remain expressed throughout life, but their adult physiological functions are still poorly understood. In the context of neurodegenerative diseases, EN1 is of particular interest since it is expressed in the mesencephalic dopaminergic (mDA) neurons of the Ventral Tegmental Area (VTA) and Substantia Nigra pars compacta (SNpc)[20,21], two cell populations that degenerate in Parkinson's Disease (PD), and promotes their survival in PD animal models[22]. In the *En1+/-* mouse (*En1*-Het), mDA neurons from the VTA and SNpc undergo progressive retrograde degeneration leading to a loss of 20% and 40% of their initial number, respectively[20,23,24]. Conversely, EN1 injection in the midbrain and its subsequent internalization by mDA neurons, thanks to its transduction properties, prevents their degeneration in rodent and macaque PD models[25,26]. Since mDA neurons express EN1, the latter observations reflect a cell-autonomous pro-survival activity of EN1. This incited us to search for other neurons that may require EN1 for their survival, either expressing *En1* or in the vicinity of *En1*-expressing cells and thus potentially exposed to secreted EN1. In the spinal cord, inhibitory V1 interneurons, including Renshaw cells (RCs), have been shown to express *En1*, and α-motoneurons (αMNs) are candidate non-cell autonomous targets owing to their direct contact with *En1*-expressing V1 interneurons[27–29].

We confirm here that *En1* is expressed in adult spinal cord V1 interneurons and now report that these interneurons do not degenerate in the *En1*-Het mouse. In contrast, αMNs of *En1*-Het mice show retrograde degeneration associated with progressive limb strength loss. This phenotype is in part the consequence of local EN1 non-cell autonomous activity since it can be partially mimicked in wild-type (WT) mice following EN1 neutralization in the spinal cord extracellular space. Furthermore, EN1 intrathecally injected at 3 months gains direct access to MNs and, in *En1*-Het mice, a single injection restores neuromuscular strength and prevents

αMN death for at least 2.5 months. This demonstrates a novel non-cell autonomous function of EN1 in the adult CNS and suggests that this HP may be of interest for developing novel therapeutic strategies and targets for diseases involving αMN degeneration.

RESULTS

ENGRAILED-1 is transferred from V1 interneurons to motoneurons in the ventral spinal cord

Previous studies reported *En1* expression in embryonic and early postnatal spinal cord V1 interneurons but, to our knowledge, no study has directly visualized EN1 in the adult mouse spinal cord. The available studies identify *En1*-expressing cells in the adult mouse using EN1-Cre-inducible reporter genes activated during development that maintain reporter gene expression throughout life[28,30,31]. In contrast with classical *in situ* hybridization (ISH) that gave very low signal in our hands, RNAscope (ISH) clearly reveals *En1* expression in adult 4.5-month-old mice V1 interneurons, including RCs identified by *Calbindin-1* expression (Figure 1A). These *En1*-expressing cells are localized dorsally (V1^P) and ventrally (V1^R, RCs) to a population of *En1*-negative neurons identified by a choline acetyltransferase (*ChAT*) ISH (RNAscope) probe, thus MNs. *En1* expression, quantified by RT-qPCR is fully maintained between 4.5 and 16 months and reduced by 2-fold in the *En1*-Het mouse (Figure 1B).

To verify EN1 expression, we used the 86/8 and the LSBio anti-EN1 antibodies that recognize EN1 C-terminal and N-terminal epitopes, respectively. The 86/8 antibody has been produced and validated in the laboratory[23], and LSBio is commercially available (Lifespan Biosciences). In 3-month-old mice, immunohistochemical staining with 86/8 failed to show a significant signal in the ventral spinal cord, including in *En1*-producing interneurons. This is in contrast with LSBio that stained both V1 interneurons (Figure 1C, left panel) and, surprisingly, *ChAT*-expressing MNs that do not express *En1* (Figure 1C, right panel). This led us to compare the two antibodies by western blot (WB) and to verify that, in the spinal cord and ventral midbrain, they recognize a major band that migrates at the same position as recombinant EN1 (Figure 1D, left panel). We then immunoprecipitated EN1 from spinal cord extracts using the 86/8 or LSBio antibody and immunoblotted with the other antibody. The results in Figure D (right panel) demonstrate that LSBio recognizes EN1 immunoprecipitated by 86/8 and vice versa. To further validate LSBio specificity, we evaluated the impact of preincubating the antibody with the EN1 protein prior to immunostaining. Figure 1E clearly demonstrates that the LSBio

staining in ChAT-expressing (ChAT+) cells (left panel) is totally lost upon preincubation of the antibody with recombinant EN1 (right panel).

This difference between the two antibodies suggests that EN1, in addition to being poorly expressed (i.e., not visible by classical ISH), adopts a conformation recognized by the LSBio but not by the 86/8 antibody. The presence of EN1 in MNs also suggests that EN1 produced by the interneurons is efficiently secreted and internalized by spinal MNs. As will be described below, EN1 intercellular transfer is further demonstrated by its extracellular neutralization through the expression of a secreted single chain antibody (scFvEN1) directed against EN1[6], resulting in an important reduction of MN staining by the LSBio antibody (Figure 4A).

Engrailed1-heterozygote mice show alpha-motoneuron retrograde degeneration

Because *En1*-expressing mDA neurons degenerate in the *En1*-Het midbrain[23], we examined whether ventral spinal cord neuronal populations, in particular interneurons that express *En1*, also degenerate in the *En1*-Het mouse. We used RNAscope labeling to compare the number of *En1*-expressing neurons in the spinal cord ventral horns of *En1*-Het mice and their wild-type (WT) littermates at 4.5 months of age. Figure 2A shows no cell loss in this population of V1 interneurons, including RCs (*Calbindin1*-expressing cells). This result and the absence of a change in *En1* expression between 4.5 and 16 months quantified by RT-qPCR (Figure 1B) demonstrate that, in contrast with mDA neurons, V1 interneuron survival is not affected by reduced *En1* expression in the *En1*-Het mouse mutant. We also quantified the number of *ChAT*-expressing (*ChAT*+) motoneurons in 4.5-month-old mice and did not observe a reduction in the *En1*-Het mouse (Figure 2A).

The ChAT-expressing population is heterogeneous with a mixture of γMNs and αMNs and the global analysis in Figure 2A may have missed degeneration, total or partial, of one of the two populations. Gamma and αMNs differ by size with the surface of γMNs ranging from 200 to 300μm^2 and that of αMNs equal or superior to 300μm^2 [32]. We counted MNs on Cresyl violet-stained and ChAT-labeled lumbar enlargement spinal cord sections (Figure 2B), taking soma size into account. We first performed this analysis at 9 months, reasoning that any degeneration would be more apparent at later ages. Cresyl violet and ChAT staining yield identical results and demonstrate a specific loss of large (>300μm^2) neurons (Figure 2C). The

concordance between data obtained from Cresyl violet and ChAT immunostaining precludes that the loss of ChAT+ cells in *En1*-Het mice is a consequence of ChAT expression down-regulation. We used Cresyl violet to quantify the number of small (100-199μm^2), medium (200-299μm^2) and large (>300m^2) neuronal populations as a function of age. Figure 2D demonstrates that, in the *En1*-Het mutant, the number of small-size (100-199μm^2) and intermediate (200-299μm^2) neurons did not decrease with age (Figure 2D, left and middle panels). In contrast, the number of αMNs strongly decreases between 3 and 4.5 months of age, with about 50% loss at 9 months and no further decrease between 9 and 15.5 months (Figure 2D, right panel). We then determined the consequences of αMN loss on muscle strength by measuring forepaw grip strength, the time holding onto an inverted grid, and the hindlimb extensor reflex. Figure 2E shows that strength loss appears between 2 and 3 months and reaches a maximum between 4.5 and 9 months of age. The loss of muscle strength between 2 and 3 months precedes αMN loss first observable at 4.5 months (Figure 2D, right panel), suggesting αMN retrograde degeneration.

This led us to analyze the organization of neuromuscular junctions (NMJs) by following the number of acetylcholine receptor (AChR) clusters, the endplate area, the percentage of endplates with perforations and that of fully occupied endplates[33]. The number of AChR clusters identified with fluorescent α-bungarotoxin (α-BTX) did not change with time (Figure 3A) and the endplate area remained stable for 9 months, with a modest 20 to 25% decrease at 15.5 months. A similar late-occurring (15.5 months) and modest effect was also observed for the percentage of endplates with perforations, a maturation index that is decreased in the mutant mice only at 15.5 months[34]. In contrast, a stronger and earlier difference was observed in the percentage of fully occupied endplates identified by the binding of fluorescent α-BTX and immunostaining of the high molecular weight neurofilament (165kDA) and the synaptic vesicle glycoprotein SV2A (Figure 3B, left panel). Fully occupied endplates were defined by an innervation occupying more than 80% of the surface and significant retrograde degeneration could already be measured in 3-month-old animals (Figure 3B, right panel).

All in all, these observations demonstrate muscle weakness and selective retrograde degeneration of αMNs in the *En1*-Het mouse.

Extracellular neutralization of ENGRAILED-1 partially reproduces the *Engrailed1-heterozygote phenotype*

Since αMNs do not express *En1*, their degeneration in the *En1-het* mouse must be a consequence of the *En1* heterozygote status of the V1 interneurons. Several mechanisms might contribute, alone or in combination, to αMN degeneration. In light of the presence of EN1 in MNs (Figure 1), of its known direct non-cell autonomous activity in several developmental phenomena[3–7] and of EN1/2 protective activity on mDA neurons[23,25,26,35,36], a parsimonious hypothesis is that EN1 exerts a direct non-cell autonomous trophic activity on αMNs.

To investigate this possibility, we used a strategy successfully developed in previous studies consisting in the expression of a secreted neutralizing anti-EN1 single-chain antibody (scFvEN1)[6] in the spinal cord of WT mice. The antibody, or its inactive control harboring a single cysteine to serine mutation that prevents disulfide bond formation and EN1 recognition, were cloned into an AAV8 under the transcriptional control of the Glial Fibrillary Acidic Protein (GFAP) promoter (Figure 4A). This promoter directs scFvEN1 expression and secretion by astrocytes, thus allowing for the neutralization of EN1 in the extracellular space. Viruses were injected intrathecally (IT) at lumbar level 5 (L5) in 1-month-old WT mice, allowing for robust scFvEN1 expression one month later (Figure 4B). The amount of EN1 in MNs was quantified 6 months post virus injection using the LSBio antibody, demonstrating a strong reduction in mice expressing scFvEN1 but not the inactive mutated form (Figure 4C). This result further confirms that LSBio recognizes MN-internalized EN1.

As shown in Figure 4D, in the three strength tests (anterior limb grasping, holding onto an inverted grid and hindlimb extensor reflex), mice expressing extracellular scFvEN1 experience gradual strength loss that begins between 6 and 8 weeks after virus injection. In contrast, the performance of mice expressing the mutated inactive antibody are indistinguishable from that of WT animals. Endplate morphology and αMN survival were evaluated 6 months after injection of the antibody-encoding viruses (i.e. at 7-months of age). The results demonstrate no effect of EN1 extracellular neutralization on the number of AChR clusters, endplate area and percentage of endplates with perforations, but a modest effect on the percentage of fully occupied endplates (Figure 4E, right panel). The analysis of total number of cells and,

separately, of the number of small, medium size and large neurons shows a small but specific loss of αMNs (Figure 4F).

Qualitatively, the latter results are similar to those obtained with the *En1*-Het mice with a small decrease in the number of fully-occupied endplates and the specific loss of large size αMNs. However, it is of note that the magnitude of the changes at 7 months was about two-fold less than in the *En1*-Het mouse with a 15% loss in fully occupied endplates and a 25% reduction in the number of αMNs, compared to 30% and 50% in 7-month-old *En1*-Het, respectively (Figures 2 and 3). These data establish that extracellular EN1 participates in the *En1*-Het phenotype but may not fully explain its magnitude.

Exogenous ENGRAILED-1 rescues the *Engrailed1*-heterozygote phenotype
We have previously demonstrated that EN1 administration in the ventral midbrain promotes mDA neuron survival in mouse and macaque PD models[23,25,26,35,36]. Since extracellular EN1 appears to exert a trophic activity on αMNs, we evaluated whether exogenous administration of EN1 could prevent αMN death in *En1*-Het mice. Therefore, human recombinant EN1 was intrathecally injected at the L5 level at a concentration of 1 µg in 5 µL in 3-month-old *En1*-Het mice (Figure 5A), a time when all αMNs are still present but muscle weakness is already noticeable, a consequence of retrograde degeneration initiated at the endplate (Figure 3). One and a half months later, the single EN1 injection had restored strength as measured by the 3 tests of anterior limb grasping, holding onto an inverted grid and hindlimb extensor reflex (Figure 5A). Accordingly, the percentage of fully occupied endplates was found to be back to WT level (Figure 5B, left panel), in parallel with the prevention of αMN death (Figure 5B, right panel). This last series of experiments establishes that extracellular EN1 exerts a curative trophic activity on αMNs of already symptomatic mice allowing for endplate reinnervation and the arrest of αMN degeneration.

The duration of the effect of EN1 injection on neuromuscular strength was evaluated over time. Figure 6A summarizes an experimental design consisting of a single EN1 injection at 3 months. Strength analysis (anterior grip, holding time on the inverted grid and extensor reflex score) demonstrates that the effect of a single injection is maintained until approximately week 12 post-injection and decreases thereafter. In spite of this decrease, 24 weeks post-

injection the strength of EN1-injected *En1*-Het mice is still superior to that of age-matched non-injected controls. Analysis of the % of fully occupied endplates and αMN number at 24 weeks confirms the long-lasting effect of a single injection. This led us to test whether a second injection 12 weeks after the first one could prolong the protective effect. Figure 6B summarizes this experimental protocol and illustrates that a second injection prolongs neuromuscular strength for another 8 to 12 weeks, depending on the test. Moreover, the percentage of fully occupied endplates and the number of αMNs at 24 weeks were identical to those observed in WT siblings.

Intrathecally injected EN1 preferentially accumulates in MNs

We then followed EN1 localization after intrathecal injection and, 24 hours after injection, we observed EN1 in the perivascular space and in ventral horn cells, primarily in MNs characterized by ChAT expression (Figure 7A, upper row). Figure 7A illustrates that EN1 staining in MNs is transient, peaking between 6 and 24 hours, still visible at 48 hours and barely so at 72 hours. The latter images were taken with the 86/8 antibody allowing us to see only the exogenous protein, but similar images were obtained with the LSBio antibody. Motoneurons identified by ChAT-staining were separated by size to distinguish between αMNs and γMNs and the kinetics of internalization were quantified in the two subpopulations (Figure 7A, bottom row), using the 86/8 (left panels) and LSBio (right panels). Indeed, the difference following internalization is much higher with the 86/8 since the basal staining due to endogenous EN1 is null with this antibody, but the time courses obtained with the two antibodies are very similar and the data obtained with the LSBio that recognizes endogenous and exogenous EN1 suggest that exogenous EN1 at its peak (24 hours) doubles the concentration of EN1 per MN (assuming that the antibody recognizes equally endogenous and exogenous EN1). All in all, the latter series of experiments demonstrates similar patterns for αMNs and γMNs, with a very rapid accumulation that peaks at 24 hours and decays rapidly, reaching nearly basal levels at 72 hours, allowing one to calculate a very short 24-hour EN1 half-time.

The preferential capture of EN1 by MNs might reflect the presence of EN1 binding sites at the MN surface. We previously identified a glycosaminoglycan (GAG) binding domain within the HP OTX2 that allows for its preferential capture by cerebral cortex parvalbumin (PV)

neurons[37]. This RK-OTX2 domain peptide (RKQRRERTTFTRAQL) competes for OTX2 binding to PV cells and antagonizes its specific internalization upon infusion or injection into the cerebral cortex[37]. A similar GAG-binding domain was recently described in EN2[38] and sequence comparisons between OTX2, EN2 and several other HPs identified putative GAG-binding sites in many of them (examples in Figure 7B). In the case of EN1, this domain (RKLKKKKNEKEDKRPRTAF), thereafter RK-EN1, is present between residues 292 and 310 of hEN1.

To test its potential role in EN1 localization, RK-EN1 was co-injected with EN1 (in a 20:1 molar ratio). Twenty-four hours after injection, in the absence of RK-EN1, the percentage of ChAT+ MNs that took up EN1 (determined with the 86/8 antibody) was about 50%, whereas when RK-EN1 was co-injected nearly 90% of the MNs did not take up EN1 (Figure 7B, central panel). In a separate experiment (Figure 7B, right panel), we tested peptide specificity by co-injecting EN1 with a peptide (AKLKAKKNEKEDKAPRTAF) in which one R and one K were replaced by alanines (AA peptide) or with a scrambled (Scr) RK-peptide (RLKNKEKPRDREKTKAKFK) composed of the same amino acids, but in a different order. As a control, 2 mice were injected with EN1 and the RK peptide, reproducing the inhibition of internalization observed in the first experiment. In contrast with the RK peptide, the two variant peptides were devoid of inhibitory activity. The absence of inhibition by the scrambled peptide demonstrates that inhibition cannot be explained only by a positive charge effect, suggesting that RK-EN1 is an addressing sequence for MNs in the ventral mouse spinal cord. It is of note that, even though only αMNs degenerate in the *En1*-Het mouse, EN1 is specifically captured by both γMNs and αMNs.

Bioinformatic analyses reveal SQTSM1/p62 as a regulated ENGRAILED-1 non-cell autonomous target in motoneurons

Retrograde degeneration of αMNs in the *En1*-Het and their long-lasting (2 months at least) rescue after a single EN1 injection is reminiscent of the situation described for mDA neurons, with the difference that EN1 activity is non-cell autonomous for αMNs and cell autonomous for mDA neurons. This similar protective effect led us to investigate if genetic pathways interacting with *En1* might be common to αMNs and mDA neurons. To that end, we compared previously acquired RNA sequencing data of the SNpc in WT and *En1*-Het mice at 6 weeks of

age, i.e. before the initiation of mDA neuron death but after the initiation of retrograde degeneration[23,24]. A significance threshold of p=0.05 was used, generating a list of differentially expressed genes[36] that was crossed with a library of MN-expressed genes[39]. Several physiological pathways were selected, indicated in the Methods section, resulting in a list of 402 genes. As in PD, there is ample evidence that the first steps in ALS occur in the distal motor neuron axon that retracts from the muscle in a ''dying back'' phenomenon[40]. To focus on relevant genes in the context of αMN retrograde degeneration, we used the STRING database to extract within the 402 genes those independently interacting with *SOD1*, *FUS*, *TARDBP-43* and *C9ORF72*, 4 genes mutated in a large majority of familial ALS (fALS)). This resulted in a short list of 20 genes modified in mDA neurons following *En1* loss of function, expressed in MNs and in interaction with one or several of the 4 fALS genes (Figure 8A).

Among these 20 genes, *SQTSM1/p62* interacts with the four fALS genes, is mutated in sporadic forms of the disease[41–44] and was recently shown to have variants in 486 fALS patients[45]. SQTSM1/p62 is a multifunctional protein that regulates the degradation of ubiquitinated proteins by the proteasome and carries ubiquitinated cargoes to the LC3 receptor at the autophagosome surface, a key step in cargo engulfment and lysosomal hydrolysis[46]. The importance of protein degradation and autophagy in protein homeostasis and age-associated pathologies[47–49] led us to compare the immunostaining for SQTSM1/p62 in lumbar αMNs (>300μm^2) and γMNs (200-299 μm^2) in different conditions. A first observation (Figure 8B) is that expression of SQTSM1/p62 in WT ventral spinal cord is very strong in MNs characterized by ChAT expression. A second one is that the intensity of SQTSM1/p62 expression increases with age in wild-type αMNs and γMNs and may thus be considered as MN physiological age markers (Figure 8C). In addition to expression, dispersion also increases with time (except between 3-and 4.5-month-old αMNs).

Comparison of SQTSM1/p62 expression with age in WT and *En1*-Het mice (Figure 8C), reveals a statistically significant difference only at 3 months in αMNs and γMNs (p=0.0499 and p=0.0536, respectively), before αMNs begin to degenerate, even though strength loss is already measurable (Figure 2C,D). Dispersion analysis yields significant differences between WT and *En1*-Het for αMNs at 3 (p<0.0001) and 4.5 months (p=0.0017) but not at 9 months (p=0.1416). The same applies to γMNs at 3 and 4.5 months (p<0.0001 and p=0.0007,

respectively), but barely at 9 months when, although significant (p<0.0001), the difference is very small. In addition, dispersion data obtained at 9 months both for αMNs and γMNs must be interpreted with care as signal saturation creates a ceiling effect, which is why we used a ß distribution adjusted between 0 and 1 (see under Methods). This increase in SQTSM1/p62 expression and dispersion at 3 months suggests accelerated aging of αMNs and γMNs in *En1*-Het mice during this period. This led us to follow SQTSM1/p62 expression and dispersion after extracellular EN1 neutralization in WT mice. Figure 8D shows that SQTSM1/p62 is significantly upregulated 6 months after injection of the scFvEN1 in one-month-old mice (p=0.0432 and 0.0392 for αMNs and γMNs, respectively), but not of its mutated inactive isoform (p=0.2961 and 0.2153 for αMNs and γMNs, respectively). Dispersion is significantly increased following EN1 extracellular neutralization (p<0.0001 for both αMNs and γMNs) with a much smaller, yet significant, effect of the mutated (inactive) antibody (p<0.0003 for both αMNs and γMNs), suggesting that a minor part of the effect on dispersion possibly results from virus-induced inflammation. When comparing the two antibodies (mutated and WT), the results are identical for αMNs and γMNs, with p=0.06 and <0.0001 for mean and dispersion, respectively.

Together, these results suggest that SQTSM1/p62 expression patterns are ageing MN marker and that extracellular EN1 neutralization in adult WT animals accelerates MN ageing.

DISCUSSION

This study establishes a novel adult function of EN1 and the importance of its expression in adult spinal cord V1 interneurons for αMN survival and neuromuscular strength. This non-cell autonomous EN1 activity involves, at least in part, EN1 secretion as shown by the deleterious effect on the neuromuscular phenotype of extracellular EN1 neutralization. Moreover, hEN1 injected intrathecally accumulates in MNs. In the *En1*-Het mouse, this extracellular gain of function, following a single intrathecal injection of hEN1, can reverse muscle denervation, arrest αMN degeneration and restore neuromuscular strength for several weeks. While EN1 injections were done using the human recombinant protein, human and mouse EN1 are 91.1% identical and show similar efficacy in mDA neuron survival experiments[25,26,35,36].

One day after injection, hEN1 is present around vascular structures probably in the perivascular space as well as within the parenchyma where it is observed within cells. Indeed,

it has been shown previously that EN1 and EN2, similarly to many HPs[19], translocate across the plasma membrane and gain access to the cell cytoplasm and nucleus by a mechanism distinct from endocytosis[14,22,50–52]. Injected hEN1 shows preferential accumulation in all ChAT-positive ventral MNs, although this does not preclude that a minority of other cell types may also capture the protein, albeit at lower levels. The deleterious effect of blocking extracellular EN1 with an anti-EN1 scFv secreted by astrocytes establishes that EN1 secreted by the V1 interneurons is internalized by αMNs and γMNs expressing EN1-binding sites and participates in several aspects of MN physiology, including αMN survival.

The latter finding is in line with the non-cell autonomous functions of EN1/2 reported in the fly wing disk and in the frog, fish and chick optic tectum[3–7]. It is also reminiscent of OTX2, another HP that specifically accumulates in PV interneurons of the cerebral cortex[13]. Prior to internalization, OTX2 binds to GAGs present at the surface of these interneurons thanks to a GAG-binding domain (RKQRRERTTFTRAQL) that overlaps with the first helix of the homeodomain[37,53]. Interestingly, similar putative GAG-binding domains are present upstream of the homeodomain in many HPs (examples in Figure 7), including EN2[38] and EN1 for which this putative GAG-binding sequence is RKLKKKKNEKEDKRP. The role of this domain in the specific targeting of EN1 and the presence of EN1 binding sites at the MN surface is illustrated by the ability of the peptide to abolish EN1 specific uptake by ventral spinal cord MNs. Although we have not directly demonstrated that EN1 binds GAGs, the conservation of the GAG binding domain and the analogy with OTX2[37] and EN2[38] support a specific association of EN1 with MN-expressed GAGs. If this is confirmed by future studies, it will be important to identify the molecular nature of this GAG as done for OTX2, EN2 and also VAX1[9]. Given the existence of a putative GAG-binding domain in many HPs, this study and those on OTX2 and VAX1 raise the issue of the existence of a sugar code for the specific recognition of their target cells by transferring HPs.

In the *En1*-Het mouse, the deficits in neuromuscular strength begin before 3 months of age. These behavioral deficits coincide with a decrease in fully occupied endplates first observed at 3 months and precede αMN loss which is initiated between 3 and 4.5 months. This temporal delay likely reflects a retrograde degeneration of αMNs that starts at the terminals, whereby peripheral strength is affected before αMN cell body loss. It might be also a consequence of

accelerated MN aging in the *En1*-Het mouse, as suggested by the enhanced p62 expression and value variance at 3 months of age in the mutant (discussed below). Retrograde degeneration also occurs in *En1*-expressing mDA neurons in the same *En1*-Het mouse line with their terminals showing signs of degeneration at 4 weeks, whereas they only start dying at 6 weeks postnatal[24].

Retrograde degeneration of αMNs is also a prominent feature of human motoneuron diseases[40]. In this context, it is of note that hEN1 injection at 3 months not only prevents lumbar αMN death for months, but also restores normal endplate innervation and muscle strength, suggesting reinnervation. The neurodegenerative process is slowed or halted on a time scale of months and a second injection 3 months after the first one prolongs rescue for another 2 to 3 months. The time course of αMN degeneration in *En1*-Het animals is rapid with approximately 70% αMNs destined to die between months 3 and 4.5. Then, the process slows down to reach a plateau with ~40% of the αMNs remaining at 15.5 months. It is thus possible that αMNs are heterogeneous in terms of their dependency on EN1. Another possibility is that when αMNs death reaches 50%, remaining MNs receive sufficient secreted EN1 to halt accelerated ageing and the ensuing degeneration (αMNs) or pathology (γMNs). This hypothesis is supported by the fact that the difference in SQTSM1/p62 expression is only visible at 3 months when MNs receive half of the normal EN1 dose but not at 4.5 months or later when almost half of the αMNs have disappeared. A strict evaluation of αMN responses, to decreasing EN1 doses will be necessary to answer the question properly.

Although EN1 activity is cell autonomous in mDA neurons and non-cell autonomous in αMNs, the similarities of their responses to *En1* hypomorphism, such as progressive retrograde degeneration preceding cell body loss, led us to interrogate the repertoire of differentially expressed genes in WT and *En1*-Het mDA neurons, and to compare it with MN pathways. This produced a short list of genes and one of them, *SQSTM1/p62*, interacts with *SOD1*, *FUS*, *TARDBP-43* and *C9ORF72*, 4 genes responsible for a majority of fALS. *SQSTM1/p62* encodes SQSTM1/p62, a known regulator of degradation by the proteasome and autophagy through its role in bringing ubiquitinated proteins to the proteasome or to LC3 at the autophagosome surface[49]. This unbiased identification of *SQSTM1/p62* in the *En1* "pathway" is particular interesting because, in addition to interacting with the 4 main fALS mutations, itself presents

variants in 486 patients with familial ALS and is also mutated in sporadic ALS cases[41–44]. We observe an upregulation of SQTSM1/p62 in the *En1*-Het mouse at 3 months or scFvEN1 injected WT mice at 7 months, thus in the surviving αMNs, and also in the γMNs which capture EN1 but do not degenerate in either mouse model. In addition, SQTSM1/p62 expression levels and dispersion in WT are increased during ageing in the absence of MN death.

These series of observations can be interpreted in two non-mutually exclusive hypotheses. A first one is that SQTSM1/p62 plays a homeostatic role in ageing neurons and possibly protects them against degeneration. A second, based on the fact that SQTSM1/p62 itself is degraded through autophagosomes, is that ageing reduces autophagy and fosters SQTSM1/p62 accumulation[54–57]. SQSTM1/p62 is a multifunctional protein with distinct domains interacting with different co-factors[58]. Thus, if a protective activity is demonstrated in further studies, specific deletions will be necessary to identify the pathways involved, in particular within the N-terminal domain, which presents a pro-survival activity through the N-kB pathway, and the C-terminal domain, which regulates proteostasis through the activation of proteasome and autophagosome activities[59].

Strikingly, although EN1 is internalized by all MNs, which show similar increases in SQTSM1/p62 expression and variance in both the *En1*-Het mouse or following EN1 extracellular neutralization, only αMNs and not γMNs degenerate in both models. This demonstrates that αMNs, contrary to γMNs, are EN1-dependent for their survival. The survival of approximately half of the αMNs may reflect a differential sensitivity of αMN subpopulations and/or the fact that, as discussed above, the surviving neurons receive higher and sufficient amounts of EN1. The absence of dependency on EN1 for γMN survival also applies to *En1*-expressing V1 interneurons that do not degenerate in the *En1*-Het mouse. This contrasts with mDA neurons that express *En1* and degenerate in the *En1*-Het mouse. EN1 survival activity is therefore independent of its cell autonomous or non-cell autonomous activity, but rather may reflect a higher sensitivity to stress of mDA neurons and αMNs.

The differential susceptibility of αMNs and γMNs to reduced *En1* expression in the *En1*-Het mouse or to EN1 extracellular availability following scFvEN1 viral expression may be explained, at least in part, by their different functions and the underlying anatomy[60]. Alpha-MNs receive

excitatory inputs mainly from local spinal circuits with an important contribution from descending upper motor neurons. They innervate extrafusal fibers and the innervation ratio (i.e., the number of muscle fibers per αMN axon) in humans can reach up to 2000[61,62]. Gamma-MNs innervate muscle spindles, receive their excitatory inputs from the reticular formation and their inhibitory ones from relay neurons mainly found in the dorsal horn that does not express EN1. They fire after αMNs at a slower rate and in humans the innervation ratio is close to parity (i.e., 1-3 muscle fibers per axon), suggesting that their metabolic demand is less than that of αMNs. Finally, while γMNs can take up exogenous EN1 similarly to αMNs, unlike them, they do not receive contacts from EN1-expressing interneurons, and the functions of EN1 might thus differ between the two MN types. A full understanding of these distinct functions requires further studies.

We observe that the deleterious phenotype is stronger in the *En1*-Het mouse than following extracellular EN1 neutralization in WT mice. This may be due to the fact that EN1 neurotrophic activity exerted on αMNs is purely non-cell autonomous but that scFvEN1 neutralizes less than 50% of extracellular EN1 (due to affinity, levels of expression and/or localization of the antibody). The near total absence of EN1 in MNs following scFvEN1 expression rather suggests a more complex explanation whereby, while scFvEN1 only targets pure non-cell autonomous EN1 activity, the *En1*-Het phenotype may result from a 50% loss in secreted EN1 and, in addition, from cell autonomous dysfunctions affecting V1 interneurons but not inducing their death. V1 interneurons are an essential component of the reciprocal inhibitory αMN circuit and loss of their inhibitory input places αMNs at risk[63–66]. Furthermore, deficits in inhibitory interneurons in the spinal cord of motoneuron disease models have been reported[67,68] and Renshaw cell pathology has been observed following αMN loss[68,69]. One possibility is that EN1 regulates the expression of classical trophic factors in spinal V1 interneurons that are secreted and support αMN survival, in conjunction with EN1 signaling (Figure 9). Such synergy between classical secreted factors and HP signaling has been reported for Engrailed and DPP in the fly wing disk[3], for EN2, adenosine and EphrinA5 in chick RGC growth cone navigation[6,70] and for PAX6 and Netrin1 in the migration of chick oligodendrocyte precursors[10].

It remains that the scFvEN1 experiment demonstrates an EN1 non-cell autonomous activity. This activity could be through extracellular EN1 acting directly on αMNs but also indirectly

through other cell types, such as astrocytes, oligodendrocytes, or microglial cells (Figure 9). However, the internalization preference of injected EN1 by MNs and the fact that injected EN1 totally restores a healthy phenotype in the *En1*-Het mouse strongly suggest that EN1 exerts its non-cell autonomous activity primarily on αMNs. This activity is reminiscent of the direct non-cell autonomous protective activity of OTX2 for RGCs upon NMDA excitotoxic insult[16]. Still in the context of the comparison between the phenotypes observed in the *En1*-Het mouse and in the scFvEN1 mouse, one must be aware of the fact that, in the *En1*-Het, EN1 levels are lower very early during development. We thus cannot exclude that extracellular EN1 acts in conjunction with compromised interneuron physiological properties in *En1*-Het animals, even if they are present in the appropriate number.

EN1 may protect αMNs through one or several of the mechanisms previously characterized for mDA neurons of the SNpc, which can be protected from oxidative stress by a single EN1 injection[26,35,36]. Among these mechanisms are the regulation of Complex I mitochondrial activity, DNA break repair, heterochromatin maintenance, and the repression of long interspersed nuclear elements (LINE-1) expression[25,35,36]. The evaluation of these possibilities in the spinal cord will be the objective of future studies but the long-lasting effect of a transient twofold increase in EN1 concentration within MNs strongly suggests an activity taking place at the chromatin level.

The *En1-Het* mouse presents muscle weaknesses, abnormal spinal reflex, NMJ denervation and αMN loss, all phenotypes reminiscent of changes in ALS patients and in some ALS mouse models. This raises the question as to whether the results reported here are relevant for αMN diseases. To our knowledge, although they may exist in some sporadic cases, as for PD[71], no studies so far associate ALS with mutations affecting *En1* integrity or *En1* expression. It is thus clear that the *En1-het* mouse cannot, as yet, be considered as an ALS mouse model. However, *En1* genetic interactions with pathways leading to ALS, and its therapeutic activity in classical mouse and non-human primate models of PD[25,26,35,36], allows us to envisage its use as a long-lasting therapeutic protein in ALS mouse models or iPSC-derived human MNs.

One main mechanism possibly linking EN1 expression to ALS is that, in many neurodegenerative diseases, including ALS, ageing is a major risk factor. Indeed, our results

indicate that SQSTM1/p62 expression is a marker of age, both in αMNs and γMNs, in agreement with its interaction with redox regulation pathways[72]. Since *En1*-Het mDA neurons are more sensitive than WT to oxidative stress and given that EN1 protects them against experimental oxidative stress[36], we speculate that the up-regulation of SQTSM1/p62 expression, either in the mutant or following extracellular loss due to scFvEN1 expression, reveals an EN1 anti-ageing activity that may explain the ALS-like phenotype of *En1* hypomorphs. Indeed, we now need to follow the effects of gain and loss of EN1 functions in *bona fide* ALS animal models.

In conclusion, we have demonstrated a novel non-cell autonomous activity for EN1 in adult spinal cord. Constitutive reduction in EN1 expression or local neutralization of extracellular EN1 cause motor endplate degeneration, muscle weakness and αMN loss. This is an example of an important non-cell autonomous activity for this transcription factor as is also the case for EN2, PAX6 and VAX1[14,22,50,51]. Given that the sequences necessary for intercellular transfer are conserved in most HP[18,73] and that such transfer has been demonstrated for about 150 of them[19], it is not unreasonable to predict that the developmental and physiological non-cell autonomous functions demonstrated for a handful of HPs only represent the tip of the iceberg.

MATERIALS AND METHODS

Animal Management

All animal treatments followed the guidelines for the care and use of laboratory animals (US National Institutes of Health), the European Directive number 86/609 (EEC Council for Animal Protection in Experimental Research and Other Scientific Utilization) and French authorizations n° 00703.01 "Therapeutic homeoproteins in Parkinson Disease" and n° APAFIS #6034-2016071110167703 v2, "Spinal cord motoneuron neuroprotection" delivered by the Minister of higher education, research and innovation. Adult mice were housed two to five per cage with *ad libitum* access to food and water and under a 12h light/dark cycle. Transgenic mouse strain *En1*-Het was bred by the Rodent Breeding Services provided by the Animal Care Services at College de France. Females and males were included in all studies. The endpoint

limit for euthanasia was a 15% or greater loss of bodyweight or signs of paralysis; in all experiments, no mouse reached these endpoints.

Behavior analyses

Mice were habituated to the behavioral room and to the experimenter 24 hours before the day of testing and again before each behavioral test. All tests were performed on the same day and behavioral assessment was carried by evaluators blind to genotype and treatment.

Forepaw Grip Strength. The Transducer (IITC Life Science Grip Strength Meter, ALMEMO 2450 AHLBORN, World Precision Instruments) was calibrated and the scale of values set to grams. Each mouse was lifted by the tail to position the front paws at the height of the bar (about 15 cm) and then moved towards the bar. When symmetric firm grip with both paws was established, the mouse was pulled backward at a constant speed until the grasp was broken and the maximal value was recorded. The test was repeated 5 times per animal with a minimal resting time of 5 minutes between tests and the mean of all values was normalized to the weight of each animal.

Inverted Grid Test. The mouse was placed on a wire grid (15x10 cm) and allowed to explore it. After 3-5 minutes, the grid was raised 30 cm above a soft surface and slowly inverted. Latency to release was recorded three times per mouse with a minimum resting time of 5 minutes between trials. The longest latency was used for analysis.

Hindlimb Extensor Reflex. Mice were suspended by the tail at a constant height (about 15 cm) and scored for hindlimb extension reflex. The scores were assigned from 0 to 3 as follows: 3 for normal symmetric extension in both hind limbs without visible tremors; 2.5 for normal extension in both hind limbs with tremor in one or both paws; 2.0 for unequal extension of the hind limbs without visible tremors; 1.5 for unequal extension in the hind limbs with tremors in one or both paws, 1.0 for extension reflex in only one hindlimb, 0.5 for minimum extension of both hindlimbs, 0 for absence of any hindlimb extension.

Tissue preparation

Spinal cord. Adult mice were euthanized by a 1 µl/g body weight dose of Dolethal (Euthasol: 4 µg/µl). Spinal cords were dissected and placed in Phosphate Buffer Saline (PBS) to remove meninges and surrounding connective tissues. Cervical and lumbar enlargements were separately placed in paraformaldehyde 4% (PFA, Thermo Scientific) for 1 hour at room temperature (RT) with gentle mixing, washed in PBS three times for 30 minutes at RT and placed in PBS with 20% sucrose overnight at 4°C. After cryoprotection, the tissue was embedded in Tissue Freezing Medium (TFM, Microm Microtech), frozen on dry ice and 30µM sections were prepared using an HM 560 Microm cryostat (Thermo Scientific).

Muscle. The lumbrical muscles were dissected in cold PBS, fixed at RT in 4% PFA for 10 minutes and washed in PBS. Muscle whole-mounts were processed *in toto* to visualize the entire innervation pattern and allow for a detailed NMJ analysis [33].

Cresyl violet staining. Slides with 30µm spinal cord sections were washed in PBS 3 times, cleared in O-Xylene (CARLO-HERBA) for 5 minutes, then hydrated in graded alcohols with increasing water and placed in Cresyl Violet acetate (MERCK). Sections were then dehydrated in increasing alcohol and mounted in Eukitt® Quick-hardening mounting medium (Sigma).

Spinal cord and muscle immunofluorescence labeling. Slides with 30µm spinal cord sections or whole-mount muscles were washed in PBS and permeabilized with 2% Triton. After 30 minutes at RT in 100µM glycine buffer, 10% Normal Goat Serum (NGS, Invitrogen) or Fetal Bovine Serum (FBS, Gibco) was added in the presence of 1% Triton before incubation with primary antibodies (sheep anti-Choline Acetyltransferase (ChAT) ABCAM 1:1000, goat anti-ChAT Millipore 1:500, rabbit anti-EN1 86/8 1:300[23], rabbit anti-EN1 LSBio (CliniSciences) 1:200, mouse anti-neurofilament 165kDa Developmental Studies Hybridoma Bank 1:50, mouse anti-synaptic vesicle glycoprotein 2A DSHB 1:100 and rabbit anti-p62 ABCAM 1:1000 overnight at 4°C, washed and further incubated with secondary antibodies for 2 hours at RT. Where indicated, anti-EN1 LSBio activity was neutralized by a 1-hour (RT) preincubation with hEN1 (1.5 EN1/LSBio molar ratio). For muscle staining, α-bungarotoxin (Alexa Fluor 488 conjugate) was included at the same time as the secondary antibodies. Slides were washed and mounted with DAPI Fluoromount-G® (Southern Biotech). Controls without primary antibodies were systematically included.

RT-qPCR

Spinal cords were removed as above and lumbar enlargements were rapidly frozen on dry ice. Total RNA was extracted (RNeasy Mini kit, Qiagen) and reverse transcribed using the QuantiTect Reverse Transcription kit (Qiagen). RT-qPCR was done using SYBR-Green (Roche Applied Science) and a Light Cycler 480 (Roche Applied Science). Data were analyzed using the "2-ddCt" method and values were normalized to *Glyceraldehyde 3-phosphate dehydrogenase* (*Gapdh*). The following primers were used: *Engrailed-1* sense: CCTGGGTCTACTGCACACG, antisense: CGCTTGTTTTGGAACCAGAT; *Gapdh* sense: TGACGTGCCGCCTGGAGAAAC, antisense: CCGGCATCGAAGGTGGAAGAG.

Protein and single-chain antibodies

Protein

Human EN1 (hEN1) was produced as described[16] and endotoxins were removed by Triton X-144 phase separation. In brief, pre-condensed 1% Triton X-144 (Sigma) was added to the protein preparation. The solution was incubated for 30 minutes at 4°C with constant stirring, transferred to 37°C for 10 minutes and centrifuged at 4000 rpm for 10 minutes at 25°C. The endotoxin-free protein was aliquoted and kept at -80°C.

Single-chain antibodies

The anti-EN1 single chain antibody plasmid was prepared from the anti-Engrailed 4G11 hybridoma (Developmental Hybridoma Bank, Iowa City, A, USA). Cloning was as described[6] with or without signal peptide[12] with 6 myc tags at the C-terminus of the antibody followed by a GFP downstream of a P2A skipping peptide. Addition of a GFAP promoter, insertion in an AAV8 backbone and production of the AAV8 were done by Vector Biolabs

Intrathecal injections

Mice were anesthetized with 1μl/g ketamine-xylazine (Imalgene1000: 10μg/μl, Rompur 2%: 0,8μg/μl) and placed on the injection platform. The tail of the animal was taken between two fingers and the spinal column was gently flattened against a padded surface with the other hand. The L3 vertebral spine was identified by palpation and a 23G x 1" needle (0.6x25 mm Terumo tip) was placed at the L1 and T13 groove and inserted through the skin at an angle of

$20°$[74]. The needle was slowly advanced into the intervertebral space until it reached the injection point, provoking a strong tail-flick reflex. Injections of 5µl were done at a rate of 1µl/min with 1µg recombinant protein, AAV8-IgK-scFvMUT-6xMyc-2A-eGFP (scFvMUT), or AAV8-IgK-scFvEN1-6xMyc-2A-eGFP (scFvEN1) (Vector Biolabs, 10^{12} GC/ml). The peptides (ProteoGenix) were co-injected with hEN1 at a 20-fold molar excess of peptides. The needle was left in place for two minutes after injection and then slowly removed. Animals were placed in a warmed chamber until recovery from anesthesia. Extensor reflex and gait analysis were examined 2 and 24 hours after injection to verify the absence of spinal cord damage.

RNAscope Fluorescent in Situ Hybridization

Mice aged 4.5 months were euthanized with Dolethal and transcardially perfused with PBS followed by 4% PFA in PBS. The lumbar region of the spinal cord was dissected, fixed overnight in 4% PFA at 4°C and cryoprotected in PBS 30% sucrose for 24 hours at 4°C. Spinal cord sections 40 µm-thick were prepared using a Leitz (1400) sliding microtome. Sections were washed in PBS, incubated with RNAscope hydrogen peroxide solution from Advanced Cell Diagnostics (ACD) for 10 min at RT, rinsed in Tris-buffered saline with Tween (50 mM Tris-Cl, pH 7.6; 150 mM NaCl, 0.1% Tween 20) at RT, collected on Super Frost plus microscope slides (Thermo Scientific), dried at RT for 1 hour, rapidly immersed in ultrapure water, dried again at RT for 1 hour, heated for 1 hour at 60°C and dried at RT overnight. The next day, sections were immersed in ultrapure water, rapidly dehydrated with 100% ethanol, incubated at 100°C for 15 minutes in RNAscope 1X Target Retrieval Reagent (ACD), washed in ultrapure water and dehydrated with 100% ethanol for 3 minutes at RT. Protease treatment was carried out using RNAscope Protease Plus solution (ACD) for 30 minutes at 40°C in a HybEZ oven (ACD). Sections were then washed in PBS before *in situ* hybridization using the RNAscope Multiplex Fluorescent V2 Assay (ACD). Probes were hybridized for 2 hours at 40°C in HybEZ oven (ACD), followed by incubation with signal amplification reagents according to the manufacturer's instructions. Probes were purchased from ACD: Mm-*En1*-C1 (catalog #442651), Mm-*Chat*-C2 (catalog #408731-C2), Mm-*Calb1*-C3 (catalog #428431-C3). The hybridized probe signal was visualized and captured on an upright Leica CFS SP5 confocal microscope with a 40x oil objective.

RNAscope and immunofluorescence co-detection

Sections prepared as described above were collected on Super Frost plus microscope slides and heated for 30 min at 60°C. RNAscope was performed according to the manufacturer's instructions (Multiplex Fluorescent V2 Assay and RNA-Protein Co-detection Ancillary Kit Cat No. 323180). In brief, sections were post-fixed in 4% PFA for 15 min at 4°C, dehydrated in 50%, 70%, 100% (twice) ethanol solutions for 5 min each, incubated with RNAscope hydrogen peroxide solution for 10 min at RT, rinsed in ultrapure water, incubated at 100°C for 5 minutes in RNAscope 1X Co-detection Target Retrieval, rinsed in ultrapure water and washed in Phosphate buffer, 0.1% tween 20 pH7.2 (PBS-T). Sections were incubated overnight at 4°C with anti-EN1 LSBio diluted (1:200) in Co-detection Antibody Diluent, washed in PBS-T, fixed in 4% PFA for 30 min at RT and washed twice in PBS-T. Sections were treated with protease 30 min at 40°C in a HybEZ oven, washed in PBS before *in situ* hybridization using the RNAscope Multiplex Fluorescent V2 Assay. Probes were hybridized for 2 hours at 40°C in HybEZ oven, followed by incubation with signal amplification reagents according to manufacturer instructions. Finally, sections were incubated with the secondary antibody diluted in the Co-detection Antibody Diluent for 30 min at room temperature.

Immunoprecipitation and Western blots

Tissue extracts

Spinal cords and ventral midbrains dissected from adult mice were kept frozen at -80 °C. After thawing on ice and 3 rinses in PBS, tissues were resuspended in 500 µl dissociation buffer (DB: 50 mM Tris-HCl pH8.0, 0.3 M NaCl, Triton X-100 1%, EDTA-free Complete Proteases inhibitor cocktail from Roche and 250U/ml of benzonase endonuclease from Sigma). After homogenization in a Dounce tissue homogenizer, 10 times with the A piston and 10 times with the B piston, lysates were incubated for 30 min on ice, sonicated for 5 min in Diagenode Bioruptor set with 30/30 seconds ON/OFF at high power, incubated on ice for 30 min and centrifuged for 10 min at 12000g. Supernatants were harvested, their protein contents measured (Pierce BCA assay)) and frozen at -80 °C.

Immunoprecipitation of EN1 from SC :

SC extracts were precleared on protein A-sepharose beads with pre-immune serum of the rabbit used to generate the antibody and submitted to immunoprecipitation with EN1 antibodies. In brief, 214 µl of extract diluted in 1ml DB and incubated with 50 µl of a pre-

immune rabbit serum on ice for 30 min before adding 100 µl of a 50/50 protein A-sepharose bead slurry in DB and incubation at 4 °C under agitation for 1 hour. Beads were centrifuged 10s at 10000g and the supernatant was kept (INPUT fraction). Five µl of 86/8 or LSBio antiserum were added to the INPUT fraction before incubation on ice for 1 hour, the addition of 100µl 50/50 protein A-sepharose slurry beads in DB and incubation at 4°C under agitation. The supernatant (Flow Through) resulting from a 10s at 10000g centrifugation was put aside, the beads were washed 3 times with the DB and eluted 3 times with 50µl Glycine 0.1M pH 3.0 (E1, E2, E3). Eluted fractions were immediately neutralized with 10µl of Tris-HCl 2 M pH 8.0, the beads were resuspended in 50 µl dissociation buffer and neutralized with the same buffer. All extracts were kept at -20 °C.

Western blots

Invitrogen Nu-PAGE 4 to 12% acrylamide 1mm SDS-PAGE gels were used to run spinal cord extracts or the hEN1recombinant protein. Extracts (2-3µg total proteins) and 2 ng of purified EN1 were diluted in 10µl Invitrogen LDS-DTT Sample Buffer, heated for 10 min at 70°C, briefly centrifuged at 10000g and loaded. For the immunoprecipitation followed by Western blot, 1/200 of total inputs or Flow-Throughs and 1/20 of the total protein A–sepharose E2 elution were used per well. SeeBlue 2 and molecular weight markers were from Invitrogen. Gels were run in MOPS buffer at 150 V for 1 hour and 15 min and the proteins transferred for 30' at 100 V on Millipore PVDF membranes (BIORAD Criterion Blotter apparatus). Membranes were blocked for 1 hour at RT with 4% non-fat dry milk (NFDM) diluted in TBS containing 0.2% Tween-20 (TBST). Antibodies to EN1 86/8 and LSBio were diluted 1/200 in TBST 4% NFDM and incubated overnight at 4 °C. Blots were rinsed 3 times for 15 min each with TBST and incubated with HRP-Labeled anti-Rabbit antibody diluted 1/2500 for 1 hour at RT. When preceded by immunoprecipitations, revelation was with HRP-protein G diluted 1/5000 for 1hour at RT followed by 3 TBST rinses of 15 min each. HRP activity was revealed with chemiluminescence substrate CLARITY from BIORAD and the light was quantified with the FUJI LAS 4000 imager. Signals were quantified with the Image Studio Lite software from Li-Cor.

Bioinformatic analysis

Differentially expressed genes (cutoff p<0.05) from the RNA-seq performed on microdissected six-week-old SNpc from WT and *En1-Het* mice (GSE72321)[36] were compared to genes

identified from an RNA-seq of microdissected MNs from WT mice (GSE38820)[41]. Transcripts present in the two lists were retained. Using STRING database, 10 biological processes of interest were selected including regulation of gene expression, cellular response to stress, cell communication, regulation of I-kappaB kinase/NF-kappaB signaling, cellular response to DNA damage stimulus, response to oxygen-containing compounds, locomotion, DNA repair, modulation of chemical synaptic transmission, and brain development. Using these pathways reduced the list of selected genes to 402. To the 402 genes, we used the STRING database to identify known interactions with the four main genes involved in familial ALS (fALS): *SOD1, FUS, C9Orf72* and *TARDBP*.

Image analyses

Cresyl violet-stained spinal cord section images were acquired with a Nikon-i90 microscope under bright field conditions at 10x with a Z-stack step of 0.5µm. RNAscope FISH images were acquired using a Yokogawa X1 Spinning Disk confocal microscope at 20x (Nikon) and acquisitions of 3D z-stack (1µm) using 491 nm, 561 nm and 642 nm laser beams. For cell quantification, at least five spinal cord sections were analyzed for each animal. Immunofluorescence-stained spinal cord sections images were acquired with a Leica SP5 inverted confocal microscope at 20x (Leica DMI6000) and acquisitions of 3D z-stack (0.5µm) were made using the UV (405 nm, 50 mW), Argon (488 nm, 200 mW) and DPSS (561 nm, 15 mW) lasers. For cell quantification, at least five spinal cord sections separated by $\geq$ 900µm were analyzed for each animal. ChAT+ cells and Cresyl violet-stained cells with a soma area above 100µm^2 were manually outlined using ImageJ software (NIH, Bethesda, MD) and their area determined. Analyses were carried out on Z-stacks through the entire 30µm thickness of the section. Cells were classified as small (100-199µm^2 cross sectional area), intermediate (200-299µm^2) or large (> 300µm^2). ChAT and Cresyl violet gave similar results for medium size (200-299µm^2) and large (>300µm^2) neurons thus allowing us to characterize αMNs based on size (>300µm^2), with or without additional ChAT staining. For p62 staining analysis, the motoneuron region of five lumbar sections were acquired at 40x for each animal. Mean intensity was measured in ChAT+ cells with a soma greater than 150µm^2 using ImageJ software. For each experiment, image acquisition was performed with the same parameter settings of the confocal microscope to allow for comparison between experimental conditions.

Lumbrical muscles were imaged with a Leica SP5 inverted confocal microscope (Leica DMI6000) with a motorized XY stage. 3D z-stack (0.5µm) acquisitions were made as above (UV, Argon and DPSS) and images analyzed using ImageJ software. Analyses were performed blind to the genotype and treatment. Endplates were categorized as fully innervated (neurofilament overlying more than 80% of the endplate), partially innervated (neurofilament overlying up to 80% of the endplate) or denervated (no neurofilament overlying the endplate). Endplate morphology was evaluated by counting the number of endplates with perforations (areas lacking α-bungarotoxin staining). For postsynaptic analysis, each endplate was manually outlined using ImageJ software and its area calculated. All analyses were performed on the entire Z-stacks through the NMJ.

Statistical analyses

Results are expressed as mean ± SD. Statistical significance was determined as follows. For RT-qPCR, WT and *En1*-Het mice were compared by Unpaired T-test with equal SD. For behavioral and NMJ analyses and αMN counting in the time-course study, groups were compared by Unpaired T-test with equal SD comparing WT with *En1*-Het for each time point. For intrathecal injections, behavioral and NMJ analyses and αMN counting, experimental data were compared by one-way ANOVA followed by a post hoc Tukey test for comparisons to WT. For behavioral analysis in the time-course protection of injected hEN1, groups were compared by Unpaired T-test with equal variances comparing WT with *En1*-Het injected at each time point.

The p62 measures are bounded between 0 and an upper quantification limit (ceiling effect), and classical normally distributed models are ill-suited to account for these two constraints. We solicited Pharmalex Belgium to develop a suitable analysis. Since it can take a wide variety of shapes and is naturally bounded in the [0;1] interval, beta-distribution was fitted on the p62 measure, for all MNs, γMNs and αMNs separately, after rescaling the original data to the [0;1] interval. The shape and scale parameters account for time, group and time*group interaction as fixed effect. It models the variance observed in the data by time, group and time*group interaction, meaning each combination has its own variability. Finally, the mouse-to-mouse variability (or between mouse variability), is modeled by time and group.

FIGURE LEGENDS

Figure 1. *En1* expression in the adult spinal cord

A*.* Triple RNAscope *in situ* hybridization showing *Engrailed-1* (*En1*), *Choline acetyltransferase* (*ChAT*) and *Calbindin-1* (*Calb1*) expression in the lumbar spinal cord. *En1* is expressed in V1 interneurons, dorsal (V1^P) and ventral (V1^R) to the main *ChAT*-expressing motoneuron pool. Ventral interneurons correspond to Renshaw cells as shown by *Calb1* expression. Scale bar: 500μm.

B. RT-qPCR of RNA from the lumbar enlargement at 4.5- and 16-months of age shows stable *En1* expression in WT at both ages and a two-fold reduction of expression in heterozygous mice. Unpaired t-test. **p<0.005; ***p<0.0005. n=3.

C. Triple staining EN1 IHC (green), *En1* RNAscope ISH (red) and *ChAT* RNAscope (blue) demonstrating the double-staining of *En1* mRNA and protein (EN1) in the V1 interneuron population (left panel insets), and the presence of EN1 protein in large cells not expressing *En1* mRNA (left panel) but expressing *ChAT* (right panel insets). Scale bar: 500μm or 30μm for high magnification insets.

D. Left panel: Western blots of spinal cord (SC) and ventral midbrain (VMB) extracts demonstrating that the 86/8 and LSBio antibodies recognize in the 2 structures the same protein migrating with recombinant EN1 velocity. Right panel: Immunoprecipitation of spinal cord extracts by LSBio or 86/8 antibody followed by Western blotting with 86/8 or LSBio antibody, respectively.

E. Left: EN1 (red) detected with the LSBio antibody is localized in ChAT-expressing neurons (green) in the ventral horns of the spinal cord. Right: EN1 signal is lost upon preincubation of the antibody with 1.5 M excess of recombinant hEN1. Scale bar: 50μm.

Figure 2. Progressive loss of αMNs and neuromuscular strength decrease in *En1*-HET mice

A. Analysis of the number of *En1+*, *Calb1+* and *ChAT+* neurons (triple RNAScope). At 4.5 months, WT and *En1*-Het mice show no difference in the number of cells expressing *En1*, *Calb1* or *ChAT*. n=5-6.

B. Cresyl violet and ChAT staining of a ventral WT spinal cord at the lumbar level. Scale bar: 100 μm.

C. Cresyl violet and ChAT staining at the lumbar level show no medium size (200-299 μm^2) cell loss (γMNs) and an about 50% decrease in the number of large size (>300 μm^2) cells (αMNs). Unpaired t-test. **p<0.005; ***p<0.0005. n=5.

D. Lumbar level Cresyl violet staining shows that, in contrast with small and medium size neurons (interneurons and γMNs, left and center panels), large size neurons (αMNs, right panel) undergo progressive death first measured at 4.5 months. Unpaired t-test with equal SD comparing WT with *En1*-Het at each time point (*p<0.05; **p<0.005; ***p<0.0005; ****p<0.0001). n=5 to 6.

E. Compared to WT mice, *En1*-Het mice experience gradual neuromuscular strength loss. This loss is observed with the forepaw grip strength (left panel), the inverted grid test (center panel and the hindlimb extensor reflex test (right panel). Strength loss is first observed between 2 and 3 months, thus before measurable αMN cell body loss.

Unpaired T-test with equal SD comparing WT with *En1*-Het at each time point (*p<0.05; **p<0.005; ***p<0.0005; ****p<0.0001). n=4 to 20.

Figure 3. Neuromuscular junction (NMJ) morphology differences between WT and *En1*-Het mice

A. NMJs of *En1*-Het and WT mice show similar numbers of AChR clusters (left panel) and late-occurring (15.5 months) decrease in endplate area (center panel) and percentage of perforated endplates (right panel). The number of AChR clusters seems to decrease between 2 and 3 months, but a 1-way ANOVA followed by Tukey corrected post hoc comparisons show that this is not significant and reflects a high variance in the analysis (p=0.071).

B. Left panel illustrates the use of Alexa Fluor 488-conjugated α-bungarotoxin (α-BTX, in green) and of neurofilament and synaptic vesicle glycoprotein antibodies (2H3 and SV2A, in red) to evaluate the percentage of fully occupied endplates (>80% occupancy). The right panel show that the % of fully occupied endplates decreases progressively in the *En1*-Het mouse, starting between 3 and 4.5 months of age. Scale bar: 100μm.

Unpaired T-test with equal SD comparing WT with *En1*-Het at each time point (*p<0.05; **p<0.005). n=4 to 8.

Figure 4. The effects of extracellular EN1 neutralization on neuromuscular strength, MN survival and NMJ morphology

A. Experimental paradigm and structure of AAV8-encoded construct containing glial fibrillary acidic protein (GFAP) promoter for expression in astrocytes, Immunoglobulin K (IgK) signal peptide for secretion, anti-ENGRAILED single-chain antibody (scFvEN1), 6 myc tags (6xMyc), skipping P2A peptide and enhanced Green Fluorescent Protein (EGFP). An inactive control antibody (scFv MUT) contains a cysteine to serine mutation that prevents disulfide bond formation between IgG chains, thus epitope recognition. The AAV8 was injected in 1-month-old WT mice and the strength phenotypes were followed for 6 months before anatomical analysis.

B. Analysis 1 month post-injection showing that scFv antibodies are expressed in astrocytes (white arrowhead) double-stained for GFAP and Myc and exported (empty arrowhead). Scale bar: 50 µm.

C. Left panel illustrates that expressing the scFvEN1, but not scFvMUT, abolishes EN1 staining by LSBio anti-EN1 antibody in ventral horn ChAT+ cells and right panel quantifies this inhibition 1-way ANOVA followed by Tukey corrected post-hoc comparisons. (n=5 mice per group, ***$p<0.0005$, ****$p<0.0001$). Scale bar: 100µm.

D. The three graphs illustrate how the WT antibody but not its mutated version leads to progressive neuromuscular strength decrease. 1-way ANOVA followed by Tukey corrected post-hoc comparisons. (*$p<0.05$; **$p<0.005$; ***$p<0.0005$; ****$p<0.0001$). n=5 per treatment.

E. Six months following infection (7-month-old mice), extracellular EN1 neutralization does not modify the number of AChR clusters, nor the endplate surface area, nor the percentage of perforated endplates. In contrast, the percentage of fully occupied endplates is diminished (right end panel). 1-way ANOVA followed by Tukey corrected post-hoc comparisons. (**$p<0.005$). n=5 per treatment.

F. Six months following infection, extracellular EN1 neutralization does not globally modify the total neuron number of cells at the lumbar level (left panel). A separate analysis of small (100-199 µm^2), medium (200-299 µm^2) and large (>300 µm^2) neurons demonstrate a specific ($p<0.0557$) loss of the latter category (αMNs). 1-way ANOVA followed by Tukey corrected post-hoc comparisons. n=5 per treatment.

Figure 5. Intrathecal hEN1injection in *En1-Het* mice restores neuromuscular strength and prevents αMN death

A. Mice were tested for neuromuscular strength at three months of age when (before the onset of αMN loss), strength has already decreased in *En1*-het mice as measured in the forepaw grip strength, inverted grid and extensor reflex tests (left side of each graph). The next day the *En1*-het mice were separated into two groups. One group received buffer and the other group recombinant hEN1 (1 μg in 5 μl), injected at the L5 level. One and a half months later (4.5 months) *En1*-het mice injected with hEN1 have recovered normal strength, in contrast with non-injected mice or mice injected with buffer. Unpaired t-test used at three months of age. ****p<0.0001. For 4.5-month comparisons, 1-way ANOVA followed by Tukey corrected post-hoc comparisons. ***p<0.0005; ****p<0.0001. n=9 to 31

B. At 4.5 months, following hEN1 injection at 3 months, the percentage of fully occupied endplates and the number of αMNs are not significantly different from control values. 1-way ANOVA followed by Tukey corrected post hoc comparisons. ***p<0.0005; ****p<0.0001. n=5 to 21

Figure 6. EN1 injection at 3 months has long-lasting effects on strength, endplate occupancy and αMN survival and the latter recoveries are prolonged by a second injection

A. Top left panel shows the single injection protocol whereby hEN1 is injected at 3 months (1μg in 5μl) intrathecally at the L5 level and mouse behavior followed for 24 weeks. The 3 time course graphs demonstrate that a single injection restores neuromuscular strength measured by the tests of grip strength, time on the inverted grid and extensor reflex and that this effect lasts for 12 weeks. After 12 weeks strength decreases progressively but, even after 24 weeks, remains superior to that of untreated *En1*-Het mice. At 24 weeks, the % of fully occupied endplates is inferior to that of WT mice, but superior to that of non-injected *En1*-Het mice. The same holds true for the number of αMNs. For behavior, the groups were compared by unpaired T-test with equal variances comparing WT with *En1*-Het injected at each time point. (*p<0.05; **p<0.005; ***p<0.0005; ****p<0.0001). For the endplate analysis and αMNs, groups were compared by 1-way ANOVA followed by Tukey corrected post-hoc comparisons (*p<0.05; **p<0.005; ***p<0.0005; ****p<0.0001). n= 4 to 10.

B. Based on the results shown in A, a new experiment was performed with a second injection 12 weeks after the first injection at 3 months of age. Again, the injection at three months of age restored neuromuscular strength and in mice receiving the second injection strength was

maintained an additional ten weeks or more compared to mice receiving a single injection. The three strength graphs demonstrate a positive effect of the second injection with values intermediate between those measured in WT mice and En1-Het mice with a single injection. The percentage of fully occupied endplates and the number of αMNs are back to wild type values in *En1*-Het mice injected twice. Groups were compared by Unpaired T-test with equal variances comparing WT with *En1*-Het injected once or twice at each time point. For behavior, the groups were compared by unpaired T-test with equal variances through 12 weeks. After, the groups were compared by 1-way ANOVA and if significant followed by Tukey corrected post-hoc comparisons (*$p<0.05$; **$p<0.005$; ***$p<0.0005$; ****$p<0.0001$). For the endplate analysis and αMNs, groups were compared by 1-way ANOVA followed by Tukey corrected post hoc comparisons (*$p<0.05$; **$p<0.005$; ***$p<0.0005$; ****$p<0.0001$). n= 4 to 10.

Figure 7. Following intrathecal injection, hEN1 is addressed specifically to all MNs

A. Top left panel shows that 24 hours after intrathecal injection (1 μg in 5μL) at the L5 level of 2-month-old mice, hEN1 (red) can be primarily visualized in ventral horn ChAT+ (green) cells Arrowheads show hEN1 internalized by MNs. Scale bar: 100 μm. Top right panels show the progressive accumulation and clearance of hEN1 in ventral horn MNs, with a pick between 6 and 24 hours. EN1 was revealed by the 86/8 antibody allowing for the visualization of exogenous EN1 only (scale bar: 100μm). Bottom panels show the quantification of EN1 in in γMNs and αMNs with the 86/8 antibody at 6, 24, 48 and 72 hours (two left panels). In the two right panels quantification is done with the LSBio antibody in non-injected mice and at 24 and 72 hours after injection. Since the LSBio sees both endogenous and exogenous EN1, the increase is only 2-fold, but qualitatively, the results are very similar, demonstrating rapid internalization and clearance of the exogenous protein and allowing one to calculate a half-life of 24 hours. 1-way ANOVA followed by Tukey corrected post-hoc comparisons (*$p<0.05$; **$p<0.005$; ***$p<0.0005$). When not significant, p values shown.

B. Left panel gives examples of putative glycosaminoglycan (GAG)-binding domain in 11 homeoprotein transcription factors. Based on the alignment and on published work on OTX2[37] and EN2[38], a putative EN1 GAG-binding domain (RK-EN1) was designed. Middle panel quantifies the inhibitory effect of RK-EN1 on hEN1 capture (86/8 antibody) by ChAT+ cells demonstrating that RK-EN1 in a 1 to 20 ratio reduces the % of EN1-postive MNs (ChAT+) from

50 to less that 10%. The right panel demonstrates that this inhibitory activity is not shared by the mutant AA peptide or by a scrambled (Scr) peptide. Unpaired T-test with equal SD (**p<0.005; ***p<0.0005). n=3-5

Figure 8. SQTSM1/p62 expression by MNs increases with age

A. The search for genes differentially expressed in WT and *En1*-Het mDA neurons and expressed in MNs, thus putative non-cell autonomous EN1 targets in MNs, allowed for the identification of 402 genes (after pathway selection). These genes were investigated for an interaction with genes mutated in the main 4 familial ALS forms. Among them, *SQTSM1/p62* (*p62*) expression is upregulated in the SNpc (RNA-seq) and in MNs of *En1*-Het mice.

B. Immunohistochemical staining shows the presence of high amounts of p62 in ChAT+ 3-month-old cell bodies. Scale bar: 50 μm.

C. Intensity measurements demonstrates that the mean of SQTSM1/p62 expression increases with age in WT γMNs (left) and αMNs (right). However, comparing WT and *En1*-Het shows a significant difference only at 3 months and not later. Dispersion also increases with time (except between 3- and 4.5-month-old γMNs) and differences between WT and *En1*-Het are seen at the 3 ages in all MNs, except at 9 months for αMNs. Statistical analysis is explained under material and methods.

D. Mean intensity and dispersion of SQTSM1/p62 expression are increased in γMNs (left) and αMNs (right) in mice expressing scFvEN1 6 months after virus injection (7-month-old mice) demonstrating that EN1 extracellular neutralization increases SQTSM1/p62 expression and dispersion. The mutated antibody (scFVMUT) does not increase p62 expression, but has a small but significant effect on dispersion.

Figure 9. Proposed schema of EN1 cell autonomous and non-cell autonomous activities

EN1 transcribed and synthesized in V1 (pink) has a cell autonomous activity in these cells contributing to the expression of several V1 factors (F_A, F_B, …) that may signal to MNs (yellow) and other cell types (blue). All the latter F factors do not necessarily require EN1 activity. EN1 after its secretion by V1 interneurons is preferentially internalized by MNs (fat red arrow) and less by other cell types (thin red arrow), if at all. These other cell types express other F Factors (F_X, F_Y, …) which are secreted and are, or not, under non-cell autonomous EN1 activity. As a result, MNs experience the signaling activity of F factors (possibly EN1-dependent as in the

scheme, but not necessarily so) and internalize EN1. Following internalization, EN1 can work alone or in synergy with the F factors to regulate transcription, translation and chromatin confirmation within MNs.

Acknowledgements

We would like to thank Ariel Di Nardo, Lizzie di Lullo and Julien Spatatza for reading the manuscript and discussing experiments and results with the authors. We are grateful for the help brought by the CIRB Imaging and animal facilities. Funding was provided by Memolife Labex PhD fellowship for SEVA, Association Nationale de la Recherche et de la Technologie (Cifre/ANRT n°2017/0488) to ML, BrainEver, Homeosign ERC-AdG n°339379 and Fondation Schueller-Bettencourt.

References

1. Gehring, W. Homeo boxes in the study of development. *Science* 236, 1245–1252 (1987).

2. Holland, P. W. H. & Takahashi, T. The evolution of homeobox genes: Implications for the study of brain development. *Brain Res Bull* 66, 484–490 (2005).

3. Layalle, S. *et al.* Engrailed homeoprotein acts as a signaling molecule in the developing fly. *Development* 138, 2315–2323 (2011).

4. Rampon, C. *et al.* Control of brain patterning by Engrailed paracrine transfer: a new function of the Pbx interaction domain. *Development* 142, 1840–1849 (2015).

5. Amblard, I. *et al.* H2O2 and Engrailed 2 paracrine activity synergize to shape the zebrafish optic tectum. *Commun Biology* 3, 536 (2020).

6. Wizenmann, A. *et al.* Extracellular Engrailed participates in the topographic guidance of retinal axons in vivo. *Neuron* 64, 355–366 (2009).

7. Brunet, I. *et al.* The transcription factor Engrailed-2 guides retinal axons. *Nature* 438, 94–98 (2005).

8. Yoon, B. C. *et al.* Local Translation of Extranuclear Lamin B Promotes Axon Maintenance. *Cell* 148, 752–764 (2012).

9. Kim, N. *et al.* Regulation of retinal axon growth by secreted Vax1 homeodomain protein. *eLife* 3, e02671 (2014).

10. Lullo, E. D. *et al.* Paracrine Pax6 activity regulates oligodendrocyte precursor cell migration in the chick embryonic neural tube. *Development* 138, 4991–5001 (2011).

11. Kaddour, H. *et al.* Extracellular Pax6 Regulates Tangential Cajal–Retzius Cell Migration in the Developing Mouse Neocortex. *Cereb Cortex* (2019) doi:10.1093/cercor/bhz098.

12. Lesaffre, B., Joliot, A., Prochiantz, A. & Volovitch, M. Direct non-cell autonomous Pax6 activity regulates eye development in the zebrafish. *Neural Development* 2, 2 (2007).

13. Sugiyama, S. *et al.* Experience-dependent transfer of Otx2 homeoprotein into the visual cortex activates postnatal plasticity. *Cell* 134, 508–520 (2008).

14. Nardo, A. A. D., Joliot, A. & Prochiantz, A. Homeoprotein transduction in neurodevelopment and physiopathology. *Sci Adv* 6, eabc6374 (2020).

15. Torero-Ibad, R. *et al.* OTX2 non-cell autonomous activity regulates inner retinal function. *Eneuro* 7, ENEURO.0012-19.2020 (2020).

16. Ibad, R. T. *et al.* Otx2 promotes the survival of damaged adult retinal ganglion cells and protects against excitotoxic loss of visual acuity in vivo. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 31, 5495–5503 (2011).

17. Pensieri, P. *et al.* Photoreceptor cKO of OTX2 enhances OTX2 intercellular transfer in the retina and causes photophobia. *Eneuro* ENEURO.0229-21.2021 (2021) doi:10.1523/eneuro.0229-21.2021.

18. Prochiantz, A. & Joliot, A. Can transcription factors function as cell–cell signalling molecules? *Nat Rev Mol Cell Bio* 4, 814–819 (2003).

19. Lee, E. J. *et al.* Global Analysis of Intercellular Homeodomain Protein Transfer. *Cell Reports* 28, 712-722.e3 (2019).

20. Sgadò, P. *et al.* Slow progressive degeneration of nigral dopaminergic neurons in postnatal Engrailed mutant mice. *Proceedings of the National Academy of Sciences* 103, 15242–15247 (2006).

21. Nardo, A. A. D. *et al.* Dendritic localization and activity-dependent translation of Engrailed1 transcription factor. *Molecular and Cellular Neuroscience* 35, 230–236 (2007).

22. Nardo, A. A. D., Fuchs, J., Joshi, R. L., Moya, K. L. & Prochiantz, A. The Physiology of Homeoprotein Transduction. *Physiol Rev* 98, 1943–1982 (2018).

23. Sonnier, L. *et al.* Progressive Loss of Dopaminergic Neurons in the Ventral Midbrain of Adult Mice Heterozygote for Engrailed1. *J Neurosci* 27, 1063–1071 (2007).

24. Nordström, U. *et al.* Progressive nigrostriatal terminal dysfunction and degeneration in the engrailed1 heterozygous mouse model of Parkinson's disease. *Neurobiology of Disease* 73, 70–82 (2015).

25. Alvarez-Fischer, D. *et al.* Engrailed protects mouse midbrain dopaminergic neurons against mitochondrial complex I insults. *Nature Neuroscience* 14, 1260–1266 (2011).

26. Thomasson, N. *et al.* Engrailed-1 induces long-lasting behavior benefit in an experimental Parkinson primate model. *Movement Disorders* 34, 1082–1084 (2019).

27. Wenner, P., O'Donovan, M. J. & Matise, M. P. Topographical and physiological characterization of interneurons that express engrailed-1 in the embryonic chick spinal cord. *Journal of neurophysiology* 84, 2651–2657 (2000).

28. Alvarez, F. J. *et al.* Postnatal phenotype and localization of spinal cord V1 derived interneurons. *J Comp Neurol* 493, 177–192 (2005).

29. Higashijima, S., Masino, M. A., Mandel, G. & Fetcho, J. R. Engrailed-1 expression marks a primitive class of inhibitory spinal interneuron. *J Neurosci Official J Soc Neurosci* 24, 5827–39 (2004).

30. Sapir, T. Pax6 and Engrailed 1 Regulate Two Distinct Aspects of Renshaw Cell Development. *Journal of Neuroscience* 24, 1255–1264 (2004).

31. Lane, A. R., Cogdell, I. C., Jessell, T. M., Bikoff, J. B. & Alvarez, F. J. Genetic targeting of adult Renshaw cells using a Calbindin 1 destabilized Cre allele for intersection with Parvalbumin or Engrailed1. *Sci Rep-uk* 11, 19861 (2021).

32. Powis, R. A. & Gillingwater, T. H. Selective loss of alpha motor neurons with sparing of gamma motor neurons and spinal cord cholinergic neurons in a mouse model of spinal muscular atrophy. *Journal of Anatomy* 228, 443–451 (2016).

33. Sleigh, J. N., Burgess, R. W., Gillingwater, T. H. & Cader, M. Z. Morphological analysis of neuromuscular junction development and degeneration in rodent lumbrical muscles. *J Neurosci Meth* 227, 159–165 (2014).

34. Comley, L. H., Nijssen, J., Frost-Nylen, J. & Hedlund, E. Cross-disease comparison of amyotrophic lateral sclerosis and spinal muscular atrophy reveals conservation of selective vulnerability but differential neuromuscular junction pathology. *Journal of Comparative Neurology* 524, 1424–1442 (2016).

35. Thé, F.-X. B. de *et al.* Engrailed homeoprotein blocks degeneration in adult dopaminergic neurons through LINE-1 repression. *The EMBO Journal* 37, (2018).

36. Rekaik, H. *et al.* Engrailed Homeoprotein Protects Mesencephalic Dopaminergic Neurons from Oxidative Stress. *CellReports* 13, 242–250 (2015).

37. Beurdeley, M. *et al.* Otx2 Binding to Perineuronal Nets Persistently Regulates Plasticity in the Mature Visual Cortex. *Journal of Neuroscience* 32, 9429–9437 (2012).

38. Cardon, S. *et al.* A cationic motif in Engrailed-2 homeoprotein controls its internalization via selective cell-surface glycosaminoglycans interactions. *Biorxiv* 2021.07.29.454375 (2021) doi:10.1101/2021.07.29.454375.

39. Bandyopadhyay, U. *et al.* RNA-Seq Profiling of Spinal Cord Motor Neurons from a Presymptomatic SOD1 ALS Mouse. *Plos One* 8, e53575 (2013).

40. Fischer, L. R. *et al.* Amyotrophic lateral sclerosis is a distal axonopathy: evidence in mice and man. *Exp Neurol* 185, 232–240 (2004).

41. Shimizu, H. *et al.* Sporadic ALS with compound heterozygous mutations in the SQSTM1 gene. *Acta Neuropathol* 126, 453–459 (2013).

42. Chen, Y. *et al.* SQSTM1 mutations in Han Chinese populations with sporadic amyotrophic lateral sclerosis. *Neurobiol Aging* 35, 726.e7-726.e9 (2014).

43. Fecto, F. *et al.* SQSTM1 Mutations in Familial and Sporadic Amyotrophic Lateral Sclerosis. *Arch Neurol-chicago* 68, 1440–1446 (2011).

44. Yang, Y. *et al.* Six SQSTM1 mutations in a Chinese amyotrophic lateral sclerosis cohort. *Amyotroph Lat Scl Fr* 16, 378–384 (2015).

45. Yilmaz, R. *et al.* SQSTM1/p62 variants in 486 familial ALS patients from Germany and Sweden. *Neurobiol Aging* (2019) doi:10.1016/j.neurobiolaging.2019.10.018.

46. Doherty, J. & Baehrecke, E. H. Life, death and autophagy. *Nature Cell Biology* 20, 1110–1117 (2018).

47. Menzies, F. M., Fleming, A. & Rubinsztein, D. C. Compromised autophagy and neurodegenerative diseases. *Nature Reviews Neuroscience* 16, 345–357 (2015).

48. Leidal, A. M., Levine, B. & Debnath, J. Autophagy and the cell biology of age-related disease. *Nature Cell Biology* 20, 1338–1348 (2018).

49. Klionsky, D. J. *et al.* Autophagy in major human diseases. *Embo J* 40, e108863 (2021).

50. Prochiantz, A. & Nardo, A. A. D. Homeoprotein signaling in the developing and adult nervous system. *Neuron* 85, 911–925 (2015).

51. Brunet, I., Nardo, A. A. D., Sonnier, L., Beurdeley, M. & Prochiantz, A. The topological role of homeoproteins in the developing central nervous system. *Trends in Neurosciences* 30, 260–267 (2007).

52. Amblard, I. *et al.* Bidirectional transfer of homeoprotein EN2 across the plasma membrane requires PIP2. *J Cell Sci* 133, jcs244327 (2020).

53. Miyata, S., Komatsu, Y., Yoshimura, Y., Taya, C. & Kitagawa, H. Persistent cortical plasticity by upregulation of chondroitin 6-sulfation. *Nat Neurosci* 15, 414–422 (2012).

54. Bjørkøy, G. *et al.* Monitoring autophagic degradation of p62/SQSTM1. *Methods Enzymol* 452, 181–97 (2009).

55. Jakobi, A. J. *et al.* Structural basis of p62/SQSTM1 helical filaments and their role in cellular cargo uptake. *Nat Commun* 11, 440 (2020).

56. Jaakkola, P. M. & Pursiheimo, J.-P. p62 degradation by autophagy: Another way for cancer cells to survive under hypoxia. *Autophagy* 5, 410–412 (2009).

57. Tai, H. *et al.* Autophagy impairment with lysosomal and mitochondrial dysfunction is an important characteristic of oxidative stress-induced senescence. *Autophagy* 13, 99–113 (2016).

58. Ma, S., Attarwala, I. Y. & Xie, X.-Q. SQSTM1/p62: A Potential Target for Neurodegenerative Disease. *Acs Chem Neurosci* 10, 2094–2114 (2019).

59. Foster, A. & Rea, S. The role of sequestosome 1/p62 protein in amyotrophic lateral sclerosis and frontotemporal dementia pathogenesis. *Neural Regen Res* 15, 2186 (2020).

60. Kanning, K. C., Kaplan, A. & Henderson, C. E. Motor Neuron Diversity in Development and Disease. *Annu Rev Neurosci* 33, 409–440 (2010).

61. Masson, G. L., Przedborski, S. & Abbott, L. F. A computational model of motor neuron degeneration. *Neuron* 83, 975–988 (2014).

62. FEINSTEIN, B., LINDEGARD, B., NYMAN, E. & WOHLFART, G. Studies on action potentials in normal human muscles. *Acta Psych Neurol* 29, 189–95 (1954).

63. Quinlan, K. A. Links between electrophysiological and molecular pathology of amyotrophic lateral sclerosis. *Integrative and comparative biology* 51, 913–925 (2011).

64. Wang, Z., Li, L., Goulding, M. & Frank, E. Early postnatal development of reciprocal Ia inhibition in the murine spinal cord. *Journal of neurophysiology* 100, 185–196 (2008).

65. Ramírez-Jarquín, U. N. & Tapia, R. Excitatory and Inhibitory Neuronal Circuits in the Spinal Cord and Their Role in the Control of Motor Neuron Function and Degeneration. *Acs Chem Neurosci* 9, 211–216 (2018).

66. Ramírez-Jarquín, U. N., Lazo-Gómez, R., Tovar-y-Romo, L. B. & Tapia, R. Spinal inhibitory circuits and their role in motor neuron degeneration. *Neuropharmacology* 82, 101–107 (2014).

67. Hossaini, M. *et al.* Spinal Inhibitory Interneuron Pathology Follows Motor Neuron Degeneration Independent of Glial Mutant Superoxide Dismutase 1 Expression in SOD1-ALS Mice. *Journal of Neuropathology and Experimental Neurology* 70, 662–677 (2011).

68. Allodi, I., Montañana-Rosell, R., Selvan, R., Löw, P. & Kiehn, O. Locomotor deficits in a mouse model of ALS are paralleled by loss of V1-interneuron connections onto fast motor neurons. *Nat Commun* 12, 3251 (2021).

69. Chang, Q. & Martin, L. J. Glycinergic innervation of motoneurons is deficient in amyotrophic lateral sclerosis mice: a quantitative confocal analysis. *The American Journal of Pathology* 174, 574–585 (2009).

70. Stettler, O. *et al.* Engrailed homeoprotein recruits the adenosine A1 receptor to potentiate ephrin A5 function in retinal growth cones. *Development* 139, 215–224 (2012).

71. Haubenberger, D. *et al.* Association of transcription factor polymorphisms PITX3 and EN1 with Parkinson's disease. *Neurobiol Aging* 32, 302–307 (2011).

72. Hensley, K. & Harris-White, M. E. Redox regulation of autophagy in healthy brain and neurodegeneration. *Neurobiol Dis* 84, 50–59 (2015).

73. Joliot, A. & Prochiantz, A. Transduction peptides: from technology to physiology. *Nature Cell Biology* 6, 189–196 (2004).

74. Hylden, J. L. K. & Wilcox, G. L. Intrathecal morphine in mice: A new technique. *Eur J Pharmacol* 67, 313–316 (1980).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

A

ALS genes	Interactions within the 402 genes
Sod1	*Xdh, Ccs, Casp9, Ogg1, Eno1, Nfe2l1, Becn1, Gsk3a, Rac3, Lrrk2, p62/Sqstm1*
Fus	*Gtf2f1, Rnps1, Ptbp1, Sf3a1, Myef2, p62/Sqstm1*
Tardbp-43	*Ube2i, Ptbp1, Phf1, Lrrk2, p62/Sqstm1*
C9Orf72	*Pde3b, Stk38l, Lrrk2, p62/Sqstm1*

Figure 8

Figure 9