Piezo1 mechanosensing regulates integrin-dependent chemotactic migration in human T cells

Chinky Shiu Chen Liu[1], Tithi Mandal[2], Parijat Biswas[3], Md. Asmaul Hoque[1,4], Purbita Bandopadhyay[1,4], Bishnu Prasad Sinha[1,4], Jafar Sarif[1,4], Ranit D'Rozario[1,4], Deepak Kumar Sinha[3], Bidisha Sinha[2], Dipyaman Ganguly[1]*

[1]IICB-Translational Research Unit of Excellence, CSIR-Indian Institute of Chemical Biology, Kolkata, India.

[2]Department of Biological Sciences, Indian Institute of Science Education and Research, Kolkata, India.

[3] Department of Biological Sciences, Indian Association for Cultivation of Science, Kolkata, India.

[4]Academy of Scientific and Innovative Research, Ghaziabad, India.

Correspondence: dipyaman@iicb.res.in

Running title: Piezo1 mechanosensing in human T cell chemotaxis

Keywords: Human T cells, Piezo1, Mechanosensing, Migration, Chemokine, Chemotaxis, Focal adhesion kinase, LFA-1, F-actin, Confocal Microscopy, Interference Reflection Microscopy

ABSTRACT

T cells are crucial for efficient antigen-specific immune responses and thus their migration within the

body, to inflamed tissues from circulating blood or to secondary lymphoid organs, play a very critical

role. T cell extravasation in inflamed tissues depends on chemotactic cues and interaction between

endothelial adhesion molecules and cellular integrins. A migrating T cell is expected to sense diverse

external and membrane-intrinsic mechano-physical cues, but molecular mechanisms of such

mechanosensing in cell migration are not established. We explored if the professional

mechanosensor Piezo1 play any role during integrin-dependent chemotaxis of human T cells. We

found that deficiency of Piezo1 in human T cells interfered with integrin-dependent cellular motility.

Piezo1 recruitment at the leading edge of moving T cells follows focal adhesion formation and local

increase in membrane tension on chemokine receptor activation. Piezo1 activation constitutes a

membrane-intrinsic component of the 'outside-in' signaling in human T cells, migrating in response

to chemokines, that mediates integrin recruitment to the leading edge.

INTRODUCTION

Efficient T cell migration from blood circulation to secondary lymphoid organs and inflamed tissue is paramount for optimal tissue antigen sampling and effector functions during adaptive immune response (1-4). The process is dependent on chemotactic cues, interaction between endothelial adhesion molecules and T cell integrins, and an intricate acto-myosin dynamics (5). Generation of mechanical force is critical to propel migrating cells (6-9). Mechanosensing of the substrate and increase in cellular membrane tension is also intuitively apparent in a migrating T cell, although mechanistic details are not explored to a great extent.

Piezo1 and Piezo2 ion channels are evolutionarily conserved professional mechanosensors that can sense increase in membrane tension mostly independent of other protein-protein interactions (10-12). Mechanosensing by Piezo1 has been demonstrated to play critical role in diverse patho-physiologic contexts given its wide tissue expression including the immunocellular compartment (13-21). We previously identified crucial role of this specialized mechanotransducer in optimal TCR triggering (14, 19). The immunocellular expression of Piezo1 thus makes it a strong candidate for mechanosensing at the plasma membrane to regulate motility of immune cells. Piezo1 shows differential involvement if cellular migration depending on context (22- 28). Previous studies point to the possibility of a role of Piezo1 in confinement sensing and integrin activation in moving cells, while it deters integrin-independent cell motility, usually referred to as amoeboid movement (29, 30).

Here we aimed at exploring if Piezo1 mechanosensing plays any role during integrin-dependent chemotactic migration of human T cell migration, a patho-physiologically critical event in an ongoing immune response as well as in the steady-state. We found that deficiency of Piezo1 expression in human CD4$^+$ T cells as well as Jurkat T cells interfered with efficient integrin-dependent cellular motility in response to a chemotactic cue. We also found that Piezo1 activation constitutes a hitherto unknown membrane-intrinsic component of the 'outside-in' signaling in human T cells in response to chemokine receptor activation, that follows activation of focal adhesion kinase leading to local increase in membrane tension and mediates local recruitment of integrin molecules at the leading edge of the moving human T cells.

METHODS

siRNA-mediated Piezo1 knockdown

Human CD4+ T lymphocytes were isolated from peripheral blood of healthy donors through
magnetic immunoselection. Cells were nucleofected with control EGFP or Piezo1 siRNA using Lonza
4D nucleofector. Knockdown was checked 72 hours post-nucleofection by quantitative PCR of Piezo1
and Piezo2 (Supplementary Figure 1A). For 2D motility assay, cells were co-nucleofected with Lonza
pmax-GFP plasmid and Piezo1 siRNA and motility difference between GFP+ and GFP+ cells were
assessed. Details of methodology are available as supplemental information.

Cell motility assays

2D non-directional motility was performed on ICAM1-coated dishes in presence of recombinant
human CCL19. 3D chemotaxis was performed using ICAM1-coated 5μm transwell inserts across
CCL19 and SDF1α gradients for CD4+ T lymphocytes and Jurkat T cells, respectively. 2D tracking
analysis was performed using Fiji. Details of methodology are available as supplemental information.

Membrane tension measurements

Membrane tension of Piezo1-GFP expressing Jurkat cells was measured using non-invasive
interference reflection microscopy on fibronectin coated dishes upon addition of SDF1α. Analysis
was performed using MATLAB. Details of methodology for experiments and analyses are available as
supplemental information.

Confocal imaging

Fixed confocal imaging for Piezo1, pFAK, Akt, pAkt, LFA-1 and actin polarity in response to
chemokine stimulation was performed using Zeiss LSM confocal setup. Live cell imaging of Piezo1
mCherry and/or actin-GFP Jurkat T cells in the presence of SDF1α was performed using the same.
Polarity analysis was done on Fiji. Details of methodology for experiments and analyses are available
as supplemental information.

RESULTS

Piezo1 deficiency abrogates integrin-dependent chemotactic motility in human T cells

We first assessed the role of Piezo1 on integrin-dependent non-directional motility of primary
human CD4+ T cells. We used GFP plasmid and Piezo1 siRNA co-transfected cells to facilitate
distinction between potentially transfected and nontransfected cells during cell tracking. We
compared the motility of GFP+ T cells (presuming the transfection efficiency would lead to co-
transfection of siRNA) with the GFP- T cells in the same culture. This was confirmed by flow

cytometric assessment of Piezo1 protein content on the T cell surface as well as the whole T cells
(Supplemental figures 1B, C). Piezo1 deficiency in T cells (potentially GFP$^+$) significantly hindered
their migration in response to CCL19 in terms of its mean-squared displacement (MSD, Figure 1A-C).
Representative cell tracks of Piezo1 deficient (GFP$^+$) and control (GFP$^-$) are shown in Figure 1B and
Supplementary Figure 1D.

ICAM1-coated transwell chemotaxis assay of human CD4$^+$ T lymphocytes and Jurkat T cells also
showed significant reduction in migration upon siRNA-mediated Piezo1 knockdown, in response to
CCL19 and SDF1α gradients, respectively (Figure. 1D, E). Piezo1 inhibition with GsMTx4 also led to
significant impairment of transwell chemotaxis (Supplementary figure 1E).

Redistribution of Piezo1 toward the leading edge of migrating T cells in response to chemotactic
cue

Cell polarity is a crucial aspect of directional cell motility in response to chemokines. CD4$^+$ T cells
seeded on ICAM-1 bed showed striking polarity in Piezo1 distribution, upon stimulation with CCL19
for 15 minutes (Figure 1F, G; Supplemental video 1). Piezo1 m-Cherry expressing Jurkat T cells also
showed striking leading edge polarity in the presence of SDF1α under ICAM1 adhesion (Figure 1H,
Supplemental Video 2). In Figure 1I, particle image velocimetry (PIV) analysis of a single
representative Jurkat cell has been shown to display the dynamically polarized Piezo1 fluorescence
upon SDF1α treatment with time, as opposed to rather uniformly distributed Piezo1 signals in a
representative untreated cell.

Piezo1 redistribution in migrating T cells follow increased membrane tension in the leading edge

Given its function, Piezo1 mechanosensors may play a role in sensing mechanical cues in a moving T
cell. We utilized the non-invasive technique of interference reflection microscopy or IRM for
studying potential redistribution of membrane tension following chemokine stimulation (31, 32).
IRM allows computation of tension magnitude from local temporal membrane fluctuations (Figure.
2A and detailed Methods in Supplementary Information). After baseline imaging, Piezo1-GFP
expressing Jurkat cells were subjected to sequential fluorescence and IRM imaging post SDF1α
addition which was followed over time. Chemokine administration to Jurkat T cells led to a decrease
in temporal fluctuations and an increase in membrane tension when computed for the whole
cellular contour (Figure 2B). The increase in membrane tension was registered within few minutes

after chemokine addition and gradually declined over time (Figure 2C). Distribution of tension and

fluctuation-amplitude in cells was also affected by the action of chemokine (Supplemental figure 2A)

which was accompanied by increase in the membrane's degree of confinement and the effective

viscosity experienced (Supplemental figure 2C & D). Moreover, the tension maps showed that the

lamellipodial structures towards the leading edges of the cells had higher tension magnitudes (Figure

2D, Supplemental figure 2B). These high-tension edges are usually further emphasized at later time-

points.

Finally, we explored if local membrane tension increase was linked to Piezo1 redistribution in the

chemokine-experienced T cells. Of note here, during acquisition, there was typically a time delay of 1

minute between measurement of tension and Piezo1-GFP fluorescence. As expected, average

tension peaked within the first few minutes of chemokine addition, followed by partial decline

(Figure 2E) as seen for the whole cell. Pixel count of Piezo1-GFP intensity showed a similar pattern,

albeit, at a delayed time, following the tension kinetics. In order to account for the delayed kinetics

of tension magnitude and Piezo1-GFP pixel count, we performed a correlation between normalised

Piezo1 pixel intensity at a particular time-frame, with normalised tension magnitude at the

preceding time frame. We observed significant positive correlation between the two parameters

(Figure 2F, G). Thus, IRM-coupled fluorescence microscopy revealed an increase in local membrane

tension in the leading edge of the moving T cells in response to chemokine receptor activation that

was associated with Piezo1 redistribution to the leading edges of the cells.

Focal adhesion following chemokine receptor activation does not depend on Piezo1

Focal adhesion formation induced by chemokine signaling is a critical step in cellular migration (33,

34). Anchoring of the local sub-membrane cytoskeleton with the extracellular substrate at sites of

focal adhesion complexes increases local membrane tension, which triggers membrane recruitment

and activation of integrins, through a mechanism of 'outside-in' signalling. This is followed by

integrin-dependent 'inside-out' signaling) driven primarily by phosphoinositide 3-kinase (PI3K)-Akt-

Rho-GTPase-actin polymerization pathway (35-38). Propelling forces generated by myosin-based

contractions and consequent F-actin retrograde flow allow the cells to move alone chemokine

gradients (6,7).

While phosphorylated FAK (pFAK) showed striking polarity upon CCL19 stimulation on ICAM-1 beds

(Figure 3A, B, Supplementary video 3), Piezo1 knockdown did not have any effect on pFAK polarity

under the same conditions (Figure 3C, D). Thus, focal adhesion formation following chemokine

receptor activation was not dependent on Piezo1. FAK activation recruits adapters like paxillin and

vinculin connecting the cortical cytoskeleton to the membrane and the extracellular matrix, thereby

increasing local membrane tension (33). Inhibition of FAK activation completely abrogated Piezo1

redistribution to the leading edge, hence the polarity (Figure 3E, F). This indicated that

mechanistically, FAK activation event precedes Piezo1 redistribution to the leading edge of the

migrating human CD4$^+$ T cells.

Membrane recruitment of LFA1 on chemokine receptor activation disrupts with Piezo1 deficiency

Integrin recruitment to the leading edge occurs downstream of FAK activation and local F-actin

polymerization. As expected, CCL19 stimulation led to LFA1 recruitment to the leading edge and,

Piezo1 and LFA1 exhibited significant colocalization upon CCL19 stimulation on ICAM1 bed (Figure

3G-I; Supplemental video 4). Piezo1 downregulation abrogated LFA-1 polarity (Figure 3J, K;

Supplemental videos 5, 6). Thus, role of Piezo1 in chemotactic migration of T cells was downstream

of FAK assembly in response to chemokine receptor signaling, but preceded integrins recruitment.

These data together point to a mechanism whereby local increase in membrane tension leading

from FAK activation and assembly drives Piezo1 activation, which in turn is linked to LFA1

recruitment to the focal adhesions.

We also looked at phosphorylation of Akt downstream of PI3K activation, which in turn is triggered

upon integrin activation. On treatment with CCL19, human CD4$^+$ T cells moving on ICAM-1-coated

plates did show a prominent leading-edge polarity of non-phosphorylated and phosphorylated Akt

(pAkt) (Figure 3L control panel). But in Piezo1-deficient cells the polarity of pAkt is abrogated (Figure

3M), not affecting the polarity of non-phosphorylated AKT (Figure 3N; Supplemental videos 7, 8),

substantiating the cellular signaling defect due to non-recruitment of LFA1 to the leading edge

membrane.

Piezo1 deficiency disrupts F-actin retrograde flow in T cells despite chemokine receptor activation

Piezo1-mediated sensing of membrane tension drives calcium ion influx, which in turn has been

shown to activate calpain and F-actin polymerization (14, 19). LFA1 recruitment is driven by local

actin scaffold formation following FAK assembly, which as our data indicates is perhaps contributed

by consolidation local actin scaffold downstream of Piezo1 activation. To confirm this, we performed

Ca^{2+} imaging in Piezo1-mCherry transfected Jurkat Cells using total internal reflection fluorescence

(TIRF) microscope in response to chemokine. We found significant colocalization of the Ca^{2+} signal

with the Piezo1 signal on the chemokine treated Jurkat cells (Figure 4A, B; Supplemental figure 3A,

B). Next to confirm that activation of calpain mediates the Piezo1 function, we observed that prior

inhibition with calpain inhibitor PD150606 abrogated CCL19-induced LFA1 recruitment to the leading

edge of the T cells. (Figure 4C, D). As expected, the upstream event of pFAK polarity was not affected

by the calpain inhibition (Supplemental figure 4C, D).

While F-actin polymerization downstream of FAK is required for LFA1 recruitment, LFA1 in turn

drives a more robust F-actin polymerization at the leading edge, through Akt/Rho GTPase activation.

Leading edge F-actin undergoes retrograde flow due to myosin contractions, thereby enabling cells

to move forward. We observed focal co-localization of LFA1 with F-actin at the leading edge, along

with a denser actin-rich protrusion in the rear edge or the uropod (Figure 4C), which was abolished

upon calpain inhibition (Figure 4C, E). Hence, LFA1 recruitment may depend on local F-actin

polarization downstream of Piezo1 activation. On the other hand, failure of LFA1 recruitment and

activation halts further F-actin polarization feeding the retrograde flow of actin during migration.

Dichotomy of Piezo1 (dense leading edge distribution) and F-actin (more dense at the rear end)

distribution was clearly demonstrated in cells stimulated with chemokine (Figure 4F and

Supplemental video 9). But, no F-actin polarity was seen on downregulating Piezo1 in the T cells

(Figure 4G, H; Supplemental videos 10, 11). Live cell imaging of Jurkat T cells expressing Piezo1-

mCherry and actin-GFP on ICAM-1 coated dishes was performed in the presence or absence of

SDF1α (Supplemental figure 4E). Chemokine addition showed significant increase in Piezo1 polarity

to the front end, as shown in the earlier experiments (Supplemental figure 4F, left). But F-actin

showed a dynamic balance at both ends as expected (Supplemental figure 4F, right). Polar plots

depicting angular distribution of Piezo1 mCherry and actin-GFP (over all time-points) in the presence

or absence of SDF1α are shown in Supplemental figure 3G and H. Representative image 4I and

supplemental video 12 shows Piezo1-mCherry distribution predominantly at the leading edge and

actin-GFP eventually accumulating at the rear end in uropod-like cellular extensions in a migrating

Jurkat T cell.

Thus, our data reveals a membrane-intrinsic event, involving a critical role of Piezo1

mechanosensors, within the outside-in signaling module in response to chemokine receptor

activation in human T cells. The mechanistic model derived from our data is depicted in Figure 4J.

The Piezo1 mechanosensing links the focal adhesion assembly to integrins recruitment at the leading

edge of the human T cells migrating in response to chemokine activation and thus play a critical role

in chemotactic migration of human T cells.

DISCUSSION

The present study explored the role of membrane-intrinsic tension generation and the crucial role of

sensing this physical cue in the context of human T cell migration in response to chemokine receptor

activation. Deficiency of Piezo1 inhibited integrin-dependent migration, demonstrating crucial role

of these mechanosensors in this mode of T cell migration. As the integrin-dependent chemotactic

migration of T cells is a critically implicated in most patho-physiologic contexts, in the present study

we focused on the potential role of Piezo1 mechanosensors in this specific context.

Notably, a previous study in vivo in mice, using genetic deletion of Piezo1 in T cells, reported no

significant effect on tissue recruitment of T cells (17). Although a functional redundancy in the role

of Piezo1-mediated mechanosensing in mouse T cells warrants exclusion, especially in case of gene

deletion in developing T cells, given the considerable expression of Piezo2 as well in mouse T cells

(39). On the other hand, a role of Piezo1 activation in fibroblastic reticular cells in Peyer's patches

has been shown to regulate lymphocyte migration through tissue conduits (40). The different tissue

distribution of Piezo1 and Piezo2 expression in humans warranted exploring the role of Piezo1 in

migrating human T cells, as in humans Piezo2 is hardly expressed in the hematopoietic cells.

Non-invasive interference reflection microscopy allowed us to demonstrate dynamic change in

membrane tension at the leading edge of T cells stimulated with chemokine on ICAM1-bed which

was closely followed by dynamic recruitment Piezo1 locally. Chemokine receptor activation leads to

focal adhesion formation at the leading edge of the migrating T cells (33, 34). Our experiments

revealed that Piezo1 recruitment at the leading edge of the cells followed focal adhesion kinase

activation. Focal adhesions do increase local membrane tension as they connect the extracellular

environment with the cortical cytoskeleton of the cell (33). Piezo1 recruitment locally seems to be a

response to this membrane tension increase at focal adhesions.

'Outside-in' signaling downstream of focal adhesion formation recruits integrins to the leading edge

of the moving T cells (35, 36) which was abrogated by Piezo1 downregulation. Piezo1 has been

shown to drive Ca^{2+} influx into the T cells leading to calpain activation (14, 19). As expected, calpain

inhibition also interfered with leading edge recruitment of the integrin LFA-1. Plausibly calpain

activation drives local F-actin polymerization at the leading edge facilitating LFA-1 recruitment. This

was also supported by loss of polarity of AKT phosphorylation with Piezo1 deficiency, which is known

to be the major downstream signaling module that succeeds LFA-1 recruitment and activation (38,

41). These dysregulations in turn abrogates the retrograde flow of actin in the moving cell that is

essential for forward propulsion of the cell body (6, 7, 37, 41). Of note here, a role of ER-resident

Piezo1, in driving R-RAS-driven activation of integrins, which was again dependent on calpain

activation, was reported even before the mechanosensing function of Piezo1 was established (30).

Thus, the present study reports a hitherto unexplored role of the professional mechanosensor
Piezo1 in integrin dependent chemotactic movement of human T cells. In the cascade of leading
edge events in a moving human T cell, following chemokine receptor activation and eventually
leading to the retrograde flow of actin, focal adhesion formation precedes Piezo1 activation, while
LFA-1 recruitment is dependent on Piezo1 function.

ACKNOWLEDGEMENTS

The study was funded by Council of Scientific and Industrial Research, India (Grant no. FBR MLP-
140). D.G. is a recipient of Swarnajayanti Fellowship from Department of Science and Technology,
Government of India. The authors express their gratitude for provision of the Piezo1 constructs by
Dr. Charles Cox, Victor Chang Institute of Cardiac Research, Australia.

The authors declare no competing interests

C.S.C.L. designed, performed and analyzed most of the experiments, prepared the figures and took
part in writing the manuscript; T.M., B.S. and C.S.C.L. performed the IRM experiments; T.M. and B.S.
analyzed IRM data; C.S.C.L., P.B. and D.K.S. performed image analyses; Pu.B., M.A.H., B.P.S., J.S. and
R.D. helped with cell culture and confocal microscopy experiments; D.G. conceptualized and
supervised the study, acquired funding, designed experiments and wrote the manuscript with input
from all authors.

REFERENCES

1. Shechter R, London A, Schwartz M. Orchestrated leukocyte recruitment to immune-
privileged sites: absolute barriers versus educational gates. Nat Rev Immunol. 2013
Mar;13(3):206-18. doi: 10.1038/nri3391.
2. Krummel MF, Bartumeus F, Gérard A. T cell migration, search strategies and mechanisms.
Nat Rev Immunol. 2016 Mar;16(3):193-201. doi: 10.1038/nri.2015.16.
3. Woodland DL, Kohlmeier JE. Migration, maintenance and recall of memory T cells in
peripheral tissues. Nat Rev Immunol. 2009 Mar;9(3):153-61. doi: 10.1038/nri2496.
4. Masopust D, Schenkel JM. The integration of T cell migration, differentiation and function.
Nat Rev Immunol. 2013 May;13(5):309-20. doi: 10.1038/nri3442.
5. Fowell DJ, Kim M. The spatio-temporal control of effector T cell migration. Nat Rev Immunol.
2021 Sep;21(9):582-596. doi: 10.1038/s41577-021-00507-0.

6. Dupré L, Houmadi R, Tang C, Rey-Barroso J. T Lymphocyte Migration: An Action Movie
Starring the Actin and Associated Actors. Front Immunol. 2015 Nov 18;6:586. doi:
10.3389/fimmu.2015.00586.

7. Nordenfelt P, Moore TI, Mehta SB, Kalappurakkal JM, Swaminathan V, Koga N, Lambert TJ,
Baker D, Waters JC, Oldenbourg R, Tani T, Mayor S, Waterman CM, Springer TA. Direction of
actin flow dictates integrin LFA-1 orientation during leukocyte migration. Nat Commun. 2017
Dec 11;8(1):2047. doi: 10.1038/s41467-017-01848-y.

8. Moreau HD, Piel M, Voituriez R, Lennon-Duménil AM. Integrating Physical and Molecular
Insights on Immune Cell Migration. Trends Immunol. 2018 Aug;39(8):632-643. doi:
10.1016/j.it.2018.04.007.

9. Shannon MJ, Pineau J, Griffié J, Aaron J, Peel T, Williamson DJ, Zamoyska R, Cope AP, Cornish
GH, Owen DM. Differential nanoscale organisation of LFA-1 modulates T-cell migration. J Cell
Sci. 2019 Oct 16;133(5):jcs232991. doi: 10.1242/jcs.232991.

10. Coste B, Mathur J, Schmidt M, Earley TJ, Ranade S, Petrus MJ, Dubin AE, Patapoutian A.
Piezo1 and Piezo2 are essential components of distinct mechanically activated cation
channels. Science. 2010 Oct 1;330(6000):55-60. doi: 10.1126/science.1193270.

11. Coste B, Xiao B, Santos JS, Syeda R, Grandl J, Spencer KS, Kim SE, Schmidt M, Mathur J, Dubin
AE, Montal M, Patapoutian A. Piezo proteins are pore-forming subunits of mechanically
activated channels. Nature. 2012 Feb 19;483(7388):176-81. doi: 10.1038/nature10812.

12. Syeda R, Florendo MN, Cox CD, Kefauver JM, Santos JS, Martinac B, Patapoutian A. Piezo1
Channels Are Inherently Mechanosensitive. Cell Rep. 2016 Nov 8;17(7):1739-1746. doi:
10.1016/j.celrep.2016.10.033.

13. Murthy SE, Dubin AE, Patapoutian A. Piezos thrive under pressure: mechanically activated
ion channels in health and disease. Nat Rev Mol Cell Biol. 2017 Dec;18(12):771-783. doi:
10.1038/nrm.2017.92. Epub 2017 Oct 4. PMID: 28974772.

14. Liu CSC, Raychaudhuri D, Paul B, Chakrabarty Y, Ghosh AR, Rahaman O, Talukdar A, Ganguly
D. Cutting Edge: Piezo1 Mechanosensors Optimize Human T Cell Activation. J Immunol. 2018
Feb 15;200(4):1255-1260. doi: 10.4049/jimmunol.1701118.

15. Atcha H, Jairaman A, Holt JR, Meli VS, Nagalla RR, Veerasubramanian PK, Brumm KT, Lim HE,
Othy S, Cahalan MD, Pathak MM, Liu WF. Mechanically activated ion channel Piezo1
modulates macrophage polarization and stiffness sensing. Nat Commun. 2021 May
31;12(1):3256. doi: 10.1038/s41467-021-23482-5.

16. Geng J, Shi Y, Zhang J, Yang B, Wang P, Yuan W, Zhao H, Li J, Qin F, Hong L, Xie C, Deng X, Sun
Y, Wu C, Chen L, Zhou D. TLR4 signalling via Piezo1 engages and enhances the macrophage

mediated host response during bacterial infection. Nat Commun. 2021 Jun 10;12(1):3519. doi: 10.1038/s41467-021-23683-y.

17. Jairaman A, Othy S, Dynes JL, Yeromin AV, Zavala A, Greenberg ML, Nourse JL, Holt JR, Cahalan SM, Marangoni F, Parker I, Pathak MM, Cahalan MD. Piezo1 channels restrain regulatory T cells but are dispensable for effector CD4+ T cell responses. Sci Adv. 2021 Jul 7;7(28):eabg5859. doi: 10.1126/sciadv.abg5859.

18. Jäntti H, Sitnikova V, Ishchenko Y, Shakirzyanova A, Giudice L, Ugidos IF, Gómez-Budia M, Korvenlaita N, Ohtonen S, Belaya I, Fazaludeen F, Mikhailov N, Gotkiewicz M, Ketola K, Lehtonen Š, Koistinaho J, Kanninen KM, Hernández D, Pébay A, Giugno R, Korhonen P, Giniatullin R, Malm T. Microglial amyloid beta clearance is driven by PIEZO1 channels. J Neuroinflammation. 2022 Jun 15;19(1):147. doi: 10.1186/s12974-022-02486-y.

19. Liu CSC, Ganguly D. Mechanical Cues for T Cell Activation: Role of Piezo1 Mechanosensors. Crit Rev Immunol. 2019;39(1):15-38. doi: 10.1615/CritRevImmunol.2019029595.

20. Ma S, Dubin AE, Zhang Y, Mousavi SAR, Wang Y, Coombs AM, Loud M, Andolfo I, Patapoutian A. A role of PIEZO1 in iron metabolism in mice and humans. Cell. 2021 Feb 18;184(4):969-982.e13. doi: 10.1016/j.cell.2021.01.024.

21. He Y, Deng B, Liu S, Luo S, Ning Y, Pan X, Wan R, Chen Y, Zhang Z, Jiang J, Xu H, Xia M, Li J. Myeloid Piezo1 Deletion Protects Renal Fibrosis by Restraining Macrophage Infiltration and Activation. Hypertension. 2022 May;79(5):918-931. doi: 10.1161/HYPERTENSIONAHA.121.18750.

22. Huang Z, Sun Z, Zhang X, Niu K, Wang Y, Zheng J, Li H, Liu Y. Loss of stretch-activated channels, PIEZOs, accelerates non-small cell lung cancer progression and cell migration. Biosci Rep. 2019 Mar 22;39(3): BSR20181679. doi: 10.1042/BSR20181679.

23. Chubinskiy-Nadezhdin VI, Vasileva VY, Vassilieva IO, Sudarikova AV, Morachevskaya EA, Negulyaev YA. Agonist-induced Piezo1 activation suppresses migration of transformed fibroblasts. Biochem Biophys Res Commun. 2019 Jun 18;514(1):173-179. doi: 10.1016/j.bbrc.2019.04.139.

24. Mousawi F, Peng H, Li J, Ponnambalam S, Roger S, Zhao H, Yang X, Jiang LH. Chemical activation of the Piezo1 channel drives mesenchymal stem cell migration via inducing ATP release and activation of P2 receptor purinergic signaling. Stem Cells. 2020 Mar;38(3):410-421. doi: 10.1002/stem.3114.

25. Wang X, Cheng G, Miao Y, Qiu F, Bai L, Gao Z, Huang Y, Dong L, Niu X, Wang X, Li Y, Tang H, Xu Y, Song X. Piezo type mechanosensitive ion channel component 1 facilitates gastric

cancer omentum metastasis. J Cell Mol Med. 2021 Feb;25(4):2238-2253. doi: 10.1111/jcmm.16217.

26. Gao L, Ji Y, Wang L, He M, Yang X, Qiu Y, Sun X, Ji Z, Yang G, Zhang J, Li S, Dai L, Zhang L. Suppression of Esophageal Squamous Cell Carcinoma Development by Mechanosensitive Protein Piezo1 Downregulation. ACS Omega. 2021 Apr 12;6(15):10196-10206. doi: 10.1021/acsomega.1c00505.

27. Holt JR, Zeng WZ, Evans EL, Woo SH, Ma S, Abuwarda H, Loud M, Patapoutian A, Pathak MM. Spatiotemporal dynamics of PIEZO1 localization controls keratinocyte migration during wound healing. Elife. 2021 Sep 27;10:e65415. doi: 10.7554/eLife.65415.

28. Velasco-Estevez M, Koch N, Klejbor I, Caratis F, Rutkowska A. Mechanoreceptor Piezo1 Is Downregulated in Multiple Sclerosis Brain and Is Involved in the Maturation and Migration of Oligodendrocytes in vitro. Front Cell Neurosci. 2022 May 26;16:914985. doi: 10.3389/fncel.2022.914985.

29. Hung WC, Yang JR, Yankaskas CL, Wong BS, Wu PH, Pardo-Pastor C, Serra SA, Chiang MJ, Gu Z, Wirtz D, Valverde MA, Yang JT, Zhang J, Konstantopoulos K. Confinement Sensing and Signal Optimization via Piezo1/PKA and Myosin II Pathways. Cell Rep. 2016 May 17;15(7):1430-1441. doi: 10.1016/j.celrep.2016.04.035.

30. McHugh BJ, Buttery R, Lad Y, Banks S, Haslett C, Sethi T. Integrin activation by Fam38A uses a novel mechanism of R-Ras targeting to the endoplasmic reticulum. J Cell Sci. 2010 Jan 1;123(Pt 1):51-61. doi: 10.1242/jcs.056424.

31. Barr VA, Bunnell SC. Interference reflection microscopy. Curr Protoc Cell Biol. 2009 Dec; Chapter 4: Unit 4.23. doi: 10.1002/0471143030.cb0423s45.

32. Chakraborty M, Mukherjee B, Nalinakshan N, Biswas A, Nayak RK, Sinha B. Effect of heterogeneous substrate adhesivity of follower cells on speed and tension profile of leader cells in primary keratocyte collective cell migration. Biol Open. 2022 Mar 15;11(3):bio058893. doi: 10.1242/bio.058893.

33. Mitra SK, Hanson DA, Schlaepfer DD. Focal adhesion kinase: in command and control of cell motility. Nat Rev Mol Cell Biol. 2005 Jan;6(1):56-68. doi: 10.1038/nrm1549.

34. Katoh K. FAK-Dependent Cell Motility and Cell Elongation. Cells. 2020 Jan 12;9(1):192. doi: 10.3390/cells9010192.

35. Yamashiro S, Watanabe N. A new link between the retrograde actin flow and focal adhesions. J Biochem. 2014 Nov;156(5):239-48. doi: 10.1093/jb/mvu053.

36. Martino F, Perestrelo AR, Vinarský V, Pagliari S, Forte G. Cellular Mechanotransduction: From Tension to Function. Front Physiol. 2018 Jul 5;9:824. doi: 10.3389/fphys.2018.00824.

386   37. Sánchez-Martín L, Sánchez-Sánchez N, Gutiérrez-López MD, Rojo AI, Vicente-Manzanares M,
387   Pérez-Alvarez MJ, Sánchez-Mateos P, Bustelo XR, Cuadrado A, Sánchez-Madrid F, Rodríguez-
388   Fernández JL, Cabañas C. Signaling through the leukocyte integrin LFA-1 in T cells induces a
389   transient activation of Rac-1 that is regulated by Vav and PI3K/Akt-1. J Biol Chem. 2004 Apr
390   16;279(16):16194-205. doi: 10.1074/jbc.M400905200.

391   38. Roy NH, Kim SHJ, Buffone A Jr, Blumenthal D, Huang B, Agarwal S, Schwartzberg PL, Hammer
392   DA, Burkhardt JK. LFA-1 signals to promote actin polymerization and upstream migration in T
393   cells. J Cell Sci. 2020 Sep 9;133(17):jcs248328. doi: 10.1242/jcs.248328.

394   39. Ganguly D. Can Piezo2 substitute Piezo1 function in the context of T cell-specific deletion in
395   mice? Sci Adv, e-Letter published Sep 8, 2021, in response to Jairaman A et al.,

396   40. Chang JE, Buechler MB, Gressier E, Turley SJ, Carroll MC. Mechanosensing by Peyer's patch
397   stroma regulates lymphocyte migration and mucosal antibody responses. Nat Immunol.
398   2019 Nov;20(11):1506-1516. doi: 10.1038/s41590-019-0505-z.

399   41. Roy NH, MacKay JL, Robertson TF, Hammer DA, Burkhardt JK. Crk adaptor proteins mediate
400   actin-dependent T cell migration and mechanosensing induced by the integrin LFA-1. Sci
401   Signal. 2018 Dec 11;11(560):eaat3178. doi: 10.1126/scisignal.aat3178.

402   42. Limozin L, Sengupta K. Quantitative reflection interference contrast microscopy (RICM) in
403   soft matter and cell adhesion. Chemphyschem. 2009 Nov 9;10(16):2752-68.

404   43. Biswas A, Alex A, Sinha B. Mapping Cell Membrane Fluctuations Reveals Their Active
405   Regulation and Transient Heterogeneities. Biophys J. 2017 Oct 17;113(8):1768-1781. doi:
406   10.1016/j.bpj.2017.08.041.

407   44. Biswas A, Kashyap P, Datta S, Sengupta T, Sinha B. Cholesterol Depletion by MβCD Enhances
408   Cell Membrane Tension and Its Variations-Reducing Integrity. Biophys J. 2019 Apr
409   23;116(8):1456-1468.

410   45. Shiba H, Noguchi H, Fournier JB. Monte Carlo study of the frame, fluctuation and internal
411   tensions of fluctuating membranes with fixed area. Soft Matter. 2016 Feb 28;12(8):2373-80.
412   doi: 10.1039/c5sm01900a.

 FIGURE LEGENDS

Figure 1. Piezo1 deficiency abrogates integrin-dependent motility in human T cells which is
mediated through redistribution of Piezo1 at the leading edge in response to chemokine
stimulation.

A. MSD versus time calculated for GFP^{+} (potential Piezo1-knockdown cells) and GFP^{-} (potential
control cells) in GFP plasmid and Piezo1 siRNA co-transfected human CD4^{+} T lymphocytes, that were
allowed to migrate in presence of recombinant CCL19 on ICAM1-coated dishes. **B.** Representative
tracks of GFP^{+} and GFP^{-} CD4^{+} T lymphocytes. Comparisons of **C.** % GFP^{+} cells after 72 hours of
nucleofection **D & E.** 3D transwell migration assay of siRNA-transfected primary human CD4^{+} T
lymphocytes (D) and Jurkat T cells (E), respectively. Data representative of at least 3 independent
experiments. **F.** Representative confocal images of Piezo1 distribution in fixed untreated and CCL19-
treated CD4^{+} T lymphocytes. 63X oil magnification. **G.** Comparison between Piezo1 polarity index
calculated for fixed, stained, human CD4^{+} T lymphocytes with or without 0.5µg/ml CCL19 treatment
for 20-30 minutes. n > 510 random cells, each. **H.** Piezo1 polarity index calculated for Jurkat cells,
expressing mCherry-tagged Piezo1 during live-cell tracking in the presence of recombinant SDF1α.
Each dot represents polarity index of each cell, averaged over all the time-frames. **I.** Representative
time kinetics of particle image velocimetry (PIV) analysis of Piezo1-mcherry transfected Jurkat cells,
allowed to move on ICAM-coated dishes in the presence of recombinant SDF1α. Top panel: No
chemokine. Bottom Panel: SDF1α.

Figure 2. Piezo1 redistribution in migrating T cells follow increased membrane tension in the
leading edge

A. Schematic illustration of interference reflection microscopy imaging setup. **B.** Probability
distribution of amplitude of temporal fluctuations (SD$_{time}$, left) and tension (right) before and after
addition of chemokines. **C.** Temporal trajectory of tension of cells before and after addition of
chemokine with ** indicating significant difference from control. **D.** Representative IRM images of
Jurkat cells (top), corresponding tension maps (middle) and R^{2} maps (bottom), before and after
chemokine treatment. **E** Correlation between normalized Piezo intensity, tension magnitudes with
time. **F.** Scatter plot to show correlation between normalized pixel count of Piezo1 and normalized
tension. Piezo1 intensity at specific time-points were correlated with tension magnitudes at the
preceding timepoints. Normalization has been done such that the maximum value is set to 1 and
others are accordingly scaled. **G.** Colour-coded Piezo1-GFP intensity map (left), epifluorescence

image (middle), and tension map (right) of representative Piezo1-GFP expressing Jurkat cell, after 30

minutes of 0.1µg/ml of SDF1α treatment.

Figure 3. Focal adhesion following chemokine receptor activation does not depend on Piezo1 while

membrane recruitment of LFA1 on chemokine receptor activation disrupts with Piezo1 deficiency.

A. Representative confocal images of human CD4$^+$ T lymphocytes, fixed and stained for phospho-FAK

(pFAK) under untreated and CCL19-treated conditions. **B.** Increased polarity of pFAK upon

chemokine stimulation of CD4$^+$ T lymphocytes as compared to unstimulated controls. N > 120 cells,

each. **C.** Representative confocal images of pFAK distribution in control and Piezo1 siRNA-transfected

CD4$^+$ T lymphocytes stimulated with chemokine. **D.** Comparison of pFAK polarity in CCL19-stimulated

control and Piezo1-knockdown CD4$^+$ T lymphocytes. n > 450 random cells, each. **E.** Representative

confocal images of stained Piezo1 in CD4$^+$ T cells stimulated with CCL19 in the presence or absence

of FAK inhibitor 14. **F.** Effect of FAK inhibition on Piezo1 polarity in CD4$^+$ T lymphocytes stimulated

with recombinant CCL19 versus untreated cells. n > 600 random cells, each. Data represented from

at least 3 independent experiments. **G.** Representative confocal images of fixed, immunostained

Piezo1 and LFA1 in unstimulated and CCL19-stimulated CD4$^+$ T lymphocytes. **H.** Increased LFA1

polarity in response to recombinant CCL19 stimulation in CD4$^+$ T cells. n > 295 random cells, each. **I.**

Manders' co-localisation analysis of Piezo1 and LFA1 in untreated and CCL19-treated CD4$^+$ T

lymphocytes. **J.** Comparison of LFA1 polarity in chemokine-treated control and Piezo1 knockdown

CD4$^+$ T lymphocytes. **K.** Representative confocal stained images of LFA1 polarity of CCL19-treated

control and Piezo1-knockdown CD4$^+$ T lymphocytes. n > 490 random cells, each. **L.** Representative

confocal images of phospho-Akt (pAkt) and Akt distribution upon chemokine treatment of control

and Piezo1-knockdown cells. **M & N.** Quantitative analyses of pAkt (M) and Akt (N) polarity in control

and Piezo1-knockdown cells after CCL19-treatment. n > 580 random cells, each. Data represented

from at least 3 independent experiments.

Figure 4. Piezo1 deficiency disrupts F-actin retrograde flow in T cells despite chemokine receptor

activation.

A. Representative TIRF image of a Jurkat cell stained with Fluo-3 AM and expressing Piezo1 mCherry.

Lower panel: zoomed in image and overlap of binary images of objects detected from Ca^{2+} and

Piezo1 signals as well as simulated randomly placed puncta having similar total (in the zoomed in

section) and average area as that of the Ca^{2+} puncta detected in the real images. Overlap RGB image

shows the overlap between Real Ca^{2+} puncta (red) and Piezo1 puncta (green), overlap puncta

(yellow), simulated Ca^{2+} puncta (blue). **B.** Comparison between percentage Ca^{2+} colocalization of

simulated data and real data. * denotes *p*-value < 0.05 calculated using Mann Whitney U test. N_{cell}=

52 (13 cells x 4 time points). N_{ROIs}= 52. **C.** Representative confocal images of LFA1 and actin

distribution in CCL19-stimulated $CD4^+$ T lymphocytes with or without inhibition of calpain.

Quantitative comparisons of **D.** LFA1 and **E.** actin polarity upon chemokine stimulation, in the

presence or absence of calpain pre-inhibition. N > 290 random cells, each. **F.** Representative fixed

confocal images of actin and Piezo1 distribution in $CD4^+$ T lymphocytes after 30 mins of 0.5µg/ml

recombinant CCL19 treatment. **G.** Quantitative comparison of actin polarity in control and Piezo1-

knockdown $CD4^+$ T lymphocytes cells after 30 mins of chemokine treatment. n > 300 random cells,

each. **H.** Representative fixed confocal images of actin distribution in chemokine-treated, control

and Piezo1-knockdown $CD4^+$ T lymphocytes. **I.** A snapshot of time-lapse imaging of Piezo1-mCherry

and actin-GFP expressing Jurkat cell treated with 0.1µg/ml of SDF1α. Image is maximum Z-projection

of the cell at 63X/1.40 magnification. **J.** Proposed model suggests chemokine receptor activation in

human T cells lead to focal adhesion kinase activation and focal adhesion formation. Focal adhesions

lead to localized increase in membrane tension at the leading edge plasma membrane which leads

to Piezo1 recruitment and activation. Piezo1 activation leads to calpain activation which potentially

drives further cytoskeletal consolidation to recruit integrin LFA1. LFA1 recruitment and activation

lead to phosphorylation of AKT and downstream signaling eventually driving the retrograde actin

flow in migrating human T cells.

Figure 1

Figure 2

Figure 3

Figure 4