Piezo1 mechanosensing regulates integrin-dependent chemotactic

migration in human T cells

Chinky Shiu Chen Liu[1], Tithi Mandal[2], Parijat Biswas[3], Md. Asmaul Hoque[1,4], Purbita Bandopadhyay[1,4],

Bishnu Prasad Sinha[1,4], Jafar Sarif[1,4], Ranit D'Rozario[1,4], Deepak Kumar Sinha[3], Bidisha Sinha[2], Dipyaman

Ganguly[1]*

[1]IICB-Translational Research Unit of Excellence, CSIR-Indian Institute of Chemical Biology, Kolkata,

India.

[2]Department of Biological Sciences, Indian Institute of Science Education and Research, Kolkata, India.

[3] Department of Biological Sciences, Indian Association for Cultivation of Science, Kolkata, India.

[4]Academy of Scientific and Innovative Research, Ghaziabad, India.

Correspondence: dipyaman@iicb.res.in

Running title: Piezo1 mechanosensing in human T cell chemotaxis

Keywords: Human T cells, Piezo1, Mechanosensing, Migration, Chemokine, Chemotaxis, Focal

adhesion kinase, LFA-1, F-actin, Confocal Microscopy, Interference Reflection Microscopy

ABSTRACT

T cells are crucial for efficient antigen-specific immune responses and thus their migration within the

body, to inflamed tissues from circulating blood or to secondary lymphoid organs, play a very critical

role. T cell extravasation in inflamed tissues depends on chemotactic cues and interaction between

endothelial adhesion molecules and cellular integrins. A migrating T cell is expected to sense diverse

external and membrane-intrinsic mechano-physical cues, but molecular mechanisms of such

mechanosensing in cell migration are not established. We explored if the professional mechanosensor

Piezo1 play any role during integrin-dependent chemotaxis of human T cells. We found that deficiency

of Piezo1 in human T cells interfered with integrin-dependent cellular motility on ICAM-1-coated

surface. Piezo1 recruitment at the leading edge of moving T cells is dependent on and follows focal

adhesion formation at the leading edge and local increase in membrane tension on chemokine

receptor activation. Piezo1 recruitment and activation, followed by calcium influx and calpain

activation, in turn are crucial for the integrin LFA-1 (CD11a/CD18) recruitment at the leading edge of

the chemotactic human T cells. Thus, we find that Piezo1 activation in response to local mechanical

cues constitutes a membrane-intrinsic component of the 'outside-in' signaling in human T cells,

migrating in response to chemokines, that mediates integrin recruitment to the leading edge.

INTRODUCTION

Efficient T cell migration from blood circulation to secondary lymphoid organs and inflamed tissue is paramount for optimal tissue antigen sampling and effector functions during adaptive immune response (1-4). The process is dependent on chemotactic cues, interaction between endothelial adhesion molecules and T cell integrins, and an intricate acto-myosin dynamics (5). Generation of mechanical force is critical to propel migrating cells (6-9). Mechanosensing of the substrate and increase in cellular membrane tension is also intuitively apparent in a migrating T cell, although mechanistic details are not explored to a great extent.

Piezo1 and Piezo2 ion channels are evolutionarily conserved professional mechanosensors that can sense increase in membrane tension mostly independent of other protein-protein interactions (10-12). Mechanosensing by Piezo1 has been demonstrated to play critical role in diverse patho-physiologic contexts given its wide tissue expression including the immunocellular compartment (13-21). We previously identified crucial role of this specialized mechanotransducer in optimal TCR triggering (14, 19). The immunocellular expression of Piezo1 thus makes it a strong candidate for mechanosensing at the plasma membrane to regulate motility of immune cells. Piezo1 shows differential involvement in cellular migration depending on context (22- 28). Previous studies point to the possibility of a role of Piezo1 in confinement sensing and integrin activation in moving cells, while it deters integrin-independent cell motility, usually referred to as amoeboid movement (29, 30).

Here we aimed at exploring if Piezo1 mechanosensing plays any role during integrin-dependent chemotactic migration of human T cell migration, a patho-physiologically critical event in an ongoing immune response as well as in the steady-state. We found that deficiency of Piezo1 expression in human CD4[+] T cells as well as Jurkat T cells interfered with efficient integrin-dependent cellular motility in response to a chemotactic cue. We also found that Piezo1 activation constitutes a hitherto unknown membrane-intrinsic component of the 'outside-in' signaling in human T cells in response to chemokine receptor activation, that follows activation of focal adhesion kinase leading to local increase in membrane tension and mediates local recruitment of integrin molecules at the leading edge of the moving human T cells.

RESULTS

Piezo1 deficiency abrogates integrin-dependent chemotactic motility in human T cells

We first assessed the role of Piezo1 on integrin-dependent, non-directional motility of primary human

CD4[+] T cells. We used GFP plasmid and Piezo1 siRNA co-transfected cells to facilitate distinction

between potentially transfected and untransfected cells during cell tracking. We compared the

motility of GFP[+] T cells (presuming they had a good transfection efficiency and thus will also have the

siRNA) with the GFP[-] T cells in the same culture. This was confirmed by flow cytometric assessment of

Piezo1 protein content on the T cell surface as well as the whole T cells (Supplemental figures 1B, C).

Piezo1 deficiency in T cells (potentially GFP[+]) significantly hindered their migration in response to

CCL19 in terms of its mean-squared displacement (MSD, Figure 1A-C). Representative cell tracks of

Piezo1 deficient (GFP[+]) and control (GFP[-]) are shown in Figure 1B, supplementary figure 1D and

supplemental video 1. These results were further confirmed in a separate experiment where Piezo1

KD cells showed significant reduction in motility compared to control siRNA-transfected cells

(Supplementary figure 1E).

ICAM1-coated transwell chemotaxis assay of human CD4[+] T lymphocytes and Jurkat T cells also

showed significant reduction in migration upon siRNA-mediated Piezo1 knockdown, in response to

CCL19 and SDF1α gradients, respectively (Figure. 1D, E). Piezo1 inhibition with GsMTx4 also led to

significant impairment of transwell chemotaxis (Supplementary figure 1F).

Redistribution of Piezo1 toward the leading edge of migrating T cells in response to chemotactic cue

Cell polarity is a crucial aspect of directional cell motility in response to chemokines. CD4[+] T cells

seeded on ICAM-1 bed showed striking polarity in Piezo1 distribution, upon stimulation with CCL19

for 15 minutes (Figure 1F, G; Supplemental video 2). Piezo1 m-Cherry expressing Jurkat T cells also

showed striking leading edge polarity in the presence of SDF1α under ICAM1 adhesion (Figure 1H). In

Figure 1I, particle image velocimetry (PIV) analysis of a single representative Jurkat cell has been

shown to display the dynamically polarized Piezo1 fluorescence upon SDF1α treatment with time, as

opposed to rather uniformly distributed Piezo1 signals in a representative untreated cell. The

continuous redistribution of Piezo1 within the motile T cells towards the leading edge can also be

appreciated in the representative moving image provided (Supplemental video 3).

Piezo1 redistribution in migrating T cells follow increased membrane tension in the leading edge

Piezo1 is a professional mechanosensor ion channel, which senses increase in plasma membrane

tension and responds by driving calcium influx into the cells (14, 19, 31). As T cells migrate on substrate

by extending membrane processes, it is intuitive to hypothesize that a component of mechanosensing

is incorporated in the mechanism of cellular movement. The data above, on a critical role of Piezo1 in

migrating human T cells, also led us to hypothesize that Piezo1 mechanosensors may play a role in

sensing mechanical cues in a moving T cell. So, first we aimed at confirming that there is polarized

increase in tension in the leading edge plasma membrane of human T cells, moving in response to a

chemotactic cue.

Cellular membrane tension can be measured by various techniques. However, most of the usual

techniques, viz. atomic force microscopy or tension measurement using optical tweezers, are invasive

and interfere with the membrane dynamics, as well as are difficult to employ on moving cells. We

utilized the non-invasive technique of interference reflection microscopy or IRM for studying potential

redistribution of membrane tension following chemokine stimulation (32, 33). In IRM technique,

steady-state local fluctuations in the plasma membrane is imaged in cells adhered to a glass coverslip.

The interference patterns generated due to continuously changing local distance between the

coverslip and the plasma membrane are computed to derive the magnitude of local tension in the

membrane – lower temporal fluctuations represent an increase in membrane tension (Figure. 2A).

After baseline imaging, Piezo1-GFP expressing Jurkat cells were subjected to sequential fluorescence

and IRM imaging post SDF1α addition which was followed over time. Chemokine administration to

Jurkat T cells led to a decrease in temporal fluctuations and an increase in membrane tension when

computed for the whole cellular contour (Figure 2B). The increase in membrane tension was

registered within few minutes after chemokine addition and gradually declined over time (Figure 2C).

Distribution of tension and fluctuation-amplitude in cells was also affected by the action of chemokine

(Supplemental figure 2A) which was accompanied by increase in the membrane's degree of

confinement and the effective viscosity experienced (Supplemental figure 2C & D). Moreover, the

tension maps showed that the lamellipodial structures towards the leading edges of the cells had

higher tension magnitudes (Figure 2D, Supplemental figure 2B). These high-tension edges are usually

further emphasized at later time-points.

Finally, we explored if local membrane tension increase was linked to Piezo1 redistribution in the

chemokine-experienced T cells. Of note here, during acquisition, there is typically a time delay of 1

minute between measurement of tension and Piezo1-GFP fluorescence. As expected, average tension

peaked within the first few minutes of chemokine addition, followed by partial decline (Figure 2E) as

seen for the whole cell. Pixel count of Piezo1-GFP intensity showed a similar pattern, albeit, at a

delayed time, following the tension kinetics. In order to account for the delayed kinetics of tension

magnitude and Piezo1-GFP pixel count, we performed a correlation between normalised Piezo1 pixel

intensity at a particular time-frame, with normalised tension magnitude at the preceding time frame.

We observed significant positive correlation between the two parameters (Figure 2F, G). Thus, IRM-

coupled fluorescence microscopy revealed an increase in local membrane tension in the leading edge

of the moving T cells in response to chemokine receptor activation that was associated with Piezo1

redistribution to the leading edges of the cells.

Focal adhesion following chemokine receptor activation does not depend on Piezo1

Focal adhesion formation is a critical step in cellular migration and activated phosphorylated focal

adhesion kinases link extracellular physical cues to the cytoskeletal actomyosin scaffold (34, 35).

Chemokine signaling induces phosphorylation of focal adhesion kinase (FAK) and membrane

recruitment of the FAK complex forming focal adhesions with the extracellular substrate. FAK recruits

adapter proteins like paxillin to anchor with the local sub-membrane cytoskeleton made by de novo

F-actin polymerization. Membrane-recruited FAK complexes anchored to cytoskeleton increases local

membrane tension as well as recruit integrins to the membrane (36, 6, 37). This so-called 'outside-in'

signaling, whereby membrane recruitment of integrins (e.g.; LFA1 in T cells) is initiated, is followed by

integrin signaling ('inside-out' signaling) driven primarily by phosphoinositide 3-kinase (PI3K)-driven

phosphorylation of Akt and downstream activation of Rho GTPases which in turn drive extensive F-

actin polymerization (38, 39). Retrograde flow of the F-actin to the rear end of the moving cells,

through myosin-based contractions, generates the propelling force for cells to move forward following

the chemokine gradient (6, 7).

Finding a critical role of Piezo1 in human T cell migration and a membrane tension-dependent

redistribution of Piezo1 to the leading edge of the moving cells led us explore if FAK recruitment to

the leading edge in response to chemokine stimulation was dependent on Piezo1. While

phosphorylated FAK (pFAK) showed striking polarity upon CCL19 stimulation on ICAM-1 beds (Figure

3A, B, Supplementary video 4), Piezo1 knockdown did not have any effect on pFAK polarity under the

same conditions (Figure 3C, D). We observed similar results upon examining distribution of paxillin

which is another critical component of focal adhesions. Paxillin showed increased polarity upon CCL19

stimulation (Figure 3E, F) and its polarity remained unaffected upon Piezo1 KD (Figure 3G, H). We also

did not observe any changes in pFAK and paxillin signal intensity upon Piezo1 knockdown in CCL19-

stimulated cells (Supplementary figure 3A, B, respectively). However, pFAK showed striking co-

localization with Piezo1 upon CCL19 stimulation (Supplementary figure 3C). Thus, focal adhesion

formation following chemokine receptor activation was not dependent on Piezo1. FAK activation

recruits adapters like paxillin and vinculin connecting the cortical cytoskeleton to the membrane and

the extracellular matrix, thereby increasing local membrane tension (34). Inhibition of FAK activation

completely abrogated Piezo1 redistribution to the leading edge, hence the polarity (Figure 3I, J). This

indicated that mechanistically, FAK activation event precedes Piezo1 redistribution to the leading edge

of the migrating human CD4$^+$ T cells.

Membrane recruitment of LFA1 on chemokine receptor activation disrupts with Piezo1 deficiency

Integrin recruitment to the leading edge occurs downstream of FAK activation and local F-actin

polymerization. As expected, CCL19 stimulation led to LFA1 recruitment to the leading edge upon

CCL19 stimulation on ICAM1 bed (Figure 4A, B; Supplemental video 5). Moreover, Piezo1 and LFA1

(CD11a) exhibited significant colocalization upon chemokine stimulation (Figure 4C). Piezo1

downregulation abrogated LFA-1 polarity (Figure 4D, E; Supplemental videos 6, 7). Thus, role of Piezo1

in chemotactic migration of T cells was downstream of FAK assembly in response to chemokine

receptor signaling, but preceded integrins recruitment. These data together point to a mechanism

whereby local increase in membrane tension leading from FAK activation and assembly drives Piezo1

activation, which in turn is linked to LFA1 recruitment to the focal adhesions.

To further confirm this critical role of Piezo1 on LFA1 recruitment, we looked at the major signaling

event downstream of LFA1 recruitment to the leading edge, i.e., phosphorylation of Akt downstream

of PI3K activation, which in turn is triggered upon integrin activation. On treatment with CCL19, human

CD4$^+$ T cells moving on ICAM-1-coated plates did show a prominent leading-edge polarity of non-

phosphorylated and phosphorylated Akt (pAkt) (Figure 4F, control panel). But in Piezo1-deficient cells

the polarity of pAkt is abrogated (Figure 4G), not affecting the polarity of non-phosphorylated AKT

(Figure 4H; Supplemental videos 8, 9), substantiating the cellular signaling defect due to non-

recruitment of LFA1 to the leading edge membrane.

Piezo1 deficiency disrupts F-actin retrograde flow in T cells despite chemokine receptor activation

Piezo1-mediated sensing of membrane tension drives calcium ion influx, which in turn has been shown

to activate calpain and F-actin polymerization (14, 19). LFA1 recruitment is driven by local actin

scaffold formation following FAK assembly, which as our data indicates is perhaps contributed by

consolidation local actin scaffold downstream of Piezo1 activation. To confirm this, we performed Ca^{2+}

imaging in Piezo1-mCherry transfected Jurkat Cells using total internal reflection fluorescence (TIRF)

microscope in response to chemokine. We found significant colocalization of the Ca^{2+} signal with the

Piezo1 signal on the chemokine treated Jurkat cells (Figure 5A, B; Supplemental figure 4A, B). Next to

confirm that activation of calpain mediates the Piezo1 function, we observed that prior inhibition with

calpain inhibitor PD150606 abrogated CCL19-induced LFA1 recruitment to the leading edge of the T

cells (Figure 5C, D). As expected, the upstream event of pFAK polarity was not affected by the calpain

inhibition (Supplemental figure 4C, D).

While F-actin polymerization downstream of FAK is required for LFA1 recruitment, LFA1 in turn drives

a more robust F-actin polymerization at the leading edge, through Akt/Rho GTPase activation. Leading

edge F-actin undergoes retrograde flow due to myosin contractions, thereby enabling cells to move

forward. Thus, both at the leading edge lamellipodia and the rear edge F-actin redistribution can be

documented in a motile cell. We observed focal co-localization of LFA1 with F-actin at the leading

edge, along with a denser actin-rich protrusion in the rear edge or the uropod (Figure 5C), which was

abolished upon calpain inhibition (Figure 5C, E). Hence, LFA1 recruitment may depend on local F-actin

polarization downstream of Piezo1 activation. On the other hand, failure of LFA1 recruitment and

activation halts further F-actin polarization feeding the retrograde flow of actin during migration.

Dichotomy of Piezo1 (dense leading edge distribution) and F-actin (more dense at the rear end)

distribution was clearly demonstrated in cells stimulated with chemokine (Figure 6A and Supplemental

video 10). But, no F-actin polarity was seen on downregulating Piezo1 in the T cells (Figure 6B, C;

Supplemental videos 11, 12). Live cell imaging of Jurkat T cells expressing Piezo1-mCherry and actin-

GFP on ICAM-1 coated dishes was performed in the presence or absence of SDF1α (Supplemental

figure 5A). Chemokine addition showed significant increase in Piezo1 polarity to the front end, as

shown in the earlier experiments (Figure 6E, Supplemental figure 5B, left) upon chemokine stimulation

when compared to untreated cells (Figure 6D, Supplemental figure 5B, left). But F-actin showed a

dynamic balance at both ends as expected (Figure 6E, Supplemental figure 5B, right). Polar plots

depicting angular distribution of Piezo1 mCherry and actin-GFP (over all time-points) in the presence

or absence of SDF1α are shown in Supplemental figure 5C. Representative image 6F and supplemental

video 13 shows Piezo1-mCherry distribution predominantly at the leading edge and actin-GFP

eventually accumulating at the rear end in uropod-like cellular extensions in a migrating Jurkat T cell.

Thus, our data reveals a membrane-intrinsic event, involving a critical role of Piezo1 mechanosensors,

within the outside-in signaling module in response to chemokine receptor activation in human T cells.

The mechanistic model derived from our data is depicted in Figure 7. The Piezo1 mechanosensing links

the focal adhesion assembly to integrins recruitment at the leading edge of the human T cells

migrating in response to chemokine activation and thus play a critical role in chemotactic migration of

human T cells.

DISCUSSION

DISCUSSION

The present study explored the role of membrane-intrinsic tension generation and the crucial role of

sensing this physical cue in the context of human T cell migration in response to chemokine receptor

activation. Deficiency of Piezo1 inhibited integrin-dependent migration, demonstrating crucial role of

these mechanosensors in this mode of T cell migration. As the integrin-dependent chemotactic

migration of T cells is a critically implicated in most patho-physiologic contexts, in the present study

we focused on the potential role of Piezo1 mechanosensors in this specific context.

Notably, a previous study in vivo in mice, using genetic deletion of Piezo1 in T cells, reported no

significant effect on tissue recruitment of T cells (17). Although a functional redundancy in the role of

Piezo1-mediated mechanosensing in mouse T cells warrants exclusion, especially in case of gene

deletion in developing T cells, given the considerable expression of Piezo2 as well in mouse T cells (40).

On the other hand, a role of Piezo1 activation in fibroblastic reticular cells in Peyer's patches has been

shown to regulate lymphocyte migration through tissue conduits (41). The different tissue distribution

of Piezo1 and Piezo2 expression in humans warranted exploring the role of Piezo1 in migrating human

T cells, as in humans Piezo2 is hardly expressed in the hematopoietic cells.

Non-invasive interference reflection microscopy allowed us to demonstrate dynamic change in

membrane tension at the leading edge of T cells stimulated with chemokine on ICAM1-bed which was

closely followed by dynamic recruitment Piezo1 locally. Chemokine receptor activation leads to focal

adhesion formation at the leading edge of the migrating T cells (34, 35). Our experiments revealed

that Piezo1 recruitment at the leading edge of the cells followed focal adhesion kinase activation.

Focal adhesions do increase local membrane tension as they connect the extracellular environment

with the cortical cytoskeleton of the cell (34). Piezo1 recruitment locally seems to be a response to

this membrane tension increase at focal adhesions. Piezo1 recruitment to focal adhesions was also

shown to play a role in enhancing glioblastoma aggression by activating integrin-FAK signalling

upstream of ECM remodelling and tissue stiffening (42). Interestingly, recent study by Yao et. al.,

showed that Piezo1 is required for formation of mature focal adhesions. Piezo1 binds to matrix

adhesion regions in a force-dependent manner via a linker domain and this binding is abrogated in

cancer cells causing transformation to malignant phenotypes (43).

'Outside-in' signaling downstream of focal adhesion formation recruits integrins to the leading edge

of the moving T cells (35, 36) which was abrogated by Piezo1 downregulation. Piezo1 has been shown

to drive Ca^{2+} influx into the T cells leading to calpain activation (14, 19). As expected, calpain inhibition

also interfered with leading edge recruitment of the integrin LFA-1. Plausibly calpain activation drives

local F-actin polymerization at the leading edge facilitating LFA-1 recruitment. This was also supported

by loss of polarity of AKT phosphorylation with Piezo1 deficiency, which is known to be the major

downstream signaling module that succeeds LFA-1 recruitment and activation (39, 44). These

dysregulations in turn abrogates the retrograde flow of actin in the moving cell that is essential for

forward propulsion of the cell body (6, 7, 37, 43). Of note here, a role of ER-resident Piezo1, in driving

R-RAS-driven activation of integrins, which was again dependent on calpain activation, was reported

even before the mechanosensing function of Piezo1 was established (30).

Thus, the present study reports a hitherto unexplored role of the professional mechanosensor Piezo1

in integrin dependent chemotactic movement of human T cells. In the cascade of leading edge events

in a moving human T cell, following chemokine receptor activation and eventually leading to the

retrograde flow of actin, focal adhesion formation precedes Piezo1 activation, while LFA-1 recruitment

is dependent on Piezo1 function.

METHODS

Human CD4$^+$ T cell isolation and transfection.

Human CD4$^+$ T lymphocytes were isolated from peripheral blood from healthy donors through

magnetic immunoselection. Peripheral blood samples from healthy human subjects were collected on

obtaining written informed consents. The studies were approved by the Human Ethics Committee of

CSIR-Indian Institute of Chemical Biology, India. Cells were transfected with either EGFP control or

Piezo1-specific siRNAs using Lonza 4D nucleofector, as per the manufacturer's protocol. Briefly, cells

were nucleofected in 100µl of supplemented P3 primary buffer containing 175ng of respective siRNAs.

Nucleofection was performed using EO-115 pulse in Lonza 4D nucleofector. Nucleofected cells were

incubated at 37°C. 5% CO_2 in RPMI media containing 10% FBS for 3 days before experiments were

conducted. Knockdown was checked by qPCR measurement of Piezo1 and Piezo2 transcript levels

(Supplementary Figure. 1A).

GFP/Piezo1 siRNA co-nucleofection

Approximately 2×10^6 human CD4+ T lymphocytes were nucleofected with 3µg of GFP-expressing

plasmid along with 175ng of Piezo1-specific siRNA. Cells were incubated for 3 days before migration

studies. EGFP esiRNA was used as control siRNA in experiments involving comparison between control

and Piezo1 siRNA-KD cells.

Analysis of Piezo1 knockdown

GFP and Piezo1 siRNA co-transfected CD4[+] T cells were stained for Piezo1 to measure downregulation.

Both surface and intracellular (0.2% triton X-100 permeabilization) Piezo1 were stained. Briefly, cells

were collected on day and fixed in 0.5% PFA for 5 minutes on ice. Cells were stained with 1.67µg/ml

of rabbit anti-Piezo1 for 45 minutes at 4°C. After washing, 1.5µg/ml of anti-rabbit alexa 568-IgG was

added and incubated for 30 minutes at 4°C. Cells were acquired on BD LSR Fortessa III after washing.

Piezo1 mean fluorescence intensity (MFI) were measured both for GFP[+] and GFP[–] cells.

2-dimensional cell tracking.

12-mm confocal dishes were coated with 4µg/ml of recombinant human ICAM-1 overnight at 4°C.

Wells were washed with 1X PBS. Tracking of GFP/Piezo1 siRNA-transfected cells were performed after

cell trace violet-labelling in the presence of recombinant CCL19 on ICAM1-coated dishes. Stained cells

were added to the wells and allowed to settle for 1 hour at 37°C. 2D tracking was performed at 37°C

after addition of 0.5µg/ml of recombinant human CCL19 and incubated for 10 minutes before

acquisition. Images were acquired at regular intervals at room temperature using Zeiss LSM setup at

20X magnification for a total duration of 10-15 minutes at 30 second interval. CD4[+] T lymphocytes

were RPMI without phenol red supplemented with 10% FBS and 25mM HEPES during the duration of

imaging.

2D cell tracking analysis

For tracking analyses of GFP and Piezo1 siRNA co-transfected cells, all cells above a specific threshold

area were used to eliminate potential debris and dead cells due to GFP plasmid transfection. Time-

based image stacks obtained were analysed using the Particle Tracker 2D/3D plugin of the FIJI Mosaic

Suite. Briefly, images were corrected for any shift in alignment during acquisition, using the StackReg

plugin. Corrected images were the thresholded to create a 0-255 binary mask which were then used

for tracking using the Particle Tracker 2D/3D plugin. Tracking provided frame-wise X-Y co-ordinates

of cells which were subsequently used to derive the following parameter to measure motility:

a. Mean-squared displacement

$MSD (n) = \sum_{i=0}^{i=n}(x_n - x_0)^2 + (y_n - y_0)^2$ where,

i = 0 is the first frame,

i = n is the n[th] frame.

Tranwell chemotaxis assay

5μm transwell inserts were coated with 4μg/ml of recombinant human ICAM-1. Equal number of primary human CD4[+] T cells and Jurkat cells nucleofected with control or Piezo1-specific siRNA were added to the ICAM-1-coated inserts. Cells were allowed to migrate in response to recombinant human CCL19 (0.5μg/ml) or SDF1α (0.1μg/ml) in the lower chamber wells for 5 hours and 1.5 hours, respectively, at 37°C, 5% CO_2. Migration was halted by placing the setup on ice. Percent migration was calculated by counting the number of cells in the lower chamber and dividing it by total number of cells that was seeded.

Overexpression of mCherry/GFP-tagged Piezo1 and Actin-GFP in Jurkat cells.

GFP and mCherry-tagged Piezo1 constructs were kindly provided by Dr. Charles Cox at Victor Chang Cardiac Research Laboratory, Darlinghurst, Australia. Approximately 2 million Jurkat cells were nucleofected with 2-3μg of fluorescent Piezo1 construct in 100μl of supplemented SE cell line buffer. Cells were pulsed using CL-120 program in Lonza 4D nucleofector. Cells were cultured in antibiotics-deficient RPMI supplemented with 20% FBS for 36 hours. Overexpression was confirmed using flow cytometry.

Interference Reflection Microscopy and Analysis

Jurkat cells were nucleofected with 2-3μg/ml of Piezo1-GFP construct. Overexpression was confirmed using flow cytometry. Transfected cells were seeded on recombinant human fibronectin glass-based dishes. Cells were allowed to completely adhere for 1 hour to allow IRM imaging.

An inverted microscope (Nikon, Japan) with 60x 1.2 NA water-immersion objective, Hg arc lamp and necessary interference filter (546 ± 12 nm) were used for the interference reflection microscopy (45, 46) and images were captured at 19.91 frames per second using a EMCCD camera (Evolve 512 Delta; Photometrics, Trenton, NJ). Cells that were significantly adhered to the glass bottom were used for imaging. Regions used for analysis were stringently chosen to avoid parts farther than ~ 100 nm from the coverslip – also called the first branch region (FBR) and the size chosen was 4x4 pixels corresponding to ~ (720 nm)^2. First, cells were imaged by IRM to take baseline reading following which chemokine (SDF1α) was added. Cells were followed till 30 min with ~ 2-3 min interval post-chemokine addition.

For stringent selection of subset of regions, extraction of parameters as well as for visualizing the amplitude of fluctuations, the methodology developed previously (46, 47) was followed. 2048 frames corresponding to ~ 102 sec were captured and used for a single measurement of tension. The analysis

348 involved calculation of standard deviation of temporal height fluctuations (SD_{time}) and estimating the

349 power spectral density (PSD) by autoregressive technique (probability of covariance or pcov) where

350 the order is chosen to match best with PSD obtained using fast fourier transform (MATLAB, Mathworks

351 Inc., USA). For deriving mechanical parameters, the PSD is fitted with a Helfrich-based model –

352
$$PSD(f) = \frac{4\eta_{eff}Ak_BT}{\pi}\int_{q_{min}}^{q_{max}}\frac{dq}{(4\eta_{eff}(2\pi f))^2 + \left[\kappa q^3 + \sigma q + \frac{\gamma}{q}\right]^2}$$

with four fitting parameters (46) using fluctuation-tension (σ), effective viscosity (η_{eff}), level of

confinement (γ) and the effective active temperature (A)). The bending rigidity (κ) was fixed at 15 k_BT.

Membrane fluctuation-tension closely follows membrane frame tension for a large range of tension

values (48). Membrane fluctuation-tension closely follows membrane frame tension for a large range

of tension values (48). For brevity, membrane fluctuation-tension has been referred to as "tension" in

the rest of the manuscript. Tension maps were created from PSDs calculated for every pixel which

were subsequently fitted and parameters extracted for every pixel. For analysing correlation between

fluorescence and tension, first the fluorescence distribution of cells was plotted. Regions of

intermediate intensity (around the first peak of the distribution) of a fixed intensity range was filtered

and the pixel count of the filtered area was plotted against the average tension in those pixels.

Normalization was done by diving by the maximum value in the time series. Normalized pixel count,

thus, reflects how area with high Piezo1-GFP at the membrane changes while avoiding a small area

fraction with very high Piezo1-GFP which might arise from internal structures. To ensure significance

of the observed differences, the statistical test undertaken was the Mann-Whitney U test.

Piezo1 and actin distribution in live cell imaging

Jurkat cells expressing mCherry-tagged Piezo1 were seeded on ICAM-1-coated dishes. 0.1μg/ml of

recombinant human SDF1α was added to the cells for 15 minutes prior to imaging. Cell tracking was

performed in EVOS FL microscope using the PE-Texas red filter, at an interval of 1 frame per minute,

at room temperature. Imaging was performed for a total duration of 15 minutes.

For Jurkat cells transfected with Piezo1-mcherry and actin-GFP, cells were treated similarly and

acquired on Zeiss LSM confocal setup. Z-stack was obtained at 1μm per stack to capture the entire

cell. Time-lapse was imaged at an interval of 30 second per frame for a total duration of 5 minutes.

Confocal imaging was performed at 37°C.

For particle image velocimetry (PIV) analysis of Piezo1-mCherry-expressing Jurkat cells was performed

using the Iterative PIV plugin of FIJI. Time-lapse image stacks were split and stacks of paired

subsequent images were formed. Each of these stacks was then corrected for any shift in alignment

using the StackReg plugin. Aligned stacks were then subjected to iterative FIJI PIV analysis using the

conventional cross-correlation method to measure flow of fluorescent Piezo1 in chemokine-treated

Jurkat cells. PIV post-processing was done with default parameters and PIV magnitude plots were

obtained from the plugin.

Measurement of cell polarity during live-cell imaging.

A custom code was written in Fiji (ImageJ) to analyze the cell image time-lapse for the calculation

Front/Back (F/B) ratio of actin and Piezo1 expression relative to the direction of cell migration. Briefly,

intensity-based thresholding was performed on the actin images to create binary masks of the cells.

Individual cell boundaries were identified in the time-lapse, which were stored in the form of polygon

ROIs. The cell ROIs enabled the computation of each cell's trajectory and displacement vectors from

the geometrical centroid at every frame of the time-lapse video. The computation of cell polarity

involved fitting each cell's ROI to an ellipse and bisecting the ellipse perpendicular to the migration

direction. The ratio of total intensities in the leading half and the trailing half of the ellipse was

calculated for actin and piezo1 images to get the F/B ratio.

For polarity measurement of fixed confocal images, cell boundaries were similarly identified. Stored

ROIs were then split into two halves perpendicular to the long axis of the cell. The polarity index was

measured by calculating the difference of total intensities of two halves of the cell divided by the sum

of their total intensities. A measurement of 1 shows complete skewing of fluorescent signals to one

half of the cell, while a value of 0 shows uniform distribution of fluorescent signals in both cell halves

Representative polarity analysis Piezo1 (fixed), Piezo1 mCherry (live), pFAK (fixed), CD11a/LFA-1

(fixed) and actin (fixed) of untreated and CCL19-stimulated CD4[+] T lymphocytes (Supplemental figure

6A-F).

Generation of Piezo1 mCherry and Actin-GFP polar plots.

For analysis of angular/spatial distribution of Piezo1 mCherry and actin-GFP with respect to direction

of cell movement, cell ROIs were applied to the actin and piezo1 channels individually, to find the

intensity-weighted centre of mass (through all the time-points). Depending on the fluorescence

intensity distribution, the intensity-weighted centre of mass is shifted away from the geometric

centroid and towards the direction of a higher protein distribution. Thus, a vector connecting the

centroid (geometric centre) to the centre of mass gives the direction of polarization of the protein of

interest. The relative angle of the polarization vector against the cell displacement vector was

calculated for each frame, and a polar histogram of the angles was plotted for actin-GFP and piezo1-
mCherry.

Fixed immunostaining for confocal imaging

Untransfected human CD4$^+$ T lymphocytes were seeded on 12-mm, ICAM-1-coated glass coverslips at
a density of about 0.2 million cells per coverslip. Cells were incubated at 37°C, 5% CO$_2$ for 1.5 hours.
0.5µg/ml of recombinant human CCL19 was added to each coverslip and incubated for 20-30 minutes
at 37°C. Post-treatment, cells were fixed in 4% paraformaldehyde for 15 minutes at room
temperature. Blocking was performed in 3% bovine serum albumin in 1X PBS containing 0.2% Triton
X-100 for 50 minutes at room temperature. Primary rabbit anti-human Piezo1 (Proteintech) was
added at a dilution of 1:100 (5µg/ml) and incubated at room temperature for 1 to 1.5 hours. Alexa
568-conjugated anti-rabbit IgG was added at a dilution of 1:350 (5.7µg/ml), and incubated for further
45-50 minutes. Nuclear staining was performed for 1 minute with Hoechst 33342 dye at a
concentration of 1µg/ml in 1X PBS. All washes between steps were carried out in 1X PBS. Stained cells
were then mounted on Vectashield® mounting media and sealed before confocal imaging. For Piezo1
and CD44 co-staining, rabbit anti-Piezo1 and mouse anti-human CD44 were used at a concentration
of 5µg/ml, each. Alexa 488-conjugated anti-rabbit IgG and alexa 633-conjugated anti-mouse IgG were
used at a concentration of 5.7µg/ml, each.

For actin imaging, LifeAct-TagGFP2 protein was added at a concentration of 0.5µg/ml, and incubated
for 45 minutes at room temperature before washing. For LFA-1 ($\alpha_L\beta2$) staining, mouse anti-human
CD11a (α_L subunit) was added at a concentration of 5µg/ml, followed by addition of anti-mouse Alexa
594-conjugated IgG. For LFA-1 and Piezo1 co-staining, primary antibodies were added at above
concentrations, with Alexa 568-conjugated anti-rabbit IgG (for Piezo1) and Alexa 488-conjuagted anti-
mouse IgG (for CD11a/LFA-1). Rabbit/mouse anti-phosphorylated FAK (Tyr-397) and mouse Paxillin
was used to stain pFAK and paxillin, respectively to detect focal adhesions. These cells were treated
with chemokine CCL19 for 10 minutes prior to fixation. While doing confocal microscopy, Z-stacks
were captured at 0.5µm intervals. Confocal images are represented are maximum intensity
projections of the Z-slices. 3Dprojectionfunction of Fiji was used to create a 3D rendering of the Z-
slices, using the brightest point method of projection.

TIRF Imaging and Ca^{2+}-Piezo1 colocalization

Ca^{2+}-Piezo1 imaging involved incubating cells with Fluo-3 AM (2.5 µM) in HBSS buffer for 30 minutes
at 37˚C, followed by washing and imaging in RPMI containing 10% FBS, and 25 mM HEPES media.
Jurkat cell were transfected with Piezo1 mCherry plasmid and only those expressing Piezo1 mCherry

were selected for the study. Dual-colour sequential TIRF imaging was employed, with Piezo1 mCherry

being imaged at 300 ms (100 frames) and Ca2+ imaging performed using a 10 ms exposure time (2000

frames). The short exposure for Ca2+ was used to capture transient local peaks of fluorescence which

would otherwise average out at longer exposure times. The maximum projection of the time series

was used to detect local peaks using adaptive local thresholding (MATLAB), and a size filter was used

to exclude large objects identified with a relatively low Ca^{2+} gradient. For each specific set of settings,

the Ca^{2+} signal objects detected were used to calculate the average cluster size and total area of

objects. Uniform random numbers were used to simulate clusters with appropriate probability to

ensure that the final area of the simulated Ca^{2+} clusters closely matched the real data. The overlap

between the real Ca^{2+} signal and Piezo1 objects was compared with the simulated data to test for

deviation from a random placement of Ca^{2+} spots. To assess the robustness of the results, different

combinations of size filters of Piezo1 and Ca^{2+} objects were used, ranging from 50 to 350 pixels.

Calpain Inhibition

PD150606 (Merck, India), a non-competitive calpain 1 and 2 inhibitors, was used at a concentration of

100μM. Untransfected CD4+ T lymphocytes were treated with PD150606 for 1 hour at 37°C prior to

treatment of chemokine. Post -chemokine treatment cells were stained either with pFAK as described.

Focal adhesion kinase Inhibition

FAK inhibitor 14 (1,2,4,5-Benzenetetraamine tetrahydrochloride) was used to selectively inhibit FAK

activity. Inhibitor was added at a final concentration 10μM, 1 hour prior to treatment with

recombinant CCL19. Post-treatment with CCL19, CD4+ T lymphocytes were stained with Piezo1 for

confocal imaging.

Piezo1 Inhibition

Primary human CD4+ T cells were incubated with 10μM of GsMTx4 for 1 hour prior to transwell

chemotaxis assay in response to recombinant CCL19.

Statistics

Student t-test was used to calculate significance.

ACKNOWLEDGEMENTS

The study was funded by Council of Scientific and Industrial Research, India (Grant no. FBR MLP-140).

D.G. is a recipient of Swarnajayanti Fellowship from Department of Science and Technology,

Government of India. The authors express their gratitude for provision of the Piezo1 constructs by Dr.

Charles Cox, Victor Chang Institute of Cardiac Research, Australia.

The authors declare no competing interests

C.S.C.L. designed, performed and analyzed most of the experiments, prepared the figures and took

part in writing the manuscript; T.M., B.S. and C.S.C.L. performed the IRM experiments; T.M. and B.S.

analyzed IRM data; C.S.C.L., P.B. and D.K.S. performed image analyses; Pu.B., M.A.H., B.P.S., J.S. and

R.D. helped with cell culture and confocal microscopy experiments; D.G. conceptualized and

supervised the study, acquired funding, designed experiments and wrote the manuscript with input

from all authors.

REFERENCES

1. Shechter R, London A, Schwartz M. Orchestrated leukocyte recruitment to immune-privileged

sites: absolute barriers versus educational gates. Nat Rev Immunol. 2013 Mar;13(3):206-18.

doi: 10.1038/nri3391.

2. Krummel MF, Bartumeus F, Gérard A. T cell migration, search strategies and mechanisms. Nat

Rev Immunol. 2016 Mar;16(3):193-201. doi: 10.1038/nri.2015.16.

3. Woodland DL, Kohlmeier JE. Migration, maintenance and recall of memory T cells in peripheral

tissues. Nat Rev Immunol. 2009 Mar;9(3):153-61. doi: 10.1038/nri2496.

4. Masopust D, Schenkel JM. The integration of T cell migration, differentiation and function. Nat

Rev Immunol. 2013 May;13(5):309-20. doi: 10.1038/nri3442.

5. Fowell DJ, Kim M. The spatio-temporal control of effector T cell migration. Nat Rev Immunol.

2021 Sep;21(9):582-596. doi: 10.1038/s41577-021-00507-0.

6. Dupré L, Houmadi R, Tang C, Rey-Barroso J. T Lymphocyte Migration: An Action Movie Starring

the Actin and Associated Actors. Front Immunol. 2015 Nov 18;6:586. doi:

10.3389/fimmu.2015.00586.

7. Nordenfelt P, Moore TI, Mehta SB, Kalappurakkal JM, Swaminathan V, Koga N, Lambert TJ,

Baker D, Waters JC, Oldenbourg R, Tani T, Mayor S, Waterman CM, Springer TA. Direction of

actin flow dictates integrin LFA-1 orientation during leukocyte migration. Nat Commun. 2017

Dec 11;8(1):2047. doi: 10.1038/s41467-017-01848-y.

8.   Moreau HD, Piel M, Voituriez R, Lennon-Duménil AM. Integrating Physical and Molecular Insights on Immune Cell Migration. Trends Immunol. 2018 Aug;39(8):632-643. doi: 10.1016/j.it.2018.04.007.

9.   Shannon MJ, Pineau J, Griffié J, Aaron J, Peel T, Williamson DJ, Zamoyska R, Cope AP, Cornish GH, Owen DM. Differential nanoscale organisation of LFA-1 modulates T-cell migration. J Cell Sci. 2019 Oct 16;133(5):jcs232991. doi: 10.1242/jcs.232991.

10. Coste B, Mathur J, Schmidt M, Earley TJ, Ranade S, Petrus MJ, Dubin AE, Patapoutian A. Piezo1 and Piezo2 are essential components of distinct mechanically activated cation channels. Science. 2010 Oct 1;330(6000):55-60. doi: 10.1126/science.1193270.

11. Coste B, Xiao B, Santos JS, Syeda R, Grandl J, Spencer KS, Kim SE, Schmidt M, Mathur J, Dubin AE, Montal M, Patapoutian A. Piezo proteins are pore-forming subunits of mechanically activated channels. Nature. 2012 Feb 19;483(7388):176-81. doi: 10.1038/nature10812.

12. Syeda R, Florendo MN, Cox CD, Kefauver JM, Santos JS, Martinac B, Patapoutian A. Piezo1 Channels Are Inherently Mechanosensitive. Cell Rep. 2016 Nov 8;17(7):1739-1746. doi: 10.1016/j.celrep.2016.10.033.

13. Murthy SE, Dubin AE, Patapoutian A. Piezos thrive under pressure: mechanically activated ion channels in health and disease. Nat Rev Mol Cell Biol. 2017 Dec;18(12):771-783. doi: 10.1038/nrm.2017.92. Epub 2017 Oct 4. PMID: 28974772.

14. Liu CSC, Raychaudhuri D, Paul B, Chakrabarty Y, Ghosh AR, Rahaman O, Talukdar A, Ganguly D. Cutting Edge: Piezo1 Mechanosensors Optimize Human T Cell Activation. J Immunol. 2018 Feb 15;200(4):1255-1260. doi: 10.4049/jimmunol.1701118.

15. Atcha H, Jairaman A, Holt JR, Meli VS, Nagalla RR, Veerasubramanian PK, Brumm KT, Lim HE, Othy S, Cahalan MD, Pathak MM, Liu WF. Mechanically activated ion channel Piezo1 modulates macrophage polarization and stiffness sensing. Nat Commun. 2021 May 31;12(1):3256. doi: 10.1038/s41467-021-23482-5.

16. Geng J, Shi Y, Zhang J, Yang B, Wang P, Yuan W, Zhao H, Li J, Qin F, Hong L, Xie C, Deng X, Sun Y, Wu C, Chen L, Zhou D. TLR4 signalling via Piezo1 engages and enhances the macrophage mediated host response during bacterial infection. Nat Commun. 2021 Jun 10;12(1):3519. doi: 10.1038/s41467-021-23683-y.

17. Jairaman A, Othy S, Dynes JL, Yeromin AV, Zavala A, Greenberg ML, Nourse JL, Holt JR, Cahalan SM, Marangoni F, Parker I, Pathak MM, Cahalan MD. Piezo1 channels restrain regulatory T cells but are dispensable for effector CD4+ T cell responses. Sci Adv. 2021 Jul 7;7(28):eabg5859. doi: 10.1126/sciadv.abg5859.

18. Jäntti H, Sitnikova V, Ishchenko Y, Shakirzyanova A, Giudice L, Ugidos IF, Gómez-Budia M,

Korvenlaita N, Ohtonen S, Belaya I, Fazaludeen F, Mikhailov N, Gotkiewicz M, Ketola K,

Lehtonen Š, Koistinaho J, Kanninen KM, Hernández D, Pébay A, Giugno R, Korhonen P,

Giniatullin R, Malm T. Microglial amyloid beta clearance is driven by PIEZO1 channels. J

Neuroinflammation. 2022 Jun 15;19(1):147. doi: 10.1186/s12974-022-02486-y.

19. Liu CSC, Ganguly D. Mechanical Cues for T Cell Activation: Role of Piezo1 Mechanosensors.

Crit Rev Immunol. 2019;39(1):15-38. doi: 10.1615/CritRevImmunol.2019029595.

20. Ma S, Dubin AE, Zhang Y, Mousavi SAR, Wang Y, Coombs AM, Loud M, Andolfo I, Patapoutian

A. A role of PIEZO1 in iron metabolism in mice and humans. Cell. 2021 Feb 18;184(4):969-

982.e13. doi: 10.1016/j.cell.2021.01.024.

21. He Y, Deng B, Liu S, Luo S, Ning Y, Pan X, Wan R, Chen Y, Zhang Z, Jiang J, Xu H, Xia M, Li J.

Myeloid Piezo1 Deletion Protects Renal Fibrosis by Restraining Macrophage Infiltration and

Activation. Hypertension. 2022 May;79(5):918-931. doi:

10.1161/HYPERTENSIONAHA.121.18750.

22. Huang Z, Sun Z, Zhang X, Niu K, Wang Y, Zheng J, Li H, Liu Y. Loss of stretch-activated channels,

PIEZOs, accelerates non-small cell lung cancer progression and cell migration. Biosci Rep. 2019

Mar 22;39(3): BSR20181679. doi: 10.1042/BSR20181679.

23. Chubinskiy-Nadezhdin VI, Vasileva VY, Vassilieva IO, Sudarikova AV, Morachevskaya EA,

Negulyaev YA. Agonist-induced Piezo1 activation suppresses migration of transformed

fibroblasts. Biochem Biophys Res Commun. 2019 Jun 18;514(1):173-179. doi:

10.1016/j.bbrc.2019.04.139.

24. Mousawi F, Peng H, Li J, Ponnambalam S, Roger S, Zhao H, Yang X, Jiang LH. Chemical

activation of the Piezo1 channel drives mesenchymal stem cell migration via inducing ATP

release and activation of P2 receptor purinergic signaling. Stem Cells. 2020 Mar;38(3):410-

421. doi: 10.1002/stem.3114.

25. Wang X, Cheng G, Miao Y, Qiu F, Bai L, Gao Z, Huang Y, Dong L, Niu X, Wang X, Li Y, Tang H, Xu

Y, Song X. Piezo type mechanosensitive ion channel component 1 facilitates gastric cancer

omentum metastasis. J Cell Mol Med. 2021 Feb;25(4):2238-2253. doi: 10.1111/jcmm.16217.

26. Gao L, Ji Y, Wang L, He M, Yang X, Qiu Y, Sun X, Ji Z, Yang G, Zhang J, Li S, Dai L, Zhang L.

Suppression of Esophageal Squamous Cell Carcinoma Development by Mechanosensitive

Protein Piezo1 Downregulation. ACS Omega. 2021 Apr 12;6(15):10196-10206. doi:

10.1021/acsomega.1c00505.

565 27. Holt JR, Zeng WZ, Evans EL, Woo SH, Ma S, Abuwarda H, Loud M, Patapoutian A, Pathak MM.
566   Spatiotemporal dynamics of PIEZO1 localization controls keratinocyte migration during
567   wound healing. Elife. 2021 Sep 27;10:e65415. doi: 10.7554/eLife.65415.

568 28. Velasco-Estevez M, Koch N, Klejbor I, Caratis F, Rutkowska A. Mechanoreceptor Piezo1 Is
569   Downregulated in Multiple Sclerosis Brain and Is Involved in the Maturation and Migration of
570   Oligodendrocytes in vitro. Front Cell Neurosci. 2022 May 26;16:914985. doi:
571   10.3389/fncel.2022.914985.

572 29. Hung WC, Yang JR, Yankaskas CL, Wong BS, Wu PH, Pardo-Pastor C, Serra SA, Chiang MJ, Gu
573   Z, Wirtz D, Valverde MA, Yang JT, Zhang J, Konstantopoulos K. Confinement Sensing and Signal
574   Optimization via Piezo1/PKA and Myosin II Pathways. Cell Rep. 2016 May 17;15(7):1430-1441.
575   doi: 10.1016/j.celrep.2016.04.035.

576 30. McHugh BJ, Buttery R, Lad Y, Banks S, Haslett C, Sethi T. Integrin activation by Fam38A uses a
577   novel mechanism of R-Ras targeting to the endoplasmic reticulum. J Cell Sci. 2010 Jan 1;123(Pt
578   1):51-61. doi: 10.1242/jcs.056424.

579 31. Xiao B. 2020. Levering Mechanically Activated Piezo Channels for Potential Pharmacological
580   Intervention. Annu Rev PharmacolToxicol. 60:195-218. doi: 10.1146/annurev-pharmtox-
581   010919-023703.

582 32. Barr VA, Bunnell SC. Interference reflection microscopy. Curr Protoc Cell Biol. 2009 Dec;
583   Chapter 4: Unit 4.23. doi: 10.1002/0471143030.cb0423s45.

584 33. Chakraborty M, Mukherjee B, Nalinakshan N, Biswas A, Nayak RK, Sinha B. Effect of
585   heterogeneous substrate adhesivity of follower cells on speed and tension profile of leader
586   cells in primary keratocyte collective cell migration. Biol Open. 2022 Mar 15;11(3):bio058893.
587   doi: 10.1242/bio.058893.

588 34. Mitra SK, Hanson DA, Schlaepfer DD. Focal adhesion kinase: in command and control of cell
589   motility. Nat Rev Mol Cell Biol. 2005 Jan;6(1):56-68. doi: 10.1038/nrm1549.

590 35. Katoh K. FAK-Dependent Cell Motility and Cell Elongation. Cells. 2020 Jan 12;9(1):192. doi:
591   10.3390/cells9010192.

592 36. Yamashiro S, Watanabe N. A new link between the retrograde actin flow and focal adhesions.
593   J Biochem. 2014 Nov;156(5):239-48. doi: 10.1093/jb/mvu053.

594 37. Martino F, Perestrelo AR, Vinarský V, Pagliari S, Forte G. Cellular Mechanotransduction: From
595   Tension to Function. Front Physiol. 2018 Jul 5;9:824. doi: 10.3389/fphys.2018.00824.

596 38. Sánchez-Martín L, Sánchez-Sánchez N, Gutiérrez-López MD, Rojo AI, Vicente-Manzanares M,
597   Pérez-Alvarez MJ, Sánchez-Mateos P, Bustelo XR, Cuadrado A, Sánchez-Madrid F, Rodríguez-
598   Fernández JL, Cabañas C. Signaling through the leukocyte integrin LFA-1 in T cells induces a

transient activation of Rac-1 that is regulated by Vav and PI3K/Akt-1. J Biol Chem. 2004 Apr
16;279(16):16194-205. doi: 10.1074/jbc.M400905200.

39. Roy NH, Kim SHJ, Buffone A Jr, Blumenthal D, Huang B, Agarwal S, Schwartzberg PL, Hammer
DA, Burkhardt JK. LFA-1 signals to promote actin polymerization and upstream migration in T
cells. J Cell Sci. 2020 Sep 9;133(17):jcs248328. doi: 10.1242/jcs.248328.

40. Ganguly D. Can Piezo2 substitute Piezo1 function in the context of T cell-specific deletion in
mice? Sci Adv, e-Letter published Sep 8, 2021, in response to Jairaman A et al.,

41. Chang JE, Buechler MB, Gressier E, Turley SJ, Carroll MC. Mechanosensing by Peyer's patch
stroma regulates lymphocyte migration and mucosal antibody responses. Nat Immunol. 2019
Nov;20(11):1506-1516. doi: 10.1038/s41590-019-0505-z.

42. Chen X, Wanggou S, Bodalia A, Zhu M, Dong W, Fan JJ, Yin WC, Min HK, Hu M, Draghici D, Dou
W, Li F, Coutinho FJ, Whetstone H, Kushida MM, Dirks PB, Song Y, Hui CC, Sun Y, Wang LY, Li
X, Huang X. A Feedforward Mechanism Mediated by Mechanosensitive Ion Channel PIEZO1
and Tissue Mechanics Promotes Glioma Aggression. Neuron. 2018 Nov 21;100(4):799-815.e7.

43. Yao M, Tijore A, Cheng D, Li JV, Hariharan A, Martinac B, Tran Van Nhieu G, Cox CD, Sheetz M.
Force- and cell state-dependent recruitment of Piezo1 drives focal adhesion dynamics and
calcium entry. Sci Adv. 2022 Nov 11;8(45):eabo1461.

44. Roy NH, MacKay JL, Robertson TF, Hammer DA, Burkhardt JK. Crk adaptor proteins mediate
actin-dependent T cell migration and mechanosensing induced by the integrin LFA-1. Sci
Signal. 2018 Dec 11;11(560):eaat3178. doi: 10.1126/scisignal.aat3178.

45. Limozin L, Sengupta K. Quantitative reflection interference contrast microscopy (RICM) in soft
matter and cell adhesion. Chemphyschem. 2009 Nov 9;10(16):2752-68.

46. Biswas A, Alex A, Sinha B. Mapping Cell Membrane Fluctuations Reveals Their Active
Regulation and Transient Heterogeneities. Biophys J. 2017 Oct 17;113(8):1768-1781. doi:
10.1016/j.bpj.2017.08.041.

47. Biswas A, Kashyap P, Datta S, Sengupta T, Sinha B. Cholesterol Depletion by MβCD Enhances
Cell Membrane Tension and Its Variations-Reducing Integrity. Biophys J. 2019 Apr
23;116(8):1456-1468.

48. Shiba H, Noguchi H, Fournier JB. Monte Carlo study of the frame, fluctuation and internal
tensions of fluctuating membranes with fixed area. Soft Matter. 2016 Feb 28;12(8):2373-80.
doi: 10.1039/c5sm01900a.

Figure 1. Piezo1 deficiency abrogates integrin-dependent motility in human T cells which is mediated through redistribution of Piezo1 at the leading edge in response to chemokine stimulation.

A. MSD versus time calculated for GFP⁺ (potential Piezo1-knockdown cells) and GFP⁻ (potential control cells) in GFP plasmid and Piezo1 siRNA co-transfected human CD4⁺ T lymphocytes, that were allowed to migrate in presence of recombinant CCL19 on ICAM1-coated dishes. **B.** Representative tracks of GFP⁺ and GFP⁻ CD4⁺ T lymphocytes. Comparisons of **C.** % GFP⁺ cells after 72 hours of nucleofection **D**

& E. 3D transwell migration assay of siRNA-transfected primary human CD4[+] T lymphocytes (D) and

Jurkat T cells (E), respectively. Data representative of at least 3 independent experiments. **F.**

Representative confocal images of Piezo1 distribution in fixed untreated and CCL19-treated CD4[+] T

lymphocytes. 63X oil magnification. **G.** Comparison between Piezo1 polarity index calculated for fixed,

stained, human CD4[+] T lymphocytes with or without 0.5μg/ml CCL19 treatment for 20-30 minutes. n

> 510 random cells, each. **H.** Piezo1 polarity index calculated for Jurkat cells, expressing mCherry-

tagged Piezo1 during live-cell tracking in the presence of recombinant SDF1α. Each dot represents

polarity index of each cell, averaged over all the time-frames. **I.** Representative time kinetics of particle

image velocimetry (PIV) analysis of Piezo1-mcherry transfected Jurkat cells, allowed to move on ICAM-

coated dishes in the presence of recombinant SDF1α. Top panel: No chemokine. Bottom Panel: SDF1α.

Figure 2

Figure 2. Piezo1 redistribution in migrating T cells follow increased membrane tension in the leading edge

A. Schematic illustration of interference reflection microscopy imaging setup. **B.** Probability distribution of amplitude of temporal fluctuations (SD_{time}, left) and tension (right) before and after addition of chemokines. **C.** Temporal trajectory of tension of cells before and after addition of chemokine with ** indicating significant difference from control. **D.** Representative IRM images of Jurkat cells (top), corresponding tension maps (middle) and R^2 maps (bottom), before and after chemokine treatment. **E** Correlation between normalized Piezo intensity, tension magnitudes with time. **F.** Scatter plot to show correlation between normalized pixel count of Piezo1 and normalized tension. Piezo1 intensity at specific time-points were correlated with tension magnitudes at the preceding timepoints. Normalization has been done such that the maximum value is set to 1 and others are accordingly scaled. **G.** Colour-coded Piezo1-GFP intensity map (left), epifluorescence image (middle), and tension map (right) of representative Piezo1-GFP expressing Jurkat cell, after 30 minutes of 0.1µg/ml of SDF1α treatment.

Figure 3

Figure 3. Focal adhesion following chemokine receptor activation does not depend on Piezo1

A. Representative confocal images of human CD4+ T lymphocytes, fixed and stained for phospho-FAK

(pFAK) under untreated and CCL19-treated conditions. **B.** Increased polarity of pFAK upon chemokine

stimulation of CD4+ T lymphocytes as compared to unstimulated controls. N > 120 cells, each. **C.**

Representative confocal images of pFAK distribution in control and Piezo1 siRNA-transfected CD4+ T

lymphocytes stimulated with chemokine. **D**. Comparison of pFAK polarity in CCL19-stimulated control

and Piezo1-knockdown CD4+ T lymphocytes. n > 450 random cells, each. **E.** Representative confocal

images of paxillin distribution in untreated versus CCL19-stimulated CD4+ T lymphocytes. **F.** Increased

paxillin polarity in response to CCL19 stimulation as compared to untreated CD4+ T lymphocytes.

Untreated n > 70, CCL19-treated n > 150. **G.** Representative confocal images of immunostained paxillin

in chemokine-stimulated, control and Piezo1-knockdown CD4+ T lymphocytes. **H**. Comparison of

paxillin polarity in CCL19-stimulated control and Piezo1-knockdown CD4+ T lymphocytes. n > 550

random cells, each. **I.** Representative confocal images of stained Piezo1 in CD4+ T cells stimulated with

CCL19 in the presence or absence of FAK inhibitor 14. **J.** Effect of FAK inhibition on Piezo1 polarity in

CD4[+] T lymphocytes stimulated with recombinant CCL19 versus untreated cells. n > 600 random cells,

each. Data represented from at least 3 independent experiments.

Figure 4

Figure 4. Membrane recruitment of LFA1 on chemokine receptor activation disrupts with Piezo1 deficiency.

A. Representative confocal images of fixed, immunostained Piezo1 and LFA1 in unstimulated and CCL19-stimulated CD4[+] T lymphocytes. **B.** Increased LFA1 polarity in response to recombinant CCL19 stimulation in CD4[+] T cells. n > 295 random cells, each. **C.** Manders' co-localisation analysis of Piezo1 and LFA1 in untreated and CCL19-treated CD4[+] T lymphocytes. **D.** Comparison of LFA1 polarity in chemokine-treated control and Piezo1 knockdown CD4[+] T lymphocytes. **E.** Representative confocal stained images of LFA1 polarity of CCL19-treated control and Piezo1-knockdown CD4[+] T lymphocytes. n > 490 random cells, each. **F.** Representative confocal images of phospho-Akt (pAkt) and Akt

721   distribution upon chemokine treatment of control and Piezo1-knockdown cells. **G & H.** Quantitative

722   analyses of pAkt (M) and Akt (N) polarity in control and Piezo1-knockdown cells after CCL19-

723   treatment. n > 580 random cells, each. Data represented from at least 3 independent experiments.

Figure 5

Figure 5. Local Ca²⁺ mobilization with Piezo1 redistribution upon chemokine stimulation.

A. Representative TIRF image of a Jurkat Cell stained with Fluo3AM and transiently expressing with

Piezo1 mCherry. Lower panel: zoomed in image and overlap of binary images of objects detected from

Calcium and Piezo channels as well as simulated randomly placed punctas having similar total (in the

zoomed in section) and average area as that of the Calcium punctas detected in the real images.

Overlap RGB image shows the overlap between Real Calcium (Red) and piezo puncta (Green), overlap

puncta (Yellow), simulated calcium puncta (blue). **B.** Comparison between percentage calcium

colocalization of simulated data and real data. * Denote p value < 0.05 calculated using Mann Whitney

U test. N cell= 52 (13 cells x 4 time points). NROIS= 52. **C.** Representative confocal images of LFA1 and

actin distribution in CCL19-stimulated CD4+ T lymphocytes with or without inhibition of calpain.

100µM of PD15606 was added to the cells 1 hour prior to addition of chemokine. Quantitative

comparisons of **D.** LFA1 and **E.** actin polarity upon chemokine stimulation, in the presence or absence

of calpain pre-inhibition. N > 290 random cells, each. Data represented is of at least 3 independent

experiments.

Figure 6

Figure 6. Piezo1 deficiency disrupts F-actin retrograde flow in T cells despite chemokine receptor

activation.

A. Representative fixed confocal images of actin and Piezo1 distribution in CD4$^+$ T lymphocytes after

30 mins of 0.5µg/ml recombinant CCL19 treatment. **B.** Quantitative comparison of actin polarity in

control and Piezo1-knockdown CD4$^+$ T lymphocytes cells after 30 mins of chemokine treatment. n >

300 random cells, each. **C.** Representative fixed confocal images of actin distribution in chemokine-

treated, control and Piezo1-knockdown CD4$^+$ T lymphocytes. Front-back (F/B) of Piezo1 and actin-GFP

in SDF1α treated **(D)** and untreated **(E)** Jurkat cells co-expressing actin-GFP and Piezo1-mCherry. **F.** A

snapshot of time-lapse imaging of Piezo1-mCherry and actin-GFP expressing Jurkat cell treated with

0.1µg/ml of SDF1α. Image is maximum Z-projection of the cell at 63X/1.40 magnification. Data

represented is of at least 3 independent experiments.

Figure 7

The mechanistic model depicting involvement of Piezo1 mechanosensing in leading edge

events in a migrating T cell.

Proposed model suggests chemokine receptor activation in human T cells lead to focal adhesion kinase

activation and focal adhesion formation. Focal adhesions lead to localized increase in membrane

tension at the leading edge plasma membrane which leads to Piezo1 recruitment and activation.

Piezo1 activation leads to calpain activation which potentially drives further cytoskeletal consolidation

to recruit integrin LFA1. LFA1 recruitment and activation lead to phosphorylation of AKT and

downstream signaling eventually driving the retrograde actin flow in migrating human T cells.

 FIGURE LEGENDS

Figure 1. Piezo1 deficiency abrogates integrin-dependent motility in human T cells which is
mediated through redistribution of Piezo1 at the leading edge in response to chemokine
stimulation.

A. MSD versus time calculated for GFP$^+$ (potential Piezo1-knockdown cells) and GFP$^-$ (potential control
cells) in GFP plasmid and Piezo1 siRNA co-transfected human CD4$^+$ T lymphocytes, that were allowed
to migrate in presence of recombinant CCL19 on ICAM1-coated dishes. **B.** Representative tracks of
GFP$^+$ and GFP$^-$ CD4$^+$ T lymphocytes. Comparisons of **C.** % GFP$^+$ cells after 72 hours of nucleofection **D**
& E. 3D transwell migration assay of siRNA-transfected primary human CD4$^+$ T lymphocytes (D) and
Jurkat T cells (E), respectively. Data representative of at least 3 independent experiments. **F.**
Representative confocal images of Piezo1 distribution in fixed untreated and CCL19-treated CD4$^+$ T
lymphocytes. 63X oil magnification. **G.** Comparison between Piezo1 polarity index calculated for fixed,
stained, human CD4$^+$ T lymphocytes with or without 0.5µg/ml CCL19 treatment for 20-30 minutes. n
> 510 random cells, each. **H.** Piezo1 polarity index calculated for Jurkat cells, expressing mCherry-
tagged Piezo1 during live-cell tracking in the presence of recombinant SDF1α. Each dot represents
polarity index of each cell, averaged over all the time-frames. **I.** Representative time kinetics of particle
image velocimetry (PIV) analysis of Piezo1-mcherry transfected Jurkat cells, allowed to move on ICAM-
coated dishes in the presence of recombinant SDF1α. Top panel: No chemokine. Bottom Panel: SDF1α.

Figure 2. Piezo1 redistribution in migrating T cells follow increased membrane tension in the leading
edge

A. Schematic illustration of interference reflection microscopy imaging setup. **B.** Probability
distribution of amplitude of temporal fluctuations (SD$_{time}$, left) and tension (right) before and after
addition of chemokines. **C.** Temporal trajectory of tension of cells before and after addition of
chemokine with ** indicating significant difference from control. **D.** Representative IRM images of
Jurkat cells (top), corresponding tension maps (middle) and R^2 maps (bottom), before and after
chemokine treatment. **E** Correlation between normalized Piezo intensity, tension magnitudes with
time. **F.** Scatter plot to show correlation between normalized pixel count of Piezo1 and normalized
tension. Piezo1 intensity at specific time-points were correlated with tension magnitudes at the
preceding timepoints. Normalization has been done such that the maximum value is set to 1 and
others are accordingly scaled. **G.** Colour-coded Piezo1-GFP intensity map (left), epifluorescence image

(middle), and tension map (right) of representative Piezo1-GFP expressing Jurkat cell, after 30 minutes

of 0.1μg/ml of SDF1α treatment.

Figure 3. Focal adhesion following chemokine receptor activation does not depend on Piezo1

A. Representative confocal images of human CD4$^+$ T lymphocytes, fixed and stained for phospho-FAK

(pFAK) under untreated and CCL19-treated conditions. **B.** Increased polarity of pFAK upon chemokine

stimulation of CD4$^+$ T lymphocytes as compared to unstimulated controls. N > 120 cells, each. **C.**

Representative confocal images of pFAK distribution in control and Piezo1 siRNA-transfected CD4$^+$ T

lymphocytes stimulated with chemokine. **D**. Comparison of pFAK polarity in CCL19-stimulated control

and Piezo1-knockdown CD4$^+$ T lymphocytes. n > 450 random cells, each. **E.** Representative confocal

images of paxillin distribution in untreated versus CCL19-stimulated CD4$^+$ T lymphocytes. **F.** Increased

paxillin polarity in response to CCL19 stimulation as compared to untreated CD4$^+$ T lymphocytes.

Untreated n > 70, CCL19-treated n > 150. **G.** Representative confocal images of immunostained paxillin

in chemokine-stimulated, control and Piezo1-knockdown CD4$^+$ T lymphocytes. **H**. Comparison of

paxillin polarity in CCL19-stimulated control and Piezo1-knockdown CD4$^+$ T lymphocytes. n > 550

random cells, each. **I.** Representative confocal images of stained Piezo1 in CD4$^+$ T cells stimulated with

CCL19 in the presence or absence of FAK inhibitor 14. **J.** Effect of FAK inhibition on Piezo1 polarity in

CD4$^+$ T lymphocytes stimulated with recombinant CCL19 versus untreated cells. n > 600 random cells,

each. Data represented from at least 3 independent experiments.

Figure 4. Membrane recruitment of LFA1 on chemokine receptor activation disrupts with Piezo1
deficiency.

A. Representative confocal images of fixed, immunostained Piezo1 and LFA1 in unstimulated and

CCL19-stimulated CD4$^+$ T lymphocytes. **B.** Increased LFA1 polarity in response to recombinant CCL19

stimulation in CD4$^+$ T cells. n > 295 random cells, each. **C.** Manders' co-localisation analysis of Piezo1

and LFA1 in untreated and CCL19-treated CD4$^+$ T lymphocytes. **D.** Comparison of LFA1 polarity in

chemokine-treated control and Piezo1 knockdown CD4$^+$ T lymphocytes. **E.** Representative confocal

stained images of LFA1 polarity of CCL19-treated control and Piezo1-knockdown CD4$^+$ T lymphocytes.

n > 490 random cells, each. **F.** Representative confocal images of phospho-Akt (pAkt) and Akt

distribution upon chemokine treatment of control and Piezo1-knockdown cells. **G & H.** Quantitative

analyses of pAkt (M) and Akt (N) polarity in control and Piezo1-knockdown cells after CCL19-

treatment. n > 580 random cells, each. Data represented from at least 3 independent experiments.

Figure 5. Local Ca²⁺ mobilization with Piezo1 redistribution upon chemokine stimulation.

A. Representative TIRF image of a Jurkat Cell stained with Fluo3AM and transiently expressing with Piezo1 mCherry. Lower panel: zoomed in image and overlap of binary images of objects detected from Calcium and Piezo channels as well as simulated randomly placed punctas having similar total (in the zoomed in section) and average area as that of the Calcium punctas detected in the real images. Overlap RGB image shows the overlap between Real Calcium (Red) and piezo puncta (Green), overlap puncta (Yellow), simulated calcium puncta (blue). **B.** Comparison between percentage calcium colocalization of simulated data and real data. * Denote p value < 0.05 calculated using Mann Whitney U test. N cell= 52 (13 cells x 4 time points). NROIS= 52. **C.** Representative confocal images of LFA1 and actin distribution in CCL19-stimulated CD4+ T lymphocytes with or without inhibition of calpain. 100μM of PD15606 was added to the cells 1 hour prior to addition of chemokine. Quantitative comparisons of **D.** LFA1 and **E.** actin polarity upon chemokine stimulation, in the presence or absence of calpain pre-inhibition. N > 290 random cells, each. Data represented is of at least 3 independent experiments.

Figure 6. Piezo1 deficiency disrupts F-actin retrograde flow in T cells despite chemokine receptor activation.

A. Representative fixed confocal images of actin and Piezo1 distribution in CD4⁺ T lymphocytes after 30 mins of 0.5μg/ml recombinant CCL19 treatment. **B.** Quantitative comparison of actin polarity in control and Piezo1-knockdown CD4⁺ T lymphocytes cells after 30 mins of chemokine treatment. n > 300 random cells, each. **C.** Representative fixed confocal images of actin distribution in chemokine-treated, control and Piezo1-knockdown CD4⁺ T lymphocytes. Front-back (F/B) of Piezo1 and actin-GFP in SDF1α treated **(D)** and untreated **(E)** Jurkat cells co-expressing actin-GFP and Piezo1-mCherry. **F.** A snapshot of time-lapse imaging of Piezo1-mCherry and actin-GFP expressing Jurkat cell treated with 0.1μg/ml of SDF1α. Image is maximum Z-projection of the cell at 63X/1.40 magnification. Data represented is of at least 3 independent experiments.

Figure 7. The mechanistic model depicting involvement of Piezo1 mechanosensing in leading edge events in a migrating T cell.

Proposed model suggests chemokine receptor activation in human T cells lead to focal adhesion kinase
activation and focal adhesion formation. Focal adhesions lead to localized increase in membrane
tension at the leading edge plasma membrane which leads to Piezo1 recruitment and activation.
Piezo1 activation leads to calpain activation which potentially drives further cytoskeletal consolidation
to recruit integrin LFA1. LFA1 recruitment and activation lead to phosphorylation of AKT and
downstream signaling eventually driving the retrograde actin flow in migrating human T cells.

Figure 7

Figure 1

Vector Units

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6