Detecting anomalous content in dairy with whole metagenome sequencing

Niina Haiminen,[a †] Kristen L. Beck,[b †] Akshay Agarwal,[b] Anna Paola Carrieri,[c] Ban Kawas,[b] Martin Wiedmann,[d] Erika Ganda[e*]

[a]IBM T. J. Watson Research Center, Yorktown Heights NY, USA; [b]IBM Almaden Research Center, San Jose CA, USA; [c]IBM Research Europe - Daresbury, Warrington, UK; [d]Cornell University, College of Agriculture and Life Sciences, Ithaca NY, USA; [e]Penn State University, Department of Animal Science, University Park PA, USA

†Niina Haiminen and Kristen L. Beck contributed equally to this work. Author order was determined by reverse alphabetic order.

ABSTRACT

The increasing knowledge of microbial ecology in food products relating to quality and safety as well as the established usefulness of machine learning algorithms for anomaly detection in multiple scenarios suggests that the application of microbiome data in food production systems for anomaly detection is an important next step toward bettering our food supply chain. To that end, the objective of the research presented here was to assess the feasibility of using whole metagenome shotgun sequencing data as input into anomaly detection algorithms using fluid milk as a model system. Contrastive PCA, cluster-based methods, and explainable AI were evaluated as potential methods for detection of two anomalous sample classes using longitudinal metagenomic profiling of fluid milk compared to baseline samples collected under comparable circumstances. Traditional methods (alpha and beta diversity, clustering-based contrastive PCA, MDS, and dendrograms) failed to differentiate anomalous sample classes; however, explainable AI was able to classify anomalous vs. baseline samples and indicate microbial drivers in association with antibiotic use e.g., *Morganella* and *Enterobacter* and microbial signatures related to milk collected from an alternative farm e.g., *Coxiella*. This work also characterizes the microbiome of fluid milk in greater depth than previously published studies. Our results indicate the application of artificial intelligence continues to hold promise in the realm of microbiome data analysis and could present further opportunity for downstream analytic automation to aid in food safety and quality.

IMPORTANCE   We evaluated the feasibility of using untargeted metagenomic sequencing of raw milk for detecting anomalous food ingredient content with artificial intelligence methods. The approach could potentially be applied in the food industry for safety and quality control.

KEYWORDS: Metagenome, Microbiome, Food, Milk, Anomaly Detection, Machine Learning, Explainable Artificial Intelligence

INTRODUCTION

Issues in food quality and safety can have rippling effects through the supply chain causing substantial health and economic damages. In this work, bovine raw milk was selected as a model ingredient as it is the sole ingredient used for production of fluid milk— a high-volume food with considerable concern for fraud and food safety issues. Beyond this, milk is used as an ingredient to make a variety of products and as other

foods, with raw milk quality having considerable impacts on finished product quality, safety, and production effciency. Other studies have aimed to characterize the microbiome of food ingredients in production settings for example in high protein powders ([1], [2]), produce ([3], [4]), and fermented foods ([5], [6], [7], [8]). These studies are useful in demonstrating the potential that metagenomics and metatranscriptomics have in advancing food safety and quality for targeted assessments as well as for improving sensitivity for regular surveillance. Metagenomic and metatranscriptomic studies have been able to describe the microbial components of food samples with observable shifts that can be related back to key attributes of metadata e.g. ingredient contamination ([1]). However, it should be noted that when studies rely on amplicon sequencing (often due to cost and resource limitations), there can be important reductions in sensitivity and taxonomic resolution.

The food supply chain is highly complex, with a multitude of touch points (e.g. farmers, suppliers, transportation, storage, etc.) existing prior to reaching a finished product, where issues occurring at each step have the potential to cause quality and safety issues. Therefore, while these early studies have set a foundation for the use of the microbiome in food production, the expansion of these analyses and applications into additional food ingredients and for different supply chain challenges will only continue to refine and increase the robustness of analysis similar to how much work has been done to build microbiome standards across human-associated ([9], [10], [11], [12]) and environmental niches ([13]).

Our objective was to expand on the growing evidence that the microbiome can be used as relevant indicator through an application in fluid milk with the hypothesis that it could be used to detect two key anomalies: the introduction of milk from an outside farm and the presence of milk from mastitis-affected cows treated with antibiotics which should instead be withheld from bulk tanks (referred to as OF and ABX, respectively). These anomalies were selected to mimic potential real issues that could be faced in the dairy industry (e.g., introduction of an unknown supplier into an ingredient stream or violation of industry regulations).

Milk was used as a model system for examining the application of metagenomic sequencing of microbial communities for food safety and quality, building on our earlier work evaluating associated DNA extraction and host depletion methods ([14]). Raw milk microbiomes have been found to be diverse ([15], [16]) and potentially have an influence on the quality of downstream processed dairy products ([17], [18]). These and other published works support utilizing milk microbiomes as a potential source of information for quality assurance and risk assessment in the food industry.

We evaluated different anomaly detection methods beginning with the classical microbiology ecological metrics of alpha and beta diversity, differential abundance, clustering, as well as ordination through contrastive PCA and MDS. However, these classical methods were limited in their ability to differentiate sample classes. In turn, a growing number of studies have demonstrated the benefit of leveraging machine learning to differentiate sample classes in microbiome studies. These include predicting the risk of type 2 diabetes ([19]), diarrhea associated with cancer treatment ([20]), and liver disease ([21]) from the gut microbiome. Additionally, when sampling the microbiome from leg skin, explainable AI was able to identify microbial drivers associated with skin hydration, age, and pre/post-menopausal status from the skin microbiome ([22]).

For this work, we collected 58 bulk tank milk samples in a block-randomized time-constrained design to assess the ability of the microbiome to indicate deviations from a baseline (BL) community related to anomalies (outside farm and antibiotic use) that

FIG 1 Overall analysis pipeline.

could be present in the food supply chain and be related to food quality issues. A
set of 33 consensus microbes were found to be stable elements in baseline shotgun
metagenomics samples with *Pseudomonas*, *Serratia*, *Cutibacterium*, and *Staphylococcus*
to be the most abundant. Traditional methods of ordination (cPCA and MDS) as well as
alpha and beta diversity were limited in their ability to fully separate sample classes
and microbial differences associated with anomalies. However, explainable AI was
able to differentiate sample classes while also identifying three key microbial drivers
which separated sample classes with significance even in this dataset, which would be
considered small in the realm of machine learning techniques studies. Nevertheless,
this study characterized raw milk metagenomes in more depth than any other pub-
lished work and provides advances in the application of machine learning that can be
expanded across the food industry.

RESULTS

**Collection and shotgun metagenome sequencing of baseline and anomalous
raw milk samples** For whole metagenome shotgun sequencing, in total 65 samples
were collected: 33 baseline (BL), 13 outside farm (OF), 12 antibiotic treated (ABX), 6
negative DNA extraction controls, and 1 sequencing blank. Anomalies were chosen
to represent potential sources of concern in a dairy processing plant (milk from an
unknown source or milk contaminated with antibiotic residues). The sampling scheme
is shown in Supplemental Figure S1.

Sampling dates were block randomized to ensure even distribution across the
sampling period for each anomaly type. A short time frame of five weeks was cho-
sen to control for seasonality.

Metadata including milk components (e.g., lactose, fat, milk urea nitrogen), so-
matic cell count, and standard plate count were also collected and provided as Supple-
mental File S1. No strong correlations were observed between the metadata features
and individual microbe reads per million (RPM) values or the summed microbe RPM
values per sample (|Spearman corr.| < 0.7).

Shotgun whole metagenome sequencing (Methods Section 3.2) with Illumina No-
vaSeq 6000 at 2 × 150bp resulted in 39.6-79.2 million read pairs per raw milk sample.
Figure 1 details the overall bioinformatics pipeline that was applied to this data.

During quality control of the reads with FastQC (23) and manual inspection, full
length junction adapters and trailing G's were observed, indicating that the DNA was
likely fragmented prior to sequencing. These artifacts were removed (Methods Section
3.3), and resulting reads were trimmed for low quality bases using TrimGalore (24). A
median of 59.3M read pairs per sample were retained as high quality with a median

of 0.5% read pairs needing removal due to quality issues (see Supplemental Figure S2
and Supplemental Table S2).

To remove bovine and potential contaminant sequence content, we employed ma-
trix filtering as reported in a recent publication on food microbiome sequencing (1).
Kraken (25) was utilized with a custom-built reference database of 31 common food
ingredient and contaminant genomes (1) including *Bos taurus* (assembly Btau_5.0.1)
and *Homo sapiens* (assembly GRCh38.p10). A large fraction, 91.3–98.7%, of reads were
discarded from subsequent microbial analysis as matrix-classified (see Supplemental
Figure S2(a) and Supplemental Table S2). Negative sequencing controls were utilized
to quantify the presence of typical laboratory contaminants that can be expected in
shotgun sequencing. With this, the background microbial contamination was iden-
tified bioinformatically using the decontam (26) R package on the genus-annotated
reads. The analysis identified 14 genera which were removed from subsequent analy-
sis: *Histophilus*, *Rahnella*, *Raoultella*, *T4virus*, *Pragia*, *C2virus*, *Methylophilus*, *Oceanobacil-
lus*, *Streptosporangium*, *Fluviicola*, *Oenococcus*, *Alkalilimnicola*, *Geminocystis*, and *Bre-
vibacillus*.

Microbiome characterization of raw bulk tank milk We utilized Kraken (25)
with the NCBI RefSeq genome collection (27) to annotate the high quality non-matrix
read pairs and summarized the classified reads at genus level (Supplemental Table S3).
Each genus observed was defined based on a minimum abundance of 0.1 Reads per
Million (RPM) as indicated in (1) resulting in 572 observed genera.

Alpha diversity was calculated using the Shannon index from the genus level table.
Four baseline and two outside farm samples (BL-04, BL-08, BL-13, BL-16, OF-09, and
OF-13) were observed to have very low diversity (Shannon index <0.4, Figure 2a), yet a
higher relative proportion of reads assigned to microbes (Supplemental Figure S2(b))
together indicating likely contamination (either at the farm or laboratory level). Be-
yond this, these samples were dominated by a single organism where 94–95% of an-
notated reads were classified as *Pseudomonas* or *Staphylococcus* (in the case of OF-13).
Therefore, these six samples were removed from further analyses.

The remaining samples had an average of 27K microbial classified reads at genus
taxonomic rank or more specific. The alpha diversity within each sample class was rela-
tively consistent (Figure 2). The average Shannon index per sample class was 1.84–1.95
with an average number of genera observed per sample class to be 143–162 with no
major differences due to sampling date. Between sample classes, the alpha diversity
shows a similar distribution with a Wilcoxon rank sum test for BL vs OF p = 0.7654 and
BL vs ABX p = 0.1194. Additionally, there was slightly lower diversity in some samples
treated with antibiotic compared to baseline or outside farm, as one would expect
(Figure 2b).

Classified reads per million quality-controlled sequenced reads (RPM) were com-
puted for each genus and a threshold of 0.1 RPM applied to define *supported* genera,
as described in Beck et al. (1). The supported genera RPM values are provided as Sup-
plemental Table S4. Figure 3 highlights the most abundant microbes observed in the
milk microbiomes. Genera with RPM greater than 5% of the supported genera RPM
total in at least one sample are shown (with remaining genera summed as "Other"). In
total, there were 12 such genera, and combined they account for 73.5–97.3% of the to-
tal supported genera RPM sum per sample. Only *Pseudomonas*, *Serratia*, *Cutibacterium*,
and *Staphylococcus* were observed to account for more than 5% of the total supported
genera RPM in every sample.

Additionally, a date-localized increase in abundance of *Cutibacterium* from 03-Oct.
to 07-Oct. was observed in baseline, outside farm, and antibiotic treated samples.

The anomalous samples were from sub-sampled dates based on our block random-
ized sampling selection (in comparison to continuous date sampling for the baseline).
Additionally, there were also a select number of microbes observed in only a few sam-
ples within one class. For example, *Klebsiella* was only observed in BL-01, BL-17, and
BL-33 as high abundance, and *Enterobacter* was uniquely observed in antibiotic treated
samples (ABX-01) as was *Elizabethkingia* (ABX-10, ABX-11).

Contrasting baseline and anomalous community profiles using traditional
methods To contrast sample classes and better understand microbial drivers asso-
ciated with anomalous sample classes, we began by exploring traditional microbial
ecology and ordination methods.

Beta diversity was calculated by abiding by principles of compositional data analy-
sis and computing an Aitchison distance as described in Beck et al. (1) and in Methods
Section 3.5. This distance was used to cluster samples as shown in Figure 4a. The sam-
ples clustered into three main clades with intermixed sample classes and sampling
dates. The first clade (furthest left, OF-11, BL-29, BL-32, BL-30, ABX-12) was driven by
the presence of *Propionibacterium*, but not with the co-occurrence with *Klebsiella* as in
BL-33. The remaining two larger clades intermixed all sample types and dates without
notable microbial differences defining their structure and separations.

Contrastive PCA (cPCA) (28) was applied to the microbial community relative abun-
dance data. However, it was not able to successfully separate the outside farm, an-
tibiotic, and baseline samples when baseline samples were used as the background
dataset (Supplemental Figure S3). The difference between PCA and cPCA is that cPCA
aims to identify enriched patterns in a target dataset (foreground dataset) by contrast-
ing it with another dataset (background dataset). This is an unsupervised technique
that uses a hyperparameter named alpha to adjust the trade off between high target
variance and low background variance. At alpha = 0, cPCA collapses to PCA on the
target dataset. At alpha = inf, it puts an infinite penalty on any direction which is not
in the null space of the background dataset.

In contrast to cPCA that operates on the microbial RPM count table, multidimen-
sional scaling ingests a pairwise distance matrix and aims to project the samples onto
a lower dimensional space while retaining their distances. Multidimensional scaling
(MDS) based on the Aitchison distance indicated some separation between the anoma-
lous sample types ABX and OF in the two-dimensional projection (Figure 4b), although
the baseline samples appear intermixed with the anomalous samples. The three classes
are significantly separated according to PERMANOVA (p = 0.0064).

Differentially abundant genera Two-sample Kolmogorov-Smirnov tests were per-
formed independently for each genus to determine statistically significant differen-
tially abundant features. After Bonferroni correction for multiple testing, the adjusted
p-values were significant ($p < 0.01$) for *Coxiella* for BL vs. OF and *Enterobacter*, *Mor-*
ganella for BL vs. ABX. Their RPM distributions are visualized in Figure 5. *Coxiella* was
observed to be increased in outside farm samples and *Enterobacter* and *Morganella*
in antibiotic treated samples. Most notably, the median RPM value of *Enterobacter* in
ABX samples was nearly 12 times that of the baseline samples (28.0 vs. 2.4 RPM).

Sample class prediction with explainable AI We employed an explainable AI
workflow, as described in a recent study on the skin microbiome (22), to perform two
separate classification tasks: BL vs. OF and BL vs. ABX. The genus-level RPM data after
removing low-diversity outlier samples and decontaminant genera was used as input
(52 samples, 572 features). For each classification task, the samples were split into
training (70%) and test (30%) sets uniformly at random, while maintaining the class
size distribution in each set. Five randomized iterations of the train/test split were

performed to obtain robust results. To evaluate and compare the predictive perfor-
mance of our machine learning model we used a state-of-art measure of accuracy, the
F1-score. The F1-score is the harmonic mean between precision and recall.

Three machine learning models (XGBoost, Random Forest, LightGBM) were hyper-
tuned on the training data to then perform the classification task on unseen samples
in the test data. After comparing the predictive performance of the tuned machine
learning models, the best model overall was selected. XGBoost, a gradient-boosted
decision tree ensemble method, consistently reported a higher F1-score on the test
data and during cross validation. XGBoost has also performed well in recent compar-
ative studies on microbiome data (29, 22, 30).

Generating explanations for the ML models' predictions is an important active
field of research. Understanding and explaining the mechanisms underlying the pre-
dictions can help validate the predictive models and reveal powerful and novel in-
sights about the connections between the samples and phenotype under investiga-
tion. Therefore, for each classification task, we generated explanations of the predic-
tions for XGBoost using an explainable AI algorithm called SHapley Additive exPlana-
tions (SHAP) (31). The SHAP algorithm assigns a SHAP value to each genus (i.e., feature)
that represents the impact, negative or positive, that the genus has in predicting a class
for a given sample. The genera are then ranked based on their average absolute SHAP
impact value across all the samples in the training dataset, to obtained a ranked list
of impactful genera (see Methods Section 3.7 for more details).

Given the number of samples per class, the random variability in the splitting of
samples into training and test sets has an effect on the predictive performance and the
ranked list of the most impactful genera (i.e, when running multiple iterations while
changing the global random seed). As such, the predictive performance and the stabil-
ity (32) of the top impactful genera from SHAP was examined across five randomized
iterations. We observed overall high variation in the order of the impactful genera
across the five randomized iteration with different random seeds. *Stable predictive*
features were defined as those being among the top three most impactful features in
at least two out of five randomized iterations.

Despite the observed variation, three stable impactful features were identified:
Coxiella for the BL vs. OF comparison as well as *Enterobacter* and *Morganella* for the BL
vs. ABX comparison. These stable features that are impactful for the prediction are
the same as the differentially abundant genera identified above with the KS-test.

In addition to confirming the predictive impact of the three statistically significant
genera, one advantage of the explainable AI algorithm, SHAP, over other feature im-
portance methods, is that it also explains how each of these impactful features is con-
tributing, positively or negatively, to the prediction of a particular class (e.g., BL) for
each samples or across a set of samples.

In Figure 6(a) for one of the five runs, *Coxiella* is shown as positively contributing
to the prediction of the class OF for those samples that have a higher abundance of
Coxiella. Similarly, in Figure 6(b) *Enterobacter* and *Morganella* are driving the prediction
of ABX for those samples with higher abundances of the genera. The SHAP plots for
the remaining four iterations are shown in Supplemental Figure S4.

For the BL vs. OF prediction, in 4/5 runs *Coxiella* was the most impactful and in 1/5
runs it was the second most impactful. The F1-score of BL vs. OF prediction ranged
from 0.833 to 0.917 (mean 0.87). For the BL vs. ABX prediction, in 4/5 runs *Morganella*
was the most impactful. In 1/5 runs *Enterobacter* was the most impactful and in 2/5
runs the second most impactful. The F1-score of BL vs. OF prediction ranged from
0.692 to 0.923 (mean 0.83).

(a)

(b)

FIG 2 Alpha diversity (Shannon index) of all raw milk samples where presence is indicated as genus RPM > 0.1. **(a)** Shannon index is shown for all 58 samples and **(b)** summarized by sample class with the six low-diversity outlier samples removed here and in all subsequent figures ("Baseline_04", "Baseline_08", "Baseline_13", "Baseline_16", "OutsideFarm_09", "OutsideFarm_13").

(a)

(b)

FIG 3 Microbial community membership is shown per sample for genera with RPM abundance greater than 5% in any sample in (**a**) standard stacked barplot and (**b**) with values scaled to 100% by sample. Genera observed in less than 5% of the summed per sample abundance are aggregated into the "Other" class.

(a)

(b)

FIG 4 Beta diversity using Aitchison distance of raw milk samples with six outlier samples removed is shown here in **(a)** hierarchical clustering and **(b)** multidimensional scaling.

FIG 5 Distribution of RPM values for each group, for the differentially abundant genera. **(a)** Outside farm vs. baseline samples. **(b)** Antibiotic treated vs. baseline samples. Unadjusted and adjusted p-values from the two-sample KS-test are shown within each plot. White circles indicate median values.

(a) Outside farm vs. baseline **(b)** Antibiotic treated vs. baseline

FIG 6 Explainable AI results, SHAP summary dot plots in one exemplar iteration of each anomaly type comparison against baseline (see Supplemental Figure S4 for the other four iterations). SHAP values indicate the importance of that feature on prediction of the sample class (see further explanation in Section 3.7). **(a)** The most impactful features predicting outside farm anomaly class. **(b)** The most impactful features predicting antibiotic treated anomaly class.

DISCUSSION

We have applied various traditional and artificial intelligence methods to test the hypothesis that the microbial community of raw milk, as characterized by shot gun metagenomics, could be used for anomaly detection based on the intuition that the microbiome would differ in relation to two anomalous states: (i) treatment with antibiotics and (ii) the presence of milk from a differing farm. The rationale for testing such anomalous states was to attempt to detect anomalies that would be meaningful to the industry (e.g., regulation violation or unknown ingredient source) (33, 34). The microbiome has been shown to be a highly dynamic ecosystem where microbial community membership and relative abundances can shift in response to a variety of perturbations (35). This has been demonstrated in health (36), the environment (37), and more recently in food systems (1, 38, 39), further substantiating our motivation.

While our study was deliberately designed to control for seasonality and thus represents a worst-case, but most realistic, scenario in terms of what would be useful for application in anomaly detection in the industry, our results indicate that under such circumstances, explainable AI applied to microbiome data might become a valuable tool for monitoring and anomaly detection in food systems in the future.

Raw milk microbial composition is dominated by few typical milk genera and overall uniform across sample types To the best of our knowledge, this study characterized raw milk metagenomes in more depth than any other published work to date and demonstrates that there is a set of consensus microbes that were found to be stable elements across samples. We observed 32 microbial genera as present (RPM > 0.1) in all samples (excluding low-diversity outliers), see Supplemental Table S5. *Pseudomonas*, *Serratia*, *Cutibacterium*, and *Staphylococcus* were the most abundant on average. This is in agreement with most of the literature, with *Pseudomonas* and *Staphylococcus* being reported in many studies that characterized the raw milk microbiome (40, 15, 41, 42). Mastitis causing bacteria, including *Streptococcus sp.*, *Staphylococcus sp.*, and coliforms such as *E. coli* and *Klebsiella* have been reported in milk microbiome studies (43, 44, 45), — all of which were detected in our dataset. Consequently, our data along with previously published data indicate that bacteria such as *Pseudomonas*, and *Staphylococcus*, among others, represent the core bulk tank microbiome.

In our study *Cutibacterium* was one of the most prevalent genera. *Cutibacterium* belongs to the family *Propionibacteriaceae*, which have been reported in the teat apex microbiota (40). A few skin-associated *Propionibacteria* have been reclassified as *Cutibacterium* (46), indicating that the teat skin microbiota might contribute to the bulk tank milk microbiota.

Traditional microbial diversity metrics, clustering, and MDS and methods fail to differentiate sample classes For the detection of anomalous samples, we evaluated several traditional methods for comparative analysis of the microbiome community. In this study clustering, cPCA, and MDS analyses did not indicate a strong separation of the three different classes of samples (baseline, outside farm, and antibiotics). This is in contrast with recently published studies (15, 16), in which milk samples were significantly different based on origin as identified by PERMANOVA and visualized with Principal Component Analysis. One explanation for the disparity is that in those studies, sampling was performed across seasons, thus increasing the variability between sample sites. In this study, we purposely sampled within a short time frame to control for seasonal variability, and thus avoid that confounder in our analyses. While our PERMANOVA results indicate that there was a significant difference in beta diversity between the three classes with multidimensional scaling ($p = 0.0064$), no clear separation could be observed between sample classes when the first two MDS

are plotted. One might argue that most antibiotic samples are closer to one another
than the other two classes, and thus that might explain our significant results within
PERMANOVA. The high degree of uniformity observed here might be explained by our
sampling strategy which was to constrain sampling to a short period of time to con-
trol for seasonality. It is possible that management practices may explain differences
between baseline and outside farm samples in our study.

Bacterial taxa can be used as biomarkers for anomaly detection Despite the
overall uniformity in our data, three genera were identified to be differentially abun-
dant between baseline and anomalous samples. *Enterobacter* have been reported to
be present in milk microbiomes in many studies (40, 15, 41, 42), and *Coxiella* and *Mor-*
ganella were identified in this study but not in many others, perhaps because of to the
large sequencing depth applied here. Nevertheless, these are environmental organ-
isms that could be found in any farm, and their relative abundances could be informa-
tive when attempting to identify anomalies in milk.

In the case of milk received from an outside farm, we aimed to identify if the mi-
crobiome could signal more subtle differences since major influences, e.g. seasonality,
region, and temperature, were all constant. Here while the cows and bulk tanks may
be similar across farms, there were differences in farm management, diet, and milking
processing protocols that one would expect to impart differences in the bulk tank milk
microbiome. While baseline samples and outside farm samples were similar in their
diversity and overall community membership, *Coxiella* was observed to have a signif-
icant increase in the outside farm samples. *Coxiella* is a known foodborne pathogen
and a target organism for defining time and temperature combinations for milk pas-
teurization (47, 48). Thus it is not an unusual finding in raw milk microbiome, reiter-
ating the fact that when using microbiome data for anomaly detection, one should
not rely only on finding which features are unique to a sample class, but also how the
relative abundances of features observed across classes might be associated with a
particular class.

In milk samples spiked with milk from cows treated with antibiotics to control mas-
titis, the overall diversity does not differ from baseline diversity in a significant manner.
Previous studies investigating bovine mastitis and antibiotic treatment have indicated
that alpha diversity is significantly lower in mastitic milk when compared to healthy
milk (49, 50). The lack of difference in diversity in our antibiotic anomalous samples
is in accordance with the design of this study, in which antibiotic anomalies were not
milk from an individual cow, but bulk tank milk spiked with 10% v/v of milk from an an-
imal whose milk should not be entering the food supply chain. This represents a trace
amount of contamination that although more diffcult to detect, would represent a
more realistic scenario. Even in such circumstances, we identified genera e.g. *Enter-*
obacter and *Morganella* that are observed with increased abundance in the antibiotic
sample class. *Enterobacter spp.* are environmental mastitis-causing pathogens (43),
hence expected in the mastitic milk from cows treated with antibiotics. *Morganella* is
an environmental organism and part of the intestinal tract of mammals, and an op-
portunistic pathogen and having been isolated from cheese and (51) and reported to
infect calves (52).

Explainable AI outperforms traditional methods in microbiome-based anomaly
detection and can predict baseline vs. anomalous sample class using combined
signal from all genera For the detection of anomalous samples, we evaluated sev-
eral traditional methods for comparative analysis of the microbiome community. Al-
though clustering, cPCA, and MDS analyses did not indicate a strong separation of the
three different classes of samples, predicting each type of anomaly versus the base-

line could be done fairly accurately with both anomaly types using explainable AI even
in our relatively small dataset. We observed that XGBoost was able to differentiate
anomalous samples from baseline and were able to quantify the impact of the dif-
ferentiating features with SHAP. Likewise, XGBoost, a gradient-boosted decision tree
ensemble method, has also performed well in recent comparative studies on micro-
biome data (29, 22, 30).

While most machine learning algorithms function in a black-box manner, the ex-
plainability algorithm we used in this work— SHAP— assigns a value to each sample for
each feature that describes the impact of that feature for predicting a specific sample
class. One could speculate that this capability will be very useful for the food industry
as evidence is amounting indicating that certain pathogens tend to co-occur with cer-
tain environmental microbes (53, 54). Thus, having the ability to "flag" microbes that
predict a specific problematic sample class and might prove useful to inform sanita-
tion and foodborne pathogen control practices. For example, *Listeria* has been shown
to co-occur with certain taxa and thus observing those taxa as impactful genera in pre-
dicting a specific location might raise awareness about potential future problems with
Listeria (53, 54).

The three genera observed to be most impactful by explainable AI were the same
as those identified with the KS-test after Bonferroni correction; however by leveraging
explainable AI, we were able to use the combined signal from all genera to assess the
impact of the differing abundance of these microbes.

Despite the high metagenomics sequencing depth, we recognize the small nature
of the sample size (58 samples) when it pertains to testing machine learning algorithms
(55, 56, 57), and that it may impart limitations on the quantitative and comparative
assessments in characterizing the microbiomes of the multiple sample classes. Never-
theless, our results provide evidence for the feasibility of this approach and indicate
that explainable AI has the potential to become a useful tool for microbiome-based
quality monitoring for the food industry.

Conclusion and future directions In raw milk and other food systems, microbes
can present important challenges to food safety in the case of pathogenic organisms
and affect food quality such as flavor and storage attributes. Characterizing the mi-
crobial composition in a diverse set of food ingredients and products is of the utmost
importance to better understand and improve the safety and quality of food. Since
the microbiome is sensitive to changes in temperature, salinity, pH, and the compo-
sition of the material that it resides on among other things, it can also be utilized as
an indicator for when food items deviate from a baseline of normality. As sequencing
technologies become more cost effective, laboratory processes are streamlined, and
larger datasets are produced, we demonstrate here that explainable AI applied to mi-
crobiome sequencing data could become a useful approach for anomaly detection in
the food industry.

MATERIALS AND METHODS

Milk sampling Baseline raw bulk tank milk samples were collected daily from the
Cornell University Ruminant Center (CURC; Hartford, NY) between September 5th and
October 7th, 2018 and are referred to by the abbreviation BL. Anomalous samples
from an outside farm (abbreviated as OF) were collected from a collaborating com-
mercial dairy farm located in the same region (< 30 miles) and over the same time pe-
riod as the Cornell Dairy. Bulk tank milk samples were collected aseptically into sterile
10 oz. vials (Capitol Plastics, Amsterdam, NY) and transported on ice to the Milk Qual-
ity Improvement Laboratory (Ithaca, NY). Anomalous samples from antibiotic-treated

cows (abbreviated as ABX) were prepared at the laboratory by spiking the baseline
sample of that day with 10% v/v with milk from an animal currently being treated with
antibiotics, which was collected through a milking system collection device into a 10
oz vial.

Milk samples were aliquoted at the laboratory and frozen at -80°C until DNA extrac-
tion. A volume of 200 µL of milk samples were used as starting material for magnetic-
based DNA extraction using a 96-well plate and CORE kit in a KingFisher instrument
(Thermo Fisher Scientific, San Jose, CA, United States). Negative DNA extraction con-
trols (reagents only) were carried out within the same plate for quality control. Ex-
tracted DNA was frozen at -80°C until library preparation and sequencing.

Shotgun metagenome sequencing Samples were quantified with a Qubit (Thermo
Fisher Scientific, San Jose, CA, United States) before library preparation. Ten nanograms
of each Qubit quantified genomic DNA was sheared with a Covaris E220 instrument
operating SonoLab v6.2.6 generating approximately 300 bp DNA fragments according
to the manufacturer's protocol. Between 10 and 100 ng of fragmented DNA was pro-
cessed into Illumina compatible sequencing libraries using sparQ DNA Library Prep
Kit (Quantabio, Beverly, MA, United States). Each library was barcoded with unique
dual index sequences (NEXTFLEX® Unique Dual Index Barcodes, BioO Scientific). Li-
brary size and amount were confirmed with a Bioanalyzer High Sensitivity DNA chip.
Polymerase chain reaction primers and reagents included in the sparQ kit were used
to perform PCR, and products were purified with AMPure XP beads. Equimolar li-
braries were pooled and subjected to Illumina NovaSeq 6000 sequencing at 2 × 150
bp (Illumina, San Diego, CA, United States). Shotgun whole metagenome sequencing
was performed at the Genome Sciences and Bioinformatics Core at the Pennsylvania
State University College of Medicine, Hershey, PA, United States. Illumina bcl2fastq
(released version 2.20.0.422) was used to extract de-multiplexed sequencing reads.

Read quality control and host filtering Reads that included full length auxil-
iary sequences (junction adapter) P5 or P7 ("CTGTCTCTTATACACATCTCCGAGCCCAC-
GAGAC" or "CTGTCTCTTATACACATCTGACGCTGCCGACGA") were removed with a cus-
tom script (as were their read pairs), since their presence indicates issues with se-
quencing those particular reads (see Figure 2(b-d) in Illumina's Sequencing Technical
Note (58)). Trailing stretches of "G", prevalent in Nextera sequences due to the two
color channel technology, were removed with a custom script. If the length of the "G"
tail was 30 or more, the read and its pair were discarded as low quality. The sequenced
reads were then processed with TrimGalore (24) for adapter and quality trimming (pa-
rameters: –trim-n –paired –length 50 –phred33). The reads were handled as pairs
through all the quality control and filtering steps.

Read filtering against the internal control PhiX and common host and contaminant
genomes was performed with Kraken (25) as in Beck et al (1). PhiX reads were filtered
against the NCBI Reference Sequence: NC_001422.1. Host reads were filtered against
a database of plant and animal sequences introduced previously for metagenomic
studies of food (1), which has been open-sourced and is available via PrecisionFDA
https://precision.fda.gov/home/assets/file-GFfjPQj0ZqJV93P00bJ3vFgG-1. Additionally,
kraken-filter with score threshold 0.1 was applied to avoid removing microbial reads.

Microbial genus profiling The reads passing quality control were classified as in
Beck et al (1), against the NCBI's RefSeq Complete (27) genome collection and corre-
sponding taxonomy of bacterial, archaeal, viral, and eukaryotic microorganisms (ap-
prox. 7,800 genomes retrieved April 2017). Kraken (25) was used with a minimum
score threshold of 0.05. Classified read counts per genus were collected as the sum
of the read counts assigned to a genus or a taxonomic level below it. Sequencing

blanks were used as negative controls to remove contaminating genera with the de-
contam R package (26) with the following parameters: threshold = 0.5 and normalize
= TRUE. From this analysis, there were 14 genera which were removed from subse-
quent analysis: *Histophilus*, *Rahnella*, *Raoultella*, *T4virus*, *Pragia*, *C2virus*, *Methylophilus*,
Oceanobacillus, *Streptosporangium*, *Fluviicola*, *Oenococcus*, *Alkalilimnicola*, *Geminocystis*,
and *Brevibacillus*. Finally, classified reads per million quality-controlled sequenced
reads (RPM) were computed for each genus and a threshold of 0.1 RPM applied to
define *supported* genera, as described in Beck et al. (1).

Community diversity Shannon diversity was calculated from the supported mi-
crobial genera table using the diversity function in the vegan R package (59) with de-
fault parameters. Beta diversity was calculated using principles of compositional data
analysis (60, 61). Therefore, read counts assigned to each genus were pseudo-counted
by adding one in advance of computation of RPM prior to calculating the Aitchison dis-
tance from the microbial table. Beta diversity was calculated using the R package rob-
Compositions (62) and hierarchical clustering was performed using base R function
hclust using the "ward.D2" method.

Contrastive PCA (28) Python implementation was run with the aim of identifying
enriched patterns in outside farm and antibiotic treated samples by contrasting them
with baseline samples, on the supported microbial genera (Supplemental Table S4).
In our target (foreground) data we kept OF, ABX and BL samples. In our background
dataset, we only kept the baseline samples. We removed the 6 low diversity outliers
from both target and background dataset. This was in an effort to uncover compo-
nents which have high variance in the target dataset but low variance in the back-
ground dataset. We tried automatic assignment of alpha values where the algorithm
generates and evaluates different alpha values and we also experimented with increas-
ing alpha values systematically.

Multidimensional scaling (MDS, Matlab function cmdscale, p=2) and permutational
multivariate analysis of variance (PERMANOVA, function f_permanova, iter=10,000, from
the Fathom toolbox (63) for MATLAB) were applied on the pairwise Aitchison distances
of all samples excluding the four baseline and two outside farm samples identified as
low-diversity outliers, on the supported microbial genera table.

Differences between baseline and outside farm, antibiotic treated samples
Two-sample Kolmogorov-Smirnov tests (MATLAB function kstest2) were performed for
each genus to determine microbes with significant differential abundance (p < 0.001)
between baseline and each type of anomaly. Bonferroni multiple-comparison correc-
tion was applied, $p' = pm$, where m is the number of genera, to obtain adjusted p-
values p'. The differentially abundant genera were visualized using Violin Plot (64) in
MATLAB.

Explainable AI As in (22) for each prediction tasks we compared the predictive
performances of various machine learning models, after hyper-tuning on the training
data, and selected the best model at predicting each phenotype under investigation.
The explanations for the predictions were then generated for the best ML model (i.e.,
XGBoost).

We used an explainable AI algorithm called SHapley Additive exPlanations (SHAP),
because of its ability to work with any machine learning model: tree-based models,
such as XGBoost and LightGBM, as well as kernel-based and deep learning models.
The explainability algorithm, SHAP, provides local explanations, i.e., interpretations of
how the model predicts a particular value for a given sample. The local explanations
show how each feature is contributing, either positively or negatively, to the prediction
of a particular instance, for example of a particular class in case of classification task.

We applied the tuned XGBoost model coupled up with SHAP to explain the predictions (e.g., baseline vs anomalous) for each sample across the entire dataset. In addition to providing the ranked list of important features for a ML model, an advantage of SHAP over other feature importance methods is that it also explains how each of these impactful features is contributing (positively or negatively) to the prediction of specific phenotypic values. If we consider a binary classification task, the SHAP explainer returns two Shapley values tables of the same dimension of the input table (number of samples x number of genera/features), respectively for the class 0 (baseline) and the class 1 (anomalous). If we examine the table for class-0 baseline, each entry in the table is the SHAP impact (positive or negative) that a given genus has for the prediction of class baseline for a given sample. The absolute SHAP impact values for each feature are then averaged out across the entire set of samples to get an indication of the overall impact of a feature for the prediction, which results in a *ranked list of the most impactful features*.

Several plots can be generated using SHAP to provide various visualisations of local explanations as well as a global view of local explanations that allow for interpretation of the entire model. The *SHAP summary dot plot*, in particular, is a visualisation of the Shapley values matrix for a particular class (e.g., baseline). The plot shows the impact that each feature has on the prediction of the class for samples that share similar feature values. The y-axis is the ordered list (descending order of importance) of impactful features in predicting the class using a specific ML model (e.g., XGBoost). Therefore each row is a feature. The dots in each row are the data points, or samples, and are colored by the original feature value, that in this case is the genus abundance. The x-axis is the SHAP value or impact. A positive SHAP value/impact of a feature for a sample (the dot is on the right side of the x-axis) indicates that the feature (e.g., genus) has a positive impact in predicting the class (e.g., baseline), while a negative SHAP impact (the dot is on the left side of the x-axis) indicated that the feature has a negative impact on the prediction of the class. For each row (feature) the yellow and green dots can form separate clusters that are positioned towards the right or left side of the x-axis. This indicates that overall the feature (e.g., genus) tend to have a similar impact (positive or negative) for samples in which it has similar feature values (e.g., high or low abundance). We used the python implementation of SHAP available at https://github.com/slundberg/shap. To obtain the appropriate SHAP explainer for XGBoost we combined the tree explainer with the best optimised model by applying the function shap.TreeExplainer(tunedmodel). We used the resulting explainer to compute the matrix of Shapley values for both the training and the test datasets.

Data availability The sequencing data generated in this study are available at the NCBI BioProject PRJNA726965 https://www.ncbi.nlm.nih.gov/bioproject/PRJNA726965. The code used to generate analyses in this study is available at https://github.com/gandalab/milk-anomaly-detection.

SUPPLEMENTAL MATERIAL

FIG S1. Sampling scheme.

FIG S2. Read counts per sample.

FIG S3. Contrastive PCA results.

FIG S4. Explainable AI results.

TABLE S1. Metadata on the raw milk samples.

TABLE S2. Summary of read counts.

TABLE S3. Read counts per genus.

TABLE S4. Supported genera RPM with contaminants removed.

TABLE S5. Core genera RPM with contaminants and outliers removed.

ACKNOWLEDGMENTS

The authors would like to thank David Chambliss at IBM Research for his support with
read filtering and data processing. This work was supported by the USDA National
Institute of Food and Agriculture and Hatch Appropriations under Project #PEN04752,
Accession #1023328 awarded to Erika Ganda. Sequencing was partially supported by
a gift from IBM to Cornell University.

REFERENCES

1. **Beck KL, Haiminen N, Chambliss D, Edlund S, Kunitomi M, Huang BC, Kong N, Ganesan B, Baker R, Markwell P, Kawas B, Davis M, Prill RJ, Krishnareddy H, Seabolt E, Marlowe CH, Pierre S, Quintanar A, Parida L, Dubois G, Kaufman J, Weimer BC**. 12 2021. Monitoring the microbiome for food safety and quality using deep shotgun sequencing. npj Sci Food 5 (1). doi:10.1038/s41538-020-00083-y.

2. **Haiminen N, Edlund S, Chambliss D, Kunitomi M, Weimer BC, Ganesan B, Baker R, Markwell P, Davis M, Huang BC, Kong N, Prill RJ, Marlowe CH, Quintanar A, Pierre S, Dubois G, Kaufman JH, Parida L, Beck KL**. 2019. Food authentication from shotgun sequencing reads with an application on high protein powders. npj Sci Food 3 (24). doi:10.1038/s41538-019-0056-6.

3. **Martins G, Miot-Sertier C, Lauga B, Claisse O, Lonvaud-Funel A, Soulas G, Masneuf-Pomarède I**. 8 2012. Grape berry bacterial microbiota: Impact of the ripening process and the farming system. Int J Food Microbiol 158:93–100. doi:10.1016/J.IJFOODMICRO.2012.06.013.

4. **Droby S, Wisniewski M**. 6 2018. The fruit microbiome: A new frontier for postharvest biocontrol and postharvest biology. Postharvest Biol Technol 140:107–112. doi:10.1016/J.POSTHARVBIO.2018.03.004.

5. **Walsh AM, Crispie F, Kilcawley K, O'Sullivan O, O'Sullivan MG, Claesson MJ, Cotter PD**. 10 2016. Microbial Succession and Flavor Production in the Fermented Dairy Beverage Kefir. mSystems 1. doi:10.1128/mSystems.00052-16.

6. **Walsh AM, Crispie F, O'Sullivan O, Finnegan L, Claesson MJ, Cotter PD**. 12 2018. Species classifier choice is a key consideration when analysing low-complexity food microbiome data. Microbiome 6:50. doi:10.1186/s40168-018-0437-0.

7. **Duru IC, Laine P, Andreevskaya M, Paulin L, Kananen S, Tynkkynen S, Auvinen P, Smolander OP**. 9 2018. Metagenomic and meta-transcriptomic analysis of the microbial community in Swiss-type Maasdam cheese during ripening. Int J Food Microbiol 281:10–22. doi:10.1016/J.IJFOODMICRO.2018.05.017.

8. **Williams AG, Choi SC, Banks JM**. 1 2002. Variability of the species and strain phenotype composition of the non-starter lactic acid bacterial population of cheddar cheese manufactured in a commercial creamery. Food Res Int 35:483–493. doi:10.1016/S0963-9969(01)00147-8.

9. **Vangay P, Burgin J, Johnston A, Beck KL, Berrios DC, Blumberg K, Canon S, Chain P, Chandonia JM, Christianson D, Costes SV, Damerow J, Duncan WD, Dundore-Arias JP, Fagnan K, Galazka JM, Gibbons SM, Hays D, Hervey J, Hu B, Hurwitz BL, Jaiswal P, Joachimiak MP, Kinkel L, Ladau J, Martin SL, McCue LA, Miller K, Mouncey N, Mungall C, Pafilis E, Reddy TBK, Richardson L, Roux S, Schriml LM, Shaffer JP, Sundaramurthi JC, Thompson LR, Timme RE, Zheng J, Wood-Charlson EM, Eloe-Fadrosh EA**. 2 2021. Microbiome Metadata Standards: Report of the National Microbiome Data Collaborative's Workshop and Follow-On Activities. mSystems 6. doi:10.1128/MSYSTEMS.01194-20.

10. **Human Microbiome Project Consortium**. 2012. A framework for human microbiome research. Nature 486:215–221. doi:10.1038/nature11209.

11. **Greathouse KL, Sinha R, Vogtmann E**. 2019. DNA extraction for human microbiome studies: The issue of standardization. Genome Biol 20:1–4. doi:10.1186/s13059-019-1843-8.

12. **Costea PI, Zeller G, Sunagawa S, Pelletier E, Alberti A, Levenez F, Tramontano M, Driessen M, Hercog R, Jung FE, Kultima JR, Hayward MR, Coelho LP, Allen-Vercoe E, Bertrand L, Blaut M, Brown JRM, Carton T, Cools-Portier S, Daigneault M, Derrien M, Druesne A, Vos WMD, Finlay BB, Flint HJ, Guarner F, Hattori M, Heilig H, Luna RA, Vlieg JVH, Junick J, Klymiuk I, Langella P, Chatelier EL, Mai V, Manichanh C, Martin JC, Mery C, Morita H, O'toole PW, Orvain C, Patil KR, Penders J, Persson S, Pons N, Popova M, Salonen A, Saulnier D, Scott KP, Singh B, Slezak K, Veiga P, Versalovic J, Zhao L, Zoetendal EG, Ehrlich SD, Dore J, Bork P**. 10 2017. Towards standards for human fecal sample processing in metagenomic studies. Nat Biotechnol 2017 35:11 35:1069–1076. doi:10.1038/nbt.3960.

13. **Thompson L, Sanders J, McDonald D, Amir A, Ladau J, Locey K, Prill R, Tripathi A, Gibbons S, Ackermann G, Navas-Molina J, Janssen S, Kopylova E, Vázquez-Baeza Y, González A, Morton J, Mirarab S, Zech Xu Z, Jiang L, Haroon M, Kanbar J, Zhu Q, Jin Song S, Kosciolek T, Bokulich N, Le er J, Brislawn C, Humphrey G, Owens S, Hampton-Marcell J, Berg-Lyons D, McKenzie V, Fierer N, Fuhrman J, Clauset A, Stevens R, Shade A, Pollard K, Goodwin K, Jansson J, Gilbert J, Knight R**. 11 2017. A communal catalogue reveals Earth's multiscale microbial diversity. Nat 2017 551:7681 551:457–463. doi:10.1038/nature24621.

14. **Ganda E, Beck KL, Haiminen N, Silverman JD, Kawas B, Cronk BD, Anderson RR, Goodman LB, Wiedmann M, Gibbons SM**. 2021. DNA Extraction and Host Depletion Methods Significantly Impact and Potentially Bias Bacterial Detection in a Biological Fluid. mSystems 6 (3):e00619–21. doi:10.1128/mSystems.00619-21.

15. **Skeie SB, Håland M, Thorsen IM, Narvhus J, Porcellato D**. 2019. Bulk tank raw milk microbiota differs within and between farms: A moving goalpost challenging quality control. J Dairy Sci 102:1959–1971. doi:10.3168/JDS.2017-14083.

16. **Yap M, Gleeson D, O'Toole PW, O'Sullivan O, Cotter PD, Björkroth J**. 2021. Seasonality and Geography Have a Greater Influence than the Use of Chlorine-Based Cleaning Agents on the Microbiota of Bulk Tank Raw Milk. Appl Environ Microbiol 87 (22):e01081–21. doi:10.1128/AEM.01081-21.

17. **Porcellato D, Aspholm M, Skeie SB, Monshaugen M, Brendehaug J, Mellegård H**. 2018. Microbial diversity of consumption milk during processing and storage. Int J Food Microbiol 266:21–30. doi:10.1016/j.ijfoodmicro.2017.11.004.

18. **McHugh AJ, Feehily C, Fenelon MA, Gleeson D, Hill C, Cotter PD, Gilbert JA**. 2020. Tracking the Dairy Microbiota from Farm Bulk Tank to Skimmed Milk Powder. mSystems 5 (2):e00226–20. doi:10.1128/mSystems.00226-20.

19. **Gou W, Ling CW, He Y, Jiang Z, Fu Y, Fengzhe X, Miao ZL, Ting-yu S, Jie-Sheng L, Zhu H, Zhou HW, Chen YM, Zheng JS**. 05 2020. Interpretable Machine Learning Algorithm Reveals Novel Gut Microbiome Features in Predicting Type 2 Diabetes. Curr Dev Nutr 4 (Supplement_-2):1559–1559. doi:10.1093/cdn/nzaa062_016.

20. **Wong CW, Yost SE, Lee JS, Gillece JD, Folkerts M, Reining L, Highlander SK, Eftekhari Z, Mortimer J, Yuan Y**. 2021. Analysis of Gut Microbiome Using Explainable Machine Learning Predicts Risk of Diarrhea Associated With Tyrosine Kinase Inhibitor Neratinib: A Pilot Study. Front Oncol 11. doi:10.3389/fonc.2021.604584.

21. **Liu Y, Mé G, Havulinna AS, Salomaa V**. 2022. Early prediction of incident liver disease using conventional risk factors and gut-microbiome-augmented gradient boosting. Cell Metab 34:719–730.e4. doi:10.1016/j.cmet.2022.03.002.

22. **Carrieri AP, Haiminen N, Maudsley-Barton S, Gardiner L, Murphy B, Mayes A, Paterson S, Grimshaw S, Winn M, Shand C, Hadjidoukas P, Rowe W, Hawkins S, MacGuire-Flanagan A, Tazzioli J, Kenny J, Parida L, Hoptroff M, Pyzer-Knapp E**. 2 2021. Explainable AI reveals changes in skin microbiome composition linked to phenotypic differences. Sci Rep. 11 (4565). doi:10.1038/s41598-021-83922-6.

23. **Andrews S**. 2010. FastQC: A Quality Control Tool for High Throughput Sequence Data http://www.bioinformatics.babraham.ac.uk/projects/fastqc.

24. **Krueger F**. 2012. Trim Galore https://github.com/FelixKrueger/TrimGalore.

25. **Wood DE, Salzberg SL**. 2014. Kraken: ultrafast metagenomic sequence classification using exact alignments. Genome Biol 15 (3):R46. doi:10.1186/gb-2014-15-3-r46.

26. **Davis NM, Proctor DM, Holmes SP, Relman DA, Callahan BJ**. 2018. Simple statistical identification and removal of contaminant sequences in marker-gene and metagenomics data. Microbiome 6 (226). doi:10.1186/S40168-018-0605-2/FIGURES/6.

27. **O'Leary NA, Wright MW, Brister JR, Ciufo S, Haddad D, McVeigh R, Rajput B, Robbertse B, Smith-White B, Ako-Adjei D, Astashyn A, Badretdin A, Bao Y, Blinkova O, Brover V, Chetvernin V, Choi J, Cox E, Ermolaeva O, Farrell CM, Goldfarb T, Gupta T, Haft D, Hatcher E, Hlavina W, Joardar VS, Kodali VK, Li W, Maglott D, Masterson P, McGarvey KM, Murphy MR, O'Neill K, Pujar S, Rangwala SH, Rausch D, Riddick LD, Schoch C, Shkeda A, Storz SS, Sun H, Thibaud-Nissen F, Tolstoy I, Tully RE, Vatsan AR, Wallin C, Webb D, Wu W, Landrum MJ, Kimchi A, Tatusova T, DiCuccio M, Kitts P, Murphy TD, Pruitt KD**. 11 2015. Reference sequence (RefSeq) database at NCBI: current status, taxonomic expansion, and functional annotation. Nucleic Acids Res 44 (D1):D733–D745. doi:10.1093/nar/gkv1189.

28. **Abid A, Zhang MJ, Bagaria VK, Zou J**. 2018. Exploring patterns enriched in a dataset with contrastive principal component analysis. Nat communications 9 (1):2134.

29. **Wang XW, Liu YY**. 2020. Comparative study of classifiers for human microbiome data. Med Microecol 4:100013. doi:10.1016/j.medmic.2020.100013.

30. **Chen Y, Wang H, Lu W, Wu T, Yuan W, Zhu J, Lee YK, Zhao J, Zhang H, Chen W**. 2022. Human gut microbiome aging clocks based on taxonomic and functional signatures through multi-view learning. Gut Microbes 14 (1):2025016. doi:10.1080/19490976.2021.2025016.

31. **Lundberg SM, Lee SI**. 2017. A unified approach to interpreting model predictions. Proceedings of the Neural Information Processing System 30:4765–4774.

32. **Jiang L, Haiminen N, Carrieri AP, Huang S, Vázquez-Baeza Y, Parida L, Kim HC, Swafford AD, Knight R, Natarajan L**. 2021. Utilizing stability criteria in choosing feature selection methods yields reproducible results in microbiome data. Biometrics doi:10.1111/biom.13481.

33. **Montgomery H, Haughey SA, Elliott CT**. 9 2020. Recent food safety and fraud issues within the dairy supply chain (2015–2019). Glob Food Secur 26:100447. doi:10.1016/J.GFS.2020.100447.

34. **FDA**. 2017. Grade "A" Pasteurized Milk Ordinance.

35. **Dini-Andreote F, Pylro VS, Baldrian P, Elsas JDV, Salles JF**. 8 2016. Ecological succession reveals potential signatures of marine–terrestrial transition in salt marsh fungal communities. The ISME J 10:1984. doi:10.1038/ISMEJ.2015.254.

36. **Leung H, Long X, Ni Y, Qian L, Nychas E, Siliceo SL, Pohl D, Hanhineva K, Liu Y, Xu A, Nielsen HB, Belda E, Clément K, Loomba R, Li H, Jia W, Panagiotou G**. 6 2022. Risk assessment with gut microbiome and metabolite markers in NAFLD development. Sci Transl Med 14:855. doi:10.1126/scitranslmed.abk0855.

37. **Dini-Andreote F, Elsas JDV, Olff H, Salles JF**. 12 2018. Dispersal-competition tradeoff in microbiomes in the quest for land colonization. Sci Reports 8. doi:10.1038/S41598-018-27783-6.

38. **Yap M, Ercolini D, Álvarez Ordóñez A, O'Toole PW, O'Sullivan O, Cotter PD**. 2022. Next-Generation Food Research: Use of Meta-Omic Approaches for Characterizing Microbial Communities Along the Food Chain. Annu Rev Food Sci Technol 13 (1):361–384. doi:10.1146/annurev-food-052720-010751.

39. **Mukherjee A, Gómez-Sala B, O'Connor EM, Kenny JG, Cotter PD**. 5 2022. Global Regulatory Frameworks for Fermented Foods: A Review. Front Nutr 0:1084. doi:10.3389/FNUT.2022.902642.

40. **Derakhshani H, Fehr KB, Sepehri S, Francoz D, De Buck J, Barkema HW, Plaizier JC, Khafipour E**. 2018. Invited review: Microbiota of the bovine udder: Contributing factors and potential implications for udder health and mastitis susceptibility. J Dairy Sci 101 (12):10605–10625. doi:https://doi.org/10.3168/jds.2018-14860.

41. **Liu J, Zhu Y, Jay-Russell M, Lemay DG, Mills DA**. 2020. Reservoirs of antimicrobial resistance genes in retail raw milk. Microbiome 8 (1):99. doi:https://doi.org/10.1186/s40168-020-00861-6.

42. **Ruegg PL**. 2022. The bovine milk microbiome – an evolving science. Domest Animal Endocrinol 79:106708. doi:https://doi.org/10.1016/j.domaniend.2021.106708.

43. **Oliveira L, Hulland C, Ruegg PL**. 2013. Characterization of clinical mastitis occurring in cows on 50 large dairy herds in Wisconsin. J Dairy Sci 96 (12):7538–7549. doi:https://doi.org/10.3168/jds.2012-6078.

44. **Bhatt VD, Ahir VB, Koringa PG, Jakhesara SJ, Rank DN, Nauriyal DS, Kunjadia AP, Joshi CG**. 2012. Milk microbiome signatures of subclinical mastitis-affected cattle analysed by shotgun sequencing. J Appl Microbiol 112 (4):639–650. doi:https://doi.org/10.1111/j.1365-2672.2012.05244.x.

45. **Hoque MN, Istiaq A, Clement RA, Sultana M, Crandall KA, Siddiki AZ, Hossain MA**. 9 2019. Metagenomic deep sequencing reveals association of microbiome signature with functional biases in bovine mastitis. Sci Reports 2019 9:1 9:1–14. doi:10.1038/s41598-019-49468-4.

46. **Scholz CFP, Kilian M**. 11 2016. The natural history of cutaneous propionibacteria, and reclassification of selected species within the genus propionibacterium to the proposed novel genera acidipropionibacterium gen. Nov., cutibacterium gen. nov. and pseudopropionibacterium gen. nov. Int J Syst Evol Microbiol 66:4422–4432. doi:10.1099/IJSEM.0.001367/CITE/REFWORKS.

47. **Commission CA, et al.**. 2004. Code of hygienic practice for milk and milk products. CAC/RCP 57-2004. Food Agric Organ Rome .

48. **Wittwer M, Hammer P, Runge M, Valentin-Weigand P, Neubauer H, Henning K, Mertens-Scholz K**. 1 2022. Inactivation Kinetics of Coxiella burnetii During High-Temperature Short-Time Pasteurization of

Milk. Front Microbiol 12:3975. doi:10.3389/fmicb.2021.753871.

49. **Ganda EK, Bisinotto RS, Lima SF, Kronauer K, Decter DH, Oikonomou G, Schukken YH, Bicalho RC**. 2016. Longitudinal metagenomic profiling of bovine milk to assess the impact of intramammary treatment using a third-generation cephalosporin. Sci Reports 6:1–13. doi:10.1038/srep37565.

50. **Ganda EK, Gaeta N, Sipka A, Pomeroy B, Oikonomou G, Schukken YH, Bicalho RC**. 2017. Normal milk microbiome is reestablished following experimental infection with Escherichia coli independent of intramammary antibiotic treatment with a third-generation cephalosporin in bovines. Microbiome 5. doi:10.1186/s40168-017-0291-5.

51. **Ryser LT, Arias-Roth E, Perreten V, Irmler S, Bruggmann R**. 2021. Genetic and Phenotypic Diversity of Morganella morganii Isolated From Cheese. Front Microbiol 12:3378. doi:10.3389/fmicb.2021.738492.

52. **Li G, Niu X, Yuan S, Liang L, Liu Y, Hu L, Liu J, Cheng Z**. 12 2018. Emergence of Morganella morganii subsp. morganii in dairy calves, China. Emerg Microbes & Infect 7. doi:10.1038/S41426-018-0173-3.

53. **Fox EM, Solomon K, Moore JE, Wall PG, Fanning S**. 2014. Phylogenetic profiles of in-house microflora in drains at a food production facility: Comparison and biocontrol implications of listeria-positive and -negative bacterial populations. Appl Environ Microbiol 80:3369–3374. doi:10.1128/AEM.00468-14.

54. **Willis C, Jørgensen F, Aird H, Elviss N, Fox A, Jenkins C, Fenelon D, Sadler-Reeves L, McLauchlin J**. 2 2018. An assessment of the microbiological quality and safety of raw drinking milk on retail sale in England. J Appl Microbiol 124:535–546. doi:10.1111/JAM.13660.

55. **Cui Z, Gong G**. 9 2018. The effect of machine learning regression algorithms and sample size on individualized behavioral prediction with functional connectivity features. NeuroImage 178:622–637. doi:10.1016/j.neuroimage.2018.06.001.

56. **Balki I, Amirabadi A, Levman J, Martel AL, Emersic Z, Meden B, Garcia-Pedrero A, Ramirez SC, Kong D, Moody AR, Tyrrell PN**. 11 2019. Sample-Size Determination Methodologies for Machine Learning in Medical Imaging Research: A Systematic Review. Can Assoc Radiol journal = J l'Association canadienne des radiologistes 70:344–353. doi:10.1016/J.CARJ.2019.06.002.

57. **Vabalas A, Gowen E, Poliakoff E, Casson AJ**. 11 2019. Machine learning algorithm validation with a limited sample size. PLoS ONE 14. doi:10.1371/journal.pone.0224365.

58. **Illumina**. 2012. Data Processing of Nextera® Mate Pair Reads on Illumina Sequencing Platforms https://www.illumina.com/documents/products/technotes/technote_nextera_matepair_data_processing.pdf.

59. **Oksanen J, Blanchet FG, Friendly M, Kindt R, Legendre P, McGlinn D, Minchin PR, O'Hara RB, Simpson GL, Solymos P, Stevens MHH, Szoecs E, Wagner H**. 2020. vegan: Community Ecology Package. https://CRAN.R-project.org/package=vegan. R package version 2.5-7.

60. **Gloor GB, Macklaim JM, Pawlowsky-Glahn V, Egozcue JJ**. 11 2017. Microbiome Datasets Are Compositional: And This Is Not Optional. Front Microbiol 8:2224. doi:10.3389/fmicb.2017.02224.

61. **Aitchison J, Barceló-Vidal C, Martín-Fernández JA, Pawlowsky-Glahn V**. 2000. Logratio Analysis and Compositional Distance. Math Geol 32:271–275. doi:10.1023/A:1007529726302.

62. **Templ M, Hron K, Filzmoser P**. 7 2011. robCompositions: An R-package for Robust Statistical Analysis of Compositional Data. John Wiley & Sons, Ltd. doi:10.1002/9781119976462.ch25.

63. **Jones DL**. 2017. Fathom Toolbox for MATLAB: software for multivariate ecological and oceanographic data analysis https://www.usf.edu/marine-science/research/matlab-resources/index.aspx.

64. **Hoffmann H**. 2015. Violin Plot. MATLAB Central File Exchange https://www.mathworks.com/matlabcentral/fileexchange/45134-violin-plot.

Adapter and quality trimming

- Custom adapter trimming
- Trailing G removal
- TrimGalore
- PhiX filtering

Host filtering

- Database of 31 food ingredient and contaminant genomes
- Kraken with score threshold

Microbial genus profiling

- NCBI RefSeq genome database
- Kraken with score threshold
- RPM (Reads Per Million) threshold

Ordination and machine learning

- Clustering
- Contrastive PCA, MDS
- Differential abundance
- Explainable AI

Supported genera microbial composition

Beta diversity clustering by sample (Aitchison distance, outliers removed)

(a) Outside farm vs. baseline

(b) Antibiotic treated vs. baseline

(a) Outside farm vs. baseline

(b) Antibiotic treated vs. baseline