The overlooked dimension: soil depth governs microbial community assembly and enzymatic activity in extreme environments

Xin Jing[1], Aimée T. Classen[2], Daijiang Li[3,4], Litao Lin[5], Mingzhen Lu[6], Nathan J. Sanders[2], Yugang Wang[7,8,9], Wenting Feng[5*]

[1] State Key Laboratory of Herbage Improvement and Grassland Agro-Ecosystems, and College of Pastoral Agriculture Science and Technology, Lanzhou University, Lanzhou, 730000, Gansu, China

[2] Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, Michigan, USA

[3] Department of Biological Sciences, Louisiana State University, Baton Rouge, Louisiana, USA

[4] Center for Computation & Technology, Louisiana State University, Baton Rouge, Louisiana, USA

[5] Institute of Agricultural Resource and Regional Planning, Chinese Academy of Agricultural Sciences, Beijing, 100081, China

[6] Santa Fe Institute, Santa Fe, NM 87501, USA

[7] State Key Laboratory of Desert and Oasis Ecology, Xinjiang Institute of Ecology

and Geography, Chinese Academy of Sciences, Urumqi, Xinjiang, 830011, China

[8] Fukang Station of Desert Ecology, Chinese Academy of Sciences, Fukang,

Xinjiang, 831505, China

[9] University of Chinese Academy of Sciences, Beijing, 100049, China

[*] **Corresponding author**

Wenting Feng, Email: fengwenting@caas.cn

Institute of Agricultural Resources and Regional Planning, Chinese Academy of

Agricultural Sciences, Beijing, 100081, China

Abstract

Soil microbes have a major impact on ecosystem functions in extreme environments, such as deserts, where environmental factors such as salinity and soil moisture act as strong filters in shaping community structure. In fact, environmental filtering in extreme environments can occur at a small scale along soil depths, which is often overlooked, but it is important for an explicit understanding of how microbial community structure relates to microbial functions. Here, we leveraged a 200-km desert salinity gradient to evaluate microbial community structure-function relationships in extreme environments, and to further assess the prime ecological processes controlling the assembly of microbial communities and the activity of carbon, nitrogen, and phosphorus cycling-relevant enzymes along soil depths. We found that differences in soil bacterial and fungal community structure were not linked to differences in enzymatic activity, implying that soil microbial communities have a high degree of functional redundancy among soil depths. We further found that soil depth (at the scale of meters) was just as important as geographic distance (at the scale of kilometers) in shaping bacterial and fungal community structure, while the vertical variability in enzymatic activity along soil depth was largely attributed to the increase in heterogeneity of soil properties, such as soil texture, water content, and pH. Our results suggest that dispersal limitation is particularly important to soil microbial community assembly, while environmental filtering is important to enzymatic activity. Such a mismatch in ecological processes could have disproportionate influences on soil microbial community structure and enzymatic

activity, leading to uncoupled structure-function relationships. Our results further

suggest that soil depth governs the assembly and enzymatic activity of soil

microbial communities. This work highlights that soil conservation efforts should

consider soil depth as a key attribute in the face of ongoing desertification in arid

ecosystems.

Keywords: ecosystem functioning; extreme environments; microbial beta-biodiversity; soil microbiome; structure-function relationships; Taklamakan desert

1. Introduction

A major goal of biodiversity-ecosystem functioning (hereafter, BEF) research

has been to understand how community structure influences a wide range of

ecosystem functions, such as aboveground net primary productivity, soil carbon

fluxes, and nutrient cycling (Tilman et al., 2014). Many studies of biodiversity

experiment find positive relationships between biodiversity and ecosystem

functions, suggesting that enhanced biodiversity would lead to enriched

ecosystem functioning (van der Plas, 2019). However, these studies have been

largely carried out in constructed communities, in which community structure is

often randomly assembled from a species pool (Loreau et al., 2001; Schmid et

al., 2002; Jochum et al., 2020). As a result of this human-driven random

assembly process, a wide range of naturally occurring biotic and abiotic

ecological filters are excluded from the processes of community structure

assembly, such as disturbance, environmental heterogeneity or extremes, and

dispersal limitation (Sundqvist et al., 2013; Talbot et al., 2014; Mori et al., 2018).

This raises an important and unresolved question: would these positive BEF

relationships that we observe in constructed communities hold in natural settings

(e.g., natural grasslands) (Wardle, 2016; Manning et al., 2019; Schmid et al.,

2022)? Put it another way, can we safely extend our current understanding of

BEF relationships from constructed communities to natural communities (Hagan

et al., 2021), including belowground microbial communities (Bell et al., 2009;

Eisenhauer et al., 2019; Berlinches de Gea et al., 2022)?

Despite the growing recognition that microbial communities in soil are

important for numerous ecosystem processes (van der Heijden et al., 2008;

Delgado-Baquerizo et al., 2020), extending BEF theory to soil microbial

communities is a relatively new endeavor (Bell et al., 2009; Bardgett and van der

Putten, 2014; Peay et al., 2016; Berlinches de Gea et al., 2022). One of the

challenges is the hyper-diversity of microorganisms that inhabit the soil matrix

(Coleman and Whitman, 2005; Nielsen et al., 2015; Díaz and Malhi, 2022), which

makes it much difficult to uncover the relationships between biodiversity and

functions; there can be 100-9000 microbial species/strains in a small soil sample

(Bardgett and van der Putten, 2014) vs. 1~32 species in a constructed grassland

biodiversity experiment (Hector et al., 1999).

One strategy to address this challenge is to study BEF in extreme

ecosystems, such as desert, glaciers, and hypersaline habitats, where soil

microbial diversity is relatively low and environmental constraints are high (Shu

and Huang, 2022). Under extreme environmental conditions, biological

community structure, the characteristics of a community that can be described by

community diversity and/or composition, is generally simple, providing

opportunities to understand how an understandable amount of biodiversity

influences ecosystem functions (Wall and Virginia, 1999; Osborne et al., 2020;

Ertekin et al., 2021; Shu and Huang, 2022). Moreover, extreme environments

provide unique opportunities to predict how biodiversity might respond in an era

of environmental extremes and dramatic land-use change (Wall et al., 2015).

Environmental filters such as salinity, soil water, and nutrient availability can

shape microbial composition and function in extreme environments (e.g., desert),

but these drivers may change with soil depth (Rath et al., 2019a; Rath et al.,

2019b). In fact, prior work highlights that soil depth influences microbial

community structure, because important drivers of microbial diversity, such as

soil organic carbon, nutrient, pH, oxygen concentration, moisture, and salinity,

also vary with soil depth (Fierer et al., 2003; Powell et al., 2015; Stone et al.,

2015; Sun et al., 2020). However, we don't have as much of an idea of how the

vertical patterns in microbial community structure are intimately linked to their

functions, even though we know that soil depth is important for microbial

functions such as organic matter decomposition and nutrient cycling.

Here, we study the relationships between soil microbial community structure

and enzymatic activity relevant to soil carbon, nitrogen, and phosphorus cycling

in soils from the Taklamakan desert, the world's 2[nd] largest shifting sand desert;

a desert area that is roughly the size of Germany. We evaluate BEF relationships

across 10 field sites situated along a unique 200-km salinity gradient (the

horizontal gradient; Fig. 1a,b). At each site we also examine BEF relationships

across a one-meter soil profile (the vertical gradient; Fig. 1a,c), a new dimension

for BEF research. We asked whether and how differences in microbial

community structure are related to differences in enzymatic activity along these

two gradients (Fig. 1a,d). We measured soil bacterial and fungal community

structure based on pairwise measures of dissimilarities in community

composition between sample units (Anderson et al., 2011), potential soil

microbial enzymatic activity, and a range of environmental variables including soil

salinity, physical and chemical properties, and nutrient pools.

We hypothesize that differences in soil bacterial and fungal community

structure would be positively associated with spatial variability in potential

microbial enzymatic activity (H1). This hypothesis is based on the assumption

that extreme environments reduce community functional redundancy (Wall and

Virginia, 1999), leading to a strong relationship between microbial function and

community structure (Peay et al., 2016). However, it remains unclear if this

relationship is a positive one (Fig. 1d) (Schimel and Gulledge, 1998), as previous

studies have found positive, neutral and negative relationships. Studies that

focus on local-scale tend to report positive microbial structure-function

relationships (Mori et al., 2016), while studies that span large spatial scale tend

to report neutral or weakly positive relationships (Talbot et al., 2014; Martinez-

Almoyna et al., 2019; Jing et al., 2021; Jing et al., 2022).

We also hypothesize that soil depth is as important as horizontal geographic

distance in shaping the horizontal and vertical structure-function relationships,

but different factors of environmental filtering modify the structure-function

relationships across spatial dimensions (H2). Specifically, environmental

conditions such as differences in soil salinity would be the main environmental

filter driving the horizontal structure-function relationships (Rath et al., 2019a),

whereas the heterogeneity of soil properties (Powell et al., 2015) and nutrients

(Fierer et al., 2003) along soil depth profiles may drive vertical structure-function

relationships.

2. Materials and Methods

2.1. Study sites and data collection

We conducted this study in shrubland vegetation located along the Tarim

desert highway, which crosses the Tarim basin of Taklamakan desert

(37.1–41.8^{o} N, 82.6–84.3^{o} E) (Fig. S1). The Taklamakan desert is arid, with a

mean annual precipitation of 24.6 mm yr^{-1} and a mean annual potential

evaporation of 3639 mm yr^{-1}. The mean annual air temperature is 2.4°C, with the

coldest monthly mean temperature reaching –8.1°C in December and the

warmest monthly mean temperature reaching 28.2°C in July. Most of the desert

is bare land without vegetation cover, except for a few extremely drought- and

salt-tolerant shrub species, such as, *Tamarix taklamakanensis* and *Calligonum*

taklamakanensis. Soils in the desert are aeolian sandy soil. The top 0–20 cm of

soil is composed of 87.3% of sand, 12.4% silt, and 0.3% of clay (Zhang et al.,

2016). The soils have extremely low soil organic matter content (0.74 g kg^{-1}) and

soil total nitrogen (0.14 g kg^{-1}) in the top 0–10 cm soil depth (Li et al., 2015).

In 2005, the shrubland vegetation was established to prevent shifting sand

dunes from damaging the Tarim desert highway. After 12 years of vegetation

development the plant community has stabilized, with *Haloxylon ammodendron*

dominant the community and *Calligonum aborescens*, *Tamarix ramosissima*,

Nitraria tangutorum, *Populus euphratica* and *Elaeagnus angustifolia* as much

lower abundance. The aboveground plant community did not vary along the 200

km salinity gradient. Moreover, the aboveground biomass of the dominant

species *H. ammodendron* did not vary across the studied sites, likely leading to

an equivalent amount of carbon inputs into soil. The groundwater level in the

Taklamakan desert ranges from 3 to 5 m, and the groundwater is charged by

water from the Tianshan and Kunlun Mountains (Fig. S1). The groundwater was

pumped for drip irrigation to water shrubs in the vegetation. The pipes for drip

irrigation run along the rows of shrubs, and the vegetation was irrigated twice per

month from March to May and from September to October, and three times per

month from June to August with ~30 L of water per plant per irrigation event (Xu

et al., 2006).

182  We surveyed ten shrubland sites along an extreme gradient of irrigation

183 water salinity that varied from 3.2 to 38.0 g L^{-1} of dissolved salts (the normal

184 range of sea water salinity is between 33-37 g L^{-1} of dissolved salts) from the

185 south (site 34) to the north sampling site (site 1) (Figs. 1b and S1). We collected

186 irrigation water and soil samples in early October of 2017, which marks the end

187 of the growing season in the desert. At each study site, irrigation water was

188 collected from running water pumped from wells with three replicates at each

189 site. Irrigation water samples were stored in 500 ml plastic bottles, sealed, and

190 taken to the laboratory for water salinity analysis. We sampled soils using soil

191 cores (5 cm in diameter) at three randomly selected plots, which were located

192 within the two rows of shrubs and 30□50 cm away from a shrub. We collected

193 soils to the depth of 100 cm with sampling intervals of 20 cm, except that we

194 segmented the topsoils to 0□10 and 10□20 cm. We collected 180 soil samples

195 (10 sites × 6 soil layers × 3 replicates). Using plastic bags and coolers, a

196 subsample (~30 g) of each soil was kept cool and immediately returned to the

197 laboratory. We sieved all soils to 2 mm removing rocks and roots by hand. We

198 stored soils for microbial sequencing at □20 ^{o}C, and soils used for assays of soil

199 enzymatic activity at 4 ^{o}C. We air-dried the remaining soils to analyze soil

200 properties, including water content, texture, salinity, pH, total organic carbon,

201 total nitrogen, and available phosphorus.

202  We analyzed the salinity of irrigation water in the following ways. We

203 analyzed the concentration of Na^{+}, K^{+}, Ca^{2+}, and Mg^{2+} using a spectrometer

(iCAP 6300-ICP-OES) with a CID detector (Thermo Scientific, MA, USA). Using

titration, we analyzed the concentration of Cl^- with $AgNO_3$, SO_4^{2-} with

ethylenediaminetetraacetic acid (EDTA), and CO_3^{2-} and HCO_3^- with sulfuric acid

until all bicarbonate was neutralized. Next, we calculated irrigation water salinity

as the sum of Na^+, K^+, Ca^{2+}, Mg^{2+}, Cl^-, SO_4^{2-}, CO_3^{2-} and HCO_3^- concentrations.

We measured soil salinity and pH in a 1:5 suspension of soil to water. We

analyzed soil salinity using the methods described above for irrigation water

salinity. We analyzed soil pH using the Mettler Toledo FiveEasy FE28

(Greifensee, Switzerland). We measured soil gravimetric water content by oven

drying 10 g fresh soils at $105^{o}C$ for 24 h. We analyzed soil organic carbon using

the Walkley and Black method of oxidizing organic carbon with $K_2Cr_2O_7$-H_2SO_4

(Nelson and Sommers, 1983). We analyzed soil total nitrogen using an

AutoKjeldahl Unit model K370 (BUCHI, Flawil, Switzerland). We analyzed soil

available phosphorus using the sodium bicarbonate extraction with Mo-Sb Anti-

spectrophotometer method. We analyzed soil texture using the laser diffraction

method (Sympatec GmbH, System-PartikelTechnik, Clausthal-Zellerfeld,

Germany) after removing the organic material by 10% hydrogen peroxide and

removing carbonates by 0.2% hydrochloric acid.

To analyze the soil bacterial and fungal diversity, we extracted soil DNA

using the PowerSoil DNA extraction kit (Qiagen, Carlsbad, CA, USA) following

the manufacturer's protocol. We amplified the V4 region of 16S rRNA gene for

soil bacteria using the primer pairs 515F/806R and the ITS1 region for soil fungi

using the primer pairs ITS5-1737F/ITS2-2043R. We performed the PCR using

the BioRad S1000 thermal cycler (Bio-Rad Laboratory, CA, USA) after mixing 25

µl of 2x Premix Taq polymerase, 2 µl of 10 mM primer pairs, 3 µl of 20 ng µl^{-1}

DNA template and 20 µl nuclease-free water. Next, we ran the thermal cycling as

follows: initial denaturation at 94°C for 5 min, followed by 30 cycles of

denaturation at 94°C for 30 s, annealing at 52°C for 30 s, and elongation at 72°C

for 30 s, with a final step of 72°C for 10 mins. We repeated this PCR procedure

three times for each sample. We next mixed the PCR products using GeneTools

Analysis Software (Version 4.03.05.1, SynGene). The process was followed by

purifying the DNA products using the EZNA Gel extraction kit (Omega Bio-Tek,

GA, USA). Following the manufacturer's instruction, we generated sequencing

libraries using the NEBNext ® Ultra™ DNA Library Prep Kit for Illumina® (New

England Biolabs, MA, USA) and we sequenced the libraries on an Illumina

HiSeq2500 platform (Illumina PE250, CA, USA).

We processed raw DNA sequences using the Trimmomatic software (V0.33,

http://www.usadellab.org/cms/?page=trimmomatic). Specifically, we excluded the

raw DNA sequences by removing paired-end reads with N, read quality <20, and

read length <100 bp. Next, we allocated the paired-end reads to corresponding

soil samples using the Mothur pipeline (V1.35.1, http://www.mothur.org). We

removed barcode and primers to derive the paired-end clean reads, which were

merged by FLASH (V1.2.11, https://ccb.jhu.edu/software/FLASH). The minimum

length of overlap was 10 bp and the maximum ratio of number of mismatches to

overlap length was 0.1. Next, we assigned these clean tags to OTUs with 97%

similarity using the USEARCH software (V8.0.1517,

http://www.drive5.com/usearch). We used representative sequences for

taxonomic classification using RDP classifier according to Greengenes database

(http://greengenes.secondgenome.com) for bacteria and Unite database

(http://unite.ut.ee/index.php) for fungi in the QIIME pipeline (version 2;

http://qiime.org/scripts/assign_taxonomy.html).

To explore microbial function, we assayed the potential activity of four

hydrolytic enzymes (β-1, 4-glucosidase, BG; β-$_D$- cellobiohydrolase, CB; β-1, 4-

N-acetyl-glucosaminidase, NAG; alkaline phosphatase, ALP) using the 96-well

micro-plate and the fluorometric technique (German et al., 2011; Bell et al.,

2013). We selected these enzymes because their potential activities are

commonly measured in soils and are used in the acquisition of carbon (BG and

CB), nitrogen (NAG) and phosphorus (ALP) (Sinsabaugh et al., 2008) and can be

considered as proxies for fast ecosystem processes to define ecosystem

multifunctionality (Garland et al., 2021). Briefly, sieved fresh soil (1.5 g) was

suspended in 150 ml of 50 mM Tris buffer (pH = 8.0). Soil slurries (200 µl) of

each sample were transferred to eight wells of the 96-well micro-plate and mixed

with 50 µl of 200 µM standard fluorometric substrates of each enzyme. The

micro-plates were incubated for 2.5 h in dark at 25^{o}C. The quantity of

fluorescence was determined at 360 nm excitation and 460 nm emission in a 96-

well micro-plate reader (Biotek Synergy 2, Winooski, VT, USA). Multiple-point

calibration curve was used to calculate enzymatic activities (Bell et al., 2013).

The units were expressed as nmol g dry weight^{-1} h^{-1}.

To measure the spatial variability in soil microbial community structure

between assemblages, we used the Podani family of Sorensen indices (Podani

and Schmera, 2011; Legendre, 2014; Schmera et al., 2020), including the

Sorensen dissimilarity index (β_{sor}), species replacement (β_{repl}) and richness

difference (β_{rich}). The Sorensen dissimilarity index quantifies the total differences

in species composition between assemblages. It can be partitioned into two

components, species replacement and richness difference. The former quantifies

species in one assemblage that are replaced by different species in another

assemblage and the latter quantifies differences in the number of species

between assemblages. We used the following equations:

$$\beta_{sor} = \frac{b + c}{2a + b + c} \qquad \text{Eqn. 1}$$

$$\beta_{repl} = \frac{2 \times min\ (b, c)}{2a + b + c} \qquad \text{Eqn. 2}$$

$$\beta_{rich} = \frac{|b - c|}{2a + b + c} \qquad \text{Eqn. 3}$$

where *a* denotes the number of species present in both assemblages, *b* denotes

the number of species present in assemblage 1 but not in assemblage 2, and *c*

denotes the number of species present in assemblage 2 but not in assemblage

1.

We considered potential enzymatic activity of BG, CB, NAG and ALP as

proxy for microbial function (Burns et al., 2013; Talbot et al., 2014; Fanin et al.,

2022). We used the Euclidean distances as the measures of spatial variability in

microbial function between soil microbial assemblages (Mori et al., 2016;

Martinez-Almoyna et al., 2019; Jing et al., 2021; Jing et al., 2022) as follows:

$$D(x, y) = \sqrt{\sum_{i=1}^{n} (x_i - y_i)^2}$$ Eqn. 4

where $D(x, y)$ denotes the Euclidean distance, n is the number of enzymes ($n = $

4), x_i and y_i denote enzymatic activity in assemblage x and assemblage y,

respectively. Before calculating the Euclidean distance, we standardized all the

variables with z-score (Peters et al., 2019).

We calculated the Euclidean distances as the measures of geographic

distance and differences in soil depth, salinity, properties, and nutrient pools

(Table S1). Specifically, the geographic distance was calculated using the

geographic coordinates of sampling sites, and it does not account for differences

in soil depth. We calculated differences in soil depth using the mean depth of

each sampling layer, and it does not account for geographic distance. We

calculated differences in soil salinity using the concentrations of soil Na^+, K^+,

Ca^{2+}, Mg^{2+}, Cl^-, SO_4^{2-}, CO_3^{2-} and HCO_3^{2-}. Heterogeneity of soil properties was

calculated using soil clay content, sand content, silt content, pH, and moisture.

Differences in nutrient pools were calculated using the data of soil organic

carbon, soil total nitrogen, and available phosphorus (Jing et al., 2022). The

variables used for calculating the Euclidean distances of soil salinity, properties,

and nutrient pools were standardized with z-score.

Widespread in global drylands, soil salinity varies among sites (horizontally)

as well as between soil layers (vertically) within a site (Fig. 1a). Here, we

developed a dimension partitioning approach to quantify the impacts of vertical

and horizontal differences in soil salinity and other abiotic factors on the spatial

variability in the structure and activity of soil bacterial and fungal communities.

We applied this approach to assess the relative importance of pure horizontal,

pure vertical, and combined vertical and horizontal differences in soil depth,

salinity, properties, nutrient pools, and spatial geographic distance on soil

microbial community structure and function in a desert system, where a gradient

of soil salinity was created by 12-years of saline water irrigation (Fig. 1a-c).

Specifically, we calculated the pair-wise dissimilarity/distance matrix between

soil microbial assemblages for all the combinations of sampling site and soil layer

(n = 60). The lower triangle of the pair-wise dissimilarity/distance matrix was next

converted into a single vector, which resulted in 1770 observations (60 × 59 / 2).

We further partitioned the single vector into three spatial dimensions, including

the horizontal, vertical, and horizontal & vertical dimensions. Differences in

structure/function within a soil layer between sampling sites were defined as the

horizontal dimension that has 270 observations (10 × 9 / 2 × 6), between soil

layers within a sampling site were defined as the vertical dimension that has 150

observations (6 × 5 / 2 × 10), and between soil layers and between sampling

sites were defined as the horizontal & vertical dimension that has 1350

observations. The dimension partitioning was carried out for all the variables,

including soil microbial community structure and function, geographic distance,

depth difference, salinity difference, and soil heterogeneity.

2.2. Statistical analyses

To test how soil microbial community structure is related to enzymatic activity

across spatial dimensions of soil depth profiles, we inspected the strength and

direction of bivariate relationships between differences in soil microbial

community structure and differences in enzymatic activity using the Spearman's

rank correlation test. Next, we used path models and hierarchical variance

partitioning analysis to estimate the relative importance of environmental

variables on the spatial variability in soil microbial community structure-function

relationships.

First, we conducted the path analysis to estimate the relative importance of

dispersal limitation and environmental filtering on soil microbial community

structure (Fig. S2). In terms of soil bacterial and fungal community structure, we

considered geographic distance and differences in soil depth as proxies for

dispersal limitation (Powell et al., 2015) and differences in soil salinity, properties,

and nutrient pools as proxies for environmental filtering (Martiny et al., 2006; Jing

et al., 2022). Using the path analysis, we also estimated the relative importance

of environmental variables and soil bacterial and fungal community structure on

differences in microbial enzymatic activity at each spatial dimension. We used

the P value of χ^2 statistic test (P >0.05), comparative fit index (CFI >0.90), root

mean square error of approximation (RMSEA <0.05), and standardized root

mean square residual (SRMR <0.10) as measures of path model evaluation and

selection (Grace, 2020). Because a large sample size (n > 100) may lead to the

P value of χ^2 statistic test that was less than 0.05 (Grace, 2020), we only used

CFI, RMSEA, and SRMR to evaluate and select models for overall and horizontal

and vertical spatial dimensions. All path models fitted the data well (Table S2).

We used 9 999 bootstrapped sampling iterations for significance tests of the path

coefficients.

Second, we conducted hierarchical variance partitioning to estimate the

proportion of spatial variability in microbial community structure that was

independently explained by the individual effects of geographic distance,

differences in soil depth, soil salinity, soil properties, and soil nutrient pools. We

also included differences in microbial community structure, soil salinity,

properties, and nutrient pools to estimate the proportion of spatial variability in

enzymatic activity that independently explained by these biotic and abiotic

factors. All combinations of explanatory variables were fitted into regression

models following the hierarchical partitioning algorithm of Chevan and Sutherland

(1991). The independent effects of each explanatory variable were calculated

based on the goodness-of-fit measures (i.e., R^2) of regression models with the

target variables (microbial community structure or enzymatic activity) as the

dependent variable. We used 9 999 bootstrapped sampling iterations to calculate

the 95% confidence interval of the independent effects of each explanatory

variable.

All statistical analyses were performed in R version 3.6.1 (R Development

Core Team, 2019) using the packages 'lavaan' (Rosseel, 2012) and 'hier.part'

(Mac Nally and Walsh, 2004). Prior to the data analyses, we removed singleton

OTUs, which are present only once across all microbial assemblages, from soil

microbial community data and pooled all the data to the level of site × soil layer to

remove the non-independent influences of sampling replicates.

3. Results

3.1. Soil microbial community structure-function relationships

Overall, differences in soil bacterial and fungal community structure were

associated with differences in enzymatic activity, but these relationships were

weak (Spearman correlation coefficient $\rho = 0.15$, $P < 0.001$; Fig. 2a,b). The weak

associations of soil microbial community structure with enzymatic activity were

consistently observed across the horizontal and vertical dimensions of soil depth

profiles ($\rho < 0.16$ in all cases; Fig. S3). Moreover, differences in replacement and

richness of soil bacterial and fungal community structure were weakly associated

with differences in enzymatic activity ($\rho < 0.16$ in all cases; Figs. S4 and S5).

When we controlled for geographic distance and differences in soil depth,

salinity, properties, and nutrient pools, differences in enzymatic activity were not

explained by differences in soil microbial community structure (standardized path

coefficients β_{std} <0.13 in all spatial dimensions; Fig. 3 and Table S3). However,

these variables separately explained between 14% and 27% of the variance in

differences in enzymatic activity, between 29% and 39% of the variance in

differences in soil bacterial community structure, and between 27% and 33% of

the variance in differences in soil fungal community structure (Fig. 3a-d).

Specifically, we found that differences in soil bacterial and fungal community

structure were most strongly associated with geographic distance (Fig. 3a,b,d),

while the effects of differences in soil depth (at the scale of meters; Fig. 3c) along

the vertical dimension were just as important as the effects of geographic

distance (on the scale of kilometers; Fig. 3a,b,d) along the horizontal dimension

on soil bacterial and fungal community structure (Appendix A). In addition, the

effects of heterogeneity in soil properties on differences in enzymatic activity

along the vertical dimension were as important as those effects at the horizontal

dimension (Appendix A).

3.2. Factors mediating microbial community structure and enzymatic activity

Geographic distance and differences in soil depth had the largest

independent effects on differences in soil bacterial and fungal community

structure along soil depth profiles (Fig. 4a,b). Geographic distance explained

20.7-25.8% of variance in differences in soil bacterial community structure and

24.1-31.9% of variance in differences in soil fungal community structure (Fig.

4a,b). Along the vertical dimension, differences in soil depth explained 21.9%

(95% confidence interval = [12.6, 32.6]) of variance in differences in soil bacterial

community structure (Fig. 4a and Table S4) and 23.8% (95% confidence interval

= [12.6, 35.6]) of variance in differences in soil fungal community structure (Fig.

4b and Table S4). The heterogeneity of soil properties further explained 11.0%

(95% confidence interval = [4.6, 21.0]) and 5.9% (95% confidence interval = [1.3,

14.3]) of variance in differences in soil bacterial and fungal community structure,

respectively (Fig. 4a,b and Table S4).

The heterogeneity of soil properties had the largest independent effects on

differences in enzymatic activity at both horizontal and vertical dimensions (Fig.

4c). The heterogeneity of soil properties explained 8.4-17.2% of variance in

differences in enzymatic activity along the overall, horizontal, and horizontal &

vertical dimensions, and explained 11.9% of variance in differences in enzymatic

activity along the vertical dimension (95% confidence interval = [3.9, 23.3]) (Table

S4). Differences in soil bacterial and fungal community structure had the least

independent effects on differences in enzymatic activity (Fig. 4c).

4. Discussion

Much of our understanding of BEF relationships was developed in temperate

ecosystems with a focus on vascular plants. We may expect these relationships

to be even stronger in extreme environments with strong environmental gradients

(Forrester and Bauhus, 2016). Along strong salinity gradients, both horizontally

and vertically, we predict that soil salinity would define structure-function

relationships horizontally among sites, whereas the heterogeneity in soil

properties and nutrients would define the structure-function relationships

vertically with soil depth. Here, the most salient result of this study is that soil

microbial community structure was decoupled from enzymatic activity relevant to

carbon, nitrogen, and phosphorus cycling along both the horizontal and vertical

dimension of soil depth profiles. This result runs counter to previous work

exploring these relationships in microbial systems (Mori et al., 2016; Delgado-

Baquerizo et al., 2020) and does not support our first hypothesis that differences

in soil microbial community structure would be positively associated with spatial

variability in enzymatic activity in extreme environments. The results of spatial

dimension partitioning partially support our second hypothesis that soil depth is

as important as horizontal geographic distance in shaping bacterial and fungal

community structure. However, our results show that the heterogeneity of soil

properties (i.e., soil texture, pH, and water) rather than soil salinity or nutrient

pools, is most strongly associated with differences in enzymatic activity across all

spatial dimensions of soil depth profiles. Our findings suggest that soil depth

governs the assembly of soil microbial community structure and the activity of

carbon, nitrogen, and phosphorus cycling enzymes. Soil depth profile thus can

serve as a test bed for predicting changes in soil microbial community structure

and enzymatic activity in real-world extreme environments.

4.1. Uncoupled microbial community structure-function relationships

Differences in soil microbial community structure did not lead to differences in

enzymatic activity in the studied hyper-saline environments. The uncoupled

relationships between soil microbial structure and function were further verified

across the horizontal and vertical dimensions of soil depth profiles. Our findings,

thus, support the idea of functional redundancy (Coleman and Whitman, 2005;

Peay et al., 2016), implying that soil microbial communities – even ones that are

in extreme environments – differ, but function in the same manner. As a result,

the losses/gains of certain species may have limited effects on the spatial

variability in enzymatic activity. This means that maintaining functionally

redundant communities is crucial to safeguard microbial function against the

negative influences of extreme environmental changes (Allison and Martiny,

2008), and thus could promote ecosystem stability and the provisioning of

multiple ecosystem services.

There are two explanations for the uncoupled structure-function relationships.

First, soil bacterial and fungal communities have likely adapted to the harsh

desert conditions, where abiotic stress is high (Jansson and Hofmockel, 2020).

As soil becomes saltier and drier, soil microorganisms become inactive because

soil carbon and nutrients availability declines. In addition, soil microorganisms

tend to minimize their activity, even becoming dormant, to conserve energy

under harsh conditions. In either case, we would observe the uncoupled

relationships between soil microbial community structure and enzymatic activity.

Second, the uncoupled structure-function relationship is likely due to the

mismatch of dominant driving factors responsible for microbial community

structure and enzymatic activity. Our results show that geographic distance and

differences in soil depth have the most strongly independent effects on microbial

community structure, while the heterogeneity in soil properties has the most

strongly independent effects on enzymatic activity. Such a mismatch in driving

factors could have disproportionate influences on soil microbial community

structure and enzymatic activity, leading to uncoupled structure-function

relationships.

4.2. Soil depth profile as a new frontier for BEF study

We found that the spatial variability in soil microbial community structure was

strongly associated with geographic distance and differences in soil depth. This

result suggests that soil bacterial and fungal communities show dissimilar

community structure with increased spatial distance both horizontally and

vertically along soil depth profiles. Moreover, our work indicates that soil depth is

as important as geographic distance in shaping soil bacterial and fungal

community structure (Fig. 4a,b). The importance of soil depth is supported by an

earlier study showing that the random assembly processes of soil microbial

communities (e.g., dispersal limitation) intensified with soil depth because soil

microorganisms disperse passively (Powell et al., 2015). The observation that

soil depth and geographic distance are equally important emerges only when the

overall spatial variability in microbial structure is partitioned into the pure vertical

and horizontal dimensions (Fig. 4a,b). Without the spatial dimension partitioning,

we would underestimate the contribution of soil depth to the assembly processes

of soil microbial communities.

Overall, our work highlights that dispersal limitation in soil microbes is

particularly important to the assembly of soil microbial communities in the vertical

and horizontal dimensions of soil depth profiles in extreme environments. To

maintain the essential and irreplaceable ecosystem functions and services

provided by belowground biodiversity in the extreme conditions or severely

disturbed conditions, a workable plan is to preserve habitats along both

horizontal and vertical habitats along soil depth profiles.

Our study further advances the understanding that the spatial variability in

soil enzymatic activity is best explained by the heterogeneity in soil texture, pH,

and water, rather than differences in soil salinity and nutrient pools across soil

depth. In the extreme environments, environmental filters (e.g., salt stress) and

resource levels (e.g., soil organic matter and microbial biomass) are expected to

be important drivers of spatial variability in soil microbial community structure and

function (Martiny et al., 2006; Rath and Rousk, 2015; Rath et al., 2019a; Rath et

al., 2019b). By contrast, our results suggest that differences in soil salinity and

nutrient pools do not explain the activity of carbon, nitrogen, and phosphorus

cycling enzymes. One possible explanation is that soil salinity is too high in our

study system and the first priority of soil microbes is to survive rather than to

thrive in the extreme saline environments, which leads soil microbes to allocate

few resources to produce enzymes (Rath and Rousk, 2015). As a result,

changes in soil salinity and nutrient pools did not alter enzymatic activity because

soil salinity and resources are not limiting factors of enzymatic activity. However,

our result indicates that reduced heterogeneity in soil properties, which could be

induced by land degradation, soil erosion, and desertification, might result in the

loss of soil microbial activity and soil functionality, soil fertility, and plant growth.

Given that the variability in soil physical and chemical properties is created by

human activities and/or by nature ubiquitously, this work highlights the

significance of analyzing the overlooked dimension of microbial structure-function

relationships at the vertical dimension along soil depth profiles, which represent a

valuable step towards soil conservation efforts in dealing with ongoing

desertification in arid ecosystems.

In recent years, microbial ecologists have focused primarily on surface

communities because they have the highest biomass density and measured

nutrient cycling activities. This study is unique in that it provides a new approach

to assess soil microbial community structure-function relationships both

horizontally and vertically along soil depths with similar range of salinity variability

in one of the extreme biomes, i.e., the Taklamakan desert. However, large-scale

studies remain the key to verify whether these findings of this study hold true

across other biomes with wide range of horizontal and vertical environmental

gradients that are created either by human activities or by nature. Further, it is

important to note that the measurements of four hydrolytic enzyme activities

capture only a few microbial potential functions that are relevant to carbon,

nitrogen, and phosphorus cycling. Future work that uses shotgun metagenomics

could quantify broad shifts in microbial community functionality, e.g., those

functions related to organotrophs and perhaps autotrophs (both by phototrophs

and chemolithotrophs such as hydrogen oxidizers), and thus advance the

understanding of structure-function relation of microbial communities in xeric or

arid soils.

5. Conclusion

This study advances the understanding of biodiversity-functioning

relationships into naturally assembled soil microbial communities in extreme

environments through partitioning soil depth profiles into the horizontal and

vertical dimensions. We demonstrate that soil microbial community structure is

decoupled from enzymatic activity along both the horizontal and vertical

dimensions of soil depth profiles. Moreover, we show that soil depth is as

important as horizontal geographic/environmental gradient in shaping soil

microbial community structure and enzymatic activity. These findings highlight

the need to analyze the new dimension of biodiversity and ecosystem

functioning, i.e., the vertical dimension. Such information is crucial to predict the

influences of global change drivers on real-world naturally assembled community

structure-function relationships across horizontal and vertical spatial scales.

Moreover, this study highlights that considering increasing desertification in arid

ecosystems, soil conservation efforts should take soil depth into consideration as

a critical factor governing the assembly of microbial communities and the activity

of enzymes.

Acknowledgments

This work was supported by National Natural Science Foundation of China

(42077023, U1803342), Intergovernmental International Cooperation on

Scientific and Technological Innovation as Part of National Key Research and

Development Program (2021YFE0114500), Agricultural Science and Technology

Innovation Program of Chinese Academy of Agricultural Sciences, and K. C.

Wong Education Foundation (GJTD-2020-14).

Author Contributions: XJ, WF and YW designed the study. YW and WF

conducted the field- and lab-work. XJ analyzed the data. XJ and WF wrote the

first draft of the manuscript. All authors contributed to the conceptualization,

review and editing of the manuscript.

Competing Interest Statement: The authors declare no competing interests.

Data availability: Sequence data were deposited in the European Nucleotide

Archive with accession number PRJEB52395

(http://www.ebi.ac.uk/ena/data/view/OW744067-OW763241) for soil bacteria,

and with accession number PRJEB46878

(http://www.ebi.ac.uk/ena/data/view/OU495976-OU501470) for soil fungi. Data

and code will be publicly available in the Zenodo repository prior to publication.

References

Allison, S.D., Martiny, J.B., 2008. Resistance, resilience, and redundancy in microbial communities. Proceedings of the National Academy of Sciences, USA 105, 11512-11519.

Anderson, M.J., Crist, T.O., Chase, J.M., Vellend, M., Inouye, B.D., Freestone, A.L., Sanders, N.J., Cornell, H.V., Comita, L.S., Davies, K.F., 2011. Navigating the multiple meanings of β diversity: a roadmap for the practicing ecologist. Ecology Letters 14, 19-28.

Bardgett, R.D., van der Putten, W.H., 2014. Belowground biodiversity and ecosystem functioning. Nature 515, 505-511.

Bell, C.W., Fricks, B.E., Rocca, J.D., Steinweg, J.M., McMahon, S.K., Wallenstein, M.D., 2013. High-throughput fluorometric measurement of potential soil extracellular enzyme activities. JoVE, e50961.

Bell, T., Gessner, M.O., Griffiths, R.I., McLaren, J.R., Morin, P.J., Van Der Heijden, M., van der Putten, W.H., 2009. Microbial biodiversity and ecosystem functioning under controlled conditions and in the wild, In: Naeem, S., E, B.D., Hector, A., Loreau, M., Perrings, C. (Eds.), Biodiversity, ecosystem functioning, human wellbeing: an ecological economic perspective. Oxford University Press, New York, USA, pp. 121-133.

Berlinches de Gea, A., Hautier, Y., Geisen, S., 2022. Interactive effects of global change drivers as determinants of the link between soil biodiversity and ecosystem functioning. Global Change Biology 00, 1-12.

Burns, R.G., DeForest, J.L., Marxsen, J., Sinsabaugh, R.L., Stromberger, M.E., Wallenstein, M.D., Weintraub, M.N., Zoppini, A., 2013. Soil enzymes in a changing environment: Current knowledge and future directions. Soil Biology and Biochemistry 58, 216-234.

Chevan, A., Sutherland, M., 1991. Hierarchical partitioning. The American Statistician 45, 90-96.

Coleman, D.C., Whitman, W.B., 2005. Linking species richness, biodiversity and ecosystem function in soil systems. Pedobiologia 49, 479-497.

Delgado-Baquerizo, M., Reich, P.B., Trivedi, C., Eldridge, D.J., Abades, S., Alfaro, F.D., Bastida, F., Berhe, A.A., Cutler, N.A., Gallardo, A., Garcia-Velazquez, L., Hart, S.C., Hayes, P.E., He, J.Z., Hseu, Z.Y., Hu, H.W., Kirchmair, M., Neuhauser, S., Perez, C.A., Reed, S.C., Santos, F., Sullivan, B.W., Trivedi, P., Wang, J.T., Weber-Grullon, L., Williams, M.A., Singh, B.K., 2020. Multiple elements of soil biodiversity drive ecosystem functions across biomes. Nature Ecology & Evolution 4, 210-220.

Díaz, S., Malhi, Y., 2022. Biodiversity: Concepts, patterns, trends, and perspectives. Annual Review of Environment and Resources 47, 31-63.

Eisenhauer, N., Schielzeth, H., Barnes, A.D., Barry, K.E., Bonn, A., Brose, U., Bruelheide, H., Buchmann, N., Buscot, F., Ebeling, A., Ferlian, O., Freschet, G.T., Giling, D.P., Hättenschwiler, S., Hillebrand, H., Hines, J., Isbell, F., Koller-France, E., König-Ries, B., de Kroon, H., Meyer, S.T., Milcu, A., Müller, J., Nock, C.A., Petermann, J.S., Roscher, C., Scherber, C., Scherer-Lorenzen, M., Schmid, B., Schnitzer, S.A., Schuldt, A., Tscharntke, T., Türke, M., van Dam, N.M., van der Plas, F., Vogel, A., Wagg, C., Wardle, D.A., Weigelt, A., Weisser, W.W., Wirth, C., Jochum, M., 2019. A multitrophic perspective on biodiversity–ecosystem functioning research, In: Eisenhauer, N., Bohan, D.A., Dumbrell, A.J. (Eds.), Advances in Ecological Research. Academic Press, pp. 1-54.

Ertekin, E., Meslier, V., Browning, A., Treadgold, J., DiRuggiero, J., 2021. Rock structure drives the taxonomic and functional diversity of endolithic microbial communities in extreme environments. Environmental Microbiology 23, 3937-3956.

Fanin, N., Mooshammer, M., Sauvadet, M., Meng, C., Alvarez, G., Bernard, L., Bertrand, I., Blagodatskaya, E., Bon, L., Fontaine, S., Niu, S.L., Lashermes, G., Maxwell, T.L., Weintraub, M.N., Wingate, L., Moorhead, D., Nottingham, A.T., 2022. Soil enzymes in response to climate warming: Mechanisms and feedbacks. Functional Ecology 36, 1378-1395.

Fierer, N., Schimel, J.P., Holden, P.A., 2003. Variations in microbial community composition through two soil depth profiles. Soil Biology and Biochemistry 35, 167-176.

Forrester, D.I., Bauhus, J., 2016. A review of processes behind diversity-productivity relationships in forests. Current Forestry Reports 2, 45-61.

Garland, G., Banerjee, S., Edlinger, A., Oliveira, E.M., Herzog, C., Wittwer, R., Philippot, L., Maestre, F.T., van der Heijden, M.G.A., 2021. A closer look at the functions behind ecosystem multifunctionality: A review. Journal of Ecology 109, 600-613.

German, D.P., Weintraub, M.N., Grandy, A.S., Lauber, C.L., Rinkes, Z.L., Allison, S.D., 2011. Optimization of hydrolytic and oxidative enzyme methods for ecosystem studies. Soil Biology and Biochemistry 43, 1387-1397.

Grace, J., 2020. A 'Weight of Evidence' approach to evaluating structural equation models. One Ecosystem 5, e50452.

Hagan, J.G., Vanschoenwinkel, B., Gamfeldt, L., 2021. We should not necessarily expect positive relationships between biodiversity and ecosystem functioning in observational field data. Ecology Letters 24, 2537-2548.

Hector, A., Schmid, B., Beierkuhnlein, C., Caldeira, M.C., Diemer, M., Dimitrakopoulos, P.G., Finn, J.A., Freitas, H., Giller, P.S., Good, J., Harris, R., Hogberg, P., Huss-Danell, K., Joshi, J., Jumpponen, A., Korner, C., Leadley, P.W., Loreau, M., Minns, A., Mulder, C.P., O'Donovan, G., Otway, S.J., Pereira, J.S., Prinz, A., Read, D.J., Scherer-Lorenzen, M., Schulze, E.D., Siamantziouras, A.S.D., Spehn, E.M., Terry, A.C., Troumbis, A.Y., Woodward, F.I., Yachi, S., Lawton, J.H., 1999. Plant diversity and productivity experiments in European grasslands. Science 286, 1123-1127.

Jansson, J.K., Hofmockel, K.S., 2020. Soil microbiomes and climate change. Nature
Reviews Microbiology 18, 35-46.

Jing, X., Prager, C.M., Borer, E.T., Gotelli, N.J., Gruner, D.S., He, J.-S., Kirkman, K.,
MacDougall, A.S., McCulley, R.L., Prober, S.M., Seabloom, E.W., Stevens, C.J.,
Classen, A.T., Sanders, N.J., 2021. Spatial turnover of multiple ecosystem functions is
more associated with plant than soil microbial β-diversity. Ecosphere 12, e03644.

Jing, X., Prager, C.M., Chen, L., Chu, H., Gotelli, N.J., He, J.S., Shi, Y., Yang, T., Zhu,
B., Classen, A.T., Sanders, N.J., 2022. The influence of aboveground and belowground
species composition on spatial turnover in nutrient pools in alpine grasslands. Global
Ecology and Biogeography 31, 486-500.

Jochum, M., Fischer, M., Isbell, F., Roscher, C., van der Plas, F., Boch, S., Boenisch,
G., Buchmann, N., Catford, J.A., Cavender-Bares, J., Ebeling, A., Eisenhauer, N.,
Gleixner, G., Holzel, N., Kattge, J., Klaus, V.H., Kleinebecker, T., Lange, M., Le Provost,
G., Meyer, S.T., Molina-Venegas, R., Mommer, L., Oelmann, Y., Penone, C., Prati, D.,
Reich, P.B., Rindisbacher, A., Schafer, D., Scheu, S., Schmid, B., Tilman, D.,
Tscharntke, T., Vogel, A., Wagg, C., Weigelt, A., Weisser, W.W., Wilcke, W., Manning,
P., 2020. The results of biodiversity-ecosystem functioning experiments are realistic.
Nature Ecology & Evolution 4, 1485-1494.

Legendre, P., 2014. Interpreting the replacement and richness difference components of
beta diversity. Global Ecology and Biogeography 23, 1324-1334.

Li, C., Lei, J., Zhao, Y., Xu, X., Li, S., 2015. Effect of saline water irrigation on soil
development and plant growth in the Taklimakan Desert Highway shelterbelt. Soil &
Tillage Research 146, 99-107.

Loreau, M., Naeem, S., Inchausti, P., Bengtsson, J., Grime, J.P., Hector, A., Hooper,
D.U., Huston, M.A., Raffaelli, D., Schmid, B., Tilman, D., Wardle, D.A., 2001. Biodiversity
and ecosystem functioning: Current knowledge and future challenges. Science 294, 804-
808.

Mac Nally, R., Walsh, C.J., 2004. Hierarchical partitioning public-domain software.
Biodiversity and Conservation 13, 659.

Manning, P., Loos, J., Barnes, A.D., Batáry, P., Bianchi, F.J.J.A., Buchmann, N., De
Deyn, G.B., Ebeling, A., Eisenhauer, N., Fischer, M., Fründ, J., Grass, I., Isselstein, J.,
Jochum, M., Klein, A.M., Klingenberg, E.O.F., Landis, D.A., Lepš, J., Lindborg, R.,
Meyer, S.T., Temperton, V.M., Westphal, C., Tscharntke, T., 2019. Transferring
biodiversity-ecosystem function research to the management of 'real-world' ecosystems,
Mechanisms underlying the relationship between biodiversity and ecosystem function.
Elsevier, pp. 323-356.

Martinez-Almoyna, C., Thuiller, W., Chalmandrier, L., Ohlmann, M., Foulquier, A.,
Clément, J.C., Zinger, L., Münkemüller, T., 2019. Multi-trophic β-diversity mediates the
effect of environmental gradients on the turnover of multiple ecosystem functions.
Functional Ecology 33, 2053-2064.

Martiny, J.B., Bohannan, B.J., Brown, J.H., Colwell, R.K., Fuhrman, J.A., Green, J.L.,
Horner-Devine, M.C., Kane, M., Krumins, J.A., Kuske, C.R., Morin, P.J., Naeem, S.,
Ovreas, L., Reysenbach, A.L., Smith, V.H., Staley, J.T., 2006. Microbial biogeography:
putting microorganisms on the map. Nature Review Microbiology 4, 102-112.

Mori, A.S., Isbell, F., Fujii, S., Makoto, K., Matsuoka, S., Osono, T., 2016. Low
multifunctional redundancy of soil fungal diversity at multiple scales. Ecology Letters 19,
249-259.

Mori, A.S., Isbell, F., Seidl, R., 2018. β-Diversity, community assembly, and ecosystem
functioning. Trends in Ecology & Evolution 33, 549-564.

Nelson, D.W., Sommers, L.E., 1983. Total Carbon, Organic Carbon, and Organic Matter,
In: Page, A.L. (Ed.), Methods of Soil Analysis. Part 2. Chemical and microbiological
properties. ASA and SSSA, Madison, WI, USA, pp. 539-579.

Nielsen, U.N., Wall, D.H., Six, J., 2015. Soil biodiversity and the environment. Annual
Review of Environment and Resources 40, 63-90.

Osborne, P., Hall, L.J., Kronfeld-Schor, N., Thybert, D., Haerty, W., 2020. A rather dry
subject; investigating the study of arid-associated microbial communities. Environ
Microbiome 15, 20.

Peay, K.G., Kennedy, P.G., Talbot, J.M., 2016. Dimensions of biodiversity in the Earth
mycobiome. Nature Reviews Microbiology 14, 434-447.

Peters, M.K., Hemp, A., Appelhans, T., Becker, J.N., Behler, C., Classen, A., Detsch, F.,
Ensslin, A., Ferger, S.W., Frederiksen, S.B., Gebert, F., Gerschlauer, F., Gütlein, A.,
Helbig-Bonitz, M., Hemp, C., Kindeketa, W.J., Kühnel, A., Mayr, A.V., Mwangomo, E.,
Ngereza, C., Njovu, H.K., Otte, I., Pabst, H., Renner, M., Röder, J., Rutten, G.,
Schellenberger Costa, D., Sierra-Cornejo, N., Vollstädt, M.G.R., Dulle, H.I., Eardley,
C.D., Howell, K.M., Keller, A., Peters, R.S., Ssymank, A., Kakengi, V., Zhang, J.,
Bogner, C., Böhning-Gaese, K., Brandl, R., Hertel, D., Huwe, B., Kiese, R., Kleyer, M.,
Kuzyakov, Y., Nauss, T., Schleuning, M., Tschapka, M., Fischer, M., Steffan-Dewenter,
I., 2019. Climate–land-use interactions shape tropical mountain biodiversity and
ecosystem functions. Nature 568, 88-92.

Podani, J., Schmera, D., 2011. A new conceptual and methodological framework for
exploring and explaining pattern in presence–absence data. Oikos 120, 1625-1638.

Powell, J.R., Karunaratne, S., Campbell, C.D., Yao, H., Robinson, L., Singh, B.K., 2015.
Deterministic processes vary during community assembly for ecologically dissimilar taxa.
Nature Communications 6, 1-10.

R Development Core Team, 2019. R: A language and environment for statistical
computing. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-
project.org/.

Rath, K.M., Fierer, N., Murphy, D.V., Rousk, J., 2019a. Linking bacterial community
composition to soil salinity along environmental gradients. The ISME Journal 13, 836-
846.

Rath, K.M., Murphy, D.N., Rousk, J., 2019b. The microbial community size, structure,
and process rates along natural gradients of soil salinity. Soil Biology and Biochemistry
138, 107607.

Rath, K.M., Rousk, J., 2015. Salt effects on the soil microbial decomposer community
and their role in organic carbon cycling: A review. Soil Biology and Biochemistry 81, 108-
123.

Rosseel, Y., 2012. lavaan: An R Package for Structural Equation Modeling. Journal of
Statistical Software 48, 1-36.

Schimel, J.P., Gulledge, J., 1998. Microbial community structure and global trace gases.
Global Change Biology 4, 745-758.

Schmera, D., Podani, J., Legendre, P., 2020. What do beta diversity components reveal
from presence-absence community data? Let us connect every indicator to an
indicandum! Ecological Indicators 117.

Schmid, B., Hector, A., Huston, M., Inchausti, P., Nijs, I., Leadley, P., Tilman, D., 2002.
The design and analysis of biodiversity experiments, In: Loreau, M., Shahid, N., Pablo, I.
(Eds.), Biodiversity and Ecosystem Functioning: Synthesis and Perspectives. Oxford
University Press, Oxford, pp. 61-78.

Schmid, B., Schmitz, M., Rzanny, M., Scherer-Lorenzen, M., Mwangi, P.N., Weisser,
W.W., Hector, A., Schmid, R., Flynn, D.F.B., 2022. Removing subordinate species in a
biodiversity experiment to mimic observational field studies. Grassland Research 1, 53-
62.

Shu, W.S., Huang, L.N., 2022. Microbial diversity in extreme environments. Nature
Reviews Microbiology 20, 219-235.

Sinsabaugh, R.L., Lauber, C.L., Weintraub, M.N., Ahmed, B., Allison, S.D., Crenshaw,
C., Contosta, A.R., Cusack, D., Frey, S., Gallo, M.E., Gartner, T.B., Hobbie, S.E.,
Holland, K., Keeler, B.L., Powers, J.S., Stursova, M., Takacs-Vesbach, C., Waldrop,
M.P., Wallenstein, M.D., Zak, D.R., Zeglin, L.H., 2008. Stoichiometry of soil enzyme
activity at global scale. Ecology Letters 11, 1252-1264.

Stone, M.M., Kan, J., Plante, A.F., 2015. Parent material and vegetation influence
bacterial community structure and nitrogen functional genes along deep tropical soil
profiles at the Luquillo Critical Zone Observatory. Soil Biology and Biochemistry 80, 273-
282.

Sun, T., Wang, Y., Hui, D., Jing, X., Feng, W., 2020. Soil properties rather than climate
and ecosystem type control the vertical variations of soil organic carbon, microbial
carbon, and microbial quotient. Soil Biology and Biochemistry 148, 107905.

Sundqvist, M.K., Sanders, N.J., Wardle, D.A., 2013. Community and ecosystem
responses to elevational gradients: Processes, mechanisms, and insights for global
change. Annual Review of Ecology, Evolution, and Systematics 44, 261-280.

Talbot, J.M., Bruns, T.D., Taylor, J.W., Smith, D.P., Branco, S., Glassman, S.I.,
Erlandson, S., Vilgalys, R., Liao, H.-L., Smith, M.E., Peay, K.G., 2014. Endemism and
functional convergence across the North American soil mycobiome. Proceedings of the
National Academy of Sciences, USA 111, 6341-6346.

Tilman, D., Isbell, F., Cowles, J.M., 2014. Biodiversity and ecosystem functioning.
Annual Review of Ecology, Evolution, and Systematics 45, 471-493.

van der Heijden, M.G.A., Bardgett, R.D., van Straalen, N.M., 2008. The unseen majority:
soil microbes as drivers of plant diversity and productivity in terrestrial ecosystems.
Ecology Letters 11, 296-310.

van der Plas, F., 2019. Biodiversity and ecosystem functioning in naturally assembled
communities. Biological Reviews of the Cambridge Philosophical Society 94, 1220-1245.

Wall, D.H., Nielsen, U.N., Six, J., 2015. Soil biodiversity and human health. Nature 528,
69-76.

Wall, D.H., Virginia, R.A., 1999. Controls on soil biodiversity: insights from extreme
environments. Applied Soil Ecology 13, 137-150.

Wardle, D.A., 2016. Do experiments exploring plant diversity–ecosystem functioning
relationships inform how biodiversity loss impacts natural ecosystems? Journal of
Vegetation Science 27, 646-653.

Xu, X., Li, B., Wang, X., 2006. Progress in study on irrigation practice with saline
groundwater on sandlands of Taklimakan Desert Hinterland. Chinese Science Bulletin
51, 161-166.

Zhang, J., Xu, X., Li, S., Zhao, Y., Zhang, A., Zhang, T., Jiang, R., 2016. Is the
Taklimakan desert highway shelterbelt sustainable to long-term drip irrigation with high
saline groundwater? Plos One 11, e0164106.

Figure legend

Fig. 1 A conceptual framework for testing the relationships between community

structure and function along the horizontal and vertical dimensions of soil depth

profiles

Fig. 2 Bivariate relationships between differences in soil microbial community

structure and differences in enzymatic activity

Fig. 3 Uncoupled microbial community structure and function relationships along

soil depth profiles

Fig. 4 Independent effects of explanatory variables on differences in soil

microbial community structure and enzymatic activity

817
818

Fig. 1 A conceptual framework for testing the relationships between community structure and function along the horizontal and vertical dimensions of soil depth profiles. a) Heatmap illustrating changes in soil salinity along the horizontal and vertical dimension of soil depth profiles. Codes of the sampling sites are shown at the x-axis, and the average sampling depth for each soil layer is shown at the y-axis. b) The variability in irrigation water salinity along the horizontal gradient (geographic distance). Sampling sites are given next to the points. Site 1 with the highest water salinity is selected as the reference site for calculating the geographic distance (km). c) Impacts of irrigation water salinity on soil salinity across soil layers of the vertical dimension.

Linear mixed-effects models are used for significance tests. Solid lines denote

significant effects of irrigation water salinity on soil salinity and dashed lines

denote insignificant effects. d) If there is low functional redundancy in extreme

environments, we predict that community structure is positively associated with

microbial function. The levels of soil/water salinity in a-c) are represented by

colors from dark-blue (high soil/water salinity) to light-blue (low soil/water

salinity).

836
837

Fig. 2 Bivariate relationships between differences in soil microbial

community structure and differences in enzymatic activity. a) bacterial

community structure-enzymatic activity relationships; b) fungal community

structure-enzymatic activity relationships. Lines denote the loess fitted curves

showing the direction of the bivariate relationships. Shaded areas denote the

95% confidence interval. Spearman correlation coefficients are used to show the

strength of the bivariate relationships. Boxplots with median, quartiles, and

extreme values denote the data distributions of differences in community

structure (top) and differences in enzymatic activity (right). Different letters

denote significant differences among spatial dimensions.

Fig. 3 Uncoupled microbial community structure and function relationships along soil depth profiles. Four path models are shown based on spatial dimensions, including a) all dimensions; b) pure horizontal dimension; c) pure vertical dimension; d) combined horizontal & vertical dimension. Arrow width is proportional to the standardized path coefficients. Black arrows denote positive path coefficients and red arrows denote negative path coefficients. Bootstraps are used for the significance test of path coefficients (see Table S3 for details). R^2 denote the variance of community structure and enzymatic activity explained by environmental factors, including geographic distance (GeoDist), differences in

859   soil depth (DepDiff), differences in soil salinity (SalDiff), heterogeneity of soil

860   properties (HeteroProp), and differences in nutrient pools (NutPool). BacStruc,

861   FunStruc, and EnzAct denote differences in bacterial community structure, fungal

862   community structure, and enzymatic activity, respectively.

Fig. 4 Independent effects of explanatory variables on differences in soil microbial community structure and enzymatic activity at each spatial dimension. a) differences in bacterial community structure; b) differences in fungal community structure; c) differences in enzymatic activity. Bars represent the pure variance explained by the explanatory variables, and the values are shown along the bars. Points with 95% confidence interval denote the variability of explained variance for each explanatory variable (see Table S4 for details).