Soil depth governs microbial community assembly and enzymatic activity in extreme environments

Running title: **microbial community structure-function**

Xin Jing[1], Aimée T. Classen[2], Daijiang Li[3,4], Litao Lin[5], Mingzhen Lu[6], Nathan J. Sanders[2], Yugang Wang[7,8,9], Wenting Feng[10*]

[1] State Key Laboratory of Herbage Improvement and Grassland Agro-Ecosystems, and College of Pastoral Agriculture Science and Technology, Lanzhou University, Lanzhou, 730000, Gansu, China

[2] Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, Michigan, USA

[3] Department of Biological Sciences, Louisiana State University, Baton Rouge, Louisiana, USA

[4] Center for Computation & Technology, Louisiana State University, Baton Rouge, Louisiana, USA

[5] State Key Laboratory of Efficient Utilization of Arid and Semi-arid Arable Land in Northern China, the Institute of Agricultural Resource and Regional Planning, Chinese Academy of Agricultural Sciences, Beijing, 100081, China

[6] Santa Fe Institute, Santa Fe, NM 87501, USA

[7] State Key Laboratory of Desert and Oasis Ecology, Xinjiang Institute of Ecology

and Geography, Chinese Academy of Sciences, Urumqi, Xinjiang, 830011, China

[8] Fukang Station of Desert Ecology, Chinese Academy of Sciences, Fukang,

Xinjiang, 831505, China

[9] University of Chinese Academy of Sciences, Beijing, 100049, China

[10] School of Grassland Science, Beijing Forestry University, Beijing, China

[*] **Corresponding author**

Wenting Feng, Email: fengwenting@caas.cn

School of Grassland Science, Beijing Forestry University, Beijing, China

Acknowledgments

This work was supported by the National Key Research and Development

Program (2021YFE0114500), the National Natural Science Foundation of China

(42077023, U1803342, 41930761), the K. C. Wong Education Foundation

(GJTD-2020-14), and the Agricultural Science and Technology Innovation

Program (ASTIP).

Abstract

Aim: A fundamental challenge in soil macroecology is to understand how microbial community structure shapes ecosystem functions along environmental gradients of land surface (i.e., horizontal dimension). However, little is known of microbial community structure-function relationships along environmental gradients of soil depth (i.e., vertical dimension) in extreme environments. A full understanding of the consequences of environmental change for microbial communities structure and subsequent changes in microbial functions could enable more accurate predictions of extreme environmental change effects. Here, we leveraged a 200-km desert soil salinity gradient that is created by a 12-year saline-water irrigation to evaluate how soil microbial community structure-function relationships change with soil salinity in the horizontal and vertical dimensions.

Location: The Tarim basin of Taklamakan desert.

Taxa: Soil bacteria and fungi.

Methods: We assessed the prime ecological processes controlling the assembly of microbial communities and the activity of enzymes relevant to carbon, nitrogen, and phosphorus cycling along soil salinity gradients across study sites (horizontal dimension) and soil depths (vertical dimension) by using the general linear model, hierarchical variance partitioning, and path model.

Results: Differences in soil depth (on the scale of meters) was as important as

geographic distance (on the scale of kilometers) in shaping the structure of

bacterial and fungal communities, while both the vertical and horizontal variability

in enzymatic activity were largely attributed to the increase in the heterogeneity

of soil properties, such as soil texture, water content, and pH.

Main conclusions: Our results suggest that dispersal limitation and

environmental heterogeneity, not soil salinization, along soil depth governs

microbial community assembly and enzymatic activity, respectively. This work

highlights that conservation efforts of soil macroecology should consider soil

depth as a key attribute in the face of ongoing salinization in arid ecosystems.

Keywords: ecosystem functioning; extreme environments; microbial beta-

biodiversity; soil macroecology; structure-function relationships; Taklamakan

desert

1. Introduction

Understanding how soil microbial community structure, i.e., community biodiversity and composition, shapes spatial variability in ecosystem functions across environmental gradients is becoming a fundamental challenge in the field of soil macroecology (Chu, Gao, Ma, Fan, & Delgado-Baquerizo, 2020; Eisenhauer et al., 2021; White et al., 2020; X. F. Xu et al., 2020). Although soil microbial biodiversity is important for numerous ecosystem functions along environmental gradients on land surface (i.e., horizontal dimension) (Bardgett & van der Putten, 2014; Delgado-Baquerizo et al., 2020; X. Jing et al., 2015; van der Heijden, Bardgett, & van Straalen, 2008; D. H. Wall, Nielsen, & Six, 2015), it is a relatively new endeavor to investigate the influences of spatial variability in soil microbial community composition on spatial variability in microbial functions (Bardgett & van der Putten, 2014; T. Bell et al., 2009; Berlinches de Gea, Hautier, & Geisen, 2022; Xin Jing et al., 2022; Martinez-Almoyna et al., 2019; Mori et al., 2016; Peay, Kennedy, & Talbot, 2016; Talbot et al., 2014). One of the grand challenges is that hyper-diverse microorganisms inhabit soil matrix (Coleman & Whitman, 2005; Díaz & Malhi, 2022; Nielsen, Wall, & Six, 2015), as there can be 100-9000 microbial species/strains in a small soil sample (Bardgett & van der Putten, 2014) vs. 1~32 species in a grassland biodiversity experiment (Hector et al., 1999). This makes it much difficult to uncover microbial community structure-function relationships (Peay et al., 2016).

One strategy to address this challenge is to study community structure-

function in extreme environments, such as desert, glaciers, and hypersaline

habitats, where environmental constraints are high and biological community

structure is relatively simple (Ertekin, Meslier, Browning, Treadgold, &

DiRuggiero, 2021; Osborne, Hall, Kronfeld-Schor, Thybert, & Haerty, 2020; Shu

& Huang, 2022; Diana H Wall & Virginia, 1999). Moreover, the study of extreme

environments provides not only useful information on how environmental

conditions shape the geographic distributions of micro- and macro-organisms,

but also unique opportunities to predict how community structure and functions

responds to accelerated environmental change (N. Fierer et al., 2012; Shu &

Huang, 2022; Diana H Wall & Virginia, 1999).

In extreme arid environments (e.g., desert), niche-based environmental filters

such as salinity, soil water, and nutrient availability can shape the composition

and function of microbial communities, but these drivers can also change with

soil depth (i.e., vertical dimension) (Rath, Fierer, Murphy, & Rousk, 2019; Rath,

Murphy, & Rousk, 2019). Soil depth is a spatial dimension that is often

overlooked in the field of soil macroecology (Eisenhauer et al., 2021; White et al.,

2020) but important for an explicit understanding of how microbial community

structure relates to microbial functions through soil depth profiles (Noah Fierer,

Schimel, & Holden, 2003). In fact, prior work highlights that soil depth influences

the diversity of microbial communities, because important drivers, such as soil

organic carbon, nutrient, salinity, oxygen concentration, and moisture, vary with

soil depth (Noah Fierer et al., 2003; Powell et al., 2015; Stone, Kan, & Plante,

2015; Sun, Wang, Hui, Jing, & Feng, 2020). However, although soil depth also

regulates microbial functions such as organic matter decomposition and nutrient

cycling, we do not know much about how the vertical patterns of species turnover

in microbial community composition, i.e., β-diversity, one of the important

components of community structure, are related to the spatial variability in

microbial functions along soil profiles.

Here, we address the question how microbial community structure affects the

spatial variation in microbial activity following a 12-year saline-water irrigation

(i.e., soil salinization) in the Taklamakan desert, the world's 2nd largest shifting

sand desert with the area of roughly the size of Germany (Fig. S1). The potential

consequences of soil salinization on microbial physiology, growth and community

structure have been extensively studied (Rath & Rousk, 2015), but long-term

impacts of salinization on soil microbial community structure and enzymatic

activity in both the horizontal and vertical dimensions have been rarely

investigated. We, therefore, proposed a spatial dimension partitioning approach

(Fig. 1) to test two hypotheses: soil salinization will enhance soil microbial

species turnover (Hypothesis 1) and mediate the microbial community structure-

enzymatic activity relationships (Hypothesis 2) in the horizontal and vertical

spatial dimensions. For hypothesis 2, we specifically predict that soil salinization

will enhance species turnover when controlling for other environmental factors,

e.g., geographic distance, differences in soil chemical properties and nutrient

pools (H2a); soil salinization will indirectly enhance enzymatic turnover via soil

bacterial and fungal species turnover (H2b). Using the analytical framework, we

evaluate soil bacterial and fungal community structure and enzymatic activity

relevant to soil carbon, nitrogen, and phosphorus cycling along a unique 200-km

salinity gradient (i.e., the horizontal dimension; Fig. S2a and S2b). At each site

we also examine bacterial and fungal community assembly and enzymatic

activity across a one-meter soil profile (i.e., the vertical dimension; Fig. S2a and

S2c), an overlooked dimension of soil macroecology. In addition to soil salinity,

we assess a range of ecological factors, i.e., soil physical and chemical

properties and nutrient pools (Table S1) that could mediate microbial community

structure-enzymatic activity relationships.

Fig. 1. Conceptual depictions of spatial dimension partitioning approach. (a) heatmap

denotes the horizontal and vertical dimension of environmental gradients, i.e., soil

salinity. The gradients of soil salinity were created by a 12-year saline-water irrigation in

the Taklamakan desert. (b) a framework for illustrating the spatial dimension partitioning

of β-diversity. β_H, β_V and $\beta_{H\&V}$ denotes species turnover along the horizontal, vertical,

and horizontal & vertical gradient of soil salinity, respectively.

2. Materials and Methods

2.1. Study sites and data collection

We conducted this study in shrubland vegetation located along the Tarim desert highway, which crosses the Tarim basin of Taklamakan desert (37.1–41.8^{o} N, 82.6–84.3^{o} E) (Fig. S1). The Taklamakan desert is arid, with a mean annual precipitation of 24.6 mm yr^{-1} and a mean annual potential evaporation of 3639 mm yr^{-1}. The mean annual air temperature is 2.4°C, with the coldest monthly mean temperature reaching –8.1°C in December and the warmest monthly mean temperature reaching 28.2°C in July. Most of the desert is bare land without vegetation cover, except for a few extremely drought- and salt-tolerant shrub species, such as, *Tamarix taklamakanensis* and *Calligonum taklamakanensis*. Soils in the desert are aeolian sandy soil. The top 0☐20 cm soil is composed of 87.3% of sand, 12.4% of silt, and 0.3% of clay (J. Zhang et al., 2016). The soils have an extremely low organic matter content (0.74 g kg^{-1}) and total nitrogen (0.14 g kg^{-1}) in the top 0☐10 cm soil depth (Li, Lei, Zhao, Xu, & Li, 2015).

In 2005, the shrubland vegetation was established to prevent shifting sand dunes from damaging the Tarim desert highway. After 12 years of vegetation development, the plant community has stabilized, with *Haloxylon ammodendron* being dominant the community and *Calligonum aborescens*, *Tamarix ramosissima*, *Nitraria tangutorum*, *Populus euphratica* and *Elaeagnus angustifolia* with much lower abundance. The aboveground plant community did

not vary along the 200 km salinity gradient of study region. Additionally, the

aboveground biomass of dominant species *H. ammodendron* did not vary across

the studied sites, probably leading to comparable amount of carbon inputs into

the soil. The groundwater level in the Taklamakan desert ranges from 3 to 5 m,

and the groundwater is charged by water from the Tianshan and Kunlun

mountains (Fig. S1). The groundwater was pumped for drip irrigation to water

shrubs in the vegetation. The drip irrigation pipes run along the rows of shrubs,

and the vegetation was irrigated twice per month from March to May and from

September to October, and three times per month from June to August with ~30

L of water per plant per irrigation event (X. Xu, Li, & Wang, 2006).

We surveyed ten shrubland sites along an extreme gradient of irrigation

water salinity using a regression design of field experiment (see Pelini et al.,

2011 for an example of the regression design). The regression design used a

gradient of water salinity that varied from 3.2 to 38.0 g L^{-1} of total dissolved salts

(the normal range of seawater salinity is between 33-37 g L^{-1} of dissolved salts)

from the south to the north sampling site (Figs. S1 and S2b). We collected

irrigation water and soil samples in early October 2017, which is the end of the

growing season in the desert. At each study site, three replicates of irrigation

water were collected from running water that pumped from wells. Irrigation water

samples were stored in 500 ml plastic bottles, sealed, and taken to the laboratory

for water salinity analysis. We sampled soils using soil cores (5 cm in diameter)

at three randomly selected plots, which were located between the two rows of

shrubs and 30□50 cm away from a shrub. We collected soils to the depth of 100

cm with sampling intervals of 20 cm, except that we segmented the topsoils to

0□10 and 10□20 cm. We collected 180 soil samples (10 sites × 6 soil layers × 3

replicates). Using plastic bags and coolers, a subsample (~30 g) of each soil was

kept cool and immediately returned to the laboratory. We sieved all soils to 2 mm

and removed rocks and roots by hand. We stored soils at □20^{o}C for microbial

sequencing and at 4^{o}C for the assays of soil enzymatic activity. We air-dried the

remaining soils to analyze soil properties, including salinity, pH, water content,

texture, total organic carbon, total nitrogen, and available phosphorus.

We analyzed the salinity of irrigation water in the following ways. We

determined the concentrations of Na^+, K^+, Ca^{2+}, and Mg^{2+} using a spectrometer

(iCAP 6300-ICP-OES) with a CID detector (Thermo Scientific, MA, USA). Using

titration, we analyzed the concentration of Cl^- with $AgNO_3$, SO_4^{2-} with

ethylenediaminetetraacetic acid (EDTA), and CO_3^{2-} and HCO_3^- with sulfuric acid

until all bicarbonate was neutralized. Next, we calculated the salinity of irrigation

water as the sum of the concentrations of Na^+, K^+, Ca^{2+}, Mg^{2+}, Cl^-, SO_4^{2-}, CO_3^{2-}

and HCO_3^-.

We measured soil salinity and pH in a 1:5 suspension of soil to water. We

analyzed soil salinity using the methods described above for irrigation water

salinity. We analyzed soil pH using the Mettler Toledo FiveEasy FE28

(Greifensee, Switzerland). We measured soil gravimetric water content by oven

drying 10 g of fresh soils at 105^{o}C for 24 h. We measured soil texture using the

laser diffraction method (Sympatec GmbH, System-PartikelTechnik, Clausthal-

Zellerfeld, Germany) after removing the organic material by 10% hydrogen

peroxide and removing carbonates by 0.2% hydrochloric acid. We analyzed soil

organic carbon using the Walkley and Black method of oxidizing organic carbon

with $K_2Cr_2O_7$-H_2SO_4 (Nelson & Sommers, 1983). We analyzed soil total nitrogen

using an AutoKjeldahl Unit model K370 (BUCHI, Flawil, Switzerland) and soil

available phosphorus using the sodium bicarbonate extraction with Mo-Sb Anti-

spectrophotometer method.

To analyze soil bacterial and fungal diversity, we extracted soil DNA using

the PowerSoil DNA extraction kit (Qiagen, Carlsbad, CA, USA) following the

manufacturer's protocol. We amplified the V4 region of 16S rRNA gene for soil

bacteria using the primer pairs 515F/806R and the ITS1 region for soil fungi

using the primer pairs ITS5-1737F/ITS2-2043R. We performed PCR using the

BioRad S1000 thermal cycler (Bio-Rad Laboratory, CA, USA) after mixing 25 µl

of 2x Premix Taq polymerase, 2 µl of 10 mM primer pairs, 3 µl of 20 ng µl^{-1} DNA

template and 20 µl nuclease-free water. Next, we ran the thermal cycling as

follows: initial denaturation at 94°C for 5 min, followed by 30 cycles of

denaturation at 94°C for 30 s, annealing at 52°C for 30 s, and elongation at 72°C

for 30 s, with a final step of 72°C for 10 min. We repeated this PCR procedure

three times for each sample. We next mixed the PCR products using GeneTools

Analysis Software (Version 4.03.05.1, SynGene). The process was followed by

purifying the DNA products using the EZNA Gel extraction kit (Omega Bio-Tek,

GA, USA). Following the manufacturer's instruction, we generated sequencing

libraries using the NEBNext ® Ultra™ DNA Library Prep Kit for Illumina® (New

England Biolabs, MA, USA) and sequenced the libraries on an Illumina

HiSeq2500 platform (Illumina PE250, CA, USA).

We processed raw DNA sequences using the Trimmomatic software (V0.33,

http://www.usadellab.org/cms/?page=trimmomatic). Specifically, we excluded the

raw DNA sequences by removing paired-end reads with N, read quality <20, and

read length <100 bp. Next, we allocated the reads on the paired end to the

corresponding soil samples using the Mothur pipeline (V1.35.1,

http://www.mothur.org). We removed the barcode and primers to derive the

paired-end clean reads, which were merged with FLASH (V1.2.11,

https://ccb.jhu.edu/software/FLASH). The minimum length of overlap was 10 bp

and the maximum ratio of number of mismatches to overlap length was 0.1. Next,

we assigned these clean tags to OTUs with 97% similarity using USEARCH

software (V8.0.1517, http://www.drive5.com/usearch). We used representative

sequences for taxonomic classification using the RDP classifier according to the

Greengenes database (http://greengenes.secondgenome.com) for bacteria and

the Unite database (http://unite.ut.ee/index.php) for fungi in the QIIME pipeline

(version 2; http://qiime.org/scripts/assign_taxonomy.html).

To explore microbial functions, we assayed the potential activities of four

hydrolytic enzymes (β-1, 4-glucosidase, BG; β-$_D$- cellobiohydrolase, CB; β-1, 4-

N-acetyl-glucosaminidase, NAG; alkaline phosphatase, ALP) using the 96-well

microplate and the fluorometric technique (C. W. Bell et al., 2013; German et al.,

2011). We selected these enzymes because their potential activities are

commonly measured in soils and used as proxies for the cycling of carbon (BG

and CB), nitrogen (NAG), and phosphorus (ALP) (Sinsabaugh et al., 2008).

Briefly, sieved fresh soil (1.5 g) was suspended in 150 ml of 50 mM Tris buffer

(pH = 8.0). Soil slurries (200 µl) of each sample were transferred to eight wells of

the 96-well microplate and mixed with 50 µl of 200 µM standard fluorometric

substrates of each enzyme. The microplates were incubated for 2.5 h in the dark

at 25^{o}C. The quantity of fluorescence was determined at 360 nm excitation and

460 nm emission in a 96-well microplate reader (Biotek Synergy 2, Winooski, VT,

USA). A multiple-point calibration curve was used to calculate enzymatic

activities (C. W. Bell et al., 2013). The units were expressed as nmol g dry

weight^{-1} h^{-1}.

2.2. Analyses and Statistics

We used a two-step approach to examine how soil salinization affects soil

microbial community structure and enzymatic activity.

Step 1, to test whether increasing soil salinity enhances species turnover

(Hypothesis 1), we developed a dimension partitioning approach (Khattar,

Macedo, Monteiro, & Peres☐Neto, 2021) to quantify the impacts of vertical and

horizontal differences in soil salinity on spatial variability in the structure of

bacterial and fungal communities. We calculated the pair-wise dissimilarity matrix

between soil microbial assemblages for all combinations of study sites and soil

layers (n = 60). The lower triangle of the pair-wise dissimilarity matrix was

converted into a single vector, resulting in 1770 observations (60 × 59 / 2). We

further partitioned the single vector into three spatial dimensions, including the

horizontal, vertical, and horizontal & vertical dimensions. Differences in

community structure of the same soil layer between study sites were defined as

the horizontal dimension (10 × 9 / 2 × 6 = 270 observations), between soil layers

within a study site were defined as the vertical dimension (6 × 5 / 2 × 10 = 150

observations), and between soil layers and between sampling sites were defined

as the horizontal & vertical dimension (1350 observations).

To measure the spatial variability in soil microbial community structure

between assemblages, we used the Podani family of Sorensen indices

(Legendre, 2014; Podani & Schmera, 2011; Schmera, Podani, & Legendre,

2020), including the Sorensen dissimilarity index (β_{sor}), species replacement

(β_{repl}), and richness difference (β_{rich}). The Sorensen dissimilarity index quantifies

the total differences in species composition between assemblages. It was

partitioned into two additive components, i.e., species replacement and richness

difference. The former quantifies the species in one assemblage that are

replaced by different species in another assemblage, and the latter estimates

differences in the number of species between assemblages. We used the

following equations:

$$\beta_{sor} = \frac{b + c}{2a + b + c} \qquad\qquad \text{Eqn. 1}$$

$$\beta_{repl} = \frac{2 \times min\ (b, c)}{2a + b + c} \qquad \text{Eqn. 2}$$

$$\beta_{rich} = \frac{|b - c|}{2a + b + c} \qquad \text{Eqn. 3}$$

where *a* denotes the number of species present in both assemblages, *b* denotes

the number of species present in assemblage 1 not in assemblage 2, and *c*

denotes the number of species present in assemblage 2 not in assemblage 1.

To compare the species turnover rate with differences in soil salinity, we

calculated the Euclidean distances as the measure of differences in soil salinity.

We calculated differences in soil salinity using the concentrations of soil Na^+, K^+,

Ca^{2+}, Mg^{2+}, Cl^-, SO_4^{2-}, and HCO_3^{2-} as follows:

$$D(x, y) = \sqrt{\sum_{i=1}^{n}(x_i - y_i)^2} \qquad \text{Eqn. 4}$$

where *D(x, y)* denotes the Euclidean distance, *n* is the number of salt ions (*n*

= 7), x_i and y_i denote the concentration of salt ions in the sample *x* and sample *y*,

respectively. We regressed pair-wise compositional dissimilarity (i.e., β-diversity)

against differences in soil salinity across the horizontal and vertical spatial

dimensions. We used the standardized regression slopes to compare the rate of

species turnover among spatial dimensions and the permutation test to examine

the significance of regression slopes.

Step 2, to test whether increasing soil salinity mediate microbial community

structure-enzymatic activity relationships (Hypothesis 2), we used hierarchical

variance partitioning and path model. We conducted hierarchical variance

partitioning to estimate the proportion of spatial variability in the structure of

microbial communities (Hypothesis 2a). We used the Euclidean distances to

calculate geographic distance and differences in soil depth, salinity, properties,

and nutrient pools (Table S1). Specifically, the geographic distance was

calculated using the geographic coordinates of the sampling sites. Differences in

soil depth were calculated using the mean depth of each sampling layer.

Differences in soil property were calculated using soil clay content, silt content,

sand content, pH, and moisture. Differences in nutrient pools were calculated

using the data of soil organic carbon, soil total nitrogen, and available

phosphorus (Xin Jing et al., 2022). All combinations of explanatory variables

were fitted into regression models following the hierarchical partitioning algorithm

of Chevan and Sutherland (1991). The independent effects of each explanatory

variable were determined based on the goodness-of-fit measures (i.e., R^2) of

regression models, with the target variables (microbial community structure or

enzymatic activity) as the dependent variable. We used 9,999 bootstrapped

sampling iterations to calculate the 95% confidence interval of the independent

effects of each explanatory variable.

We used the path model to estimate the indirect effects of soil salinity on

enzymatic turnover via bacterial and fungal species turnover (Hypothesis 2b). We

considered the potential enzymatic activities of BG, CB, NAG and ALP as a

proxy for microbial functions (Burns et al., 2013; Fanin et al., 2022; Talbot et al.,

2014). We used the Euclidean distances as the measures of spatial variability in

microbial functions, i.e., enzymatic turnover or differences in enzymatic activity

(Xin Jing et al., 2021; Xin Jing et al., 2022; Martinez-Almoyna et al., 2019; Mori et

al., 2016). The path model was also used to assess the relative importance of

dispersal limitation and environmental filtering on the structure of soil microbial

communities (Xin Jing et al., 2021; Xin Jing et al., 2022). For soil bacterial and

fungal community structure, we considered geographic distance and differences

in soil depth as the proxies for dispersal limitation (Powell et al., 2015) and

differences in soil salinity, properties, and nutrient pools as the proxies for

environmental filtering (Xin Jing et al., 2022; Martiny et al., 2006). Using path

model, we also estimated the relative importance of environmental variables and

community structure of soil bacteria and fungi on differences in microbial

enzymatic activity. We used the P value of χ^2 statistic test (P >0.05), comparative

fit index (CFI >0.90), root mean square error of approximation (RMSEA <0.05),

and standardized root mean square residual (SRMR <0.10) as measures of path

model evaluation and selection (Grace, 2020). Since a large sample size (n >

100) can lead to the P value of χ^2 statistic test that was less than 0.05 (Grace,

2020), we only used CFI, RMSEA, and SRMR to evaluate and select models for

overall and horizontal and vertical spatial dimensions. We used 9,999

bootstrapped sampling iterations to test the significance of the path coefficients.

All path models fitted the data well (Table S2).

The dimension partitioning was carried out for all the variables, including soil

microbial community structure and enzymatic activity, geographic distance, depth

difference, salinity difference, property difference, and nutrient pool difference.

Prior to data analyses, we removed singleton OTUs, which are present only once

across all microbial assemblages, from soil microbial community data and pooled

all the data to the level of site × soil layer to remove the non-independent

influences of sample replicates. All statistical analyses were performed in R

version 3.6.1 (R Development Core Team, 2019) using the packages 'lmPerm'

(Wheeler, Torchiano, & Torchiano, 2016), 'hier.part' (Nally & Walsh, 2004), and

'lavaan' (Rosseel, 2012).

3. Results

Bacterial total β-diversity (β_{sor}) increased with differences in soil salinity

across horizontal, vertical, and horizontal & vertical spatial dimensions (Fig. 2a,

2c, and 2e). Bacterial richness difference (β_{rich}) showed weakly positive relations

with differences in soil salinity, while bacterial replacement was low at high level

of differences in soil salinity across the horizontal and horizontal & vertical spatial

dimensions (Fig. 2a and 2e). Fungal β_{sor} decreased with differences in soil

salinity in the horizontal spatial dimension (Fig. 2b). Fungal β_{repl} decreased with

differences in soil salinity in the horizontal spatial dimension but increased with

differences in soil salinity in the vertical spatial dimension (Fig. 2b and 2d).

Fungal β_{rich} increased with differences in soil salinity in the horizontal spatial

dimension, while had no relations with differences in soil salinity in other spatial

dimensions (Fig. 2b, 2d and 2f).

Fig. 2. Relationships between differences in microbial β-diversity and soil salinity

across the horizontal and vertical spatial dimensions. Solid lines denote

significant regressions (if †, 0.05 < P < 0.10, otherwise P < 0.05) and dashed

lines denote nonsignificant regressions (P > 0.10). Significant standardized

regression coefficients are shown at the top right side of each panel and

nonsignificant standardized regression coefficients are shown as '--'. β_{sor} =

Sorensen dissimilarity index (blue); β_{repl} = replacement component of β_{sor} (dark

gray); β_{rich} = richness difference component of β_{sor} (gray).

Differences in soil salinity did not exert significant independent effects on soil

microbial β_{sor}, β_{repl}, and β_{rich} (Figs. 3, S3 and S4; Table S3). By contrast,

geographic distance and differences in soil depth had the largest independent

effects on differences in soil bacterial and fungal β_{sor} in the horizontal or vertical

spatial dimensions (Fig. 3a-3f). Geographic distance explained 20.7-25.2% of the

variance in differences in soil bacterial β_{sor} and 24.1-25.3% of the variance in

differences in soil fungal β_{sor} (Fig. 3a, 3c, 3d and 3f; Table S3). Throughout the

vertical dimension, the differences in soil depth explained 21.9% (95%

confidence interval [12.4, 32.7]) of the variance in differences in soil bacterial β_{sor}

(Fig. 3b Table S4) and 23.7% [12.7, 35.8] of the variance in differences in soil

fungal community structure (Fig. 3b and 3e; Table S3).

Fig. 3. Independent effects of explanatory variables on microbial β-diversity

across the horizontal and vertical spatial dimensions by hierarchical variance

partitioning. Bars represent the pure variance explained by the explanatory

variables, and the proportion of explained variance is shown along the bars. The

95% confidence interval is given. We show only the Sorensen dissimilarity index,

see Figs. S3-S4 and Table S2 for the summary of the replacement and richness

difference component of Sorensen dissimilarity index.

Path analysis showed that geographic distance and differences in soil depth,

salinity, properties, and nutrient pools separately explained 14%□27% of the

variance in differences in enzymatic activity, 29%□39% of the variance in

differences in soil bacterial community structure, and 27%⎕32% of the variance

in differences in soil fungal community structure (Fig. 4a-4c). However,

differences in enzymatic activity were not explained by differences in soil

microbial community structure (standardized path coefficients $β_{std}$ < 0.13 in all

spatial dimensions; Figs. 4 and S5-S10; Table S4). Differences in soil properties

but not soil salinity had the largest effects on differences in enzymatic activity in

the vertical dimension, which were as important as the effects in the horizontal

dimension (Fig. 4). In addition, differences in soil bacterial and fungal community

structure were weakly associated with differences in soil salinity (Fig. 4). The

effects of differences in soil depth (Fig. 4b) on soil bacterial and fungal $β_{sor}$ along

the vertical dimension were as important as the effects of geographic distance

(Fig. 4a and 4c) along the horizontal dimension on soil bacterial and fungal $β_{sor}$.

Fig. 4. Path model shows the direct and indirect effects of soil salinity on

microbial community structure and enzymatic activity across the horizontal and

vertical spatial dimensions. Three path models are shown based on spatial

dimensions, including a) pure horizontal dimension; b) pure vertical dimension; c)

combined horizontal & vertical dimension. The width of the arrows is proportional

to the standardized path coefficients. Black arrows denote positive path

coefficients and red arrows denote negative path coefficients. Bootstraps are

used for the significance test of path coefficients (see Tables S3-S4 for details).

R^2 denotes the variance in community structure and enzymatic activity explained

by environmental factors, including geographic distance (GeoDist), differences in

soil depth (DepDiff), differences in soil salinity (SalDiff), differences in soil

properties (PropDiff), and differences in nutrient pools (NutPool). BacStruc,

FunStruc, and EnzAct denote differences in bacterial community structure, fungal

community structure, and enzymatic activity, respectively. We show only the

Sorensen dissimilarity index, see Figs. S10-S11 and Tables S3-S4 for summary

of the replacement and richness difference component of Sorensen dissimilarity

index.

4. Discussion

Here, we used a unique environmental (i.e., salinity) gradient that display

spatial variation in both the horizonal and vertical dimensions. Using this salinity

gradient, we were able to separate the influences of horizontal soil salinization

(on the scale of kilometers) on microbial community assembly and enzymatic

activity from the influences of vertical soil salinization (one the scale of meters),

which is an overlooked dimension of soil macroecology. Our first hypothesis that

increasing soil salinity will lead to enhanced microbial species turnover was

weakly supported by fungal species turnover in the vertical spatial dimension.

Instead, we found that richness difference contributed to variation in bacterial and

fungal species composition in both the horizontal and vertical spatial dimensions

(Fig. 2). Our second hypothesis was rejected because geographic distance in the horizontal dimension and difference in soil depth in the vertical dimension, rather than differences in soil salinity, consistently determined variation in bacterial and fungal species composition (Fig. 3). Differences in soil properties rather than soil salinity enhanced the spatial turnover in enzymatic activity (Fig. 4). Our findings suggest that soil depth serves as a test bed for predicting changes in soil microbial community structure and enzymatic activity in extreme environments.

4.1. Soil depth as a new frontier for studies in soil macroecology

In this work, we found that spatial variability in the structure of soil microbial communities was strongly associated with geographic distance and differences in soil depth rather than differences in soil salinity. This result suggests that soil bacterial and fungal communities are resistant to changes in soil salinity but show a dissimilar community structure with an increase in spatial distance both horizontally and vertically along soil depth profile. Our finding that soil microbial community structure is resistant to changes in soil salinity is in contrast with previous work (Rath, Fierer, et al., 2019; Van Horn et al., 2014; K. Zhang et al., 2019). One reason must be that soil salinity is extremely high after 12-year of saline-water irrigation, which created a long-lasting harsh environmental condition (Schradin, Makuya, Pillay, & Rimbach, 2023). Soil microbes may have adapted well to the harsh environment, as low species turnover was associated with differences in soil salinity (Fig. 2). However, our work indicates that soil depth is as important as geographic distance in shaping the structure of bacterial

and fungal community in the saline desert soil (Figs. 3 and 4). The importance of

soil depth is supported by an early study showing that random assembly

processes of soil microbial communities (e.g., dispersal limitation) intensified with

soil depth because soil microorganisms disperse passively (Powell et al., 2015).

Our results indicate that soil depth and geographic distance are equally important

when the overall spatial variability in microbial structure is partitioned into the

pure vertical and horizontal spatial dimensions (Fig. 3a,b). Without spatial

dimension partitioning, we will underestimate the contribution of soil depth to the

assembly processes of soil microbial communities. Together, our work highlights

that dispersal limitation in soil microbes is particularly important to the assembly

of microbial communities in the vertical and horizontal dimensions of soil depth

profiles in extreme environments. To maintain the essential and irreplaceable

ecosystem functions and services provided by belowground biodiversity in

extreme conditions or severely disturbed conditions, a workable plan is to

preserve habitats along both the horizontal and vertical habitats along soil depth

profiles.

Our study further indicates that the spatial variability in soil enzymatic activity

is best explained by the heterogeneity in soil texture, pH, and water rather than

differences in soil salinity and nutrient pools across soil depths. In extreme

environments, environmental filters (e.g., salt stress) and resource levels (e.g.,

soil organic matter and microbial biomass) are expected to be the drivers of

spatial variability in soil microbial community structure and function (Martiny et

al., 2006; Rath, Fierer, et al., 2019; Rath, Murphy, et al., 2019; Rath & Rousk,

2015). On the contrary, our results suggest that differences in soil salinity and

nutrient pools do not explain the activity of carbon, nitrogen, and phosphorus

cycling related enzymes. It could be that soil salinity is too high in our study

system and the priority of soil microbes is to survive rather than to thrive in such

extreme saline environments, which does not allow soil microbes to produce

energy-expensive enzymes (Rath & Rousk, 2015). As a result, soil salinity and

resources are not limiting factors of enzymatic activity, and changes in soil

salinity and nutrient pools did not alter enzymatic activity. However, our result

indicates that reduced heterogeneity in soil properties, which could be induced by

land degradation, soil erosion, and desertification, might result in loss of soil

microbial activity and soil functionality, soil fertility, and plant growth. Since the

variability in soil physical and chemical properties is created by human activities

and nature ubiquitously, we should analyze the overlooked dimension of soil

macroecology in the vertical dimension along soil depth profiles, which represent

a valuable step towards soil conservation efforts in dealing with ongoing

desertification in arid ecosystems.

4.2. Uncoupled microbial community structure-function relationships

The most salient result of this study is that differences in soil microbial

community structure did not lead to differences in enzymatic activity in the hyper-

saline environments. This result runs counter to previous work exploring these

relationships in microbial systems with a focus on surface soils (Delgado-

Baquerizo et al., 2020; Xin Jing et al., 2022; Mori et al., 2016). The uncoupled

relationships between microbial structure and function were further verified

across the horizontal and vertical dimensions of soil depth profile. Thus, our

findings support functional redundancy (Coleman & Whitman, 2005; Peay et al.,

2016), implying that even in extreme environments the structure of soil microbial

communities differ, but their function in the same manner. As a result, the

losses/gains of certain species may have limited effects on the spatial variability

in enzymatic activity. This means that functionally redundant communities may

safeguard ecosystem functions against the negative impacts of extreme

environmental changes (Allison & Martiny, 2008).

There are two explanations for the uncoupled structure-function relationships

under soil salinization. First, soil bacterial and fungal communities have likely

adapted to the extreme conditions in desert, where the abiotic stress is high

(Jansson & Hofmockel, 2020). As the soil becomes saltier and drier, soil

microorganisms become inactive because the availability of carbon and nutrients

in the soil decreases. In addition, soil microorganisms tend to minimize their

activity, even becoming dormant, to conserve energy under harsh conditions. In

either case, we would observe the uncoupled relationships between soil

microbial community structure and enzymatic activity. Second, the uncoupled

structure-function relationship is likely due to the mismatch of dominant driving

factors responsible for microbial community structure and enzymatic activity. Our

results show that geographic distance and differences in soil depth have the

strongest independent effects on the structure of microbial communities, while

differences in soil properties have the strongest independent effects on

enzymatic activity. Different driving factors could have disproportionate

influences on microbial community structure and enzymatic activity in soil,

leading to uncoupled structure-function relationships.

In recent years, soil macroecologists have focused on microbial communities

in surface soils because they have high biomass density and nutrient cycling

activities (X. F. Xu et al., 2020). This study is unique in that it provides a new

approach to assess microbial community structure-function relationships both

horizontally and vertically along soil depth profiles with the same range of salinity

variability in a typical extreme biome, i.e., the Taklamakan desert. However,

large-scale studies remain the key to verify whether the findings hold true in other

biomes, which may have a wider range of horizontal and vertical gradients that

are induced by human activities or by nature under the context of soil

degradation, e.g., soil salinization, acidification, and erosion. Further, it is

important to note that the measure of four hydrolytic enzyme activities capture

only a few microbial functions that are relevant to carbon, nitrogen, and

phosphorus cycling. Future work that uses shotgun metagenomics could quantify

broad shifts in microbial community functionality, e.g., the functions related to

organotrophs and autotrophs (both by phototrophs and chemolithotrophs),

advancing our understanding of the structure-function relationships of microbial

communities in xeric or arid soils.

5. Conclusions

We find that environmental gradients along soil depth profile is as important

as horizontal geographic distance in shaping microbial community structure and

enzymatic activity. These findings highlight the need to analyze the overlooked

dimension of soil macroecology, i.e., the vertical spatial dimension (Eisenhauer

et al., 2021; White et al., 2020). Such information is crucial to predict the

influences of global change drivers on real-world community structure-function

relationships across horizontal and vertical spatial scales. Moreover, this study

highlights that given the increase in desertification in arid ecosystems, soil

conservation efforts should take soil depth into account as a critical factor

governing the assembly of microbial communities and the activity of enzymes.

References

Allison, S. D., & Martiny, J. B. (2008). Resistance, resilience, and redundancy in

microbial communities. *Proceedings of the National Academy of Sciences, USA,*

105(Supplement 1), 11512-11519.

Bardgett, R. D., & van der Putten, W. H. (2014). Belowground biodiversity and

ecosystem functioning. *Nature, 515*(7528), 505-511. doi:10.1038/nature13855

Bell, C. W., Fricks, B. E., Rocca, J. D., Steinweg, J. M., McMahon, S. K., & Wallenstein,

M. D. (2013). High-throughput fluorometric measurement of potential soil

extracellular enzyme activities. *Journal of Visualized Experiments*(81), e50961.

doi:doi:10.3791/50961

Bell, T., Gessner, M. O., Griffiths, R. I., McLaren, J. R., Morin, P. J., Van Der Heijden,

M., & van der Putten, W. H. (2009). Microbial biodiversity and ecosystem

functioning under controlled conditions and in the wild. In S. Naeem, B. D. E, A.

Hector, M. Loreau, & C. Perrings (Eds.), *Biodiversity, ecosystem functioning,*

human wellbeing: an ecological economic perspective (pp. 121-133). New York,

USA: Oxford University Press.

Berlinches de Gea, A., Hautier, Y., & Geisen, S. (2022). Interactive effects of global

change drivers as determinants of the link between soil biodiversity and

ecosystem functioning. *Global Change Biology, 00*, 1-12. doi:10.1111/gcb.16471

Burns, R. G., DeForest, J. L., Marxsen, J., Sinsabaugh, R. L., Stromberger, M. E.,

Wallenstein, M. D., . . . Zoppini, A. (2013). Soil enzymes in a changing

environment: Current knowledge and future directions. *Soil Biology and*

Biochemistry, 58, 216-234. doi:10.1016/j.soilbio.2012.11.009

Chevan, A., & Sutherland, M. (1991). Hierarchical partitioning. *The American*

Statistician, 45(2), 90-96.

Chu, H., Gao, G. F., Ma, Y., Fan, K., & Delgado-Baquerizo, M. (2020). Soil microbial

biogeography in a changing world: Recent advances and future perspectives.

mSystems, 5(2), e00803-00819. doi:10.1128/mSystems.00803-19

Coleman, D. C., & Whitman, W. B. (2005). Linking species richness, biodiversity and

ecosystem function in soil systems. *Pedobiologia, 49*(6), 479-497.

Delgado-Baquerizo, M., Reich, P. B., Trivedi, C., Eldridge, D. J., Abades, S., Alfaro, F.

D., . . . Singh, B. K. (2020). Multiple elements of soil biodiversity drive ecosystem

functions across biomes. *Nature Ecology & Evolution, 4*(2), 210-220.

doi:10.1038/s41559-019-1084-y

Díaz, S., & Malhi, Y. (2022). Biodiversity: Concepts, patterns, trends, and perspectives.

Annual Review of Environment and Resources, 47(1), 31-63.

doi:10.1146/annurev-environ-120120-054300

Eisenhauer, N., Buscot, F., Heintz-Buschart, A., Jurburg, S. D., Kusel, K., Sikorski, J., . .

. Guerra, C. A. (2021). The multidimensionality of soil macroecology. *Glob Ecol*

Biogeogr, 30(1), 4-10. doi:10.1111/geb.13211

Ertekin, E., Meslier, V., Browning, A., Treadgold, J., & DiRuggiero, J. (2021). Rock

structure drives the taxonomic and functional diversity of endolithic microbial

communities in extreme environments. *Environmental Microbiology, 23*(7), 3937-

3956. doi:10.1111/1462-2920.15287

Fanin, N., Mooshammer, M., Sauvadet, M., Meng, C., Alvarez, G., Bernard, L., . . .

Nottingham, A. T. (2022). Soil enzymes in response to climate warming:

Mechanisms and feedbacks. *Functional Ecology, 36*, 1378-1395.

doi:10.1111/1365-2435.14027

Fierer, N., Leff, J. W., Adams, B. J., Nielsen, U. N., Bates, S. T., Lauber, C. L., . . .

Caporaso, J. G. (2012). Cross-biome metagenomic analyses of soil microbial

communities and their functional attributes. *Proceedings of the National*

Academy of Sciences, USA, 109(52), 21390-21395.

doi:10.1073/pnas.1215210110

Fierer, N., Schimel, J. P., & Holden, P. A. (2003). Variations in microbial community

composition through two soil depth profiles. *Soil Biology and Biochemistry, 35*(1),

167-176. doi:http://dx.doi.org/10.1016/S0038-0717(02)00251-1

German, D. P., Weintraub, M. N., Grandy, A. S., Lauber, C. L., Rinkes, Z. L., & Allison,

S. D. (2011). Optimization of hydrolytic and oxidative enzyme methods for

ecosystem studies. *Soil Biology and Biochemistry, 43*(7), 1387-1397.

doi:10.1016/j.soilbio.2011.03.017

Grace, J. (2020). A 'Weight of Evidence' approach to evaluating structural equation

models. *One Ecosystem, 5*, e50452. doi:10.3897/oneeco.5.e50452

Hector, A., Schmid, B., Beierkuhnlein, C., Caldeira, M. C., Diemer, M., Dimitrakopoulos,

P. G., . . . Lawton, J. H. (1999). Plant diversity and productivity experiments in

European grasslands. *Science, 286*(5442), 1123-1127.

doi:10.1126/science.286.5442.1123

Jansson, J. K., & Hofmockel, K. S. (2020). Soil microbiomes and climate change. *Nature*

Reviews Microbiology, 18(1), 35-46. doi:10.1038/s41579-019-0265-7

Jing, X., Prager, C. M., Borer, E. T., Gotelli, N. J., Gruner, D. S., He, J.-S., . . . Sanders,

N. J. (2021). Spatial turnover of multiple ecosystem functions is more associated

with plant than soil microbial β-diversity. *Ecosphere, 12*(7), e03644.

doi:10.1002/ecs2.3644

Jing, X., Prager, C. M., Chen, L., Chu, H., Gotelli, N. J., He, J. S., . . . Sanders, N. J.

(2022). The influence of aboveground and belowground species composition on

spatial turnover in nutrient pools in alpine grasslands. *Global Ecology and*

Biogeography, 31(3), 486-500. doi:10.1111/geb.13442

Jing, X., Sanders, N. J., Shi, Y., Chu, H. Y., Classen, A. T., Zhao, K., . . . He, J.-S.

(2015). The links between ecosystem multifunctionality and above- and

belowground biodiversity are mediated by climate. *Nature Communications, 6*,

8159. doi:10.1038/ncomms9159

Khattar, G., Macedo, M., Monteiro, R., & Peres-Neto, P. (2021). Determinism and

stochasticity in the spatial–temporal continuum of ecological communities: the

case of tropical mountains. *Ecography, 44*(9), 1391-1402.

doi:10.1111/ecog.05726

Legendre, P. (2014). Interpreting the replacement and richness difference components

of beta diversity. *Global Ecology and Biogeography, 23*(11), 1324-1334.

doi:10.1111/geb.12207

Li, C., Lei, J., Zhao, Y., Xu, X., & Li, S. (2015). Effect of saline water irrigation on soil

development and plant growth in the Taklimakan Desert Highway shelterbelt. *Soil*

and Tillage Research, 146, 99-107.

Martinez-Almoyna, C., Thuiller, W., Chalmandrier, L., Ohlmann, M., Foulquier, A.,

Clément, J. C., . . . Münkemüller, T. (2019). Multi-trophic β-diversity mediates the

effect of environmental gradients on the turnover of multiple ecosystem functions.

Functional Ecology, 33, 2053-2064. doi:10.1111/1365-2435.13393

Martiny, J. B., Bohannan, B. J., Brown, J. H., Colwell, R. K., Fuhrman, J. A., Green, J. L.,

. . . Staley, J. T. (2006). Microbial biogeography: putting microorganisms on the

map. *Nature Review Microbiology, 4*(2), 102-112. doi:10.1038/nrmicro1341

Mori, A. S., Isbell, F., Fujii, S., Makoto, K., Matsuoka, S., & Osono, T. (2016). Low

multifunctional redundancy of soil fungal diversity at multiple scales. *Ecology*

Letters, 19(3), 249-259. doi:10.1111/ele.12560

Nally, R. M., & Walsh, C. J. (2004). Hierarchical partitioning public-domain software.

Biodiversity and Conservation, 13(3), 659.

Nelson, D. W., & Sommers, L. E. (1983). Total Carbon, Organic Carbon, and Organic

Matter. In A. L. Page (Ed.), *Methods of Soil Analysis. Part 2. Chemical and*

microbiological properties. (pp. 539-579). Madison, WI, USA: ASA and SSSA.

Nielsen, U. N., Wall, D. H., & Six, J. (2015). Soil biodiversity and the environment.

Annual Review of Environment and Resources, 40(1), 63-90.

doi:10.1146/annurev-environ-102014-021257

Osborne, P., Hall, L. J., Kronfeld-Schor, N., Thybert, D., & Haerty, W. (2020). A rather

dry subject; investigating the study of arid-associated microbial communities.

Environmental Microbiome, 15(1), 20. doi:10.1186/s40793-020-00367-6

Peay, K. G., Kennedy, P. G., & Talbot, J. M. (2016). Dimensions of biodiversity in the

Earth mycobiome. *Nature Reviews Microbiology, 14*(7), 434-447.

doi:10.1038/nrmicro.2016.59

Pelini, S. L., Bowles, F. P., Ellison, A. M., Gotelli, N. J., Sanders, N. J., & Dunn, R. R.

(2011). Heating up the forest: open-top chamber warming manipulation of

arthropod communities at Harvard and Duke Forests. *Methods in Ecology and*

Evolution, 2(5), 534-540. doi:10.1111/j.2041-210X.2011.00100.x

Podani, J., & Schmera, D. (2011). A new conceptual and methodological framework for

exploring and explaining pattern in presence–absence data. *Oikos, 120*(11),

1625-1638. doi:10.1111/j.1600-0706.2011.19451.x

Powell, J. R., Karunaratne, S., Campbell, C. D., Yao, H., Robinson, L., & Singh, B. K.

(2015). Deterministic processes vary during community assembly for ecologically

dissimilar taxa. *Nature Communications, 6*(1), 1-10.

R Development Core Team. (2019). *R: A language and environment for statistical*

computing. R Foundation for Statistical Computing, Vienna, Austria.

https://www.R-project.org/.

Rath, K. M., Fierer, N., Murphy, D. V., & Rousk, J. (2019). Linking bacterial community
composition to soil salinity along environmental gradients. *The ISME Journal,*
13(3), 836-846. doi:10.1038/s41396-018-0313-8
Rath, K. M., Murphy, D. N., & Rousk, J. (2019). The microbial community size, structure,
and process rates along natural gradients of soil salinity. *Soil Biology and*
Biochemistry, 138, 107607. doi:https://doi.org/10.1016/j.soilbio.2019.107607
Rath, K. M., & Rousk, J. (2015). Salt effects on the soil microbial decomposer
community and their role in organic carbon cycling: A review. *Soil Biology and*
Biochemistry, 81, 108-123. doi:https://doi.org/10.1016/j.soilbio.2014.11.001
Rosseel, Y. (2012). lavaan: An R Package for Structural Equation Modeling. *Journal of*
Statistical Software, 48(2), 1-36. doi:10.18637/jss.v048.i02
Schmera, D., Podani, J., & Legendre, P. (2020). What do beta diversity components
reveal from presence-absence community data? Let us connect every indicator
to an indicandum! *Ecological Indicators, 117*, 106540.
doi:10.1016/j.ecolind.2020.106540
Schradin, C., Makuya, L., Pillay, N., & Rimbach, R. (2023). Harshness is not stress.
Trends in Ecology & Evolution. doi:10.1016/j.tree.2022.12.005
Shu, W. S., & Huang, L. N. (2022). Microbial diversity in extreme environments. *Nature*
Reviews Microbiology, 20(4), 219-235. doi:10.1038/s41579-021-00648-y
Sinsabaugh, R. L., Lauber, C. L., Weintraub, M. N., Ahmed, B., Allison, S. D., Crenshaw,
C., . . . Zeglin, L. H. (2008). Stoichiometry of soil enzyme activity at global scale.
Ecology Letters, 11(11), 1252-1264. doi:10.1111/j.1461-0248.2008.01245.x
Stone, M. M., Kan, J., & Plante, A. F. (2015). Parent material and vegetation influence
bacterial community structure and nitrogen functional genes along deep tropical
soil profiles at the Luquillo Critical Zone Observatory. *Soil Biology and*
Biochemistry, 80, 273-282.
Sun, T., Wang, Y., Hui, D., Jing, X., & Feng, W. (2020). Soil properties rather than
climate and ecosystem type control the vertical variations of soil organic carbon,
microbial carbon, and microbial quotient. *Soil Biology and Biochemistry, 148*,
107905.
Talbot, J. M., Bruns, T. D., Taylor, J. W., Smith, D. P., Branco, S., Glassman, S. I., . . .
Peay, K. G. (2014). Endemism and functional convergence across the North

American soil mycobiome. *Proceedings of the National Academy of Sciences,*
USA, 111(17), 6341-6346. doi:10.1073/pnas.1402584111

van der Heijden, M. G. A., Bardgett, R. D., & van Straalen, N. M. (2008). The unseen
majority: soil microbes as drivers of plant diversity and productivity in terrestrial
ecosystems. *Ecology Letters, 11*, 296-310. doi:10.1111/j.1461-
0248.2007.01139.x

Van Horn, D. J., Okie, J. G., Buelow, H. N., Gooseff, M. N., Barrett, J. E., & Takacs-
Vesbach, C. D. (2014). Soil microbial responses to increased moisture and
organic resources along a salinity gradient in a polar desert. *Applied and*
Environmental Microbiology, 80(10), 3034-3043. doi:10.1128/AEM.03414-13

Wall, D. H., Nielsen, U. N., & Six, J. (2015). Soil biodiversity and human health. *Nature,*
528(7580), 69-76. doi:10.1038/nature15744

Wall, D. H., & Virginia, R. A. (1999). Controls on soil biodiversity: insights from extreme
environments. *Applied Soil Ecology, 13*(2), 137-150.

Wheeler, B., Torchiano, M., & Torchiano, M. M. (2016). lmPerm: Permutation tests for
linear models. *R Package Version 2.1.0. https://CRAN.R-*
project.org/package=lmPerm.

White, H. J., León-Sánchez, L., Burton, V. J., Cameron, E. K., Caruso, T., Cunha, L., . . .

Schrodt, F. (2020). Methods and approaches to advance soil macroecology.
Global Ecology and Biogeography, 29(10), 1674-1690. doi:10.1111/geb.13156

Xu, X., Li, B., & Wang, X. (2006). Progress in study on irrigation practice with saline
groundwater on sandlands of Taklimakan Desert Hinterland. *Chinese Science*
Bulletin, 51(1), 161-166.

Xu, X. F., Wang, N. N., Lipson, D., Sinsabaugh, R., Schimel, J., He, L. Y., . . . Tedersoo,
L. (2020). Microbial macroecology: In search of mechanisms governing microbial
biogeographic patterns. *Global Ecology and Biogeography, 29,* 1870-1886.
doi:10.1111/geb.13162

Zhang, J., Xu, X., Li, S., Zhao, Y., Zhang, A., Zhang, T., & Jiang, R. (2016). Is the
Taklimakan desert highway shelterbelt sustainable to long-term drip irrigation
with high saline groundwater? *Plos One, 11*(10), e0164106.

Zhang, K., Shi, Y., Cui, X., Yue, P., Li, K., Liu, X., . . . Chu, H. (2019). Salinity Is a key
determinant for soil microbial communities in a desert ecosystem. *mSystems,*
4(1). doi:10.1128/mSystems.00225-18

Author Contributions: XJ, WF and YW designed the study. YW and WF

conducted the field- and lab-work. XJ analyzed the data. XJ and WF wrote the

first draft of the manuscript. All authors contributed to the conceptualization,

review and editing of the manuscript.

Conflict of Interest: The authors declare that they have no competing interests.

Data Availability: Sequence data were deposited in the European Nucleotide

Archive with accession number PRJEB52395

(http://www.ebi.ac.uk/ena/data/view/OW744067-OW763241) for soil bacteria,

and with accession number PRJEB46878

(http://www.ebi.ac.uk/ena/data/view/OU495976-OU501470) for soil fungi. The

data and code will be publicly available in the Zenodo repository prior to

publication.

Biosketch

Xin Jing is a professor in the College of Pastoral Agriculture Science and

Technology at Lanzhou University. He is interested in understanding the causes

of biodiversity change and its consequences for ecosystem structure and

function.