A Positive Correlation between CRISPR-Cas Spacer Abundance and Cell Cycle Duration Across the Bacteria Domain

Zhi-Ling Liu, En-Ze Hu, and Deng-Ke Niu*

MOE Key Laboratory for Biodiversity Science and Ecological Engineering and Beijing Key Laboratory of Gene Resource and Molecular Development, College of Life Sciences, Beijing Normal University, Beijing, 100875, China

*Corresponding author.

Mailing address for all the authors:

College of Life Sciences, Beijing Normal University, Beijing, 100875, China

Telephones and Emails:

Zhi-Ling Liu, ++8618801280771, 201931200042@mail.bnu.edu.cn

En-Ze Hu, ++8615810572691, enzehu@126.com

Deng-Ke Niu, ++8613683295521, dkniu@bnu.edu.cn, dengkeniu@hotmail.com

Abstract

CRISPR-Cas systems provide adaptive immunity for prokaryotic cells by recognizing and eliminating the recurrent genetic invaders whose sequences had been captured in a prior infection and stored in the CRISPR arrays as spacers. However, the biological factors determining the efficiency of this adaptive immune system have not been fully characterized. Recent studies in cultured bacteria showed that slowing the growth rate of bacterial cells could promote their acquisition of novel spacers. This paper reports a positive correlation between the minimal doubling time and the spacer abundance across the bacteria domain. The correlation is still significant after controlling for the effect of growth temperature. This observation provides evidence for the existence of the trade-off between bacterial growth and defense proposed in the "Killing the Winner" hypothesis.

Keywords

CRISPR-Cas, cell cycle, bacteria, phage, growth rate.

CRISPR-Cas systems provide adaptive immunity for prokaryotic cells against invasive

genetic elements. Accumulating evidence indicates that slowing down host cell growth rate

could enhance spacer acquisition probably because the slow growth affects virus replication

more than the cell's response to the invaders. Using a fluorescent CRISPR adaptation reporter

method, Amlinger et al. (1) quantified the adaptation efficiency of Type II CRISPR in the

different growth phases of *Escherichia coli*. They found that spacer acquisition occurs

predominantly during the late exponential/early stationary phase. In line with this study,

McKenzie et al. (2) observed that the *E. coli* cells that adapted and cleared the target DNA

earlier grew significantly slower than the population means. The soil-dwelling bacterium

Pseudomonas aeruginosa can also infect the plant and human bodies. It grows slowly in the

cooler temperatures of the soil or during plant infection and rapidly under the higher

temperature of human bodies. Høyland-Kroghsbo et al. (3) found that decreasing the growth

rate could increase the CRISPR adaptation frequency of the bacterium. Dimitriu et al. (4)

recently found that the bacteriostatic agents that slowing down the growth of *P. aeruginosa*

could promote the phage-derived acquisition of novel spacers into the CRISPR array.

Furthermore, they showed that limited carbon sources associated with slow bacterial growth

could also enhance the evolution of CRISPR immunity. The latter result is in line with the

previous studies that observed increased CRISPR-Cas immunity in bacteria grown in a

nutrient-limited medium (4). Dormancy could be regarded as a prolonged growth state of the

prokaryotic cells. In the response of the type VI CRISPR-Cas system to RNA virus infection,

the Cas13 enzymes of the bacteria *Listeria ivanovii* were found to destruct both the RNA

virus genomes and the host mRNA molecules, which further induce dormancy of the host

bacterial cells (5). In the same study, the immunity of the bacterial cells against the unrelated

virus was enhanced in the type-VI-CRISPR-Cas-system-induced dormant state.

All these results were obtained from analyzing a few cultured bacteria in laboratories. We

examined whether there is a universal relationship between growth rate and the adaptive

immunity across the bacteria domain.

We used the phylogenetic generalized least squares (PGLS) regression analysis to

examine the relationship between spacer abundance and growth rate across the bacterial and

archaeal domains. A significant positive correlation was observed in the bacterial domain

when the growth rates were measured by the predicted minimal doubling times ($n = 4143$,

slope $= 1.75$, $P = 9 \times 10^{-5}$). When the growth rates were measured by the empirical minimal

doubling times, we observed a positive correlation at a marginally significant level ($n = 262$,

slope $= 1.37$, $P = 0.056$). To examine whether the empirical minimal doubling times and the

predicted minimal doubling times lead to the same conclusion, we examined the 252 bacteria

with both empirical minimal doubling times and predicted minimal doubling times. Among

these bacteria, the spacer abundance and growth rate are significantly correlated for both

cases: slope $= 10.6$, $P = 3 \times 10^{-8}$ when the growth rate was measured by predicted minimal

doubling times, and slope $= 1.56$, $P = 0.026$ when the growth rate was measured by empirical

minimal doubling times. The empirical minimal doubling times and the predicted minimal

doubling times gave the same conclusion. However, the predicted minimal doubling time

strongly correlates with the spacer abundance.

Previous studies observed that bacterial CRISPR abundance is positively correlated with

growth temperatures (6-9), with an abrupt jump at around 45°C (10). Here, using spacer

abundance as the dependent variable, the minimal doubling time and the growth temperature

as the independent variables, we performed multiple PGLS regression analyses. The results

show that the minimal doubling time and the growth temperature are independently correlated

with spacer abundance (Table 1).

In archaea, we did not observe a significant correlation between spacer abundance and

growth rate ($P > 0.100$), no matter when the growth rate was measured by the empirical

minimal doubling times ($n = 104$) or the predicted minimal doubling times ($n = 256$). Multiple

PGLS regression analyses showed that archaeal spacer abundance is not significantly

correlated with either minimal doubling time or growth temperature ($P > 0.10$ for both cases).

The authors reporting the enhancing effects of slow growth on spacer acquisition in

cultured bacterial species generally explain their observation by the mechanisms of spacer

acquisition (1, 3, 11). The mechanistic explanation could be applied to the correlation between

spacer acquisition and growth rate across the bacteria domain. Furthermore, we could put the

spacer acquisition into an ecological context. Studies on natural bacterial communities

support the existence of trade-offs between competition ability (faster growth) and

invulnerability to predation (including virus resistance) (12-14). With the limited resources

available in natural environments, the bacteria that allocate more resources to defense have to

grow slowly, and the bacteria with high competition ability allocate less resource to virus

resistance. In addition, there is evidence that prophages were more frequent in bacteria with

small minimal doubling times (15). Targeting prophages might lead to auto-immunity and

produce a selective force driving the loss of CRISPR–Cas systems or against spacer

acquisition (16, 17).

MATERIALS AND METHODS

The datasets of prokaryotic empirical minimal doubling times and predicted minimal

doubling times were retrieved from reference (18) and the EGGO database (accessed Nov 26,

2021) (19), respectively. To cover more prokaryotes, we used the optimal growth

temperatures predicted by the program Tome (version 1.0.0) (20). The phylogenetic trees and

taxonomic information of bacteria and archaea were retrieved from the Genome Taxonomy

Database (GTDB, accessed Apr 8, 2022). The prokaryotic genome assembly files were

downloaded from the GenBank database (accessed Apr 26, 2021) (21). The CRISPR-Cas

systems of the prokaryotic genomes were annotated using CRISPRCasFinder (version 4.2.20) (22). The avoid the outlier effects on the statistical results, we only retained the minimal doubling times shorter than 60h. To avoid underestimation of the CRISPR-Cas systems, partially assembled genomes have been discarded, and only those with assembly levels of "complete genome" or "chromosome" were retained in our analyses. Because there is no phylogenetic divergence information of different strains of the same prokaryotic species in GTDB (23), we used the representative genome of each species in this study. In addition, all the suppressed genomes of the GenBank database were discarded.

We estimated the phylogenetic signals (λ) of the analyzed characters using the R (Version 4.0.2) package phytools (Version 0.7-47) (24). All the analyzed characters exhibit significant phylogenetic signals, indicating that phylogenetic comparative methods are required in the analyses to control the effect of common ancestry. The correlations between different characters were estimated by the PGLS regression using the R (Version 4.0.2) package phylolm (version 2.6.2) (25). Pagel's λ model has been applied in this study.

Acknowledgments

This work was supported by the National Natural Science Foundation of China (grant number 31671321).

References

1. Amlinger L, Hoekzema M, Wagner EGH, Koskiniemi S, Lundgren M. 2017. Fluorescent CRISPR Adaptation Reporter for rapid quantification of spacer acquisition. Scientific Reports 7:10392.

2. McKenzie RE, Keizer EM, Vink JNA, van Lopik J, Büke F, Kalkman V, Fleck C, Tans

SJ, Brouns SJJ. 2022. Single cell variability of CRISPR-Cas interference and

adaptation. Molecular Systems Biology 18:e10680.

3.   Høyland-Kroghsbo NM, Muñoz KA, Bassler BL. 2018. Temperature, by controlling

growth rate, regulates CRISPR-Cas activity in *Pseudomonas aeruginosa*. mBio

9:e02184-18.

4.   Westra ER, van Houte S, Oyesiku-Blakemore S, Makin B, Broniewski JM, Best A,

Bondy-Denomy J, Davidson A, Boots M, Buckling A. 2015. Parasite exposure drives

selective evolution of constitutive versus inducible defense. Current Biology

25:1043-1049.

5.   Meeske AJ, Nakandakari-Higa S, Marraffini LA. 2019. Cas13-induced cellular

dormancy prevents the rise of CRISPR-resistant bacteriophage. Nature 570:241-245.

6.   Makarova KS, Wolf YI, Snir S, Koonin EV. 2011. Defense islands in bacterial and

archaeal genomes and prediction of novel defense systems. Journal of Bacteriology

193:6039-6056.

7.   Anderson RE, Brazelton WJ, Baross JA. 2011. Using CRISPRs as a metagenomic tool

to identify microbial hosts of a diffuse flow hydrothermal vent viral assemblage.

FEMS Microbiology Ecology 77:120-133.

8.   Weinberger AD, Wolf YI, Lobkovsky AE, Gilmore MS, Koonin EV. 2012. Viral

diversity threshold for adaptive immunity in prokaryotes. mBio 3:e00456-12.

9.   Gophna U, Kristensen DM, Wolf YI, Popa O, Drevet C, Koonin EV. 2015. No

evidence of inhibition of horizontal gene transfer by CRISPR–Cas on evolutionary

timescales. The ISME Journal 9:2021-2027.

10. Lan X-R, Liu Z-L, Niu D-K. 2022. Precipitous increase of bacterial CRISPR-Cas

155    abundance at around 45°C. Frontiers in Microbiology 13:773114.

156 11. Dimitriu T, Kurilovich E, Lapinska U, Severinov K, Pagliara S, Szczelkun MD,

157    Westra ER. 2022. Bacteriostatic antibiotics promote CRISPR-Cas adaptive immunity

158    by enabling increased spacer acquisition. Cell host & microbe 30:1-10.

159 12. Cram JA, Parada AE, Fuhrman JA. 2016. Dilution reveals how viral lysis and grazing

160    shape microbial communities. Limnology and Oceanography 61:889-905.

161 13. Yang JW, Chang F-H, Yeh Y-C, Tsai A-Y, Chiang K-P, Shiah F-K, Gong G-C, Hsieh

162    C-h. 2021. Trade-off between competition ability and invulnerability to predation in

163    marine microbes; protist grazing versus viral lysis effects. BioRxiv

164    doi:10.1101/2021.10.09.463787.

165 14. Winter C, Bouvier T, Weinbauer MG, Thingstad TF. 2010. Trade-offs between

166    competition and defense specialists among unicellular planktonic organisms: the

167    "killing the winner" hypothesis revisited. Microbiology and Molecular Biology

168    Reviews 74:42-57.

169 15. Touchon M, Bernheim A, Rocha EPC. 2016. Genetic and life-history traits associated

170    with the distribution of prophages in bacteria. The ISME Journal 10:2744-2754.

171 16. Rollie C, Chevallereau A, Watson BNJ, Chyou T-y, Fradet O, McLeod I, Fineran PC,

172    Brown CM, Gandon S, Westra ER. 2020. Targeting of temperate phages drives loss of

173    type I CRISPR–Cas systems. Nature 578:149-153.

174 17. Nobrega FL, Walinga H, Dutilh BE, Brouns SJJ. 2020. Prophages are associated with

175    extensive CRISPR–Cas auto-immunity. Nucleic Acids Research 48:12074-12084.

18.   Madin JS, Nielsen DA, Brbic M, Corkrey R, Danko D, Edwards K, Engqvist MKM,

Fierer N, Geoghegan JL, Gillings M, Kyrpides NC, Litchman E, Mason CE, Moore L,

Nielsen SL, Paulsen IT, Price ND, Reddy TBK, Richards MA, Rocha EPC, Schmidt

TM, Shaaban H, Shukla M, Supek F, Tetu SG, Vieira-Silva S, Wattam AR, Westfall

DA, Westoby M. 2020. A synthesis of bacterial and archaeal phenotypic trait data.

Scientific Data 7:170.

19.   Weissman JL, Hou S, Fuhrman JA. 2021. Estimating maximal microbial growth rates

from cultures, metagenomes, and single cells via codon usage patterns. Proceedings of

the National Academy of Sciences 118:e2016810118.

20.   Li G, Rabe KS, Nielsen J, Engqvist MKM. 2019. Machine learning applied to

predicting microorganism growth temperatures and enzyme catalytic optima. ACS

Synthetic Biology 8:1411-1420.

21.   Sayers EW, Cavanaugh M, Clark K, Pruitt KD, Schoch CL, Sherry Stephen T,

Karsch-Mizrachi I. 2021. GenBank. Nucleic Acids Research 50:D161-D164.

22.   Couvin D, Bernheim A, Toffano-Nioche C, Touchon M, Michalik J, Néron B, Rocha

EPC, Vergnaud G, Gautheret D, Pourcel C. 2018. CRISPRCasFinder, an update of

CRISRFinder, includes a portable version, enhanced performance and integrates

search for Cas proteins. Nucleic Acids Research 46:W246-W251.

23.   Parks DH, Chuvochina M, Rinke C, Mussig AJ, Chaumeil P-A, Hugenholtz P. 2021.

GTDB: an ongoing census of bacterial and archaeal diversity through a

phylogenetically consistent, rank normalized and complete genome-based taxonomy.

Nucleic Acids Research 50:D785-D794.

24.   Revell LJ. 2012. phytools: an R package for phylogenetic comparative biology (and

other things). Methods in Ecology and Evolution 3:217-223.

25.   Ho LST, Ane C. 2014. A linear-time algorithm for Gaussian and non-Gaussian trait

evolution models. Systematic Biology 63:397-408.

Table 1. Correlations of bacterial CRISPR spacer abundance with the minimal doubling time

and the growth temperature.

		MDT		Growth temperature	
	N	Slope	P	Slope	P
Spacer abundance[1]	262	1.84	0.011	2.06	0.001
Spacer abundance[2]	4142	1.66	2×10^{-4}	1.43	4×10^{-8}

207 The correlations were examined using multiple PGLS regression analyses, with the positive

208 sign of the regression slope to indicate the positive correlation. [1] Correlation of spacer

209 abundance with empirical MDT and optimal growth temperature. [2] Correlation of spacer

210 abundance with predicted MDT and optimal growth temperature. MDT, minimal doubling

211 time.