1

Disentangling the effects of eutrophication and natural variability on macrobenthic communities across French coastal lagoons

Auriane G. Jones[*][1], Gauthier Schaal[2], Aurélien Boyé[3], Marie
Creemers[1], Valérie Derolez[1], Nicolas Desroy[4], Annie Fiandrino[1],
Théophile L. Mouton[1], Monique Simier[1], Niamh Smith[1], Vincent
Ouisse[1]

[1] MARBEC, Univ. Montpellier, CNRS, Ifremer, IRD, Sète, France

[2] Univ Brest, CNRS, IRD, Ifremer, LEMAR, 29280 Plouzané, France

[3] IFREMER, Centre de Bretagne, DYNECO, Laboratoire d'Ecologie Benthique Côtière (LEBCO), 29280 Plouzané,
France

[4] IFREMER, Laboratoire Environnement et Ressources Bretagne nord, 35801 Dinard cedex, France

[*]Corresponding author

Correspondence: jones.ecology@gmail.com

ABSTRACT

26Coastal lagoons are transitional ecosystems that host a unique diversity of species and support 27many ecosystem services. Owing to their position at the interface between land and sea, they 28are also subject to increasing human impacts, which alter their ecological functioning. Because 29coastal lagoons are naturally highly variable in their environmental conditions, disentangling 30the effects of anthropogenic disturbances like eutrophication from those of natural variability is 31a challenging, yet necessary issue to address. Here, we analyze a dataset composed of 32macrobenthic invertebrate abundances and environmental variables (hydro-morphology, 33water, sediment and macrophytes) gathered across 29 Mediterranean coastal lagoons located 34in France, to characterize the main drivers of community composition and structure. Using 35correlograms, linear models and variance partitioning, we found that lagoon hydro-morphology 36(connection to the sea and lagoon surface), which affects the level of environmental variability 37(salinity and temperature), as well as lagoon-scale benthic habitat diversity (using macrophyte 38morphotypes) seemed to regulate macrofauna distribution, while eutrophication and 39associated stressors like low dissolved oxygen, acted upon the existing communities, mainly by 40reducing species richness and diversity. Furthermore, M-AMBI, a multivariate index composed 41of species richness, Shannon diversity and AMBI (AZTI's Marine Biotic Index) and currently used 42to evaluate the ecological state of French coastal lagoons, was more sensitive to eutrophication 43(18%) than to natural variability (9%), with nonetheless 49% of its variability explained jointly 44by both. To improve the robustness of benthic indicators like M-AMBI and increase the 45effectiveness of lagoon benthic habitat management, we call for a revision of the ecological 46groups at the base of the AMBI index and of the current lagoon typology which could be 47inspired by the lagoon-sea connection levels used in this study.

49*Keywords:* Mediterranean Sea, macrofauna, taxonomic diversity, M-AMBI, macrophyte morphotypes, 50variance partitioning

                                        Introduction

53   Coastal lagoons are inland, shallow and transitional water bodies, usually oriented parallel to the coast,
54separated from the sea by a barrier, connected to the sea by one or more restricted inlets which remain
55open at least intermittently (Kjerfve, 1994). These ecosystems represent 13% of the world's coastline
56(Barnes, 1980) and provide essential ecosystem services, especially through nutrient regulation and
57sequestration as well as climate regulation (Kermagoret et al., 2019; Levin et al., 2001). These ecosystems are
58characterized by strong salinity variations at different temporal scales, ranging from fortnights to seasons
59and decades (Pérez-Ruzafa et al., 2007a), that are mainly governed by short-term freshwater inflow, seasonal
60temperature variations, the degree of connection of each water mass with the sea and decadal climatic
61variations (rainfall and temperatures) (Barbone and Basset, 2010; De Casabianca, 1996; Zaldívar et al., 2008).
62Oxygen saturation naturally fluctuates in lagoons, due to water temperature and re-aeration processes
63controlled by wind conditions (Chapelle et al., 2001; Derolez et al., 2020a). Oxygen dynamics are further
64affected by primary production, which is mainly governed, in oligo and mesotrophic lagoons by seasonal
65benthic macrophyte growth and decay cycles, and in hyper and eutrophic lagoons by pelagic phytoplankton
66dynamics (Zaldívar et al., 2008). Overall, coastal lagoons are naturally highly dynamic systems in terms of
67environmental factors, representing a form of natural environmental stress for any organism living in these
68systems like fish and benthic invertebrates (Pérez-Ruzafa et al., 2007b; Zaldívar et al., 2008). Consequently,
69lagoon benthic communities, just like estuarine ones, generally present low species richness and high
70abundances of tolerant species, characteristics typical of disturbed or stressed coastal ecosystems, giving rise
71to the "transitional waters quality paradox" (Zaldívar et al., 2008), a generalization of the "Estuarine quality
72paradox" (Elliott and Quintino, 2007). These paradoxes highlight the problems faced by transitional waters
73concerning the development of methodologies permitting the evaluation of their ecological quality status,
74because it results from both anthropogenic pressures and natural characteristics.

75   Coastal lagoons have also been intensely used and modified by humans for food (fishing and
76aquaculture), transportation (canals, ports and inlet modification), livelihood (land reclamation for
77construction) and recreation (fishing, sailing, swimming) (Lasserre, 1979; Pérez-Ruzafa et al., 2011a). The
78position of coastal lagoons at the interface between land and sea further exposes them to human land-
79derived pollution such as nitrogen, phosphorous, pesticide, or heavy metal pollution (Caliceti et al., 2002;
80Derolez et al., 2019; Souchu et al., 2010). The oldest and most studied is nitrogen and phosphorous pollution
81linked to agriculture and urban development (Bendoricchio et al., 1993; Kjerfve, 1994; Lacoste et al., 2023;
82Tournoud et al., 2006), which can lead to eutrophication of coastal lagoons, a phenomenon associated to an
83increase in phytoplankton biomass (Cloern, 2001; Zaldívar et al., 2008), a replacement of long-lived
84submerged macrophytes like marine plants, by emergent free-living benthic macroalgae and even to the
85disappearance of all benthic macrophytes (Le Fur et al., 2019, 2018). The high light availability linked to
86shallow depths and the naturally high nutrient concentration of water and sediments (*i.e.* coastal lagoons can
87be considered as "naturally eutrophic" (De Casabianca, 1996; Zaldívar et al., 2008)) promote the
88development of structurally complex benthic macrophytes, which are characteristic features of coastal
89lagoons. Macrophyte presence, cover and composition create a highly heterogeneous aquatic landscape
90(Arocena, 2007; Gamito et al., 2005), which is also determined by the sediment nature (*i.e.* hard bottom,
91gravel, sand, mud) and organic matter content (Menu et al., 2019).

92   Benthic macroinvertebrates are a key biotic compartment of coastal lagoons, through their role in water
93filtration, food web dynamics, recycling of organic matter and nutrients and sediment bioturbation (Barnes,
941980; Gamito et al., 2005; Levin et al., 2001; Zaldívar et al., 2008). These benthic communities are reported
95to be shaped mainly by natural environmental variability (*i.e.* environmental filtering) and passive diffusion
96(mainly as planktonic larvae) (Basset et al., 2007). At the intra-lagoon scale, the spatial distribution of

97macrobenthic invertebrates is tightly linked to the renewal of marine-originating molecules, which is directly
98linked to the connection level with the open sea, a result used to define the notion of confinement
99(Guelorget and Perthuisot, 1983). Superimposed onto the previously mentioned "natural environmental
100stress" (*i.e.* strong fluctuations of abiotic parameters like salinity and temperature) are a number of human
101related disturbances like anthropogenic eutrophication, which can further impact lagoonal organisms (Elliott
102and Quintino, 2007; Rossi et al., 2006). Most benthic macroinvertebrates are exposed to multiple human-
103induced disturbances as they generally live for several years. Furthermore, these organisms tend to present,
104as adults, a low dispersal potential (*i.e.* below a few kilometers), meaning that faced with a disturbance, each
105individual will either resist, become accustomed or die, with overall very little chance of escaping the
106disturbance, especially if it affects the entire lagoon (Cognetti and Maltagliati, 2000). These characteristics
107make benthic macroinvertebrates good theoretical indicators of the ecological health status of coastal
108lagoons and since 2000, this biological compartment is officially included in the assessment of European
109transitional water bodies carried out within the Water Framework Directive (WFD) (Reizopolou et al., 2018).
110However, although they are in theory good ecological indicators, benthic macroinvertebrates are also
111influenced by "natural environmental stress", hence it appears crucial to disentangle the effect of natural
112environmental variability from the effect of variability linked to human disturbances on this compartment in
113hyper-variable ecosystems like coastal lagoons (Pitacco et al., 2019). Most of the published work on
114Mediterranean coastal lagoon macroinvertebrates have either been descriptive studies focusing on one or a
115few lagoons (Amanieu et al., 1977; Arocena, 2007; Bachelet et al., 2000; Carlier et al., 2007; Guelorget et al.,
1161994; Guelorget and Michel, 1979; Magni et al., 2022; Pérez-Ruzafa et al., 2007a; Rossi et al., 2006),
117theoretical oriented studies testing ecological hypotheses like environmental filtering (Basset et al., 2007) or
118species-area relations (Sabetta et al., 2007) or very applied studies related to the WFD and focusing on
119lagoon typology definition, indicator development or indicator reference values (Barbone et al., 2012; Basset
120et al., 2012; Reizopoulou et al., 2014). Disentangling the respective effect of natural vs human-induced
121disturbances on lagoon communities requires a statistical power which can be reached by using extensive
122datasets, spanning a variety of ecological conditions (*i.e.* several lagoon systems) and a thorough
123characterization of both sources of disturbance. In this study, we make use of the full potential of large-scale
124environmental surveys of the various physical and biological compartments (water column, sediments,
125macrophytes and benthic macroinvertebrates) of French Mediterranean coastal lagoons: *(i)* to disentangle
126the effects of an anthropogenic disturbance, here eutrophication, from those of natural environmental
127variability on benthic macroinvertebrates and *(ii)* to understand the links between the different
128environmental variables characterizing coastal lagoons, that affect benthic macroinvertebrates. This study,
129based on a dataset including 29 French Mediterranean coastal lagoons (ranging from oligo- to eu-haline, with
130diverse environmental and anthropic settings), focuses on multiple levels of biological complexity; first the
131broad and complex community level, then simple diversity indices like species richness and Shannon diversity
132index and finally, a multimetric and WFD-compliant index.
133

134   **Methods**

135**Study sites**
136   The French Mediterranean coast features a great diversity of lagoons characterized by different hydro-
137climatic contexts (*e.g.* temperature, wind, rainfall). Some of these lagoons are under the influence of large
138watersheds (up to 1 500 km²) while others are connected to more restricted basins that may be 300 times
139smaller (*ca.* 5 km²). These lagoons also present variable connections to the sea, with both natural vs man-
140made channels, and continuous vs temporary channels. Human land-use in this region is particularly high,
141associated to both urban and agricultural development (Derolez et al., 2019). Most lagoons in continental

142France are located near large cities like Narbonne or Montpellier and/or in areas intensively cultivated for
143wine, vegetables and cereals (CORINE Land Cover, 2018) whereas the Corsican lagoons of Palo, Urbino and
144Diana are relatively preserved from direct urban pressures. Differently, the Corsican lagoon of Biguglia is
145located just south of the urban hub of Bastia and in a growing agricultural plain (CORINE Land Cover, 2018;
146Pasqualini et al., 2017). Overall, French Mediterranean lagoons face various degrees of anthropogenic
147pressures.

148   A total of 29 nanotidal (tide < 0.5 m) coastal lagoons located in the Gulf of Lion (north-west
149Mediterranean Sea) and along the east coast of Corsica were selected for this study and sampled across 41
150stations (between one and three in each lagoon) (Figure 1). The sampling was carried out once a time in 2006
151or 2009 (depending on the station) during the "Réseau de Suivi Lagunaire" and WFD campaigns. Infaunal
152benthic macroinvertebrates were sampled in May, except four lagoons sampled at the end of April (La
153Marette, Ponant, Berre and Vaine) and one lagoon at the end of June (Bolmon). The studied lagoons
154encompass a wide range of hydro-morphological characteristics in terms of surface, depth and connection to
155the sea (Table 1) with the lagoon-sea connection level expressed as weak, moderate or strong depending on
156the differences in salinity variations between the lagoon and a neighboring marine station (Derolez et al.,
1572012). Lagoon surfaces vary from 1 km^2 (Campignol) to 133 km^2 (Berre) (Menu et al., 2019) and station depth
158is comprised between 0.1 m (La Palme south station) and 9.5 m (Urbino) with most stations less than 3 m
159deep (Table 1).

Figure 1 - Map of the 41 stations sampled in the 29 lagoons located along the French Mediterranean coast (Gulf of Lion and Corsica). Each station's salinity type (according to *their median salinity and Venice classification (Anonymous, 1958) in* Le Fur et al. (2018)) and group membership (K-means partitioning) based on the benthic macrofauna Hellinger-transformed mean abundances are indicated using a symbol (triangle: oligo- and meso-haline and circle: poly- and eu-haline) and a color (red: group A and blue: group B), respectively (see section *Macrobenthic community structure* for the explanation on this analysis). See Table 1 for the full lagoon names and corresponding stations.

The different hydro-morphological characteristics of the studied lagoons lead to a great diversity of
salinity regimes with 8 oligo- or meso-haline lagoons and 21 poly- or eu-haline lagoons (according to Le
Fur et al., (2018), Figure 1) and to large discrepancies between lagoons in terms of annual variability of
temperature, salinity and oxygen saturation. Overall, the physico-chemical status of the studied lagoons
between 2004 and 2009 was evaluated according to the WFD methodology, as poor (15 lagoons), bad (3
lagoons), moderate (4 lagoons) or good (7 lagoons) (Table 1). This evaluation takes into account the
concentrations of total and dissolved inorganic forms of nutrients, turbidity and oxygen saturation during
the summer months of the 2004-2009 period and is conducted according to the WFD quality grid (Andral
and Sargian, 2010a, 2010b).

Table 1 - Hydro-morphological characteristics (surface, depth and connection level to the sea) and
water physico-chemical status of the 29 lagoons as evaluated by the water framework directive
(green = good, yellow = moderate, orange = bad, red = poor). The physico-chemical status is based
on concentrations of total and dissolved inorganic forms of nutrients, turbidity and oxygen
saturation during the summer months of the 2004-2009 period (Andral and Sargian, 2010a; Andral
and Sargian, 2010b). Depth, sediment organic matter content (OMC, % of dry sediment weight),
beta diversity of macrophyte morphotypes (β_{MM}, no unit – see section *Estimating the macrophyte*
morphotype beta diversity) of the 41 stations sampled for benthic macrofauna either in 2006 (#) or
in 2009 (*) is also indicated. For stations sampled in the same lagoon, each one is labeled using a 3
letter code with the last letter indicating the geographical location of the station in the lagoon
(North (N), Middle (M), South (S), East (E), West (W), except THP which indicates the station's local
name Pisse-Saumes). The lagoons are listed from west to east along the Gulf of Lion (see Figure 1
for their geographical location).

Lagoon characteristics				Benthic station characteristics			
Lagoon	Surface (km²)	Lagoon-sea connection	Physico-chemical status	Station code	Station depth (m)	OMC (%)	β_{MM}
Canet*	6.43	weak	🟥	CAN	0.9	5.6	0.13
Leucate*	52.88	strong	🟩	LEN	1.9	8.2	0.71
				LES	3.3	7.2	0.67
La Palme#	6.94	moderate	🟩	LPN	0.6	4.68	0.6
				LPS	0.1	1.61	0.53
Bages-Sigean*	37.05	moderate	🟩	BGN	1.9	11.5	0.65
				BGM	2.2	5.9	0.8
				BGS	1.3	7.1	0.53
Campignol*	1	weak	🟥	CAM	0.8	3.8	1
Gruissan*	1.31	strong	🟩	GRU	0.9	3.4	0.61

Vendres*	6.6	weak		VEN	0.8	22.8	0.6
Grand Bagnas#	1.69	weak		GBA	0.5	5.38	0
Thau#	67.7	strong		THE	8.2	11.38	0.55
				THW	4	9.9	0.68
				THP	3.8	5.64	0.68
Ingril North#	2.28	strong		INN	1.1	12.48	0.4
Ingril South#	3.17	strong		INS	1.2	7.01	0.25
Vic#	12.96	moderate		VIC	1.5	13.05	0.69
Pierre-Blanche#	3.14	moderate		PBE	0.7	1.04	0.33
				PBW	0.7	9.93	0.67
Arnel#	4.42	moderate		ARN	0.8	7.88	0.29
Prévost#	2.46	strong		PRE	0.7	9.25	0.4
				PRW	0.7	8.4	0.6
Méjean#	7.12	moderate		MEE	1	7.67	0
				MEW	0.9	12.17	0
Grec#	1.08	moderate		GRE	0.7	8.49	0.33
Or (Mauguio)*	31.31	weak		ORE	1.4	10.9	0.75
				ORW	1.3	12.5	0.54
Ponant*	1.92	moderate		PON	3	10.9	0.11
La Marette*	1.3	weak		MRN	0.7	15.9	0
				MRS	0.6	14.2	0
Vaccares*	65.9	weak		VAC	1.9	5.8	0.23
La Palissade*	2.95	weak		PLS	0.6	1.5	0.6
Berre*	132.71	strong		BRN	5.9	11.8	0.39

			🟩	BRS	8.9	14.5	0.38
Vaine*	22.77	moderate	🟨	VAI	5.1	12.7	0.73
Bolmon*	6.01	weak	🟥	BOL	1.4	14.8	0.30
Biguglia*	13.75	weak	🟧	BIG	1	7.9	0.25
Diana*	5.4	strong	🟩	DIA	5.8	10.5	0.39
Urbino*	7.58	moderate	🟨	URB	9.5	16.3	0.63
Palo*	1.08	weak	🟨	PAL	1	12.5	0

190 **Sampling**

191 *Benthic macroinvertebrates*

192 At each of the 41 stations, three replicates (*ca.* 10 m apart) of unvegetated sediment were sampled. 193 Each replicate was made up of four pooled Ekman-Birge grabs, representing a total surface of 0.09 m^2 per 194 replicate and 0.27 m^2 per station. Samples were sieved on a 1 mm mesh, fixed in a 5 to 7% formaldehyde/ 195 seawater solution and stained with Rose Bengal for latter identification. The sediments were then sorted 196 in the laboratory and all organisms identified if possible to the species level, following the nomenclature 197 of the European Register of Marine Species (ERMS). Taxa are indicated using WORMS reference (https:// 198 www.marinespecies.org/aphia.php?p=match). Prior to the data analyses, we removed meiofauna taxa 199 (Ostracoda, Nematoda, Harpacticoida and Collembola) and organisms present in only one lagoon and 200 determined at the order (Decapoda, Coleoptera, Nudibranchia), class (Insecta, Ascidiacea) or subclass 201 (Oligochaeta) level. We kept Actiniaria, Nemertea and Polycladida as they were identified across multiple 202 lagoons.

203 *Sediment parameters*

204 At each station, an additional grab sample was collected to evaluate sediment organic matter and 205 grain-size distribution. Sediment samples were sieved on a 2 mm mesh to remove large macrophytes 206 debris, and the filtrate was weighted and dried until stable weight (110°C for 24 to 48h) in the laboratory. 207 Organic matter content (OMC, % of dry sediment weight) was calculated as the loss of weight after 208 combustion for 12h at 450°C of the dry sediment fraction below 2 mm. Because of technical problems, no 209 OMC was available for the stations ARN, LPN, LPS, PRW and VIC in 2006. To estimate these missing 210 values, we used OMC values measured in spring of 2006 (ARN, PRW, VIC) or spring of 2007 (LPN and LPS) 211 for the WFD sediment quality evaluation. For ARN, LPS, LPN and VIC, we estimated OMC by averaging 212 over the values measured in the three or four stations closest to the macrofauna station (distance 213 between 285 m and 707 m). For PRW, we used the OMC value measured at the closest station as this 214 station was located only 114 m from the macrofauna station (Supplementary material 1 and 2). Since 215 laboratory protocols changed between years (2006 and 2009) and between lagoons regarding the 216 sediment particle sizes considered for the grain-size analysis, we chose not to consider grain-size 217 parameters.

218*Water physico-chemical properties*

219   Water column physico-chemical properties of the 29 lagoons are measured monthly during summer
220(from June to August or September) at least since 2003, following a standardized protocol implemented
221by the WFD (Derolez et al., 2019). Salinity (Sal), temperature (Temp) and oxygen saturation (O_2 sat) were
222measured using a portable field probe (WTW LF 197, WTW-GmbH, Weilheim, Germany) and turbidity
223(Turb) was measured in the laboratory using an optical turbidimeter (2100N IS turbidimeter ISO 7027).
224Chlorophyll *a* concentration ([Chla]) was measured in 50 mL samples of water, first filtered on a
225Whatman GF/F filter (0.7 μm). Pigments were later extracted in the laboratory in a 90% acetone +
226distilled water solution during 24 hours (4°C, dark conditions after 10 minutes of centrifugation). The
227supernatant was then dosed using a spectrofluorometer following Neveux and Lantoine (1993). The full
228description of the protocol followed to measure [Chla], total nitrogen concentration [TN], ammonium
229concentration [NH_4^+], sum of nitrate plus nitrite concentrations [NO_x], total phosphorus concentration
230[TP] and dissolved inorganic phosphorus concentration [DIP] is described in Bec et al. (2011) and Souchu
231et al. (2010). Only sub-surface data were available for all stations except the THW, THE, BRN, BRS stations
232located in the deep Thau and Berre lagoons for which sub-surface and bottom data were available. In
233these four cases, we chose to consider the bottom data.

234*Estimating the macrophyte morphotype beta diversity*

235   Lagoons are characterized by a mosaic of benthic habitats, determined in particular by macrophyte
236diversity which can affect benthic macroinvertebrates (Arocena, 2007; Cardoso et al., 2004; Magni and
237Gravina, 2023; Ouisse et al., 2011) by acting as food (Jones et al., 2021; Quillien et al., 2016; Vizzini and
238Mazzola, 2006), refuge (Mancinelli and Rossi, 2001; Nordström and Booth, 2007; Ware et al., 2019)
239and/or means of transport (Norkko et al., 2000). Although sampling was carried out on locally
240unvegetated sediments to avoid biases linked to macrophytes and associated faunal assemblages, we
241postulate that the presence and composition of macrophyte assemblages within each lagoon can affect
242the composition of infaunal benthic communities. We developed an index (later beta diversity of
243macrophyte morphotypes, β_{MM}) calculated as the average Bray-Curtis dissimilarity at the lagoon or sub-
244lagoon scale (when several macrofauna stations were sampled in one lagoon) based on a station by
245macrophyte morphotype matrix (presence/absence). Each macrophyte station is coded as bare sediment
246if the macrophyte cover is less than 25% or as one or several of seven macrophyte morphotypes (*i.e.*
247mushroom, feather, hair, ball, branched, sheet and meadow) based on the thallus morphology of the
248dominant or co-dominant macrophyte taxa. This index varies between 0 if all the considered stations are
249dominated or co-dominated by the same macrophyte morphotype(s) or if they all have bare sediment
250and 1 if all the considered stations are dominated or co-dominated by different macrophyte morphotypes
251(Appendices 1 and 2). We gathered the data used to calculate this index from the WFD macrophyte
252dataset (all information regarding sampling protocols are available in Le Fur et al., 2017).

253**Data analyses**

254   We extracted all the data used in this study from the French database Quadrige (Ifremer, http://
255quadrige.eaufrance.fr/). All the raw data used in this study (macrofauna, environmental data, GIS data)
256are available online along with all the scripts and code used to analyze the data (Jones et al., 2023). We
257performed all analyses using R version 3.6.3 (R Core Team, 2020) and the packages vegan (*v2.5-6*;
258Oksanen et al., 2019), missMDA (Josse and Husson, 2016), labdsv (*v2.0-1*; Roberts, 2019), olsrr (v0.5.3;
259Hebbali, 2020), MASS (Venables and Ripley, 2002), corrplot (*v0.84*; Wei and Simko, 2017) and rstatix
260(v0.6.0; Kassambara, 2020). We considered a significance level of 0.05 for all statistical tests.

Environmental variables

262   Water column parameters were measured in summer whereas macrofauna was sampled in spring
263which prevented us from relating these two datasets. Nonetheless, the presence and abundance of
264macroinvertebrates in a lagoon year y is influenced not only by the environmental conditions prevailing
265that year but also those prevailing the previous years especially for pluri-annual species. As such, we
266associated the macrofauna data collected year y to the mean and coefficient of variation (CV) of the
267water column parameters measured years y-2, y-1 and y, as done by Le Fur et al. (2019) (Table 2).

268   We classified the parameters presented in Tables 1 and 2 into two categories. The first category
269included 11 physical and biotic factors used as a proxy for the natural variability of environmental
270conditions (lagoon surface, lagoon-sea connection, station depth, β_{MM}, OMC, mean and CV of Sal, Temp,
271O_2 sat). The second category included 14 biogeochemical variables that either cause eutrophication (*i.e.*
272mean and CV of [TN], [NO_x], [NH_4^+], [TP], [DIP]) or are the direct consequence of eutrophication (*i.e.* mean
273and CV of [TN], [TP], [Chla], turbidity) (Souchu et al., 2010). Because of technical problems, no [NO_x] and
274[DIP] data were available for the stations BOL and PLS. Consequently, we imputed the missing mean and
275CV values for these two parameters at both stations with the 'imputePCA' function. This function
276estimates the missing parameters by first performing a principal component analysis on all the
277environmental variables and then by using the regularized fitted matrix. These steps are performed
278iteratively until convergence is reached (Josse and Husson, 2012).

Table 2 - Environmental parameters and their respective code used to characterize the water. For
each water parameter, the mean and coefficient of variation (CV) were calculated between the
years y-2 and y (with y the macrofauna sampling year) and the overall median, minimal and
maximal values across the 41 stations are indicated. The station with the minimal and maximal
value of each parameter is also indicated between brackets, as well as the transformation applied
to each parameter. See Table 1 for the station codes.

Parameter (unit)	Code	Measure	Median	Min	Max	Transf.
Salinity (PSU)	Sal	mean	32.39	2.83 (PLS)	42.30 (PBW)	/
		CV	13.52	2.49 (DIA)	82.24 (CAM)	$\log_e$
Oxygen saturation (%)	O_2 sat	mean	100.88	56.61 (BRN, BRS)	190.40 (BOL)	$\log_e$
		CV	27.38	0.82 (PLS)	67.33 (PRE)	$\sqrt{}$
Temperature (°C)	Temp	mean	24.08	19.63 (BRN, BRS)	27.09 (URB, BIG)	/
		CV	9.05	1.82 (CAN)	22.81 (PON)	$\sqrt{}$
Chlorophyll *a* concentration (µg.L^{-1})	[Chla]	mean	9.43	0.70 (BGS)	144.20 (MEW)	$\log_e$
		CV	71.04	25.42 (BOL)	178.73 (BIG)	$\log_e$
Total nitrogen concentration (µmol.L^{-1})	[TN]	mean	52.05	11.08 (DIA)	266.03 (MEW)	$\log_e$
		CV	27.77	6.94 (VAC)	89.47 (BIG)	$\log_e$

Ammonium concentration (µmol.L^{-1})	[NH$_4^+$]	mean	1.47	0.06 (PLS)	31.62 (GRE)	log$_e$
		CV	106.88	28.65 (CAN)	257.93 (THE)	√
Nitrate plus nitrite concentration (µmol.L^{-1})	[NO$_x$]	mean	0.37	0.06 (GBA)	22.81 (CAM)	log$_e$
		CV	94.87	43.30 (CAN)	227.37 (URB)	log$_e$
Total phosphorous concentration (µmol.L^{-1})	[TP]	mean	2.51	0.64 (LEN)	31.19 (CAN)	log$_e$
		CV	36.75	8.91 (VAC)	106.12 (ARN)	log$_e$
Phosphate concentration (µmol.L^{-1})	[DIP]	mean	0.24	0.02 (LES)	14.53 (CAN)	log$_e$
		CV	93.10	40.97 (THW)	187.55 (INN)	log$_e$
Turbidity (NTU)	Turb	mean	5.20	1.14 (BGS)	99.65 (PLS)	log$_e$
		CV	68.30	1.92 (PLS)	207.54 (ARN)	√

285Transformation (Transf.): / = no transformation, log$_e$ = natural logarithm, √ = square-root

286*Macrobenthic community structure*

287   First, we calculated the mean macroinvertebrate abundance at the station scale by averaging over 288the three replicates and transformed it using a Hellinger transformation, to give more weight to common 289taxa and less weight to highly abundant, rare and under-sampled taxa (Legendre and Legendre, 2012). 290We then classified the 41 stations based on these transformed mean abundances, using K-means 291partitioning and the Calinski-Harabasz criterion with 100 iterations ('cascadeKM' function). Our goal was 292to highlight similarities between lagoons based on common species without focusing on rare taxa, as 293each coastal lagoon can present unique species assemblages (Basset et al., 2007), and to help with the 294result interpretation thanks to a simple classification. We tested for significant differences in the 295composition of our macrofauna groups using the analysis of similarity (ANOSIM) test with 9999 296permutations and Euclidean distances ('anosim' function). Finally, we characterized each group using the 297indicator species metric ('indval' function, 9999 permutations). A high value of this metric for a given 298taxon associated to a given group means this taxon is consistently more present (fidelity) and abundant 299(specificity) in the stations composing that group than in stations composing the other groups (Dufrêne 300and Legendre, 1997). For this last analysis, we adjusted p-values using the False Discovery Rate correction 301method (FDR; Benjamini and Hochberg, 1995).

302*Taxonomy-based indices*

303   To characterize the macrobenthic community structure, we calculated the commonly used M-AMBI 304(Multivariate AZTI Marine Biotic Index), the three indices composing the M-AMBI - species richness (SR), 305Shannon diversity index (H') and AMBI (Muxika et al., 2007) - plus the inverse of Simpson's index (N2) and 306the total macrofauna density. The AMBI is an index based on the classification of benthic species into 307ecological groups, from the most sensitive to disturbances (i.e. organic matter enrichment) to the most 308opportunistic ones (Borja et al., 2000). The raw (i.e. untransformed) macrofauna abundances were used 309to calculate all these indices. For AMBI and M-AMBI calculations, we removed high order taxa without an 310ecological group value (Anemonia and Polycladida), planktonic (Mysidae) and hard-substrate taxa

311(Serpula and Serpulidae) (Borja et al., 2005). We computed M-AMBI using the R function available in
312Sigovini et al. (2013) and we defined the good reference conditions as the highest values of SR and H' (46
313and 3.73, respectively) and lowest value of AMBI (0.31) measured in the dataset, irrespective of the WFD
314physico-chemical status described above. All indices were computed at the replicate level ; the mean and
315standard deviation (SD) values were then calculated for each station. Finally, we tested for differences in
316total macrofauna density, SR, H', N2, AMBI and M-AMBI between previously defined groups of stations
317(see section *Macrobenthic community structure*) using non-parametric Kruskal-Wallis rank sum tests
318(Breslow, 1970).

319*Linking macrobenthic communities and environmental variables*
320   To assess how natural environmental variations correlated with anthropogenic pressures
321(explanatory variables) and how each explanatory variable correlated with macrobenthic indices, we built
322a correlogram showing the significant Spearman correlations ('cor_pmat' function) between all the raw
323variables. We also tested for differences in the variables considered (explanatory variables and
324macrobenthic indices) between lagoon-sea connection levels using non-parametric Kruskal-Wallis rank
325sum tests (Breslow, 1970), followed by pairwise Wilcoxon tests for multiple comparison, while adjusting
326p-values with FDR (Benjamini and Hochberg, 1995). Then, we transformed (natural logarithm or square
327root transformation), when necessary, the environmental variables and macrobenthic indices to
328approximate normal distributions and reduce data spread, before standardizing all the explanatory
329variables to facilitate the interpretation of the multiple linear regression coefficients (Schielzeth, 2010).
330We $\log_e$-transformed the lagoon surface, station depth, SR, N2 and macrofauna density (see Table 2 for
331the rest of the transformations). To limit multicollinearity and avoid artificially inflating the fit of linear
332models, we reduced the number of environmental variables (see section *Environmental variables*) for all
333future analyses by considering Pearson correlations < $|0.6|$, corresponding to a p-value < 5.10^{-5} and
334leading to 19 remaining environmental variables.
335   To determine which environmental variables most strongly influence lagoon macrobenthic
336communities, we performed a redundancy analysis (RDA, 'rda' function) between the 19 previously
337selected environmental variables and the Hellinger-transformed mean taxa abundances. We computed
338the variance inflation factor (VIF, 'vif.cca' function) to check for multicollinearity and sequentially
339removed variables with a VIF > 5 (James et al., 2013). We verified the significance of the full RDA model
340using the 'anova.cca' function (1000 steps) and we performed a forward model selection procedure
341('ordiR2step' function) based on the adjusted R2 of the full model and a p-value of 0.05 (9999
342permutations) (Blanchet et al., 2008). We then computed the marginal effect of each selected
343explanatory variable as the adjusted R^2 of the constrained ordination containing only the given variable as
344the predictor. The conditional effect of a variable was equal, during the forward selection procedure, to
345the additional amount of variance in species communities explained by the corresponding variable at the
346time it was included into the model. Finally, we performed a partial RDA using the variables previously
347selected, to isolate the impact of eutrophication from natural environmental variability on lagoon
348macrobenthic communities (Legendre and Legendre, 2012).
349   In the same way as for the constrained multivariate analysis, we implemented a multiple linear
350regression approach (without any interactions between variables) to investigate which variables related
351to eutrophication and natural variability most strongly influenced the macrobenthic indices (SR, H', N2,
352total macrofauna density, AMBI and M-AMBI). As there were still 19 potential explanatory variables, we
353performed a preliminary selection by including all the quantitative variables which alone had a significant
354effect on the index as first or second-degree orthogonal polynomial and the lagoon-sea connection if at
355least one of the three categories (weak, moderate, strong) had a significant effect on the index ('lm'
356function). We included second-degree polynomials to account for potential non-linear relations (Pérez-

357Ruzafa et al., 2007b) and verified that the selected variables had VIFs inferior to 5. We also checked the
358model hypotheses using the residuals vs fitted values plot for linearity, the Shapiro-Wilk test for residual
359normality, the Breush Pagan test for residual homoscedasticity ('ols_test_breusch_pagan' function) and
360the 'acf' function for the detection of residual spatial autocorrelation based on the station coordinates.
361Then, we ran a stepwise model selection using the 'stepAIC' function set in both directions and the BIC
362criterion. Finally, we checked the hypotheses of the selected model as previously explained, we
363investigated potential high leverage points (Cook's distance > 0.5) and we reran the selected model
364without these points to verify its robustness. To compare the relative effect of each selected explanatory
365variable on a given macrobenthic index, we expressed the slope estimates of the standardized variables
366as effect sizes when the variable was quantitative and as mean standardized effects when the variable
367was categorical (Schielzeth, 2010).

368   Lastly, we implemented a variance partitioning procedure using the 'varpart' function, to assess how
369much community variation or macrobenthic index variation could be: (i) uniquely attributed to
370eutrophication when the effect of natural variability had been removed, (ii) uniquely attributed to natural
371variability when the effect of eutrophication had been removed or (iii) jointly attributed to both effects
372(Legendre and Legendre, 2012). Prior to this analysis, we reran the variable selection procedures
373implemented at the community and macrobenthic index levels considering alternatively only the
374eutrophication variables or only the variables related to natural variability. We then included the
375variables selected by the aforementioned method in the variance partitioning, which was based on
376partial RDAs or on partial linear regressions and we presented the results using the adjusted R^2 values.

377   **Results**

378**Inter-lagoon environmental variability**
379   The dataset considered in this study encompasses a wide diversity of lagoons in terms of physico-
380chemical state according to the WFD and environmental parameters associated to natural variability and
381eutrophication (Tables 1 and 2). First, only lagoons with a strong (Leucate, Gruissan, Thau, Berre and
382Diana) or a moderate (La Palme and Bages-Sigean) connection to the sea were in a good physico-chemical
383state (green in Table 1). All the stations sampled in these lagoons presented mean [Chla] below 7 µg.L^{-1}
384and all of them except the ones located in the deep Berre and Diana lagoons (station depth > 5.8 m)
385presented a high beta diversity of macrophyte morphotypes (β_{MM} > 0.53). Five stations sampled in these
386lagoons presented the lowest value for a eutrophication related variable (DIA: mean [TN], BGS: mean
387[Chla] and mean Turb, LEN : mean [TP], LES: mean [DIP], THW: CV [DIP]). However, the THE station (Thau)
388had the most variable [NH$_4^+$] and the BRN and BRS stations (Berre) had the lowest mean O$_2$ sat of the
389dataset. Seven lagoons were in a moderate (yellow in Table 1) or in a bad physico-chemical state (orange
390in Table 1) and were either moderately (Ponant, Vaine, Urbino) or weakly (Grand Bagnas, Vaccarès,
391Biguglia and Palo) connected to the sea. In this group, the GBA and VAC stations respectively had the
392lowest mean [NO$_x$] and the lowest CV [TN] of the dataset whereas the URB and BIG stations respectively
393had the highest CV [NO$_x$] and CV [Chla]. All the remaining weakly connected lagoons were in a poor
394physico-chemical state (Canet, Campignol, Vendres, La Marette, La Palissade, Bolmon). In this group, the
395CAN station had both the highest mean [TP] and [DIP] and the lowest CV [NH$_4^+$] and [NO$_x$] of the dataset,
396while the CAM and BOL stations presented respectively the highest mean [NO$_x$] and CV [Chla] of the
397dataset. Finally, all the lagoons between the cities of Sète and Montpellier (Figure 1) were in a poor
398condition, despite being strongly (Ingril North, Ingril South, Prévost) or moderately (Vic, Pierre-Blanche,
399Arnel, Méjean and Grec) connected to the sea. All the stations located in these lagoons except PBE, had
400OMC above 7% and up to 13%. In this group, the MEW station presented the highest mean [Chla] and

401[TN] of the dataset while the GRE station had the highest mean [NH$_4^+$], the ARN station the highest CV
402[TP] and Turb and the INN station the highest CV [DIP] of the dataset.

403   Based on the significant Spearman correlations between the environmental variables (Figure 2),
404stations in the larger lagoons were significantly deeper, cooler in summer and less variable in terms of
405summer salinity, presented higher β$_{MM}$ and lower values for 8 eutrophication related variables (*e.g.* mean
406[Chla], mean [NO$_x$] and CV [NO$_x$]). The deepest stations were also the ones with the highest OMC. Two
407variables related to natural environmental variability were also significantly different according to the
408lagoon-sea connection level (KW test, *p* < 0.05 and post-hoc pairwise Wilcoxon tests *p* < 0.05): mean Sal
409(moderate > weak and strong > weak) and CV Sal (weak > moderate > strong).

Figure 2 - Correlogram showing only significant positive (red toned circles) and negative (grey toned
circles) Spearman correlations (*p* < 0.05) between the untransformed taxonomic indices (in black,
SR: species richness, H': Shannon diversity, N2: inverse of Simpson's dominance, M-AMBI:
Multivariate AMBI), the untransformed environmental variables related to the lagoon's natural
variability (in blue, β$_{MM}$: beta diversity of the macrophyte morphotypes, OMC: sediment organic
matter content) and the untransformed environmental variables related to eutrophication
expressed as means and coefficient of variations (in green, [Chla]: chlorophyll *a* concentration, [TN]:

total nitrogen concentration, [NH$_4^+$]: ammonium concentration, [NO$_x$]: nitrate plus nitrite
concentration, [TP]: total phosphorous concentration, [DIP]: dissolved inorganic phosphorous
concentration) calculated across 41 stations located in 29 lagoons along the French Mediterranean
coast.

Macrofauna communities and macrobenthic indices across French Mediterranean lagoons

A total of 42 465 benthic macroinvertebrates were counted and identified across all samples,
belonging to 227 different taxa with most organisms identified to species level (74.5%), others to genus
(17.2%), family (2.6%), order (2.6%), subclass (0.4%), class (1.8%) or phylum (0.9%). Across the 227 taxa,
1.3% were identified in over 50% of the stations, 90% in less than eight stations and 40% in only one
station. No taxon was present in all 41 stations. Polychaetes were by far the most diversified group (102
taxa), followed by Malacostraca crustaceans (54 taxa), bivalves (25 taxa) and gastropods (19 taxa).
Among the three taxa with a frequency of occurrence above 50%, two were polychaete species (*Hediste*
diversicolor (66%) and *Heteromastus filiformis* (51%)), and the third was the bivalve *Cerastoderma*
glaucum (51%). These three species also presented an important inter-station variability in abundance.
For all the following analyses, we considered 217 taxa (see section *Benthic macroinvertebrates*), except
for the AMBI and M-AMBI calculation, for which we considered only 213 taxa (see section *Taxonomy-*
based indices).

The K-means clustering delineated two groups of stations with distinct macrobenthic communities
(minimal difference in Calinsky-Harabasz criterion = 1.4), a result confirmed by the analysis of similarity
test (ANOSIM statistic = 0.51, *p* < 0.001). These groups did not correspond to any obvious spatial or
salinity pattern, as exemplified by the four Corsican stations and the seven oligo-mesohaline stations
which were classified in both groups (Figure 1). The polychaete *Hediste diversicolor* (*p* = 0.022) was an
indicator taxa of group A stations (red in Figure 1) along with the bivalve *Abra segmentum* (*p* = 0.022),
Chironomidae (*p* = 0.029) and the amphipod *Gammarus aequicauda* (*p* = 0.038). Conversely, group B
stations (blue in Figure 1) were significantly characterized by the bivalve *Scrobicularia cottardii* (*p* =
0.048), closely followed by another bivalve, *Loripes orbiculatus* (*p* = 0.065).

Finally, we found significant differences among the two groups in terms of SR and M-AMBI; group A
was composed of stations with lower SR and weaker M-AMBI values than group B (SR: $\chi^2(1)$ = 6.41, *p* =
0.011; M-AMBI: $\chi^2(1)$ = 5.76, *p* = 0.016). Group B was also characterized by stations with higher H' and N2
values and higher maximal macroinvertebrate densities than stations composing group A, albeit these
tendencies were not significant (Appendix 3).

Relationship between macrofauna communities and environmental variables

In all the following analyses, six variables were not considered because they were highly correlated
with another variable (|r| ≥ 0.6 ; *p* < 5. 10^{-5}): station depth (to lagoon surface), CV Sal (to mean Sal), mean
[TN], mean [TP], mean Turb (to mean [Chla]) and CV [TP] (to CV [TN]). Redundancy analysis (RDA) was
used to relate the macrofauna communities recorded within the lagoons to their natural environmental
variability and their eutrophication levels (Figure 3). Mean Sal was also not considered because of a VIF >
5 linked to significant differences between lagoon-sea connection levels (see section *Inter-lagoon*
environmental variability). The final RDA model, determined using a forward model selection procedure,
explained 22% of the inter-station community variability (model adj. R^2) and included six variables related
to natural environmental variability (lagoon-sea connection level (*p* < 0.001), mean Temp (*p* < 0.001),
lagoon surface (*p* = 0.0076), CV Temp (*p* = 0.013), mean O$_2$ sat (*p* = 0.026) and OMC (*p* = 0.041) and two
eutrophication related variables (mean [Chla] (*p* < 0.001) and mean [DIP] (*p* = 0.024)). Regarding the
individual effect of environmental variables, mean [Chla] (7.22%), lagoon surface (5.90%), lagoon-sea
connection (5.30%) and mean Temp (5.04%) were the most influential variables according to their

464marginal effect. Based on conditional effects, the most influential variables were mean [Chla] (7.22%),
465lagoon-sea connection (4.64%), mean Temp (3.62%) and lagoon surface (1.69%) (Appendix 4).

466   The first two axes accounted for 33.8 and 17.8% of the RDA model variance, respectively. First, the
467stations from group A and B were clearly separated along the first axis of the RDA, except for LPN, LPS,
468PLS and INS (Figure 3a). On the ordination of the environmental variables (Figure 3b), lagoon surface and
469CV Temp grouped on the positive side of the first axis while mean [Chla], mean [DIP] and mean Temp
470grouped on the negative side of the first axis. Consequently, group B macrobenthic communities were
471linked to large lagoons (also relatively deep), several of which presented high summer temperature
472variations. They were characterized by higher relative abundances of the bivalves *S. cottardii* and *L.*
473*orbiculatus* (Figure 3c). Conversely, group A communities were associated to smaller lagoons, subject to
474higher mean summer temperatures and high eutrophication levels. They were characterized by higher
475relative abundances of the polychaete *H. diversicolor* and Chironomidae (Figure 3c).

476   The second axis mainly corresponded to a confinement and also to an organic matter content
477gradient, mostly separating each group into two sub-groups with the lagoons weakly connected to the
478sea presenting sediments enriched in organic matter located towards the positive side of the second axis
479and the moderately and strongly connected lagoons located towards the negative side of the second axis.
480As mean Sal and CV Sal were correlated to lagoon-sea connection level (see section *Inter-lagoon*
481*environmental variability*), this axis also corresponded to a salinity gradient with low and variable salinity
482stations characterized by a higher

Figure 3 - Redundancy analysis plots (RDA, axis 1 and 2) focusing on (a) the benthic macrofauna sampled across 41 stations located in 29 lagoons along the French Mediterranean coast (see Table 1 for the full lagoon names and corresponding stations) as explained by the set of environmental variables related to eutrophication and natural variability composing the most parsimonious model,

(b) the selected environmental variables, where mean [Chla], lagoon surface, mean O_2 sat and mean [DIP] were $\log_e$ transformed and CV Temp was square-root transformed (see Tables 1 and 2 for the full environmental variable names and values) and (c) the macrofauna taxa, where only taxa with a goodness of fit (cumulative proportion of variance explained by the first two axis) above 0.25 are plotted. In (a), inter-station distances are euclidean while in (b) and (c) angles between all vectors reflect linear correlations. Across all plots, each station (point) is represented using a symbol for its salinity type (triangle: oligo- and meso-haline and circle: poly- and eu-haline) and a color (red: group A and blue: group B) for its group membership (K-means partitioning) based on Hellinger-transformed mean abundances.

497 abundance of *Chironomus salinarius* and *Streblospio shrubsolii* (Figure 3c). All the oligo- and meso-haline 498 stations (triangles on Figure 3) grouped on the upper part of the RDA space and most poly- and eu-haline 499 stations (circles on Figure 3) grouped on the lower part of the figure. Furthermore, communities of 500 lagoons weakly connected to the sea were characterized by higher abundances of the bivalve 501 *Scrobicularia cottardii* for group B, while communities from lagoons moderately to strongly connected to 502 the sea were characterized either by higher abundances of *L. orbiculatus* for group B or by a diversified 503 suite of species like the decapod *Carcinus aestuarii* or the polychaete *Capitella capitata* for group A.

504 Once the effect of natural environmental variability on macrofauna communities was removed 505 through partial RDA and the only explanatory variables left to vary were mean [Chla] and mean [DIP] 506 (eutrophication variables), no differences could be made between the communities of groups A and B 507 (Appendix 5). This partial RDA explained only 6.4% of the inter-station community variability (model adj. 508 R^2).

510 **Relationship between macrobenthic indices and environmental conditions**

511 Based on the significant Spearman correlations between the raw environmental variables and raw 512 macrobenthic indices (Figure 2), SR, H', N2 and M-AMBI were significantly higher at stations where mean 513 Temp was lower, salinity was more stable and eutrophication related variables were lower, especially 514 mean [Chla]. Furthermore, H', N2 and M-AMBI were significantly higher at stations where mean Sal and 515 β_{MM} were higher while SR and M-AMBI were also significantly higher at deeper stations located in larger 516 lagoons. AMBI appeared to be significantly correlated positively only with mean [NO_x] while total 517 macrofauna density was significantly higher at stations with lower but more variable summer water 518 temperatures, lower salinity and lower mean [DIP].

519 The results of the multiple linear regressions identifying which variables influenced the macrobenthic 520 indices are presented in Table 3. The most parsimonious models contained between two and four 521 predictors and explained between 27% and 74% of the macrobenthic indices' variability. All the models 522 contained at least one predictor associated to eutrophication and at least one predictor associated to 523 natural variability except the N2 and AMBI models, which only contained predictors associated to natural 524 variability and eutrophication respectively. Lagoon-sea connection was selected in three models, H', N2 525 and M-AMBI, and in all three, the response variable was significantly higher in strongly connected 526 lagoons. Differently, H' and N2 were significantly higher and lower in moderately connected lagoons, 527 respectively, while M-AMBI was significantly lower in weakly connected lagoons. Mean Temp and mean 528 Sal were selected in two models, SR and macrofauna density. Mean Temp negatively influenced SR and 529 macrofauna density while mean Sal positively influenced SR and had a quadratic effect (concave form) on 530 macrofauna density. β_{MM} was selected in two models, H' and N2, and positively influenced both response 531 variables. The AMBI model was composed of mean [TN] (positive effect) and CV [NH_4^+] (quadratic effect, 532 concave form). Finally, three other eutrophication related variables, CV [TN], mean [NH_4^+] and CV [NO_x], 533 were selected respectively in H', macrofauna density and M-AMBI models where they had either a 534 negative effect (M-AMBI) or a quadratic effect (concave form, H' and density).

To isolate the independent effect of eutrophication on macrobenthic indices, we also plotted the
partial effect of mean [Chla] on SR, H' and M-AMBI once the effect of the other predictors retained in the
minimal adequate models (Table 3) had been removed (Appendix 6). After controlling for the effect of
mean Sal and mean Temp, the simple linear regression model between mean [Chla] and SR was still
significant (adj. $R^2 = 0.32$, $F(1, 39) = 19.55$, $p < 0.001$). It was also the case for the model between mean
[Chla] and H' after controlling for the effect of lagoon-sea connection, CV [TN], CV $[TN]^2$ and β_{MM} (adj. $R_2 =$
0.14, $F(1, 39) = 7.33$, $p = 0.01$) and for the model between mean [Chla] and M-AMBI after controlling for
lagoon-sea connection and CV $[NO_x]$ (adj. $R^2 = 0.26$, $F(1, 39) = 15.41$, $p < 0.001$).

Table 3 - Most parsimonious models (ordinary least-square regressions) explaining the standardized
macrobenthic taxonomic indices (species richness (SR), Shannon diversity (H'), inverse of Simpson's
dominance (N2), AZTI's Marine Biotic Index (AMBI), Multivariate AMBI (M-AMBI)). SR, H', N2 and
total macrofauna density were $\log_e$ transformed prior to standardization. Both eutrophication
related and natural variability related variables are considered as explanatory variables and the
slope estimates are expressed either as effect sizes when the variable is quantitative or as mean
standardized effects when the variable is categorical.

Model terms		Estimate	SE	p-value
SR: $F(3,37) = 18.18$; RSE = 0.66 ; adj. $R^2 = 0.56$				
[Chla]	mean	-0.50	0.12	< 0.001
Temperature	mean	-0.34	0.11	0.005
Salinity	mean	0.27	0.11	0.018
H': $F(6,34) = 13.57$; RSE = 0.59 ; adj. $R^2 = 0.65$				
Lagoon-sea connection	strong	0.61	0.19	< 0.001
	moderate	0.39	0.15	0.016
	weak	-0.16	0.17	0.32
[Chla]	mean	-0.31	0.12	0.016
[TN]	CV	-0.02	0.11	0.89
$[TN]^2$		-0.30	0.10	0.004
β_{MM}		0.23	0.10	0.035
N2: $F(3,37) = 11.18$; RSE = 0.75 ; adj. $R^2 = 0.43$				
Lagoon-sea connection	strong	0.79	0.21	< 0.001
	moderate	-0.58	0.19	0.005

	weak	-0.27	0.22	0.11
β_{MM}		0.31	0.12	0.016
Total macrofauna density: $F(5,35) = 8.48$; RSE = 0.72 ; adj. $R^2 = 0.48$				
Temperature	mean	-0.36	0.12	0.006
Salinity	mean	-0.23	0.12	0.071
Salinity^2		-0.30	0.13	0.023
[NH$_4^+$]	mean	0.21	0.12	0.10
[NH$_4^+$]^2		-0.30	0.13	0.030
AMBI: $F(3,37) = 5.84$; RSE = 0.86 ; adj. $R^2 = 0.27$				
[TN]	mean	0.34	0.14	0.017
[NH$_4^+$]	CV	0.11	0.14	0.41
[NH$_4^+$]^2		-0.43	0.13	0.003
M-AMBI: $F(4,36) = 29.04$; RSE = 0.51 ; adj. $R^2 = 0.74$				
Lagoon-sea connection	strong	0.59	0.16	0.005
	moderate	-0.06	0.14	0.67
	weak	-0.52	0.15	0.028
[NO$_x$]	CV	-0.37	0.09	< 0.001
[Chla]	mean	-0.35	0.09	< 0.001

551 RSE: Residual standard error, CV: coefficient of variation, [Chla]: chlorophyll *a* concentration in the water, [TN]: total nitrogen 552 concentration in the water, β_{MM}: beta diversity of the macrophyte morphotypes, [NH$_4^+$]: ammonium concentration in the water and [NO$_x$]: 553 nitrate plus nitrite concentration in the water

555 To synthesize our results, we plotted for each station the M-AMBI along with the three macrobenthic 556 indices composing it (SR, H' and AMBI) (mean ± SD) as a function of mean [Chla] (Figure 4). Mean [Chla] 557 was chosen as a synthetic indicator of eutrophication because it is (*i*) significantly and positively 558 correlated to 9 out of the 13 eutrophication related variables (Figure 2), (*ii*) the eutrophication related 559 variable (with mean [TN]) that presents the strongest correlations with SR, H' and M-AMBI (Figure 2), (*iii*) 560 the most frequently selected eutrophication related variable in the macrobenthic index models (Table 3) 561 and (*iv*) a relevant indicator of eutrophication status according to Bec et al. (2011) and Cloern (2001).

Figure 4 - Mean (± SD) of the species richness (SR), Shannon diversity (H'), AMBI and M-AMBI indices as a

function of mean chlorophyll a concentration ($\log_e$ scale) across 41 stations located in 29 lagoons along the

French Mediterranean coast (see Table 1 for the full lagoon names and corresponding stations). Across all

plots, each station (point) is represented using a symbol for its salinity type (triangle: oligo- and meso-

haline and circle: poly- and eu-haline) and a color (red: group A and blue: group B) for its group

membership (K-means partitioning) based on Hellinger-transformed mean abundances. The dashed lines

correspond to loess regressions.

SR, H' and M-AMBI were significantly correlated to mean [Chla] (r (Spearman) > |0.50|) while AMBI was

not (r = 0.12). Based only on the mean [Chla], most stations from group B (in blue) grouped towards

higher levels of mean SR, H' and M-AMBI than stations from group A (in red). Nonetheless, we identified

an intermediary zone corresponding to mean [Chla] between 6.3 µg.L^{-1} and 14.2 µg.L^{-1} and composed of

stations from both group located between the INN (SR = 12 ± 3, H' = 3.06 ± 0.42 and M-AMBI = 0.57 ± 0.1)

and ORE (SR = 9 ± 1, H' = 1.84 ± 0.07 and M-AMBI = 0.31 ± 0.01) stations. In the line with these

observations, the LPN, LPS and BOL stations appeared as outliers, as did LPN and LPS in the RDA (Figure

3). Finally, mean [Chla] was also significantly correlated (r (Spearman) = -0.40, *p* = 0.009) to the intra-

station SR variability expressed using the standard deviation.

Disentangling natural from anthropogenic factors

581 Based on the eutrophication and natural variability variables independently selected, we performed a 582variance partitioning, to investigate the amount of community and macrobenthic index variability 583uniquely linked to each variable category (Figure 5). Macrobenthic community and indices can be 584grouped into three categories according to the relative importance of natural variability alone, 585eutrophication alone, and the two together, in explaining their respective variability. (*i*) Macrofauna 586density was first influenced uniquely by natural variability (29.0%), then jointly by natural variability and 587eutrophication (11.5%) and finally uniquely by eutrophication (7.9%). (*ii*) Macrobenthic community, N2 588and H' were first jointly influenced by natural variability and eutrophication (community: 8.2%, H': 33.6%, 589N2: 22.1%), and then uniquely by natural variability (community: 8.0%, H': 17.8%, N2: 21.1%) and finally 590uniquely by eutrophication (community: 3.7%, H': 13.9%, N2: 2.9%). (*iii*) SR and M-AMBI were first jointly 591influenced by natural variability and eutrophication (SR: 30.7%, M-AMBI: 49.0%), and then uniquely by 592eutrophication (SR: 20.3%, M-AMBI: 17.6%) and finally uniquely by natural variability (SR: 5.8%, M-AMBI: 59338.9%).

Figure 5 - On the left, variance (adjusted R2) of the macrofauna community data (community), total macrofauna density (density), inverse of Simpson's dominance (N2), Shannon diversity (H'), species richness (SR) and multivariate AMBI (M-AMBI) uniquely attributed to eutrophication (green), uniquely attributed to natural variability (blue) and jointly attributed to both effects (red). The residual variance is shown in white. On the right, eutrophication variables and variables related to natural variability considered in the variance partitioning of the macrofauna community and macrobenthic taxonomic indices.

603 Using figure 2, we can identify variables related to eutrophication or to natural variability that 604significantly co-vary and are jointly implicated in a macrobenthic index model (Table 3) and/or in a 605variance partitioning (Figure 5). Mean [Chla] was significantly correlated with lagoon surface (-0.35), β_{MM} 606(-0.37), mean Temp (0.40) and CV O_2 sat (0.40). Mean [NO_x] was significantly correlated with lagoon 607surface (-0.38), mean Sal (-0.49) and CV O_2 sat (-0.33) while CV [NO_x] was significantly correlated with 608lagoon surface (-0.37), β_{MM} (-0.36) and mean Temp (0.53). Finally, mean [DIP] and lagoon surface were

609significantly correlated (-0.32). Lagoon-sea connection, a qualitative variable, was another source of 610confounding effects between natural variability and eutrophication on the macrobenthic indices and 611community. Indeed, there was a significant difference between the lagoon-sea connection and: *(i)* mean 612[Chla] (weak > strong connection); *(ii)* mean [NO_x] (weak > strong connection) and *(iii)* CV [NO_x] (moderate 613> strong).

614   **Discussion**

615   As outlined by Cloern (2001), eutrophication is a complex phenomenon with direct (*e.g.* changes in 616dissolved nutrient contents) and indirect (*e.g.* changes in community) responses that are modulated by 617attributes of each recipient system. This study is a first step towards better understanding how hydro-618morphology, natural environmental variability and eutrophication jointly shape macrobenthic 619communities in Mediterranean coastal lagoon ecosystems.
620
621**Macrobenthic communities are driven by lagoon hydro-morphology and natural environmental** 622**(in)stability**
623   Lagoon macrobenthic communities appear to be first determined by lagoon hydro-morphology and 624natural environmental stability. When investigating the independent effect of each environmental 625parameter on macrobenthic communities, lagoon-sea connection level was the variable explaining the 626most of H', N2 and M-AMBI variability and the second most important variable explaining community 627composition. This qualitative variable is a proxy for marine water renewal (*i.e.* residence time), a gradient 628well known for characterizing coastal lagoon biological communities (Barbone and Basset, 2010; Perez-629Ruzafa and Marcos-Diego, 1992; Tagliapietra et al., 2009), and closely linked to mean Sal (moderate > 630weak and strong > weak) and CV Sal (weak > moderate > strong). Salinity, historically used to classify 631coastal lagoons (Anonymous, 1959), was also an important driver of SR (linear positive effect) and total 632macrofauna density (concave effect). Indeed, simple and multimetric descriptors of lagoon macrobenthic 633communities like species richness are known to be highly and positively influenced by mean water salinity 634(Barbone et al., 2012; Magni et al., 2022; Pitacco et al., 2019; Reizopoulou et al., 2014). Differently from 635us, Barbone et al. (2012) reported a macrofauna density decrease from oligo- to eu-haline lagoons, 636maybe because only undisturbed lagoons were considered in their study. Furthermore, the effect of 637lagoon-sea connection on N2, a diversity index sensitive to high abundance taxa (Morris et al., 2014), 638indicates that a strong connection to the sea promotes benthic communities with significantly more 639relatively abundant taxa whereas moderate and weak connections promote communities dominated by a 640few abundant and in our case, euryhaline taxa (*e.g. Hediste diversicolor, Streblospio shrubsolii*), a 641characteristic typical of naturally stressed environments like lagoons subject to strong salinity variations 642(Elliott and Quintino, 2007). This result seems to confirm that moderately and weakly connected lagoons 643are highly selective environments for benthic organisms at least partly because of their strongly 644fluctuating environmental conditions (Barnes, 1980; Provost et al., 2012; Zaldívar et al., 2008). The 645negative effect of decreasing salinity and lower connection levels to the sea on SR, N2 and macrofauna 646density can also be linked to the physiological stress that natural and abrupt salinity variations (*e.g.* 647droughts, heavy rainfall) can represent for larvae, juveniles and adults of marine-originating benthic 648organisms inhabiting coastal lagoons (Cognetti and Maltagliati, 2000; Magalhães et al., 2019; Vernberg, 6491982). Overall and based on other published evidence, our results could be explained by *(i)* water 650renewal from marine origin, *(ii)* primary colonization and/or post-disturbance recolonization of lagoons 651by marine-originating larvae through dispersal and recruitment and *(iii)* environmental (in)stability, which 652are known to strongly shape lagoon benthic communities (Basset et al., 2006; Pérez-Ruzafa et al., 2007b;

653Pérez-Ruzafa and Marcos-Diego, 1992), as proposed in the confinement theory (Guelorget and
654Perthuisot, 1983).

655   Water temperature was another important driver of lagoon macrobenthic communities, SR and
656macrofauna density in our study. Temperature had a negative effect on these communities, probably
657directly through thermal stress, and indirectly through its effect on oxygen dynamics. Higher and more
658variable summer temperatures directly represent an abiotic constraint for benthic organisms by
659increasing their oxygen demand and reducing their mobility (Vernberg, 1982). At the same time, high
660summer temperatures decrease oxygen solubility and increase the probability of hypoxia and anoxia
661(Derolez et al., 2020a) while also reducing the tolerance of benthic organisms to low dissolved oxygen
662(Vaquer-Sunyer and Duarte, 2011). This cocktail of effects can ultimately cause the death of sensitive
663taxa, such as large sized and long-lived species (Quillien et al., 2015; Reizopoulou and Nicolaidou, 2007;
664Sigala et al., 2012) and reduce the ability to move of more tolerant taxa (Vernberg, 1982), hence
665increasing their susceptibility to predation and decreasing their possibility to search for food and/or for
666less stressful environmental conditions (Díaz and Rosenberg, 1995; Rosenberg et al., 1992; Weissberger
667et al., 2009).

668   The diversity of macrophyte morphotypes (β_{MM}) also affected positively H' and N2, and was positively
669correlated to all macrobenthic indices except density and AMBI. Simple taxonomic indices like lagoon
670macrobenthic species richness, biomass density and numerical density are locally sensitive to vegetation
671presence and type (Arocena, 2007; Barbone et al., 2012, Magni and Gravina, 2023). Conversely, we show
672here the probable effect of benthic habitat heterogeneity expressed at the lagoon or sub-lagoon scale on
673the structure of benthic communities at a local scale (*ca.* 50 m^2 station), an effect mainly visible when
674focusing on intermediate and high abundance taxa (Morris et al., 2014). Focusing on macrobenthic fauna
675at the lagoon scale, β_{MM} informs us on the number of potential (*i*) environmental niches that can be filled
676by the species pool and (*ii*) refuges (Magni and Gravina, 2023; Nordström and Booth, 2007; Ware et al.,
6772019) from which these organisms can (re)colonize habitat patches after mortality events caused by
678temporary unfavorable conditions. A model of local extinctions linked to adverse environmental
679conditions, followed by recolonization events via pelagic larvae was suggested by Vergara-Chen et al.
680(2013) to explain the small-scale genetic structure of the cockle *Cerastoderma glaucum* inside the Mar
681Menor lagoon (Spain). The positive effect of β_{MM} on macrobenthic diversity also provides us with a few
682leads on how macrophytes can structure lagoon macrobenthic communities. Macrophytes can act as (*i*)
683temporary refuges from vagile predators for small mobile benthic organisms like amphipods (Nordström
684and Booth, 2007; Ware et al., 2019), (*ii*) a food source for grazers and detritus-feeders (Carlier et al.,
6852007; Lepoint et al., 2000; Ouisse et al., 2011; Vizzini and Mazzola, 2006) and (*iii*) settlement promoters
686for pelagic particles like phytoplankton and larvae (Colden et al., 2016; Donadi et al., 2014; González-Ortiz
687et al., 2014), especially in the case of meadow-forming species like *Zostera*. All these local effects are
688likely to spill-over to adjacent unvegetated sediments (Boström et al., 2006; Magni et al., 2017).

689   Overall, natural environmental (in)stability, connection level with adjacent water systems and/or
690number of potential environmental niches seem to be the main drivers of lagoon macrobenthic
691communities and to be stronger drivers of benthic macrofauna assemblages than eutrophication, which
692probably represents an additional source of stress for benthic invertebrates (Barbone and Basset, 2010).

693   The environmental variables considered in the manuscript do not unfortunately explain the
694communities described at the LPN and LPS stations. These two macrofauna stations sampled were
695actually classified in a group generally associated to small eutrophic lagoons and mainly composed of
696stations sampled in lagoons evaluated by the WFD as being in a poor or in a bad physico-chemical status
697even through is considered in a good physico-chemical status according to Chla concentration. This could
698be the result of the strong variations in salinity recorded throughout the year, which we have not been
699considered here. In this lagoon, marked annual salinity variations are observed (between 3.6 and 76.8 in

700Wilke and Boutiere (2000)), variations indetectable with the available summer salinity data. 701Consequently, the eight environmental variables selected to explain the overall macrofauna communities 702were not able to explain the one characterizing this lagoon. This result also reinforces the importance of 703natural environmental instability probably has on structuring benthic macrofauna communities.
704

705**Eutrophication affects lagoon macrobenthic communities through interconnections with natural** 706**environmental variability**

707   Eutrophication appears to act as an additional disturbance for lagoon macrobenthic communities via 708direct (*e.g.* toxicity of inorganic nitrogenous compounds) and indirect mechanisms (*e.g.* hypoxic stress). 709The three groups derived from the variance partitioning approach highlight that the most influential 710category of variables (*i.e.* natural variability alone, eutrophication alone, natural variability and 711eutrophication together) is not the same depending on the level of biological organization we focus on. 712Indeed, the benthic invertebrate density was first explained by natural variability alone and more 713precisely by physiologically constraining variables (salinity and temperature) whereas macrobenthic 714community composition, SR, H' and N2 were first explained jointly by natural variability and 715eutrophication. With these joint effects, we probably measure *(i)* how lagoon-sea connection modulates 716the susceptibility (*i.e.* propensity) of each lagoon to eutrophication, *(ii)* how higher water temperatures 717promote phytoplankton blooms and *(iii)* how eutrophication reduces benthic habitat heterogeneity, 718followed by their respective effects on benthic community diversity.

719   *(i)* Lagoon-sea connection informs us on each lagoon's water residence time and consequently, on its 720retention potential regarding nutrients and pollutants (Chapelle et al., 2001; Cloern, 2001; Fiandrino et 721al., 2017). As such, watershed related nutrients and their direct effect on primary producers are buffered 722by each lagoon's degree of connection to the sea, as indicated by the significantly higher mean [Chla] and 723mean [NO$_x$] in lagoons weakly connected to the sea compared with strongly connected ones. *(ii)* 724Independently from lagoon hydro-morphology, the positive correlation between mean [Chla] and mean 725temperature illustrates how higher water temperatures increase phytoplankton growth rates which can 726lead to blooms (Rose and Caron, 2007), as recently demonstrated in Thau (Trombetta et al., 2019), a large 727lagoon strongly connected to the sea. Nonetheless, it seems that eutrophication is buffered in larger 728lagoons where water temperatures increase less in summer, as suggested by the negative correlation 729between lagoon surface and mean summer temperature. *(iii)* Furthermore, one of the direct responses of 730Mediterranean coastal lagoons to changes in nutrient inputs is an increase in pelagic primary production 731(Souchu et al., 2010), which in turn decreases water transparency (Cloern, 2001) and limits light 732availability for benthic primary producers (Le Fur et al., 2019, 2018). Ultimately, sustained eutrophication 733of coastal lagoons can cause a regime shift from a primary producer community dominated by perennial 734macrophytes presenting different morphotypes, to a community dominated by free-floating and 735opportunistic macroalgae like *Ulva rigida* and finally, to a phytoplankton-dominated community (Le Fur 736et al., 2019). The negative correlation between several eutrophication variables (*e.g.* mean [Chla], mean 737turbidity) and β_{MM} highlights the simplification of benthic habitats caused by lagoon eutrophication 738(Cloern, 2001; Le Fur et al., 2019, 2018) that shift from a mosaic of different macrophyte morphotypes to 739a sediment void of macrophytes or with one macrophyte morphotype. Overall, the multiple 740interconnections between natural environmental variability and eutrophication likely explain the high 741amount of macrobenthic community and index variance jointly attributed to both effects, as underlined 742in the conceptual model developed by Cloern (2001).

743   The unique effect of eutrophication variables on each taxonomic component also informs us about 744which of these variables likely have a direct effect on macrobenthic invertebrates without being 745modulated by natural variability. Indeed, benthic organisms can be affected by eutrophication directly 746through the toxicity of inorganic nitrogenous compounds like ammonium, nitrites and nitrates (Jessen et

747al., 2015). Our results indicate that high NH_4^+ concentrations seem to have a very small negative effect on
748total macrofauna density while NO_x toxicity could explain the strong unique effect of eutrophication on
749SR and M-AMBI, and both NH_4^+ and NO_x toxicity could be responsible for the very weak unique effect of
750eutrophication on macrobenthic communities. These results are probably also linked to indirect effects
751high eutrophication levels can have on macrobenthic communities via increased water turbidity and
752sediment bio-geochemistry modifications (*e.g.* anoxic superficial sediments (Zilius et al., 2015)). Indeed,
753water NH_4^+ and NO_x concentrations in excess can be the sign of highly eutrophic lagoons (Souchu et al.,
7542010). Overall, experimental work on a few well-selected species and populations sampled in lagoons
755presenting different levels of natural variability would greatly improve our understanding of the effect of
756eutrophication on lagoon benthic invertebrates via inorganic nitrogenous compound toxicity. Even trough
757marine animals appear to be less sensitive to nitrate than freshwater animals, nitrate toxicity to benthic
758invertebrate increases with increasing nitrate concentrations and exposure times (Alonso and Camargo,
7592006; Boardman et al., 2004; Camargo et al., 2005) Nitrate toxicity may also increase with decreasing
760body size, water salinity, and environmental adaptation. In lagoon environments, where organisms are
761often smaller in size, this could also be of importance.

762   Furthermore, mean [Chla] appears to have a strong negative effect on macrobenthic communities by
763impacting the diversity of low and intermediate abundance taxa (i.e. SR and H' are diversity indices that
764give more weight respectively, to low abundance/rare and intermediate abundance taxa) and this effect
765is at least partly independent from the selected natural variability variables (*i.e.* mean temperature, mean
766salinity for SR and lagoon-sea connection, β_{MM} for H'). Two non-exclusive hypotheses could explain this
767result: small-scale habitat homogenization and hypoxic stress. Despite not being the appropriate index to
768investigate beta diversity, the standard deviation of SR at the intra-station scale was significantly
769correlated negatively to mean [Chla], which seems to point towards a homogenization of benthic
770communities at the meter scale (i.e. a few meters separate each replicate of a given station) linked to
771eutrophication. The loss of benthic taxa presenting restricted environmental niches and dependent upon
772small-scale habitat diversity for reproduction, recruitment, refuge and/or food (Brauns et al., 2007;
773Harman, 1972; Stendera and Johnson, 2008) could explain this result. Secondly, low to zero dissolved
774oxygen (*i.e.* hypoxia and anoxia) is another well-known symptom of eutrophication in transitional waters
775like coastal lagoons (Chapelle et al., 2001; Cladas et al., 2016; Vignes et al., 2010). Oxygen depletion
776generally takes place during the summer and is maximal at the end of the night/early morning (Newton
777et al., 2010), often starting near the bottom because of benthic organic matter mineralization (*e.g.*
778sedimented phytoplankton, dead macroalgae) and low water renewal (Rigaud et al., 2021; Souchu et al.,
7791998). In the worst cases, these low to zero oxygen zones can spread to the entire water column and last
780for several days (Chapelle et al., 2001; Cladas et al., 2016), as seen during the 2018 dystrophic crisis in
781Thau lagoon (Derolez et al., 2020a). These crises are known to impact macrobenthic community first in
782terms of abundance and then in terms of composition and species richness depending on the sensitivity
783of each species to reduced oxygen (Díaz and Rosenberg, 1995; Riedel et al., 2012). Consequently, lagoon
784macrobenthic invertebrates are often reported as being less abundant and less diverse in fall than in
785spring because of summer hypoxic stress (Barbone et al., 2012; Basset et al., 2012; Vanda Orro and
786Cabana, 2021). The negative impact of eutrophication via hypoxic/anoxic stress on lagoon macrobenthic
787communities is suggested by the RDA results where high mean O_2 sat (*i.e.* hyper-saturation linked to
788strong pelagic primary production) is associated to group A stations that present significantly lower
789species richness than group B stations. Nonetheless, oxygen related variables were not selected in the
790other models, a result probably linked to when oxygen was measured (late morning) and at what depth it
791was measured (generally sub-surface). Indeed, these features limit our ability to detect potential short
792and long-term effects of hypoxic stress on macrobenthic communities. Bottom hypoxia/anoxia is also
793known to predominantly take place in deep lagoons like Berre and Thau (Derolez et al., 2020a; Rigaud et

794al., 2021; Souchu et al., 1998), where wind derived water mixing is limited by depth. Nonetheless, anoxic
795crises inside coastal lagoons are expected to increase in frequency and intensity with climate change
796(Derolez et al., 2020a; Miyamoto et al., 2019). Consequently, it appears paramount to (*i*) study the
797sensitivity of lagoon benthic organisms to low dissolved oxygen, not only of freshwater and marine taxa
798(Díaz and Rosenberg, 1995; Landman et al., 2005) and (*ii*) monitor more closely summer bottom dissolved
799oxygen concentrations, especially at higher frequency (O_2 being very variable) and at temporal scales
800relevant to these organisms (*i.e.* across several days starting when bottom hypoxia is detected (Riedel et
801al., 2012)).

802   Overall, lagoon hydro-morphology (connection to the sea, surface and depth) which affects the level
803of environmental variability (salinity and temperature), as well as lagoon-scale benthic habitat diversity
804(β_{MM}) seem to regulate the distribution of macrobenthic species (Bachelet et al., 2000), while
805eutrophication and associated stressors like low dissolved oxygen act upon the existing communities,
806mainly by reducing taxa richness and diversity (SR and H'). Disentangling the effects of anthropogenic
807stressors and natural environmental variables remains nevertheless challenging, especially as stressors
808and variables can co-vary, even in the absence of anthropogenic impact. Moreover, it is often impossible
809to measure effects of every single possible stressor on macrobenthic communites. In our study, grain-size
810could not be included to differences in laboratory protocols between lagoons and between years and
811heavy metal contaminations were not available at the same temporal and spatial scale (one
812measurement per lagoon every ten years). It should be relevant to re-run the analyses when such data
813would be available at the macrofauna sampling sites to test the effect of this supplementary
814anthropogenic driver to check if it would be discriminating benthic communities.
815

816**The M-AMBI and recommendations for its use as an indicator for Mediterranean coastal lagoons**

817   With this study, we investigate for the first time the joint sensitivity of the M-AMBI index to natural
818and anthropogenic stressors across 29 lagoons covering most of the French Mediterranean coastline. The
819M-AMBI (Muxika et al., 2007) is currently used to evaluate the WFD ecological state of poly- and eu-
820haline French Mediterranean lagoons based on soft sediment benthic invertebrate, with reference
821conditions using minimally impacted sites (Reizopolou et al., 2018). This index has previously been shown
822to be sensitive to pressures related to eutrophication, like concentration in chlorophyll *a* and total
823nitrogen (Derolez et al., 2014) but it is also known to be sensitive to natural environmental variability
824(*e.g.* water salinity) and lagoon hydro-morphology (*e.g.* surface and depth) (Barbone et al., 2012).

825   Our results show that the M-AMBI appears to be a sensitive indicator of eutrophication through the
826species richness and Shannon diversity components of the index but not the AMBI one. First, the strong
827correlation between M-AMBI and mean [Chla] was linked only to the significant correlation between SR
828and mean [Chla] and between H' and mean [Chla] while AMBI was not significantly linked to most
829eutrophication related variables (except mean [NO_x]) nor to chlorophyll *a* concentration, a common
830eutrophication indicator (Bec et al., 2011; Cloern, 2001). Importantly, these linear relations remained
831significant once the effect of identified covariables was removed. Regarding AMBI, its current application
832in transitional waters like coastal lagoons suffers from methodological limitations. Indeed, this index is
833based on macroinvertebrate sensitivity/tolerance to organic matter enrichment following Pearson and
834Rosenberg (1978) and was developed for coastal waters (Borja et al., 2000), not for coastal lagoons,
835which are known to sometimes present sediments naturally enriched in organic matter linked to the
836decay of meadow-forming macrophytes (Bachelet et al., 2000), watershed inputs and the overall low
837organic matter exportation (except during floods) (Perthuisot and Guelorget, 1983). These results
838highlight the limits of indirectly including AMBI in the evaluation of the ecological state of Mediterranean
839coastal lagoons. Nonetheless, its absence of sensitivity to natural variability and lagoon hydro-
840morphology suggests an index based on redefined ecological groups could be promising (Robertson et al.,

8412016), especially if we aim towards a more specific macrofauna-based index to evaluate eutrophication. 842Overall, increasing the robustness of M-AMBI would likely necessitate additional work to classify lagoon 843macroinvertebrates according to other eutrophication related stressors like low oxygen, nitrogenous 844compound toxicity and/or hydrogen sulfide toxicity.

845   Secondly, M-AMBI variability was two times more strongly attributed uniquely to eutrophication 846related variables than uniquely to natural variability ones, with nonetheless a very strong joint effect of 847the two types of variables. Our study reinforce the difficulty of M-AMBI to clearly identify natural 848disturbance but also the high sensitivity of this multimetric index to anthropogenic pressures such as 849eutrophication or metal contaminations and pesticides in transitional waters (Paul et al., 2023; Pelletier 850and Charpentier, 2023; Pollice et al., 2015). The strong joint effect is probably related, as previously 851discussed, to the same interconnections between lagoon-sea connection that buffers eutrophication, 852temperature that promotes phytoplankton blooms and eutrophication that homogenizes benthic 853habitats. Several modifications could help limit the sensitivity of M-AMBI to natural variability. First, as 854recommended by Borja et al. (2012) who pointed that the inability of this index to detect anthropogenic 855responses is often linked to the use of inappropriate reference conditions, the currently used poly- and 856eu-haline vs oligo- and meso-haline categories (Provost et al., 2012; Reizopolou et al., 2018) could be 857replaced by one based on the three lagoon-sea connection levels that strongly influence the M-AMBI or 858on another typology like the choked vs restricted typology (Barbone et al., 2012; Basset et al., 2013), with 859the test of new reference lagoons (Reizopolou et al., 2018). Secondly, the replacement of a sampling 860done with an Ekman-Birge grab by a diver-operated sampling would limit heterogeneity between 861replicats by (i) verifying that only unvegetated sediments are being collected and (ii) describing the 862presence and type of macrophyte near the sampling site (*e.g.* 10 meter radius). This could represent a 863bias in many homogeneous environments where random selection of the sampling zone is quite possible 864but macrophytes are often heterogeneously distributed in coastal lagoon and areas of unvegetated 865sediments are rare. Selection and description of the sampling area by divers could therefore be relevant 866in these shallow environments even if this often means additional human and financial costs. This last 867modification would likely help reduce the effect of local benthic habitat heterogeneity on H' and M-AMBI 868(Magni and Gravina, 2023; de Paz et al., 2008).

869   Finally, several studies have underlined the limits of using the M-AMBI based on abundance to 870evaluate the ecological state of transitional ecosystems like coastal lagoons, as opportunistic and tolerant 871species are generally numerically dominant in these systems (Mistri et al., 2018; Pitacco et al., 2019; 872Puente and Diaz, 2008). Other indicators based on the size spectra and biomass of benthic 873macroinvertebrates like the biomass based M-AMBI (M-bAMBI (Mistri et al., 2018)) and the index of size 874distribution (ISD (Reizopoulou and Nicolaidou, 2007)) have showed promising results and applying these 875indices to French coastal lagoons would represent an additional trial in the context of historically highly 876eutrophic lagoons currently undergoing a re-oligotrophication trajectory (Derolez et al., 2020a, 2020b; Le 877Fur et al., 2019; Leruste et al., 2016). Another interesting avenue would be to investigate more closely the 878spatial scales at which changes in macroinvertebrate abundance and diversity can be detected and link 879these spatial changes to localized anthropogenic disturbances like heavy metals (Magalhães et al., 2019) 880while also considering small-scale biotic interactions such as competition, predation and facilitation. 881Indeed, sediment contaminants like biotic interactions are likely to affect benthic macrofauna at fine 882spatial scales (*ca.* 1 km) that are more relevant to low mobility macroinvertebrates (Berthelsen et al., 8832018; Brauko et al., 2020, 2015; Carvalho et al., 2006).

884   In a nutshell, to improve the robustness of M-AMBI, we call for a revision of the ecological groups at 885the base of the AMBI index and of the sampling protocol (*e.g.* using divers) along with a re-evaluation of 886the lagoon typology. More generally, it appears key to increase our understanding of the effects of other 887disturbances like hypoxia and heavy metals on lagoon invertebrates and to reevaluate the small-scale

888distribution of benthic organisms in French coastal lagoons, more than 30 years after the historical
889studies by Amanieu et al. (1977), Guelorget et al. (1994) and Guelorget and Michel (1979).
890

891Overcoming taxonomic limitations

892   A large part of the identified macroinvertebrates had a very restricted geographical range, preventing
893us from building a general species-based framework linking eutrophication and macrobenthic
894communities. Indeed, we found a very high level of lagoon-specificity for benthic macroinvertebrates
895across French Mediterranean coastal lagoons with no taxon present across all 41 stations, 40% of the 227
896taxa identified in only one station and only 1.3% recorded in over half of the stations. This result is
897probably explained by environmental filtering resulting from very contrasted conditions across the
898different sampled systems, passive larval diffusion (Basset et al., 2007) and the randomness of the
899colonization process by species of marine origin (Pérez-Ruzafa et al., 2011b), a phenomenon previously
900observed for benthic macroinvertebrates in Italian coastal lagoons (Basset et al., 2007), fish in Atlantic
901and Mediterranean coastal lagoons (Pérez-Ruzafa et al., 2007b) and macrophytes in estuaries and
902lagoons (Pérez-Ruzafa et al., 2011c). Consequently, unique assemblages of benthic macroinvertebrates
903are present in each lagoon and even at smaller spatial scales (*i.e.* stations inside a lagoon), which could
904explain why we were able to explain only 22% of the macrobenthic community variability. This high level
905of lagoon-specificity present across multiple biological compartments in coastal lagoons warrants the
906need to go beyond a species-based approach and towards a trait-based approach (Lacoste et al., 2023),
907as previously outlined using macrofauna size (Basset et al., 2008; Brauko et al., 2020; Sigala et al., 2012).
908Such an approach could build upon the saprobity concept (Tagliapietra et al., 2012) to select relevant
909biological traits and ultimately help us build a more general framework around macrobenthic community
910assembly rules in the line of what Pérez-Ruzafa et al. (2013) did for fish by revisiting the r and K-
911strategies. Nonetheless, for such an approach to be relevant, we also need to increase our knowledge of
912the life-cycle of lagoon benthic invertebrates (Brauko et al., 2020) and to create a harmonized
913Mediterranean wide trait database for these organisms (Lacoste et al., 2023).

914   Another interesting avenue to pursue could be to investigate intra-specific biological trait and/or
915physiological response variability (*i.e.* oxydative stress) of species like the polychaete *Hediste diversicolor*
916along eutrophication gradients. Indeed, this species is known to display a great diversity of life history
917characteristics along with morphological, biochemical and physiological differences according to local
918environmental conditions (Scaps, 2002). For example, the feeding behavior and diet of this species has
919been shown to vary according to sewage pollution levels across English estuaries (Aberson et al., 2016).
920Another study has related oxydative stress of the polychaete *Nephtys cirrosa* to seasonal variations of
921abiotic parameters like temperature and salinity in the Ria de Aveiro (Magalhães et al., 2019).
922Furthermore, *H. diversicolor*, a species we found to be associated to eutrophic lagoons, is reported to
923present a great diversity of feeding modalities (*i.e.* predator, grazer, suspension-feeder and deposit-
924feeder (Faulwetter et al., 2014)), whereas the species associated to oligotrophic lagoons, like *Loripes*
925*orbiculatus*, present less diverse feeding modes (*i.e.* mainly based on sulfo-oxydizing bacteria (Roques et
926al., 2020)). This flexibility could allow organisms to better tolerate the fluctuating abiotic conditions that
927characterize small eutrophic lagoons, stressing that environmental stability is probably a strong driver of
928macrobenthic communities in coastal lagoons, as previously shown for macrophytes (Pérez-Ruzafa et al.,
9292011c). Overall, the plasticity several species display in environmentally constraining conditions stresses
930the difficulty to implement trait-based approaches in ultra-variable environmental systems like coastal
931lagoons, where realized trait modalities may differ significantly from the ones available in the literature,
932commonly acquired from non-transitional coastal ecosystems (Martini et al., 2021). Finally, a functional
933approach based on well-informed biological traits and biomass data could help overcome the identified
934taxonomic limitations and build a more holistic framework linking natural variability and eutrophication

935of coastal lagoons to macrobenthic communities but the intra-specific plasticity for many biological traits
936makes this approach questionable today.

Conclusion

938Coastal lagoons are transitional ecosystems naturally characterized by a strong variability in abiotic
939parameters like salinity and temperature. Superimposed onto this natural environmental variability are
940numerous anthropogenic disturbances, eutrophication being a historical one targeted by the European
941Water Framework Directive. By jointly analyzing water column, sediment and macrophyte parameters,
942our study shows that macrobenthic communities across 29 French Mediterranean coastal lagoons are
943primarily determined by (*i*) lagoon hydro-morphology (connection level with the sea, surface and depth)
944which further affects natural environmental variability in terms of salinity and temperature, and (*ii*)
945lagoon-scale benthic habitat diversity, as estimated through macrophyte morphotypes. Eutrophication
946and associated stressors like low dissolved oxygen act upon the existing communities by reducing
947taxonomic richness and diversity. Eutrophication appeared to act on lagoon macrobenthos through
948relatively direct mechanisms like toxicity of inorganic nitrogenous compounds, as well as indirectly via for
949example, the simplification of benthic habitats and sediment bio-geochemistry modifications.
950Furthermore, our study highlights the complex interconnections between lagoon hydro-morphology,
951natural environmental variability and eutrophication and their joint effect on benthic communities.
952Indeed, lagoon-sea connection modulates the susceptibility of each lagoon to eutrophication, high water
953temperatures promote eutrophication related phytoplankton blooms while eutrophication reduces
954benthic habitat heterogeneity, all of which affect benthic communities. Finally, benthic taxa richness and
955diversity also proved to be indicators sensitive to eutrophication in coastal lagoons while an AMBI like
956index based on redefined ecological groups for coastal lagoons, which can be naturally enriched in
957organic matter, has the potential of being a specific eutrophication indicator.
958
959

961 **Appendices**

962 **Appendix 1 - Procedure used to calculate the beta diversity of macrophyte morphotypes (β$_{MM}$)**
963 **corresponding to each macrofauna station**
964

Step 1: Select the macrophyte stations corresponding to each macrofauna station

• If one macrofauna station is considered in a given lagoon, then all the macrophyte stations of
that given lagoon are taken into account.

• If two or three macrofauna stations are considered in a given lagoon, then all the macrophyte
stations attributed by the WFD protocol to the same section of the lagoon as the macrofauna
station are taken into account.
971

Step 2 : Build the macrophyte station by macrophyte morphotype matrix for each macrofauna station
For each macrophyte station of the dataset:

• If the macrophyte cover is below 25% or equal to 25%, a value of 1 is entered in the sediment
column and 0 in all the others (line 1 of the example matrix below).

• If the cover is above 25%, the dominant or co-dominant taxa are determined based on the taxa
specific cover data or the taxa specific biomass data, then these taxa are recoded into a
morphotype (see Appendix 2) and finally, a value of 1 is entered in all the cells corresponding to
the given morphotypes of the dominant or co-dominant taxa and a value of 0 is entered in the
remaining cells (lines 2 and 3 of the example matrix below).
981

982 *Step 3 : Calculate the index value corresponding to each macrofauna station*

983 • Calculate the Bray-Curtis dissimilarity between all the station pairs using the previously built
984 station by macrophyte morphotype matrix (presence/absence)

985 • Average the Bray-Curtis dissimilarities
986

987 Example of a macrophyte station by macrophyte morphotype matrix used to calculate the β$_{MM}$

Station	mushroom	feather	hair	ball	branched	sheet	meadow	sediment
1	0	0	0	0	0	0	0	1
2	0	0	0	0	1	0	1	0
3	0	0	0	0	0	1	0	0
...								
n-1								
n								

988
989
990
991

Appendix 2 - Correspondence between macrophyte taxa and morphotypes

Macrophyte taxa	Morphotype
Acetabularia acetabulum	mushroom
Aglaothamnion pseudobyssoides	
Alsidium corallinum	
Cladophora vagabunda	feather
Dasyaceae	
Spyridia filamentosa	
Chaetomorpha aerea	hair
Ulva clathrata	
Cladophora glomerata	ball
Valonia aegagropila	
Gracilaria dura	
Gracilaria gracilis	branched
Halopithys incurva	
Ulva rigida	sheet
Cymodocea nodosa	
Ruppia cirrhosa	
Stuckenia pectinata	meadow
Zostera marina	
Zostera noltei	

995**Appendix 3**

996Mean ± SD of the species richness (SR), Shannon diversity (H'), inverse of Simpson dominance index (N2), 997macrofauna density, AMBI (AZTI's Marine Biotic Index) and M-AMBI (Multivariate AMBI) calculated for 998the A and B station groups determined using a K-means partitioning based on the benthic macrofauna 999Hellinger-transformed mean abundances. The results of the non-parametric Kruskal-Wallis rank sum test 1000comparing the simple and multimetric macrobenthic taxonomic indices between the A and B groups are 1001also indicated.

	Group A	Group B	Kruskal-Wallis rank sum test
SR	9.30 ± 3.48	17.48 ± 11.41	$\chi^2(1) = 6.41$, $p = 0.011$
H'	2.00 ± 0.57	2.36 ± 0.69	$\chi^2(1) = 2.29$, $p = 0.13$
N2	3.30 ± 1.35	3.83 ± 1.76	$\chi^2(1) = 0.74$, $p = 0.39$
Macrofauna density (ind.m^{-2})	2 280 ± 1 372	5 113 ± 5 205	$\chi^2(1) = 0.98$, $p = 0.32$
AMBI	3.03 ± 0.53	2.87 ± 1.08	$\chi^2(1) = 0.01$, $p = 0.92$
M-AMBI	0.43 ± 0.08	0.53 ± 0.14	$\chi^2(1) = 5.76$, $p = 0.016$

1002
1003

1004**Appendix 4**

1005Coordinates (biplot scores for quantitative variables and centroids for lagoon-sea connection) on the first 1006and second axis of the redundancy analysis (RDA) plots, marginal effects and conditional effects of the 8 1007environmental variables selected to explain the macrobenthic community variability. After the 1008abbreviated name of each environmental variable is indicated the type of transformation used to 1009normalize it (log_e: natural logarithm and sqrt: square-root). [Chla] and [DIP] is the water concentration in 1010chlorophyll *a* (μg.L^{-1}) and dissolved inorganic phosphorous (μmol.L^{-1}), respectively. Temp (°C) is the lagoon 1011water temperature. O_2 sat (%) is the water oxygen saturation level. OMC (%) is the sediment organic 1012matter content. CV stands for coefficient of variation.

Selected environmental variable	RDA1	RDA2	Marginal effect (%)	Conditional effect (%)
Mean [Chla] (log_e)	-0.76	0.23	7.22	7.22
Lagoon surface (log_e)	0.73	-0.15	5.90	1.69
Weak lagoon-sea connection	-0.10	0.49		
Moderate lagoon-sea connection	-0.073	-0.18	5.30	4.64
Strong lagoon-sea connection	0.18	-0.29		
Mean Temp	-0.61	-0.24	5.04	3.62
Mean [DIP] (log_e)	-0.51	0.092	3.16	1.31
CV Temp (sqrt)	0.44	0.15	2.20	1.53
OMC	0.11	0.27	0.50	1.20
Mean O_2 sat (log_e)	-0.16	-0.063	0.47	1.28

1029**Appendix 5**

1030Partial redundancy analysis plots (pRDA, axis 1 and 2) focusing on (a) the benthic macrofauna sampled 1031across 41 stations located in 29 lagoons along the French Mediterranean coast (see Table 1 for the full 1032lagoon names and corresponding stations) as explained by keeping constant the natural environmental 1033variability variables (lagoon-sea connection, lagoon surface, mean and CV water temperature, mean 1034oxygen saturation and sediment organic matter content) and letting the eutrophication variables vary 1035(mean concentration in chlorophyll *a* (mean [Chla]) and mean dissolved inorganic phosphorous (mean 1036[DIP]), (b) the environmental variables left to vary where mean [Chla] and mean [DIP] are $\log_e$ 1037transformed (see Tables 1 and 2 for the full environmental variable names and values) and (c) the 1038macrofauna taxa, where only taxa with a goodness of fit (cumulative proportion of variance explained by 1039the first two axis) above 0.23 are plotted. In (a), inter-station distances are euclidean while in (b) and (c) 1040angles between all vectors reflect linear correlations. Across all plots, each station (point) is represented 1041using a symbol for its salinity type (triangle: oligo- and meso-haline and circle: poly- and eu-haline) and a 1042color (red: group A and blue: group B) for its group membership (K-means partitioning) based on 1043Hellinger-transformed mean abundances.

1044

1045

a

1046

b

c

1047

1049Partial effect of mean chlorophyll *a* concentration ([Chla]) on (a) species richness (SR) once the effect of 1050mean temperature and mean salinity has been removed, (b) Shannon diversity (H') once the effect of 1051lagoon-sea connection, CV [TN], CV [TN]^2 and β_{MM} has been removed and (c) M-AMBI once the effect of 1052lagoon-sea connection and CV [NO$_x$] has been removed (see Table 3 for the parsimonious linear models) 1053across 41 stations located in 29 lagoons along the French Mediterranean coast (see Table 1 for the full 1054lagoon names and corresponding stations). The dashed lines correspond to simple linear regression 1055models and the shaded zone to the confidence interval. Each station (point) is represented using a 1056symbol for its salinity type (triangle: oligo- and meso-haline and circle: poly- and eu-haline) and a color 1057(red: group A and blue: group B) for its group membership (K-means partitioning) based on Hellinger-1058transformed mean abundances.

1059

1060   **Acknowledgements**

1061The authors would like to thank all the members of the LERLR and LERPAC Ifremer labs as well as all the
1062managers of the different French Mediterranean lagoons for sampling and analyzing the water column
1063parameters, the sediment parameters and the macrofauna. A special thanks to Corinne Pélaprat
1064(STARESO), Sébastien Thorin (CREOCEAN), Thibault Schwartz (CREOCEAN) and Daniel Martin (Centre
1065d'Estudis Avançats de Blanes – CEAB) for sampling, identifying, and analyzing the macrofauna. The
1066authors are also extremely grateful to Aurélie Fouveau (Ifremer, LERBN) for extracting the ecological
1067group values, Olivier Boutron (Tour du Valat) for the information on the Vaccarès lagoon system and
1068Marion Richard (Ifremer, LERLR) for the fruitful discussions. Finally, we would like to thank all our close
1069colleagues from the LERLR (Ophélie Serais, Elise Bellamy, Aurélien Bouquet, Elise Caillard, Nicolas
1070Cimiterra, Christine Felix, Elise Foucault, Camille Gianaroli, Yannick Gueguen, Elise Lacoste, Franck
1071Lagarde, Julie Le Ray, Danièle Martin, Grégory Messian, Dominique Munaron, Hervé Violette) for creating
1072such a cheerful, dynamic, and enthusiastic work environment, despite all the COVID-19 ups and downs.

1073   **Data, scripts, code, and supplementary information availability**

1074   Data, scripts, and code are available online on Zenodo: https://doi.org/10.1101/2022.08.18.504439

1075   Supplementary information is available online on biorxiv as part of the preprint:
1076https://www.biorxiv.org/content/10.1101/2022.08.18.504439v1

1077   **Conflict of interest disclosure**

1078   The authors declare that they comply with the PCI rule of having no financial conflicts of interest in
1079relation to the content of the article.

1080   **Funding**

1081   This work is part of the MALAG project (Effect of eutrophication on the biodiversity of Mediterranean
1082coastal lagoon benthic macrofauna) funded by "Office Français de la Biodiversité" and Ifremer. A. G. J., M.
1083C. and T. L. M. received funding through the MALAG project. All the data presented in this study was
1084collected during the projects "Réseau de Suivi Lagunaire" and Water Framework Directive, which
1085received financial and technical support from Ifremer, "Agence de l'Eau Rhône Méditerranée Corse",
1086"Région Languedoc-Roussillon/Occitanie" and technical support from "Cépralmar".

1087

1088   **References**

1089Aberson, M.J.R., Bolam, S.G., Hughes, R.G., 2016. The effect of sewage pollution on the feeding behaviour
1090   and diet of *Hediste* (*Nereis diversicolor* (O.F. Müller, 1776)) in three estuaries in south-east England,
1091   with implications for saltmarsh erosion. Marine Pollution Bulletin 105, 150–160.
1092   https://doi.org/10.1016/j.marpolbul.2016.02.033
1093Alonso, A., Camargo, J.A., 2006. Toxicity of nitrite to three species of freshwater invertebrates.
1094   Environmental Toxicology 21, 90–94. https://doi.org/10.1002/tox.20155
1095Amanieu, M., Guelorget, O., Michel, P., 1977. Richesse et diversité de la macrofaune benthique d'une
1096   lagune littorale méditerranéenne. Vie et Milieu XXVII, 85–109.
1097Andral, B., Sargian, P., 2010a. Directive Cadre Eau. District Rhône et Côtiers méditerranéens. Contrôle de
surveillance/opérationnel.   Campagne   DCE   2009.   RST.DOP/LER-PAC/10-19.
https://archimer.ifremer.fr/doc/00038/14923/

Andral, B., Sargian, P., 2010b. Directive Cadre Eau. District Corse. Contrôle de surveillance/opérationnel.
Campagne DCE 2009. RST.DOPLER/PAC/10-20. https://archimer.ifremer.fr/doc/00028/13918/

Anonymous, 1958. The Venice System for the classification of marine waters according to salinity.
Limnology and Oceanography 3: 346–347.

Anonymous, 1959. Symposium on the classification of brackish waters. Presented at the Archivio di
Oceanografia e Limnologia, Venice, 8–14 April 1958.

Arocena, R., 2007. Effects of submerged aquatic vegetation on macrozoobenthos in a coastal lagoon of
the southwestern atlantic. International Review of Hydrobiology 92, 33–47.
https://doi.org/10.1002/iroh.200610881

Bachelet, G., de Montaudouin, X., Auby, I., Labourg, P.-J., 2000. Seasonal changes in macrophyte and
macrozoobenthos assemblages in three coastal lagoons under varying degrees of eutrophication. ICES
Journal of Marine Science 57, 1495–1506. https://doi.org/10.1006/jmsc.2000.0902

Barbone, E., Basset, A., 2010. Hydrological constraints to macrobenthic fauna biodiversity in transitional
waters ecosystems. Rend. Fis. Acc. Lincei 21, 301–314. https://doi.org/10.1007/s12210-010-0090-4

Barbone, E., Rosati, I., Reizopoulou, S., Basset, A., 2012. Linking classification boundaries to sources of
natural variability in transitional waters: A case study of benthic macroinvertebrates. Ecological
Indicators, Marine Benthic Indicators 12, 105–122. https://doi.org/10.1016/j.ecolind.2011.04.014

Barnes, R.S.K., 1980. Coastal lagoons: the natural history of a neglected habitat, Cambridge University
Press. ed. Cambridge.

Basset, A., Barbone, E., Borja, Á., Elliott, M., Jona-Lasinio, G., Marques, J., Mazik, K., Muxika, I., Neto, J.,
Reizopoulou, S., Rosati, I., Teixeira, H., 2012. Natural variability and reference conditions: setting type-
specific classification boundaries for lagoon macroinvertebrates in the Mediterranean and Black Seas.
Hydrobiologia. https://doi.org/10.1007/s10750-012-1273-z

Basset, A., Barbone, E., Elliott, M., Li, B.-L., Jorgensen, S.E., Lucena-Moya, P., Pardo, I., Mouillot, D., 2013.
A unifying approach to understanding transitional waters: Fundamental properties emerging from
ecotone ecosystems. Estuarine, Coastal and Shelf Science, Estuarine and lagoon biodiversity and their
natural goods and services 132, 5–16. https://doi.org/10.1016/j.ecss.2012.04.012

Basset, A., Galuppo, N., Sabetta, L., 2007. Environmental heterogeneity and benthic macroinvertebrate
guilds in italian lagoons. Transitional Waters Bulletin 1, 48–63.
https://doi.org/10.1285/i1825226Xv1n1p48

Basset, A., Sabetta, L., Fonnesu, A., Mouillot, D., Do Chi, T., Viaroli, P., Giordani, G., Reizopoulou, S.,
Abbiati, M., Carrada, G.C., 2006. Typology in Mediterranean transitional waters: new challenges and
perspectives. Aquatic Conservation: Marine and Freshwater Ecosystems 16, 441–455. https://doi.org/
10.1002/aqc.767

Basset, A., Sabetta, L., Sangiorgio, F., Pinna, M., Migoni, D., Fanizzi, F., Barbone, E., Galuppo, N., Fonda
Umani, S., Reizopoulou, S., Nicolaidou, A., Arvanitidis, C., Moncheva, S., Trajanova, A., Georgescu, L.,
Beqiraj, S., 2008. Biodiversity conservation in Mediterranean and Black Sea lagoons: a trait-oriented
approach to benthic invertebrate guilds. Aquatic Conservation: Marine and Freshwater Ecosystems
18, S4–S15. https://doi.org/10.1002/aqc.979

Bec, B., Collos, Y., Souchu, P., Vaquer, A., Lautier, J., Fiandrino, A., Benau, L., Orsoni, V., Laugier, T., 2011.
Distribution of picophytoplankton and nanophytoplankton along an anthropogenic eutrophication
gradient in French Mediterranean coastal lagoons. Aquatic Microbial Ecology 63, 29–45.
https://doi.org/10.3354/ame01480

Bendoricchio, G., Di Luzio, M., Baschieri, P., Capodaglio, A.G., 1993. Diffuse pollution in the lagoon of
Venice. Water Science and Technology 28, 69–78. https://doi.org/10.2166/wst.1993.0405

Benjamini, Y., Hochberg, Y., 1995. Controlling the false discovery rate: a practical and powerful approach
to multiple testing. Journal of the Royal Statistical Society. Series B (Methodological) 57, 289–300.
https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Berthelsen, A., Atalah, J., Clark, D., Goodwin, E., Patterson, M., Sinner, J., 2018. Relationships between
biotic indices, multiple stressors and natural variability in New Zealand estuaries. Ecological Indicators
85, 634–643. https://doi.org/10.1016/j.ecolind.2017.10.060

Blanchet, F.G., Legendre, P., Borcard, D., 2008. Forward selection of explanatory variables. Ecology 89,
2623–2632. https://doi.org/10.1890/07-0986.1

Boardman, G.D., Starbuck, S.M., Hudgins, D.B., Li, X., Kuhn, D.D., 2004. Toxicity of ammonia to three
marine fish and three marine invertebrates. Environmental Toxicology 19, 134–142.
https://doi.org/10.1002/tox.20006

Borja, A., Franco, J., Pérez, V., 2000. A marine biotic index to establish the ecological quality of soft-
bottom benthos within European estuarine and coastal environments. Marine Pollution Bulletin 40,
1100–1114. https://doi.org/10.1016/S0025-326X(00)00061-8

Borja, A., Muxika, I., 2005. Guidelines for the use of AMBI (AZTI's Marine Biotic Index) in the assessment
of the benthic ecological quality. Marine Pollution Bulletin 50, 787–789.
https://doi.org/10.1016/j.marpolbul.2005.04.040

Boström, C., Jackson, E.L., Simenstad, C.A., 2006. Seagrass landscapes and their effects on associated
fauna: A review. Estuarine, Coastal and Shelf Science, Ecological and Management Implications on
Seagrass Landscapes 68, 383–403. https://doi.org/10.1016/j.ecss.2006.01.026

Brauko, K.M., Cerveira, A.F., Freitas, R.F., Hadlich, H.L., Beras, T., Fonseca, A.L., Pagliosa, P.R., 2020.
Comparing structural and functional approaches to distinguish benthic communities from eutrophic
and non-eutrophic estuaries. Marine Pollution Bulletin 157, 111290.
https://doi.org/10.1016/j.marpolbul.2020.111290

Brauko, M.K., de Souza, F.M., Muniz, P., de Camargo, M.G., Lana, P. da C., 2015. Spatial variability of
three benthic indices for marine quality assessment in a subtropical estuary of Southern Brazil. Marine
Pollution Bulletin 91, 454–460. https://doi.org/10.1016/j.marpolbul.2014.10.025

Brauns, M., Garcia, X.-F., Walz, N., Pusch, M.T., 2007. Effects of human shoreline development on littoral
macroinvertebrates in lowland lakes. Journal of Applied Ecology 44, 1138–1144.
https://doi.org/10.1111/j.1365-2664.2007.01376.x

Breslow, N., 1970. A generalized Kruskal-Wallis test for comparing k samples subject to unequal patterns
of censorship. Biometrika 57, 579–594. https://doi.org/10.2307/2334776

Caliceti, M., Argese, E., Sfriso, A., Pavoni, B., 2002. Heavy metal contamination in the seaweeds of the
Venice lagoon. Chemosphere 47, 443–454. https://doi.org/10.1016/S0045-6535(01)00292-2

Camargo, J.A., Alonso, A., Salamanca, A., 2005. Nitrate toxicity to aquatic animals: a review with new data
for freshwater invertebrates. Chemosphere 58, 1255–1267.
https://doi.org/10.1016/j.chemosphere.2004.10.044

Cardoso, P.G., Pardal, M.A., Raffaelli, D., Baeta, A., Marques, J.C., 2004. Macroinvertebrate response to
different species of macroalgal mats and the role of disturbance history. Journal of Experimental
Marine Biology and Ecology 308, 207–220. https://doi.org/10.1016/j.jembe.2004.02.018

Carlier, A., Riera, P., Amouroux, J.-M., Bodiou, J.-Y., Escoubeyrou, K., Desmalades, M., Caparros, J.,
Grémare, A., 2007. A seasonal survey of the food web in the La Palme lagoon (northwestern
Mediterranean) assessed by carbon and nitrogen stable isotope analysis. Estuarine, Coastal and Shelf
Science 73, 299–315. https://doi.org/10.1016/j.ecss.2007.01.012

Carvalho, S., Gaspar, M.B., Moura, A., Vale, C., Antunes, P., Gil, O., da Fonseca, L.C., Falcão, M., 2006. The
use of the marine biotic index AMBI in the assessment of the ecological status of the Obidos lagoon
(Portugal). Marine Pollution Bulletin 52, 1414–1424. https://doi.org/10.1016/j.marpolbul.2006.04.004

Chapelle, A., Lazure, P., Souchu, P., 2001. Modélisation numérique des crises anoxiques (malaïgues) dans la lagune de Thau (France). Oceanologica Acta Supplement 1, 87–97. https://doi.org/10.1016/S0399-1784(00)01109-9

Cladas, Y., Papantoniou, G., Bekiari, V., Fragkopoulu, N., 2016. Dystrophic crisis event in Papas Lagoon, Araxos Cape, Western Greece in the summer 2012. Mediterranean Marine Science 17, 32–38. https://doi.org/10.12681/mms.1409

Cloern, J.E., 2001. Our evolving conceptual model of the coastal eutrophication problem. Marine Ecology Progress Series 210, 223–253. https://doi.org/10.3354/meps210223

Cognetti, G., Maltagliati, F., 2000. Biodiversity and adaptive mechanisms in brackish water fauna. Marine Pollution Bulletin 40, 7–14. https://doi.org/10.1016/S0025-326X(99)00173-3

Colden, A.M., Fall, K.A., Cartwright, G.M., Friedrichs, C.T., 2016. Sediment suspension and deposition across restored oyster reefs of varying orientation to flow: implications for restoration. Estuaries and Coasts 39, 1435–1448. https://doi.org/10.1007/s12237-016-0096-y

CORINE Land Cover [WWW Document], 2018. https://land.copernicus.eu/pan-european/corine-land-cover (accessed 12.14.21).

De Casabianca, M.-L., 1996. France — The Mediterranean lagoons, in: Schramm, W., Nienhuis, P.H. (Eds.), Marine benthic vegetation: Recent changes and the effects of eutrophication, Ecological Studies. Springer, Berlin, Heidelberg, pp. 307–329. https://doi.org/10.1007/978-3-642-61398-2_14

de Paz, L., Patrício, J., Marques, J.C., Borja, A., Laborda, A.J., 2008. Ecological status assessment in the lower Eo estuary (Spain). The challenge of habitat heterogeneity integration: a benthic perspective. Marine Pollution Bulletin 56, 1275–1283. https://doi.org/10.1016/j.marpolbul.2008.04.027

Derolez, V., Bec, B., Munaron, D., Fiandrino, A., Pete, R., Simier, M., Souchu, P., Laugier, T., Aliaume, C., Malet, N., 2019. Recovery trajectories following the reduction of urban nutrient inputs along the eutrophication gradient in French Mediterranean lagoons. Ocean & Coastal Management 171, 1–10. https://doi.org/10.1016/j.ocecoaman.2019.01.012

Derolez, V., Fiandrino, A., Munaron, D., Cadoret, M., 2014. Bilan sur les principales pressions pesant sur les lagunes méditerranéennes et leurs liens avec l'état DCE. RST-LER/LR 14-20. https://archimer.ifremer.fr/doc/00254/36574/

Derolez, V., Leurion, A., Fiandrino, A., 2012. Directive Cadre Eau. Estimation du degré d'échange avec la mer pour les lagunes du bassin Rhône Méditerranée & Corse. Agence de l'eau, Ref. RST. DOP/LER-LR/ 12-10, 36p. https://archimer.ifremer.fr/doc/00094/20558/

Derolez, V., Malet, N., Fiandrino, A., Lagarde, F., Richard, M., Ouisse, V., Bec, B., Aliaume, C., 2020a. Fifty years of ecological changes: Regime shifts and drivers in a coastal Mediterranean lagoon during oligotrophication. Science of The Total Environment 732, 139292. https://doi.org/10.1016/j.scitotenv.2020.139292

Derolez, V., Soudant, D., Malet, N., Chiantella, C., Richard, M., Abadie, E., Aliaume, C., Bec, B., 2020b. Two decades of oligotrophication: Evidence for a phytoplankton community shift in the coastal lagoon of Thau (Mediterranean Sea, France). Estuarine, Coastal and Shelf Science 241, 106810. https://doi.org/10.1016/j.ecss.2020.106810

Díaz, R., Rosenberg, R., 1995. Marine benthic hypoxia: A review of its ecological effects and the behavioural responses of benthic macrofauna, in: Oceanography and Marine Biology : An Annual Review. University College London Press, London, UK, pp. 245–303.

Donadi, S., van der Zee, E.M., van der Heide, T., Weerman, E.J., Piersma, T., van de Koppel, J., Olff, H., Bartelds, M., van Gerwen, I., Eriksson, B.K., 2014. The bivalve loop: Intra-specific facilitation in burrowing cockles through habitat modification. Journal of Experimental Marine Biology and Ecology 461, 44–52. https://doi.org/10.1016/j.jembe.2014.07.019

Dufrêne, M., Legendre, P., 1997. Species assemblages and indicator species:the need for a flexible
asymmetrical approach. Ecological Monographs 67, 345–366. https://doi.org/10.1890/0012-
9615(1997)067[0345:SAAIST]2.0.CO;2

Elliott, M., Quintino, V., 2007. The estuarine quality paradox, environmental homeostasis and the
difficulty of detecting anthropogenic stress in naturally stressed areas. Marine Pollution Bulletin 54,
640–645. https://doi.org/10.1016/j.marpolbul.2007.02.003

Faulwetter, S., Markantonatou, V., Pavloudi, C., Papageorgiou, N., Keklikoglou, K., Chatzinikolaou, E.,
Pafilis, E., Chatzigeorgiou, G., Vasileiadou, K., Dailianis, T., Fanini, L., Koulouri, P., Arvanitidis, C., 2014.
Polytraits: A database on biological traits of marine polychaetes. Biodiversity Data Journal 2, e1024.
https://doi.org/10.3897/BDJ.2.e1024

Fiandrino, A., Ouisse, V., Dumas, F., Lagarde, F., Pete, R., Malet, N., Le Noc, S., de Wit, R., 2017. Spatial
patterns in coastal lagoons related to the hydrodynamics of seawater intrusion. Marine Pollution
Bulletin 119, 132–144. https://doi.org/10.1016/j.marpolbul.2017.03.006

Gamito, S., Gilabert, J., Marcos Diego, C., Pérez-Ruzafa, A., 2005. Effects of changing environmental
conditions on lagoon ecology, in: Gonenc, I., Wolfin, J. (Eds.), Coastal Lagoons – Ecosystem Processes
and Modeling for Sustainable Use and Development. CRC Press, pp. 193–229.

Giampaoletti, J., Sbrana, A., Magni, P., Gravina, M.F., 2023. Macrobenthos of the Tortolì lagoon: a
peculiar case of high benthic biodiversity among Mediterranean lagoons. Diversity 15, 783.
https://www.mdpi.com/1424-2818/15/6/783

González-Ortiz, V., Egea, L.G., Jiménez-Ramos, R., Moreno-Marín, F., Pérez-Lloréns, J.L., Bouma, T.J., Brun,
F.G., 2014. Interactions between seagrass complexity, hydrodynamic flow and biomixing alter food
availability for associated filter-feeding organisms. PLOS ONE 9, e104949.
https://doi.org/10.1371/journal.pone.0104949

Guelorget, O., Michel, P., 1979. Les peuplements benthiques d'un étang littoral Languedocien, l'étang du
Prévost (Hérault): 2. Étude quantitative de la macrofaune des sables. Téthys.

Guelorget, O., Perthuisot, J., Lamy, N., Lefebvre, A., 1994. Structure et organisation de l'étang de Thau
d'après la faune benthique (macrofaune, méiofaune). Relations avec le confinement. Oceanologica
Acta 17, 105–114.

Guelorget, O., Perthuisot, J.P., 1983. Le domaine paralique : expressions géologiques, biologiques et
économiques du confinement, Travaux du laboratoire de géologie, n°16. Presses de l'école normale
supérieure, Paris.

Harman, W.N., 1972. Benthic substrates: their effect on fresh-water mollusca. Ecology 53, 271–277.
https://doi.org/10.2307/1934081

Hebbali, A., 2020. olsrr: tools for building ols regression models.

Kjerfve, B., 1994. Coastal lagoon processes. Elsevier Oceanography Series 60. Elsevier ed. Amsterdam,
London, New York, Tokyo. 577 p.

James, G., Witten, D., Hastie, T., and Tibshirani, R. (eds.). (2013). An introduction to statistical learning:
with applications in R. New York: Springer.

Jessen, C., Bednarz, V., Rix, L., Teichberg, M., Wild, C., 2015. Marine Eutrophication, in: Environmental
Indicators. p. 27. https://doi.org/10.1007/978-94-017-9499-2_11

Jones, A.G., Dubois, S.F., Desroy, N., Fournier, J., 2021. Intertidal ecosystem engineer species promote
benthic-pelagic coupling and diversify trophic pathways. Marine Ecology Progress Series 660, 119–
139. https://doi.org/10.3354/meps13600

Jones, A.G., Schaal, G., Boyé, A., Creemers, M., Derolez, V., Desroy, N., Fiandrino, A., Mouton, T.L., Simier,
M., Smith, N., Ouisse, V., 2023. Data, scripts and code for the preprint "Disentangling the effects of
eutrophication and natural variability on macrobenthic communities across French coastal lagoons."
https://doi.org/10.1101/2022.08.18.504439

Josse, J., Husson, F., 2016. missMDA: a package for handling missing values in multivariate data analysis.
Journal of Statistical Software 70, 1–31. https://doi.org/10.18637/jss.v070.i01

Josse, J., Husson, F., 2012. Handling missing values in exploratory multivariate data analysis methods.
Journal de la Société Française de Statistique 153, 79–99.

Kassambara, A., 2020. rstatix: pipe-friendly framework for basic statistical tests.

Kermagoret, C., Claudet, J., Derolez, V., Nugues, M.M., Ouisse, V., Quillien, N., Baulaz, Y., Le Mao, P.,
Scemama, P., Vaschalde, D., Bailly, D., Mongruel, R., 2019. How does eutrophication impact bundles of
ecosystem services in multiple coastal habitats using state-and-transition models. Ocean & Coastal
Management 174, 144–153. https://doi.org/10.1016/j.ocecoaman.2019.03.028

Kjerfve, B., 1994. Chapter 1 Coastal Lagoons, in: Kjerfve, B. (Ed.), Elsevier Oceanography Series, Coastal
Lagoon Processes. Elsevier, pp. 1–8. https://doi.org/10.1016/S0422-9894(08)70006-0

Lacoste, E., Jones, A., Callier, M., Klein, J., Lagarde, F., Derolez, V., 2023. A review of knowledge on the
impacts of multiple anthropogenic pressures on the soft-bottom benthic ecosystem in Mediterranean
coastal lagoons. Estuaries and Coasts. https://doi.org/10.1007/s12237-023-01188-9

Landman, M.J., Van Den Heuvel, M.R., Ling, N., 2005. Relative sensitivities of common freshwater fish and
invertebrates to acute hypoxia. New Zealand Journal of Marine and Freshwater Research 39, 1061–
1067. https://doi.org/10.1080/00288330.2005.9517375

Lasserre, P., 1979. Coastal lagoons: sanctuary ecosystems, cradles of culture, targets for economic
growth. Nature and Resources.

Le Fur, I., De Wit, R., Plus, M., Oheix, J., Simier, M., Ouisse, V., 2018. Submerged benthic macrophytes in
Mediterranean lagoons: distribution patterns in relation to water chemistry and depth. Hydrobiologia
808, 175–200. https://doi.org/10.1007/s10750-017-3421-y

Le Fur, I., Wit, R.D., Plus, M., Oheix, J., Derolez, V., Simier, M., Malet, N., Ouisse, V., 2019. Re-
oligotrophication trajectories of macrophyte assemblages in Mediterranean coastal lagoons based on
17-year time-series. Marine Ecology Progress Series 608, 13–32. https://doi.org/10.3354/meps12814

Legendre, P., Legendre, L., 2012. Numerical ecology, 3rd ed. Elsevier.

Lepoint, G., Nyssen, F., Gobert, S., Dauby, P., Bouquegneau, J.-M., 2000. Relative impact of a seagrass bed
and its adjacent epilithic algal community in consumer diets. Marine Biology 136, 513–518.
https://doi.org/10.1007/s002270050711

Leruste, A., Malet, N., Munaron, D., Derolez, V., Hatey, E., Collos, Y., De Wit, R., Bec, B., 2016. First steps
of ecological restoration in Mediterranean lagoons: Shifts in phytoplankton communities. Estuarine,
Coastal and Shelf Science 180, 190–203. https://doi.org/10.1016/j.ecss.2016.06.029

Levin, L.A., Boesch, D.F., Covich, A., Dahm, C., Erséus, C., Ewel, K.C., Kneib, R.T., Moldenke, A., Palmer,
M.A., Snelgrove, P., Strayer, D., Weslawski, J.M., 2001. The function of marine critical transition zones
and the importance of sediment biodiversity. Ecosystems 4, 430–451.
https://doi.org/10.1007/s10021-001-0021-4

Magalhães, L., Pires, A., Velez, C., Martins, R., Figueira, E., Soares, A.M.V.M., Freitas, R., 2019. Seasonal
and spatial alterations in macrofaunal communities and in *Nephtys cirrosa* (Polychaeta) oxidative
stress under a salinity gradient: A comparative field monitoring approach. Ecological Indicators 96,
192–201. https://doi.org/10.1016/j.ecolind.2018.08.045

Magni, P., Como, S., Kamijo, A., Montani, S., 2017. Effects of *Zostera marina* on the patterns of spatial
distribution of sediments and macrozoobenthos in the boreal lagoon of Furen (Hokkaido, Japan).
Marine Environmental Research 131, 90–102. https://doi.org/10.1016/j.marenvres.2017.09.013

Magni, P., Semprucci, F., Gravina, M.F., 2022. Joint analysis of macrofaunal and meiofaunal assemblages
improves the assessment of lagoonal environmental heterogeneity. Estuarine, Coastal and Shelf
Science 266, 107740. https://doi.org/10.1016/j.ecss.2021.107740

Magni, P., Gravina, M.F., 2023. Macrobenthos of lagoon ecosystems: a comparison in vegetated and bare
sediments. Advances in Oceanography and Limnology 14. https://doi.org/10.4081/aiol.2023.11124

Mancinelli, G., Rossi, L., 2001. Indirect, size-dependent effects of crustacean mesograzers on the
Rhodophyta *Gracilaria verrucosa* (Hudson) Papenfuss: evidence from a short-term study in the Lesina
Lagoon (Italy). Marine Biology 138, 1163–1173. https://doi.org/10.1007/s002270100545

Martini, S., Larras, F., Boyé, A., Faure, E., Aberle, N., Archambault, P., Bacouillard, L., Beisner, B.E., Bittner,
L., Castella, E., Danger, M., Gauthier, O., Karp-Boss, L., Lombard, F., Maps, F., Stemmann, L., Thiébaut,
E., Usseglio-Polatera, P., Vogt, M., Laviale, M., Ayata, S.-D., 2021. Functional trait-based approaches as
a common framework for aquatic ecologists. Limnology and Oceanography 66, 965–994.
https://doi.org/10.1002/lno.11655

Menu, M., Sandrine, V., Bajjouk, T., Derolez, V., Fiandrino, A., Giraud, A., Grillas, P., Ouisse, V., 2019.
Rapport final du projet CHAMILA (Cartographie des habitats en milieu lagunaire méditerranéen)
R.ODE/UL/LER/LR/19.34. https://doi.org/10.13155/70545

Mistri, M., Borja, A., Aleffi, I.F., Lardicci, C., Tagliapietra, D., Munari, C., 2018. Assessing the ecological
status of Italian lagoons using a biomass-based index. Marine Pollution Bulletin 126, 600–605. https://
doi.org/10.1016/j.marpolbul.2017.09.048

Miyamoto, Y., Nakano, T., Yamada, K., Hatakeyama, K., Hamaguchi, M., 2019. Combined effects of drift
macroalgal bloom and warming on occurrence and intensity of diel-cycling hypoxia in a eutrophic
coastal lagoon. Estuaries and Coasts 42, 494–503. https://doi.org/10.1007/s12237-018-0484-6

Morris, E.K., Caruso, T., Buscot, F., Fischer, M., Hancock, C., Maier, T.S., Meiners, T., Müller, C.,
Obermaier, E., Prati, D., Socher, S.A., Sonnemann, I., Wäschke, N., Wubet, T., Wurst, S., Rillig, M.C.,
2014. Choosing and using diversity indices: insights for ecological applications from the German
Biodiversity Exploratories. Ecol Evol 4, 3514–3524. https://doi.org/10.1002/ece3.1155

Muxika, I., Borja, Á., Bald, J., 2007. Using historical data, expert judgement and multivariate analysis in
assessing reference conditions and benthic ecological status, according to the European Water
Framework Directive. Marine Pollution Bulletin, Implementation of the Water Framework Directive in
European marine waters 55, 16–29. https://doi.org/10.1016/j.marpolbul.2006.05.025

Neveux, J., Lantoine, F., 1993. Spectrofluorometric assay of chlorophylls and phaeopigments using the
least squares approximation technique. Deep Sea Research Part I: Oceanographic Research Papers 40,
1747–1765. https://doi.org/10.1016/0967-0637(93)90030-7

Newton, A., Oliveira, P.S., Icely, J.D., Foster, P.A., 2010. Monitoring of oxygen condition in the Ria
Formosa coastal lagoon, Portugal. J Environ Monit 12, 355–360. https://doi.org/10.1039/b914015h

Nordström, M., Booth, D.M., 2007. Drift algae reduce foraging efficiency of juvenile flatfish. Journal of Sea
Research 58, 335–341. https://doi.org/10.1016/j.seares.2007.08.001

Norkko, J., Bonsdorff, E., Norkko, A., 2000. Drifting algal mats as an alternative habitat for benthic
invertebrates: Species specific responses to a transient resource. Journal of Experimental Marine
Biology and Ecology 248, 79–104. https://doi.org/10.1016/S0022-0981(00)00155-6

Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.R., O'Hara, R.B.,
Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H., 2019. vegan: Community Ecology
Package.

Ouisse, V., Riera, P., Migné, A., Leroux, C., Davoult, D., 2011. Freshwater seepages and ephemeral
macroalgae proliferation in an intertidal bay: I Effect on benthic community structure and food web.
Estuarine, Coastal and Shelf Science 91, 272–281. https://doi.org/10.1016/j.ecss.2010.10.034

Pasqualini, V., Derolez, V., Garrido, M., Orsoni, V., Baldi, Y., Etourneau, S., Leoni, V., Rébillout, P., Laugier,
T., Souchu, P., Malet, N., 2017. Spatiotemporal dynamics of submerged macrophyte status and
watershed exploitation in a Mediterranean coastal lagoon: Understanding critical factors in ecosystem

degradation and restoration. Ecological Engineering 102, 1–14. https://doi.org/10.1016/j.ecoleng.2017.01.027

Paul, J., Nestlerode, J.A., Jarvis, B.M., 2023. Evaluating the effectiveness of M-AMBI with other biotic indexes in a temperate estuary. Marine Pollution Bulletin, 193: 115194. https://doi.org/10.1016/j.marpolbul.2023.115194

Pearson, T.H., Rosenberg, R., 1978. Macrobenthic succession in relation to organic enrichment and pollution of the marine environment. Oceanography and Marine Biology: An Annual Review 16, 229–311. https://doi.org/10.2983/035.034.0121u1.10

Pelletier, M.C., Charpentier, M., 2023. Assessing relative importance of stressors to the benthic index, M-AMBI: An example from U.S. estuaries. Marine Pollution Bulletin, 186:114456. https://doi.org/10.1016/j.marpolbul.2022.114456

Pérez-Ruzafa, A., Marcos, C., Pérez-Ruzafa, I.M., 2011b. Recent advances in coastal lagoons ecology: evolving old ideas and assumptions. Transitional Waters Bulletin 5, 50–74. https://doi.org/10.1285/i1825229Xv5n1p50

Pérez-Ruzafa, A., Marcos, C., Pérez-Ruzafa, I.M., 2011a. Mediterranean coastal lagoons in an ecosystem and aquatic resources management context. Physics and Chemistry of the Earth, Parts A/B/C, Water in the Mediterranean Basin 36, 160–166. https://doi.org/10.1016/j.pce.2010.04.013

Pérez-Ruzafa, A., Marcos, C., Pérez-Ruzafa, I.M., Barcala, E., Hegazi, M.I., Quispe, J., 2007a. Detecting changes resulting from human pressure in a naturally quick-changing and heterogeneous environment: Spatial and temporal scales of variability in coastal lagoons. Estuarine, Coastal and Shelf Science, Biodiversity and Ecosystem Functioning in Coastal and Transitional Waters 75, 175–188. https://doi.org/10.1016/j.ecss.2007.04.030

Pérez-Ruzafa, A., Marcos, C., Pérez-Ruzafa, I.M., Pérez-Marcos, M., 2013. Are coastal lagoons physically or biologically controlled ecosystems ? Revisiting r vs. K strategies in coastal lagoons and estuaries. Estuarine, Coastal and Shelf Science, Estuarine and lagoon biodiversity and their natural goods and services 132, 17–33. https://doi.org/10.1016/j.ecss.2012.04.011

Pérez-Ruzafa, Angel, Marcos, C., Pérez-Ruzafa, I.M., Pérez-Marcos, M., 2011c. Coastal lagoons: "transitional ecosystems" between transitional and coastal waters. Journal of Coastal Conservation 15, 369–392. https://doi.org/10.1007/s11852-010-0095-2

Perez-Ruzafa, A., Marcos-Diego, C., 1992. Colonization rates and dispersal as essential parameters in the confinement theory to explain the structure and horizontal zonation of lagoon benthic assemblages. Technical Report Rapport Commission International Mer Medit 33.

Pérez-Ruzafa, Angel, Mompeán, M.C., Marcos, C., 2007b. Hydrographic, geomorphologic and fish assemblage relationships in coastal lagoons. Hydrobiologia 577, 107–125. https://doi.org/10.1007/s10750-006-0421-8

Perthuisot, J.-P., Guelorget, O., 1983. Le confinement, paramètre essentiel de la dynamique biologique du domaine paralique. Sciences Géologiques, bulletins et mémoires 36, 239–248. https://doi.org/10.3406/sgeol.1983.1642

Pitacco, V., Reizopoulou, S., Sfriso, Adriano, Sfriso, Andrea, Mistri, M., Munari, C., 2019. The difficulty of disentangling natural from anthropogenic forcing factors makes the evaluation of ecological quality problematic: A case study from Adriatic lagoons. Marine Environmental Research 150, 104756. https://doi.org/10.1016/j.marenvres.2019.104756

Pollice, A., Arima, S., Lasinio G.J., Basset, A., Rosati, I., 2015. Bayesian analysis of three indices for lagoons ecological status evaluation. Stochastic Environmental Research and Risk Assessment, 29: 477-485. DOI: 10.1007/s00477-014-0885-4

Provost, C., Gevrey, M., Argillier, C., 2012. La macrofaune benthique indicatrice de l'état des lagunes oligohalines méditerranéennes ? [Rapport de recherche] IRSTEA. 2012, pp.37. https://hal.science/hal-02597249/

Puente, A., Diaz, R.J., 2008. Is it possible to assess the ecological status of highly stressed natural estuarine environments using macroinvertebrates indices? Marine Pollution Bulletin 56, 1880–1889. https://doi.org/10.1016/j.marpolbul.2008.07.016

Quillien, N., Nordström, M.C., Gauthier, O., Bonsdorff, E., Paulet, Y.-M., Grall, J., 2015. Effects of macroalgal accumulations on the variability in zoobenthos of high-energy macrotidal sandy beaches. Marine Ecology Progress Series 522, 97–114. https://doi.org/10.3354/meps11151

Quillien, N., Nordström, M.C., Schaal, G., Bonsdorff, E., Grall, J., 2016. Opportunistic basal resource simplifies food web structure and functioning of a highly dynamic marine environment. Journal of Experimental Marine Biology and Ecology 477, 92–102. https://doi.org/10.1016/j.jembe.2016.01.010

R Core Team, 2020. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Reizopolou, S., Penna, M., Boix, D., Buchet, R., Costas, N., Derolez, V., Gascon, S., Gifre, J., Martinoy, M., Pardo, I., Quintana, X., Trabucco, B., Sala, J., Salas Herrero, F., 2018. Transitional waters Mediterranean geographic intercalibration group: Benthic invertebrates fauna ecological assessment methods, EUR 29561 EN, Publications Office of the European Union, Luxembourg, 2018, ISBN 978-92-79-98373-3, JRC114720. https://doi.org/10.2760/625400

Reizopoulou, S., Nicolaidou, A., 2007. Index of size distribution (ISD): a method of quality assessment for coastal lagoons. Hydrobiologia 577, 141–149. https://doi.org/10.1007/s10750-006-0423-6

Reizopoulou, S., Simboura, N., Sigala, K., Barbone, E., Aleffi, F., Kaisakis, G., Rosati, I., Basset, A., Nicolaidou, A., 2014. Assessing the ecological status of Mediterranean coastal lagoons using macroinvertebrates. Comparison of the most commonly used methods. Mediterranean Marine Science 15, 602–612. https://doi.org/10.12681/mms.606

Riedel, B., Zuschin, M., Stachowitsch, M., 2012. Tolerance of benthic macrofauna to hypoxia and anoxia in shallow coastal seas: a realistic scenario. Marine Ecology Progress Series 458, 39–52. https://doi.org/10.3354/meps09724

Rigaud, S., Deflandre, B., Grenz, C., Cesbron, F., Pozzato, L., Voltz, B., Grémare, A., Romero-Ramirez, A., Mirleau, P., Meulé, S., Faure, V., Mayot, N., Michotey, V., Bonin, P., Pascal, L., Cordier, M.-A., Lamarque, B., Tenório, M., Radakovitch, O., 2021. Benthic oxygen dynamics and implication for the maintenance of chronic hypoxia and ecosystem degradation in the Berre lagoon (France). Estuarine, Coastal and Shelf Science 258, 107437. https://doi.org/10.1016/j.ecss.2021.107437

Roberts, D.W., 2019. labdsv: ordination and multivariate analysis for ecology.

Robertson, B.P., Savage, C., Gardner, J.P.A., Robertson, B.M., Stevens, L.M., 2016. Optimising a widely-used coastal health index through quantitative ecological group classifications and associated thresholds. Ecological Indicators 69, 595–605. https://doi.org/10.1016/j.ecolind.2016.04.003

Roques, C., Grousset, E., Troussellier, M., Hermet, S., Le Carrer, J., Sar, C., Caro, A., 2020. A trade-off between mucocytes and bacteriocytes in *Loripes orbiculatus* gills (Bivalvia, Lucinidae): a mixotrophic adaptation to seasonality and reproductive status in a symbiotic species? Marine Biology 167. https://doi.org/10.1007/s00227-020-03768-w

Rose, J.M., Caron, D.A., 2007. Does low temperature constrain the growth rates of heterotrophic protists? Evidence and implications for algal blooms in cold waters. Limnology and Oceanography 52, 886–895. https://doi.org/10.4319/lo.2007.52.2.0886

Rosenberg, R., Loo, L.-O., Möller, P., 1992. Hypoxia, salinity and temperature as structuring factors for marine benthic communities in a eutrophic area. Netherlands Journal of Sea Research 30, 121–129. https://doi.org/10.1016/0077-7579(92)90051-F

Rossi, F., Castelli, A., Lardicci, C., 2006. Distribution of macrobenthic assemblages along a marine gradient
in Mediterranean eutrophic coastal lagoons. Marine ecology 27, 66–75.
https://doi.org/10.1111/j.1439-0485.2006.00078.x

Sabetta, L., Barbone, E., Giardino, A., Galuppo, N., Basset, A., 2007. Species–area patterns of benthic
macro-invertebrates in Italian lagoons. Hydrobiologia 577, 127–139. https://doi.org/10.1007/s10750-
006-0422-7

Scaps, P., 2002. A review of the biology, ecology and potential use of the common ragworm *Hediste*
diversicolor (O.F. Müller) (Annelida: Polychaeta). Hydrobiologia 470, 203–218.
https://doi.org/10.1023/A:1015681605656

Schielzeth, H., 2010. Simple means to improve the interpretability of regression coefficients. Methods in
Ecology and Evolution 1, 103–113. https://doi.org/10.1111/j.2041-210X.2010.00012.x

Sigala, K., Reizopoulou, S., Basset, A., Nicolaidou, A., 2012. Functional diversity in three Mediterranean
transitional water ecosystems. https://doi.org/10.1016/J.ECSS.2012.06.002

Sigovini, M., Keppel, E., Tagliapietra, D., 2013. M-AMBI revisited: looking inside a widely-used benthic
index. Hydrobiologia 717, 41–50. https://doi.org/10.1007/s10750-013-1565-y

Souchu, P., Bec, B., Smith, V.H., Laugier, T., Fiandrino, A., Benau, L., Orsoni, V., Collos, Y., Vaquer, A.,
2010. Patterns in nutrient limitation and chlorophyll a along an anthropogenic eutrophication gradient
in French Mediterranean coastal lagoons. Can. J. Fish. Aquat. Sci. 67, 743–753.
https://doi.org/10.1139/F10-018

Souchu, P., Gasc, A., Collos, Y., Vaquer, A., Tournier, H., Bibent, B., Deslous-Paoli, J.-M., 1998.
Biogeochemical aspects of bottom anoxia in a Mediterranean lagoon (Thau, France). Marine Ecology-
progress Series 164, 135–146. https://doi.org/10.3354/meps164135

Stendera, S., Johnson, R.K., 2008. Habitat-specific stability and persistence of benthic invertebrate
communities in boreal lakes. https://doi.org/10.1127/1863-9135/2008/0171-0311

Tagliapietra, D., Sigovini, M., Ghirardini, A., 2009. A review of terms and definitions to categorise
estuaries, lagoons and associated environments. https://doi.org/10.1071/MF08088

Tagliapietra, D., Sigovini, M., Magni, P., 2012. Saprobity: a unified view of benthic succession models for
coastal lagoons. Hydrobiologia 686, 15–28. https://doi.org/10.1007/s10750-012-1001-8

Tournoud, M.-G., Payraudeau, S., Cernesson, F., Salles, C., 2006. Origins and quantification of nitrogen
inputs into a coastal lagoon: Application to the Thau lagoon (France). Ecological Modelling, Special
Issue on Southern European Coastal Lagoons 193, 19–33.
https://doi.org/10.1016/j.ecolmodel.2005.07.038

Trombetta, T., Vidussi, F., Mas, S., Parin, D., Simier, M., Mostajir, B., 2019. Water temperature drives
phytoplankton blooms in coastal waters. PLOS ONE 14, e0214933.
https://doi.org/10.1371/journal.pone.0214933

Vanda Orro, C., Cabana, D., 2021. Seasonal implications of the benthic ecological status assessment in a
protected Mediterranean coastal lagoon: Stagno Longu of Posada (Sardinia, Italy).
https://doi.org/10.1016/J.RSMA.2021.101759

Vaquer-Sunyer, R., Duarte, C.M., 2011. Temperature effects on oxygen thresholds for hypoxia in marine
benthic organisms. Global Change Biology 17, 1788–1797. https://doi.org/10.1111/j.1365-
2486.2010.02343.x

Venables, W.N., Ripley, B.D., 2002. Modern Applied Statistics with S, Fourth. ed. Springer, New York.

Vergara-Chen, C., González-Wangüemert, M., Marcos, C., Pérez-Ruzafa, Á., 2013. Small-scale genetic
structure of *Cerastoderma glaucum* in a lagoonal environment: potential significance of habitat
discontinuity and unstable population dynamics. Journal of Molluscan Studies 79, 230–240.
https://doi.org/10.1093/mollus/eyt015

Vernberg, F.J., 1982. Environmental adaptation to lagoon systems. Oceanologica Acta, Special issue.

Vignes, F., Barbone, E., Breber, P., D'Adamo, R., Leonilde, R., Ungaro, N., Focardi, S., Renzi, M., Basset, A.,
2010. Spatial and temporal description of the dystrophic crisis in Lesina lagoon during summer 2008.
Transitional Waters Bulletin 3, 47–62. https://doi.org/10.1285/i1825229Xv3n2p47

Vizzini, S., Mazzola, A., 2006. Sources and transfer of organic matter in food webs of a Mediterranean
coastal environment: Evidence for spatial variability. Estuarine, Coastal and Shelf Science 66, 459–467.
https://doi.org/10.1016/j.ecss.2005.10.004

Ware, C., Dijkstra, J.A., Mello, K., Stevens, A., O'Brien, B., Ikedo, W., 2019. A novel three-dimensional
analysis of functional architecture that describes the properties of macroalgae as a refuge. Marine
Ecology Progress Series 608, 93–103. https://doi.org/10.3354/meps12800

Wei, T., Simko, V., 2017. corrplot: visualization of a correlation matrix.

Weissberger, E.J., Coiro, L.L., Davey, E.W., 2009. Effects of hypoxia on animal burrow construction and
consequent effects on sediment redox profiles. Journal of Experimental Marine Biology and Ecology
371, 60–67. https://doi.org/10.1016/j.jembe.2009.01.005

Wilke, M., Boutiere, H., 2000. Hydrobiological, physical and chemical characteristics and spatio-temporal
dynamics of an oligotrophic mediterranean lagoon: the étang de la Palme (France). Vie et Milieu / Life
& Environment, 101-115

Zaldívar, J., Viaroli, P., Newton, A., Wit, R., Ibañez, C., Reizopoulou, S., Somma, F., Razinkovas, A., Basset,
A., Holmer, M., Murray, N., 2008. Eutrophication in transitional waters: an overview.
https://doi.org/10.1285/I18252273V2N1P1

Zilius, M., Giordani, G., Petkuviene, J., Lubiene, I., Ruginis, T., Bartoli, M., 2015. Phosphorus mobility
under short-term anoxic conditions in two shallow eutrophic coastal systems (Curonian and Sacca di
Goro   lagoons).   Estuarine,   Coastal   and   Shelf   Science   164,   134–146.
https://doi.org/10.1016/j.ecss.2015.07.004

1539