1

Disentangling the effects of eutrophication and natural variability on macrobenthic communities across French coastal lagoons

Auriane G. Jones*[1], Gauthier Schaal[2], Aurélien Boyé[3], Marie Creemers[1], Valérie Derolez[1], Nicolas Desroy[4], Annie Fiandrino[1], Théophile L. Mouton[1], Monique Simier[1], Niamh Smith[1], Vincent Ouisse[1]

[1] MARBEC, Univ. Montpellier, CNRS, Ifremer, IRD, Sète, France

[2] Univ Brest, CNRS, IRD, Ifremer, LEMAR, 29280 Plouzané, France

[3] IFREMER, Centre de Bretagne, DYNECO, Laboratoire d'Ecologie Benthique Côtière (LEBCO), 29280 Plouzané, France

[4] IFREMER, Laboratoire Environnement et Ressources Bretagne nord, 35801 Dinard cedex, France

*Corresponding author

Correspondence: jones.ecology@gmail.com

ABSTRACT

Coastal lagoons are transitional ecosystems that host a unique diversity of species and support many ecosystem services. Owing to their position at the interface between land and sea, they are also subject to increasing human impacts, which alter their ecological functioning. Because coastal lagoons are naturally highly variable in their environmental conditions, disentangling the effects of anthropogenic disturbances like eutrophication from those of natural variability is a challenging, yet necessary issue to address. Here, we analyze a dataset composed of macrobenthic invertebrate abundances and environmental variables (hydro-morphology, water, sediment and macrophytes) gathered across 29 Mediterranean coastal lagoons located in France, to characterize the main drivers of community composition and structure. Using correlograms, linear models and variance partitioning, we found that lagoon hydro-morphology (connection to the sea and lagoon surface), which affects the level of environmental variability (salinity and temperature), as well as lagoon-scale benthic habitat diversity (using macrophyte morphotypes) seemed to regulate macrofauna distribution, while eutrophication and associated stressors like low dissolved oxygen, acted upon the existing communities, mainly by reducing species richness and diversity. Furthermore, M-AMBI, a multivariate index composed of species richness, Shannon diversity and AMBI (AZTI's Marine Biotic Index) and currently used to evaluate the ecological state of French coastal lagoons, was more sensitive to eutrophication (18%) than to natural variability (9%), with nonetheless 49% of its variability explained jointly by both. To improve the robustness of benthic indicators like M-AMBI and increase the effectiveness of lagoon benthic habitat management, we call for a revision of the ecological groups at the base of the AMBI index and of the current lagoon typology which could be inspired by the lagoon-sea connection levels used in this study.

Keywords: Mediterranean Sea, macrofauna, taxonomic diversity, M-AMBI, macrophyte morphotypes, variance partitioning

Introduction

Coastal lagoons are inland, shallow and transitional water bodies, usually oriented parallel to the coast, separated from the sea by a barrier, connected to the sea by one or more restricted inlets which remain open at least intermittently (Kjerfve, 1994). These ecosystems represent 13% of the world's coastline (Barnes, 1980) and provide essential ecosystem services, especially through nutrient regulation and sequestration as well as climate regulation (Kermagoret et al., 2019; Levin et al., 2001). These ecosystems are characterized by strong salinity variations at different temporal scales, ranging from fortnights to seasons and decades (Pérez-Ruzafa et al., 2007a), that are mainly governed by short-term freshwater inflow, seasonal temperature variations, the degree of connection of each water mass with the sea and decadal climatic variations (rainfall and temperatures) (Barbone and Basset, 2010; De Casabianca, 1996; Zaldívar et al., 2008). Oxygen saturation naturally fluctuates in lagoons, due to water temperature and re-aeration processes controlled by wind conditions (Chapelle et al., 2001; Derolez et al., 2020a). Oxygen dynamics are further affected by primary production, which is mainly governed, in oligo and mesotrophic lagoons by seasonal benthic macrophyte growth and decay cycles, and in hyper and eutrophic lagoons by pelagic phytoplankton dynamics (Zaldívar et al., 2008). Overall, coastal lagoons are naturally highly dynamic systems in terms of environmental factors, representing a form of natural environmental stress for any organism living in these systems like fish and benthic invertebrates (Pérez-Ruzafa et al., 2007b; Zaldívar et al., 2008). Consequently, lagoon benthic communities, just like estuarine ones, generally present low species richness and high abundances of tolerant species, characteristics typical of disturbed or stressed coastal ecosystems, giving rise to the "transitional waters quality paradox" (Zaldívar et al., 2008), a generalization of the "Estuarine quality paradox" (Elliott and Quintino, 2007). These paradoxes highlight the problems faced by transitional waters concerning the development of methodologies permitting the evaluation of their ecological quality status, because it results from both anthropogenic pressures and natural characteristics.

Coastal lagoons have also been intensely used and modified by humans for food (fishing and aquaculture), transportation (canals, ports and inlet modification), livelihood (land reclamation for construction) and recreation (fishing, sailing, swimming) (Lasserre, 1979; Pérez-Ruzafa et al., 2011a). The position of coastal lagoons at the interface between land and sea further exposes them to human land-derived pollution such as nitrogen, phosphorous, pesticide, or heavy metal pollution (Caliceti et al., 2002; Derolez et al., 2019; Souchu et al., 2010). The oldest and most studied is nitrogen and phosphorous pollution linked to agriculture and urban development (Bendoricchio et al., 1993; Kjerfve, 1994; Lacoste et al., 2023; Tournoud et al., 2006), which can lead to eutrophication of coastal lagoons, a phenomenon associated to an increase in phytoplankton biomass (Cloern, 2001; Zaldívar et al., 2008), a replacement of long-lived submerged macrophytes like marine plants, by emergent free-living benthic macroalgae and even to the disappearance of all benthic macrophytes (Le Fur et al., 2019, 2018). The high light availability linked to shallow depths and the naturally high nutrient concentration of water and sediments (*i.e.* coastal lagoons can be considered as "naturally eutrophic" (De Casabianca, 1996; Zaldívar et al., 2008)) promote the development of structurally complex benthic macrophytes, which are characteristic features of coastal lagoons. Macrophyte presence, cover and composition create a

highly heterogeneous aquatic landscape (Arocena, 2007; Gamito et al., 2005), which is
also determined by the sediment nature (*i.e.* hard bottom, gravel, sand, mud) and
organic matter content (Menu et al., 2019).
Benthic macroinvertebrates are a key biotic compartment of coastal lagoons,
through their role in water filtration, food web dynamics, recycling of organic matter
and nutrients and sediment bioturbation (Barnes, 1980; Gamito et al., 2005; Levin et
al., 2001; Zaldívar et al., 2008). These benthic communities are reported to be shaped
mainly by natural environmental variability (*i.e.* environmental filtering) and passive
diffusion (mainly as planktonic larvae) (Basset et al., 2007). At the intra-lagoon scale,
the spatial distribution of macrobenthic invertebrates is tightly linked to the renewal of
marine-originating molecules, which is directly linked to the connection level with the
open sea, a result used to define the notion of confinement (Guelorget and Perthuisot,
1983). Superimposed onto the previously mentioned "natural environmental stress"
(*i.e.* strong fluctuations of abiotic parameters like salinity and temperature) are a
number of human related disturbances like anthropogenic eutrophication, which can
further impact lagoonal organisms (Elliott and Quintino, 2007; Rossi et al., 2006). Most
benthic macroinvertebrates are exposed to multiple human-induced disturbances as
they generally live for several years. Furthermore, these organisms tend to present, as
adults, a low dispersal potential (*i.e.* below a few kilometers), meaning that faced with a
disturbance, each individual will either resist, become accustomed or die, with overall
very little chance of escaping the disturbance, especially if it affects the entire lagoon
(Cognetti and Maltagliati, 2000). These characteristics make benthic
macroinvertebrates good theoretical indicators of the ecological health status of coastal
lagoons and since 2000, this biological compartment is officially included in the
assessment of European transitional water bodies carried out within the Water
Framework Directive (WFD) (European Commission, 2018). However, although they are
in theory good ecological indicators, benthic macroinvertebrates are also influenced by
"natural environmental stress", hence it appears crucial to disentangle the effect of
natural environmental variability from the effect of variability linked to human
disturbances on this compartment in hyper-variable ecosystems like coastal lagoons
(Pitacco et al., 2019). Most of the published work on Mediterranean coastal lagoon
macroinvertebrates have either been descriptive studies focusing on one or a few
lagoons (Amanieu et al., 1977; Arocena, 2007; Bachelet et al., 2000; Carlier et al., 2007;
Guelorget et al., 1994; Guelorget and Michel, 1979; Magni et al., 2022; Pérez-Ruzafa et
al., 2007a; Rossi et al., 2006), theoretical oriented studies testing ecological hypotheses
like environmental filtering (Basset et al., 2007) or species-area relations (Sabetta et
al., 2007) or very applied studies related to the WFD and focusing on lagoon typology
definition, indicator development or indicator reference values (Barbone et al., 2012;
Basset et al., 2013a; Reizopoulou et al., 2014). Disentangling the respective effect of
natural vs human-induced disturbances on lagoon communities requires a statistical
power which can be reached by using extensive datasets, spanning a variety of
ecological conditions (*i.e.* several lagoon systems) and a thorough characterization of
both sources of disturbance. In this study, we make use of the full potential of large-
scale environmental surveys of the various physical and biological compartments
(water column, sediments, macrophytes and benthic macroinvertebrates) of French
Mediterranean coastal lagoons: *(i)* to disentangle the effects of an anthropogenic
disturbance, here eutrophication, from those of natural environmental variability on
benthic macroinvertebrates and *(ii)* to understand the links between the different
environmental variables characterizing coastal lagoons, that affect benthic

macroinvertebrates. This study, based on a dataset including 29 French Mediterranean
coastal lagoons (ranging from oligo- to eu-haline, with diverse environmental and
anthropic settings), focuses on multiple levels of biological complexity; first the broad
and complex community level, then simple diversity indices like species richness and
Shannon diversity index and finally, a multimetric and WFD-compliant index.
162

Methods

Study sites

The French Mediterranean coast features a great diversity of lagoons characterized
by different hydro-climatic contexts (*e.g.* temperature, wind, rainfall). Some of these
lagoons are under the influence of large watersheds (up to 1 500 km²) while others are
connected to more restricted basins that may be 300 times smaller (*ca.* 5 km²). These
lagoons also present variable connections to the sea, with both natural vs man-made
channels, and continuous vs temporary channels. Human land-use in this region is
particularly high, associated to both urban and agricultural development (Derolez et al.,
2019). Most lagoons in continental France are located near large cities like Narbonne or
Montpellier and/or in areas intensively cultivated for wine, vegetables and cereals
(CORINE Land Cover, 2018) whereas the Corsican lagoons of Palo, Urbino and Diana are
relatively preserved from direct urban pressures. Differently, the Corsican lagoon of
Biguglia is located just south of the urban hub of Bastia and in a growing agricultural
plain (CORINE Land Cover, 2018; Pasqualini et al., 2017). Overall, French Mediterranean
lagoons face various degrees of anthropogenic pressures.
A total of 29 nanotidal (tide < 0.5 m) coastal lagoons located in the Gulf of Lion
(north-west Mediterranean Sea) and along the east coast of Corsica were selected for
this study and sampled across 41 stations (between one and three in each lagoon)
(Figure 1). The sampling was carried out once a time in 2006 or 2009 (depending on the
station) during the "Réseau de Suivi Lagunaire" and WFD campaigns. Infaunal benthic
macroinvertebrates were sampled in May, except four lagoons sampled at the end of
April (La Marette, Ponant, Berre and Vaine) and one lagoon at the end of June (Bolmon).
The studied lagoons encompass a wide range of hydro-morphological characteristics in
terms of surface, depth and connection to the sea (Table 1) with the lagoon-sea
connection level expressed as weak, moderate or strong depending on the differences
in salinity variations between the lagoon and a neighboring marine station (Derolez et
al., 2012). Lagoon surfaces vary from 1 km² (Campignol) to 133 km² (Berre) (Menu et
al., 2019) and station depth is comprised between 0.1 m (La Palme south station) and
9.5 m (Urbino) with most stations less than 3 m deep (Table 1).

Figure 1 - Map of the 41 stations sampled in the 29 lagoons located along the French Mediterranean coast (Gulf of Lion and Corsica). Each station's salinity type (according to *their median salinity and Venice classification (Anonymous, 1958) in* Le Fur et al. (2018)) and group membership (K-means partitioning) based on the benthic macrofauna Hellinger-transformed mean abundances are indicated using a symbol (triangle: oligo- and meso-haline and circle: poly- and eu-haline) and a color (red: group A and blue: group B), respectively (see section *Macrobenthic community structure* for the explanation on this analysis). See Table 1 for the full lagoon names and corresponding stations.

The different hydro-morphological characteristics of the studied lagoons lead to a
great diversity of salinity regimes with 8 oligo- or meso-haline lagoons and 21 poly-
or eu-haline lagoons (according to Le Fur et al., (2018), Figure 1) and to large
discrepancies between lagoons in terms of annual variability of temperature, salinity
and oxygen saturation. Overall, the physico-chemical status of the studied lagoons
between 2004 and 2009 was evaluated according to the WFD methodology, as poor
(15 lagoons), bad (3 lagoons), moderate (4 lagoons) or good (7 lagoons) (Table 1).
This evaluation takes into account the concentrations of total and dissolved inorganic
forms of nutrients, turbidity and oxygen saturation during the summer months of the
2004-2009 period and is conducted according to the WFD quality grid (Andral and
Sargian, 2010a, 2010b).

Table 1 - Hydro-morphological characteristics (surface, depth and connection
level to the sea) and water physico-chemical status of the 29 lagoons as
evaluated by the water framework directive (green = good, yellow = moderate,
orange = bad, red = poor). The physico-chemical status is based on
concentrations of total and dissolved inorganic forms of nutrients, turbidity and
oxygen saturation during the summer months of the 2004-2009 period (Andral
and Sargian, 2010a; Andral and Sargian, 2010b). Depth, sediment organic
matter content (OMC, % of dry sediment weight), beta diversity of macrophyte
morphotypes (β_{MM}, no unit – see section *Estimating the macrophyte morphotype*
beta diversity) of the 41 stations sampled for benthic macrofauna either in 2006
(#) or in 2009 (*) is also indicated. For stations sampled in the same lagoon,
each one is labeled using a 3 letter code with the last letter indicating the
geographical location of the station in the lagoon (North (N), Middle (M), South
(S), East (E), West (W), except THP which indicates the station's local name
Pisse-Saumes). The lagoons are listed from west to east along the Gulf of Lion
(see Figure 1 for their geographical location).
227

	Lagoon characteristics			Benthic station characteristics			
Lagoon	**Surface (km²)**	**Lagoon-sea connection**	**Physico-chemical status**	**Station code**	**Station depth (m)**	**OMC (%)**	**β_{MM}**
Canet*	6.43	weak	🟥	CAN	0.9	5.6	0.13
Leucate*	52.88	strong	🟩	LEN	1.9	8.2	0.71
				LES	3.3	7.2	0.67
La Palme#	6.94	moderate	🟩	LPN	0.6	4.68	0.6
				LPS	0.1	1.61	0.53
Bages-Sigean*	37.05	moderate	🟩	BGN	1.9	11.5	0.65
				BGM	2.2	5.9	0.8
				BGS	1.3	7.1	0.53

Site	Value	Strength		Code			
			🟩				
Campignol*	1	weak	🟥	CAM	0.8	3.8	1
Gruissan*	1.31	strong	🟩	GRU	0.9	3.4	0.61
Vendres*	6.6	weak	🟥	VEN	0.8	22.8	0.6
Grand Bagnas#	1.69	weak	🟧	GBA	0.5	5.38	0
Thau#	67.7	strong	🟩	THE	8.2	11.38	0.55
			🟩	THW	4	9.9	0.68
			🟩	THP	3.8	5.64	0.68
Ingril North#	2.28	strong	🟥	INN	1.1	12.48	0.4
Ingril South#	3.17	strong	🟥	INS	1.2	7.01	0.25
Vic#	12.96	moderate	🟥	VIC	1.5	13.05	0.69
Pierre-Blanche#	3.14	moderate	🟥	PBE	0.7	1.04	0.33
			🟥	PBW	0.7	9.93	0.67
Arnel#	4.42	moderate	🟥	ARN	0.8	7.88	0.29
Prévost#	2.46	strong	🟥	PRE	0.7	9.25	0.4
			🟥	PRW	0.7	8.4	0.6
Méjean#	7.12	moderate	🟥	MEE	1	7.67	0
			🟥	MEW	0.9	12.17	0
Grec#	1.08	moderate	🟥	GRE	0.7	8.49	0.33
Or (Mauguio)*	31.31	weak	🟥	ORE	1.4	10.9	0.75
			🟥	ORW	1.3	12.5	0.54
Ponant*	1.92	moderate	🟧	PON	3	10.9	0.11

La Marette*	1.3	weak		MRN	0.7	15.9	0
				MRS	0.6	14.2	0
Vaccares*	65.9	weak		VAC	1.9	5.8	0.23
La Palissade*	2.95	weak		PLS	0.6	1.5	0.6
Berre*	132.71	strong		BRN	5.9	11.8	0.39
				BRS	8.9	14.5	0.38
Vaine*	22.77	moderate		VAI	5.1	12.7	0.73
Bolmon*	6.01	weak		BOL	1.4	14.8	0.30
Biguglia*	13.75	weak		BIG	1	7.9	0.25
Diana*	5.4	strong		DIA	5.8	10.5	0.39
Urbino*	7.58	moderate		URB	9.5	16.3	0.63
Palo*	1.08	weak		PAL	1	12.5	0

Sampling

Benthic macroinvertebrates

At each of the 41 stations, three replicates (*ca.* 10 m apart) of unvegetated
sediment were sampled. Each replicate was made up of four pooled Ekman-Birge
grabs, representing a total surface of 0.09 m^2 per replicate and 0.27 m^2 per station.
Samples were sieved on a 1 mm mesh, fixed in a 5 to 7% formaldehyde/seawater
solution and stained with Rose Bengal for latter identification. The sediments were
then sorted in the laboratory and all organisms identified if possible to the species
level, following the nomenclature of the European Register of Marine Species
(ERMS). Taxa are indicated using WORMS reference
(https://www.marinespecies.org/aphia.php?p=match). Prior to the data analyses, we
removed meiofauna taxa (Ostracoda, Nematoda, Harpacticoida and Collembola) and
organisms present in only one lagoon and determined at the order (Decapoda,
Coleoptera, Nudibranchia), class (Insecta, Ascidiacea) or subclass (Oligochaeta)
level. We kept Actiniaria, Nemertea and Polycladida as they were identified across
multiple lagoons.

Sediment parameters

At each station, an additional grab sample was collected to evaluate sediment
organic matter and grain-size distribution. Sediment samples were sieved on a 2 mm
mesh to remove large macrophytes debris, and the filtrate was weighted and dried
until stable weight (110°C for 24 to 48h) in the laboratory. Organic matter content

(OMC, % of dry sediment weight) was calculated as the loss of weight after
combustion for 12h at 450°C of the dry sediment fraction below 2 mm. Because of
technical problems, no OMC was available for the stations ARN, LPN, LPS, PRW and
VIC in 2006. To estimate these missing values, we used OMC values measured in
spring of 2006 (ARN, PRW, VIC) or spring of 2007 (LPN and LPS) for the WFD
sediment quality evaluation. For ARN, LPS, LPN and VIC, we estimated OMC by
averaging over the values measured in the three or four stations closest to the
macrofauna station (distance between 285 m and 707 m). For PRW, we used the
OMC value measured at the closest station as this station was located only 114 m
from the macrofauna station (Supplementary material 1 and 2). Since laboratory
protocols changed between years (2006 and 2009) and between lagoons regarding
the sediment particle sizes considered for the grain-size analysis, we chose not to
consider grain-size parameters.

Water physico-chemical properties
Water column physico-chemical properties of the 29 lagoons are measured
monthly during summer (from June to August or September) at least since 2003,
following a standardized protocol implemented by the WFD (Derolez et al., 2019).
Salinity (Sal), temperature (Temp) and oxygen saturation (O_2 sat) were measured
using a portable field probe (WTW LF 197, WTW-GmbH, Weilheim, Germany) and
turbidity (Turb) was measured in the laboratory using an optical turbidimeter (2100N
IS turbidimeter ISO 7027). Chlorophyll *a* concentration ([Chla]) was measured in 50
mL samples of water, first filtered on a Whatman GF/F filter (0.7 µm). Pigments were
later extracted in the laboratory in a 90% acetone + distilled water solution during
24 hours (4°C, dark conditions after 10 minutes of centrifugation). The supernatant
was then dosed using a spectrofluorometer following Neveux and Lantoine (1993).
The full description of the protocol followed to measure [Chla], total nitrogen
concentration [TN], ammonium concentration [NH_4^+], sum of nitrate plus nitrite
concentrations [NO_x], total phosphorus concentration [TP] and dissolved inorganic
phosphorus concentration [DIP] is described in Bec et al. (2011) and Souchu et al.
(2010). Only sub-surface data were available for all stations except the THW, THE,
BRN, BRS stations located in the deep Thau and Berre lagoons for which sub-surface
and bottom data were available. In these four cases, we chose to consider the
bottom data.

Estimating the macrophyte morphotype beta diversity
Lagoons are characterized by a mosaic of benthic habitats, determined in
particular by macrophyte diversity which can affect benthic macroinvertebrates
(Arocena, 2007; Cardoso et al., 2004; Magni and Gravina, 2023; Ouisse et al., 2011)
by acting as food (Jones et al., 2021; Quillien et al., 2016; Vizzini and Mazzola, 2006),
refuge (Mancinelli and Rossi, 2001; Nordström and Booth, 2007; Ware et al., 2019)
and/or means of transport (Norkko et al., 2000). Although sampling was carried out
on locally unvegetated sediments to avoid biases linked to macrophytes and
associated faunal assemblages, we postulate that the presence and composition of
macrophyte assemblages within each lagoon can affect the composition of infaunal
benthic communities. We developed an index (later beta diversity of macrophyte
morphotypes, β_{MM}) calculated as the average Bray-Curtis dissimilarity at the lagoon
or sub-lagoon scale (when several macrofauna stations were sampled in one lagoon)
based on a station by macrophyte morphotype matrix (presence/absence). Each
macrophyte station is coded as bare sediment if the macrophyte cover is less than

25% or as one or several of seven macrophyte morphotypes (*i.e.* mushroom, feather,
hair, ball, branched, sheet and meadow) based on the thallus morphology of the
dominant or co-dominant macrophyte taxa. This index varies between 0 if all the
considered stations are dominated or co-dominated by the same macrophyte
morphotype(s) or if they all have bare sediment and 1 if all the considered stations
are dominated or co-dominated by different macrophyte morphotypes (Appendices 1
and 2). We gathered the data used to calculate this index from the WFD macrophyte
dataset (all information regarding sampling protocols are available in Le Fur et al.,
2017).

Data analyses

We extracted all the data used in this study from the French database Quadrige
(Ifremer, http:// quadrige.eaufrance.fr/). All the raw data used in this study
(macrofauna, environmental data, GIS data) are available online along with all the
scripts and code used to analyze the data (Jones et al., 2024). We performed all
analyses using R version 3.6.3 (R Core Team, 2020) and the packages vegan (*v2.5*-6;
Oksanen et al., 2019), missMDA (Josse and Husson, 2016), labdsv (*v2.0*-1; Roberts,
2019), olsrr (v0.5.3; Hebbali, 2020), MASS (Venables and Ripley, 2002), corrplot
(*v0.84*; Wei and Simko, 2017) and rstatix (v0.6.0; Kassambara, 2020). We considered
a significance level of 0.05 for all statistical tests.

Environmental variables

Water column parameters were measured in summer whereas macrofauna was
sampled in spring which prevented us from relating these two datasets. Nonetheless,
the presence and abundance of macroinvertebrates in a lagoon year y is influenced
not only by the environmental conditions prevailing that year but also those
prevailing the previous years especially for pluri-annual species. As such, we
associated the macrofauna data collected year y to the mean and coefficient of
variation (CV) of the water column parameters measured years y-2, y-1 and y, as
done by Le Fur et al. (2019) (Table 2).

We classified the parameters presented in Tables 1 and 2 into two categories. The
first category included 11 physical and biotic factors used as a proxy for the natural
variability of environmental conditions (lagoon surface, lagoon-sea connection,
station depth, β_{MM}, OMC, mean and CV of Sal, Temp, O_2 sat). The second category
included 14 biogeochemical variables that either cause eutrophication (*i.e.* mean
and CV of [TN], [NO$_x$], [NH$_4^+$], [TP], [DIP]) or are the direct consequence of
eutrophication (*i.e.* mean and CV of [TN], [TP], [Chla], turbidity) (Souchu et al.,
2010). Because of technical problems, no [NO$_x$] and [DIP] data were available for the
stations BOL and PLS. Consequently, we imputed the missing mean and CV values
for these two parameters at both stations with the 'imputePCA' function. This
function estimates the missing parameters by first performing a principal component
analysis on all the environmental variables and then by using the regularized fitted
matrix. These steps are performed iteratively until convergence is reached (Josse
and Husson, 2012).

Table 2 - Environmental parameters and their respective code used to
characterize the water. For each water parameter, the mean and coefficient of
variation (CV) were calculated between the years y-2 and y (with y the
macrofauna sampling year) and the overall median, minimal and maximal
values across the 41 stations are indicated. The station with the minimal and

maximal value of each parameter is also indicated between brackets, as well as
the transformation applied to each parameter. See Table 1 for the station codes.

Parameter (unit)	Code	Measure	Median	Min	Max	Transf.
Salinity (PSU)	Sal	mean	32.39	2.83 (PLS)	42.30 (PBW)	/
		CV	13.52	2.49 (DIA)	82.24 (CAM)	log_e
Oxygen saturation (%)	O_2 sat	mean	100.88	56.61 (BRN, BRS)	190.40 (BOL)	log_e
		CV	27.38	0.82 (PLS)	67.33 (PRE)	$\sqrt{}$
Temperature (°C)	Temp	mean	24.08	19.63 (BRN, BRS)	27.09 (URB, BIG)	/
		CV	9.05	1.82 (CAN)	22.81 (PON)	$\sqrt{}$
Chlorophyll a concentration (µg.L^{-1})	[Chla]	mean	9.43	0.70 (BGS)	144.20 (MEW)	log_e
		CV	71.04	25.42 (BOL)	178.73 (BIG)	log_e
Total nitrogen concentration (µmol.L^{-1})	[TN]	mean	52.05	11.08 (DIA)	266.03 (MEW)	log_e
		CV	27.77	6.94 (VAC)	89.47 (BIG)	log_e
Ammonium concentration (µmol.L^{-1})	[NH$_4^+$]	mean	1.47	0.06 (PLS)	31.62 (GRE)	log_e
		CV	106.88	28.65 (CAN)	257.93 (THE)	$\sqrt{}$
Nitrate plus nitrite concentration (µmol.L^{-1})	[NO$_x$]	mean	0.37	0.06 (GBA)	22.81 (CAM)	log_e
		CV	94.87	43.30 (CAN)	227.37 (URB)	log_e
Total phosphorous concentration (µmol.L^{-1})	[TP]	mean	2.51	0.64 (LEN)	31.19 (CAN)	log_e
		CV	36.75	8.91 (VAC)	106.12 (ARN)	log_e
Phosphate concentration (µmol.L^{-1})	[DIP]	mean	0.24	0.02 (LES)	14.53 (CAN)	log_e
		CV	93.10	40.97 (THW)	187.55 (INN)	log_e
Turbidity (NTU)	Turb	mean	5.20	1.14 (BGS)	99.65 (PLS)	log_e
		CV	68.30	1.92 (PLS)	207.54 (ARN)	$\sqrt{}$

347   Transformation (Transf.): / = no transformation, loge = natural logarithm, √ = square-root

Macrobenthic community structure

First, we calculated the mean macroinvertebrate abundance at the station scale by averaging over the three replicates and transformed it using a Hellinger transformation, to give more weight to common taxa and less weight to highly abundant, rare and under-sampled taxa (Legendre and Legendre, 2012). We then classified the 41 stations based on these transformed mean abundances, using K-means partitioning and the Calinski-Harabasz criterion with 100 iterations ('cascadeKM' function). Our goal was to highlight similarities between lagoons based on common species without focusing on rare taxa, as each coastal lagoon can present unique species assemblages (Basset et al., 2007), and to help with the result interpretation thanks to a simple classification. We tested for significant differences in the composition of our macrofauna groups using the analysis of similarity (ANOSIM) test with 9999 permutations and Euclidean distances ('anosim' function). Finally, we characterized each group using the indicator species metric ('indval' function, 9999 permutations). A high value of this metric for a given taxon associated to a given group means this taxon is consistently more present (fidelity) and abundant (specificity) in the stations composing that group than in stations composing the other groups (Dufrêne and Legendre, 1997). For this last analysis, we adjusted p-values using the False Discovery Rate correction method (FDR; Benjamini and Hochberg, 1995).

Taxonomy-based indices

To characterize the macrobenthic community structure, we calculated the commonly used M-AMBI (Multivariate AZTI Marine Biotic Index), the three indices composing the M-AMBI - species richness (SR), Shannon diversity index (H') and AMBI (Muxika et al., 2007) - plus the inverse of Simpson's index (N2) and the total macrofauna density. The AMBI is an index based on the classification of benthic species into ecological groups, from the most sensitive to disturbances (i.e. organic matter enrichment) to the most opportunistic ones (Borja et al., 2000). The raw (i.e. untransformed) macrofauna abundances were used to calculate all these indices. For AMBI and M-AMBI calculations, we removed high order taxa without an ecological group value (Anemonia and Polycladida), planktonic (Mysidae) and hard-substrate taxa (Serpula and Serpulidae) (Borja et al., 2005). We computed M-AMBI using the R function available in Sigovini et al. (2013) and we defined the good reference conditions as the highest values of SR and H' (46 and 3.73, respectively) and lowest value of AMBI (0.31) measured in the dataset, irrespective of the WFD physico-chemical status described above. The results are perfectly similar whether the reference values used are those described above, those from the French government or those from the reference lagoons (see supplementary material 3). All indices were computed at the replicate level (calculation of M-AMBI at this scale does not affect the results, see supplementary material 4); the mean and standard deviation (SD) values were then calculated for each station. Finally, we tested for differences in total macrofauna density, SR, H', N2, AMBI and M-AMBI between previously defined groups of stations (see section *Macrobenthic community structure*) using non-parametric Kruskal-Wallis rank sum tests (Breslow, 1970).

Linking macrobenthic communities and environmental variables

To assess how natural environmental variations correlated with anthropogenic pressures (explanatory variables) and how each explanatory variable correlated with

macrobenthic indices, we built a correlogram showing the significant Spearman
correlations ('cor_pmat' function) between all the raw variables. We also tested for
differences in the variables considered (explanatory variables and macrobenthic
indices) between lagoon-sea connection levels using non-parametric Kruskal-Wallis
rank sum tests (Breslow, 1970), followed by pairwise Wilcoxon tests for multiple
comparison, while adjusting p-values with FDR (Benjamini and Hochberg, 1995).
Then, we transformed (natural logarithm or square root transformation), when
necessary, the environmental variables and macrobenthic indices to approximate
normal distributions and reduce data spread, before standardizing all the
explanatory variables to facilitate the interpretation of the multiple linear regression
coefficients (Schielzeth, 2010). We $\log_e$-transformed the lagoon surface, station
depth, SR, N2 and macrofauna density (see Table 2 for the rest of the
transformations). To limit multicollinearity and avoid artificially inflating the fit of
linear models, we reduced the number of environmental variables (see section
Environmental variables) for all future analyses by considering Pearson correlations
< |0.6|, corresponding to a p-value < 5. 10^{-5} and leading to 19 remaining
environmental variables.
To determine which environmental variables most strongly influence lagoon
macrobenthic communities, we performed a redundancy analysis (RDA, 'rda'
function) between the 19 previously selected environmental variables and the
Hellinger-transformed mean taxa abundances. We computed the variance inflation
factor (VIF, 'vif.cca' function) to check for multicollinearity and sequentially removed
variables with a VIF > 5 (James et al., 2013). We verified the significance of the full
RDA model using the 'anova.cca' function (1000 steps) and we performed a forward
model selection procedure ('ordiR2step' function) based on the adjusted R2 of the
full model and a p-value of 0.05 (9999 permutations) (Blanchet et al., 2008). We
then computed the marginal effect of each selected explanatory variable as the
adjusted R^2 of the constrained ordination containing only the given variable as the
predictor. The conditional effect of a variable was equal, during the forward selection
procedure, to the additional amount of variance in species communities explained by
the corresponding variable at the time it was included into the model. Finally, we
performed a partial RDA using the variables previously selected, to isolate the
impact of eutrophication from natural environmental variability on lagoon
macrobenthic communities (Legendre and Legendre, 2012).
In the same way as for the constrained multivariate analysis, we implemented a
multiple linear regression approach (without any interactions between variables) to
investigate which variables related to eutrophication and natural variability most
strongly influenced the macrobenthic indices (SR, H', N2, total macrofauna density,
AMBI and M-AMBI). As there were still 19 potential explanatory variables, we
performed a preliminary selection by including all the quantitative variables which
alone had a significant effect on the index as first or second-degree orthogonal
polynomial and the lagoon-sea connection if at least one of the three categories
(weak, moderate, strong) had a significant effect on the index ('lm' function). We
included second-degree polynomials to account for potential non-linear relations
(Pérez-Ruzafa et al., 2007b) and verified that the selected variables had VIFs inferior
to 5. We also checked the model hypotheses using the residuals vs fitted values plot
for linearity, the Shapiro-Wilk test for residual normality, the Breush Pagan test for
residual homoscedasticity ('ols_test_breusch_pagan' function) and the 'acf' function
for the detection of residual spatial autocorrelation based on the station coordinates.

Then, we ran a stepwise model selection using the 'stepAIC' function set in both
directions and the BIC criterion. Finally, we checked the hypotheses of the selected
model as previously explained, we investigated potential high leverage points
(Cook's distance > 0.5) and we reran the selected model without these points to
verify its robustness. To compare the relative effect of each selected explanatory
variable on a given macrobenthic index, we expressed the slope estimates of the
standardized variables as effect sizes when the variable was quantitative and as
mean standardized effects when the variable was categorical (Schielzeth, 2010).
Lastly, we implemented a variance partitioning procedure using the 'varpart'
function, to assess how much community variation or macrobenthic index variation
could be: (i) uniquely attributed to eutrophication when the effect of natural
variability had been removed, (ii) uniquely attributed to natural variability when the
effect of eutrophication had been removed or (iii) jointly attributed to both effects
(Legendre and Legendre, 2012). Prior to this analysis, we reran the variable selection
procedures implemented at the community and macrobenthic index levels
considering alternatively only the eutrophication variables or only the variables
related to natural variability. We then included the variables selected by the
aforementioned method in the variance partitioning, which was based on partial
RDAs or on partial linear regressions and we presented the results using the adjusted
R² values.

464   **Results**

Inter-lagoon environmental variability
The dataset considered in this study encompasses a wide diversity of lagoons in
terms of physico-chemical state according to the WFD and environmental
parameters associated to natural variability and eutrophication (Tables 1 and 2).
First, only lagoons with a strong (Leucate, Gruissan, Thau, Berre and Diana) or a
moderate (La Palme and Bages-Sigean) connection to the sea were in a good
physico-chemical state (green in Table 1). All the stations sampled in these lagoons
presented mean [Chla] below 7 µg.L^{-1} and all of them except the ones located in the
deep Berre and Diana lagoons (station depth > 5.8 m) presented a high beta
diversity of macrophyte morphotypes (β_{MM} > 0.53). Five stations sampled in these
lagoons presented the lowest value for a eutrophication related variable (DIA: mean
[TN], BGS: mean [Chla] and mean Turb, LEN : mean [TP], LES: mean [DIP], THW: CV
[DIP]). However, the THE station (Thau) had the most variable [NH$_4^+$] and the BRN
and BRS stations (Berre) had the lowest mean O$_2$ sat of the dataset. Seven lagoons
were in a moderate (yellow in Table 1) or in a bad physico-chemical state (orange in
Table 1) and were either moderately (Ponant, Vaine, Urbino) or weakly (Grand
Bagnas, Vaccarès, Biguglia and Palo) connected to the sea. In this group, the GBA
and VAC stations respectively had the lowest mean [NO$_x$] and the lowest CV [TN] of
the dataset whereas the URB and BIG stations respectively had the highest CV [NO$_x$]
and CV [Chla]. All the remaining weakly connected lagoons were in a poor physico-
chemical state (Canet, Campignol, Vendres, La Marette, La Palissade, Bolmon). In
this group, the CAN station had both the highest mean [TP] and [DIP] and the lowest
CV [NH$_4^+$] and [NO$_x$] of the dataset, while the CAM and BOL stations presented
respectively the highest mean [NO$_x$] and CV [Chla] of the dataset. Finally, all the
lagoons between the cities of Sète and Montpellier (Figure 1) were in a poor
condition, despite being strongly (Ingril North, Ingril South, Prévost) or moderately

(Vic, Pierre-Blanche, Arnel, Méjean and Grec) connected to the sea. All the stations
located in these lagoons except PBE, had OMC above 7% and up to 13%. In this
group, the MEW station presented the highest mean [Chla] and [TN] of the dataset
while the GRE station had the highest mean [NH$_4^+$], the ARN station the highest CV
[TP] and Turb and the INN station the highest CV [DIP] of the dataset.

Based on the significant Spearman correlations between the environmental
variables (Figure 2), stations in the larger lagoons were significantly deeper, cooler in
summer and less variable in terms of summer salinity, presented higher β_{MM} and
lower values for 8 eutrophication related variables (*e.g.* mean [Chla], mean [NO$_x$]
and CV [NO$_x$]). The deepest stations were also the ones with the highest OMC. Two
variables related to natural environmental variability were also significantly different
according to the lagoon-sea connection level (KW test, $p < 0.05$ and post-hoc
pairwise Wilcoxon tests $p < 0.05$): mean Sal (moderate > weak and strong > weak)
and CV Sal (weak > moderate > strong).

Figure 2 - Correlogram showing only significant positive (red toned circles) and
negative (grey toned circles) Spearman correlations ($p < 0.05$) between the

untransformed taxonomic indices (in black, SR: species richness, H': Shannon
diversity, N2: inverse of Simpson's dominance, M-AMBI: Multivariate AMBI), the
untransformed environmental variables related to the lagoon's natural
variability (in blue, β_{MM}: beta diversity of the macrophyte morphotypes, OMC:
sediment organic matter content) and the untransformed environmental
variables related to eutrophication expressed as means and coefficient of
variations (in green, [Chla]: chlorophyll *a* concentration, [TN]: total nitrogen
concentration, [NH_4^+]: ammonium concentration, [NO_x]: nitrate plus nitrite
concentration, [TP]: total phosphorous concentration, [DIP]: dissolved inorganic
phosphorous concentration) calculated across 41 stations located in 29 lagoons
along the French Mediterranean coast.

Macrofauna communities and macrobenthic indices across French
Mediterranean lagoons

A total of 42 465 benthic macroinvertebrates were counted and identified across
all samples, belonging to 227 different taxa with most organisms identified to
species level (74.5%), others to genus (17.2%), family (2.6%), order (2.6%), subclass
(0.4%), class (1.8%) or phylum (0.9%). Across the 227 taxa, 1.3% were identified in
over 50% of the stations, 90% in less than eight stations and 40% in only one
station. No taxon was present in all 41 stations. Polychaetes were by far the most
diversified group (102 taxa), followed by Malacostraca crustaceans (54 taxa),
bivalves (25 taxa) and gastropods (19 taxa). Among the three taxa with a frequency
of occurrence above 50%, two were polychaete species (*Hediste diversicolor* (66%)
and *Heteromastus filiformis* (51%)), and the third was the bivalve *Cerastoderma*
glaucum (51%). These three species also presented an important inter-station
variability in abundance. For all the following analyses, we considered 217 taxa (see
section *Benthic macroinvertebrates*), except for the AMBI and M-AMBI calculation, for
which we considered only 213 taxa (see section *Taxonomy-based indices*).

The K-means clustering delineated two groups of stations with distinct
macrobenthic communities (minimal difference in Calinsky-Harabasz criterion = 1.4),
a result confirmed by the analysis of similarity test (ANOSIM statistic = 0.51, $p <$
0.001). These groups did not correspond to any obvious spatial or salinity pattern, as
exemplified by the four Corsican stations and the seven oligo-mesohaline stations
which were classified in both groups (Figure 1). The polychaete *Hediste diversicolor*
($p = 0.022$) was an indicator taxa of group A stations (red in Figure 1) along with the
bivalve *Abra segmentum* ($p = 0.022$), Chironomidae ($p = 0.029$) and the amphipod
Gammarus aequicauda ($p = 0.038$). Conversely, group B stations (blue in Figure 1)
were significantly characterized by the bivalve *Scrobicularia cottardii* ($p = 0.048$),
closely followed by another bivalve, *Loripes orbiculatus* ($p = 0.065$).

Finally, we found significant differences among the two groups in terms of SR and
M-AMBI; group A was composed of stations with lower SR and weaker M-AMBI values
than group B (SR: $\chi^2(1) = 6.41$, $p = 0.011$; M-AMBI: $\chi^2(1) = 5.76$, $p = 0.016$). Group B
was also characterized by stations with higher H' and N2 values and higher maximal
macroinvertebrate densities than stations composing group A, albeit these
tendencies were not significant (Appendix 3).

Relationship between macrofauna communities and environmental
variables

In all the following analyses, six variables were not considered because they were
highly correlated with another variable (|r| ≥ 0.6 ; $p < 5. 10^{-5}$): station depth (to
lagoon surface), CV Sal (to mean Sal), mean [TN], mean [TP], mean Turb (to mean

[Chla]) and CV [TP] (to CV [TN]). Redundancy analysis (RDA) was used to relate the
macrofauna communities recorded within the lagoons to their natural environmental
variability and their eutrophication levels (Figure 3). Mean Sal was also not
considered because of a VIF > 5 linked to significant differences between lagoon-sea
connection levels (see section *Inter-lagoon environmental variability*). The final RDA
model, determined using a forward model selection procedure, explained 22% of the
inter-station community variability (model adj. R^2) and included six variables related
to natural environmental variability (lagoon-sea connection level ($p < 0.001$), mean
Temp ($p < 0.001$), lagoon surface ($p = 0.0076$), CV Temp ($p = 0.013$), mean O_2 sat
($p = 0.026$) and OMC ($p = 0.041$) and two eutrophication related variables (mean
[Chla] ($p < 0.001$) and mean [DIP] ($p = 0.024$)). Regarding the individual effect of
environmental variables, mean [Chla] (7.22%), lagoon surface (5.90%), lagoon-sea
connection (5.30%) and mean Temp (5.04%) were the most influential variables
according to their marginal effect. Based on conditional effects, the most influential
variables were mean [Chla] (7.22%), lagoon-sea connection (4.64%), mean Temp
(3.62%) and lagoon surface (1.69%) (Appendix 4).
The first two axes accounted for 33.8 and 17.8% of the RDA model variance,
respectively. First, the stations from group A and B were clearly separated along the
first axis of the RDA, except for LPN, LPS, PLS and INS (Figure 3a). On the ordination
of the environmental variables (Figure 3b), lagoon surface and CV Temp grouped on
the positive side of the first axis while mean [Chla], mean [DIP] and mean Temp
grouped on the negative side of the first axis. Consequently, group B macrobenthic
communities were linked to large lagoons (also relatively deep), several of which
presented high summer temperature variations. They were characterized by higher
relative abundances of the bivalves *S. cottardii* and *L. orbiculatus* (Figure 3c).
Conversely, group A communities were associated to smaller lagoons, subject to
higher mean summer temperatures and high eutrophication levels. They were
characterized by higher relative abundances of the polychaete *H. diversicolor* and
Chironomidae (Figure 3c).
The second axis mainly corresponded to a confinement and also to an organic
matter content gradient, mostly separating each group into two sub-groups with the
lagoons weakly connected to the sea presenting sediments enriched in organic
matter located towards the positive side of the second axis and the moderately and
strongly connected lagoons located towards the negative side of the second axis. As
mean Sal and CV Sal were correlated to lagoon-sea connection level (see section
Inter-lagoon environmental variability), this axis also corresponded to a salinity
gradient with low and variable salinity stations characterized by a higher

Figure 3 - Redundancy analysis plots (RDA, axis 1 and 2) focusing on (a) the benthic macrofauna sampled across 41 stations located in 29 lagoons along the French Mediterranean coast (see Table 1 for the full lagoon names and corresponding stations) as explained by the set of environmental variables related to eutrophication and natural variability composing the most

parsimonious model, (b) the selected environmental variables, where mean [Chla], lagoon surface, mean O_2 sat and mean [DIP] were $\log_e$ transformed and CV Temp was square-root transformed (see Tables 1 and 2 for the full environmental variable names and values) and (c) the macrofauna taxa, where only taxa with a goodness of fit (cumulative proportion of variance explained by the first two axis) above 0.25 are plotted. In (a), inter-station distances are euclidean while in (b) and (c) angles between all vectors reflect linear correlations. Across all plots, each station (point) is represented using a symbol for its salinity type (triangle: oligo- and meso-haline and circle: poly- and eu-haline) and a color (red: group A and blue: group B) for its group membership (K-means partitioning) based on Hellinger-transformed mean abundances.

abundance of *Chironomus salinarius* and *Streblospio shrubsolii* (Figure 3c). All the oligo- and meso-haline stations (triangles on Figure 3) grouped on the upper part of the RDA space and most poly- and eu-haline stations (circles on Figure 3) grouped on the lower part of the figure. Furthermore, communities of lagoons weakly connected to the sea were characterized by higher abundances of the bivalve *Scrobicularia cottardii* for group B, while communities from lagoons moderately to strongly connected to the sea were characterized either by higher abundances of *L. orbiculatus* for group B or by a diversified suite of species like the decapod *Carcinus aestuarii* or the polychaete *Capitella capitata* for group A.

Once the effect of natural environmental variability on macrofauna communities was removed through partial RDA and the only explanatory variables left to vary were mean [Chla] and mean [DIP] (eutrophication variables), no differences could be made between the communities of groups A and B (Appendix 5). This partial RDA explained only 6.4% of the inter-station community variability (model adj. R^2).

Relationship between macrobenthic indices and environmental conditions

Based on the significant Spearman correlations between the raw environmental variables and raw macrobenthic indices (Figure 2), SR, H', N2 and M-AMBI were significantly higher at stations where mean Temp was lower, salinity was more stable and eutrophication related variables were lower, especially mean [Chla]. Furthermore, H', N2 and M-AMBI were significantly higher at stations where mean Sal and β_{MM} were higher while SR and M-AMBI were also significantly higher at deeper stations located in larger lagoons. AMBI appeared to be significantly correlated positively only with mean [NO_x] while total macrofauna density was significantly higher at stations with lower but more variable summer water temperatures, lower salinity and lower mean [DIP].

The results of the multiple linear regressions identifying which variables influenced the macrobenthic indices are presented in Table 3. The most parsimonious models contained between two and four predictors and explained between 27% and 74% of the macrobenthic indices' variability. All the models contained at least one predictor associated to eutrophication and at least one predictor associated to natural variability except the N2 and AMBI models, which only contained predictors associated to natural variability and eutrophication respectively. Lagoon-sea connection was selected in three models, H', N2 and M-AMBI, and in all three, the response variable was significantly higher in strongly connected lagoons. Differently, H' and N2 were significantly higher and lower in moderately connected lagoons, respectively, while M-AMBI was significantly lower in weakly connected lagoons. Mean Temp and mean Sal were selected in two models, SR and macrofauna density. Mean Temp negatively influenced SR and macrofauna

density while mean Sal positively influenced SR and had a quadratic effect (concave form) on macrofauna density. β_{MM} was selected in two models, H' and N2, and positively influenced both response variables. The AMBI model was composed of mean [TN] (positive effect) and CV [NH_4^+] (quadratic effect, concave form). Finally, three other eutrophication related variables, CV [TN], mean [NH_4^+] and CV [NO_x], were selected respectively in H', macrofauna density and M-AMBI models where they had either a negative effect (M-AMBI) or a quadratic effect (concave form, H' and density).

To isolate the independent effect of eutrophication on macrobenthic indices, we also plotted the partial effect of mean [Chla] on SR, H' and M-AMBI once the effect of the other predictors retained in the minimal adequate models (Table 3) had been removed (Appendix 6). After controlling for the effect of mean Sal and mean Temp, the simple linear regression model between mean [Chla] and SR was still significant (adj. R^2 = 0.32, F(1, 39) = 19.55, p < 0.001). It was also the case for the model between mean [Chla] and H' after controlling for the effect of lagoon-sea connection, CV [TN], CV [TN]^2 and β_{MM} (adj. R_2 = 0.14, F(1, 39) = 7.33, p = 0.01) and for the model between mean [Chla] and M-AMBI after controlling for lagoon-sea connection and CV [NO_x] (adj. R^2 = 0.26, F(1, 39) = 15.41, p < 0.001).

Table 3 - Most parsimonious models (ordinary least-square regressions) explaining the standardized macrobenthic taxonomic indices (species richness (SR), Shannon diversity (H'), inverse of Simpson's dominance (N2), AZTI's Marine Biotic Index (AMBI), Multivariate AMBI (M-AMBI)). SR, H', N2 and total macrofauna density were log_e transformed prior to standardization. Both eutrophication related and natural variability related variables are considered as explanatory variables and the slope estimates are expressed either as effect sizes when the variable is quantitative or as mean standardized effects when the variable is categorical.

Model terms		Estimate	SE	p-value
SR: F(3,37) = 18.18 ; RSE = 0.66 ; adj. R^2 = 0.56				
[Chla]	mean	-0.50	0.12	< 0.001
Temperature	mean	-0.34	0.11	0.005
Salinity	mean	0.27	0.11	0.018
H': F(6,34) = 13.57 ; RSE = 0.59 ; adj. R^2 = 0.65				
Lagoon-sea connection	strong	0.61	0.19	< 0.001
	moderate	0.39	0.15	0.016
	weak	-0.16	0.17	0.32
[Chla]	mean	-0.31	0.12	0.016

[TN]	CV	-0.02	0.11	0.89
[TN]^2		-0.30	0.10	0.004
β_{MM}		0.23	0.10	0.035

N2: $F_{(3,37)}$ = 11.18 ; RSE = 0.75 ; adj. R^2 = 0.43				
Lagoon-sea connection	strong	0.79	0.21	< 0.001
	moderate	-0.58	0.19	0.005
	weak	-0.27	0.22	0.11
β_{MM}		0.31	0.12	0.016

Total macrofauna density: $F_{(5,35)}$ = 8.48 ; RSE = 0.72 ; adj. R^2 = 0.48				
Temperature	mean	-0.36	0.12	0.006
Salinity	mean	-0.23	0.12	0.071
Salinity^2		-0.30	0.13	0.023
[NH$_4^+$]	mean	0.21	0.12	0.10
[NH$_4^+$]^2		-0.30	0.13	0.030

AMBI: $F_{(3,37)}$ = 5.84 ; RSE = 0.86 ; adj. R^2 = 0.27				
[TN]	mean	0.34	0.14	0.017
[NH$_4^+$]	CV	0.11	0.14	0.41
[NH$_4^+$]^2		-0.43	0.13	0.003

M-AMBI: $F_{(4,36)}$ = 29.04 ; RSE = 0.51 ; adj. R^2 = 0.74				
Lagoon-sea connection	strong	0.59	0.16	0.005
	moderate	-0.06	0.14	0.67
	weak	-0.52	0.15	0.028
[NO$_x$]	CV	-0.37	0.09	< 0.001

[Chla]	mean	-0.35	0.09	< 0.001

RSE: Residual standard error, CV: coefficient of variation, [Chla]: chlorophyll *a* concentration in the water, [TN]:
total nitrogen concentration in the water, β_{MM}: beta diversity of the macrophyte morphotypes, [NH$_4^+$]:
ammonium concentration in the water and [NO$_x$]: nitrate plus nitrite concentration in the water

To synthesize our results, we plotted for each station the M-AMBI along with the
three macrobenthic indices composing it (SR, H' and AMBI) (mean ± SD) as a
function of mean [Chla] (Figure 4). Mean [Chla] was chosen as a synthetic indicator
of eutrophication because it is (*i*) significantly and positively correlated to 9 out of
the 13 eutrophication related variables (Figure 2), (*ii*) the eutrophication related
variable (with mean [TN]) that presents the strongest correlations with SR, H' and M-
AMBI (Figure 2), (*iii*) the most frequently selected eutrophication related variable in
the macrobenthic index models (Table 3) and (*iv*) a relevant indicator of
eutrophication status according to Bec et al. (2011) and Cloern (2001).

Figure 4 - Mean (± SD) of the species richness (SR), Shannon diversity (H'), AMBI and M-AMBI indices as a function of mean chlorophyll a concentration ($\log_e$ scale) across 41 stations located in 29 lagoons along the French Mediterranean coast (see Table 1 for the full lagoon names and corresponding stations). Across all plots, each station (point) is represented using a symbol for its salinity type (triangle: oligo- and meso-haline and circle: poly- and eu-haline) and a color (red: group A and blue: group B) for its group membership (K-means partitioning) based on Hellinger-transformed mean abundances. The dashed lines correspond to loess regressions.

SR, H' and M-AMBI were significantly correlated to mean [Chla] (r (Spearman) > | 0.50|) while AMBI was not (r = 0.12). Based only on the mean [Chla], most stations from group B (in blue) grouped towards higher levels of mean SR, H' and M-AMBI than stations from group A (in red). Nonetheless, we identified an intermediary zone corresponding to mean [Chla] between 6.3 µg.L^{-1} and 14.2 µg.L^{-1} and composed of stations from both group located between the INN (SR = 12 ± 3, H' = 3.06 ± 0.42 and M-AMBI = 0.57 ± 0.1) and ORE (SR = 9 ± 1, H' = 1.84 ± 0.07 and M-AMBI = 0.31 ± 0.01) stations. In the line with these observations, the LPN, LPS and BOL stations appeared as outliers, as did LPN and LPS in the RDA (Figure 3). Finally, mean [Chla] was also significantly correlated (r (Spearman) = -0.40, *p* = 0.009) to the intra-station SR variability expressed using the standard deviation.

Disentangling natural from anthropogenic factors

Based on the eutrophication and natural variability variables independently selected, we performed a variance partitioning, to investigate the amount of community and macrobenthic index variability uniquely linked to each variable category (Figure 5). Macrobenthic community and indices can be grouped into three categories according to the relative importance of natural variability alone, eutrophication alone, and the two together, in explaining their respective variability. (*i*) Macrofauna density was first influenced uniquely by natural variability (29.0%), then jointly by natural variability and eutrophication (11.5%) and finally uniquely by eutrophication (7.9%). (*ii*) Macrobenthic community, N2 and H' were first jointly influenced by natural variability and eutrophication (community: 8.2%, H': 33.6%, N2: 22.1%), and then uniquely by natural variability (community: 8.0%, H': 17.8%, N2: 21.1%) and finally uniquely by eutrophication (community: 3.7%, H': 13.9%, N2: 2.9%). (*iii*) SR and M-AMBI were first jointly influenced by natural variability and eutrophication (SR: 30.7%, M-AMBI: 49.0%), and then uniquely by eutrophication (SR: 20.3%, M-AMBI: 17.6%) and finally uniquely by natural variability (SR: 5.8%, M-AMBI: 8.9%).

Figure 5 - On the left, variance (adjusted R2) of the macrofauna community
data (community), total macrofauna density (density), inverse of Simpson's
dominance (N2), Shannon diversity (H'), species richness (SR) and multivariate
AMBI (M-AMBI) uniquely attributed to eutrophication (green), uniquely attributed
to natural variability (blue) and jointly attributed to both effects (red). The
residual variance is shown in white. On the right, eutrophication variables and
variables related to natural variability considered in the variance partitioning of
the macrofauna community and macrobenthic taxonomic indices.

Using figure 2, we can identify variables related to eutrophication or to natural
variability that significantly co-vary and are jointly implicated in a macrobenthic
index model (Table 3) and/or in a variance partitioning (Figure 5). Mean [Chla] was
significantly correlated with lagoon surface (-0.35), β_{MM} (-0.37), mean Temp (0.40)
and CV O_2 sat (0.40). Mean [NO_x] was significantly correlated with lagoon surface (-
0.38), mean Sal (-0.49) and CV O_2 sat (-0.33) while CV [NO_x] was significantly
correlated with lagoon surface (-0.37), β_{MM} (-0.36) and mean Temp (0.53). Finally,
mean [DIP] and lagoon surface were significantly correlated (-0.32). Lagoon-sea
connection, a qualitative variable, was another source of confounding effects
between natural variability and eutrophication on the macrobenthic indices and
community. Indeed, there was a significant difference between the lagoon-sea
connection and: *(i)* mean [Chla] (weak > strong connection); *(ii)* mean [NO_x] (weak >
strong connection) and *(iii)* CV [NO_x] (moderate > strong).

Discussion

As outlined by Cloern (2001), eutrophication is a complex phenomenon with direct
(*e.g.* changes in dissolved nutrient contents) and indirect (*e.g.* changes in
community) responses that are modulated by attributes of each recipient system.
This study is a first step towards better understanding how hydro-morphology,
natural environmental variability and eutrophication jointly shape macrobenthic
communities in Mediterranean coastal lagoon ecosystems.

Macrobenthic communities are driven by lagoon hydro-morphology and natural environmental (in)stability

Lagoon macrobenthic communities appear to be first determined by lagoon hydro-morphology and natural environmental stability. When investigating the independent effect of each environmental parameter on macrobenthic communities, lagoon-sea connection level was the variable explaining the most of H', N2 and M-AMBI variability and the second most important variable explaining community composition. This qualitative variable is a proxy for marine water renewal (*i.e.* residence time), a gradient well known for characterizing coastal lagoon biological communities (Barbone and Basset, 2010; Perez-Ruzafa and Marcos-Diego, 1992; Tagliapietra et al., 2009), and closely linked to mean Sal (moderate > weak and strong > weak) and CV Sal (weak > moderate > strong). Salinity, historically used to classify coastal lagoons (Anonymous, 1959), was also an important driver of SR (linear positive effect) and total macrofauna density (concave effect). Indeed, simple and multimetric descriptors of lagoon macrobenthic communities like species richness are known to be highly and positively influenced by mean water salinity (Barbone et al., 2012; Magni et al., 2022; Pitacco et al., 2019; Reizopoulou et al., 2014). Differently from us, Barbone et al. (2012) reported a macrofauna density decrease from oligo- to eu-haline lagoons, maybe because only undisturbed lagoons were considered in their study. Furthermore, the effect of lagoon-sea connection on N2, a diversity index sensitive to high abundance taxa (Morris et al., 2014), indicates that a strong connection to the sea promotes benthic communities with significantly more relatively abundant taxa whereas moderate and weak connections promote communities dominated by a few abundant and in our case, euryhaline taxa (*e.g. Hediste diversicolor, Streblospio shrubsolii*), a characteristic typical of naturally stressed environments like lagoons subject to strong salinity variations (Elliott and Quintino, 2007). This result seems to confirm that moderately and weakly connected lagoons are highly selective environments for benthic organisms at least partly because of their strongly fluctuating environmental conditions (Barnes, 1980; Provost et al., 2012; Zaldívar et al., 2008). The negative effect of decreasing salinity and lower connection levels to the sea on SR, N2 and macrofauna density can also be linked to the physiological stress that natural and abrupt salinity variations (*e.g.* droughts, heavy rainfall) can represent for larvae, juveniles and adults of marine-originating benthic organisms inhabiting coastal lagoons (Cognetti and Maltagliati, 2000; Magalhães et al., 2019; Vernberg, 1982). Overall and based on other published evidence, our results could be explained by (*i*) water renewal from marine origin, (*ii*) primary colonization and/or post-disturbance recolonization of lagoons by marine-originating larvae through dispersal and recruitment and (*iii*) environmental (in)stability, which are known to strongly shape lagoon benthic communities (Basset et al., 2006; Pérez-Ruzafa et al., 2007b; Pérez-Ruzafa and Marcos-Diego, 1992), as proposed in the confinement theory (Guelorget and Perthuisot, 1983).

Water temperature was another important driver of lagoon macrobenthic communities, SR and macrofauna density in our study. Temperature had a negative effect on these communities, probably directly through thermal stress, and indirectly through its effect on oxygen dynamics. Higher and more variable summer temperatures directly represent an abiotic constraint for benthic organisms by increasing their oxygen demand and reducing their mobility (Vernberg, 1982). At the same time, high summer temperatures decrease oxygen solubility and increase the

probability of hypoxia and anoxia (Derolez et al., 2020a) while also reducing the
tolerance of benthic organisms to low dissolved oxygen (Vaquer-Sunyer and Duarte,
2011). This cocktail of effects can ultimately cause the death of sensitive taxa, such
as large sized and long-lived species (Quillien et al., 2015; Reizopoulou and
Nicolaidou, 2007; Sigala et al., 2012) and reduce the ability to move of more tolerant
taxa (Vernberg, 1982), hence increasing their susceptibility to predation and
decreasing their possibility to search for food and/or for less stressful environmental
conditions (Díaz and Rosenberg, 1995; Rosenberg et al., 1992; Weissberger et al.,
2009).
The diversity of macrophyte morphotypes (β_{MM}) also affected positively H' and N2,
and was positively correlated to all macrobenthic indices except density and AMBI.
Simple taxonomic indices like lagoon macrobenthic species richness, biomass
density and numerical density are locally sensitive to vegetation presence and type
(Arocena, 2007; Barbone et al., 2012, Magni and Gravina, 2023). Conversely, we
show here the probable effect of benthic habitat heterogeneity expressed at the
lagoon or sub-lagoon scale on the structure of benthic communities at a local scale
(*ca.* 50 m^2 station), an effect mainly visible when focusing on intermediate and high
abundance taxa (Morris et al., 2014). Focusing on macrobenthic fauna at the lagoon
scale, β_{MM} informs us on the number of potential (*i*) environmental niches that can be
filled by the species pool and (*ii*) refuges (Magni and Gravina, 2023; Nordström and
Booth, 2007; Ware et al., 2019) from which these organisms can (re)colonize habitat
patches after mortality events caused by temporary unfavorable conditions. A model
of local extinctions linked to adverse environmental conditions, followed by
recolonization events via pelagic larvae was suggested by Vergara-Chen et al. (2013)
to explain the small-scale genetic structure of the cockle *Cerastoderma glaucum*
inside the Mar Menor lagoon (Spain). The positive effect of β_{MM} on macrobenthic
diversity also provides us with a few leads on how macrophytes can structure lagoon
macrobenthic communities. Macrophytes can act as (*i*) temporary refuges from
vagile predators for small mobile benthic organisms like amphipods (Nordström and
Booth, 2007; Ware et al., 2019), (*ii*) a food source for grazers and detritus-feeders
(Carlier et al., 2007; Lepoint et al., 2000; Ouisse et al., 2011; Vizzini and Mazzola,
2006) and (*iii*) settlement promoters for pelagic particles like phytoplankton and
larvae (Colden et al., 2016; Donadi et al., 2014; González-Ortiz et al., 2014),
especially in the case of meadow-forming species like *Zostera.* All these local effects
are likely to spill-over to adjacent unvegetated sediments (Boström et al., 2006;
Magni et al., 2017).
Overall, natural environmental (in)stability, connection level with adjacent water
systems and/or number of potential environmental niches seem to be the main
drivers of lagoon macrobenthic communities and to be stronger drivers of benthic
macrofauna assemblages than eutrophication, which probably represents an
additional source of stress for benthic invertebrates (Barbone and Basset, 2010).
The environmental variables considered in the manuscript do not unfortunately
explain the communities described at the LPN and LPS stations. These two
macrofauna stations sampled were actually classified in a group generally associated
to small eutrophic lagoons and mainly composed of stations sampled in lagoons
evaluated by the WFD as being in a poor or in a bad physico-chemical status even
through is considered in a good physico-chemical status according to Chla
concentration. This could be the result of the strong variations in salinity recorded
throughout the year, which we have not been considered here. In this lagoon,

marked annual salinity variations are observed (between 3.6 and 76.8 in Wilke and
Boutiere (2000)), variations indetectable with the available summer salinity data.
Consequently, the eight environmental variables selected to explain the overall
macrofauna communities were not able to explain the one characterizing this
lagoon. This result also reinforces the importance of natural environmental instability
probably has on structuring benthic macrofauna communities.
864
Eutrophication affects lagoon macrobenthic communities through
interconnections with natural environmental variability
Eutrophication appears to act as an additional disturbance for lagoon
macrobenthic communities via direct (*e.g.* toxicity of inorganic nitrogenous
compounds) and indirect mechanisms (*e.g.* hypoxic stress). The three groups derived
from the variance partitioning approach highlight that the most influential category
of variables (*i.e.* natural variability alone, eutrophication alone, natural variability
and eutrophication together) is not the same depending on the level of biological
organization we focus on. Indeed, the benthic invertebrate density was first
explained by natural variability alone and more precisely by physiologically
constraining variables (salinity and temperature) whereas macrobenthic community
composition, SR, H' and N2 were first explained jointly by natural variability and
eutrophication. With these joint effects, we probably measure *(i)* how lagoon-sea
connection modulates the susceptibility (*i.e.* propensity) of each lagoon to
eutrophication, *(ii)* how higher water temperatures promote phytoplankton blooms
and *(iii)* how eutrophication reduces benthic habitat heterogeneity, followed by their
respective effects on benthic community diversity.
(i) Lagoon-sea connection informs us on each lagoon's water residence time and
consequently, on its retention potential regarding nutrients and pollutants (Chapelle
et al., 2001; Cloern, 2001; Fiandrino et al., 2017). As such, watershed related
nutrients and their direct effect on primary producers are buffered by each lagoon's
degree of connection to the sea, as indicated by the significantly higher mean [Chla]
and mean [NO$_x$] in lagoons weakly connected to the sea compared with strongly
connected ones. *(ii)* Independently from lagoon hydro-morphology, the positive
correlation between mean [Chla] and mean temperature illustrates how higher water
temperatures increase phytoplankton growth rates which can lead to blooms (Rose
and Caron, 2007), as recently demonstrated in Thau (Trombetta et al., 2019), a large
lagoon strongly connected to the sea. Nonetheless, it seems that eutrophication is
buffered in larger lagoons where water temperatures increase less in summer, as
suggested by the negative correlation between lagoon surface and mean summer
temperature. *(iii)* Furthermore, one of the direct responses of Mediterranean coastal
lagoons to changes in nutrient inputs is an increase in pelagic primary production
(Souchu et al., 2010), which in turn decreases water transparency (Cloern, 2001) and
limits light availability for benthic primary producers (Le Fur et al., 2019, 2018)*.
Ultimately, sustained eutrophication of coastal lagoons can cause a regime shift from
a primary producer community dominated by perennial macrophytes presenting
different morphotypes, to a community dominated by free-floating and opportunistic
macroalgae like *Ulva rigida* and finally, to a phytoplankton-dominated community (Le
Fur et al., 2019). The negative correlation between several eutrophication variables
(*e.g.* mean [Chla], mean turbidity) and β_{MM} highlights the simplification of benthic
habitats caused by lagoon eutrophication (Cloern, 2001; Le Fur et al., 2019, 2018)
that shift from a mosaic of different macrophyte morphotypes to a sediment void of

macrophytes or with one macrophyte morphotype. Overall, the multiple
interconnections between natural environmental variability and eutrophication likely
explain the high amount of macrobenthic community and index variance jointly
attributed to both effects, as underlined in the conceptual model developed by
Cloern (2001).

The unique effect of eutrophication variables on each taxonomic component also
informs us about which of these variables likely have a direct effect on macrobenthic
invertebrates without being modulated by natural variability. Indeed, benthic
organisms can be affected by eutrophication directly through the toxicity of
inorganic nitrogenous compounds like ammonium, nitrites and nitrates (Jessen et al.,
2015). Our results indicate that high NH_4^+ concentrations seem to have a very small
negative effect on total macrofauna density while NO_x toxicity could explain the
strong unique effect of eutrophication on SR and M-AMBI, and both NH_4^+ and NO_x
toxicity could be responsible for the very weak unique effect of eutrophication on
macrobenthic communities. These results are probably also linked to indirect effects
high eutrophication levels can have on macrobenthic communities via increased
water turbidity and sediment bio-geochemistry modifications (*e.g.* anoxic superficial
sediments (Zilius et al., 2015)). Indeed, water NH_4^+ and NO_x concentrations in excess
can be the sign of highly eutrophic lagoons (Souchu et al., 2010). Overall,
experimental work on a few well-selected species and populations sampled in
lagoons presenting different levels of natural variability would greatly improve our
understanding of the effect of eutrophication on lagoon benthic invertebrates via
inorganic nitrogenous compound toxicity. Even trough marine animals appear to be
less sensitive to nitrate than freshwater animals, nitrate toxicity to benthic
invertebrate increases with increasing nitrate concentrations and exposure times
(Alonso and Camargo, 2006; Boardman et al., 2004; Camargo et al., 2005) Nitrate
toxicity may also increase with decreasing body size, water salinity, and
environmental adaptation. In lagoon environments, where organisms are often
smaller in size, this could also be of importance.

Furthermore, mean [Chla] appears to have a strong negative effect on
macrobenthic communities by impacting the diversity of low and intermediate
abundance taxa (i.e. SR and H' are diversity indices that give more weight
respectively, to low abundance/rare and intermediate abundance taxa) and this
effect is at least partly independent from the selected natural variability variables
(*i.e.* mean temperature, mean salinity for SR and lagoon-sea connection, β_{MM} for H').
Two non-exclusive hypotheses could explain this result: small-scale habitat
homogenization and hypoxic stress. Despite not being the appropriate index to
investigate beta diversity, the standard deviation of SR at the intra-station scale was
significantly correlated negatively to mean [Chla], which seems to point towards a
homogenization of benthic communities at the meter scale (i.e. a few meters
separate each replicate of a given station) linked to eutrophication. The loss of
benthic taxa presenting restricted environmental niches and dependent upon small-
scale habitat diversity for reproduction, recruitment, refuge and/or food (Brauns et
al., 2007; Harman, 1972; Stendera and Johnson, 2008) could explain this result.
Secondly, low to zero dissolved oxygen (*i.e.* hypoxia and anoxia) is another well-
known symptom of eutrophication in transitional waters like coastal lagoons
(Chapelle et al., 2001; Cladas et al., 2016; Vignes et al., 2010). Oxygen depletion
generally takes place during the summer and is maximal at the end of the
night/early morning (Newton et al., 2010), often starting near the bottom because of

benthic organic matter mineralization (*e.g.* sedimented phytoplankton, dead
macroalgae) and low water renewal (Rigaud et al., 2021; Souchu et al., 1998). In the
worst cases, these low to zero oxygen zones can spread to the entire water column
and last for several days (Chapelle et al., 2001; Cladas et al., 2016), as seen during
the 2018 dystrophic crisis in Thau lagoon (Derolez et al., 2020a). These crises are
known to impact macrobenthic community first in terms of abundance and then in
terms of composition and species richness depending on the sensitivity of each
species to reduced oxygen (Díaz and Rosenberg, 1995; Riedel et al., 2012).
Consequently, lagoon macrobenthic invertebrates are often reported as being less
abundant and less diverse in fall than in spring because of summer hypoxic stress
(Barbone et al., 2012; Basset et al., 2013a; Orro and Cabana, 2021). The negative
impact of eutrophication via hypoxic/anoxic stress on lagoon macrobenthic
communities is suggested by the RDA results where high mean O_2 sat (*i.e.* hyper-
saturation linked to strong pelagic primary production) is associated to group A
stations that present significantly lower species richness than group B stations.
Nonetheless, oxygen related variables were not selected in the other models, a
result probably linked to when oxygen was measured (late morning) and at what
depth it was measured (generally sub-surface). Indeed, these features limit our
ability to detect potential short and long-term effects of hypoxic stress on
macrobenthic communities. Bottom hypoxia/anoxia is also known to predominantly
take place in deep lagoons like Berre and Thau (Derolez et al., 2020a; Rigaud et al.,
2021; Souchu et al., 1998), where wind derived water mixing is limited by depth.
Nonetheless, anoxic crises inside coastal lagoons are expected to increase in
frequency and intensity with climate change (Derolez et al., 2020a; Miyamoto et al.,
2019). Consequently, it appears paramount to (*i*) study the sensitivity of lagoon
benthic organisms to low dissolved oxygen, not only of freshwater and marine taxa
(Díaz and Rosenberg, 1995; Landman et al., 2005) and (*ii*) monitor more closely
summer bottom dissolved oxygen concentrations, especially at higher frequency (O_2
being very variable) and at temporal scales relevant to these organisms (*i.e.* across
several days starting when bottom hypoxia is detected (Riedel et al., 2012)).
Overall, lagoon hydro-morphology (connection to the sea, surface and depth)
which affects the level of environmental variability (salinity and temperature), as
well as lagoon-scale benthic habitat diversity (β_{MM}) seem to regulate the distribution
of macrobenthic species (Bachelet et al., 2000), while eutrophication and associated
stressors like low dissolved oxygen act upon the existing communities, mainly by
reducing taxa richness and diversity (SR and H'). Disentangling the effects of
anthropogenic stressors and natural environmental variables remains nevertheless
challenging, especially as stressors and variables can co-vary, even in the absence
of anthropogenic impact. Moreover, it is often impossible to measure effects of every
single possible stressor on macrobenthic communites. In our study, grain-size could
not be included to differences in laboratory protocols between lagoons and between
years and heavy metal contaminations were not available at the same temporal and
spatial scale (one measurement per lagoon every ten years). It should be relevant to
re-run the analyses when such data would be available at the macrofauna sampling
sites to test the effect of this supplementary anthropogenic driver to check if it
would be discriminating benthic communities.
1002
The M-AMBI and recommendations for its use as an indicator for
Mediterranean coastal lagoons

With this study, we investigate for the first time the joint sensitivity of the M-AMBI
index to natural and anthropogenic stressors across 29 lagoons covering most of the
French Mediterranean coastline. The M-AMBI (Muxika et al., 2007) is currently used
to evaluate the WFD ecological state of poly- and eu-haline French Mediterranean
lagoons based on soft sediment benthic invertebrate, with reference conditions using
minimally impacted sites (European Commission, 2018). This index has previously
been shown to be sensitive to pressures related to eutrophication, like concentration
in chlorophyll *a* and total nitrogen (Derolez et al., 2014) but it is also known to be
sensitive to natural environmental variability (*e.g.* water salinity) and lagoon hydro-
morphology (*e.g.* surface and depth) (Barbone et al., 2012).
Our results show that the M-AMBI appears to be a sensitive indicator of
eutrophication through the species richness and Shannon diversity components of
the index but not the AMBI one. First, the strong correlation between M-AMBI and
mean [Chla] was linked only to the significant correlation between SR and mean
[Chla] and between H' and mean [Chla] while AMBI was not significantly linked to
most eutrophication related variables (except mean [NO_x]) nor to chlorophyll *a*
concentration, a common eutrophication indicator (Bec et al., 2011; Cloern, 2001).
Importantly, these linear relations remained significant once the effect of identified
covariables was removed. Regarding AMBI, its current application in transitional
waters like coastal lagoons suffers from methodological limitations. Indeed, this
index is based on macroinvertebrate sensitivity/tolerance to organic matter
enrichment following Pearson and Rosenberg (1978) and was developed for coastal
waters (Borja et al., 2000), not for coastal lagoons, which are known to sometimes
present sediments naturally enriched in organic matter linked to the decay of
meadow-forming macrophytes (Bachelet et al., 2000), watershed inputs and the
overall low organic matter exportation (except during floods) (Perthuisot and
Guelorget, 1983). These results highlight the limits of indirectly including AMBI in the
evaluation of the ecological state of Mediterranean coastal lagoons. Nonetheless, its
absence of sensitivity to natural variability and lagoon hydro-morphology suggests
an index based on redefined ecological groups could be promising (Robertson et al.,
2016), especially if we aim towards a more specific macrofauna-based index to
evaluate eutrophication. Overall, increasing the robustness of M-AMBI would likely
necessitate additional work to classify lagoon macroinvertebrates according to other
eutrophication related stressors like low oxygen, nitrogenous compound toxicity
and/or hydrogen sulfide toxicity.
Secondly, M-AMBI variability was two times more strongly attributed uniquely to
eutrophication related variables than uniquely to natural variability ones, with
nonetheless a very strong joint effect of the two types of variables. Our study
reinforce the difficulty of M-AMBI to clearly identify natural disturbance but also the
high sensitivity of this multimetric index to anthropogenic pressures such as
eutrophication or metal contaminations and pesticides in transitional waters (Paul et
al., 2023; Pelletier and Charpentier, 2023; Pollice et al., 2015). The strong joint effect
is probably related, as previously discussed, to the same interconnections between
lagoon-sea connection that buffers eutrophication, temperature that promotes
phytoplankton blooms and eutrophication that homogenizes benthic habitats.
Several modifications could help limit the sensitivity of M-AMBI to natural variability.
First, as recommended by Borja et al. (2012) who pointed that the inability of this
index to detect anthropogenic responses is often linked to the use of inappropriate
reference conditions, the currently used poly- and eu-haline vs oligo- and meso-

haline categories (Provost et al., 2012; European Commission, 2018) could be
replaced by one based on the three lagoon-sea connection levels that strongly
influence the M-AMBI or on another typology like the choked vs restricted typology
(Barbone et al., 2012; Basset et al., 2013b), with the test of new reference lagoons
(European Commission, 2018). Secondly, the replacement of a sampling done with
an Ekman-Birge grab by a diver-operated sampling would limit heterogeneity between
replicats by (i) verifying that only unvegetated sediments are being collected and (ii)
describing the presence and type of macrophyte near the sampling site (*e.g.* 10
meter radius). This could represent a bias in many homogeneous environments
where random selection of the sampling zone is quite possible but macrophytes are
often heterogeneously distributed in coastal lagoon and areas of unvegetated
sediments are rare. Selection and description of the sampling area by divers could
therefore be relevant in these shallow environments even if this often means
additional human and financial costs. This last modification would likely help reduce
the effect of local benthic habitat heterogeneity on H' and M-AMBI (Magni and
Gravina, 2023; de Paz et al., 2008).

Finally, several studies have underlined the limits of using the M-AMBI based on
abundance to evaluate the ecological state of transitional ecosystems like coastal
lagoons, as opportunistic and tolerant species are generally numerically dominant in
these systems (Mistri et al., 2018; Pitacco et al., 2019; Puente and Diaz, 2008). Other
indicators based on the size spectra and biomass of benthic macroinvertebrates like
the biomass based M-AMBI (M-bAMBI (Mistri et al., 2018)) and the index of size
distribution (ISD (Reizopoulou and Nicolaidou, 2007)) have showed promising results
and applying these indices to French coastal lagoons would represent an additional
trial in the context of historically highly eutrophic lagoons currently undergoing a re-
oligotrophication trajectory (Derolez et al., 2020a, 2020b; Le Fur et al., 2019; Leruste
et al., 2016). Another interesting avenue would be to investigate more closely the
spatial scales at which changes in macroinvertebrate abundance and diversity can
be detected and link these spatial changes to localized anthropogenic disturbances
like heavy metals (Magalhães et al., 2019) while also considering small-scale biotic
interactions such as competition, predation and facilitation. Indeed, sediment
contaminants like biotic interactions are likely to affect benthic macrofauna at fine
spatial scales (*ca.* 1 km) that are more relevant to low mobility macroinvertebrates
(Berthelsen et al., 2018; Brauko et al., 2020, 2015; Carvalho et al., 2006).

In a nutshell, to improve the robustness of M-AMBI, we call for a revision of the
ecological groups at the base of the AMBI index and of the sampling protocol (*e.g.*
using divers) along with a re-evaluation of the lagoon typology. More generally, it
appears key to increase our understanding of the effects of other disturbances like
hypoxia and heavy metals on lagoon invertebrates and to reevaluate the small-scale
distribution of benthic organisms in French coastal lagoons, more than 30 years after
the historical studies by Amanieu et al. (1977), Guelorget et al. (1994) and Guelorget
and Michel (1979).
1096

Overcoming taxonomic limitations
1098   A large part of the identified macroinvertebrates had a very restricted
1099 geographical range, preventing us from building a general species-based framework
1100 linking eutrophication and macrobenthic communities. Indeed, we found a very high
1101 level of lagoon-specificity for benthic macroinvertebrates across French
1102 Mediterranean coastal lagoons with no taxon present across all 41 stations, 40% of

the 227 taxa identified in only one station and only 1.3% recorded in over half of the stations. This result is probably explained by environmental filtering resulting from very contrasted conditions across the different sampled systems, passive larval diffusion (Basset et al., 2007) and the randomness of the colonization process by species of marine origin (Pérez-Ruzafa et al., 2011b), a phenomenon previously observed for benthic macroinvertebrates in Italian coastal lagoons (Basset et al., 2007), fish in Atlantic and Mediterranean coastal lagoons (Pérez-Ruzafa et al., 2007b) and macrophytes in estuaries and lagoons (Pérez-Ruzafa et al., 2011c). Consequently, unique assemblages of benthic macroinvertebrates are present in each lagoon and even at smaller spatial scales (*i.e.* stations inside a lagoon), which could explain why we were able to explain only 22% of the macrobenthic community variability. This high level of lagoon-specificity present across multiple biological compartments in coastal lagoons warrants the need to go beyond a species-based approach and towards a trait-based approach (Lacoste et al., 2023), as previously outlined using macrofauna size (Basset et al., 2008; Brauko et al., 2020; Sigala et al., 2012). Such an approach could build upon the saprobity concept (Tagliapietra et al., 2012) to select relevant biological traits and ultimately help us build a more general framework around macrobenthic community assembly rules in the line of what Pérez-Ruzafa et al. (2013) did for fish by revisiting the r and K-strategies. Nonetheless, for such an approach to be relevant, we also need to increase our knowledge of the life-cycle of lagoon benthic invertebrates (Brauko et al., 2020) and to create a harmonized Mediterranean wide trait database for these organisms (Lacoste et al., 2023).

Another interesting avenue to pursue could be to investigate intra-specific biological trait and/or physiological response variability (*i.e.* oxydative stress) of species like the polychaete *Hediste diversicolor* along eutrophication gradients. Indeed, this species is known to display a great diversity of life history characteristics along with morphological, biochemical and physiological differences according to local environmental conditions (Scaps, 2002). For example, the feeding behavior and diet of this species has been shown to vary according to sewage pollution levels across English estuaries (Aberson et al., 2016). Another study has related oxydative stress of the polychaete *Nephtys cirrosa* to seasonal variations of abiotic parameters like temperature and salinity in the Ria de Aveiro (Magalhães et al., 2019). Furthermore, *H. diversicolor,* a species we found to be associated to eutrophic lagoons, is reported to present a great diversity of feeding modalities (*i.e.* predator, grazer, suspension-feeder and deposit-feeder (Faulwetter et al., 2014)), whereas the species associated to oligotrophic lagoons, like *Loripes orbiculatus,* present less diverse feeding modes (*i.e.* mainly based on sulfo-oxydizing bacteria (Roques et al., 2020)). This flexibility could allow organisms to better tolerate the fluctuating abiotic conditions that characterize small eutrophic lagoons, stressing that environmental stability is probably a strong driver of macrobenthic communities in coastal lagoons, as previously shown for macrophytes (Pérez-Ruzafa et al., 2011c). Overall, the plasticity several species display in environmentally constraining conditions stresses the difficulty to implement trait-based approaches in ultra-variable environmental systems like coastal lagoons, where realized trait modalities may differ significantly from the ones available in the literature, commonly acquired from non-transitional coastal ecosystems (Martini et al., 2021). Finally, a functional approach based on well-informed biological traits and biomass data could help overcome the identified taxonomic limitations and build a more holistic framework linking natural variability

1152 and eutrophication of coastal lagoons to macrobenthic communities but the intra-
1153 specific plasticity for many biological traits makes this approach questionable today.

Conclusion

Coastal lagoons are transitional ecosystems naturally characterized by a strong
variability in abiotic parameters like salinity and temperature. Superimposed onto
this natural environmental variability are numerous anthropogenic disturbances,
eutrophication being a historical one targeted by the European Water Framework
Directive. By jointly analyzing water column, sediment and macrophyte parameters,
our study shows that macrobenthic communities across 29 French Mediterranean
coastal lagoons are primarily determined by (*i*) lagoon hydro-morphology
(connection level with the sea, surface and depth) which further affects natural
environmental variability in terms of salinity and temperature, and (*ii*) lagoon-scale
benthic habitat diversity, as estimated through macrophyte morphotypes.
Eutrophication and associated stressors like low dissolved oxygen act upon the
existing communities by reducing taxonomic richness and diversity. Eutrophication
appeared to act on lagoon macrobenthos through relatively direct mechanisms like
toxicity of inorganic nitrogenous compounds, as well as indirectly via for example,
the simplification of benthic habitats and sediment bio-geochemistry modifications.
Furthermore, our study highlights the complex interconnections between lagoon
hydro-morphology, natural environmental variability and eutrophication and their
joint effect on benthic communities. Indeed, lagoon-sea connection modulates the
susceptibility of each lagoon to eutrophication, high water temperatures promote
eutrophication related phytoplankton blooms while eutrophication reduces benthic
habitat heterogeneity, all of which affect benthic communities. Finally, benthic taxa
richness and diversity also proved to be indicators sensitive to eutrophication in
coastal lagoons while an AMBI like index based on redefined ecological groups for
coastal lagoons, which can be naturally enriched in organic matter, has the potential
of being a specific eutrophication indicator.
1180
1181

1182

Appendices

Appendix 1 - Procedure used to calculate the beta diversity of macrophyte morphotypes (β_{MM}) corresponding to each macrofauna station

Step 1: Select the macrophyte stations corresponding to each macrofauna station
- If one macrofauna station is considered in a given lagoon, then all the macrophyte stations of that given lagoon are taken into account.
- If two or three macrofauna stations are considered in a given lagoon, then all the macrophyte stations attributed by the WFD protocol to the same section of the lagoon as the macrofauna station are taken into account.

Step 2 : Build the macrophyte station by macrophyte morphotype matrix for each macrofauna station
For each macrophyte station of the dataset:
- If the macrophyte cover is below 25% or equal to 25%, a value of 1 is entered in the sediment column and 0 in all the others (line 1 of the example matrix below).
- If the cover is above 25%, the dominant or co-dominant taxa are determined based on the taxa specific cover data or the taxa specific biomass data, then these taxa are recoded into a morphotype (see Appendix 2) and finally, a value of 1 is entered in all the cells corresponding to the given morphotypes of the dominant or co-dominant taxa and a value of 0 is entered in the remaining cells (lines 2 and 3 of the example matrix below).

Step 3 : Calculate the index value corresponding to each macrofauna station
- Calculate the Bray-Curtis dissimilarity between all the station pairs using the previously built station by macrophyte morphotype matrix (presence/absence)
- Average the Bray-Curtis dissimilarities

Example of a macrophyte station by macrophyte morphotype matrix used to calculate the β_{MM}

Station	mushroom	feather	hair	ball	branched	sheet	meadow	sediment
1	0	0	0	0	0	0	0	1
2	0	0	0	0	1	0	1	0
3	0	0	0	0	0	1	0	0
…								
n-1								
n								

1215

1216

1217

1218 **Appendix 2 - Correspondence between macrophyte taxa and morphotypes**

Macrophyte taxa	Morphotype
Acetabularia acetabulum	mushroom
Aglaothamnion pseudobyssoides	
Alsidium corallinum	
Cladophora vagabunda	feather
Dasyaceae	
Spyridia filamentosa	
Chaetomorpha aerea	hair
Ulva clathrata	
Cladophora glomerata	ball
Valonia aegagropila	
Gracilaria dura	
Gracilaria gracilis	branched
Halopithys incurva	
Ulva rigida	sheet
Cymodocea nodosa	
Ruppia cirrhosa	
Stuckenia pectinata	meadow
Zostera marina	
Zostera noltei	

1219

1220

Appendix 3
Mean ± SD of the species richness (SR), Shannon diversity (H'), inverse of Simpson
dominance index (N2), macrofauna density, AMBI (AZTI's Marine Biotic Index) and M-
AMBI (Multivariate AMBI) calculated for the A and B station groups determined using
a K-means partitioning based on the benthic macrofauna Hellinger-transformed
mean abundances. The results of the non-parametric Kruskal-Wallis rank sum test
comparing the simple and multimetric macrobenthic taxonomic indices between the
A and B groups are also indicated.

	Group A	Group B	Kruskal-Wallis rank sum test
SR	9.30 ± 3.48	17.48 ± 11.41	$\chi^2(1) = 6.41$, $p = 0.011$
H'	2.00 ± 0.57	2.36 ± 0.69	$\chi^2(1) = 2.29$, $p = 0.13$
N2	3.30 ± 1.35	3.83 ± 1.76	$\chi^2(1) = 0.74$, $p = 0.39$
Macrofauna density (ind.m^{-2})	2 280 ± 1 372	5 113 ± 5 205	$\chi^2(1) = 0.98$, $p = 0.32$
AMBI	3.03 ± 0.53	2.87 ± 1.08	$\chi^2(1) = 0.01$, $p = 0.92$
M-AMBI	0.43 ± 0.08	0.53 ± 0.14	$\chi^2(1) = 5.76$, $p = 0.016$

1229
1230

Appendix 4

Coordinates (biplot scores for quantitative variables and centroids for lagoon-sea connection) on the first and second axis of the redundancy analysis (RDA) plots, marginal effects and conditional effects of the 8 environmental variables selected to explain the macrobenthic community variability. After the abbreviated name of each environmental variable is indicated the type of transformation used to normalize it ($\log_e$: natural logarithm and sqrt: square-root). [Chla] and [DIP] is the water concentration in chlorophyll *a* ($\mu g.L^{-1}$) and dissolved inorganic phosphorous ($\mu mol.L^{-1}$), respectively. Temp (°C) is the lagoon water temperature. O_2 sat (%) is the water oxygen saturation level. OMC (%) is the sediment organic matter content. CV stands for coefficient of variation.

Selected environmental variable	RDA1	RDA2	Marginal effect (%)	Conditional effect (%)
Mean [Chla] ($\log_e$)	-0.76	0.23	7.22	7.22
Lagoon surface ($\log_e$)	0.73	-0.15	5.90	1.69
Weak lagoon-sea connection	-0.10	0.49		
Moderate lagoon-sea connection	-0.073	-0.18	5.30	4.64
Strong lagoon-sea connection	0.18	-0.29		
Mean Temp	-0.61	-0.24	5.04	3.62
Mean [DIP] ($\log_e$)	-0.51	0.092	3.16	1.31
CV Temp (sqrt)	0.44	0.15	2.20	1.53
OMC	0.11	0.27	0.50	1.20
Mean O_2 sat ($\log_e$)	-0.16	-0.063	0.47	1.28

1256
1257

Appendix 5

Partial redundancy analysis plots (pRDA, axis 1 and 2) focusing on (a) the benthic macrofauna sampled across 41 stations located in 29 lagoons along the French Mediterranean coast (see Table 1 for the full lagoon names and corresponding stations) as explained by keeping constant the natural environmental variability variables (lagoon-sea connection, lagoon surface, mean and CV water temperature, mean oxygen saturation and sediment organic matter content) and letting the eutrophication variables vary (mean concentration in chlorophyll *a* (mean [Chla]) and mean dissolved inorganic phosphorous (mean [DIP]), (b) the environmental variables left to vary where mean [Chla] and mean [DIP] are $\log_e$ transformed (see Tables 1 and 2 for the full environmental variable names and values) and (c) the macrofauna taxa, where only taxa with a goodness of fit (cumulative proportion of variance explained by the first two axis) above 0.23 are plotted. In (a), inter-station distances are euclidean while in (b) and (c) angles between all vectors reflect linear correlations. Across all plots, each station (point) is represented using a symbol for its salinity type (triangle: oligo- and meso-haline and circle: poly- and eu-haline) and a color (red: group A and blue: group B) for its group membership (K-means partitioning) based on Hellinger-transformed mean abundances.

Appendix 6

Partial effect of mean chlorophyll *a* concentration ([Chla]) on (a) species richness (SR) once the effect of mean temperature and mean salinity has been removed, (b) Shannon diversity (H') once the effect of lagoon-sea connection, CV [TN], CV [TN]^2 and β_{MM} has been removed and (c) M-AMBI once the effect of lagoon-sea connection and CV [NO$_x$] has been removed (see Table 3 for the parsimonious linear models) across 41 stations located in 29 lagoons along the French Mediterranean coast (see Table 1 for the full lagoon names and corresponding stations). The dashed lines correspond to simple linear regression models and the shaded zone to the confidence interval. Each station (point) is represented using a symbol for its salinity type (triangle: oligo- and meso-haline and circle: poly- and eu-haline) and a color (red: group A and blue: group B) for its group membership (K-means partitioning) based on Hellinger-transformed mean abundances.

1293

Acknowledgements

The authors would like to thank all the members of the LERLR and LERPAC Ifremer
labs as well as all the managers of the different French Mediterranean lagoons for
sampling and analyzing the water column parameters, the sediment parameters and
the macrofauna. A special thanks to Corinne Pélaprat (STARESO), Sébastien Thorin
(CREOCEAN), Thibault Schwartz (CREOCEAN) and Daniel Martin (Centre d'Estudis
Avançats de Blanes – CEAB) for sampling, identifying, and analyzing the macrofauna.
The authors are also extremely grateful to Aurélie Fouveau (Ifremer, LERBN) for
extracting the ecological group values, Olivier Boutron (Tour du Valat) for the
information on the Vaccarès lagoon system and Marion Richard (Ifremer, LERLR) for

the fruitful discussions. Finally, we would like to thank all our close colleagues from the LERLR (Ophélie Serais, Elise Bellamy, Aurélien Bouquet, Elise Caillard, Nicolas Cimiterra, Christine Felix, Elise Foucault, Camille Gianaroli, Yannick Gueguen, Elise Lacoste, Franck Lagarde, Julie Le Ray, Danièle Martin, Grégory Messian, Dominique Munaron, Hervé Violette) for creating such a cheerful, dynamic, and enthusiastic work environment, despite all the COVID-19 ups and downs.

Data, scripts, code, and supplementary information availability

Data, scripts, and code are available online on Zenodo: https://doi.org/10.5281/zenodo.10825165

Supplementary information is available online on biorxiv as part of the preprint: https://www.biorxiv.org/content/10.1101/2022.08.18.504439

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

Funding

This work is part of the MALAG project (Effect of eutrophication on the biodiversity of Mediterranean coastal lagoon benthic macrofauna) funded by "Office Français de la Biodiversité" and Ifremer. A. G. J., M. C. and T. L. M. received funding through the MALAG project. All the data presented in this study was collected during the projects "Réseau de Suivi Lagunaire" and Water Framework Directive, which received financial and technical support from Ifremer, "Agence de l'Eau Rhône Méditerranée Corse", "Région Languedoc-Roussillon/Occitanie" and technical support from "Cépralmar".

References

Aberson, M.J.R., Bolam, S.G., Hughes, R.G., 2016. The effect of sewage pollution on the feeding behaviour and diet of *Hediste* (*Nereis diversicolor* (O.F. Müller, 1776)) in three estuaries in south-east England, with implications for saltmarsh erosion. Marine Pollution Bulletin 105, 150–160. https://doi.org/10.1016/j.marpolbul.2016.02.033

Alonso, A., Camargo, J.A., 2006. Toxicity of nitrite to three species of freshwater invertebrates. Environmental Toxicology 21, 90–94. https://doi.org/10.1002/tox.20155

Amanieu, M., Guelorget, O., Michel, P., 1977. Richesse et diversité de la macrofaune benthique d'une lagune littorale méditerranéenne. Vie et Milieu XXVII, 85–109.

Andral, B., Sargian, P., 2010a. Directive Cadre Eau. District Rhône et Côtiers méditerranéens. Contrôle de surveillance/opérationnel. Campagne DCE 2009. RST.DOP/LER-PAC/10-19. https://archimer.ifremer.fr/doc/00038/14923/

Andral, B., Sargian, P., 2010b. Directive Cadre Eau. District Corse. Contrôle de surveillance/opérationnel. Campagne DCE 2009. RST.DOPLER/PAC/10-20. https://archimer.ifremer.fr/doc/00028/13918/

Anonymous, 1959. Symposium on the classification of brackish waters. Presented at the Archivio di Oceanografia e Limnologia, Venice, 8–14 April 1958.

Anonymous, 1958. The Venice System for the classification of marine waters according to salinity. Limnology and Oceanography 3: 346–347.

Arocena, R., 2007. Effects of submerged aquatic vegetation on macrozoobenthos in a coastal lagoon of the southwestern atlantic. International Review of Hydrobiology 92, 33–47. https://doi.org/10.1002/iroh.200610881

Bachelet, G., de Montaudouin, X., Auby, I., Labourg, P.-J., 2000. Seasonal changes in macrophyte and macrozoobenthos assemblages in three coastal lagoons under varying degrees of eutrophication. ICES Journal of Marine Science 57, 1495–1506. https://doi.org/10.1006/jmsc.2000.0902

Barbone, E., Basset, A., 2010. Hydrological constraints to macrobenthic fauna biodiversity in transitional waters ecosystems. Rend. Fis. Acc. Lincei 21, 301–314. https://doi.org/10.1007/s12210-010-0090-4

Barbone, E., Rosati, I., Reizopoulou, S., Basset, A., 2012. Linking classification boundaries to sources of natural variability in transitional waters: A case study of benthic macroinvertebrates. Ecological Indicators, Marine Benthic Indicators 12, 105–122. https://doi.org/10.1016/j.ecolind.2011.04.014

Barnes, R.S.K., 1980. Coastal lagoons: the natural history of a neglected habitat, Cambridge University Press. ed. Cambridge.

Basset, A., Barbone, E., Borja, Á., Elliott, M., Jona-Lasinio, G., Marques, J., Mazik, K., Muxika, I., Neto, J., Reizopoulou, S., Rosati, I., Teixeira, H., 2013a. Natural variability and reference conditions: setting type-specific classification boundaries for lagoon macroinvertebrates in the Mediterranean and Black Seas. Hydrobiologia. https://doi.org/10.1007/s10750-012-1273-z

Basset, A., Barbone, E., Elliott, M., Li, B.-L., Jorgensen, S.E., Lucena-Moya, P., Pardo, I., Mouillot, D., 2013b. A unifying approach to understanding transitional waters: Fundamental properties emerging from ecotone ecosystems. Estuarine, Coastal and Shelf Science, Estuarine and lagoon biodiversity and their natural goods and services 132, 5–16. https://doi.org/10.1016/j.ecss.2012.04.012

Basset, A., Galuppo, N., Sabetta, L., 2007. Environmental heterogeneity and benthic macroinvertebrate guilds in italian lagoons. Transitional Waters Bulletin 1, 48–63. https://doi.org/10.1285/i1825226Xv1n1p48

Basset, A., Sabetta, L., Fonnesu, A., Mouillot, D., Do Chi, T., Viaroli, P., Giordani, G., Reizopoulou, S., Abbiati, M., Carrada, G.C., 2006. Typology in Mediterranean transitional waters: new challenges and perspectives. Aquatic Conservation: Marine and Freshwater Ecosystems 16, 441–455. https://doi.org/10.1002/aqc.767

Basset, A., Sabetta, L., Sangiorgio, F., Pinna, M., Migoni, D., Fanizzi, F., Barbone, E., Galuppo, N., Fonda Umani, S., Reizopoulou, S., Nicolaidou, A., Arvanitidis, C., Moncheva, S., Trajanova, A., Georgescu, L., Beqiraj, S., 2008. Biodiversity conservation in Mediterranean and Black Sea lagoons: a trait-oriented approach to benthic invertebrate guilds. Aquatic Conservation: Marine and Freshwater Ecosystems 18, S4–S15. https://doi.org/10.1002/aqc.979

Bec, B., Collos, Y., Souchu, P., Vaquer, A., Lautier, J., Fiandrino, A., Benau, L., Orsoni, V., Laugier, T., 2011. Distribution of picophytoplankton and nanophytoplankton along an anthropogenic eutrophication gradient in French Mediterranean coastal lagoons. Aquatic Microbial Ecology 63, 29–45. https://doi.org/10.3354/ame01480

Bendoricchio, G., Di Luzio, M., Baschieri, P., Capodaglio, A.G., 1993. Diffuse pollution
in the lagoon of Venice. Water Science and Technology 28, 69–78.
https://doi.org/10.2166/wst.1993.0405
Benjamini, Y., Hochberg, Y., 1995. Controlling the false discovery rate: a practical
and powerful approach to multiple testing. Journal of the Royal Statistical Society.
Series B (Methodological) 57, 289–300. https://doi.org/10.1111/j.2517-
6161.1995.tb02031.x
Berthelsen, A., Atalah, J., Clark, D., Goodwin, E., Patterson, M., Sinner, J., 2018.
Relationships between biotic indices, multiple stressors and natural variability in
New Zealand estuaries. Ecological Indicators 85, 634–643.
https://doi.org/10.1016/j.ecolind.2017.10.060
Blanchet, F.G., Legendre, P., Borcard, D., 2008. Forward selection of explanatory
variables. Ecology 89, 2623–2632. https://doi.org/10.1890/07-0986.1
Boardman, G.D., Starbuck, S.M., Hudgins, D.B., Li, X., Kuhn, D.D., 2004. Toxicity of
ammonia to three marine fish and three marine invertebrates. Environmental
Toxicology 19, 134–142. https://doi.org/10.1002/tox.20006
Borja, A., Franco, J., Pérez, V., 2000. A marine biotic index to establish the ecological
quality of soft-bottom benthos within European estuarine and coastal
environments. Marine Pollution Bulletin 40, 1100–1114.
https://doi.org/10.1016/S0025-326X(00)00061-8
Borja, A., Muxika, I., 2005. Guidelines for the use of AMBI (AZTI's Marine Biotic Index)
in the assessment of the benthic ecological quality. Marine Pollution Bulletin 50,
787–789. https://doi.org/10.1016/j.marpolbul.2005.04.040
Boström, C., Jackson, E.L., Simenstad, C.A., 2006. Seagrass landscapes and their
effects on associated fauna: A review. Estuarine, Coastal and Shelf Science,
Ecological and Management Implications on Seagrass Landscapes 68, 383–403.
https://doi.org/10.1016/j.ecss.2006.01.026
Brauko, K.M., Cerveira, A.F., Freitas, R.F., Hadlich, H.L., Beras, T., Fonseca, A.L.,
Pagliosa, P.R., 2020. Comparing structural and functional approaches to
distinguish benthic communities from eutrophic and non-eutrophic estuaries.
Marine Pollution Bulletin 157, 111290.
https://doi.org/10.1016/j.marpolbul.2020.111290
Brauko, M.K., de Souza, F.M., Muniz, P., de Camargo, M.G., Lana, P. da C., 2015.
Spatial variability of three benthic indices for marine quality assessment in a
subtropical estuary of Southern Brazil. Marine Pollution Bulletin 91, 454–460.
https://doi.org/10.1016/j.marpolbul.2014.10.025
Brauns, M., Garcia, X.-F., Walz, N., Pusch, M.T., 2007. Effects of human shoreline
development on littoral macroinvertebrates in lowland lakes. Journal of Applied
Ecology 44, 1138–1144. https://doi.org/10.1111/j.1365-2664.2007.01376.x
Breslow, N., 1970. A generalized Kruskal-Wallis test for comparing k samples subject
to unequal patterns of censorship. Biometrika 57, 579–594.
https://doi.org/10.2307/2334776
Caliceti, M., Argese, E., Sfriso, A., Pavoni, B., 2002. Heavy metal contamination in the
seaweeds of the Venice lagoon. Chemosphere 47, 443–454.
https://doi.org/10.1016/S0045-6535(01)00292-2
Camargo, J.A., Alonso, A., Salamanca, A., 2005. Nitrate toxicity to aquatic animals: a
review with new data for freshwater invertebrates. Chemosphere 58, 1255–1267.
https://doi.org/10.1016/j.chemosphere.2004.10.044

Cardoso, P.G., Pardal, M.A., Raffaelli, D., Baeta, A., Marques, J.C., 2004. Macroinvertebrate response to different species of macroalgal mats and the role of disturbance history. Journal of Experimental Marine Biology and Ecology 308, 207–220. https://doi.org/10.1016/j.jembe.2004.02.018

Carlier, A., Riera, P., Amouroux, J.-M., Bodiou, J.-Y., Escoubeyrou, K., Desmalades, M., Caparros, J., Grémare, A., 2007. A seasonal survey of the food web in the La Palme lagoon (northwestern Mediterranean) assessed by carbon and nitrogen stable isotope analysis. Estuarine, Coastal and Shelf Science 73, 299–315. https://doi.org/10.1016/j.ecss.2007.01.012

Carvalho, S., Gaspar, M.B., Moura, A., Vale, C., Antunes, P., Gil, O., da Fonseca, L.C., Falcão, M., 2006. The use of the marine biotic index AMBI in the assessment of the ecological status of the Obidos lagoon (Portugal). Marine Pollution Bulletin 52, 1414–1424. https://doi.org/10.1016/j.marpolbul.2006.04.004

Chapelle, A., Lazure, P., Souchu, P., 2001. Modélisation numérique des crises anoxiques (malaïgues) dans la lagune de Thau (France). Oceanologica Acta Supplement 1, 87–97. https://doi.org/10.1016/S0399-1784(00)01109-9

Cladas, Y., Papantoniou, G., Bekiari, V., Fragkopoulu, N., 2016. Dystrophic crisis event in Papas Lagoon, Araxos Cape, Western Greece in the summer 2012. Mediterranean Marine Science 17, 32–38. https://doi.org/10.12681/mms.1409

Cloern, J.E., 2001. Our evolving conceptual model of the coastal eutrophication problem. Marine Ecology Progress Series 210, 223–253. https://doi.org/10.3354/meps210223

Cognetti, G., Maltagliati, F., 2000. Biodiversity and adaptive mechanisms in brackish water fauna. Marine Pollution Bulletin 40, 7–14. https://doi.org/10.1016/S0025-326X(99)00173-3

Colden, A.M., Fall, K.A., Cartwright, G.M., Friedrichs, C.T., 2016. Sediment suspension and deposition across restored oyster reefs of varying orientation to flow: implications for restoration. Estuaries and Coasts 39, 1435–1448. https://doi.org/10.1007/s12237-016-0096-y

CORINE Land Cover [WWW Document], 2018. https://land.copernicus.eu/pan-european/corine-land-cover (accessed 12.14.21).

De Casabianca, M.-L., 1996. France — The Mediterranean lagoons, in: Schramm, W., Nienhuis, P.H. (Eds.), Marine benthic vegetation: Recent changes and the effects of eutrophication, Ecological Studies. Springer, Berlin, Heidelberg, pp. 307–329. https://doi.org/10.1007/978-3-642-61398-2_14

de Paz, L., Patrício, J., Marques, J.C., Borja, A., Laborda, A.J., 2008. Ecological status assessment in the lower Eo estuary (Spain). The challenge of habitat heterogeneity integration: a benthic perspective. Marine Pollution Bulletin 56, 1275–1283. https://doi.org/10.1016/j.marpolbul.2008.04.027

Derolez, V., Bec, B., Munaron, D., Fiandrino, A., Pete, R., Simier, M., Souchu, P., Laugier, T., Aliaume, C., Malet, N., 2019. Recovery trajectories following the reduction of urban nutrient inputs along the eutrophication gradient in French Mediterranean lagoons. Ocean & Coastal Management 171, 1–10. https://doi.org/10.1016/j.ocecoaman.2019.01.012

Derolez, V., Fiandrino, A., Munaron, D., Cadoret, M., 2014. Bilan sur les principales pressions pesant sur les lagunes méditerranéennes et leurs liens avec l'état DCE. RST-LER/LR 14-20. https://archimer.ifremer.fr/doc/00254/36574/

Derolez, V., Leurion, A., Fiandrino, A., 2012. Directive Cadre Eau. Estimation du degré d'échange avec la mer pour les lagunes du bassin Rhône Méditerranée &

Corse. Agence de l'eau, Ref. RST. DOP/LER-LR/12-10, 36p.
https://archimer.ifremer.fr/doc/00094/20558/

Derolez, V., Malet, N., Fiandrino, A., Lagarde, F., Richard, M., Ouisse, V., Bec, B.,
Aliaume, C., 2020a. Fifty years of ecological changes: Regime shifts and drivers in
a coastal Mediterranean lagoon during oligotrophication. Science of The Total
Environment 732, 139292. https://doi.org/10.1016/j.scitotenv.2020.139292

Derolez, V., Soudant, D., Malet, N., Chiantella, C., Richard, M., Abadie, E., Aliaume,
C., Bec, B., 2020b. Two decades of oligotrophication: Evidence for a phytoplankton
community shift in the coastal lagoon of Thau (Mediterranean Sea, France).
Estuarine, Coastal and Shelf Science 241, 106810.
https://doi.org/10.1016/j.ecss.2020.106810

Díaz, R., Rosenberg, R., 1995. Marine benthic hypoxia: A review of its ecological
effects and the behavioural responses of benthic macrofauna, in: Oceanography
and Marine Biology : An Annual Review. University College London Press, London,
UK, pp. 245–303.

Donadi, S., van der Zee, E.M., van der Heide, T., Weerman, E.J., Piersma, T., van de
Koppel, J., Olff, H., Bartelds, M., van Gerwen, I., Eriksson, B.K., 2014. The bivalve
loop: Intra-specific facilitation in burrowing cockles through habitat modification.
Journal of Experimental Marine Biology and Ecology 461, 44–52.
https://doi.org/10.1016/j.jembe.2014.07.019

Dufrêne, M., Legendre, P., 1997. Species assemblages and indicator species:the
need for a flexible asymmetrical approach. Ecological Monographs 67, 345–366.
https://doi.org/10.1890/0012-9615(1997)067[0345:SAAIST]2.0.CO;2

Elliott, M., Quintino, V., 2007. The estuarine quality paradox, environmental
homeostasis and the difficulty of detecting anthropogenic stress in naturally
stressed areas. Marine Pollution Bulletin 54, 640–645.
https://doi.org/10.1016/j.marpolbul.2007.02.003

European Commission: Joint Research Centre, Gifre, J., Reizopolou, S., Penna, M.,
Costas, N., Trabucco, B., Salas Herrero, F., Pardo, I., Derolez, V., Quintana, X.,
Buchet, R., Gascon, S., Boix, D., Martinoy, M., & Sala, J. (2018). Transitional waters
Mediterranean geographic intercalibration group : benthic invertebrates fauna
ecological assessment methods, Publications Office.
https://data.europa.eu/doi/10.2760/625400

Faulwetter, S., Markantonatou, V., Pavloudi, C., Papageorgiou, N., Keklikoglou, K.,
Chatzinikolaou, E., Pafilis, E., Chatzigeorgiou, G., Vasileiadou, K., Dailianis, T.,
Fanini, L., Koulouri, P., Arvanitidis, C., 2014. Polytraits: A database on biological
traits of marine polychaetes. Biodiversity Data Journal 2, e1024.
https://doi.org/10.3897/BDJ.2.e1024

Fiandrino, A., Ouisse, V., Dumas, F., Lagarde, F., Pete, R., Malet, N., Le Noc, S., de
Wit, R., 2017. Spatial patterns in coastal lagoons related to the hydrodynamics of
seawater intrusion. Marine Pollution Bulletin 119, 132–144.
https://doi.org/10.1016/j.marpolbul.2017.03.006

Gamito, S., Gilabert, J., Marcos Diego, C., Pérez-Ruzafa, A., 2005. Effects of changing
environmental conditions on lagoon ecology, in: Gonenc, I., Wolfin, J. (Eds.),
Coastal Lagoons – Ecosystem Processes and Modeling for Sustainable Use and
Development. CRC Press, pp. 193–229.

Giampaoletti, J., Sbrana, A., Magni, P., Gravina, M.F., 2023. Macrobenthos of the
Tortolì lagoon: a peculiar case of high benthic biodiversity among Mediterranean
lagoons. Diversity 15, 783. https://www.mdpi.com/1424-2818/15/6/783

González-Ortiz, V., Egea, L.G., Jiménez-Ramos, R., Moreno-Marín, F., Pérez-Lloréns, J.L., Bouma, T.J., Brun, F.G., 2014. Interactions between seagrass complexity, hydrodynamic flow and biomixing alter food availability for associated filter-feeding organisms. PLOS ONE 9, e104949. https://doi.org/10.1371/journal.pone.0104949

Guelorget, O., Michel, P., 1979. Les peuplements benthiques d'un étang littoral Languedocien, l'étang du Prévost (Hérault): 2. Étude quantitative de la macrofaune des sables. Téthys.

Guelorget, O., Perthuisot, J.P., 1983. Le domaine paralique : expressions géologiques, biologiques et économiques du confinement, Travaux du laboratoire de géologie, n°16. Presses de l'école normale supérieure, Paris.

Guelorget, O., Perthuisot, J., Lamy, N., Lefebvre, A., 1994. Structure et organisation de l'étang de Thau d'après la faune benthique (macrofaune, méiofaune). Relations avec le confinement. Oceanologica Acta 17, 105–114.

Harman, W.N., 1972. Benthic substrates: their effect on fresh-water mollusca. Ecology 53, 271–277. https://doi.org/10.2307/1934081

Hebbali, A., 2020. olsrr: tools for building ols regression models.

James, G., Witten, D., Hastie, T., and Tibshirani, R. (eds.). (2013). An introduction to statistical learning: with applications in R. New York: Springer.

Jessen, C., Bednarz, V., Rix, L., Teichberg, M., Wild, C., 2015. Marine Eutrophication, in: Environmental Indicators. p. 27. https://doi.org/10.1007/978-94-017-9499-2_11

Jones, A.G., Dubois, S.F., Desroy, N., Fournier, J., 2021. Intertidal ecosystem engineer species promote benthic-pelagic coupling and diversify trophic pathways. Marine Ecology Progress Series 660, 119–139. https://doi.org/10.3354/meps13600

Jones, A. G., Schaal, G., Boyé, A., Creemers, M., Derolez, V., Desroy, N., Fiandrino, A., Mouton, T., Simier, M., Smith, N., Ouisse, V. (2024). Disentangling the effects of eutrophication and natural variability on macrobenthic communities across French coastal lagoons (Version 2) [Data set]. Zenodo. https://doi.org/10.5281/zenodo.10825165

Josse, J., Husson, F., 2016. missMDA: a package for handling missing values in multivariate data analysis. Journal of Statistical Software 70, 1–31. https://doi.org/10.18637/jss.v070.i01

Josse, J., Husson, F., 2012. Handling missing values in exploratory multivariate data analysis methods. Journal de la Société Française de Statistique 153, 79–99.

Kassambara, A., 2020. rstatix: pipe-friendly framework for basic statistical tests.

Kermagoret, C., Claudet, J., Derolez, V., Nugues, M.M., Ouisse, V., Quillien, N., Baulaz, Y., Le Mao, P., Scemama, P., Vaschalde, D., Bailly, D., Mongruel, R., 2019. How does eutrophication impact bundles of ecosystem services in multiple coastal habitats using state-and-transition models. Ocean & Coastal Management 174, 144–153. https://doi.org/10.1016/j.ocecoaman.2019.03.028

Kjerfve, B., 1994. Chapter 1 Coastal Lagoons, in: Kjerfve, B. (Ed.), Elsevier Oceanography Series, Coastal Lagoon Processes. Elsevier, pp. 1–8. https://doi.org/10.1016/S0422-9894(08)70006-0

Lacoste, E., Jones, A., Callier, M., Klein, J., Lagarde, F., Derolez, V., 2023. A review of knowledge on the impacts of multiple anthropogenic pressures on the soft-bottom benthic ecosystem in Mediterranean coastal lagoons. Estuaries and Coasts. https://doi.org/10.1007/s12237-023-01188-9

Landman, M.J., Van Den Heuvel, M.R., Ling, N., 2005. Relative sensitivities of common freshwater fish and invertebrates to acute hypoxia. New Zealand Journal

of Marine and Freshwater Research 39, 1061–1067.
https://doi.org/10.1080/00288330.2005.9517375

Lasserre, P., 1979. Coastal lagoons: sanctuary ecosystems, cradles of culture,
targets for economic growth. Nature and Resources.

Le Fur, I., Wit, R.D., Plus, M., Oheix, J., Derolez, V., Simier, M., Malet, N., Ouisse, V.,
2019. Re-oligotrophication trajectories of macrophyte assemblages in
Mediterranean coastal lagoons based on 17-year time-series. Marine Ecology
Progress Series 608, 13–32. https://doi.org/10.3354/meps12814

Le Fur, I., De Wit, R., Plus, M., Oheix, J., Simier, M., Ouisse, V., 2018. Submerged
benthic macrophytes in Mediterranean lagoons: distribution patterns in relation to
water chemistry and depth. Hydrobiologia 808, 175–200.
https://doi.org/10.1007/s10750-017-3421-y

Legendre, P., Legendre, L., 2012. Numerical ecology, 3rd ed. Elsevier.

Lepoint, G., Nyssen, F., Gobert, S., Dauby, P., Bouquegneau, J.-M., 2000. Relative
impact of a seagrass bed and its adjacent epilithic algal community in consumer
diets. Marine Biology 136, 513–518. https://doi.org/10.1007/s002270050711

Leruste, A., Malet, N., Munaron, D., Derolez, V., Hatey, E., Collos, Y., De Wit, R., Bec,
B., 2016. First steps of ecological restoration in Mediterranean lagoons: Shifts in
phytoplankton communities. Estuarine, Coastal and Shelf Science 180, 190–203.
https://doi.org/10.1016/j.ecss.2016.06.029

Levin, L.A., Boesch, D.F., Covich, A., Dahm, C., Erséus, C., Ewel, K.C., Kneib, R.T.,
Moldenke, A., Palmer, M.A., Snelgrove, P., Strayer, D., Weslawski, J.M., 2001. The
function of marine critical transition zones and the importance of sediment
biodiversity. Ecosystems 4, 430–451. https://doi.org/10.1007/s10021-001-0021-4

Magalhães, L., Pires, A., Velez, C., Martins, R., Figueira, E., Soares, A.M.V.M., Freitas,
R., 2019. Seasonal and spatial alterations in macrofaunal communities and in
Nephtys cirrosa (Polychaeta) oxidative stress under a salinity gradient: A
comparative field monitoring approach. Ecological Indicators 96, 192–201.
https://doi.org/10.1016/j.ecolind.2018.08.045

Magni, P., Como, S., Kamijo, A., Montani, S., 2017. Effects of *Zostera marina* on the
patterns of spatial distribution of sediments and macrozoobenthos in the boreal
lagoon of Furen (Hokkaido, Japan). Marine Environmental Research 131, 90–102.
https://doi.org/10.1016/j.marenvres.2017.09.013

Magni, P., Gravina, M.F., 2023. Macrobenthos of lagoon ecosystems: a comparison in
vegetated and bare sediments. Advances in Oceanography and Limnology 14.
https://doi.org/10.4081/aiol.2023.11124

Magni, P., Semprucci, F., Gravina, M.F., 2022. Joint analysis of macrofaunal and
meiofaunal assemblages improves the assessment of lagoonal environmental
heterogeneity. Estuarine, Coastal and Shelf Science 266, 107740.
https://doi.org/10.1016/j.ecss.2021.107740

Mancinelli, G., Rossi, L., 2001. Indirect, size-dependent effects of crustacean
mesograzers on the Rhodophyta *Gracilaria verrucosa* (Hudson) Papenfuss:
evidence from a short-term study in the Lesina Lagoon (Italy). Marine Biology 138,
1163–1173. https://doi.org/10.1007/s002270100545

Martini, S., Larras, F., Boyé, A., Faure, E., Aberle, N., Archambault, P., Bacouillard, L.,
Beisner, B.E., Bittner, L., Castella, E., Danger, M., Gauthier, O., Karp-Boss, L.,
Lombard, F., Maps, F., Stemmann, L., Thiébaut, E., Usseglio-Polatera, P., Vogt, M.,
Laviale, M., Ayata, S.-D., 2021. Functional trait-based approaches as a common

framework for aquatic ecologists. Limnology and Oceanography 66, 965–994.
https://doi.org/10.1002/lno.11655
Menu, M., Sandrine, V., Bajjouk, T., Derolez, V., Fiandrino, A., Giraud, A., Grillas, P.,
Ouisse, V., 2019. Rapport final du projet CHAMILA (Cartographie des habitats en
milieu lagunaire méditerranéen) R.ODE/UL/LER/LR/19.34.
https://doi.org/10.13155/70545
Mistri, M., Borja, A., Aleffi, I.F., Lardicci, C., Tagliapietra, D., Munari, C., 2018.
Assessing the ecological status of Italian lagoons using a biomass-based index.
Marine Pollution Bulletin 126, 600–605.
https://doi.org/10.1016/j.marpolbul.2017.09.048
Miyamoto, Y., Nakano, T., Yamada, K., Hatakeyama, K., Hamaguchi, M., 2019.
Combined effects of drift macroalgal bloom and warming on occurrence and
intensity of diel-cycling hypoxia in a eutrophic coastal lagoon. Estuaries and
Coasts 42, 494–503. https://doi.org/10.1007/s12237-018-0484-6
Morris, E.K., Caruso, T., Buscot, F., Fischer, M., Hancock, C., Maier, T.S., Meiners, T.,
Müller, C., Obermaier, E., Prati, D., Socher, S.A., Sonnemann, I., Wäschke, N.,
Wubet, T., Wurst, S., Rillig, M.C., 2014. Choosing and using diversity indices:
insights for ecological applications from the German Biodiversity Exploratories.
Ecol Evol 4, 3514–3524. https://doi.org/10.1002/ece3.1155
Muxika, I., Borja, Á., Bald, J., 2007. Using historical data, expert judgement and
multivariate analysis in assessing reference conditions and benthic ecological
status, according to the European Water Framework Directive. Marine Pollution
Bulletin, Implementation of the Water Framework Directive in European marine
waters 55, 16–29. https://doi.org/10.1016/j.marpolbul.2006.05.025
Neveux, J., Lantoine, F., 1993. Spectrofluorometric assay of chlorophylls and
phaeopigments using the least squares approximation technique. Deep Sea
Research Part I: Oceanographic Research Papers 40, 1747–1765.
https://doi.org/10.1016/0967-0637(93)90030-7
Newton, A., Oliveira, P.S., Icely, J.D., Foster, P.A., 2010. Monitoring of oxygen
condition in the Ria Formosa coastal lagoon, Portugal. J Environ Monit 12, 355–
360. https://doi.org/10.1039/b914015h
Nordström, M., Booth, D.M., 2007. Drift algae reduce foraging efficiency of juvenile
flatfish. Journal of Sea Research 58, 335–341.
https://doi.org/10.1016/j.seares.2007.08.001
Norkko, J., Bonsdorff, E., Norkko, A., 2000. Drifting algal mats as an alternative
habitat for benthic invertebrates: Species specific responses to a transient
resource. Journal of Experimental Marine Biology and Ecology 248, 79–104.
https://doi.org/10.1016/S0022-0981(00)00155-6
Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin,
P.R., O'Hara, R.B., Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E.,
Wagner, H., 2019. vegan: Community Ecology Package.
Orro, C., Cabana, D., 2021. Seasonal implications of the benthic ecological status
assessment in a protected Mediterranean coastal lagoon: Stagno Longu of Posada
(Sardinia, Italy). Regional Studies in Marine Science 44.
https://doi.org/10.1016/J.RSMA.2021.101759
Ouisse, V., Riera, P., Migné, A., Leroux, C., Davoult, D., 2011. Freshwater seepages
and ephemeral macroalgae proliferation in an intertidal bay: I Effect on benthic
community structure and food web. Estuarine, Coastal and Shelf Science 91, 272–
281. https://doi.org/10.1016/j.ecss.2010.10.034

Pasqualini, V., Derolez, V., Garrido, M., Orsoni, V., Baldi, Y., Etourneau, S., Leoni, V., Rébillout, P., Laugier, T., Souchu, P., Malet, N., 2017. Spatiotemporal dynamics of submerged macrophyte status and watershed exploitation in a Mediterranean coastal lagoon: Understanding critical factors in ecosystem degradation and restoration. Ecological Engineering 102, 1–14. https://doi.org/10.1016/j.ecoleng.2017.01.027

Paul, J., Nestlerode, J.A., Jarvis, B.M., 2023. Evaluating the effectiveness of M-AMBI with other biotic indexes in a temperate estuary. Marine Pollution Bulletin, 193: 115194. https://doi.org/10.1016/j.marpolbul.2023.115194

Pearson, T.H., Rosenberg, R., 1978. Macrobenthic succession in relation to organic enrichment and pollution of the marine environment. Oceanography and Marine Biology: An Annual Review 16, 229–311. https://doi.org/10.2983/035.034.0121u1.10

Pelletier, M.C., Charpentier, M., 2023. Assessing relative importance of stressors to the benthic index, M-AMBI: An example from U.S. estuaries. Marine Pollution Bulletin, 186:114456. https://doi.org/10.1016/j.marpolbul.2022.114456

Perez-Ruzafa, A., Marcos-Diego, C., 1992. Colonization rates and dispersal as essential parameters in the confinement theory to explain the structure and horizontal zonation of lagoon benthic assemblages. Technical Report Rapport Commission International Mer Medit 33.

Pérez-Ruzafa, A., Marcos, C., Pérez-Ruzafa, I.M., 2011b. Recent advances in coastal lagoons ecology: evolving old ideas and assumptions. Transitional Waters Bulletin 5, 50–74. https://doi.org/10.1285/i1825229Xv5n1p50

Pérez-Ruzafa, A., Marcos, C., Pérez-Ruzafa, I.M., 2011a. Mediterranean coastal lagoons in an ecosystem and aquatic resources management context. Physics and Chemistry of the Earth, Parts A/B/C, Water in the Mediterranean Basin 36, 160–166. https://doi.org/10.1016/j.pce.2010.04.013

Pérez-Ruzafa, A., Marcos, C., Pérez-Ruzafa, I.M., Barcala, E., Hegazi, M.I., Quispe, J., 2007a. Detecting changes resulting from human pressure in a naturally quick-changing and heterogeneous environment: Spatial and temporal scales of variability in coastal lagoons. Estuarine, Coastal and Shelf Science, Biodiversity and Ecosystem Functioning in Coastal and Transitional Waters 75, 175–188. https://doi.org/10.1016/j.ecss.2007.04.030

Pérez-Ruzafa, A., Marcos, C., Pérez-Ruzafa, I.M., Pérez-Marcos, M., 2013. Are coastal lagoons physically or biologically controlled ecosystems? Revisiting r vs. K strategies in coastal lagoons and estuaries. Estuarine, Coastal and Shelf Science, Estuarine and lagoon biodiversity and their natural goods and services 132, 17–33. https://doi.org/10.1016/j.ecss.2012.04.011

Pérez-Ruzafa, Angel, Marcos, C., Pérez-Ruzafa, I.M., Pérez-Marcos, M., 2011c. Coastal lagoons: "transitional ecosystems" between transitional and coastal waters. Journal of Coastal Conservation 15, 369–392. https://doi.org/10.1007/s11852-010-0095-2

Pérez-Ruzafa, Angel, Mompeán, M.C., Marcos, C., 2007b. Hydrographic, geomorphologic and fish assemblage relationships in coastal lagoons. Hydrobiologia 577, 107–125. https://doi.org/10.1007/s10750-006-0421-8

Perthuisot, J.-P., Guelorget, O., 1983. Le confinement, paramètre essentiel de la dynamique biologique du domaine paralique. Sciences Géologiques, bulletins et mémoires 36, 239–248. https://doi.org/10.3406/sgeol.1983.1642

Pitacco, V., Reizopoulou, S., Sfriso, Adriano, Sfriso, Andrea, Mistri, M., Munari, C., 2019. The difficulty of disentangling natural from anthropogenic forcing factors makes the evaluation of ecological quality problematic: A case study from Adriatic lagoons. Marine Environmental Research 150, 104756. https://doi.org/10.1016/j.marenvres.2019.104756

Pollice, A., Arima, S., Lasinio G.J., Basset, A., Rosati, I., 2015. Bayesian analysis of three indices for lagoons ecological status evaluation. Stochastic Environmental Research and Risk Assessment, 29: 477-485. DOI: 10.1007/s00477-014-0885-4

Provost, C., Gevrey, M., Argillier, C., 2012. La macrofaune benthique indicatrice de l'état des lagunes oligohalines méditerranéennes ? [Rapport de recherche] IRSTEA. 2012, pp.37. https://hal.science/hal-02597249/

Puente, A., Diaz, R.J., 2008. Is it possible to assess the ecological status of highly stressed natural estuarine environments using macroinvertebrates indices? Marine Pollution Bulletin 56, 1880–1889. https://doi.org/10.1016/j.marpolbul.2008.07.016

Quillien, N., Nordström, M.C., Gauthier, O., Bonsdorff, E., Paulet, Y.-M., Grall, J., 2015. Effects of macroalgal accumulations on the variability in zoobenthos of high-energy macrotidal sandy beaches. Marine Ecology Progress Series 522, 97–114. https://doi.org/10.3354/meps11151

Quillien, N., Nordström, M.C., Schaal, G., Bonsdorff, E., Grall, J., 2016. Opportunistic basal resource simplifies food web structure and functioning of a highly dynamic marine environment. Journal of Experimental Marine Biology and Ecology 477, 92–102. https://doi.org/10.1016/j.jembe.2016.01.010

R Core Team, 2020. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Reizopoulou, S., Nicolaidou, A., 2007. Index of size distribution (ISD): a method of quality assessment for coastal lagoons. Hydrobiologia 577, 141–149. https://doi.org/10.1007/s10750-006-0423-6

Reizopoulou, S., Simboura, N., Sigala, K., Barbone, E., Aleffi, F., Kaisakis, G., Rosati, I., Basset, A., Nicolaidou, A., 2014. Assessing the ecological status of Mediterranean coastal lagoons using macroinvertebrates. Comparison of the most commonly used methods. Mediterranean Marine Science 15, 602–612. https://doi.org/10.12681/mms.606

Riedel, B., Zuschin, M., Stachowitsch, M., 2012. Tolerance of benthic macrofauna to hypoxia and anoxia in shallow coastal seas: a realistic scenario. Marine Ecology Progress Series 458, 39–52. https://doi.org/10.3354/meps09724

Rigaud, S., Deflandre, B., Grenz, C., Cesbron, F., Pozzato, L., Voltz, B., Grémare, A., Romero-Ramirez, A., Mirleau, P., Meulé, S., Faure, V., Mayot, N., Michotey, V., Bonin, P., Pascal, L., Cordier, M.-A., Lamarque, B., Tenório, M., Radakovitch, O., 2021. Benthic oxygen dynamics and implication for the maintenance of chronic hypoxia and ecosystem degradation in the Berre lagoon (France). Estuarine, Coastal and Shelf Science 258, 107437. https://doi.org/10.1016/j.ecss.2021.107437

Roberts, D.W., 2019. labdsv: ordination and multivariate analysis for ecology.

Robertson, B.P., Savage, C., Gardner, J.P.A., Robertson, B.M., Stevens, L.M., 2016. Optimising a widely-used coastal health index through quantitative ecological group classifications and associated thresholds. Ecological Indicators 69, 595–605. https://doi.org/10.1016/j.ecolind.2016.04.003

Roques, C., Grousset, E., Troussellier, M., Hermet, S., Le Carrer, J., Sar, C., Caro, A., 2020. A trade-off between mucocytes and bacteriocytes in *Loripes orbiculatus* gills (Bivalvia, Lucinidae): a mixotrophic adaptation to seasonality and reproductive status in a symbiotic species? Marine Biology 167. https://doi.org/10.1007/s00227-020-03768-w

Rose, J.M., Caron, D.A., 2007. Does low temperature constrain the growth rates of heterotrophic protists? Evidence and implications for algal blooms in cold waters. Limnology and Oceanography 52, 886–895. https://doi.org/10.4319/lo.2007.52.2.0886

Rosenberg, R., Loo, L.-O., Möller, P., 1992. Hypoxia, salinity and temperature as structuring factors for marine benthic communities in a eutrophic area. Netherlands Journal of Sea Research 30, 121–129. https://doi.org/10.1016/0077-7579(92)90051-F

Rossi, F., Castelli, A., Lardicci, C., 2006. Distribution of macrobenthic assemblages along a marine gradient in Mediterranean eutrophic coastal lagoons. Marine ecology 27, 66–75. https://doi.org/10.1111/j.1439-0485.2006.00078.x

Sabetta, L., Barbone, E., Giardino, A., Galuppo, N., Basset, A., 2007. Species–area patterns of benthic macro-invertebrates in Italian lagoons. Hydrobiologia 577, 127–139. https://doi.org/10.1007/s10750-006-0422-7

Scaps, P., 2002. A review of the biology, ecology and potential use of the common ragworm *Hediste diversicolor* (O.F. Müller) (Annelida: Polychaeta). Hydrobiologia 470, 203–218. https://doi.org/10.1023/A:1015681605656

Schielzeth, H., 2010. Simple means to improve the interpretability of regression coefficients. Methods in Ecology and Evolution 1, 103–113. https://doi.org/10.1111/j.2041-210X.2010.00012.x

Sigala, K., Reizopoulou, S., Basset, A., Nicolaidou, A., 2012. Functional diversity in three Mediterranean transitional water ecosystems. https://doi.org/10.1016/J.ECSS.2012.06.002

Sigovini, M., Keppel, E., Tagliapietra, D., 2013. M-AMBI revisited: looking inside a widely-used benthic index. Hydrobiologia 717, 41–50. https://doi.org/10.1007/s10750-013-1565-y

Souchu, P., Bec, B., Smith, V.H., Laugier, T., Fiandrino, A., Benau, L., Orsoni, V., Collos, Y., Vaquer, A., 2010. Patterns in nutrient limitation and chlorophyll a along an anthropogenic eutrophication gradient in French Mediterranean coastal lagoons. Can. J. Fish. Aquat. Sci. 67, 743–753. https://doi.org/10.1139/F10-018

Souchu, P., Gasc, A., Collos, Y., Vaquer, A., Tournier, H., Bibent, B., Deslous-Paoli, J.-M., 1998. Biogeochemical aspects of bottom anoxia in a Mediterranean lagoon (Thau, France). Marine Ecology-progress Series 164, 135–146. https://doi.org/10.3354/meps164135

Stendera, S., Johnson, R.K., 2008. Habitat-specific stability and persistence of benthic invertebrate communities in boreal lakes. https://doi.org/10.1127/1863-9135/2008/0171-0311

Tagliapietra, D., Sigovini, M., Ghirardini, A., 2009. A review of terms and definitions to categorise estuaries, lagoons and associated environments. https://doi.org/10.1071/MF08088

Tagliapietra, D., Sigovini, M., Magni, P., 2012. Saprobity: a unified view of benthic succession models for coastal lagoons. Hydrobiologia 686, 15–28. https://doi.org/10.1007/s10750-012-1001-8

Tournoud, M.-G., Payraudeau, S., Cernesson, F., Salles, C., 2006. Origins and
quantification of nitrogen inputs into a coastal lagoon: Application to the Thau
lagoon (France). Ecological Modelling, Special Issue on Southern European Coastal
Lagoons 193, 19–33. https://doi.org/10.1016/j.ecolmodel.2005.07.038

Trombetta, T., Vidussi, F., Mas, S., Parin, D., Simier, M., Mostajir, B., 2019. Water
temperature drives phytoplankton blooms in coastal waters. PLOS ONE 14,
e0214933. https://doi.org/10.1371/journal.pone.0214933

Vaquer-Sunyer, R., Duarte, C.M., 2011. Temperature effects on oxygen thresholds for
hypoxia in marine benthic organisms. Global Change Biology 17, 1788–1797.
https://doi.org/10.1111/j.1365-2486.2010.02343.x

Venables, W.N., Ripley, B.D., 2002. Modern Applied Statistics with S, Fourth. ed.
Springer, New York.

Vergara-Chen, C., González-Wangüemert, M., Marcos, C., Pérez-Ruzafa, Á., 2013.
Small-scale genetic structure of *Cerastoderma glaucum* in a lagoonal
environment: potential significance of habitat discontinuity and unstable
population dynamics. Journal of Molluscan Studies 79, 230–240.
https://doi.org/10.1093/mollus/eyt015

Vernberg, F.J., 1982. Environmental adaptation to lagoon systems. Oceanologica
Acta, Special issue.

Vignes, F., Barbone, E., Breber, P., D'Adamo, R., Leonilde, R., Ungaro, N., Focardi, S.,
Renzi, M., Basset, A., 2010. Spatial and temporal description of the dystrophic
crisis in Lesina lagoon during summer 2008. Transitional Waters Bulletin 3, 47–62.
https://doi.org/10.1285/i1825229Xv3n2p47

Vizzini, S., Mazzola, A., 2006. Sources and transfer of organic matter in food webs of
a Mediterranean coastal environment: Evidence for spatial variability. Estuarine,
Coastal and Shelf Science 66, 459–467. https://doi.org/10.1016/j.ecss.2005.10.004

Ware, C., Dijkstra, J.A., Mello, K., Stevens, A., O'Brien, B., Ikedo, W., 2019. A novel
three-dimensional analysis of functional architecture that describes the properties
of macroalgae as a refuge. Marine Ecology Progress Series 608, 93–103.
https://doi.org/10.3354/meps12800

Wei, T., Simko, V., 2017. corrplot: visualization of a correlation matrix.

Weissberger, E.J., Coiro, L.L., Davey, E.W., 2009. Effects of hypoxia on animal burrow
construction and consequent effects on sediment redox profiles. Journal of
Experimental Marine Biology and Ecology 371, 60–67.
https://doi.org/10.1016/j.jembe.2009.01.005

Wilke, M., Boutiere, H., 2000. Hydrobiological, physical and chemical characteristics
and spatio-temporal dynamics of an oligotrophic mediterranean lagoon: the étang
de la Palme (France). Vie et Milieu / Life & Environment, 101-115

Zaldívar, J., Viaroli, P., Newton, A., Wit, R., Ibañez, C., Reizopoulou, S., Somma, F.,
Razinkovas, A., Basset, A., Holmer, M., Murray, N., 2008. Eutrophication in
transitional waters: an overview. https://doi.org/10.1285/I18252273V2N1P1

Zilius, M., Giordani, G., Petkuviene, J., Lubiene, I., Ruginis, T., Bartoli, M., 2015.
Phosphorus mobility under short-term anoxic conditions in two shallow eutrophic
coastal systems (Curonian and Sacca di Goro lagoons). Estuarine, Coastal and
Shelf Science 164, 134–146. https://doi.org/10.1016/j.ecss.2015.07.004

1875