1
2

Disentangling the effects of eutrophication and natural variability on macrobenthic communities across French coastal lagoons

Auriane G. Jones*[1], Gauthier Schaal[2], Aurelien Boye[3], Marie Creemers[1], Valerie Derolez[1], Nicolas Desroy[4], Annie Fiandrino[1], Theophile L. Mouton[1,5], Monique Simier[1], Niamh Smith[1,6], Vincent Ouisse[1]

[1] MARBEC, Univ. Montpellier, CNRS, Ifremer, IRD, Sète, France
[2] Univ Brest, CNRS, IRD, Ifremer, LEMAR, 29280 Plouzané, France
[3] IFREMER, Centre de Bretagne, DYNECO, Laboratoire d'Ecologie Benthique Côtière (LEBCO), 29280 Plouzané, France
[4] IFREMER, Laboratoire Environnement et Ressources Bretagne nord, 35801 Dinard cedex, France
[5] FRB-CESAB, Institut Buisson Bertrand, 5 Rue de l'Ecole de Médecine, 34000 Montpellier, France
[6] IFREMER, Centre Atlantique, Département Ecologie et Modelés pour l'Halieutique (EMH), Rue de l'Ile d'Yeu, BP 21105, 44311 Nantes cedex 3, France

*Corresponding author
Correspondence: jones.ecology@gmail.com

ABSTRACT

Coastal lagoons are transitional ecosystems that host a unique diversity of species and support many ecosystem services. Owing to their position at the interface between land and sea, they are also subject to increasing human impacts, which alter their ecological functioning. Because coastal lagoons are naturally highly variable in their environmental conditions, disentangling the effects of anthropogenic disturbances like eutrophication from those of natural variability is a challenging, yet necessary issue to address. Here, we analyze a dataset composed of macrobenthic invertebrate abundances and environmental variables (hydro-morphology, water, sediment and macrophytes) gathered across 29 Mediterranean coastal lagoons located in France, to characterize the main drivers of community composition and structure. Using correlograms, linear models and variance partitioning, we found that lagoon hydro-morphology (connection to the sea and lagoon surface), which affects the level of environmental variability (salinity and temperature), as well as lagoon-scale benthic habitat diversity (using macrophyte morphotypes) seemed to regulate macrofauna distribution, while eutrophication and associated stressors like low dissolved oxygen, acted upon the existing communities, mainly by reducing species richness and diversity. Furthermore, M-AMBI, a multivariate index composed of species richness, Shannon diversity and AMBI (AZTI's Marine Biotic Index) and currently used to evaluate the ecological state of French coastal lagoons, was more sensitive to eutrophication (18%) than to natural variability (9%), with nonetheless 49% of its variability explained jointly

by both. Conversely, AMBI was the only tested index that uniquely responded to
eutrophication variables, which nonetheless explained less than a third of its variability. To
improve the robustness of benthic indicators like M-AMBI and increase the effectiveness of
lagoon benthic habitat management, we call for a revision of the ecological groups at the base
of the AMBI index and of the current lagoon typology which could be inspired by the lagoon-
sea connection levels used in this study.
50

51 *Keywords:* Mediterranean Sea, macrofauna, taxonomic diversity, M-AMBI, macrophyte morphotypes,
52 variance partitioning
53

 Introduction

Coastal lagoons are transitional ecosystems between land and sea that represent 13% of the world's
coastline (Barnes, 1980) and provide essential ecosystem services, especially through nutrient regulation and
sequestration as well as climate regulation (Kermagoret et al., 2019; Levin et al., 2001). These ecosystems are
characterized by strong salinity variations at different temporal scales, ranging from fortnights to seasons
and decades (Pérez-Ruzafa et al., 2007a), that are mainly governed by short-term freshwater inflow, seasonal
temperature variations, the degree of connection of each water mass with the sea and decadal climatic
variations (rainfall and temperatures) (Barbone and Basset, 2010; De Casabianca, 1996; Zaldívar et al., 2008).
Oxygen saturation naturally fluctuates in lagoons, due to water temperature and re-aeration processes
controlled by wind conditions (Chapelle et al., 2001; Derolez et al., 2020a). Oxygen dynamics are further
affected by primary production, which is mainly governed, in oligo and mesotrophic lagoons by seasonal
benthic macrophyte growth and decay cycles, and in hyper and eutrophic lagoons by pelagic phytoplankton
dynamics (Zaldívar et al., 2008). Overall, coastal lagoons are naturally highly dynamic systems in terms of
environmental factors, representing a form of natural environmental stress for any organism living in these
systems like fish and benthic invertebrates (Pérez-Ruzafa et al., 2007b; Zaldívar et al., 2008). Consequently,
lagoonal, just like estuarine benthic communities, generally present low species richness and high
abundances of tolerant species, characteristics typical of disturbed or stressed coastal ecosystems, giving rise
to the "transitional waters quality paradox" (Zaldívar et al., 2008), a generalization of the "Estuarine quality
paradox" (Elliott and Quintino, 2007).

Coastal lagoons have also been intensely used and modified by humans for food (fishing and
aquaculture), transportation (canals, ports and inlet modification), livelihood (land reclamation for
construction) and recreation (fishing, sailing, swimming) (Lasserre, 1979; Pérez-Ruzafa et al., 2011a). The
position of coastal lagoons at the interface between land and sea further exposes them to human land-
derived pollution such as nitrogen, phosphorous, pesticide, or heavy metal pollution (Caliceti et al., 2002;
Derolez et al., 2019; Souchu et al., 2010). The oldest and most studied is nitrogen and phosphorous pollution
linked to agriculture and urban development (Bendoricchio et al., 1993; Kjerfve, 1994; Lacoste et al., 2023;
Tournoud et al., 2006), which can lead to eutrophication of coastal lagoons, a phenomenon associated to an
increase in phytoplankton biomass (Cloern, 2001; Zaldívar et al., 2008), a replacement of long-lived
submerged macrophytes like marine plants, by emergent free-living benthic macroalgae and even to the
disappearance of all benthic macrophytes (Le Fur et al., 2019, 2018). The high light availability linked to
shallow depths and the naturally high nutrient concentration of water and sediments (*i.e.* coastal lagoons can
be considered as "naturally eutrophic" (De Casabianca, 1996; Zaldívar et al., 2008)) promote the
development of structurally complex benthic macrophytes, which are characteristic features of coastal
lagoons. Macrophyte presence, cover and composition create a highly heterogeneous aquatic landscape
(Arocena, 2007; Gamito et al., 2005), which is also determined by the sediment nature (*i.e.* hard bottom,
gravel, sand, mud) and organic matter content (Menu et al., 2019).

Benthic macroinvertebrates are a key biotic compartment of coastal lagoons, through their role in water
filtration, food web dynamics, recycling of organic matter and nutrients and sediment bioturbation (Barnes,
1980; Gamito et al., 2005; Levin et al., 2001; Zaldívar et al., 2008). These benthic communities are reported
to be shaped mainly by natural environmental variability (*i.e.* environmental filtering) and passive diffusion
(mainly as planktonic larvae) (Basset et al., 2007). At the intra-lagoon scale, the spatial distribution of
macrobenthic invertebrates is tightly linked to the renewal of marine-originating molecules, which is directly
linked to the connection level with the open sea, a result used to define the notion of confinement
(Guelorget and Perthuisot, 1983). Superimposed onto the previously mentioned "natural environmental
stress" (*i.e.* strong fluctuations of abiotic parameters like salinity and temperature) are a number of human
related disturbances like anthropogenic eutrophication, which can further impact lagoonal organisms (Elliott
and Quintino, 2007; Rossi et al., 2006). Most benthic macroinvertebrates are exposed to multiple human-
induced disturbances as they generally live for several years. Furthermore, these organisms tend to present,
as adults, a low dispersal potential (*i.e.* below a few kilometers), meaning that faced with a disturbance, each
individual will either resist, become accustomed or die, with overall very little chance of escaping the
disturbance, especially if it affects the entire lagoon (Cognetti and Maltagliati, 2000). These characteristics
make benthic macroinvertebrates good theoretical indicators of the ecological health status of coastal
lagoons and since 2000, this biological compartment is officially included in the assessment of European

transitional water bodies carried out within the Water Framework Directive (WFD) (Reizopolou et al., 2018).
However, although they are in theory good ecological indicators, benthic macroinvertebrates are also
influenced by "natural environmental stress", hence it appears crucial to disentangle the effect of natural
environmental variability from the effect of variability linked to human disturbances on this compartment in
hyper-variable ecosystems like coastal lagoons (Pitacco et al., 2019). Most of the published work on
Mediterranean coastal lagoon macroinvertebrates have either been descriptive studies focusing on one or a
few lagoons (Amanieu et al., 1977; Arocena, 2007; Bachelet et al., 2000; Carlier et al., 2007; Guelorget et al.,
1994; Guelorget and Michel, 1979; Magni et al., 2022; Pérez-Ruzafa et al., 2007a; Rossi et al., 2006),
theoretical oriented studies testing ecological hypotheses like environmental filtering (Basset et al., 2007) or
species-area relations (Sabetta et al., 2007) or very applied studies related to the WFD and focusing on
lagoon typology definition, indicator development or indicator reference values (Barbone et al., 2012; Basset
et al., 2012; Reizopoulou et al., 2014). Disentangling the respective effect of natural vs human-induced
disturbances on lagoon communities requires a statistical power which can be reached by using extensive
datasets, spanning a variety of ecological conditions (*i.e.* several lagoon systems) and a thorough
characterization of both sources of disturbance. In this study, we make use of the full potential of large-scale
environmental surveys of the various compartments of French Mediterranean coastal lagoons (water
column, sediments, macrophytes and benthic macroinvertebrates): *(i)* to disentangle the effects of an
anthropogenic disturbance, here eutrophication, from those of natural environmental variability on benthic
macroinvertebrates and *(ii)* to understand the links between the different environmental variables
characterizing coastal lagoons, that affect benthic macroinvertebrates. This study, based on a dataset
including 29 French Mediterranean coastal lagoons (ranging from oligo- to eu-haline, with diverse
environmental and anthropic settings), focuses on multiple levels of biological complexity; first the broad and
complex community level, then simple diversity indices like species richness and Shannon diversity index and
finally, a multimetric and WFD-compliant index.
131

Methods

Study sites

The French Mediterranean coast features a great diversity of lagoons characterized by different hydro-
climatic contexts (*e.g.* temperature, wind, rainfall). Some of these lagoons are under the influence of large
watersheds (up to 1 500 km²) while others are connected to more restricted basins that may be 300 times
smaller (*ca.* 5 km²). These lagoons also present variable connections to the sea, with both natural vs man-
made channels, and continuous vs temporary channels. Human land-use in this region is particularly high,
associated to both urban and agricultural development (Derolez et al., 2019). Most lagoons in continental
France are located near large cities like Narbonne or Montpellier and/or in areas intensively cultivated for
wine, vegetables and cereals (CORINE Land Cover, 2018) whereas the Corsican lagoons of Palo, Urbino and
Diana are relatively preserved from direct urban pressures. Differently, the Corsican lagoon of Biguglia is
located just south of the urban hub of Bastia and in a growing agricultural plain (CORINE Land Cover, 2018;
Pasqualini et al., 2017). Overall, French Mediterranean lagoons face various degrees of anthropogenic
pressures.
A total of 29 nanotidal (tide < 0.5 m) coastal lagoons located in the Gulf of Lion (north-west
Mediterranean Sea) and along the east coast of Corsica were selected for this study and sampled across 41
stations (between one and three in each lagoon) (Figure 1). The sampling was carried out in 2006 or 2009
during the "Réseau de Suivi Lagunaire" and WFD campaigns. Infaunal benthic macroinvertebrates were
sampled in May, except four lagoons sampled at the end of April (La Marette, Ponant, Berre and Vaine) and
one lagoon at the end of June (Bolmon). The studied lagoons encompass a wide range of hydro-
morphological characteristics in terms of surface, depth and connection to the sea (Table 1) with the lagoon-
sea connection level expressed as weak, moderate or strong depending on the differences in salinity
variations between the lagoon and a neighboring marine station (Derolez et al., 2012). Lagoon surfaces vary
from 1 km² (Campignol) to 133 km² (Berre) (Menu et al., 2019) and station depth is comprised between 0.1
m (La Palme south station) and 9.5 m (Urbino) with most stations less than 3 m deep (Table 1).
157
158

159
160

161
Figure 1 - Map of the 41 stations sampled in the 29 lagoons located along the French Mediterranean coast (Gulf of Lion and Corsica). Each station's salinity type and
group membership (K-means partitioning) based on the benthic macrofauna Hellinger-transformed mean abundances are indicated using a symbol (triangle: oligo- and
meso-haline and circle: poly- and eu-haline) and a color (red: group A and blue: group B), respectively (see section *Macrobenthic community structure* for the
explanation on this analysis). See Table 1 for the full lagoon names and corresponding stations.

The various hydro-morphological characteristics of the studied lagoons lead to a great diversity of
salinity regimes with 8 oligo- or meso-haline lagoons and 21 poly- or eu-haline lagoons (Figure 1) and to
large inter-lagoon discrepancies in terms of annual variability of temperature, salinity and oxygen
saturation. Overall, the physico-chemical status of the studied lagoons between 2004 and 2009 was
evaluated according to the WFD methodology, as poor (15 lagoons), bad (3 lagoons), moderate (4
lagoons) or good (7 lagoons) (Table 1). This evaluation takes into account the concentrations of total and
dissolved inorganic forms of nutrients, turbidity and oxygen saturation during the summer months of the
2004-2009 period and is conducted according to the WFD quality grid (Andral and Sargian, 2010a,
2010b).

Table 1 - Hydro-morphological characteristics (surface, depth and connection level to the sea) and
water physico-chemical status of the 29 lagoons as evaluated by the water framework directive
(green = good, yellow = moderate, orange = bad, red = poor). The physico-chemical status is based
on concentrations of total and dissolved inorganic forms of nutrients, turbidity and oxygen
saturation during the summer months of the 2004-2009 period (Andral and Sargian, 2010a; Andral
and Sargian, 2010b). Depth, sediment organic matter content (OMC, % of dry sediment weight),
beta diversity of macrophyte morphotypes (β_{MM}, no unit – see section *Estimating the macrophyte*
morphotype beta diversity) of the 41 stations sampled for benthic macrofauna either in 2006 (#) or
in 2009 (*) is also indicated. For stations sampled in the same lagoon, each one is labeled using a 3
letter code with the last letter indicating the geographical location of the station in the lagoon
(North (N), Middle (M), South (S), East (E), West (W), except THP which indicates the station's local
name Pisse-Saumes). The lagoons are listed from west to east along the Gulf of Lion (see Figure 1
for their geographical location).
188

Lagoon characteristics				Benthic station characteristics			
Lagoon	Surface (km^2)	Lagoon-sea connection	Physico-chemical status	Station code	Station depth (m)	OMC (%)	β_{MM}
Canet*	6.43	weak	red	CAN	0.9	5.6	0.13
Leucate*	52.88	strong	green	LEN	1.9	8.2	0.71
				LES	3.3	7.2	0.67
La Palme#	6.94	moderate	green	LPN	0.6	4.68	0.6
				LPS	0.1	1.61	0.53
Bages-Sigean*	37.05	moderate	green	BGN	1.9	11.5	0.65
				BGM	2.2	5.9	0.8
				BGS	1.3	7.1	0.53
Campignol*	1	weak	red	CAM	0.8	3.8	1
Gruissan*	1.31	strong	green	GRU	0.9	3.4	0.61
Vendres*	6.6	weak	red	VEN	0.8	22.8	0.6
Grand Bagnas#	1.69	weak	orange	GBA	0.5	5.38	0
Thau#	67.7	strong	green	THE	8.2	11.38	0.55
				THW	4	9.9	0.68
				THP	3.8	5.64	0.68
Ingril North#	2.28	strong	red	INN	1.1	12.48	0.4
Ingril South#	3.17	strong	red	INS	1.2	7.01	0.25
Vic#	12.96	moderate	red	VIC	1.5	13.05	0.69
Pierre-Blanche#	3.14	moderate	red	PBE	0.7	1.04	0.33
				PBW	0.7	9.93	0.67
Arnel#	4.42	moderate	red	ARN	0.8	7.88	0.29
Prévost#	2.46	strong	red	PRE	0.7	9.25	0.4
				PRW	0.7	8.4	0.6

Méjean#	7.12	moderate		MEE	1	7.67	0
				MEW	0.9	12.17	0
Grec#	1.08	moderate		GRE	0.7	8.49	0.33
Or (Mauguio)*	31.31	weak		ORE	1.4	10.9	0.75
				ORW	1.3	12.5	0.54
Ponant*	1.92	moderate		PON	3	10.9	0.11
La Marette*	1.3	weak		MRN	0.7	15.9	0
				MRS	0.6	14.2	0
Vaccares*	65.9	weak		VAC	1.9	5.8	0.23
La Palissade*	2.95	weak		PLS	0.6	1.5	0.6
Berre*	132.71	strong		BRN	5.9	11.8	0.39
				BRS	8.9	14.5	0.38
Vaine*	22.77	moderate		VAI	5.1	12.7	0.73
Bolmon*	6.01	weak		BOL	1.4	14.8	0.30
Biguglia*	13.75	weak		BIG	1	7.9	0.25
Diana*	5.4	strong		DIA	5.8	10.5	0.39
Urbino*	7.58	moderate		URB	9.5	16.3	0.63
Palo*	1.08	weak		PAL	1	12.5	0

Sampling

Benthic macroinvertebrates

At each of the 41 stations, three replicates (*ca.* 10 m apart) of unvegetated sediment were sampled.
Each replicate was made up of four pooled Ekman-Birge grabs, representing a total surface of 0.09 m^2 per
replicate and 0.27 m^2 per station. Samples were sieved on a 1 mm mesh, fixed in a 5 to 7%
formaldehyde/seawater solution and stained with Rose Bengal for latter identification. The sediments
were then sorted in the laboratory and all organisms identified if possible to the species level, following
the nomenclature of the European Register of Marine Species (ERMS). Taxa are indicated using WORMS
reference (https://www.marinespecies.org/aphia.php?p=match). Prior to the data analyses, we removed
meiofauna taxa (Ostracoda, Nematoda, Harpacticoida and Collembola) and organisms present in only one
lagoon and determined at the order (Decapoda, Coleoptera, Nudibranchia), class (Insecta, Ascidiacea) or
subclass (Oligochaeta) level. We kept Actiniaria, Nemertea and Polycladida as they were identified across
multiple lagoons.

Sediment parameters

At each station, an additional grab sample was collected to evaluate sediment organic matter and
grain-size distribution. Sediment samples were sieved on a 2 mm mesh to remove large macrophytes
debris, and the filtrate was weighted and dried until stable weight (110°C for 24 to 48h) in the laboratory.
Organic matter content (OMC, % of dry sediment weight) was calculated as the loss of weight after
combustion for 12h at 450°C of the dry sediment fraction below 2 mm. Because of technical problems, no
OMC was available for the stations ARN, LPN, LPS, PRW and VIC in 2006. To estimate these missing
values, we used OMC values measured in spring of 2006 (ARN, PRW, VIC) or spring of 2007 (LPN and LPS)
for the WFD sediment quality evaluation. For ARN, LPS, LPN and VIC, we estimated OMC by averaging
over the values measured in the three or four stations closest to the macrofauna station (distance
between 285 m and 707 m). For PRW, we used the OMC value measured at the closest station as this
station was located only 114 m from the macrofauna station (Supplementary material 1 and 2). Since
laboratory protocols changed between years (2006 and 2009) and between lagoons regarding the
sediment particle sizes considered for the grain-size analysis, we chose not to consider grain-size
parameters.

Water physico-chemical properties

Water column physico-chemical properties of the 29 lagoons are measured monthly during summer (from June to August or September) at least since 2003, following a standardized protocol implemented by the WFD (Derolez et al., 2019). Salinity (Sal), temperature (Temp) and oxygen saturation (O_2 sat) were measured using a portable field probe (WTW LF 197, WTW-GmbH, Weilheim, Germany) and turbidity (Turb) was measured in the laboratory using an optical turbidimeter (2100N IS turbidimeter ISO 7027). Chlorophyll *a* concentration ([Chla]) was measured in 50 mL samples of water, first filtered on a Whatman GF/F filter (0.7 μm). Pigments were later extracted in the laboratory in a 90% acetone + distilled water solution during 24 hours (4°C, dark conditions after 10 minutes of centrifugation). The supernatant was then dosed using a spectrofluorometer following Neveux and Lantoine (1993). The full description of the protocol followed to measure [Chla], total nitrogen concentration [TN], ammonium concentration [NH_4^+], sum of nitrate plus nitrite concentrations [NO_x], total phosphorus concentration [TP] and dissolved inorganic phosphorus concentration [DIP] is described in Bec et al. (2011) and Souchu et al. (2010). Only sub-surface data were available for all stations except the THW, THE, BRN, BRS stations located in the deep Thau and Berre lagoons for which sub-surface and bottom data were available. In these four cases, we chose to consider the bottom data.

Estimating the macrophyte morphotype beta diversity

Lagoons are characterized by a mosaic of benthic habitats, determined in particular by macrophyte diversity which can affect benthic macroinvertebrates (Arocena, 2007; Cardoso et al., 2004; Magni and Gravina, 2023; Ouisse et al., 2011) by acting as food (Jones et al., 2021; Quillien et al., 2016; Vizzini and Mazzola, 2006), refuge (Mancinelli and Rossi, 2001; Nordström and Booth, 2007; Ware et al., 2019) and/or means of transport (Norkko et al., 2000). Although sampling was carried out on locally unvegetated sediments to avoid biases linked to macrophytes and associated faunal assemblages, we postulate that the presence and composition of macrophyte assemblages within each lagoon can affect the composition of infaunal benthic communities. We developed an index (later beta diversity of macrophyte morphotypes, β_{MM}) calculated as the average Bray-Curtis dissimilarity at the lagoon or sub-lagoon scale (when several macrofauna stations were sampled in one lagoon) based on a station by macrophyte morphotype matrix (presence/absence). Each macrophyte station is coded as bare sediment if the macrophyte cover is less than 25% or as one or several of seven macrophyte morphotypes (*i.e.* mushroom, feather, hair, ball, branched, sheet and meadow) based on the thallus morphology of the dominant or co-dominant macrophyte taxa. This index varies between 0 if all the considered stations are dominated or co-dominated by the same macrophyte morphotype(s) or if they all have bare sediment and 1 if all the considered stations are dominated or co-dominated by different macrophyte morphotypes (Appendices 1 and 2). We gathered the data used to calculate this index from the WFD macrophyte dataset (all information regarding sampling protocols are available in Le Fur et al., 2018).

Data analyses

We extracted all the data used in this study from the French database Quadrige (Ifremer, http://quadrige.eaufrance.fr/). All the raw data used in this study (macrofauna, environmental data, GIS data) are available online along with all the scripts and code used to analyze the data (Jones et al., 2023). We performed all analyses using R version 3.6.3 (R Core Team, 2020) and the packages vegan (*v2.5*-6; Oksanen et al., 2019), missMDA (Josse and Husson, 2016), labdsv (*v2.0*-1; Roberts, 2019), olsrr (v0.5.3; Hebbali, 2020), MASS (Venables and Ripley, 2002), corrplot (*v0.84*; Wei and Simko, 2017) and rstatix (v0.6.0; Kassambara, 2020). We considered a significance level of 0.05 for all statistical tests.

Environmental variables

Water column parameters were measured in summer whereas macrofauna was sampled in spring which prevented us from relating these two datasets. Nonetheless, the presence and abundance of macroinvertebrates in a lagoon year y is influenced not only by the environmental conditions prevailing that year but also those prevailing the previous years especially for pluri-annual species. As such, we associated the macrofauna data collected year y to the mean and coefficient of variation (CV) of the water column parameters measured years y-2, y-1 and y, as done by Le Fur et al. (2019) (Table 2).

We classified the parameters presented in Tables 1 and 2 into two categories. The first category
included 11 physical and biotic factors used as a proxy for the natural variability of environmental
conditions (lagoon surface, lagoon-sea connection, station depth, β_{MM}, OMC, mean and CV of Sal, Temp,
O_2 sat). The second category included 14 biogeochemical variables that either cause eutrophication (*i.e.*
mean and CV of [TN], [NO$_x$], [NH$_4^+$], [TP], [DIP]) or are the direct consequence of eutrophication (*i.e.*
mean and CV of [TN], [TP], [Chla], turbidity) (Souchu et al., 2010). Because of technical problems, no
[NO$_x$] and [DIP] data were available for the stations BOL and PLS. Consequently, we imputed the missing
mean and CV values for these two parameters at both stations with the 'imputePCA' function. This
function estimates the missing parameters by first performing a principal component analysis on all the
environmental variables and then by using the regularized fitted matrix. These steps are performed
iteratively until convergence is reached (Josse and Husson, 2012).

Table 2 - Environmental parameters and their respective code used to characterize the water. For
each water parameter, the mean and coefficient of variation (CV) were calculated between the
years y-2 and y (with y the macrofauna sampling year) and the overall median, minimal and
maximal values across the 41 stations are indicated. The station with the minimal and maximal
value of each parameter is also indicated between brackets, as well as the transformation applied
to each parameter. See Table 1 for the station codes.

Parameter (unit)	Code	Measure	Median	Min	Max	Transf.
Salinity (PSU)	Sal	mean	32.39	2.83 (PLS)	42.30 (PBW)	/
		CV	13.52	2.49 (DIA)	82.24 (CAM)	$\log_e$
Oxygen saturation (%)	O_2 sat	mean	100.88	56.61 (BRN, BRS)	190.40 (BOL)	$\log_e$
		CV	27.38	0.82 (PLS)	67.33 (PRE)	$\sqrt{}$
Temperature (°C)	Temp	mean	24.08	19.63 (BRN, BRS)	27.09 (URB, BIG)	/
		CV	9.05	1.82 (CAN)	22.81 (PON)	$\sqrt{}$
Chlorophyll *a* concentration (µg.L^{-1})	[Chla]	mean	9.43	0.70 (BGS)	144.20 (MEW)	$\log_e$
		CV	71.04	25.42 (BOL)	178.73 (BIG)	$\log_e$
Total nitrogen concentration (µmol.L^{-1})	[TN]	mean	52.05	11.08 (DIA)	266.03 (MEW)	$\log_e$
		CV	27.77	6.94 (VAC)	89.47 (BIG)	$\log_e$
Ammonium concentration (µmol.L^{-1})	[NH$_4^+$]	mean	1.47	0.06 (PLS)	31.62 (GRE)	$\log_e$
		CV	106.88	28.65 (CAN)	257.93 (THE)	$\sqrt{}$
Nitrate plus nitrite concentration (µmol.L^{-1})	[NO$_x$]	mean	0.37	0.06 (GBA)	22.81 (CAM)	$\log_e$
		CV	94.87	43.30 (CAN)	227.37 (URB)	$\log_e$
Total phosphorous concentration (µmol.L^{-1})	[TP]	mean	2.51	0.64 (LEN)	31.19 (CAN)	$\log_e$
		CV	36.75	8.91 (VAC)	106.12 (ARN)	$\log_e$
Phosphate concentration (µmol.L^{-1})	[DIP]	mean	0.24	0.02 (LES)	14.53 (CAN)	$\log_e$
		CV	93.10	40.97 (THW)	187.55 (INN)	$\log_e$
Turbidity (NTU)	Turb	mean	5.20	1.14 (BGS)	99.65 (PLS)	$\log_e$
		CV	68.30	1.92 (PLS)	207.54 (ARN)	$\sqrt{}$

285   Transformation (Transf.): / = no transformation, $\log_e$ = natural logarithm, $\sqrt{}$ = square-root

Macrobenthic community structure

First, we calculated the mean macroinvertebrate abundance at the station scale by averaging over
the three replicates and transformed it using a Hellinger transformation, to give more weight to common
taxa and less weight to highly abundant, rare and under-sampled taxa (Legendre and Legendre, 2012).
We then classified the 41 stations based on these transformed mean abundances, using K-means
partitioning and the Calinski-Harabasz criterion with 100 iterations ('cascadeKM' function). Our goal was
to highlight similarities between lagoons based on common species without focusing on rare taxa, as
each coastal lagoon can present unique species assemblages (Basset et al., 2007), and to help with the

result interpretation thanks to a simple classification. We tested for significant differences in the
composition of our macrofauna groups using the analysis of similarity (ANOSIM) test with 9999
permutations and Euclidean distances ('anosim' function). Finally, we characterized each group using the
indicator species metric ('indval' function, 9999 permutations). A high value of this metric for a given
taxon associated to a given group means this taxon is consistently more present (fidelity) and abundant
(specificity) in the stations composing that group than in stations composing the other groups (Dufrêne
and Legendre, 1997). For this last analysis, we adjusted p-values using the False Discovery Rate correction
method (FDR; Benjamini and Hochberg, 1995).

Taxonomy-based indices
To characterize the macrobenthic community structure of each station, we calculated the commonly
used M-AMBI (Multivariate AZTI Marine Biotic Index), the three indices composing the M-AMBI - species
richness (SR), Shannon diversity index (H') and AMBI (Muxika et al., 2007) - plus the inverse of Simpson's
index (N2) and the total macrofauna density. The AMBI is an index based on the classification of benthic
species into ecological groups, from the most sensitive to disturbances (i.e. organic matter enrichment)
to the most opportunistic ones (Borja et al., 2000). Following recommendations of Borja et al. (2005), we
calculated AMBI and M-AMBI at the replicate scale and then averaged them at the station scale.
Similarly, we calculated SR, H', N2 and total macrofauna density by averaging over the three replicates
and computed the corresponding standard deviations (SD). For AMBI and M-AMBI calculations, we
removed high order taxa without an ecological group value (Anemonia and Polycladida), planktonic
(Mysidae) and hard-substrate taxa (Serpula and Serpulidae) (Borja et al., 2005). We computed M-AMBI
using the R function available in Sigovini et al. (2013) and we defined the good reference conditions as
the highest values of SR and H' and lowest value of AMBI measured in the dataset, irrespective of the
WFD physico-chemical status described above. Finally, we tested for differences in total macrofauna
density, SR, H', N2, AMBI and M-AMBI between previously defined groups of stations (see section
Macrobenthic community structure) using non-parametric Kruskal-Wallis rank sum tests (Breslow, 1970).

Linking macrobenthic communities and environmental variables
To assess how natural environmental variations correlated with anthropogenic pressures
(explanatory variables) and how each explanatory variable correlated with macrobenthic indices, we built
a correlogram showing the significant Spearman correlations ('cor_pmat' function) between all the raw
variables. We also tested for differences in the variables considered (explanatory variables and
macrobenthic indices) between lagoon-sea connection levels using non-parametric Kruskal-Wallis rank
sum tests (Breslow, 1970), followed by pairwise Wilcoxon tests for multiple comparison, while adjusting
p-values with FDR (Benjamini and Hochberg, 1995). Then, we transformed (natural logarithm or square
root transformation), when necessary, the environmental variables and macrobenthic indices to
approximate normal distributions and reduce data spread, before standardizing all the explanatory
variables to facilitate the interpretation of the multiple linear regression coefficients (Schielzeth, 2010).
We log_e-transformed the lagoon surface, station depth, SR, N2 and macrofauna density (see Table 2 for
the rest of the transformations). To limit multicollinearity and avoid artificially inflating the fit of linear
models, we reduced the number of environmental variables (see section *Environmental variables*) for all
future analyses by considering Pearson correlations < |0.6|, leading to 19 remaining environmental
variables.
To determine which environmental variables most strongly influence lagoon macrobenthic
communities, we performed a redundancy analysis (RDA, 'rda' function) between the 19 previously
selected environmental variables and the Hellinger-transformed mean taxa abundances. We computed
the variance inflation factor (VIF, 'vif.cca' function) to check for multicollinearity and sequentially
removed variables with a VIF > 5. We verified the significance of the full RDA model using the 'anova.cca'
function (1000 steps) and we performed a forward model selection procedure ('ordiR2step' function)
based on the adjusted R2 of the full model and a p-value of 0.05 (9999 permutations) (Blanchet et al.,
2008). We then computed the marginal effect of each selected explanatory variable as the adjusted R^2 of
the constrained ordination containing only the given variable as the predictor. The conditional effect of a
variable was equal, during the forward selection procedure, to the additional amount of variance in
species communities explained by the corresponding variable at the time it was included into the model.

Finally, we performed a partial RDA using the variables previously selected, to isolate the impact of
eutrophication from natural environmental variability on lagoon macrobenthic communities (Legendre
and Legendre, 2012).
In the same way as for the constrained multivariate analysis, we implemented a multiple linear
regression approach (without any interactions between variables) to investigate which variables related
to eutrophication and natural variability most strongly influenced the macrobenthic indices (SR, H', N2,
total macrofauna density, AMBI and M-AMBI). As there were still 19 potential explanatory variables, we
performed a preliminary selection by including all the quantitative variables which alone had a significant
effect on the index as first or second-degree orthogonal polynomial and the lagoon-sea connection if at
least one of the three categories (weak, moderate, strong) had a significant effect on the index ('lm'
function). We included second-degree polynomials to account for potential non-linear relations (Pérez-
Ruzafa et al., 2007b) and verified that the selected variables had VIFs inferior to 5. We also checked the
model hypotheses using the residuals vs fitted values plot for linearity, the Shapiro-Wilk test for residual
normality, the Breush Pagan test for residual homoscedasticity ('ols_test_breusch_pagan' function) and
the 'acf' function for the detection of residual spatial autocorrelation based on the station coordinates.
Then, we ran a stepwise model selection using the 'stepAIC' function set in both directions and the BIC
criterion. Finally, we checked the hypotheses of the selected model as previously explained, we
investigated potential high leverage points (Cook's distance > 0.5) and we reran the selected model
without these points to verify its robustness. To compare the relative effect of each selected explanatory
variable on a given macrobenthic index, we expressed the slope estimates of the standardized variables
as effect sizes when the variable was quantitative and as mean standardized effects when the variable
was categorical (Schielzeth, 2010).
Lastly, we implemented a variance partitioning procedure using the 'varpart' function, to assess how
much community variation or macrobenthic index variation could be: (i) uniquely attributed to
eutrophication when the effect of natural variability had been removed, (ii) uniquely attributed to natural
variability when the effect of eutrophication had been removed or (iii) jointly attributed to both effects
(Legendre and Legendre, 2012). Prior to this analysis, we reran the variable selection procedures
implemented at the community and macrobenthic index levels considering alternatively only the
eutrophication variables or only the variables related to natural variability. We then included the
variables selected by the aforementioned method in the variance partitioning, which was based on
partial RDAs or on partial linear regressions and we presented the results using the adjusted R^2 values.

Results

Inter-lagoon environmental variability

The dataset considered in this study encompasses a wide diversity of lagoons in terms of physico-
chemical state according to the WFD and environmental parameters associated to natural variability and
eutrophication (Tables 1 and 2). First, only lagoons with a strong (Leucate, Gruissan, Thau, Berre and
Diana) or a moderate (La Palme and Bages-Sigean) connection to the sea were in a good physico-
chemical state (green in Table 1). All the stations sampled in these lagoons presented mean [Chla] below
7 µg.L^{-1} and all of them except the ones located in the deep Berre and Diana lagoons (station depth > 5.8
m) presented a high beta diversity of macrophyte morphotypes (β_{MM} > 0.53). Five stations sampled in
these lagoons presented the lowest value for a eutrophication related variable (DIA: mean [TN], BGS:
mean [Chla] and mean Turb, LEN : mean [TP], LES: mean [DIP], THW: CV [DIP]). However, the THE station
(Thau) had the most variable [NH_4^+] and the BRN and BRS stations (Berre) had the lowest mean O_2 sat of
the dataset. Seven lagoons were in a moderate (yellow in Table 1) or in a bad physico-chemical state
(orange in Table 1) and were either moderately (Ponant, Vaine, Urbino) or weakly (Grand Bagnas,
Vaccarès, Biguglia and Palo) connected to the sea. In this group, the GBA and VAC stations respectively
had the lowest mean [NO_x] and the lowest CV [TN] of the dataset whereas the URB and BIG stations
respectively had the highest CV [NO_x] and CV [Chla]. All the remaining weakly connected lagoons were in
a poor physico-chemical state (Canet, Campignol, Vendres, La Marette, La Palissade, Bolmon). In this
group, the CAN station had both the highest mean [TP] and [DIP] and the lowest CV [NH_4^+] and [NO_x] of
the dataset, while the CAM and BOL stations presented respectively the highest mean [NO_x] and CV
[Chla] of the dataset. Finally, all the lagoons between the cities of Sète and Montpellier (Figure 1) were in

a poor condition, despite being strongly (Ingril North, Ingril South, Prévost) or moderately (Vic, Pierre-
Blanche, Arnel, Méjean and Grec) connected to the sea. All the stations located in these lagoons except
PBE, had OMC above 7% and up to 13%. In this group, the MEW station presented the highest mean
[Chla] and [TN] of the dataset while the GRE station had the highest mean [NH₄⁺], the ARN station the
highest CV [TP] and Turb and the INN station the highest CV [DIP] of the dataset.

Based on the significant Spearman correlations between the environmental variables (Figure 2),
stations in the larger lagoons were significantly deeper, cooler in summer and less variable in terms of
summer salinity, presented higher β_{MM} and lower values for 8 eutrophication related variables (*e.g.* mean
[Chla], mean [NOₓ] and CV [NOₓ]). The deepest stations were also the ones with the highest OMC. Two
variables related to natural environmental variability were also significantly different according to the
lagoon-sea connection level (KW test, $p < 0.05$ and post-hoc pairwise Wilcoxon tests $p < 0.05$): mean Sal
(moderate > weak and strong > weak) and CV Sal (weak > moderate > strong).
410

Figure 2 - Correlogram showing only significant positive (red toned circles) and negative (grey toned
circles) Spearman correlations ($p < 0.05$) between the untransformed taxonomic indices (in black,
SR: species richness, H': Shannon diversity, N2: inverse of Simpson's dominance, M-AMBI:
Multivariate AMBI), the untransformed environmental variables related to the lagoon's natural
variability (in blue, β_{MM}: beta diversity of the macrophyte morphotypes, OMC: sediment organic
matter content) and the untransformed environmental variables related to eutrophication
expressed as means and coefficient of variations (in green, [Chla]: chlorophyll *a* concentration, [TN]:

total nitrogen concentration, [NH$_4^+$]: ammonium concentration, [NO$_x$]: nitrate plus nitrite
concentration, [TP]: total phosphorous concentration, [DIP]: dissolved inorganic phosphorous
concentration) calculated across 41 stations located in 29 lagoons along the French Mediterranean
coast.

Macrofauna communities and macrobenthic indices across French Mediterranean lagoons

423
A total of 42 465 benthic macroinvertebrates were counted and identified across all samples,
belonging to 227 different taxa with most organisms identified to species level (74.5%), others to genus
(17.2%), family (2.6%), order (2.6%), subclass (0.4%), class (1.8%) or phylum (0.9%). Across the 227 taxa,
1.3% were identified in over 50% of the stations, 90% in less than eight stations and 40% in only one
station. No taxon was present in all 41 stations. Polychaetes were by far the most diversified group (102
taxa), followed by Malacostraca crustaceans (54 taxa), bivalves (25 taxa) and gastropods (19 taxa).
Among the three taxa with a frequency of occurrence above 50%, two were polychaete species (*Hediste*
diversicolor (66%) and *Heteromastus filiformis* (51%)), and the third was the bivalve *Cerastoderma*
glaucum (51%). These three species also presented an important inter-station variability in abundance.
For all the following analyses, we considered 217 taxa (see section *Benthic macroinvertebrates*), except
for the AMBI and M-AMBI calculation, for which we considered only 213 taxa (see section *Taxonomy-*
based indices).
The K-means clustering delineated two groups of stations with distinct macrobenthic communities
(minimal difference in Calinsky-Harabasz criterion = 1.4), a result confirmed by the analysis of similarity
test (ANOSIM statistic = 0.51, *p* < 0.001). These groups did not correspond to any obvious spatial or
salinity pattern, as exemplified by the four Corsican stations and the seven oligo-mesohaline stations
which were classified in both groups (Figure 1). The polychaete *Hediste diversicolor* (*p* = 0.022) was an
indicator taxa of group A stations (red in Figure 1) along with the bivalve *Abra segmentum* (*p* = 0.022),
Chironomidae (*p* = 0.029) and the amphipod *Gammarus aequicauda* (*p* = 0.038). Conversely, group B
stations (blue in Figure 1) were significantly characterized by the bivalve *Scrobicularia cottardii* (*p* =
0.048), closely followed by another bivalve, *Loripes orbiculatus* (*p* = 0.065).
Finally, we found significant differences among the two groups in terms of SR and M-AMBI; group A
was composed of stations with lower SR and weaker M-AMBI values than group B (SR: $\chi^2(1)$ = 6.41, *p* =
0.011; M-AMBI: $\chi^2(1)$ = 5.76, *p* = 0.016). Group B was also characterized by stations with higher H' and N2
values and higher maximal macroinvertebrate densities than stations composing group A, albeit these
tendencies were not significant (Appendix 3).

Relationship between macrofauna communities and environmental variables

450
In all the following analyses, six variables were not considered because they were highly correlated
with another variable (|r| ≥ 0.6): station depth (to lagoon surface), CV Sal (to mean Sal), mean [TN],
mean [TP], mean Turb (to mean [Chla]) and CV [TP] (to CV [TN]). Redundancy analysis (RDA) was used to
relate the macrofauna communities recorded within the lagoons to their natural environmental
variability and their eutrophication levels (Figure 3). Mean Sal was also not considered because of a VIF >
5 linked to significant differences between lagoon-sea connection levels (see section *Inter-lagoon*
environmental variability). The final RDA model, determined using a forward model selection procedure,
explained 22% of the inter-station community variability (model adj. R^2) and included six variables related
to natural environmental variability (lagoon-sea connection level (*p* < 0.001), mean Temp (*p* < 0.001),
lagoon surface (*p* = 0.0076), CV Temp (*p* = 0.013), mean O$_2$ sat (*p* = 0.026) and OMC (*p* = 0.041) and two
eutrophication related variables (mean [Chla] (*p* < 0.001) and mean [DIP] (*p* = 0.024)). Regarding the
individual effect of environmental variables, mean [Chla] (7.22%), lagoon surface (5.90%), lagoon-sea
connection (5.30%) and mean Temp (5.04%) were the most influential variables according to their
marginal effect. Based on conditional effects, the most influential variables were mean [Chla] (7.22%),
lagoon-sea connection (4.64%), mean Temp (3.62%) and lagoon surface (1.69%) (Appendix 4).
The first two axes accounted for 33.8 and 17.8% of the RDA model variance, respectively. First, the
stations from group A and B were clearly separated along the first axis of the RDA, except for LPN, LPS,
PLS and INS (Figure 3a). On the ordination of the environmental variables (Figure 3b), lagoon surface and
CV Temp grouped on the positive side of the first axis while mean [Chla], mean [DIP] and mean Temp
grouped on the negative side of the first axis. Consequently, group B macrobenthic communities were

linked to large lagoons (also relatively deep), several of which presented high summer temperature
variations. They were characterized by higher relative abundances of the bivalves *S. cottardii* and *L.*
orbiculatus (Figure 3c). Conversely, group A communities were associated to smaller lagoons, subject to
higher mean summer temperatures and high eutrophication levels. They were characterized by higher
relative abundances of the polychaete *H. diversicolor* and Chironomidae (Figure 3c).
The second axis mainly corresponded to a confinement and also to an organic matter content
gradient, mostly separating each group into two sub-groups with the lagoons weakly connected to the
sea presenting sediments enriched in organic matter located towards the positive side of the second axis
and the moderately and strongly connected lagoons located towards the negative side of the second axis.
As mean Sal and CV Sal were correlated to lagoon-sea connection level (see section *Inter-lagoon*
environmental variability), this axis also corresponded to a salinity gradient with low and variable salinity
stations characterized by a higher

Figure 3 - Redundancy analysis plots (RDA, axis 1 and 2) focusing on (a) the benthic macrofauna
sampled across 41 stations located in 29 lagoons along the French Mediterranean coast (see Table 1
for the full lagoon names and corresponding stations) as explained by the set of environmental
variables related to eutrophication and natural variability composing the most parsimonious model,
(b) the selected environmental variables, where mean [Chla], lagoon surface, mean O_2 sat and

abundance of *Chironomus salinarius* and *Streblospio shrubsolii* (Figure 3c). All the oligo- and meso-haline stations (triangles on Figure 3) grouped on the upper part of the RDA space and most poly- and eu-haline stations (circles on Figure 3) grouped on the lower part of the figure. Furthermore, communities of lagoons weakly connected to the sea were characterized by higher abundances of the bivalve *Scrobicularia cottardii* for group B, while communities from lagoons moderately to strongly connected to the sea were characterized either by higher abundances of *L. orbiculatus* for group B or by a diversified suite of species like the decapod *Carcinus aestuarii* or the polychaete *Capitella capitata* for group A.

Once the effect of natural environmental variability on macrofauna communities was removed through partial RDA and the only explanatory variables left to vary were mean [Chla] and mean [DIP] (eutrophication variables), no differences could be made between the communities of groups A and B (Appendix 5). This partial RDA explained only 6.4% of the inter-station community variability (model adj. R^2).

Relationship between macrobenthic indices and environmental conditions

Based on the significant Spearman correlations between the raw environmental variables and raw macrobenthic indices (Figure 2), SR, H', N2 and M-AMBI were significantly higher at stations where mean Temp was lower, salinity was more stable and eutrophication related variables were lower, especially mean [Chla]. Furthermore, H', N2 and M-AMBI were significantly higher at stations where mean Sal and β_{MM} were higher while SR and M-AMBI were also significantly higher at deeper stations located in larger lagoons. AMBI appeared to be significantly correlated positively only with mean [NO$_x$] while total macrofauna density was significantly higher at stations with lower but more variable summer water temperatures, lower salinity and lower mean [DIP].

The results of the multiple linear regressions identifying which variables influenced the macrobenthic indices are presented in Table 3. The most parsimonious models contained between two and four predictors and explained between 27% and 74% of the macrobenthic indices' variability. All the models contained at least one predictor associated to eutrophication and at least one predictor associated to natural variability except the N2 and AMBI models, which only contained predictors associated to natural variability and eutrophication respectively. Lagoon-sea connection was selected in three models, H', N2 and M-AMBI, and in all three, the response variable was significantly higher in strongly connected lagoons. Differently, H' and N2 were significantly higher and lower in moderately connected lagoons, respectively, while M-AMBI was significantly lower in weakly connected lagoons. Mean Temp and mean Sal were selected in two models, SR and macrofauna density. Mean Temp negatively influenced SR and macrofauna density while mean Sal positively influenced SR and had a quadratic effect (concave form) on macrofauna density. B_{MM} was selected in two models, H' and N2, and positively influenced both response variables. The AMBI model was composed of mean [TN] (positive effect) and CV [NH$_4^+$] (quadratic effect, concave form). Finally, three other eutrophication related variables, CV [TN], mean [NH$_4^+$] and CV [NO$_x$], were selected respectively in H', macrofauna density and M-AMBI models where they had either a negative effect (M-AMBI) or a quadratic effect (concave form, H' and density).

To isolate the independent effect of eutrophication on macrobenthic indices, we also plotted the partial effect of mean [Chla] on SR, H' and M-AMBI once the effect of the other predictors retained in the minimal adequate models (Table 3) had been removed (Appendix 6). After controlling for the effect of mean Sal and mean Temp, the simple linear regression model between mean [Chla] and SR was still significant (adj. R^2 = 0.32, $F(1, 39)$ = 19.55, $p < 0.001$). It was also the case for the model between mean [Chla] and H' after controlling for the effect of lagoon-sea connection, CV [TN], CV [TN]^2 and β_{MM} (adj. R_2 = 0.14, $F(1, 39)$ = 7.33, $p = 0.01$) and for the model between mean [Chla] and M-AMBI after controlling for lagoon-sea connection and CV [NO$_x$] (adj. R^2 = 0.26, $F(1, 39)$ = 15.41, $p < 0.001$).

Table 3 - Most parsimonious models (ordinary least-square regressions) explaining the standardized macrobenthic taxonomic indices (species richness (SR), Shannon diversity (H'), inverse of Simpson's dominance (N2), AZTI's Marine Biotic Index (AMBI), Multivariate AMBI (M-AMBI)). SR, H', N2 and total macrofauna density were $\log_e$ transformed prior to standardization. Both eutrophication related and natural variability related variables are considered as explanatory variables and the slope estimates are expressed either as effect sizes when the variable is quantitative or as mean standardized effects when the variable is categorical.

Model terms		Estimate	SE	p-value
SR: $F_{(3,37)} = 18.18$; RSE = 0.66 ; adj. $R^2 = 0.56$				
[Chla]	mean	-0.50	0.12	< 0.001
Temperature	mean	-0.34	0.11	0.005
Salinity	mean	0.27	0.11	0.018
H': $F_{(6,34)} = 13.57$; RSE = 0.59 ; adj. $R^2 = 0.65$				
Lagoon-sea connection	strong	0.61	0.19	< 0.001
	moderate	0.39	0.15	0.016
	weak	-0.16	0.17	0.32
[Chla]	mean	-0.31	0.12	0.016
[TN]	CV	-0.02	0.11	0.89
[TN]^2		-0.30	0.10	0.004
β_{MM}		0.23	0.10	0.035
N2: $F_{(3,37)} = 11.18$; RSE = 0.75 ; adj. $R^2 = 0.43$				
Lagoon-sea connection	strong	0.79	0.21	< 0.001
	moderate	-0.58	0.19	0.005
	weak	-0.27	0.22	0.11
β_{MM}		0.31	0.12	0.016
Total macrofauna density: $F_{(5,35)} = 8.48$; RSE = 0.72 ; adj. $R^2 = 0.48$				
Temperature	mean	-0.36	0.12	0.006
Salinity	mean	-0.23	0.12	0.071
Salinity^2		-0.30	0.13	0.023
[NH$_4^+$]	mean	0.21	0.12	0.10
[NH$_4^+$]^2		-0.30	0.13	0.030
AMBI: $F_{(3,37)} = 5.84$; RSE = 0.86 ; adj. $R^2 = 0.27$				
[TN]	mean	0.34	0.14	0.017
[NH$_4^+$]	CV	0.11	0.14	0.41
[NH$_4^+$]^2		-0.43	0.13	0.003
M-AMBI: $F_{(4,36)} = 29.04$; RSE = 0.51 ; adj. $R^2 = 0.74$				
Lagoon-sea connection	strong	0.59	0.16	0.005
	moderate	-0.06	0.14	0.67
	weak	-0.52	0.15	0.028
[NO$_x$]	CV	-0.37	0.09	< 0.001
[Chla]	mean	-0.35	0.09	< 0.001

RSE: Residual standard error, CV: coefficient of variation, [Chla]: chlorophyll a concentration in the water, [TN]: total nitrogen concentration in the water, β_{MM}: beta diversity of the macrophyte morphotypes, [NH$_4^+$]: ammonium concentration in the water and [NO$_x$]: nitrate plus nitrite concentration in the water

To synthesize our results, we plotted for each station the M-AMBI along with the three macrobenthic
indices composing it (SR, H' and AMBI) (mean ± SD) as a function of mean [Chla] (Figure 4). Mean [Chla]
was chosen as a synthetic indicator of eutrophication because it is (*i*) significantly and positively
correlated to 9 out of the 13 eutrophication related variables (Figure 2), (*ii*) the eutrophication related
variable (with mean [TN]), that presents the strongest correlations with SR, H' and M-AMBI (Figure 2), (*iii*)
the most frequently selected eutrophication related variable in the macrobenthic index models (Table 3)
and (*iv*) a relevant indicator of eutrophication status according to Bec et al. (2011) and Cloern (2001).

Figure 4 - Mean (± SD) of the species richness (SR), Shannon diversity (H'), AMBI and M-AMBI
indices as a function of mean chlorophyll a concentration ($\log_e$ scale) across 41 stations located in
29 lagoons along the French Mediterranean coast (see Table 1 for the full lagoon names and
corresponding stations). Across all plots, each station (point) is represented using a symbol for its
salinity type (triangle: oligo- and meso-haline and circle: poly- and eu-haline) and a color (red: group
A and blue: group B) for its group membership (K-means partitioning) based on Hellinger-
transformed mean abundances. The dashed lines correspond to loess regressions.

SR, H' and M-AMBI were significantly correlated to mean [Chla] (r (Spearman) > |0.50|) while AMBI was
not (r = 0.12). Based only on the mean [Chla], most stations from group B (in blue) grouped towards
higher levels of mean SR, H' and M-AMBI than stations from group A (in red). Nonetheless, we identified
an intermediary zone corresponding to mean [Chla] between 6.3 µg.L[-1] and 14.2 µg.L[-1] and composed of

stations from both group located between the INN (SR = 12 ± 3, H' = 3.06 ± 0.42 and M-AMBI = 0.57 ±
0.1) and ORE (SR = 9 ± 1, H' = 1.84 ± 0.07 and M-AMBI = 0.31 ± 0.01) stations. In the line with these
observations, the LPN, LPS and BOL stations appeared as outliers, as did LPN and LPS in the RDA (Figure
3). Finally, mean [Chla] was also significantly correlated (r (Spearman) = -0.40, p = 0.009) to the intra-
station SR variability expressed using the standard deviation.
580
Disentangling natural from anthropogenic factors
Based on the eutrophication and natural variability variables independently selected, we performed a
variance partitioning, to investigate the amount of community and macrobenthic index variability
uniquely linked to each variable category (Figure 5). Macrobenthic community and indices can be
grouped into three categories according to the relative importance of natural variability alone,
eutrophication alone, and the two together, in explaining their respective variability. (*i*) Macrofauna
density was first influenced uniquely by natural variability (29.0%), then jointly by natural variability and
eutrophication (11.5%) and finally uniquely by eutrophication (7.9%). (*ii*) Macrobenthic community, N2
and H' were first jointly influenced by natural variability and eutrophication (community: 8.2%, H': 33.6%,
N2: 22.1%), and then uniquely by natural variability (community: 8.0%, H': 17.8%, N2: 21.1%) and finally
uniquely by eutrophication (community: 3.7%, H': 13.9%, N2: 2.9%). (*iii*) SR and M-AMBI were first jointly
influenced by natural variability and eutrophication (SR: 30.7%, M-AMBI: 49.0%), and then uniquely by
eutrophication (SR: 20.3%, M-AMBI: 17.6%) and finally uniquely by natural variability (SR: 5.8%, M-AMBI:
8.9%).
595

Figure 5 - On the left, variance (adjusted R2) of the macrofauna community data (community), total
macrofauna density (density), inverse of Simpson's dominance (N2), Shannon diversity (H'), species
richness (SR) and multivariate AMBI (M-AMBI) uniquely attributed to eutrophication (green),
uniquely attributed to natural variability (blue) and jointly attributed to both effects (red). The
residual variance is shown in white. On the right, eutrophication variables and variables related to
natural variability considered in the variance partitioning of the macrofauna community and
macrobenthic taxonomic indices.

Using figure 2, we can identify variables related to eutrophication or to natural variability that
significantly co-vary and are jointly implicated in a macrobenthic index model (Table 3) and/or in a
variance partitioning (Figure 5). Mean [Chla] was significantly correlated with lagoon surface (-0.35), β_{MM}
(-0.37), mean Temp (0.40) and CV O_2 sat (0.40). Mean [NO$_x$] was significantly correlated with lagoon
surface (-0.38), mean Sal (-0.49) and CV O_2 sat (-0.33) while CV [NO$_x$] was significantly correlated with

lagoon surface (-0.37), β_{MM} (-0.36) and mean Temp (0.53). Finally, mean [DIP] and lagoon surface were
significantly correlated (-0.32). Lagoon-sea connection, a qualitative variable, was another source of
confounding effects between natural variability and eutrophication on the macrobenthic indices and
community. Indeed, there was a significant difference between the lagoon-sea connection and: *(i)* mean
[Chla] (weak > strong connection); *(ii)* mean [NO$_x$] (weak > strong connection) and *(iii)* CV [NO$_x$]
(moderate > strong).

Discussion

616   As outlined by Cloern (2001), eutrophication is a complex phenomenon with direct (*e.g.* changes in
617 dissolved nutrient contents) and indirect (*e.g.* changes in community) responses that are modulated by
618 attributes of each recipient system. This study is a first step towards better understanding how hydro-
619 morphology, natural environmental variability and eutrophication jointly shape macrobenthic
620 communities in Mediterranean coastal lagoon ecosystems.
621
Macrobenthic communities are driven by lagoon hydro-morphology and natural environmental
(in)stability
Lagoon macrobenthic communities appear to be first determined by lagoon hydro-morphology and
natural environmental stability. When investigating the independent effect of each environmental
parameter on macrobenthic communities, lagoon-sea connection level was the variable explaining the
most of H', N2 and M-AMBI variability and the second most important variable explaining community
composition. This qualitative variable is a proxy for marine water renewal (*i.e.* residence time), a gradient
well known for characterizing coastal lagoon biological communities (Barbone and Basset, 2010; Perez-
Ruzafa and Marcos-Diego, 1992; Tagliapietra et al., 2009), and closely linked to mean Sal (moderate >
weak and strong > weak) and CV Sal (weak > moderate > strong). Salinity, historically used to classify
coastal lagoons (Anonymous, 1959), was also an important driver of SR (linear positive effect) and total
macrofauna density (concave effect). Indeed, simple and multimetric descriptors of lagoon macrobenthic
communities like species richness are known to be highly and positively influenced by mean water
salinity (Barbone et al., 2012; Magni et al., 2022; Pitacco et al., 2019; Reizopoulou et al., 2014).
Differently from us, Barbone et al. (2012) reported a macrofauna density decrease from oligo- to eu-
haline lagoons, maybe because only undisturbed lagoons were considered in their study. Furthermore,
the effect of lagoon-sea connection on N2, a diversity index sensitive to high abundance taxa (Morris et
al., 2014), indicates that a strong connection to the sea promotes benthic communities with significantly
more relatively abundant taxa whereas moderate and weak connections promote communities
dominated by a few abundant and in our case, euryhaline taxa (*e.g. Hediste diversicolor, Streblospio*
shrubsolii), a characteristic typical of naturally stressed environments like lagoons subject to strong
salinity variations (Elliott and Quintino, 2007). This result seems to confirm that moderately and weakly
connected lagoons are highly selective environments for benthic organisms at least partly because of
their strongly fluctuating environmental conditions (Barnes, 1980; Provost et al., 2012; Zaldívar et al.,
2008). The negative effect of decreasing salinity and lower connection levels to the sea on SR, N2 and
macrofauna density can also be linked to the physiological stress that natural and abrupt salinity
variations (*e.g.* droughts, heavy rainfall) can represent for larvae, juveniles and adults of marine-
originating benthic organisms inhabiting coastal lagoons (Cognetti and Maltagliati, 2000; Magalhães et
al., 2019; Vernberg, 1982). Overall, our results confirm that *(i)* water renewal from marine origin, *(ii)*
primary colonization and/or post-disturbance recolonization of lagoons by marine-originating larvae
through dispersal and recruitment and *(iii)* environmental (in)stability, strongly shape lagoon benthic
communities (Basset et al., 2006; Pérez-Ruzafa et al., 2007b; Pérez-Ruzafa and Marcos-Diego, 1992), as
proposed in the confinement theory (Guelorget and Perthuisot, 1983).
Water temperature was another important driver of macrobenthic communities, SR and macrofauna
density. Temperature had a negative effect on these communities, probably directly through thermal
stress, and indirectly through its effect on oxygen dynamics. Higher and more variable summer
temperatures directly represent an abiotic constraint for benthic organisms by increasing their oxygen
demand and reducing their mobility (Vernberg, 1982). At the same time, high summer temperatures
decrease oxygen solubility and increase the probability of hypoxia and anoxia (Derolez et al., 2020a)

while also reducing the tolerance of benthic organisms to low dissolved oxygen (Vaquer-Sunyer and Duarte, 2011). This cocktail of effects can ultimately cause the death of sensitive taxa, such as large sized and long-lived species (Quillien et al., 2015; Reizopoulou and Nicolaidou, 2007; Sigala et al., 2012) and reduce the ability to move of more tolerant taxa (Vernberg, 1982), hence increasing their susceptibility to predation and decreasing their possibility to search for food and/or for less stressful environmental conditions (Díaz and Rosenberg, 1995; Rosenberg et al., 1992; Weissberger et al., 2009).

The diversity of macrophyte morphotypes (β_{MM}) also affected positively H' and N2, and was positively correlated to all macrobenthic indices except density and AMBI. Simple taxonomic indices like lagoon macrobenthic species richness, biomass density and numerical density are locally sensitive to vegetation presence and type (Arocena, 2007; Barbone et al., 2012, Magni and Gravina, 2023). Conversely, we show here the probable effect of benthic habitat heterogeneity expressed at the lagoon or sub-lagoon scale on the structure of benthic communities at a local scale (*ca.* 50 m^2 station), an effect mainly visible when focusing on intermediate and high abundance taxa (Morris et al., 2014). Focusing on macrobenthic fauna at the lagoon scale, β_{MM} informs us on the number of potential *(i)* environmental niches that can be filled by the species pool and *(ii)* refuges (Magni and Gravina, 2023; Nordström and Booth, 2007; Ware et al., 2019) from which these organisms can (re)colonize habitat patches after mortality events caused by temporary unfavorable conditions. A model of local extinctions linked to adverse environmental conditions, followed by recolonization events via pelagic larvae was suggested by Vergara-Chen et al. (2013) to explain the small-scale genetic structure of the cockle *Cerastoderma glaucum* inside the Mar Menor lagoon (Spain). The positive effect of β_{MM} on macrobenthic diversity also provides us with a few leads on how macrophytes can structure lagoon macrobenthic communities. Macrophytes can act as *(i)* temporary refuges from vagile predators for small mobile benthic organisms like amphipods (Nordström and Booth, 2007; Ware et al., 2019), *(ii)* a food source for grazers and detritus-feeders (Carlier et al., 2007; Lepoint et al., 2000; Ouisse et al., 2011; Vizzini and Mazzola, 2006) and *(iii)* settlement promoters for pelagic particles like phytoplankton and larvae (Colden et al., 2016; Donadi et al., 2014; González-Ortiz et al., 2014), especially in the case of meadow-forming species like *Zostera.* All these local effects are likely to spill-over to adjacent unvegetated sediments (Boström et al., 2006; Magni et al., 2017).

Finally, the two macrofauna stations sampled in La Palme, a lagoon considered in a good physico-chemical status, were classified, based only on macrofauna, in a group generally associated to small eutrophic lagoons and mainly composed of stations sampled in lagoons evaluated by the WFD as being in a poor or in a bad physico-chemical status. Consequently, the eight environmental variables selected to explain the macrofauna communities were not able to explain the one characterizing this lagoon, which also presents marked seasonal salinity variations (between 3.6 and 76.8 in Wilke and Boutiere (2000)), variations indetectable with the available summer salinity data. Overall, natural environmental (in)stability, connection level with adjacent water systems and/or number of potential environmental niches seem to be the main drivers of lagoon macrobenthic communities and to be stronger drivers of benthic macrofauna assemblages than eutrophication, which probably represents an additional source of stress for benthic invertebrates (Barbone and Basset, 2010).

Eutrophication affects lagoon macrobenthic communities through interconnections with natural environmental variability

Eutrophication appears to act as an additional disturbance for lagoon macrobenthic communities via direct (*e.g.* toxicity of inorganic nitrogenous compounds) and indirect mechanisms (*e.g.* hypoxic stress). The three groups derived from the variance partitioning approach highlight that the most influential category of variables (*i.e.* natural variability alone, eutrophication alone, natural variability and eutrophication together) is not the same depending on the level of biological organization we focus on. Indeed, the benthic invertebrate density was first explained by natural variability alone and more precisely by physiologically constraining variables (salinity and temperature) whereas macrobenthic community composition, SR, H' and N2 were first explained jointly by natural variability and eutrophication. With these joint effects, we probably measure *(i)* how lagoon-sea connection modulates the susceptibility (*i.e.* propensity) of each lagoon to eutrophication, *(ii)* how higher water temperatures promote phytoplankton blooms and *(iii)* how eutrophication reduces benthic habitat heterogeneity, followed by their respective effects on benthic community diversity.

(i) Lagoon-sea connection informs us on each lagoon's water residence time and consequently, on its
retention potential regarding nutrients and pollutants (Chapelle et al., 2001; Cloern, 2001; Fiandrino et
al., 2017). As such, watershed related nutrients and their direct effect on primary producers are buffered
by each lagoon's degree of connection to the sea, as indicated by the significantly higher mean [Chla] and
mean [NO$_x$] in lagoons weakly connected to the sea compared with strongly connected ones. (ii)
Independently from lagoon hydro-morphology, the positive correlation between mean [Chla] and mean
temperature illustrates how higher water temperatures increase phytoplankton growth rates which can
lead to blooms (Rose and Caron, 2007), as recently demonstrated in Thau (Trombetta et al., 2019), a
large lagoon strongly connected to the sea. Nonetheless, it seems that eutrophication is buffered in
larger lagoons where water temperatures increase less in summer, as suggested by the negative
correlation between lagoon surface and mean summer temperature. (iii) Furthermore, one of the direct
responses of Mediterranean coastal lagoons to changes in nutrient inputs is an increase in pelagic
primary production (Souchu et al., 2010), which in turn decreases water transparency (Cloern, 2001) and
limits light availability for benthic primary producers (Le Fur et al., 2019, 2018). Ultimately, sustained
eutrophication of coastal lagoons can cause a regime shift from a primary producer community
dominated by perennial macrophytes presenting different morphotypes, to a community dominated by
free-floating and opportunistic macroalgae like *Ulva rigida* and finally, to a phytoplankton-dominated
community (Le Fur et al., 2019). The negative correlation between several eutrophication variables (*e.g.*
mean [Chla], mean turbidity) and β$_{MM}$ highlights the simplification of benthic habitats caused by lagoon
eutrophication (Cloern, 2001; Le Fur et al., 2019, 2018) that shift from a mosaic of different macrophyte
morphotypes to a sediment void of macrophytes or with one macrophyte morphotype. Overall, the
multiple interconnections between natural environmental variability and eutrophication likely explain the
high amount of macrobenthic community and index variance jointly attributed to both effects, as
underlined in the conceptual model developed by Cloern (2001).
The unique effect of eutrophication variables on each taxonomic component also informs us about
which of these variables likely have a direct effect on macrobenthic invertebrates without being
modulated by natural variability. Indeed, benthic organisms can be affected by eutrophication directly
through the toxicity of inorganic nitrogenous compounds like ammonium, nitrites and nitrates (Jessen et
al., 2015). Our results indicate that high NH$_4^+$ concentrations seem to have a very small negative effect on
total macrofauna density while NO$_x$ toxicity could explain the strong unique effect of eutrophication on
SR and M-AMBI, and both NH$_4^+$ and NO$_x$ toxicity could be responsible for the very weak unique effect of
eutrophication on macrobenthic communities. These results are probably also linked to indirect effects
high eutrophication levels can have on macrobenthic communities via increased water turbidity and
sediment bio-geochemistry modifications (*e.g.* anoxic superficial sediments (Zilius et al., 2015)). Indeed,
water NH$_4^+$ and NO$_x$ concentrations in excess can be the sign of highly eutrophic lagoons (Souchu et al.,
2010). Overall, experimental work on a few well-selected species and populations sampled in lagoons
presenting different levels of natural variability would greatly improve our understanding of the effect of
eutrophication on lagoon benthic invertebrates via inorganic nitrogenous compound toxicity, as
previously done for freshwater and marine organisms (Alonso and Camargo, 2006; Boardman et al.,
2004; Camargo et al., 2005).
Furthermore, mean [Chla] appears to have a strong negative effect on macrobenthic communities by
impacting the diversity of low and intermediate abundance taxa (SR and H') and this effect is at least
partly independent from the selected natural variability variables (*i.e.* mean temperature, mean salinity
for SR and lagoon-sea connection, β$_{MM}$ for H'). Two non-exclusive hypotheses could explain this result:
small-scale habitat homogenization and hypoxic stress. Despite not being the appropriate index to
investigate beta diversity, the standard deviation of SR at the intra-station scale was significantly
correlated negatively to mean [Chla], which seems to point towards a homogenization of benthic
communities at the decimeter scale linked to eutrophication. The loss of benthic taxa presenting
restricted environmental niches and dependent upon small-scale habitat diversity for reproduction,
recruitment, refuge and/or food (Brauns et al., 2007; Harman, 1972; Stendera and Johnson, 2008) could
explain this result. Secondly, low to zero dissolved oxygen (*i.e.* hypoxia and anoxia) is another well-known
symptom of eutrophication in transitional waters like coastal lagoons (Chapelle et al., 2001; Cladas et al.,
2016; Vignes et al., 2010). Oxygen depletion generally takes place during the summer and is maximal at
the end of the night/early morning (Newton et al., 2010), often starting near the bottom because of

benthic organic matter mineralization (*e.g.* sedimented phytoplankton, dead macroalgae) and low water
renewal (Rigaud et al., 2021; Souchu et al., 1998). In the worst cases, these low to zero oxygen zones can
spread to the entire water column and last for several days (Chapelle et al., 2001; Cladas et al., 2016), as
seen during the 2018 dystrophic crisis in Thau lagoon (Derolez et al., 2020a). These crises are known to
impact macrobenthic community first in terms of abundance and then in terms of composition and
species richness depending on the sensitivity of each species to reduced oxygen (Díaz and Rosenberg,
1995; Riedel et al., 2012). Consequently, lagoon macrobenthic invertebrates are often reported as being
less abundant and less diverse in fall than in spring because of summer hypoxic stress (Barbone et al.,
2012; Basset et al., 2012; Vanda Orro and Cabana, 2021). The negative impact of eutrophication via
hypoxic/anoxic stress on lagoon macrobenthic communities is suggested by the RDA results where high
mean O_2 sat (*i.e.* hyper-saturation linked to strong pelagic primary production) is associated to group A
stations that present significantly lower species richness than group B stations. Nonetheless, oxygen
related variables were not selected in the other models, a result probably linked to when oxygen was
measured (late morning) and at what depth it was measured (generally sub-surface). Indeed, these
features limit our ability to detect potential short and long-term effects of hypoxic stress on
macrobenthic communities. Bottom hypoxia/anoxia is also known to predominantly take place in deep
lagoons like Berre and Thau (Derolez et al., 2020a; Rigaud et al., 2021; Souchu et al., 1998), where wind
derived water mixing is limited by depth. Nonetheless, anoxic crises inside coastal lagoons are expected
to increase in frequency and intensity with climate change (Derolez et al., 2020a; Miyamoto et al., 2019).
Consequently, it appears paramount to (*i*) study the sensitivity of lagoon benthic organisms to low
dissolved oxygen, not only of freshwater and marine taxa (Díaz and Rosenberg, 1995; Landman et al.,
2005) and (*ii*) monitor more closely summer bottom dissolved oxygen concentrations, especially at higher
frequency (O_2 being very variable) and at temporal scales relevant to these organisms (*i.e.* across several
days starting when bottom hypoxia is detected (Riedel et al., 2012)).
Overall, lagoon hydro-morphology (connection to the sea, surface and depth) which affects the level
of environmental variability (salinity and temperature), as well as lagoon-scale benthic habitat diversity
(β_{MM}) seem to regulate the distribution of macrobenthic species (Bachelet et al., 2000), while
eutrophication and associated stressors like low dissolved oxygen act upon the existing communities,
mainly by reducing taxa richness and diversity (SR and H').
797
The M-AMBI and recommendations for its use as an indicator for Mediterranean coastal lagoons
With this study, we investigate for the first time the joint sensitivity of the M-AMBI index to natural
and anthropogenic stressors across 29 lagoons covering most of the French Mediterranean coastline. The
M-AMBI (Muxika et al., 2007) is currently used to evaluate the WFD ecological state of poly- and eu-
haline lagoons based on soft sediment benthic invertebrates (Reizopolou et al., 2018). This index has
previously been shown to be sensitive to pressures related to eutrophication, like concentration in
chlorophyll *a* and total nitrogen (Derolez et al., 2014) but it is also known to be sensitive to natural
environmental variability (*e.g.* water salinity) and lagoon hydro-morphology (*e.g.* surface and depth)
(Barbone et al., 2012).
Our results show that the M-AMBI appears to be a sensitive indicator of eutrophication through the
species richness and Shannon diversity components of the index but not the AMBI one. First, the strong
correlation between M-AMBI and mean [Chla] was linked only to the significant correlation between SR
and mean [Chla] and between H' and mean [Chla] while AMBI was not significantly linked to most
eutrophication related variables (except mean [NO_x]) nor to chlorophyll *a* concentration, a common
eutrophication indicator (Bec et al., 2011; Cloern, 2001). Importantly, these linear relations remained
significant once the effect of identified covariables was removed. Regarding AMBI, its current application
in transitional waters like coastal lagoons suffers from methodological limitations. Indeed, this index is
based on macroinvertebrate sensitivity/tolerance to organic matter enrichment following Pearson and
Rosenberg (1978) and was developed for coastal waters (Borja et al., 2000), not for coastal lagoons,
which are known to sometimes present sediments naturally enriched in organic matter linked to the
decay of meadow-forming macrophytes (Bachelet et al., 2000), watershed inputs and the overall low
organic matter exportation (except during floods) (Perthuisot and Guelorget, 1983). These results
highlight the limits of indirectly including AMBI in the evaluation of the ecological state of Mediterranean
coastal lagoons. Nonetheless, its absence of sensitivity to natural variability and lagoon hydro-

morphology suggests an index based on redefined ecological groups could be promising (Robertson et al., 2016), especially if we aim towards a more specific macrofauna-based index to evaluate eutrophication. Overall, increasing the robustness of M-AMBI would likely necessitate additional work to classify lagoon macroinvertebrates according to other eutrophication related stressors like low oxygen, nitrogenous compound toxicity and/or hydrogen sulfide toxicity.

Secondly, M-AMBI variability was two times more strongly attributed uniquely to eutrophication related variables than uniquely to natural variability ones, with nonetheless a very strong joint effect of the two types of variables. This strong joint effect is probably related, as previously discussed, to the same interconnections between lagoon-sea connection that buffers eutrophication, temperature that promotes phytoplankton blooms and eutrophication that homogenizes benthic habitats. Several modifications could help limit the sensitivity of M-AMBI to natural variability: (*i*) the replacement of the currently used poly- and eu-haline vs oligo- and meso-haline categories (Provost et al., 2012; Reizopolou et al., 2018) by one based on the three lagoon-sea connection levels that strongly influence the M-AMBI or on another typology like the choked vs restricted typology (Barbone et al., 2012; Basset et al., 2013), with the definition of new reference lagoons; (*ii*) the replacement of a sampling done with an Ekman-Birge grab by a diver-operated sampling in order to verify that only unvegetated sediments are being collected and to note the presence and type of macrophyte near the sampling site (*e.g.* 10 meter radius). This last modification would likely help reduce the effect of local benthic habitat heterogeneity on H' and M-AMBI (Magni and Gravina, 2023; de Paz et al., 2008).

Finally, several studies have underlined the limits of using the M-AMBI based on abundance to evaluate the ecological state of transitional ecosystems like coastal lagoons, as opportunistic and tolerant species are generally numerically dominant in these systems (Mistri et al., 2018; Pitacco et al., 2019; Puente and Diaz, 2008). Other indicators based on the size spectra and biomass of benthic macroinvertebrates like the biomass based M-AMBI (M-bAMBI (Mistri et al., 2018)) and the index of size distribution (ISD (Reizopoulou and Nicolaidou, 2007)) have showed promising results and applying these indices to French coastal lagoons would represent an additional trial in the context of historically highly eutrophic lagoons currently undergoing a re-oligotrophication trajectory (Derolez et al., 2020a, 2020b; Le Fur et al., 2019; Leruste et al., 2016). Another interesting avenue would be to investigate more closely the spatial scales at which changes in macroinvertebrate abundance and diversity can be detected and link these spatial changes to localized anthropogenic disturbances like heavy metals (Magalhães et al., 2019) while also considering small-scale biotic interactions such as competition, predation and facilitation. Indeed, sediment contaminants like biotic interactions are likely to affect benthic macrofauna at fine spatial scales (*ca.* 1 km) that are more relevant to low mobility macroinvertebrates (Berthelsen et al., 2018; Brauko et al., 2020, 2015; Carvalho et al., 2006).

In a nutshell, to improve the robustness of M-AMBI, we call for a revision of the ecological groups at the base of the AMBI index and of the sampling protocol (*e.g.* using divers) along with a re-evaluation of the lagoon typology. More generally, it appears key to increase our understanding of the effects of other disturbances like hypoxia and heavy metals on lagoon invertebrates and to reevaluate the small-scale distribution of benthic organisms in French coastal lagoons, more than 30 years after the historical studies by Amanieu et al. (1977), Guelorget et al. (1994) and Guelorget and Michel (1979).

Overcoming taxonomic limitations

A large part of the identified macroinvertebrates had a very restricted geographical range, preventing us from building a general species-based framework linking eutrophication and macrobenthic communities. Indeed, we found a very high level of lagoon-specificity for benthic macroinvertebrates across French Mediterranean coastal lagoons with no taxon present across all 41 stations, 40% of the 227 taxa identified in only one station and only 1.3% recorded in over half of the stations. This result is probably explained by environmental filtering resulting from very contrasted conditions across the different sampled systems, passive larval diffusion (Basset et al., 2007) and the randomness of the colonization process by species of marine origin (Pérez-Ruzafa et al., 2011b), a phenomenon previously observed for benthic macroinvertebrates in Italian coastal lagoons (Basset et al., 2007), fish in Atlantic and Mediterranean coastal lagoons (Pérez-Ruzafa et al., 2007b) and macrophytes in estuaries and lagoons (Pérez-Ruzafa et al., 2011c). Consequently, unique assemblages of benthic macroinvertebrates are present in each lagoon and even at smaller spatial scales (*i.e.* stations inside a lagoon), which could

explain why we were able to explain only 22% of the macrobenthic community variability. This high level
of lagoon-specificity present across multiple biological compartments in coastal lagoons warrants the
need to go beyond a species-based approach and towards a trait-based approach (Lacoste et al., 2023),
as previously outlined using macrofauna size (Basset et al., 2008; Brauko et al., 2020; Sigala et al., 2012).
Such an approach could build upon the saprobity concept (Tagliapietra et al., 2012) to select relevant
biological traits and ultimately help us build a more general framework around macrobenthic community
assembly rules in the line of what Pérez-Ruzafa et al. (2013) did for fish by revisiting the r and K-
strategies. Nonetheless, for such an approach to be relevant, we also need to increase our knowledge of
the life-cycle of lagoon benthic invertebrates (Brauko et al., 2020) and to create a harmonized
Mediterranean wide trait database for these organisms (Lacoste et al., 2023).
Another interesting avenue to pursue could be to investigate intra-specific biological trait and/or
physiological response variability (*i.e.* oxydative stress) of species like the polychaete *Hediste diversicolor*
along eutrophication gradients. Indeed, this species is known to display a great diversity of life history
characteristics along with morphological, biochemical and physiological differences according to local
environmental conditions (Scaps, 2002). For example, the feeding behavior and diet of this species has
been shown to vary according to sewage pollution levels across English estuaries (Aberson et al., 2016).
Another study has related oxydative stress of the polychaete *Nephtys cirrosa* to seasonal variations of
abiotic parameters like temperature and salinity in the Ria de Aveiro (Magalhães et al., 2019).
Furthermore, *H. diversicolor,* a species we found to be associated to eutrophic lagoons, is reported to
present a great diversity of feeding modalities (*i.e.* predator, grazer, suspension-feeder and deposit-
feeder (Faulwetter et al., 2014)), whereas the species associated to oligotrophic lagoons, like *Loripes
orbiculatus,* present less diverse feeding modes (*i.e.* mainly based on sulfo-oxydizing bacteria (Roques et
al., 2020)). This flexibility could allow organisms to better tolerate the fluctuating abiotic conditions that
characterize small eutrophic lagoons, stressing that environmental stability is probably a strong driver of
macrobenthic communities in coastal lagoons, as previously shown for macrophytes (Pérez-Ruzafa et al.,
2011c). Overall, the plasticity several species display in environmentally constraining conditions stresses
the difficulty to implement trait-based approaches in ultra-variable environmental systems like coastal
lagoons, where realized trait modalities may differ significantly from the ones available in the literature,
commonly acquired from non-transitional coastal ecosystems (Martini et al., 2021). Finally, a functional
approach based on well-informed biological traits and biomass data could help overcome the identified
taxonomic limitations and build a more holistic framework linking natural variability and eutrophication
of coastal lagoons to macrobenthic communities.

Conclusion

Coastal lagoons are transitional ecosystems naturally characterized by a strong variability in abiotic
parameters like salinity and temperature. Superimposed onto this natural environmental variability are
numerous anthropogenic disturbances, eutrophication being a historical one targeted by the European
Water Framework Directive. By jointly analyzing water column, sediment and macrophyte parameters,
our study shows that macrobenthic communities across 29 French Mediterranean coastal lagoons are
primarily determined by (*i*) lagoon hydro-morphology (connection level with the sea, surface and depth)
which further affects natural environmental variability in terms of salinity and temperature, and (*ii*)
lagoon-scale benthic habitat diversity, as estimated through macrophyte morphotypes. Eutrophication
and associated stressors like low dissolved oxygen act upon the existing communities by reducing
taxonomic richness and diversity. Eutrophication appeared to act on lagoon macrobenthos through
relatively direct mechanisms like toxicity of inorganic nitrogenous compounds, as well as indirectly via for
example, the simplification of benthic habitats and sediment bio-geochemistry modifications.
Furthermore, our study highlights the complex interconnections between lagoon hydro-morphology,
natural environmental variability and eutrophication and their joint effect on benthic communities.
Indeed, lagoon-sea connection modulates the susceptibility of each lagoon to eutrophication, high water
temperatures promote eutrophication related phytoplankton blooms while eutrophication reduces
benthic habitat heterogeneity, all of which affect benthic communities. Finally, benthic taxa richness and
diversity also proved to be indicators sensitive to eutrophication in coastal lagoons while an AMBI like

index based on redefined ecological groups for coastal lagoons, which can be naturally enriched in
organic matter, has the potential of being a specific eutrophication indicator.
929
930
931
932

933 **Appendices**

934 **Appendix 1 - Procedure used to calculate the beta diversity of macrophyte morphotypes (β_{MM})**
935 **corresponding to each macrofauna station**
936
Step 1: Select the macrophyte stations corresponding to each macrofauna station
• If one macrofauna station is considered in a given lagoon, then all the macrophyte stations of
that given lagoon are taken into account.
• If two or three macrofauna stations are considered in a given lagoon, then all the macrophyte
stations attributed by the WFD protocol to the same section of the lagoon as the macrofauna
station are taken into account.
943
Step 2 : Build the macrophyte station by macrophyte morphotype matrix for each macrofauna station
For each macrophyte station of the dataset:
• If the macrophyte cover is below 25% or equal to 25%, a value of 1 is entered in the sediment
column and 0 in all the others (line 1 of the example matrix below).
• If the cover is above 25%, the dominant or co-dominant taxa are determined based on the taxa
specific cover data or the taxa specific biomass data, then these taxa are recoded into a
morphotype (see Appendix 2) and finally, a value of 1 is entered in all the cells corresponding to
the given morphotypes of the dominant or co-dominant taxa and a value of 0 is entered in the
remaining cells (lines 2 and 3 of the example matrix below).
953
954 *Step 3 : Calculate the index value corresponding to each macrofauna station*
955 • Calculate the Bray-Curtis dissimilarity between all the station pairs using the previously built
956 station by macrophyte morphotype matrix (presence/absence)
957 • Average the Bray-Curtis dissimilarities
958
959 Example of a macrophyte station by macrophyte morphotype matrix used to calculate the β_{MM}

Station	mushroom	feather	hair	ball	branched	sheet	meadow	sediment
1	0	0	0	0	0	0	0	1
2	0	0	0	0	1	0	1	0
3	0	0	0	0	0	1	0	0
...								
n-1								
n								

976 **Appendix 2 - Correspondence between macrophyte taxa and morphotypes**

Macrophyte taxa	Morphotype
Acetabularia acetabulum	mushroom
Aglaothamnion pseudobyssoides *Alsidium corallinum* *Cladophora vagabunda* *Dasyaceae* *Spyridia filamentosa*	feather
Chaetomorpha aerea *Ulva clathrata*	hair
Cladophora glomerata *Valonia aegagropila*	ball
Gracilaria dura *Gracilaria gracilis* *Halopithys incurva*	branched
Ulva rigida	sheet
Cymodocea nodosa *Ruppia cirrhosa* *Stuckenia pectinata* *Zostera marina* *Zostera noltei*	meadow

977
Appendix 3
Mean ± SD of the species richness (SR), Shannon diversity (H'), inverse of Simpson dominance index (N2),
macrofauna density, AMBI (AZTI's Marine Biotic Index) and M-AMBI (Multivariate AMBI) calculated for
the A and B station groups determined using a K-means partitioning based on the benthic macrofauna
Hellinger-transformed mean abundances. The results of the non-parametric Kruskal-Wallis rank sum test
comparing the simple and multimetric macrobenthic taxonomic indices between the A and B groups are
also indicated.

	Group A	Group B	Kruskal-Wallis rank sum test
SR	9.30 ± 3.48	17.48 ± 11.41	$\chi^2(1) = 6.41$, $p = 0.011$
H'	2.00 ± 0.57	2.36 ± 0.69	$\chi^2(1) = 2.29$, $p = 0.13$
N2	3.30 ± 1.35	3.83 ± 1.76	$\chi^2(1) = 0.74$, $p = 0.39$
Macrofauna density (ind.m^{-2})	2 280 ± 1 372	5 113 ± 5 205	$\chi^2(1) = 0.98$, $p = 0.32$
AMBI	3.03 ± 0.53	2.87 ± 1.08	$\chi^2(1) = 0.01$, $p = 0.92$
M-AMBI	0.43 ± 0.08	0.53 ± 0.14	$\chi^2(1) = 5.76$, $p = 0.016$

Appendix 4
Coordinates (biplot scores for quantitative variables and centroids for lagoon-sea connection) on the first
and second axis of the redundancy analysis (RDA) plots, marginal effects and conditional effects of the 8
environmental variables selected to explain the macrobenthic community variability. After the
abbreviated name of each environmental variable is indicated the type of transformation used to
normalize it ($\log_e$: natural logarithm and sqrt: square-root). [Chla] and [DIP] is the water concentration in
chlorophyll a ($\mu g.L^{-1}$) and dissolved inorganic phosphorous ($\mu mol.L^{-1}$), respectively. Temp (°C) is the
lagoon water temperature. O_2 sat (%) is the water oxygen saturation level. OMC (%) is the sediment
organic matter content. CV stands for coefficient of variation.

Selected environmental variable	RDA1	RDA2	Marginal effect (%)	Conditional effect (%)
Mean [Chla] ($\log_e$)	-0.76	0.23	7.22	7.22
Lagoon surface ($\log_e$)	0.73	-0.15	5.90	1.69
Weak lagoon-sea connection	-0.10	0.49		
Moderate lagoon-sea connection	-0.073	-0.18	5.30	4.64
Strong lagoon-sea connection	0.18	-0.29		
Mean Temp	-0.61	-0.24	5.04	3.62
Mean [DIP] ($\log_e$)	-0.51	0.092	3.16	1.31
CV Temp (sqrt)	0.44	0.15	2.20	1.53
OMC	0.11	0.27	0.50	1.20
Mean O_2 sat ($\log_e$)	-0.16	-0.063	0.47	1.28

Appendix 5
Partial redundancy analysis plots (pRDA, axis 1 and 2) focusing on (a) the benthic macrofauna sampled
across 41 stations located in 29 lagoons along the French Mediterranean coast (see Table 1 for the full
lagoon names and corresponding stations) as explained by keeping constant the natural environmental
variability variables (lagoon-sea connection, lagoon surface, mean and CV water temperature, mean
oxygen saturation and sediment organic matter content) and letting the eutrophication variables vary
(mean concentration in chlorophyll a (mean [Chla]) and mean dissolved inorganic phosphorous (mean
[DIP]), (b) the environmental variables left to vary where mean [Chla] and mean [DIP] are $\log_e$
transformed (see Tables 1 and 2 for the full environmental variable names and values) and (c) the

macrofauna taxa, where only taxa with a goodness of fit (cumulative proportion of variance explained by the first two axis) above 0.23 are plotted. In (a), inter-station distances are euclidean while in (b) and (c) angles between all vectors reflect linear correlations. Across all plots, each station (point) is represented using a symbol for its salinity type (triangle: oligo- and meso-haline and circle: poly- and eu-haline) and a color (red: group A and blue: group B) for its group membership (K-means partitioning) based on Hellinger-transformed mean abundances.

1042
Appendix 6

Partial effect of mean chlorophyll *a* concentration ([Chla]) on (a) species richness (SR) once the effect of
mean temperature and mean salinity has been removed, (b) Shannon diversity (H') once the effect of
lagoon-sea connection, CV [TN], CV [TN]^2 and β_{MM} has been removed and (c) M-AMBI once the effect of
lagoon-sea connection and CV [NO$_x$] has been removed (see Table 3 for the parsimonious linear models)
across 41 stations located in 29 lagoons along the French Mediterranean coast (see Table 1 for the full
lagoon names and corresponding stations). The dashed lines correspond to simple linear regression
models and the shaded zone to the confidence interval. Each station (point) is represented using a
symbol for its salinity type (triangle: oligo- and meso-haline and circle: poly- and eu-haline) and a color
(red: group A and blue: group B) for its group membership (K-means partitioning) based on Hellinger-
transformed mean abundances.
1054
1055

Acknowledgements

The authors would like to thank all the members of the LERLR and LERPAC Ifremer labs as well as all the managers of the different French Mediterranean lagoons for sampling and analyzing the water column parameters, the sediment parameters and the macrofauna. A special thanks to Corinne Pélaprat (STARESO), Sébastien Thorin (CREOCEAN), Thibault Schwartz (CREOCEAN) and Daniel Martin (Centre d'Estudis Avançats de Blanes – CEAB) for sampling, identifying, and analyzing the macrofauna. The authors are also extremely grateful to Aurélie Fouveau (Ifremer, LERBN) for extracting the ecological group values, Olivier Boutron (Tour du Valat) for the information on the Vaccarès lagoon system and Marion Richard (Ifremer, LERLR) for the fruitful discussions. Finally, we would like to thank all our close colleagues from the LERLR (Ophélie Serais, Elise Bellamy, Aurélien Bouquet, Elise Caillard, Nicolas Cimiterra, Christine Felix, Elise Foucault, Camille Gianaroli, Yannick Gueguen, Elise Lacoste, Franck Lagarde, Julie Le Ray, Danièle Martin, Grégory Messian, Dominique Munaron, Hervé Violette) for creating such a cheerful, dynamic, and enthusiastic work environment, despite all the COVID-19 ups and downs.

Data, scripts, code, and supplementary information availability

Data, scripts, and code are available online on Zenodo: https://doi.org/10.1101/2022.08.18.504439
Supplementary information is available online on biorxiv as part of the preprint: https://www.biorxiv.org/content/10.1101/2022.08.18.504439v1

Conflict of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

Funding

This work is part of the MALAG project (Effect of eutrophication on the biodiversity of Mediterranean coastal lagoon benthic macrofauna) funded by "Office Français de la Biodiversité" and Ifremer. A. G. J., M. C. and T. L. M. received funding through the MALAG project. All the data presented in this study was collected during the projects "Réseau de Suivi Lagunaire" and Water Framework Directive, which received financial support from Ifremer, "Agence de l'Eau Rhône Méditerranée Corse", "Région Languedoc-Roussillon/Occitanie" and "Cépralmar".

References

Aberson, M.J.R., Bolam, S.G., Hughes, R.G., 2016. The effect of sewage pollution on the feeding behaviour and diet of *Hediste* (*Nereis diversicolor* (O.F. Müller, 1776)) in three estuaries in south-east England, with implications for saltmarsh erosion. Marine Pollution Bulletin 105, 150–160. https://doi.org/10.1016/j.marpolbul.2016.02.033

Alonso, A., Camargo, J.A., 2006. Toxicity of nitrite to three species of freshwater invertebrates. Environmental Toxicology 21, 90–94. https://doi.org/10.1002/tox.20155

Amanieu, M., Guelorget, O., Michel, P., 1977. Richesse et diversité de la macrofaune benthique d'une lagune littorale méditerranéenne. Vie et Milieu XXVII, 85–109.

Andral, B., Sargian, P., 2010a. Directive Cadre Eau. District Rhône et Côtiers méditerranéens. Contrôle de surveillance/opérationnel. Campagne DCE 2009. RST.DOP/LER-PAC/10-19. https://archimer.ifremer.fr/doc/00038/14923/

Andral, B., Sargian, P., 2010b. Directive Cadre Eau. District Corse. Contrôle de surveillance/opérationnel. Campagne DCE 2009. RST.DOPLER/PAC/10-20. https://archimer.ifremer.fr/doc/00028/13918/

Anonymous, 1959. Symposium on the classification of brackish waters. Presented at the Archivio di
Oceanografia e Limnologia, Venice, 8–14 April 1958.
Arocena, R., 2007. Effects of submerged aquatic vegetation on macrozoobenthos in a coastal lagoon of
the southwestern atlantic. International Review of Hydrobiology 92, 33–47.
https://doi.org/10.1002/iroh.200610881
Bachelet, G., de Montaudouin, X., Auby, I., Labourg, P.-J., 2000. Seasonal changes in macrophyte and
macrozoobenthos assemblages in three coastal lagoons under varying degrees of eutrophication. ICES
Journal of Marine Science 57, 1495–1506. https://doi.org/10.1006/jmsc.2000.0902
Barbone, E., Basset, A., 2010. Hydrological constraints to macrobenthic fauna biodiversity in transitional
waters ecosystems. Rend. Fis. Acc. Lincei 21, 301–314. https://doi.org/10.1007/s12210-010-0090-4
Barbone, E., Rosati, I., Reizopoulou, S., Basset, A., 2012. Linking classification boundaries to sources of
natural variability in transitional waters: A case study of benthic macroinvertebrates. Ecological
Indicators, Marine Benthic Indicators 12, 105–122. https://doi.org/10.1016/j.ecolind.2011.04.014
Barnes, R.S.K., 1980. Coastal lagoons: the natural history of a neglected habitat, Cambridge University
Press. ed. Cambridge.
Basset, A., Barbone, E., Borja, Á., Elliott, M., Jona-Lasinio, G., Marques, J., Mazik, K., Muxika, I., Neto, J.,
Reizopoulou, S., Rosati, I., Teixeira, H., 2012. Natural variability and reference conditions: setting type-
specific classification boundaries for lagoon macroinvertebrates in the Mediterranean and Black Seas.
Hydrobiologia. https://doi.org/10.1007/s10750-012-1273-z
Basset, A., Barbone, E., Elliott, M., Li, B.-L., Jorgensen, S.E., Lucena-Moya, P., Pardo, I., Mouillot, D., 2013.
A unifying approach to understanding transitional waters: Fundamental properties emerging from
ecotone ecosystems. Estuarine, Coastal and Shelf Science, Estuarine and lagoon biodiversity and their
natural goods and services 132, 5–16. https://doi.org/10.1016/j.ecss.2012.04.012
Basset, A., Galuppo, N., Sabetta, L., 2007. Environmental heterogeneity and benthic macroinvertebrate
guilds in italian lagoons. Transitional Waters Bulletin 1, 48–63.
https://doi.org/10.1285/i1825226Xv1n1p48
Basset, A., Sabetta, L., Fonnesu, A., Mouillot, D., Do Chi, T., Viaroli, P., Giordani, G., Reizopoulou, S.,
Abbiati, M., Carrada, G.C., 2006. Typology in Mediterranean transitional waters: new challenges and
perspectives. Aquatic Conservation: Marine and Freshwater Ecosystems 16, 441–455.
https://doi.org/10.1002/aqc.767
Basset, A., Sabetta, L., Sangiorgio, F., Pinna, M., Migoni, D., Fanizzi, F., Barbone, E., Galuppo, N., Fonda
Umani, S., Reizopoulou, S., Nicolaidou, A., Arvanitidis, C., Moncheva, S., Trajanova, A., Georgescu, L.,
Beqiraj, S., 2008. Biodiversity conservation in Mediterranean and Black Sea lagoons: a trait-oriented
approach to benthic invertebrate guilds. Aquatic Conservation: Marine and Freshwater Ecosystems
18, S4–S15. https://doi.org/10.1002/aqc.979
Bec, B., Collos, Y., Souchu, P., Vaquer, A., Lautier, J., Fiandrino, A., Benau, L., Orsoni, V., Laugier, T., 2011.
Distribution of picophytoplankton and nanophytoplankton along an anthropogenic eutrophication
gradient in French Mediterranean coastal lagoons. Aquatic Microbial Ecology 63, 29–45.
https://doi.org/10.3354/ame01480
Bendoricchio, G., Di Luzio, M., Baschieri, P., Capodaglio, A.G., 1993. Diffuse pollution in the lagoon of
Venice. Water Science and Technology 28, 69–78. https://doi.org/10.2166/wst.1993.0405
Benjamini, Y., Hochberg, Y., 1995. Controlling the false discovery rate: a practical and powerful approach
to multiple testing. Journal of the Royal Statistical Society. Series B (Methodological) 57, 289–300.
https://doi.org/10.1111/j.2517-6161.1995.tb02031.x
Berthelsen, A., Atalah, J., Clark, D., Goodwin, E., Patterson, M., Sinner, J., 2018. Relationships between
biotic indices, multiple stressors and natural variability in New Zealand estuaries. Ecological Indicators
85, 634–643. https://doi.org/10.1016/j.ecolind.2017.10.060
Blanchet, F.G., Legendre, P., Borcard, D., 2008. Forward selection of explanatory variables. Ecology 89,
2623–2632. https://doi.org/10.1890/07-0986.1
Boardman, G.D., Starbuck, S.M., Hudgins, D.B., Li, X., Kuhn, D.D., 2004. Toxicity of ammonia to three
marine fish and three marine invertebrates. Environmental Toxicology 19, 134–142.
https://doi.org/10.1002/tox.20006

Borja, A., Franco, J., Pérez, V., 2000. A marine biotic index to establish the ecological quality of soft-bottom benthos within European estuarine and coastal environments. Marine Pollution Bulletin 40, 1100–1114. https://doi.org/10.1016/S0025-326X(00)00061-8

Borja, A., Muxika, I., 2005. Guidelines for the use of AMBI (AZTI's Marine Biotic Index) in the assessment of the benthic ecological quality. Marine Pollution Bulletin 50, 787–789. https://doi.org/10.1016/j.marpolbul.2005.04.040

Boström, C., Jackson, E.L., Simenstad, C.A., 2006. Seagrass landscapes and their effects on associated fauna: A review. Estuarine, Coastal and Shelf Science, Ecological and Management Implications on Seagrass Landscapes 68, 383–403. https://doi.org/10.1016/j.ecss.2006.01.026

Brauko, K.M., Cerveira, A.F., Freitas, R.F., Hadlich, H.L., Beras, T., Fonseca, A.L., Pagliosa, P.R., 2020. Comparing structural and functional approaches to distinguish benthic communities from eutrophic and non-eutrophic estuaries. Marine Pollution Bulletin 157, 111290. https://doi.org/10.1016/j.marpolbul.2020.111290

Brauko, M.K., de Souza, F.M., Muniz, P., de Camargo, M.G., Lana, P. da C., 2015. Spatial variability of three benthic indices for marine quality assessment in a subtropical estuary of Southern Brazil. Marine Pollution Bulletin 91, 454–460. https://doi.org/10.1016/j.marpolbul.2014.10.025

Brauns, M., Garcia, X.-F., Walz, N., Pusch, M.T., 2007. Effects of human shoreline development on littoral macroinvertebrates in lowland lakes. Journal of Applied Ecology 44, 1138–1144. https://doi.org/10.1111/j.1365-2664.2007.01376.x

Breslow, N., 1970. A generalized Kruskal-Wallis test for comparing k samples subject to unequal patterns of censorship. Biometrika 57, 579–594. https://doi.org/10.2307/2334776

Caliceti, M., Argese, E., Sfriso, A., Pavoni, B., 2002. Heavy metal contamination in the seaweeds of the Venice lagoon. Chemosphere 47, 443–454. https://doi.org/10.1016/S0045-6535(01)00292-2

Camargo, J.A., Alonso, A., Salamanca, A., 2005. Nitrate toxicity to aquatic animals: a review with new data for freshwater invertebrates. Chemosphere 58, 1255–1267. https://doi.org/10.1016/j.chemosphere.2004.10.044

Cardoso, P.G., Pardal, M.A., Raffaelli, D., Baeta, A., Marques, J.C., 2004. Macroinvertebrate response to different species of macroalgal mats and the role of disturbance history. Journal of Experimental Marine Biology and Ecology 308, 207–220. https://doi.org/10.1016/j.jembe.2004.02.018

Carlier, A., Riera, P., Amouroux, J.-M., Bodiou, J.-Y., Escoubeyrou, K., Desmalades, M., Caparros, J., Grémare, A., 2007. A seasonal survey of the food web in the La Palme lagoon (northwestern Mediterranean) assessed by carbon and nitrogen stable isotope analysis. Estuarine, Coastal and Shelf Science 73, 299–315. https://doi.org/10.1016/j.ecss.2007.01.012

Carvalho, S., Gaspar, M.B., Moura, A., Vale, C., Antunes, P., Gil, O., da Fonseca, L.C., Falcão, M., 2006. The use of the marine biotic index AMBI in the assessment of the ecological status of the Obidos lagoon (Portugal). Marine Pollution Bulletin 52, 1414–1424. https://doi.org/10.1016/j.marpolbul.2006.04.004

Chapelle, A., Lazure, P., Souchu, P., 2001. Modélisation numérique des crises anoxiques (malaïgues) dans la lagune de Thau (France). Oceanologica Acta Supplement 1, 87–97. https://doi.org/10.1016/S0399-1784(00)01109-9

Cladas, Y., Papantoniou, G., Bekiari, V., Fragkopoulu, N., 2016. Dystrophic crisis event in Papas Lagoon, Araxos Cape, Western Greece in the summer 2012. Mediterranean Marine Science 17, 32–38. https://doi.org/10.12681/mms.1409

Cloern, J.E., 2001. Our evolving conceptual model of the coastal eutrophication problem. Marine Ecology Progress Series 210, 223–253. https://doi.org/10.3354/meps210223

Cognetti, G., Maltagliati, F., 2000. Biodiversity and adaptive mechanisms in brackish water fauna. Marine Pollution Bulletin 40, 7–14. https://doi.org/10.1016/S0025-326X(99)00173-3

Colden, A.M., Fall, K.A., Cartwright, G.M., Friedrichs, C.T., 2016. Sediment suspension and deposition across restored oyster reefs of varying orientation to flow: implications for restoration. Estuaries and Coasts 39, 1435–1448. https://doi.org/10.1007/s12237-016-0096-y

CORINE Land Cover [WWW Document], 2018. https://land.copernicus.eu/pan-european/corine-land-cover (accessed 12.14.21).

De Casabianca, M.-L., 1996. France — The Mediterranean lagoons, in: Schramm, W., Nienhuis, P.H. (Eds.), Marine benthic vegetation: Recent changes and the effects of eutrophication, Ecological Studies. Springer, Berlin, Heidelberg, pp. 307–329. https://doi.org/10.1007/978-3-642-61398-2_14

de Paz, L., Patrício, J., Marques, J.C., Borja, A., Laborda, A.J., 2008. Ecological status assessment in the lower Eo estuary (Spain). The challenge of habitat heterogeneity integration: a benthic perspective. Marine Pollution Bulletin 56, 1275–1283. https://doi.org/10.1016/j.marpolbul.2008.04.027

Derolez, V., Bec, B., Munaron, D., Fiandrino, A., Pete, R., Simier, M., Souchu, P., Laugier, T., Aliaume, C., Malet, N., 2019. Recovery trajectories following the reduction of urban nutrient inputs along the eutrophication gradient in French Mediterranean lagoons. Ocean & Coastal Management 171, 1–10. https://doi.org/10.1016/j.ocecoaman.2019.01.012

Derolez, V., Fiandrino, A., Munaron, D., Cadoret, M., 2014. Bilan sur les principales pressions pesant sur les lagunes méditerranéennes et leurs liens avec l'état DCE. RST-LER/LR 14-20. https://archimer.ifremer.fr/doc/00254/36574/

Derolez, V., Leurion, A., Fiandrino, A., 2012. Directive Cadre Eau. Estimation du degré d'échange avec la mer pour les lagunes du bassin Rhône Méditerranée & Corse. Agence de l'eau, Ref. RST. DOP/LER-LR/12-10, 36p. https://archimer.ifremer.fr/doc/00094/20558/

Derolez, V., Malet, N., Fiandrino, A., Lagarde, F., Richard, M., Ouisse, V., Bec, B., Aliaume, C., 2020a. Fifty years of ecological changes: Regime shifts and drivers in a coastal Mediterranean lagoon during oligotrophication. Science of The Total Environment 732, 139292. https://doi.org/10.1016/j.scitotenv.2020.139292

Derolez, V., Soudant, D., Malet, N., Chiantella, C., Richard, M., Abadie, E., Aliaume, C., Bec, B., 2020b. Two decades of oligotrophication: Evidence for a phytoplankton community shift in the coastal lagoon of Thau (Mediterranean Sea, France). Estuarine, Coastal and Shelf Science 241, 106810. https://doi.org/10.1016/j.ecss.2020.106810

Díaz, R., Rosenberg, R., 1995. Marine benthic hypoxia: A review of its ecological effects and the behavioural responses of benthic macrofauna, in: Oceanography and Marine Biology: An Annual Review. University College London Press, London, UK, pp. 245–303.

Donadi, S., van der Zee, E.M., van der Heide, T., Weerman, E.J., Piersma, T., van de Koppel, J., Olff, H., Bartelds, M., van Gerwen, I., Eriksson, B.K., 2014. The bivalve loop: Intra-specific facilitation in burrowing cockles through habitat modification. Journal of Experimental Marine Biology and Ecology 461, 44–52. https://doi.org/10.1016/j.jembe.2014.07.019

Dufrêne, M., Legendre, P., 1997. Species assemblages and indicator species:the need for a flexible asymmetrical approach. Ecological Monographs 67, 345–366. https://doi.org/10.1890/0012-9615(1997)067[0345:SAAIST]2.0.CO;2

Elliott, M., Quintino, V., 2007. The estuarine quality paradox, environmental homeostasis and the difficulty of detecting anthropogenic stress in naturally stressed areas. Marine Pollution Bulletin 54, 640–645. https://doi.org/10.1016/j.marpolbul.2007.02.003

Faulwetter, S., Markantonatou, V., Pavloudi, C., Papageorgiou, N., Keklikoglou, K., Chatzinikolaou, E., Pafilis, E., Chatzigeorgiou, G., Vasileiadou, K., Dailianis, T., Fanini, L., Koulouri, P., Arvanitidis, C., 2014. Polytraits: A database on biological traits of marine polychaetes. Biodiversity Data Journal 2, e1024. https://doi.org/10.3897/BDJ.2.e1024

Fiandrino, A., Ouisse, V., Dumas, F., Lagarde, F., Pete, R., Malet, N., Le Noc, S., de Wit, R., 2017. Spatial patterns in coastal lagoons related to the hydrodynamics of seawater intrusion. Marine Pollution Bulletin 119, 132–144. https://doi.org/10.1016/j.marpolbul.2017.03.006

Gamito, S., Gilabert, J., Marcos Diego, C., Pérez-Ruzafa, A., 2005. Effects of changing environmental conditions on lagoon ecology, in: Gonenc, I., Wolfin, J. (Eds.), Coastal Lagoons – Ecosystem Processes and Modeling for Sustainable Use and Development. CRC Press, pp. 193–229.

Giampaoletti, J., Sbrana, A., Magni, P., Gravina, M.F., 2023. Macrobenthos of the Tortolì lagoon: a peculiar case of high benthic biodiversity among Mediterranean lagoons. Diversity 15, 783. https://www.mdpi.com/1424-2818/15/6/783

González-Ortiz, V., Egea, L.G., Jiménez-Ramos, R., Moreno-Marín, F., Pérez-Lloréns, J.L., Bouma, T.J., Brun, F.G., 2014. Interactions between seagrass complexity, hydrodynamic flow and biomixing alter food availability for associated filter-feeding organisms. PLOS ONE 9, e104949. https://doi.org/10.1371/journal.pone.0104949

Guelorget, O., Michel, P., 1979. Les peuplements benthiques d'un étang littoral Languedocien, l'étang du Prévost (Hérault): 2. Étude quantitative de la macrofaune des sables. Téthys.

Guelorget, O., Perthuisot, J., Lamy, N., Lefebvre, A., 1994. Structure et organisation de l'étang de Thau
d'après la faune benthique (macrofaune, méiofaune). Relations avec le confinement. Oceanologica
Acta 17, 105–114.
Guelorget, O., Perthuisot, J.P., 1983. Le domaine paralique: expressions géologiques, biologiques et
économiques du confinement, Travaux du laboratoire de géologie, n°16. Presses de l'école normale
supérieure, Paris.
Harman, W.N., 1972. Benthic substrates: their effect on fresh-water mollusca. Ecology 53, 271–277.
https://doi.org/10.2307/1934081
Hebbali, A., 2020. olsrr: tools for building ols regression models.
Jessen, C., Bednarz, V., Rix, L., Teichberg, M., Wild, C., 2015. Marine Eutrophication, in: Environmental
Indicators. p. 27. https://doi.org/10.1007/978-94-017-9499-2_11
Jones, A.G., Dubois, S.F., Desroy, N., Fournier, J., 2021. Intertidal ecosystem engineer species promote
benthic-pelagic coupling and diversify trophic pathways. Marine Ecology Progress Series 660, 119–
139. https://doi.org/10.3354/meps13600
Jones, A.G., Schaal, G., Boyé, A., Creemers, M., Derolez, V., Desroy, N., Fiandrino, A., Mouton, T.L., Simier,
M., Smith, N., Ouisse, V., 2023. Data, scripts and code for the preprint "Disentangling the effects of
eutrophication and natural variability on macrobenthic communities across French coastal lagoons."
https://doi.org/10.1101/2022.08.18.504439
Josse, J., Husson, F., 2016. missMDA: a package for handling missing values in multivariate data analysis.
Journal of Statistical Software 70, 1–31. https://doi.org/10.18637/jss.v070.i01
Josse, J., Husson, F., 2012. Handling missing values in exploratory multivariate data analysis methods.
Journal de la Société Française de Statistique 153, 79–99.
Kassambara, A., 2020. rstatix: pipe-friendly framework for basic statistical tests.
Kermagoret, C., Claudet, J., Derolez, V., Nugues, M.M., Ouisse, V., Quillien, N., Baulaz, Y., Le Mao, P.,
Scemama, P., Vaschalde, D., Bailly, D., Mongruel, R., 2019. How does eutrophication impact bundles
of ecosystem services in multiple coastal habitats using state-and-transition models. Ocean & Coastal
Management 174, 144–153. https://doi.org/10.1016/j.ocecoaman.2019.03.028
Kjerfve, B., 1994. Chapter 1 Coastal Lagoons, in: Kjerfve, B. (Ed.), Elsevier Oceanography Series, Coastal
Lagoon Processes. Elsevier, pp. 1–8. https://doi.org/10.1016/S0422-9894(08)70006-0
Lacoste, E., Jones, A., Callier, M., Klein, J., Lagarde, F., Derolez, V., 2023. A review of knowledge on the
impacts of multiple anthropogenic pressures on the soft-bottom benthic ecosystem in Mediterranean
coastal lagoons. Estuaries and Coasts. https://doi.org/10.1007/s12237-023-01188-9
Landman, M.J., Van Den Heuvel, M.R., Ling, N., 2005. Relative sensitivities of common freshwater fish and
invertebrates to acute hypoxia. New Zealand Journal of Marine and Freshwater Research 39, 1061–
1067. https://doi.org/10.1080/00288330.2005.9517375
Lasserre, P., 1979. Coastal lagoons: sanctuary ecosystems, cradles of culture, targets for economic
growth. Nature and Resources.
Le Fur, I., De Wit, R., Plus, M., Oheix, J., Simier, M., Ouisse, V., 2018. Submerged benthic macrophytes in
Mediterranean lagoons: distribution patterns in relation to water chemistry and depth. Hydrobiologia
808, 175–200. https://doi.org/10.1007/s10750-017-3421-y
Le Fur, I., Wit, R.D., Plus, M., Oheix, J., Derolez, V., Simier, M., Malet, N., Ouisse, V., 2019. Re-
oligotrophication trajectories of macrophyte assemblages in Mediterranean coastal lagoons based on
17-year time-series. Marine Ecology Progress Series 608, 13–32. https://doi.org/10.3354/meps12814
Legendre, P., Legendre, L., 2012. Numerical ecology, 3rd ed. Elsevier.
Lepoint, G., Nyssen, F., Gobert, S., Dauby, P., Bouquegneau, J.-M., 2000. Relative impact of a seagrass bed
and its adjacent epilithic algal community in consumer diets. Marine Biology 136, 513–518.
https://doi.org/10.1007/s002270050711
Leruste, A., Malet, N., Munaron, D., Derolez, V., Hatey, E., Collos, Y., De Wit, R., Bec, B., 2016. First steps
of ecological restoration in Mediterranean lagoons: Shifts in phytoplankton communities. Estuarine,
Coastal and Shelf Science 180, 190–203. https://doi.org/10.1016/j.ecss.2016.06.029
Levin, L.A., Boesch, D.F., Covich, A., Dahm, C., Erséus, C., Ewel, K.C., Kneib, R.T., Moldenke, A., Palmer,
M.A., Snelgrove, P., Strayer, D., Weslawski, J.M., 2001. The function of marine critical transition zones
and the importance of sediment biodiversity. Ecosystems 4, 430–451.
https://doi.org/10.1007/s10021-001-0021-4

Magalhães, L., Pires, A., Velez, C., Martins, R., Figueira, E., Soares, A.M.V.M., Freitas, R., 2019. Seasonal
and spatial alterations in macrofaunal communities and in *Nephtys cirrosa* (Polychaeta) oxidative
stress under a salinity gradient: A comparative field monitoring approach. Ecological Indicators 96,
192–201. https://doi.org/10.1016/j.ecolind.2018.08.045
Magni, P., Como, S., Kamijo, A., Montani, S., 2017. Effects of *Zostera marina* on the patterns of spatial
distribution of sediments and macrozoobenthos in the boreal lagoon of Furen (Hokkaido, Japan).
Marine Environmental Research 131, 90–102. https://doi.org/10.1016/j.marenvres.2017.09.013
Magni, P., Semprucci, F., Gravina, M.F., 2022. Joint analysis of macrofaunal and meiofaunal assemblages
improves the assessment of lagoonal environmental heterogeneity. Estuarine, Coastal and Shelf
Science 266, 107740. https://doi.org/10.1016/j.ecss.2021.107740
Magni, P., Gravina, M.F., 2023. Macrobenthos of lagoon ecosystems: a comparison in vegetated and bare
sediments. Advances in Oceanography and Limnology 14. https://doi.org/10.4081/aiol.2023.11124
Mancinelli, G., Rossi, L., 2001. Indirect, size-dependent effects of crustacean mesograzers on the
Rhodophyta *Gracilaria verrucosa* (Hudson) Papenfuss: evidence from a short-term study in the Lesina
Lagoon (Italy). Marine Biology 138, 1163–1173. https://doi.org/10.1007/s002270100545
Martini, S., Larras, F., Boyé, A., Faure, E., Aberle, N., Archambault, P., Bacouillard, L., Beisner, B.E., Bittner,
L., Castella, E., Danger, M., Gauthier, O., Karp-Boss, L., Lombard, F., Maps, F., Stemmann, L., Thiébaut,
E., Usseglio-Polatera, P., Vogt, M., Laviale, M., Ayata, S.-D., 2021. Functional trait-based approaches as
a common framework for aquatic ecologists. Limnology and Oceanography 66, 965–994.
https://doi.org/10.1002/lno.11655
Menu, M., Sandrine, V., Bajjouk, T., Derolez, V., Fiandrino, A., Giraud, A., Grillas, P., Ouisse, V., 2019.
Rapport final du projet CHAMILA (Cartographie des habitats en milieu lagunaire méditerranéen)
R.ODE/UL/LER/LR/19.34. https://doi.org/10.13155/70545
Mistri, M., Borja, A., Aleffi, I.F., Lardicci, C., Tagliapietra, D., Munari, C., 2018. Assessing the ecological
status of Italian lagoons using a biomass-based index. Marine Pollution Bulletin 126, 600–605.
https://doi.org/10.1016/j.marpolbul.2017.09.048
Miyamoto, Y., Nakano, T., Yamada, K., Hatakeyama, K., Hamaguchi, M., 2019. Combined effects of drift
macroalgal bloom and warming on occurrence and intensity of diel-cycling hypoxia in a eutrophic
coastal lagoon. Estuaries and Coasts 42, 494–503. https://doi.org/10.1007/s12237-018-0484-6
Morris, E.K., Caruso, T., Buscot, F., Fischer, M., Hancock, C., Maier, T.S., Meiners, T., Müller, C.,
Obermaier, E., Prati, D., Socher, S.A., Sonnemann, I., Wäschke, N., Wubet, T., Wurst, S., Rillig, M.C.,
2014. Choosing and using diversity indices: insights for ecological applications from the German
Biodiversity Exploratories. Ecol Evol 4, 3514–3524. https://doi.org/10.1002/ece3.1155
Muxika, I., Borja, Á., Bald, J., 2007. Using historical data, expert judgement and multivariate analysis in
assessing reference conditions and benthic ecological status, according to the European Water
Framework Directive. Marine Pollution Bulletin, Implementation of the Water Framework Directive in
European marine waters 55, 16–29. https://doi.org/10.1016/j.marpolbul.2006.05.025
Neveux, J., Lantoine, F., 1993. Spectrofluorometric assay of chlorophylls and phaeopigments using the
least squares approximation technique. Deep Sea Research Part I: Oceanographic Research Papers 40,
1747–1765. https://doi.org/10.1016/0967-0637(93)90030-7
Newton, A., Oliveira, P.S., Icely, J.D., Foster, P.A., 2010. Monitoring of oxygen condition in the Ria
Formosa coastal lagoon, Portugal. J Environ Monit 12, 355–360. https://doi.org/10.1039/b914015h
Nordström, M., Booth, D.M., 2007. Drift algae reduce foraging efficiency of juvenile flatfish. Journal of
Sea Research 58, 335–341. https://doi.org/10.1016/j.seares.2007.08.001
Norkko, J., Bonsdorff, E., Norkko, A., 2000. Drifting algal mats as an alternative habitat for benthic
invertebrates: Species specific responses to a transient resource. Journal of Experimental Marine
Biology and Ecology 248, 79–104. https://doi.org/10.1016/S0022-0981(00)00155-6
Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., Minchin, P.R., O'Hara, R.B.,
Simpson, G.L., Solymos, P., Stevens, M.H.H., Szoecs, E., Wagner, H., 2019. vegan: Community Ecology
Package.
Ouisse, V., Riera, P., Migné, A., Leroux, C., Davoult, D., 2011. Freshwater seepages and ephemeral
macroalgae proliferation in an intertidal bay: I Effect on benthic community structure and food web.
Estuarine, Coastal and Shelf Science 91, 272–281. https://doi.org/10.1016/j.ecss.2010.10.034

Pasqualini, V., Derolez, V., Garrido, M., Orsoni, V., Baldi, Y., Etourneau, S., Leoni, V., Rébillout, P., Laugier,
T., Souchu, P., Malet, N., 2017. Spatiotemporal dynamics of submerged macrophyte status and
watershed exploitation in a Mediterranean coastal lagoon: Understanding critical factors in ecosystem
degradation and restoration. Ecological Engineering 102, 1–14.
https://doi.org/10.1016/j.ecoleng.2017.01.027

Pearson, T.H., Rosenberg, R., 1978. Macrobenthic succession in relation to organic enrichment and
pollution of the marine environment. Oceanography and Marine Biology: An Annual Review 16, 229–
311. https://doi.org/10.2983/035.034.0121u1.10

Pérez-Ruzafa, A., Marcos, C., Pérez-Ruzafa, I.M., 2011b. Recent advances in coastal lagoons ecology:
evolving old ideas and assumptions. Transitional Waters Bulletin 5, 50–74.
https://doi.org/10.1285/i1825229Xv5n1p50

Pérez-Ruzafa, A., Marcos, C., Pérez-Ruzafa, I.M., 2011a. Mediterranean coastal lagoons in an ecosystem
and aquatic resources management context. Physics and Chemistry of the Earth, Parts A/B/C, Water
in the Mediterranean Basin 36, 160–166. https://doi.org/10.1016/j.pce.2010.04.013

Pérez-Ruzafa, A., Marcos, C., Pérez-Ruzafa, I.M., Barcala, E., Hegazi, M.I., Quispe, J., 2007a. Detecting
changes resulting from human pressure in a naturally quick-changing and heterogeneous
environment: Spatial and temporal scales of variability in coastal lagoons. Estuarine, Coastal and Shelf
Science, Biodiversity and Ecosystem Functioning in Coastal and Transitional Waters 75, 175–188.
https://doi.org/10.1016/j.ecss.2007.04.030

Pérez-Ruzafa, A., Marcos, C., Pérez-Ruzafa, I.M., Pérez-Marcos, M., 2013. Are coastal lagoons physically
or biologically controlled ecosystems? Revisiting r vs. K strategies in coastal lagoons and estuaries.
Estuarine, Coastal and Shelf Science, Estuarine and lagoon biodiversity and their natural goods and
services 132, 17–33. https://doi.org/10.1016/j.ecss.2012.04.011

Pérez-Ruzafa, Angel, Marcos, C., Pérez-Ruzafa, I.M., Pérez-Marcos, M., 2011c. Coastal lagoons:
"transitional ecosystems" between transitional and coastal waters. Journal of Coastal Conservation
15, 369–392. https://doi.org/10.1007/s11852-010-0095-2

Perez-Ruzafa, A., Marcos-Diego, C., 1992. Colonization rates and dispersal as essential parameters in the
confinement theory to explain the structure and horizontal zonation of lagoon benthic assemblages.
Technical Report Rapport Commission International Mer Medit 33.

Pérez-Ruzafa, Angel, Mompeán, M.C., Marcos, C., 2007b. Hydrographic, geomorphologic and fish
assemblage relationships in coastal lagoons. Hydrobiologia 577, 107–125.
https://doi.org/10.1007/s10750-006-0421-8

Perthuisot, J.-P., Guelorget, O., 1983. Le confinement, paramètre essentiel de la dynamique biologique
du domaine paralique. Sciences Géologiques, bulletins et mémoires 36, 239–248.
https://doi.org/10.3406/sgeol.1983.1642

Pitacco, V., Reizopoulou, S., Sfriso, Adriano, Sfriso, Andrea, Mistri, M., Munari, C., 2019. The difficulty of
disentangling natural from anthropogenic forcing factors makes the evaluation of ecological quality
problematic: A case study from Adriatic lagoons. Marine Environmental Research 150, 104756.
https://doi.org/10.1016/j.marenvres.2019.104756

Provost, C., Gevrey, M., Argillier, C., 2012. La macrofaune benthique indicatrice de l'état des lagunes
oligohalines méditerranéennes ? [Rapport de recherche] IRSTEA. 2012, pp.37. https://hal.science/hal-
02597249/

Puente, A., Diaz, R.J., 2008. Is it possible to assess the ecological status of highly stressed natural
estuarine environments using macroinvertebrates indices? Marine Pollution Bulletin 56, 1880–1889.
https://doi.org/10.1016/j.marpolbul.2008.07.016

Quillien, N., Nordström, M.C., Gauthier, O., Bonsdorff, E., Paulet, Y.-M., Grall, J., 2015. Effects of
macroalgal accumulations on the variability in zoobenthos of high-energy macrotidal sandy beaches.
Marine Ecology Progress Series 522, 97–114. https://doi.org/10.3354/meps11151

Quillien, N., Nordström, M.C., Schaal, G., Bonsdorff, E., Grall, J., 2016. Opportunistic basal resource
simplifies food web structure and functioning of a highly dynamic marine environment. Journal of
Experimental Marine Biology and Ecology 477, 92–102. https://doi.org/10.1016/j.jembe.2016.01.010

R Core Team, 2020. R: A language and environment for statistical computing. R Foundation for Statistical
Computing, Vienna, Austria.

Reizopolou, S., Penna, M., Boix, D., Buchet, R., Costas, N., Derolez, V., Gascon, S., Gifre, J., Martinoy, M., Pardo, I., Quintana, X., Trabucco, B., Sala, J., Salas Herrero, F., 2018. Transitional waters Mediterranean geographic intercalibration group: Benthic invertebrates fauna ecological assessment methods, EUR 29561 EN, Publications Office of the European Union, Luxembourg, 2018, ISBN 978-92-79-98373-3, JRC114720. https://doi.org/10.2760/625400

Reizopoulou, S., Nicolaidou, A., 2007. Index of size distribution (ISD): a method of quality assessment for coastal lagoons. Hydrobiologia 577, 141–149. https://doi.org/10.1007/s10750-006-0423-6

Reizopoulou, S., Simboura, N., Sigala, K., Barbone, E., Aleffi, F., Kaisakis, G., Rosati, I., Basset, A., Nicolaidou, A., 2014. Assessing the ecological status of Mediterranean coastal lagoons using macroinvertebrates. Comparison of the most commonly used methods. Mediterranean Marine Science 15, 602–612. https://doi.org/10.12681/mms.606

Riedel, B., Zuschin, M., Stachowitsch, M., 2012. Tolerance of benthic macrofauna to hypoxia and anoxia in shallow coastal seas: a realistic scenario. Marine Ecology Progress Series 458, 39–52. https://doi.org/10.3354/meps09724

Rigaud, S., Deflandre, B., Grenz, C., Cesbron, F., Pozzato, L., Voltz, B., Grémare, A., Romero-Ramirez, A., Mirleau, P., Meulé, S., Faure, V., Mayot, N., Michotey, V., Bonin, P., Pascal, L., Cordier, M.-A., Lamarque, B., Tenório, M., Radakovitch, O., 2021. Benthic oxygen dynamics and implication for the maintenance of chronic hypoxia and ecosystem degradation in the Berre lagoon (France). Estuarine, Coastal and Shelf Science 258, 107437. https://doi.org/10.1016/j.ecss.2021.107437

Roberts, D.W., 2019. labdsv: ordination and multivariate analysis for ecology.

Robertson, B.P., Savage, C., Gardner, J.P.A., Robertson, B.M., Stevens, L.M., 2016. Optimising a widely-used coastal health index through quantitative ecological group classifications and associated thresholds. Ecological Indicators 69, 595–605. https://doi.org/10.1016/j.ecolind.2016.04.003

Roques, C., Grousset, E., Troussellier, M., Hermet, S., Le Carrer, J., Sar, C., Caro, A., 2020. A trade-off between mucocytes and bacteriocytes in Loripes orbiculatus gills (Bivalvia, Lucinidae): a mixotrophic adaptation to seasonality and reproductive status in a symbiotic species? Marine Biology 167. https://doi.org/10.1007/s00227-020-03768-w

Rose, J.M., Caron, D.A., 2007. Does low temperature constrain the growth rates of heterotrophic protists? Evidence and implications for algal blooms in cold waters. Limnology and Oceanography 52, 886–895. https://doi.org/10.4319/lo.2007.52.2.0886

Rosenberg, R., Loo, L.-O., Möller, P., 1992. Hypoxia, salinity and temperature as structuring factors for marine benthic communities in a eutrophic area. Netherlands Journal of Sea Research 30, 121–129. https://doi.org/10.1016/0077-7579(92)90051-F

Rossi, F., Castelli, A., Lardicci, C., 2006. Distribution of macrobenthic assemblages along a marine gradient in Mediterranean eutrophic coastal lagoons. Marine ecology 27, 66–75. https://doi.org/10.1111/j.1439-0485.2006.00078.x

Sabetta, L., Barbone, E., Giardino, A., Galuppo, N., Basset, A., 2007. Species–area patterns of benthic macro-invertebrates in Italian lagoons. Hydrobiologia 577, 127–139. https://doi.org/10.1007/s10750-006-0422-7

Scaps, P., 2002. A review of the biology, ecology and potential use of the common ragworm Hediste diversicolor (O.F. Müller) (Annelida: Polychaeta). Hydrobiologia 470, 203–218. https://doi.org/10.1023/A:1015681605656

Schielzeth, H., 2010. Simple means to improve the interpretability of regression coefficients. Methods in Ecology and Evolution 1, 103–113. https://doi.org/10.1111/j.2041-210X.2010.00012.x

Sigala, K., Reizopoulou, S., Basset, A., Nicolaidou, A., 2012. Functional diversity in three Mediterranean transitional water ecosystems. https://doi.org/10.1016/J.ECSS.2012.06.002

Sigovini, M., Keppel, E., Tagliapietra, D., 2013. M-AMBI revisited: looking inside a widely-used benthic index. Hydrobiologia 717, 41–50. https://doi.org/10.1007/s10750-013-1565-y

Souchu, P., Bec, B., Smith, V.H., Laugier, T., Fiandrino, A., Benau, L., Orsoni, V., Collos, Y., Vaquer, A., 2010. Patterns in nutrient limitation and chlorophyll a along an anthropogenic eutrophication gradient in French Mediterranean coastal lagoons. Can. J. Fish. Aquat. Sci. 67, 743–753. https://doi.org/10.1139/F10-018

Souchu, P., Gasc, A., Collos, Y., Vaquer, A., Tournier, H., Bibent, B., Deslous-Paoli, J.-M., 1998. Biogeochemical aspects of bottom anoxia in a Mediterranean lagoon (Thau, France). Marine Ecology-progress Series 164, 135–146. https://doi.org/10.3354/meps164135

Stendera, S., Johnson, R.K., 2008. Habitat-specific stability and persistence of benthic invertebrate communities in boreal lakes. https://doi.org/10.1127/1863-9135/2008/0171-0311

Tagliapietra, D., Sigovini, M., Ghirardini, A., 2009. A review of terms and definitions to categorise estuaries, lagoons and associated environments. https://doi.org/10.1071/MF08088

Tagliapietra, D., Sigovini, M., Magni, P., 2012. Saprobity: a unified view of benthic succession models for coastal lagoons. Hydrobiologia 686, 15–28. https://doi.org/10.1007/s10750-012-1001-8

Tournoud, M.-G., Payraudeau, S., Cernesson, F., Salles, C., 2006. Origins and quantification of nitrogen inputs into a coastal lagoon: Application to the Thau lagoon (France). Ecological Modelling, Special Issue on Southern European Coastal Lagoons 193, 19–33. https://doi.org/10.1016/j.ecolmodel.2005.07.038

Trombetta, T., Vidussi, F., Mas, S., Parin, D., Simier, M., Mostajir, B., 2019. Water temperature drives phytoplankton blooms in coastal waters. PLOS ONE 14, e0214933. https://doi.org/10.1371/journal.pone.0214933

Vanda Orro, C., Cabana, D., 2021. Seasonal implications of the benthic ecological status assessment in a protected Mediterranean coastal lagoon: Stagno Longu of Posada (Sardinia, Italy). https://doi.org/10.1016/J.RSMA.2021.101759

Vaquer-Sunyer, R., Duarte, C.M., 2011. Temperature effects on oxygen thresholds for hypoxia in marine benthic organisms. Global Change Biology 17, 1788–1797. https://doi.org/10.1111/j.1365-2486.2010.02343.x

Venables, W.N., Ripley, B.D., 2002. Modern Applied Statistics with S, Fourth. ed. Springer, New York.

Vergara-Chen, C., González-Wangüemert, M., Marcos, C., Pérez-Ruzafa, Á., 2013. Small-scale genetic structure of Cerastoderma glaucum in a lagoonal environment: potential significance of habitat discontinuity and unstable population dynamics. Journal of Molluscan Studies 79, 230–240. https://doi.org/10.1093/mollus/eyt015

Vernberg, F.J., 1982. Environmental adaptation to lagoon systems. Oceanologica Acta, Special issue.

Vignes, F., Barbone, E., Breber, P., D'Adamo, R., Leonilde, R., Ungaro, N., Focardi, S., Renzi, M., Basset, A., 2010. Spatial and temporal description of the dystrophic crisis in Lesina lagoon during summer 2008. Transitional Waters Bulletin 3, 47–62. https://doi.org/10.1285/i1825229Xv3n2p47

Vizzini, S., Mazzola, A., 2006. Sources and transfer of organic matter in food webs of a Mediterranean coastal environment: Evidence for spatial variability. Estuarine, Coastal and Shelf Science 66, 459–467. https://doi.org/10.1016/j.ecss.2005.10.004

Ware, C., Dijkstra, J.A., Mello, K., Stevens, A., O'Brien, B., Ikedo, W., 2019. A novel three-dimensional analysis of functional architecture that describes the properties of macroalgae as a refuge. Marine Ecology Progress Series 608, 93–103. https://doi.org/10.3354/meps12800

Wei, T., Simko, V., 2017. corrplot: visualization of a correlation matrix.

Weissberger, E.J., Coiro, L.L., Davey, E.W., 2009. Effects of hypoxia on animal burrow construction and consequent effects on sediment redox profiles. Journal of Experimental Marine Biology and Ecology 371, 60–67. https://doi.org/10.1016/j.jembe.2009.01.005

Wilke, M., Boutiere, H., 2000. Hydrobiological, physical and chemical characteristics and spatio-temporal dynamics of an oligotrophic mediterranean lagoon: the étang de la Palme (France). Vie et Milieu / Life & Environment, 101-115

Zaldívar, J., Viaroli, P., Newton, A., Wit, R., Ibañez, C., Reizopoulou, S., Somma, F., Razinkovas, A., Basset, A., Holmer, M., Murray, N., 2008. Eutrophication in transitional waters: an overview. https://doi.org/10.1285/l18252273V2N1P1

Zilius, M., Giordani, G., Petkuviene, J., Lubiene, I., Ruginis, T., Bartoli, M., 2015. Phosphorus mobility under short-term anoxic conditions in two shallow eutrophic coastal systems (Curonian and Sacca di Goro lagoons). Estuarine, Coastal and Shelf Science 164, 134–146. https://doi.org/10.1016/j.ecss.2015.07.004

	Eutrophication	**Natural variability**
Community	Mean [Chla] + mean [NO$_x$] + CV [NO$_x$] + mean [NH$_4^+$]	Lagoon-sea connection + lagoon surface + CV O$_2$ sat + mean Temp
Density	Mean [NH$_4^+$] + mean [NH$_4^+$]^2	Mean Temp + mean Sal + mean Sal^2
N2	Mean [Chla] + CV [TN] + CV [TN]^2	Lagoon-sea connection + β$_{MM}$
H'	Mean [Chla] + CV [TN] + CV [TN]^2	Lagoon-sea connection + β$_{MM}$
SR	Mean [Chla] + mean [NO$_x$]	Mean Temp + mean Sal
M-AMBI	Mean [Chla] + mean [NO$_x$] + CV [NO$_x$]	Lagoon-sea connection + β$_{MM}$ + mean Temp