1 Hypoxia truncates and constitutively activates the key
2 cholesterol synthesis enzyme squalene monooxygenase

Hudson W. Coates[1], Ellen M. Olzomer[1], Ximing Du[1], Rhonda Farrell[2,3], Hongyuan Yang[1],
Frances L. Byrne[1], and Andrew J. Brown[1]*

[1]School of Biotechnology and Biomolecular Sciences, UNSW Sydney, Sydney NSW 2052,
Australia

[2]Chris O'Brien Lifehouse, Camperdown, NSW 2050, Australia

[3]Prince of Wales Private Hospital, Randwick, NSW 2031, Australia

*Corresponding author. Email: aj.brown@unsw.edu.au

10 Abstract

Cholesterol synthesis is both energy- and oxygen-intensive, yet relatively little is
known of the regulatory effects of hypoxia on pathway enzymes. We previously showed that
the rate-limiting and first oxygen-requiring enzyme of the committed cholesterol synthesis
pathway, squalene monooxygenase (SM), can undergo partial proteasomal degradation that
renders it constitutively active. Here, we show that hypoxia is the physiological trigger for this
truncation, which occurs through a two-part mechanism: (1) increased targeting of SM to the
proteasome *via* stabilization of the E3 ubiquitin ligase MARCHF6, and (2) accumulation of the
SM substrate, squalene, which impedes the complete degradation of SM and liberates its
truncated form. Truncation of SM permits its localization to lipid droplets and is increased in
hypoxic endometrial cancer tissues. These results uncover a feedforward mechanism that
enables SM to accommodate fluctuations in substrate levels yet is also a likely contributor to
its widely reported oncogenic properties.

Keywords: squalene monooxygenase / cholesterol / hypoxia / proteasome / cancer

24 Introduction

Cholesterol is an essential component of mammalian cell membranes, yet its aberrant
accumulation is detrimental [1]. Most cellular cholesterol arises from an energetically
expensive biosynthetic pathway requiring eleven oxygen molecules and over one hundred ATP
equivalents per molecule of product [2]. Furthermore, many intermediates of this pathway are
toxic in excess [3]. Coordinated regulation of cholesterol synthesis enzymes is therefore vital
to ensure that the pathway is active only when required, and that sufficient substrates and
cofactors are available to maintain flux through the full length of the pathway.

Squalene monooxygenase (SM, also known as squalene epoxidase or SQLE,
EC:1.14.14.17) catalyzes the rate-limiting conversion of squalene to monooxidosqualene in the
committed cholesterol synthesis pathway [4]. This reaction is the first of the pathway to require
molecular oxygen, with the introduced epoxide group ultimately forming the signature C3-
hydroxyl group of cholesterol. SM can also act a second time on monooxidosqualene to
produce dioxidosqualene, the precursor of the potent regulatory oxysterol 24(*S*),25-
epoxycholesterol [5]. As a flux-controlling enzyme, SM is subject to metabolic regulation at
both the transcriptional level *via* sterol regulatory element-binding proteins [6] and the post-
translational level *via* ubiquitination and proteasomal degradation [4]. The latter is mediated
by the N-terminal regulatory domain of SM (SM-N100), which senses lipid levels in the
endoplasmic reticulum (ER) membrane and accelerates or attenuates SM degradation in
response to excess cholesterol or squalene, respectively [7, 8]. These reciprocal feedback and
feedforward loops thus fine-tune SM activity according to metabolic demand. SM is typically
fully degraded by the proteasome; however, incomplete proteolysis produces a truncated form
of SM (trunSM) that lacks a large portion of the lipid-sensing SM-N100 domain but retains the
full catalytic domain [9]. This renders trunSM cholesterol-resistant and therefore constitutively
active. Although truncation is induced by the SM inhibitor NB-598, human cell lines express
similar levels of full-length and truncated SM [9]. This points to the existence of an unknown
physiological trigger for truncation.

Clarifying the mechanisms of SM regulation is particularly pertinent given the
importance of the enzyme, and cholesterol more generally [10], in oncogenesis.
Overexpression of the SM gene *SQLE* is associated with greater invasiveness and lethality in
breast [11], prostate [12, 13], and pancreatic cancer [14], amongst others. *SQLE* also shares its
genomic locus with the oncogene *MYC*, with which it is frequently co-amplified in cancer [11,
15]. Moreover, myc itself transcriptionally upregulates *SQLE* expression [16], while the tumor
suppressor protein p53 downregulates its expression [17]. At the protein level, aberrant SM
expression is implicated in colorectal cancer progression [18, 19] and the development of both
nonalcoholic steatohepatitis and hepatocellular carcinoma [20, 21]. Given its key role in
oxygen-dependent cholesterol synthesis, SM may be particularly critical for cancer cell
survival during hypoxia, which is common in the poorly vascularized cores of solid tumors and
often associated with poor prognosis [22]. In support of this idea, SM inhibition sensitizes
breast and colorectal cancer cells to hypoxia-induced cell death [23]. Although hypoxic cells
tend to accumulate cholesterol, there are conflicting reports on changes in biosynthetic flux
[24–26]. Furthermore, with the notable exception of the early pathway enzyme 3-hydroxy-
3-methylglutaryl-CoA reductase (HMGCR) [27], the effects of hypoxia on individual
biosynthetic enzymes are unknown. It is also unclear whether these might be perturbed in a
tumor context to favor continued cholesterol synthesis and cell proliferation.

Here, we show that hypoxic conditions induce the truncation of SM in a variety of cell
lines through a combination of accelerated proteasomal degradation and inhibition of its
complete proteolysis. This occurs due to the accumulation of both MARCHF6, the major E3
ubiquitin ligase for SM, and squalene, which impedes SM degradation through a mechanism
dependent on the SM-N100 regulatory domain. We also show that trunSM, but not full-length

SM, is capable of partitioning to lipid droplets, and that SM truncation is increased and
correlates with the magnitude of hypoxia in endometrial cancer tissues. Taken together, our
findings point towards a role for the constitutively active trunSM in adaptations to hypoxic
conditions and suggest that it contributes to the oncogenic effects of SM activity.

Results

Oxygen availability regulates SM truncation

We previously showed that SM is post-translationally regulated by both its substrate,
squalene, and its pathway end-product, cholesterol [4, 8]. The enzyme also undergoes partial
proteasomal degradation of its N-terminus to liberate a truncated protein (trunSM) that is
cholesterol-resistant and thus constitutively active (Fig. 1A) [9], although the physiological
trigger is unknown. As SM is a rate-limiting enzyme of cholesterol synthesis and catalyzes its
first oxygen-dependent reaction, we next tested if SM protein levels are affected by oxygen
availability. Incubation of HEK293T cells under hypoxic conditions (1% O_2) stabilized
hypoxia-inducible factor-1α (HIF1α; Fig. 1B) and upregulated its target genes *VEGF* and *CA9*
(Supplementary Fig. S1A), confirming the induction of a hypoxic response. We also noted a
striking increase in SM truncation under these conditions, as indicated by the disappearance of
full-length SM and accumulation of trunSM (Fig. 1B). Hypoxia-induced truncation of SM
increased over time (Fig. 1C) and with the magnitude of oxygen deprivation (Fig. 1D).
Notably, trunSM accumulation was greater under the severely hypoxic conditions
characteristic of solid tumors (0.5–2% O_2) than the 'physoxic' conditions experienced by
normal human tissues *in situ* (3–7.5% O_2) [28]. This suggested that increased SM truncation is
a feature of pathophysiological hypoxia. The net result of hypoxia-induced truncation was a
progressive increase in the levels of total SM over time (expressed as the sum of full-length
SM and trunSM levels; Supplementary Fig. S1B).

We also surveyed SM levels in a panel of cell lines and found that hypoxia-induced
accumulation of trunSM was generalizable to all, although full-length SM levels did not
decrease in MDA-MB-231 breast cancer cells (Supplementary Fig. S1C). As HIF1α and
hypoxia-inducible factor-2α (HIF2α) transcriptionally regulate the cellular response to
hypoxia, we next tested if their activity is required for SM truncation. However, knockdown of
HIF1A and *HIF2A* gene expression in HEK293T cells had no effect on the hypoxia-induced
truncation of SM (Supplementary Fig. S1D, E), ruling out the involvement of these
transcription factors and their target genes in the phenomenon.

Figure 1. Oxygen availability regulates SM truncation.

(A) Simplified overview of SM truncation [9]. Full-length SM contains an N-terminal domain responsible for feedback regulation by cholesterol. Ubiquitinated SM is targeted to the

proteasome, where proteolysis is prematurely halted within the N-terminal regulatory domain.
This liberates a truncated protein (trunSM) that no longer responds to cholesterol and is
therefore constitutively active. **(B)** HEK293T cells were incubated under normoxic (21% O_2)
or hypoxic (1% O_2) conditions for 24 h. **(C)** HEK293T cells were incubated under normoxic
or hypoxic conditions for the indicated times. **(D)** HEK293T cells were incubated under the
indicated oxygen concentrations (0.5–21% O_2) for 24 h. **(B–D)** Immunoblotting was
performed for SM and truncated SM (trunSM, red). Graphs depict densitometric quantification
of protein levels normalized to the normoxic condition, which was set to 1 (dotted line). In (C),
oxygen concentrations considered hypoxic (Hyp.), 'physoxic' (Phys.) or normoxic (Norm.)
[28] are indicated in blue. Where error bars are not visible, variance is too low to be depicted.
Data presented as mean ± SEM from $n \geq 3$ independent experiments (**, $p \leq 0.01$; two-tailed
one-sample *t*-test vs. hypothetical mean of 1).

Supplementary Figure S1. Hypoxia-induced truncation of SM is generalizable and independent of hypoxia-inducible factors.

(A) HEK293T cells were incubated under normoxic (21% O_2) or hypoxic (1% O_2) conditions for 24 h. Levels of the indicated HIF1α target gene transcripts were quantified, normalized to

the levels of three housekeeping transcripts and adjusted relative to the normoxic condition,
which was set to 1 (dotted line). **(B)** Quantification of total SM levels, expressed as the sum of
SM and trunSM levels, in Fig. 1C. **(C)** The indicated cell lines were incubated under normoxic
or hypoxic conditions for 24 h. **(D)** HEK293T cells were transfected with the indicated siRNAs
for 24 h and incubated under normoxic or hypoxic conditions for 24 h. Levels of siRNA target
transcripts in normoxic cells were quantified, normalized to the levels of the *PBGD*
housekeeping transcript and adjusted relative to the control siRNA condition, which was set to
1 (dotted line). **(E)** HEK293T cells were treated as described in (B). **(B, C, E)** Graphs depict
densitometric quantification of protein levels normalized to the respective normoxic
conditions, which were set to 1 (dotted line). **(A–E)** Data presented as mean ± SEM from $n \geq 3$
independent experiments (*, $p \leq 0.05$; **, $p \leq 0.01$; [A, C, D] two-tailed one-sample *t*-test vs.
hypothetical mean of 1; [E] two-tailed ratio paired *t*-test vs. control siRNA condition).

Hypoxia transcriptionally and post-translationally reduces full-length SM levels

As the truncation of SM results from its partial proteasomal degradation [9], we
reasoned that hypoxia promotes truncation through a two-part mechanism: (1) targeting of full-
length SM to the proteasome, and (2) inhibition of its complete degradation. To confirm the
first step of this mechanism, we investigated the reason for the decline in full-length SM levels
during hypoxia. *SQLE* transcripts were downregulated in hypoxic HEK293T cells, as were
transcripts encoding the upstream cholesterol and isoprenoid synthesis enzyme HMGCR
(Fig. 2A). Downregulation of *SQLE* transcripts was not observed in MDA-MB-231 cells
(Supplementary Fig. S2A), accounting for the unchanged full-length SM levels in this cell line.
Although the reduction in *SQLE* and *HMGCR* transcripts in HEK293T cells likely reflected a
broad transcriptional suppression of cholesterol synthesis during hypoxia, as reported
previously [29, 30], the magnitude of *SQLE* downregulation was unlikely to fully explain the
large reduction in SM protein levels (Fig. 1B). Moreover, levels of a constitutively expressed
SM construct were markedly reduced during extended hypoxic incubations, with no associated
changes in mRNA levels (Fig. 2B, Supplementary Fig. S2B). We concluded that hypoxia
reduces the levels of full-length SM through both transcriptional downregulation and
accelerated post-translational degradation.

The basal and metabolically-regulated degradation of SM occurs through the ubiquitin-
proteasome system and is mediated by the SM-N100 regulatory domain [7, 8]. Therefore, we
tested the effect of hypoxia on HEK293 cells stably overexpressing an SM-N100 fusion protein
(SM-N100-GFP-V5). Like full-length SM, levels of SM-N100-GFP-V5 were reduced under
hypoxic conditions (Fig. 2C). Proteasomal inhibition using MG132 rescued SM and
SM-N100-GFP-V5 from this reduction, confirming that hypoxia-induced degradation occurs
via the proteasome. We also noted that the hypoxia-induced accumulation of trunSM was
ablated by MG132 (Supplementary Fig. S2C), consistent with the protein arising from partial
proteasomal proteolysis of SM [9]. Although hypoxia can trigger autophagy [31], this did not
play a role in SM degradation as inhibition of lysosomal acidification using bafilomycin A1
had no additive effect with MG132 (Fig. 2C). To identify residues required for hypoxia-
induced degradation of SM, we utilized protein constructs with mutations of previously
identified ubiquitination sites. Degradation was blunted by the disruption of Lys-82/90/100, a
cluster of redundant ubiquitination sites previously found to promote truncation [9], but not by
the disruption of Lys-290 [32] (Fig. 2D). Non-canonical cysteine, serine and threonine
ubiquitination sites required for the cholesterol-induced degradation of SM [33] also
contributed to hypoxia-induced degradation, suggesting that multiple ubiquitin signals are
involved.

To investigate the mechanism by which hypoxia promotes SM ubiquitination, we
considered the possible role of proline hydroxylation. This oxygen-dependent modification,
catalyzed by prolyl hydroxylases, is required for the ubiquitination and degradation of HIF1α
under normoxic conditions [34], although there is conflicting evidence for the existence of
substrates beyond the HIF proteins [35]. Indeed, treatment with prolyl hydroxylase inhibitors
had no effect on SM nor SM-N100 levels despite stabilizing HIF1α (Supplementary Fig. S2D).
SM and SM-N100 are targeted for proteasomal degradation by the E3 ubiquitin ligase

MARCHF6 [36]; therefore, we next tested if increased MARCHF6 activity could account for
the hypoxia-induced degradation of SM. In support of this idea, *MARCHF6* knockdown
reversed the hypoxic decline in full-length SM levels (Fig. 2E). The relative accumulation of
trunSM was also blunted (Supplementary Fig. S2E), consistent with truncation occurring post-
ubiquitination by MARCHF6. This effect was not abolished, however, indicating that SM can
be truncated even when targeted to the proteasome by other mechanisms. Surprisingly, there
was no significant effect of *MARCHF6* knockdown on hypoxia-induced degradation of
SM-N100-GFP-V5 (Fig. 2E), suggesting that full-length SM and the isolated SM-N100
domain are degraded through different proteasome-dependent routes under these conditions.
As truncation involves the full-length protein [9], we elected to further investigate the
MARCHF6-mediated regulation of SM.

Figure 2. Hypoxia transcriptionally and post-translationally reduces full-length SM levels.

(A) HEK293T cells were incubated under normoxic or hypoxic conditions for 24 h. Levels of the indicated transcripts were quantified, normalized to the levels of three housekeeping transcripts and adjusted relative to the normoxic condition, which was set to 1 (dotted line). **(B)** HEK293T cells were transfected with (HA)$_3$-SM-V5 for 24 h and incubated under

normoxic or hypoxic conditions for the indicated times. **(C)** HEK SM-N100-GFP-V5 cells
were treated with or without 20 µM MG132 and 20 nM bafilomycin A1 under normoxic or
hypoxic conditions for 16 h. **(D)** HEK293T cells were transfected with the indicated constructs
for 24 h and incubated under normoxic or hypoxic conditions for 16 h. **(E)** HEK
SM-N100-GFP-V5 cells were transfected with control or *MARCHF6* siRNA for 24 h and
incubated under normoxic or hypoxic conditions for 16 h. **(B–E)** Graphs depict densitometric
quantification of protein levels normalized to the respective normoxic conditions, which were
set to 1 (dotted line). **(A–E)** Data presented as mean $\pm$ SEM from $n \geq 3$ independent
experiments (*, $p \leq 0.05$; **, $p \leq 0.01$; [A, B] two-tailed one-sample t-test vs. hypothetical
mean of 1; [C–E] two-tailed ratio paired t-test vs. vehicle, wild-type [WT] or control siRNA
condition).

Supplementary Figure S2. Hypoxia-induced degradation of full-length SM is independent of proline hydroxylation.

(A) MDA-MB-321 cells were incubated under normoxic or hypoxic conditions for 24 h. Levels of *SQLE* transcripts were quantified, normalized to the levels of three housekeeping transcripts and adjusted relative to the normoxic condition, which was set to 1 (dotted line). **(B)** HEK293T cells were treated as described in Fig. 2B. Levels of *SQLE* transcripts were quantified, normalized to the levels of three housekeeping transcripts and adjusted relative to the normoxic condition, which was set to 1 (dotted line). **(C)** Quantification of trunSM levels in Fig. 2C. **(D)** HEK SM-N100-GFP-V5 cells were treated with or without 1 mM DMOG or 25 μM FG-4592 under normoxic or hypoxic conditions for 16 h. **(E)** Quantification of trunSM levels in Fig. 2E. **(C–E)** Graphs depict densitometric quantification of protein levels normalized to the respective

normoxic conditions, which were set to 1 (dotted line). **(A–E)** Data presented as mean ± SEM
from $n \geq 3$ independent experiments (*, $p \leq 0.05$; [A, B] two-tailed one-sample *t*-test vs.
hypothetical mean of 1; [C–E] two-tailed ratio paired *t*-test vs. vehicle or control siRNA
condition).

Hypoxia stabilizes the E3 ubiquitin ligase MARCHF6

To study MARCHF6 levels in hypoxic cells, we utilized a previously generated
HEK293 cell line stably overexpressing a V5-tagged form of the protein. This construct was
used due to the poor performance of endogenous MARCHF6 antibodies [37], and also to
eliminate any transcriptional contribution to protein levels. We first examined the response of
MARCHF6-V5 to hypoxia and found that it accumulated during prolonged hypoxic
incubations, correlating with the maximal decline in full-length SM levels (Fig. 3A).
Previously, we showed that cholesterol stabilizes MARCHF6 by interfering with its
autoubiquitination and proteasomal degradation [37]. To determine if a similar mechanism
accounted for the accumulation of MARCHF6 in hypoxia, we treated MARCHF6-V5 cells
with CB-5083, an inhibitor of the ER-associated degradation cofactor valosin-containing
protein, or the proteasome inhibitor MG132. Both treatments stabilized MARCHF6, with
CB-5083 having a greater effect, as described previously [37]. Importantly, both CB-5083 and
MG132 blocked the hypoxia-induced accumulation of MARCHF6 (Fig. 3B), consistent with a
mechanism involving reduced proteasomal degradation. We also tested the effect of hypoxia
on two MARCHF6 mutants: C9A, an inactive variant lacking a functional RING domain [38],
or N890A, a variant that is defective in auto-ubiquitination [39]. Both mutants showed reduced
accumulation during hypoxia (Fig. 3C), indicating that the effect on wild-type MARCHF6-V5
is due to impaired auto-ubiquitination.

Given that Asn-890 is required for the accumulation of MARCHF6 during hypoxia, we
next considered whether this residue might be modified by asparagine hydroxylation. This is
an oxygen-dependent modification known to control processes such as HIF1α target gene
transactivation [40] and protein ubiquitination [41]. To determine if a similar mechanism
regulates MARCHF6 in hypoxia, we used siRNA to deplete the asparaginyl hydroxylase
FIH-1. Although this increased basal MARCHF6-V5 levels and blunted its hypoxia-induced
accumulation (Supplementary Fig. S3A, B), confirmatory experiments using the FIH-1
inhibitor DM-NOFD [42] did not reproduce the effect (Supplementary Fig. S3C). We
concluded that asparagine hydroxylation is unrelated to the reduced autoubiquitination of
MARCHF6 in hypoxia, with the true mechanism currently unknown.

Figure 3. Hypoxia stabilizes the E3 ubiquitin ligase MARCHF6.

(A) HEK MARCHF6-V5 cells were incubated under normoxic or hypoxic conditions for the indicated times. **(B)** HEK MARCHF6-V5 cells were treated with or without 5 µM CB-5083 or 20 µM MG132 under normoxic or hypoxic conditions for 16 h. **(C)** The indicated HEK MARCHF6-V5 cell lines were incubated under normoxic or hypoxic conditions for 16 h. **(A–C)** Graphs depict densitometric quantification of MARCHF6-V5 levels normalized to the respective normoxic conditions, which were set to 1 (dotted line). Data presented as mean ± SEM from $n \geq 3$ independent experiments (*, $p \leq 0.05$; [A] two-tailed one-sample *t*-test vs. hypothetical mean of 1; [B, C] two-tailed ratio paired *t*-test vs. vehicle or WT condition).

Supplementary Figure S3. Hypoxia-induced stabilization of MARCHF6 is independent of asparagine hydroxylation.

(A) HEK293T cells were transfected with control or *FIH-1* siRNA for 24 h and incubated under normoxic or hypoxic conditions for 24 h. Levels of *FIH-1* target transcripts in normoxic cells were quantified, normalized to the levels of the *PBGD* housekeeping transcript and adjusted relative to the control siRNA condition, which was set to 1 (dotted line). **(B)** HEK MARCHF6-V5 cells were transfected with control or *FIH-1* siRNA for 24 h and incubated under normoxic or hypoxic conditions for 16 h. Graphs depict densitometric quantification of MARCHF6-V5 levels normalized to the (left) vehicle or (right) normoxic condition, which was set to 1 (dotted line). **(C)** HEK MARCHF6-V5 cells were treated with 500 μM DM-NOFD under normoxic or hypoxic conditions for 16 h. **(B, C)** Graphs depict densitometric quantification of MARCHF6-V5 levels normalized to the respective normoxic conditions, which were set to 1 (dotted line). **(A–C)** Data presented as mean $\pm$ SEM from $n \geq 3$ independent experiments (*, $p \leq 0.05$; **, $p \leq 0.01$; two-tailed one-sample *t*-test vs. hypothetical mean of 1 or [B, C] two-tailed ratio paired *t*-test vs. control siRNA or vehicle condition).

Hypoxia-induced squalene accumulation promotes partial degradation of SM

Having established that SM undergoes increased proteasomal degradation during
hypoxia, we next investigated how low oxygen levels favor its partial rather than complete
proteolysis to produce trunSM. As there is extensive precedent for metabolic regulation of
cholesterol synthesis enzymes and the pathway contains multiple oxygen-dependent reactions,
we considered whether the accumulation of a pathway intermediate might be responsible for
this phenomenon. Hypoxia-induced accumulation of trunSM occurred in cells incubated under
both lipoprotein-replete and lipoprotein-deficient conditions, in which the cholesterol synthesis
pathway is active (Fig. 4A). However, its accumulation was abolished when lipoprotein-
deficient cells were co-treated with a statin to inhibit HMGCR and the early cholesterol
synthesis pathway. By contrast, there was no effect of sterol depletion on the disappearance of
full-length SM. This indicated that an intermediate or end-product of cholesterol synthesis
promotes the partial rather than complete degradation of SM during hypoxia. We therefore
turned our attention to the SM substrate squalene, given that it allosterically regulates SM
degradation [8] and its conversion to monooxidosqualene is the first oxygen-dependent
reaction of cholesterol synthesis. Squalene accumulated over the course of a hypoxic
incubation (Fig. 4B, Supplementary Fig. S4), consistent with reduced SM activity under low-
oxygen conditions. Moreover, this accumulation of squalene was strikingly well-correlated
with the previously observed increase in trunSM levels (Fig. 4C), suggesting that the two
responses may be linked.

Delivery of exogenous squalene mimicked the hypoxia-induced accumulation of
trunSM in normoxic HEK293T (Fig. 4D, Supplementary Fig. S5A) and Huh7 cells
(Supplementary Fig. S5B), confirming its ability to promote partial degradation of SM.
Accumulation of trunSM was also induced by the oxygenated squalene derivatives
monooxidosqualene and dioxidosqualene (Fig. 4D) but not by its saturated analogue squalane
(Supplementary Fig. S5C), which has similar biophysical properties [43]. This indicated that
truncation is promoted by squalene and its structurally related molecules in a specific manner,
rather than through bulk membrane effects caused by lipid accumulation. To address the
possibility that exogenous squalene is converted to a downstream product responsible for
truncation, we performed squalene treatment of *SQLE*-knockout HEK293T cells
(Supplementary Fig. S6A–C) transfected with a catalytically inactive SM Y195F mutant [44]
to prevent the metabolism of added squalene. The truncated form of the Y195F mutant
accumulated upon squalene treatment in *SQLE*-knockout cells, confirming that squalene alone
can directly induce truncation (Supplementary Fig. S5D). There was no significant
accumulation of the truncated fragment in cells transfected with wild-type SM, likely due to
the clearance of exogenous squalene by the overexpressed protein and downstream enzymes.

To confirm that endogenously synthesized squalene is sufficient to trigger SM
truncation, cells were treated with inhibitors of the relevant cholesterol synthesis enzymes
(Supplementary Fig. S5A). The SM inhibitor NB-598 was excluded because of its ability to
induce truncation through direct binding and stabilization of the SM catalytic domain that
renders it resistant to proteasomal unfolding [8, 45]. Inhibition of squalene synthesis from
farnesyl diphosphate (TAK-475) abolished the hypoxia-induced accumulation of trunSM,

whereas elevated protein levels were still observed under conditions where squalene remained
able to accumulate: inhibition of lanosterol synthesis from monooxidosqualene (BIBB 515), or
inhibition of lanosterol demethylation (GR70585X) (Supplementary Fig. S5E). This result
confirmed that lanosterol, which also accumulates during hypoxia [27], has no effect on SM
truncation. We further noted that the inhibition of squalene or lanosterol synthesis, but not
lanosterol demethylation, increased the levels of full-length SM and SM-N100-GFP-V5 under
basal conditions. This was in line with our previous finding that farnesyl-containing molecules
such as monooxidosqualene, dioxidosqualene and a squalene-derived probe can each stabilize
SM *via* its regulatory domain in a similar manner to squalene itself [8]. Interestingly, the basal
levels of trunSM were also increased in the presence of these inhibitors (Supplementary
Fig. S5E), suggesting that all farnesyl-containing intermediates of cholesterol synthesis are
capable of inducing SM truncation. Nevertheless, as the primary substrate of SM, squalene is
likely to be the major driver of this process under hypoxic conditions.

SM contains two known squalene binding sites: the SM-N100 regulatory domain and
the active site of the catalytic domain [8]. As direct binding of the SM inhibitor NB-598 to the
catalytic domain results in truncation-promoting stabilization [9], we next considered whether
squalene also exerts its effect on truncation through direct binding and stabilization of the
catalytic domain. To eliminate the contribution of the SM-N100 domain, we transfected
SQLE-knockout cells with an ectopic form of trunSM (SM[ΔN65]-V5) that was stabilized by
NB-598 (Fig. 4E), consistent with past findings [8, 44]. By contrast, squalene treatment of cells
expressing an inactive SM(ΔN65)-V5 mutant (Y195F) did not lead to significant protein
stabilization. We concluded that squalene promotes SM truncation *via* the SM-N100 regulatory
domain, rather than through direct binding and stabilization of the SM catalytic domain.
Squalene treatment also did not increase MARCHF6-V5 levels (Supplementary Fig. S5F),
suggesting that the stabilization of the E3 ubiquitin ligase during hypoxia is unrelated to
squalene accumulation. This observation was consistent with the hypoxia-induced and
MARCHF6-mediated degradation of full-length SM being unaffected by the activity of the
cholesterol synthesis pathway (Fig. 4A). Together, this reinforced that squalene exerts its
truncation-promoting effects post-ubiquitination, during the proteasomal-degradation of SM.

Figure 4. Hypoxia-induced squalene accumulation promotes partial degradation of SM.

(A) HEK293T cells were incubated in medium containing fetal calf serum (FCS), lipoprotein-deficient FCS (LPDS) or LPDS containing 5 µM mevastatin and 50 µM mevalonolactone (LPDS + statin) for 8 h, refreshed in their respective media and incubated under normoxic or hypoxic conditions for 16 h. **(B)** HEK293T cells were incubated under normoxic or hypoxic conditions for the indicated times. Non-saponifiable lipids were extracted, and squalene levels were determined using gas chromatography-mass spectrometry and adjusted relative to the normoxic condition, which was set to 1 (dotted line). The maximal squalene level detected was 0.66 ± 0.12 ng per µg of total protein. **(C)** Pearson correlation between squalene levels in (B) and trunSM levels in Fig. 1C. Blue line indicates linear regression. **(D)** HEK293T cells were treated with or without 300 µM squalene (Squ.), monooxidosqualene (MOS) or dioxidosqualene (DOS) for 16 h. **(E)** HEK293T *SQLE*-knockout (*SQLE*-KO) clone 10 (c10) cells were transfected with the indicated constructs for 24 h, then treated with or without 1 µM

NB-598 or 300 µM squalene for 16 h. **(A, D, E)** Graphs depict densitometric quantification of

trunSM or truncated protein levels normalized to the (A) respective normoxic conditions or

(D, E) respective vehicle conditions, which were set to 1 (dotted line). **(A–E)** Data presented

as mean ± SEM from $n \geq 3$ independent experiments (*, $p \leq 0.05$; **, $p \leq 0.01$; [A] two-tailed

ratio paired *t*-test vs. FCS condition; [D, E] two-tailed one-sample *t*-test vs. hypothetical mean

of 1).

Supplementary Figure S4. Hypoxia induces squalene accumulation.

Representative selective ion monitoring chromatogram traces from gas chromatography-mass spectrometry analysis of squalene levels in Fig. 4B. Abundance was normalized to the 5α-cholestane internal standard, which was set to 1. Dagger indicates a non-specific analyte.

Supplementary Figure S5. Farnesyl-containing cholesterol synthesis intermediates
specifically promote partial degradation of SM.

(A) Simplified schematic of the cholesterol synthesis pathway depicting activities of squalene
synthase (SQS), SM, lanosterol synthase (LSS) and lanosterol 14α-demethylase (LDM),
alongside chemical structures of squalene, monooxidosqualene (MOS), dioxidosqualene
(DOS), squalane, and farnesyl diphosphate. Double-headed arrows indicate multiple enzymatic
steps, red labels indicate inhibitors of cholesterol synthesis, and green and blue labels indicate
molecules that were determined to promote or not to promote trunSM accumulation,
respectively. **(B)** Huh7 cells were treated with or without 300 μM squalene for 16 h. **(C)**
HEK293T cells were treated with or without 300 μM squalane for 16 h. **(D)** Parental (WT) or
HEK293T *SQLE*-knockout (*SQLE*-KO) clone 10 (c10) cells were transfected with the indicated
constructs for 24 h, then treated with or without 300 μM squalene for 16 h. **(E)** HEK SM-N100-
GFP-V5 cells were treated with the indicated compounds under normoxic or hypoxic
conditions for 16 h. **(F)** HEK MARCHF6-V5 cells were treated with or without 300 μM
squalene or squalane for 16 h. **(B–F)** Graphs depict densitometric quantification of protein
levels normalized to the respective vehicle or (E) respective normoxic conditions, which were
set to 1 (dotted line). Data presented as mean ± SEM from $n \geq 3$ independent experiments (*,
$p \leq 0.05$; two-tailed one-sample *t*-test vs. hypothetical mean of 1).

Supplementary Figure S6. Generation of *SQLE*-knockout HEK293T cells.

(A) HEK293T cells were transfected with CRISPR/Cas9 guide RNAs (gRNAs) targeting the proximal promoter and first exon of the *SQLE* gene as indicated, and three clonal populations of *SQLE*-KO cells were isolated and expanded for each parental cell line. The target region was amplified from genomic DNA and sequenced, which found that all three clones contained a genomic deletion between the gRNA target sites. **(B)** *SQLE* transcript levels were quantified in HEK293T *SQLE*-KO clones, normalized to the *PBGD* housekeeping transcript, and adjusted relative to the parental (WT) cell line, which was set to 1 (dotted line). Data presented as mean ± SEM for technical triplicates from a single experiment. **(C)** Protein lysates from the HEK293T WT cell line and *SQLE*-KO clones were immunoblotted for SM and trunSM.

trunSM uniquely localizes to lipid droplets

Accumulated squalene induces ER stress and is lipotoxic, necessitating its sequestration to lipid droplets [46, 47]. These organelles are also known to form under hypoxic conditions [48], prompting us to examine possible links with trunSM. Several proteomic studies have established that SM localizes to lipid droplets [49–51], but these analyses were unable to distinguish between the full-length and truncated forms of SM. Furthermore, we previously found that trunSM adopts a peripheral rather than integral ER membrane topology, which may improve its compatibility with the lipid droplet monolayer [9]. To expand on these findings, immunofluorescence and lipid droplet staining were used to investigate the localization of the (HA)$_3$-SM-V5 construct in oleate-loaded HeLa cells. The N-terminal (HA)$_3$ tag of the construct, which is present only in the full-length protein (Supplementary Fig. S7A), showed minimal association with the lipid droplet surface (Fig. 5A, C), in contrast to the lipid droplet-localized protein ORP5 (Supplementary Fig. S7B, E) [52]. However, the C-terminal V5 tag, which is present in both the full-length and truncated proteins, was associated with lipid droplets (Fig. 5B, C). The SM(ΔN65)-V5 construct showed a similar association, confirming that the truncated form of SM can partition to lipid droplets. A truncation-resistant (HA)$_3$-SM-V5 mutant (Δ81–120) did not show lipid droplet association of its N- nor C-terminal tags (Supplementary Fig. S7C–E), further demonstrating that full-length SM is excluded from this compartment. Finally, we examined the localization of the SM-N100-GFP-V5 construct, which contains a membrane-embedded re-entrant loop eliminated by truncation [9]. This construct also did not associate with lipid droplets (Fig. 5B, C; Supplementary Fig. S7B, E), indicating that the SM-N100 re-entrant loop is responsible for retaining full-length SM in the ER.

As an orthogonal approach to confirm the lipid droplet association of trunSM, we also fractionated lysates from oleate-loaded cells and performed immunoblotting. Consistent with our immunofluorescence data, endogenous trunSM, but not full-length SM, was present in the floating lipid droplet fraction alongside the marker protein ABHD5 (Fig. 5D). Likewise, the truncated form of the (HA)$_3$-SM-V5 construct showed a greater partitioning to lipid droplets than its full-length counterpart. In stark contrast, SM-N100-GFP-V5 did not partition to lipid droplets. These results complemented our immunofluorescence approach and confirmed that the loss of the SM-N100 re-entrant loop renders trunSM uniquely capable of localization to lipid droplets.

Figure 5. trunSM uniquely localizes to lipid droplets.

HeLa cells were transfected with the indicated constructs for 24 h and treated with 300 μM
oleate-bovine serum albumin complexes for 16 h. **(A, B)** Cells were fixed, lipid droplets (LDs)
were stained with BODIPY 493/503, and (A) anti-HA or (B) anti-V5 immunofluorescence was
performed (scale bar: 10 μm). **(C)** Quantification of LD association in (A) and (B), expressed
as the percentage of imaged transfected cells (*n*) with LD-proximal localization of the protein
and epitope of interest. Data were collected across two independent experiments. **(D)** LD,
cytosol (Cyto.), pellet, and total fractions were collected and adjusted for protein content, and
immunoblotting was performed for calnexin (endoplasmic reticulum marker), ABHD5 (LD
marker), and GAPDH (cytosol marker). Graph depicts quantification of LD partitioning,
expressed as the ratio between protein levels in LD and membrane fractions normalized to that
of ABHD5. Data presented as mean ± SEM from $n \geq 3$ independent experiments (*, $p \leq 0.05$;
two-tailed ratio paired *t*-test vs. full-length endogenous [endog.] SM or full-length [HA]$_3$-SM-
V5).

Supplementary Figure S7. Non-truncated SM does not associate with lipid droplets.

HeLa cells were transfected with the indicated constructs for 24 h and treated with 300 μM
oleate-bovine serum albumin complexes for 16 h. **(A)** Immunoblotting was performed for
N-terminal (HA)$_3$ and C-terminal V5 tags. Dagger indicates a non-specific band. **(B)** Cells were
fixed and lipid droplets (LDs) were stained with LipidTOX Deep Red (scale bar: 10 μm). **(C,**
D) Cells were fixed, LDs were stained with BODIPY 493/503, and (C) anti-HA or (D) anti-V5
immunofluorescence was performed (scale bar: 10 μm). **(E)** Quantification of LD association
in (B–D), expressed as the percentage of imaged transfected cells (*n*) with LD-proximal
localization of the protein and epitope of interest. Data were collected across two independent
experiments.

SM truncation is increased in hypoxic endometrial cancer tissues.

Aberrant SM expression and activity is an emerging hallmark of numerous malignancies [21, 53], yet the contribution of the constitutively active trunSM to oncogenesis is unknown. Therefore, we performed immunoblotting of paired tumor and adjacent benign tissue lysates from cohorts of lean and obese endometrial cancer patients. This cancer type was selected because of its close links with obesity [54, 55] and signaling by cholesterol-derived estrogens [56], as well as the unstudied status of SM in endometrial cancer. Similar trends were observed across both cohorts (Fig. 6A, Supplementary Fig. S8A–C), and the data were pooled. This showed that levels of HIF1α were increased in tumor tissues compared with adjacent benign tissues (Fig. 6B), consistent with the hypoxic conditions that occur within solid tumors [22]. Total SM levels were reduced in tumor tissues (Fig. 6B), yet this was accompanied by a dramatic increase in the proportion of SM that was truncated (Fig. 6C). Moreover, we observed a striking correlation between HIF1α levels and SM truncation across all tissues (Fig. 6D), further supporting the notion that hypoxia and truncation are linked. These results confirmed that the hypoxia-induced truncation of SM is a physiologically relevant phenomenon in human tissues and may contribute to endometrial tumorigenesis.

Figure 6. SM truncation is increased in hypoxic endometrial cancer tissues.

(A) Paired tumor (T) and adjacent benign (B) tissue lysates from lean and obese endometrial cancer patients were analyzed by immunoblotting. **(B, C)** Graphs depict densitometric quantification of protein levels from (A) normalized to the 14-3-3 housekeeping protein, and (C) expressed as a proportion of total SM. Data presented as mean ± SD from $n = 14$ paired tissue sets (*, $p \leq 0.05$; **, $p \leq 0.01$; two-tailed paired t-test vs. adjacent benign tissue; normality of distributions confirmed by D'Agostino & Pearson testing [57]). **(D)** Pearson correlation between HIF1α levels in (B) and trunSM levels in (C). Blue line indicates linear regression.

Supplementary Figure S8. SM truncation is increased in endometrial cancer tissues from both lean and obese patients.

Separate quantification of **(A)** HIF1α, **(B)** total SM, and **(C)** trunSM protein levels in tumor (T) and adjacent benign (B) tissues from lean and obese cohorts of endometrial cancer patients in Fig. 6A. Data presented as mean ± SD from $n = 7$ paired tissue sets in each cohort (*, $p \leq 0.05$; two-tailed Wilcoxon matched-pairs signed rank test vs. adjacent benign tissue).

Discussion

Cholesterol synthesis is tightly regulated by metabolic supply and demand, with the lipid-sensing and rate-limiting enzyme SM being a key point at which this regulation is exerted. We previously showed that partial proteasomal degradation of SM produces a truncated and constructively active form of the enzyme [9]. In this study, we identified hypoxia as the physiological trigger for trunSM formation in both cell culture and human tissues *in situ* (Fig. 1). Hypoxia-induced truncation occurs through a two-part mechanism: (1) an increase in the levels of MARCHF6, the E3 ubiquitin ligase that targets SM to the proteasome (Fig. 2, Fig. 3), and (2) the accumulation of squalene, which impedes the complete degradation of SM and yields trunSM (Fig. 4). Beyond these mechanistic insights, we found that truncation enables SM to partition to lipid droplets (Fig. 5) and is dramatically increased in endometrial cancer tissues (Fig. 6). Taken together, our results point towards the truncation of SM as an adaptive mechanism to promote squalene clearance during hypoxia, as well as a likely contributor to the widely reported oncogenic properties of SM.

Cholesterol synthesis during hypoxia

Hypoxia places great strain on metabolic processes and necessitates the strict allotment of available oxygen and energy reserves. Cholesterol synthesis is a particularly resource-intensive pathway, requiring eleven oxygen molecules and over one hundred ATP equivalents per molecule of product, yet there are conflicting reports on changes in overall flux from acetyl-CoA to cholesterol during hypoxia [24, 25], indicative of cell type-specific responses. Nevertheless, the small number of studies into individual biosynthetic enzymes indicate that their activity is suppressed by hypoxia at multiple regulatory levels, which is supported by the accumulation of various pathway intermediates [26, 27, 58]. Lanosterol demethylase, which requires three oxygen molecules for catalysis, is transcriptionally downregulated by HIF2α and the hypoxia-induced long non-coding RNA *lincNORS*, contributing to the characteristic accumulation of lanosterol under hypoxic conditions [26, 59]. Lanosterol in turn triggers the ubiquitin-dependent degradation of the early cholesterol synthesis enzyme HMGCR [27], suppressing further oxygen consumption by the pathway. Our study expands this understanding of hypoxic adaptations by establishing that oxygen availability also regulates SM, a rate-limiting enzyme of cholesterol synthesis and the first to require molecular oxygen.

We found that SM is transcriptionally downregulated in hypoxic HEK293T cells but not MDA-MB-231 cells, consistent with previously reported cell-type specific changes in *SQLE* expression [23]. This accounted in part for the decline in full-length SM levels, although reduced *SQLE* translation through mechanisms such as mTOR suppression [60] cannot be ruled out as a contributing factor. We also found that, like HMGCR, the translated SM protein is targeted for proteasomal degradation during hypoxia. However, in stark contrast to HMGCR, the net effect of SM degradation under these conditions is the maintenance of, or even an increase in, the number of enzyme molecules available for catalysis. This is due to the increased partial proteolysis of SM to form trunSM, which lacks degron features within the SM-N100 regulatory domain and has a dramatically extended half-life yet remains catalytically active

[9]. Disruption of the SM-N100 domain also renders trunSM resistant to cholesterol-induced
degradation, which is an key element in the metabolic regulation of full-length SM [4, 9].
Therefore, the hypoxia-induced truncation of SM ensures that total enzyme levels remain
constant even under cholesterol-replete conditions that would typically reduce its abundance.

The most plausible explanation for the seemingly paradoxical stabilization of SM
during hypoxia is that this can compensate for reduced oxygen availability and allow enzyme
activity to be maintained (Fig. 7). There are numerous ways in which the truncation and
constitutive activity of SM is beneficial. During transient or low-level hypoxia, it may enable
continued cholesterol synthesis so that cell growth and survival is not compromised by the
oxygen shortfall. In support of this idea, hypoxia-induced cell death is exacerbated by SM
inhibition [23]. Furthermore, the longevity of trunSM is likely to enable rapid resumption of
pathway activity when normal oxygen levels are restored. During long-term or severe hypoxia,
where there is insufficient oxygen for flux through downstream cholesterol synthesis, the role
of trunSM in hypoxia may shift towards the efficient clearance of squalene. While generally
considered an inert intermediate, excess squalene induces ER stress and is generally toxic in
cells lacking SM activity [47]. A secondary effect of squalene clearance is its downstream
conversion to lanosterol, which promotes the shutdown of cholesterol synthesis *via* HMGCR
degradation. This mechanism may explain why squalene accumulation lags behind that of
lanosterol during the acute phase of hypoxia [27, 58]. Although we detected increased squalene
levels at early hypoxic timepoints, a previous study of sterol-depleted cells did not [27], which
is likely attributable to the basal upregulation of SM expression under these conditions.
Elevated SM activity during hypoxia may also favor the synthesis of dioxidosqualene and
ultimately 24(*S*),25-epoxycholesterol, a potent suppressor of cholesterol accretion [61].
However, this oxysterol is yet to be studied in a low-oxygen context.

Molecular mechanism of truncation

The first step of SM truncation is its delivery to the proteasome [9]. This is enhanced
during hypoxia by a novel ubiquitin signal at the Lys-82/90/100 cluster, as well as the
accumulation of the E3 ubiquitin ligase MARCHF6, which facilitates the basal and
metabolically-regulated degradation of SM [36]. Non-canonical ubiquitination sites required
for cholesterol-induced degradation of SM [33] also contribute to its hypoxic degradation,
which can be reconciled by the combination of increased MARCHF6 levels and the sterol-
replete conditions under which these experiments were performed. Hypoxic accumulation of
MARCHF6 occurs due to reduced auto-ubiquitination, as this effect is blocked by inhibition
of the ER-associated degradation pathway or by mutation of residues essential for MARCHF6
activity. The precise mechanism by which low oxygen levels impede MARCHF6
auto-ubiquitination remains unclear, although the involvement of asparagine hydroxylation or
accumulated squalene was ruled out. We previously found that cholesterol and several of its
biosynthetic intermediates, including lanosterol, also impede MARCHF6 degradation [37].
This raises the possibility that a lanosterol-dependent mechanism promotes MARCHF6
accumulation during hypoxia, which is particularly intriguing given that MARCHF6 also
targets lanosterol demethylase for degradation [62] and likely contributes to lanosterol

accumulation. It remains to be determined how increased MARCHF6 levels influence the
activity of its other metabolic substrates. These include the lipid droplet biogenesis protein
perilipin-2 [63] and the thyroid hormone-activating deiodinase DIO2 [64], which both function
in pathways perturbed by hypoxia [48, 65]. Unexpectedly, MARCHF6 levels were slightly but
significantly reduced upon treatment with exogenous squalene. This was also observed for the
saturated analogue squalane, and therefore is not a squalene-specific effect. Rather, it may
suggest that the stability of the 14-transmembrane domain MARCHF6 protein is sensitive to
membrane perturbations resulting from the delivery of large amounts of exogenous lipid.

A key finding of this study is that upon the hypoxia-induced delivery of SM to the
proteasome, accumulated squalene inhibits its complete degradation and liberates the
constitutively active trunSM. This feedforward mechanism, by which the substrate of SM
preserves its activity, is mediated by the SM-N100 regulatory domain and functional under
normoxic conditions. Here, it likely buffers against transient increases in squalene levels.
Accumulation of trunSM occurs even when MARCHF6 is depleted, indicating that squalene
similarly prevents the complete degradation of SM molecules targeted to the proteasome by
other E3 ubiquitin ligases. These additional regulators, and the factors controlling their
ubiquitination of SM, await future discovery and may shed light on how the SM-N100
regulatory domain undergoes MARCHF6-independent proteasomal degradation during
hypoxia. Although hypoxia and lipid accumulation can trigger ER stress and altered
proteostasis [66], the ability of squalene and farnesyl-containing compounds to allosterically
bind the SM-N100 domain [8] strongly indicates that squalene induces SM truncation directly,
rather than *via* bulk membrane effects. The specificity of this response is further supported by
the inability of its saturated analog squalane to induce truncation. While squalene is the most
likely of the farnesyl-containing cholesterol synthesis intermediates to accumulate under
normal conditions, owing to the rate-limiting and oxygen-dependent nature of SM activity, our
current and previous [8] data suggest that a farnesyl group alone is sufficient to promote
truncation. Precisely how this lipophilic moiety interferes with SM degradation is unknown,
although interaction with the membrane-embedded SM-N100 re-entrant loop [67] is a likely
possibility. SM also contains a short intrinsically disordered domain (residues 81–120) that is
required for its partial degradation and dictates the location of the truncation site [9], although
interaction of a farnesyl group with this region is more difficult to envisage.

Previously, we reported that squalene and farnesyl-containing compounds also blunt
the MARCHF6-mediated ubiquitination of the SM-N100 domain, leading to stabilization of
full-length SM [8]. The existence of dual mechanisms enabling squalene to sustain SM activity
is intriguing, particularly as trunSM formation is itself dependent on ubiquitination [9]. One
possibility is that truncation is stimulated at a lower squalene threshold than the inhibition of
SM ubiquitination, allowing for a biphasic response to accumulating substrate. Another
possibility is that truncation is a 'failsafe' mechanism to preserve the activity of the SM
enzyme. This may be necessary when a reduced frequency of MARCHF6 ubiquitination is
insufficient to completely prevent targeting of SM to the proteasome. It may also apply in
situations where SM ubiquitination is promoted by other cellular stimuli. For instance, during

hypoxia, the stabilization of MARCHF6 may override the inhibitory effects of accumulated
squalene.

Pathophysiological consequences of truncation

Beyond hypoxia, our results point to additional contexts in which SM truncation has
physiological relevance. One of these is lipid droplets, which serve important functions in
triglyceride and cholesteryl ester storage as well as the sequestration of lipotoxic compounds,
including excess squalene [47, 68]. Lipid droplets can also serve as energy depots for fueling
cancer cell growth [69]. Although several proteomic studies have identified SM in lipid droplet
fractions [49–51], the detected peptides were unable to distinguish between its full-length and
truncated forms. Using immunofluorescence as well as immunoblotting, which remains the
most robust means of differentiating the two proteins, we found that only trunSM is capable of
localizing to lipid droplets. The absence of the membrane-embedded SM-N100 re-entrant loop,
which allows trunSM to adopt a peripheral ER membrane topology [9], also accounts for its
movement to the lipid droplet monolayer. By contrast, the C-terminal membrane-associated
helices of SM [44] appear to have a highly conserved role in lipid droplet localization, given
that they are also required for the partitioning of the yeast orthologue Erg1p [70, 71].

Further study is required to determine the functional consequences of lipid droplet-
localized trunSM. Curiously, the yeast orthologue of SM is inactive on lipid droplets due to the
absence of a microsomal electron donor [71]. Should this lack of activity hold for trunSM, lipid
droplets may nevertheless serve as a depot of stable enzyme that is inaccessible to polytopic
ER-associated degradation cofactors such as MARCHF6 [38], yet able to return to the ER *via*
contact sites [72, 73]. It is interesting to note that several other cholesterol synthesis enzymes
also partition to lipid droplets, most notably lanosterol synthase, the enzyme immediately
downstream of SM [49–51]. The yeast orthologue of this protein remains active on lipid
droplets [74], but whether this holds true in mammalian cells is unknown. The shuttling of
cholesterol synthesis intermediates such as squalene and lanosterol between the ER and lipid
droplets is an unexplored area with the potential for major consequences in the metabolic
feedback regulation of the pathway.

Both hypoxia and squalene accumulation trigger lipid droplet formation [48, 75], and
it is likely that these two phenomena are linked. As SM truncation occurs in the ER, the amount
of trunSM produced under squalene-rich conditions may be coupled with the ability of a cell
to adequately sequester this intermediate to lipid droplets. In cells with a low capacity for lipid
droplet biogenesis, rapid accumulation of squalene in the ER is expected to induce SM
truncation and facilitate its clearance. In cells with a greater capacity for lipid droplet
biogenesis, accumulation of squalene in the ER is likely to be less dramatic, resulting in a lower
level of truncation that is sufficient to clear this smaller quantity. Supporting this idea, a high
rate of lipid droplet biogenesis confers resistance to the toxic effects of SM inhibition in
neuroendocrine cancer cell lines [47].

Overexpression and overactivity of SM occurs in a wide range of malignancies
including hepatocellular carcinoma and prostate cancer, where it is positively correlated with

severity and lethality [21, 53]. Despite this, a specific examination of trunSM levels had not
been performed in any cancer type prior to our study. We found that total SM levels were
reduced in endometrial tumor tissues, which was unexpected given that high *SQLE* mRNA
expression is a risk factor for progression of this cancer [76]. However, the reduction in total
SM was accompanied by a marked increase in the proportion of the stable and constitutively
active trunSM. This was also well-correlated with the levels of HIF1α, suggesting that hypoxia-
induced truncation of SM occurs *in vivo*. An increased proportion of trunSM was observed in
endometrial tumors from both lean and obese patients, indicating that the phenomenon is likely
not influenced by obesity or circulating lipid levels. Our data indicate that trunSM may be a
contributing factor to cell growth and survival in the hypoxic tumor microenvironment, which
is noteworthy given that severe hypoxia is often associated with a poorer cancer prognosis [22].
It would be worthwhile for future studies to quantify the truncation of SM in cancer types
where its elevated protein expression has previously been identified, as well as in those that
show a particular propensity for hypoxia, such as prostate and pancreatic cancer [28]. Although
trunSM retains full catalytic activity and can likely fulfil the cholesterol synthesis-dependent
roles of SM in oncogenesis, further work is needed to confirm that it can enable the suite of
other oncogenic functions previously reported for full-length SM. To date, these include the
activation of ERK signaling [18] as well as interactions with carbonic anhydrase 3 [20], GSK3β
and p53 [19]. These cholesterol-independent effects of SM are likely to be most important
during severe hypoxia, where cholesterol synthesis cannot proceed.

Finally, although this study was largely performed using the extremely low oxygen
levels characteristic of pathophysiological hypoxia, we observed that SM truncation was
responsive to a range of 'physoxic' oxygen levels that occur within normal tissues *in situ*. This
reinforces the concept of truncation as a buffering mechanism to accommodate normal
substrate fluctuations, which may be important for preventing squalene-induced dysfunction in
tissues with lower oxygen perfusion, such as the brain [28]. Squalene fluctuations also occur
diurnally [77] and with changing sterol status [4], and its levels can vary dramatically between
different tissues [78]. Thus, the ability of squalene to stimulate formation of the rate-limiting
trunSM may play an important role in regulating flux through cholesterol synthesis under
varying physiological conditions.

Figure 7. Model of hypoxia-induced SM truncation.

(A, B) Hypoxic conditions inhibit the auto-ubiquitination of the E3 ubiquitin ligase MARCHF6, resulting in its stabilization. Simultaneously, the SM-catalyzed conversion of squalene to monooxidosqualene is inhibited, leading to squalene accumulation. **(C)** Increased MARCHF6 activity promotes the targeting of SM to the proteasome. **(D)** Squalene binds the SM-N100 domain and impedes the complete proteasomal degradation of SM, leading to **(E)** the formation of the constitutively active trunSM, which is capable of localizing to lipid droplets. During transient or low-level hypoxia, trunSM activity may facilitate continued cholesterol synthesis to compensate for the oxygen shortfall. During long-term or severe

700   hypoxia, trunSM activity may reduce squalene-induced toxicity while also promoting the
701   formation of lanosterol, which suppresses an early step of the cholesterol synthesis pathway.
702   During pathophysiological hypoxia, trunSM may enable continued cell growth and survival
703   *via* both cholesterol synthesis-dependent and -independent oncogenic properties.

Materials and methods

Reagents and cell lines

Fetal calf serum (FCS), newborn calf serum (NCS), high-glucose Dulbecco's Modified Eagle's Medium (DMEM-HG), penicillin/streptomycin, Opti-MEM reduced serum medium, RNAiMAX transfection reagent, Lipofectamine 3000 transfection reagent, TRI reagent, the SuperScript III First-Strand Synthesis kit, the PureLink Genomic DNA Mini kit, HCS LipidTOX Deep Red Neutral Lipid Stain, and BODIPY 493/503 were from Thermo Fisher (Carlsbad, CA, USA). Lipoprotein-deficient serum (LPDS, 30 mg/mL protein) was prepared from FCS by density gradient centrifugation and dialysis, as described in [79]. Primers, small interfering RNA (siRNA), protease inhibitor cocktail, and Tween-20 were from Sigma-Aldrich (St Louis, MO, USA). Tris-glycine SDS-PAGE gels were prepared in-house. Immobilon Western chemiluminescent HRP substrate and nitrocellulose membranes were from Millipore (Burlington, MA, USA). The QuantiNova SYBR Green PCR kit was from Qiagen (Hilden, GE). Skim milk powder was from Fonterra (Richmond, VIC, AU). The Lipid Droplet Isolation Kit was from Cell Biolabs Inc. (San Diego, CA, USA). Chemicals were from the following suppliers: 2,3,22,23-dioxidosqualene (Echelon Biosciences S-0302), 2,3-oxidosqualene (monooxidosqualene; Echelon Biosciences S-0301), 5α-cholestane (Sigma-Aldrich C8003), bafilomycin A1 (Sigma-Aldrich B1793), BIBB 515 (Cayman Chemical 10010517), CB-5083 (Cayman Chemical 16276), DM-NOFD (Sigma-Aldrich SML1874), DMOG (Sigma-Aldrich D3695), FG-4592 (Cayman Chemical 15294), GR70585X (GlaxoSmithKlein), mevalonolactone (Sigma-Aldrich M4667), mevastatin (Sigma-Aldrich M2537), MG132 (Sigma-Aldrich C2211), NB-598 (Chemscene CS-1274), oleate-bovine serum albumin complexes (Sigma-Aldrich O3008), squalane (Sigma-Aldrich 234311), squalene (Sigma-Aldrich S3626), TAK-475 (Sigma-Aldrich SML2168).

HEK293T cells were a gift from the UNSW School of Medical Sciences (UNSW Sydney, NSW, Australia), HCT116 cells were a gift from Dr Ewa Goldys (UNSW Sydney, NSW, Australia), Huh7 cells were a gift from the Centre for Cardiovascular Research (UNSW Sydney NSW, Australia), MBA-MB-231 and HeLaT cells were gifts from Dr Louise Lutze-Mann and Dr Noel Whitaker (UNSW Sydney, NSW, Australia), and HeLa cells were obtained from the American Type Culture Collection. HEK293 cells stably expressing SM-N100-GFP-V5, MARCHF6-V5, MARCHF6-V5 C9A, or MARCHF6-V5 N890A were generated previously using the Flp-In T-Rex system [36, 37].

Cell culture

Cells were maintained in a humidified Heraeus BB 15 incubator at 37 °C, 5% CO_2, and 21% O_2 (normoxia) in maintenance medium (DMEM-HG, 10% [v/v] FCS, 100 U/mL penicillin, 100 μg/mL streptomycin). To improve HEK293 and HEK293T cell surface adhesion, culture vessels were treated with 25 μg/mL polyethyleneimine in phosphate-buffered saline (PBS) for 15 min at 37 °C prior to cell seeding. Plasmid and siRNA transfections were performed in maintenance medium lacking penicillin and streptomycin. Sterol depletions were

performed in maintenance medium containing LPDS. Hypoxic incubations at 0.5–10% O_2
were performed in a humidified Binder CB 150 incubator at 37 °C and 5% CO_2. For all
treatments, appropriate solvent controls were used (water [DMOG]; ethanol (mevalonolactone,
GR70585X); dimethyl sulfoxide [CB-5083, MG132, bafilomycin A1, DM-NOFD, mevastatin,
TAK-475, BIBB-515, NB-598]; dimethyl sulfoxide containing 1% [v/v] Tween-20 [squalene,
monooxidosqualene, dioxidosqualene, squalane]) and the final concentration of solvent did not
exceed 1% (v/v) in cell culture medium. Treatments were delivered in full medium refreshes,
and all experiments were 48–72 h in duration.

Plasmids

Plasmids encoding Cas9 and *SQLE*-targeting guide RNAs were generated by *BbsI*
cloning into a PX458 vector as described in [80]. Amino acid substitutions (SM Y195F) within
expression vectors were generated using the overlap extension cloning method, as described
previously [81]. The identity of all plasmids was confirmed *via* Sanger dideoxy sequencing.
The plasmids used in this study are listed in Supplementary Table 1, and the primer sequences
used for DNA cloning are listed in Supplementary Table 2.

siRNA and plasmid transfection

To downregulate gene expression or transiently overexpress proteins, cells were seeded
into 12-well plates. The next day, cells were transfected with 15 pmol siRNA using RNAiMAX
(Invitrogen; 15 pmol siRNA : 2 µL reagent) or 1 µg expression vector using Lipofectamine
3000 (Invitrogen; 1 µg DNA : 2 µL reagent with 2 µL P3000 supplemental reagent), delivered
in Opti-MEM. After 24 h, cells were refreshed in maintenance medium and treated as specified
in figure legends. The siRNAs and plasmids used in this study are listed in Supplementary
Table 1.

Protein harvest and immunoblotting

To quantify protein levels, cells were seeded into 6- or 12-well plates and treated as
specified in figure legends. For detection of SM, total protein was harvested in 2% SDS lysis
buffer (10 mM Tris-HCl [pH 7.6], 2% [w/v] SDS, 100 mM NaCl, 2% [v/v] protease inhibitor
cocktail), passed through a 21-gauge needle until homogenous, and vortexed at room
temperature for 20 min. For detection of MARCHF6, cells were scraped in ice-cold PBS,
pelleted, and lysed in modified RIPA buffer (50 mM Tris-HCl [pH 8.0], 0.1% [w/v] SDS,
1.5% [w/v] IGEPAL CA-630, 0.5% [w/v] sodium deoxycholate, 150 mM NaCl, 2 mM $MgCl_2$,
2% [v/v] protease inhibitor cocktail), passed 20 times through a 22-gauge needle, rotated at
4 °C for 30 min, and centrifuged at 17,000 *g* and 4 °C for 15 min to obtain the supernatant.
Lysate protein content was quantified using the bicinchoninic acid assay (Thermo Fisher), and
sample concentrations were normalized by dilution in the appropriate lysis buffer and 0.25 vol
5× Laemmli buffer (250 mM Tris-HCl [pH 6.8], 10% [w/v] SDS, 25% [v/v] glycerol,

0.2% [w/v] bromophenol blue, 5% [v/v] β-mercaptoethanol). For SM detection, normalized

samples were heated at 95°C for 5 min.

Proteins were separated by 10% (w/v) Tris-glycine SDS-PAGE, electroblotted onto

nitrocellulose membranes, and blocked in 5% (w/v) skim milk powder in PBS containing

0.1% (v/v) Tween-20 (PBST). Immunoblotting was performed using the following antibodies:

rabbit polyclonal anti-HIF1α (Proteintech 20960-1-AP; 1:1,000 at room temperature for 1 h),

rabbit polyclonal anti-SM(SQLE) (Proteintech 12544-1-AP; 1:2,500 at 4 °C for 16 h), rabbit

monoclonal anti-GAPDH (Cell Signaling Technology 2118; 1:2,000 at 4 °C for 16 h), mouse

monoclonal anti-V5 (Thermo Fisher R960-25; 1:5,000 at room temperature for 1 h), rabbit

monoclonal anti-HA (Cell Signaling Technology 3724; 1:2,000 at 4 °C for 16 h), rabbit

monoclonal anti-calnexin (Cell Signaling Technology 2679; 1:1,000 at 4 °C for 16 h), rabbit

polyclonal anti-ABHD5 (Proteintech 12201-1-AP; 1:1,000 at 4 °C for 16 h), mouse

monoclonal anti-pan-14-3-3 (Santa Cruz Biotechnology sc-1657 ; 1:1,000 at 4 °C for 16 h),

IRDye 680RD donkey anti-rabbit IgG (LI-COR Biosciences LCR-926-68073; 1:5,000), IRDye

800CW donkey anti-mouse IgG (LI-COR Biosciences LCR-926-32212; 1:10,000 at room

temperature for 1 h), peroxidase-conjugated AffiniPure donkey anti-rabbit IgG (Jackson

ImmunoResearch Laboratories 711-035-152; 1:10,000 at room temperature for 1 h), and

peroxidase-conjugated AffiniPure donkey anti-mouse IgG (Jackson ImmunoResearch

Laboratories 715-035-150; 1:10,000 at room temperature for 1 h). All antibodies were diluted

in 5% (w/v) bovine serum albumin in PBST containing 0.02% (w/v) sodium azide, except for

anti-HIF1α and peroxidase-conjugated antibodies, which were diluted in 5% (w/v) skim milk

powder in PBST. Fluorescence-based detection of SM and trunSM, GAPDH, and $(HA)_3$-

SM-V5 was performed using an Odyssey CLx imager (LI-COR Biosciences), and enhanced

chemiluminescence-based detection of HIF1α, MARCHF6-V5, SM and trunSM (fractionated

and tissue-derived lysates), calnexin, ABHD5, GAPDH (fractionated lysates) and 14-3-3 was

performed using Immobilon Western chemiluminescent HRP substrate (Millipore) and an

ImageQuant LAS 500 imager (Cytiva Life Sciences). Densitometry analysis was performed

using Image Studio Lite v5.2.5 (LI-COR Biosciences).

RNA harvest and qRT-PCR

To quantify gene expression, cells were seeded into 12-well plates and treated as

specified in figure legends. Total RNA was harvested using TRI reagent (Sigma-Aldrich) and

polyadenylated RNA was reverse transcribed using the SuperScript III First Strand Synthesis

kit (Invitrogen). cDNA products were used as the template for quantitative reverse

transcription-PCR (qRT-PCR) in technical triplicate using the QuantiNova SYBR Green PCR

kit (Qiagen). mRNA levels were normalized to the geometric mean of *RPL11*, *GAPDH* and

ACTB for hypoxia experiments, or *PBGD* for validation of siRNA knockdowns and gene

knockout, using the comparative C_T method [82]. Normalized data were adjusted relative to

the control condition, as specified in figure legends. The primer sequences used for qRT-PCR

in this study are listed in Supplementary Table 2.

Gas chromatography-mass spectrometry

To quantify cellular squalene levels, gas chromatography-mass spectrometry was performed as described previously [8]. Briefly, cells in 6-well plates were lysed in 0.05 M NaOH, total protein was quantified using the bicinchoninic acid assay, and samples were adjusted to the lowest protein concentration using 0.05 M NaOH plus 4 µg 5α-cholestane as an internal standard, in a total volume of 1 mL. Lysates were mixed with 1 mL 100% (v/v) ethanol, 500 µL 75% (w/v) KOH, 1 µL 20 mM butylated hydroxytoluene, and 20 µL 20 mM EDTA, and saponified at 70 °C for 1 h. Non-saponifiable lipids were extracted by mixing with 1 mL 100% (v/v) ethanol and 2.5 mL hexane, centrifuging at 4,000 g for 5 min, and collection of 2 mL of the organic phase. Lipids were dried in a vacuum centrifuge, resuspended in 50 µL N,O-bis(trimethylsilyl)trifluoroacetamide, and derivatized at 60 °C for 1 h.

Derivatized lipids (1.5 µL) were injected *via* a heated (300 °C) splitless with surge (38.0 psi for 0.50 min) inlet into a Thermo Trace gas chromatograph fitted with a Trace TR-50MS GC column (60 m × 0.25 mm, 0.25 µm film thickness, Thermo Fisher). Analytes were separated with helium as the carrier gas at a constant flow of 1.2 ml/min with vacuum compensation, and temperature programming as follows: 70 °C for 0.70 min, 20 °C/min to 250 °C, 3 °C/min to 270 °C, 1.5 °C/min to 315 °C, then hold for 10 min. The GC column was coupled to a Thermo DSQIII mass spectrometer, with a transfer line temperature of 320 °C and an ion source temperature of 250 °C. For mass spectrometry analysis, the electron energy was 70 eV, the emission current was 130 µA and the detector gain was 3.0×10^5. Squalene and 5α-cholestane standards were analyzed in scan mode (34–600 Da) to identify peaks and retention times, and identity was confirmed using the National Institute of Standards and Technology databases. Experimental samples were analyzed in selective ion monitoring mode to detect squalene ($m/z = 81.0$, 410.4) and 5α-cholestane ($m/z = 149.1$, 217.2, 372.4), with a detection width of 0.1 and dwell time of 200 ms. Chromatograph peaks were integrated using Thermo Xcalibur software (v2.2 SP1.48) and the peak area of squalene was normalized to the 5α-cholestane internal standard. A squalene standard curve ranging from 3.125–100 ng/µL was used to quantify squalene levels, with data adjusted to the total protein content of the cell lysate.

Generation of SM knockout cells

To knock out SM using CRISPR/Cas9, guide RNAs targeting the *SQLE* proximal promoter (hg38 chr8:124998048–124998067; AATGGAAACGTTCCGACCCG) and first exon (hg38 chr8:124999588–124999607; ATCCGAGAAGAGGGCGAACT) were designed using CHOPCHOP [83] and cloned into the Cas9- and GFP-encoding PX598 vector using BbsI restriction enzyme cloning, as described in [80]. HEK293T cells were seeded into 10 cm dishes and transfected with 5 µg of both vectors using Lipofectamine 3000, as described above. After 24 h, cells were trypsinized, washed with PBS and resuspended in FACS buffer (5% [v/v] FCS and 10 mM EDTA in PBS). Cells were sorted based on GFP fluorescence using a BD FACSAria III at the UNSW Flow Cytometry Facility. GFP-positive cells were seeded into 10 cm dishes at a low density (6,000–12,000 cells/dish) and allowed to adhere for 2–3 weeks. Single colonies were picked, expanded, and screened for *SQLE* mRNA expression and SM

858   protein expression *via* qRT-PCR and immunoblotting, as described above. To confirm the
859   genomic lesion, genomic DNA was isolated from clones using the PureLink Genomic DNA
860   Mini kit (Thermo Fisher). The CRISPR/Cas9 target region was amplified and cloned into a
861   pcDNA3.1 vector for Sanger dideoxy sequencing. The primer sequences used for guide RNA
862   generation are listed in Supplementary Table 2.

863   Immunofluorescence and lipid droplet staining

To detect protein lipid droplet localization, cells were seeded onto coverslips in
6-well plates and treated as specified in figure legends. Cells were fixed with 4% (w/v)
paraformaldehyde for 15 min, permeabilized in 0.1% (w/v) Triton X-100 for 10 min and
blocked in 3% (w/v) bovine serum albumin in PBS for 1 h at room temperature. If required,
coverslips were incubated in anti-HA (Cell Signaling Technology 3724; 1:100) or anti-V5
(Thermo Fisher R960-25; 1:500) antibody diluted in blocking solution for 16 h at 4 °C,
followed by incubation in goat anti-rabbit Alexa Fluor 568 (Thermo Fisher A-11011; 1:1,000)
or goat anti-mouse Alexa Fluor 568 (Thermo Fisher A-11004; 1:1,000) in blocking solution
for 1 h at room temperature. To stain lipid droplets, coverslips were incubated in 1 µg/mL
BODIPY 493/503 in PBS for 10 min at room temperature, or HCS LipidTOX Deep Red
Neutral Lipid Stain (1:500 in PBS) for 1 h. Coverslips were mounted in ProLong Gold Antifade
Mountant with DAPI and confocal microscopy was performed using a Zeiss LSM 900 confocal
microscope system equipped with a 63×/1.4 oil objective (Zeiss, Jena, Germany). The
fluorochromes used were Alexa Fluor 488 (GFP, BODIPY 493/503), Alexa Fluor 568 (V5-
and HA-tagged proteins), Cy5 (LipidTOX Deep Red) and DAPI. Images were inspected using
Fiji software and lipid droplet association was quantified as the percentage of transfected cells
showing lipid droplet-proximal localization of the protein or epitope of interest.

881   Lipid droplet fractionation

To isolate lipid droplets, cells were seeded into 10 cm dishes and treated as specified in
figure legends. Cell lysates were collected, a small aliquot was taken for total protein analysis,
and the remainder was fractionated using the Lipid Droplet Isolation Kit (Cell Biolabs Inc.).
The floating lipid droplet fraction and an intermediate cytosolic fraction were collected, while
the pellet was resuspended in 2% SDS lysis buffer and passed through a 21-gauge needle until
homogenous. Total protein within each fraction was quantified using the bicinchoninic acid
assay and equal amounts were subjected to immunoblotting. Lipid droplet partitioning was
expressed as the ratio between protein levels in the lipid droplet and pellet fractions, and
normalized to that of the lipid droplet marker ABHD5.

891   Endometrial cancer tissues

Matched tumor and adjacent benign endometrial tissues were obtained from lean (body
mass index <27 kg/m^2; $n = 7$) and obese (body mass index >30 kg/m^2; $n = 7$) postmenopausal
endometrioid adenocarcinoma patients recruited at the Royal Hospital for Women and Prince

of Wales Private Hospital (Randwick, Sydney, Australia), as described in [84]. Consent was
received from all patients prior to sample collection, and all processing and experiments were
approved by the Human Research Ethics Committee (HREC) of the South Eastern Sydney
Local Health District (HREC 15/339). Protein was isolated from powdered tissues using the
AllPrep DNA/RNA/Protein Mini Kit (Qiagen). Protein pellets were lysed in HES-SDS buffer
(20 mM HEPES [pH 7.4], 2% [w/v] SDS, 250 mM sucrose, 1 mM EDTA) and mixed with
1 vol 10% (w/v) SDS. Samples were homogenized using a handheld homogenizer and
centrifuged at 15,000 g for 5 min. The protein content of the clarified supernatant was
quantified using the bicinchoninic acid assay. Samples were diluted in 0.25 vol 5× Laemmli
buffer and heated at 65 °C for 5 min prior to SDS-PAGE and immunoblotting.

Data analysis and presentation

Data were normalized as described in figure legends. All data were obtained from $n \geq 2$
independent experiments, and visualization and statistical testing were performed using
GraphPad Prism v9.0 (GraphPad Software Inc.) as specified in figure legends. Where multiple
statistical tests were conducted within a single experiment, p-values were corrected using the
Benjamini-Hochberg method [85] with a false discovery threshold of 5%. Thresholds for
statistical significance were defined as: *, $p \leq 0.05$; **, $p \leq 0.01$. Values of $0.05 < p < 0.075$
are indicated in text on graphs. Figures were assembled using Adobe Illustrator v26.3 (Adobe
Inc.).

Acknowledgements

We thank Isabelle Capell-Hattam for assistance with CRISPR/Cas9, Dr Martin
Bucknall for technical assistance with GC/MS, and the members of the Brown laboratory for
critically reviewing this manuscript. We are also grateful to the patients from whom tissue
samples were collected, and staff at the Royal Hospital for Women and Prince of Wales Private
Hospital (Randwick, NSW, Australia) who assisted with surgical procedures. H.W.C. is
supported by an Australian Research Training Program scholarship, R.F. is supported by a
RANZCOG Mary Elizabeth Courier Research Scholarship, F.L.B. is supported by a Cancer
Institute NSW Career Development Fellowship (2021/CDF1120), and the Brown laboratory is
supported by Australian Research Council Grant DP170101178 and an NSW Health
Investigator Development Grant.

Author contributions

H.W.C.: conceptualization, methodology, investigation, visualization, writing –
original draft. E.M.O.: investigation. X.D.: methodology, investigation, visualization. R.F.:
resources, funding acquisition. H.Y.: resources, funding acquisition. F.L.B.: resources, writing

– review and editing, funding acquisition. A.J.B.: conceptualization, methodology, resources,
writing – review and editing, supervision, funding acquisition.

Competing interests

The authors declare that there are no competing interests associated with the
manuscript.

References

1. Baigent, C., Blackwell, L., Emberson, J., Holland, L. E., Reith, C., Bhala, N., Peto, R.,
Barnes, E. H., Keech, A., Simes, J., Collins, R., De Lemos, J., Braunwald, E., Blazing,
M., Murphy, S., Downs, J. R., Gotto, A., Clearfield, M., Holdaas, H., Gordon, D., Davis,
B., Koren, M., Dahlof, B., Poulter, N., Sever, P., Knopp, R. H., Fellström, B., Jardine,
A., Schmieder, R., Zannad, F., Goldbourt, U., Kaplinsky, E., Colhoun, H. M.,
Betteridge, D. J., Durrington, P. N., Hitman, G. A., Fuller, J., Neil, A., Wanner, C.,
Krane, V., Sacks, F., Moyé, L., Pfeffer, M., Hawkins, C. M., Barter, P., Tavazzi, L.,
Maggioni, A., Marchioli, R., Tognoni, G., Franzosi, M. G., Bloomfield, H., Robins, S.,
Armitage, J., Parish, S., Sleight, P., Pedersen, T. R., Ridker, P. M., Holman, R., Meade,
T., MacMahon, S., Marschner, I., Tonkin, A., Shaw, J., Serruys, P. W., Nakamura, H.,
Knatterud, G., Furberg, C., Byington, R., Macfarlane, P., Cobbe, S., Ford, I., Murphy,
M., Blauw, G. J., Packard, C., Shepherd, J., Kjekshus, J., Wilhelmsen, L., Cannon, C.,
Bowman, L., Baxter, A., Landray, M., La Rosa, J., Rossouw, J., Probstfield, J., Flather,
M., Kastelein, J., Newman, C., Shear, C., Tobert, J., Varigos, J., White, H., Yusuf, S.,
Mellies, M., McGovern, M., Barclay, J., Belder, R., Mitchel, Y., Musliner, T., Ansquer,
J. C., Llewellyn, M., Bortolini, M., Brandrup-Wognsen, G., Bryzinski, B., Olsson, G.,
Pears, J., De Micco, D., Buck, G., Herrington, W. G., Kearney, P. M., Kirby, A., Lewis,
D. A., Pollicino, C., and Sourjina, T. (2010) Efficacy and safety of more intensive
lowering of LDL cholesterol: a meta-analysis of data from 170 000 participants in 26
randomised trials. *Lancet*. **376**, 1670–1681

2. Brown, A. J., Coates, H. W., and Sharpe, L. J. (2021) Cholesterol synthesis. in
Biochemistry of Lipids, Lipoproteins and Membranes, 7th Ed. (Ridgway, N. D., and
McLeod, R. S. eds), pp. 317–355, Elsevier Science

3. Porter, F. D., and Herman, G. E. (2011) Malformation syndromes caused by disorders
of cholesterol synthesis. *J. Lipid Res.* **52**, 6–34

4. Gill, S., Stevenson, J., Kristiana, I., and Brown, A. J. (2011) Cholesterol-dependent
degradation of squalene monooxygenase, a control point in cholesterol synthesis beyond
HMG-CoA reductase. *Cell Metab.* **13**, 260–273

5. Bai, M., Xiao, X., and Prestwich, G. (1992) Epoxidation of 2,3-oxidosqualene to
2,3;22,23-squalene dioxide by squalene epoxidase. *Biochem. Biophys. Res. Commun.*
185, 323–329

6. Horton, J. D., Shah, N. A., Warrington, J. A., Anderson, N. N., Park, S. W., Brown, M.
S., and Goldstein, J. L. (2003) Combined analysis of oligonucleotide microarray data

from transgenic and knockout mice identifies direct SREBP target genes. *Proc. Natl.*
Acad. Sci. **100**, 12027–12032

7.   Chua, N. K., Howe, V., Jatana, N., Thukral, L., and Brown, A. J. (2017) A conserved
degron containing an amphipathic helix regulates the cholesterol-mediated turnover of
human squalene monooxygenase, a rate-limiting enzyme in cholesterol synthesis. *J.*
Biol. Chem. **292**, 19959–19973

8.   Yoshioka, H., Coates, H. W., Chua, N. K., Hashimoto, Y., Brown, A. J., and Ohgane,
K. (2020) A key mammalian cholesterol synthesis enzyme, squalene monooxygenase,
is allosterically stabilized by its substrate. *Proc. Natl. Acad. Sci.* **117**, 7150–7158

9.   Coates, H. W., Capell-Hattam, I. M., and Brown, A. J. (2021) The mammalian
cholesterol synthesis enzyme squalene monooxygenase is proteasomally truncated to a
constitutively active form. *J. Biol. Chem.* **296**, 100731

10.   Kuzu, O. F., Noory, M. A., and Robertson, G. P. (2016) The role of cholesterol in cancer.
Cancer Res. **76**, 2063–2070

11.   Brown, D. N., Caffa, I., Cirmena, G., Piras, D., Garuti, A., Gallo, M., Alberti, S.,
Nencioni, A., Ballestrero, A., and Zoppoli, G. (2016) Squalene epoxidase is a bona fide
oncogene by amplification with clinical relevance in breast cancer. *Sci. Rep.* **6**, 19435

12.   Stopsack, K. H., Gerke, T. A., Sinnott, J. A., Penney, K. L., Tyekucheva, S., Sesso, H.
D., Andersson, S.-O. O., Cerhan, R., Giovannucci, E. L., Mucci, L. A., Rider, J. R.,
Andrén, O., Cerhan, J. R., Giovannucci, E. L., Mucci, L. A., and Rider, J. R. (2016)
Cholesterol metabolism and prostate cancer lethality. *Cancer Res.* **76**, 4785–4790

13.   Pudova, E. A., Krasnov, G. S., Kobelyatskaya, A. A., Savvateeva, M. V., Fedorova, M.
S., Pavlov, V. S., Nyushko, K. M., Kaprin, A. D., Alekseev, B. Y., Trofimov, D. Y.,
Sukhikh, G. T., Snezhkina, A. V., and Kudryavtseva, A. V. (2021) Gene expression
changes and associated pathways involved in the progression of prostate cancer
advanced stages. *Front. Genet.* **11**, 613162

14.   Bai, R., Rebelo, A., Kleeff, J., and Sunami, Y. (2021) Identification of prognostic lipid
droplet-associated genes in pancreatic cancer patients via bioinformatics analysis. *Lipids*
Health Dis. **20**, 58

15.   Parris, T. Z., Kovacs, A., Hajizadeh, S., Nemes, S., Semaan, M., Levin, M., Karlsson,
P., and Helou, K. (2014) Frequent MYC coamplification and DNA hypomethylation of
multiple genes on 8q in 8p11-p12-amplified breast carcinomas. *Oncogenesis.* **3**, e95

16.   Yang, F., Kou, J., Liu, Z., Li, W., and Du, W. (2021) MYC enhances cholesterol
biosynthesis and supports cell proliferation through SQLE. *Front. Cell Dev. Biol.* **9**,
655889

17.   Sun, H., Li, L., Li, W., Yang, F., Zhang, Z., Liu, Z., and Du, W. (2021) p53
transcriptionally regulates SQLE to repress cholesterol synthesis and tumor growth.
EMBO Rep. **22**, e52537

18.   He, L., Li, H., Pan, C., Hua, Y., Peng, J., Zhou, Z., Zhao, Y., and Lin, M. (2021)
Squalene epoxidase promotes colorectal cancer cell proliferation through accumulating
calcitriol and activating CYP24A1-mediated MAPK signaling. *Cancer Commun.* **41**,

726–746

19. Jun, S. Y., Brown, A. J., Chua, N. K., Yoon, J.-Y., Lee, J.-J., Yang, J. O., Jang, I., Jeon,
S.-J., Choi, T.-I., Kim, C.-H., and Kim, N.-S. (2021) Reduction of squalene epoxidase
by cholesterol accumulation accelerates colorectal cancer progression and metastasis.
Gastroenterology. **160**, 1194-1207.e28

20. Liu, D., Wong, C. C., Zhou, Y., Li, C., Chen, H., Ji, F., Go, M. Y. Y., Wang, F., Su, H.,
Wei, H., Cai, Z., Wong, N., Wong, V. W. S., and Yu, J. (2021) Squalene epoxidase
induces nonalcoholic steatohepatitis via binding to carbonic anhydrase 3 and is a
therapeutic target. *Gastroenterology*. 10.1053/j.gastro.2021.02.051

21. Liu, D., Wong, C. C., Fu, L., Chen, H., Zhao, L., Li, C., Zhou, Y., Zhang, Y., Xu, W.,
Yang, Y., Wu, B., Cheng, G., Lai, P. B.-S., Wong, N., Sung, J. J., and Yu, J. (2018)
Squalene epoxidase drives NAFLD-induced hepatocellular carcinoma and is a direct
pharmaceutical target. *Sci. Transl. Med.* **10**, eaap9840

22. Rankin, E. B., and Giaccia, A. J. (2016) Hypoxic control of metastasis. *Science*. **352**,
175–180

23. Haider, S., McIntyre, A., van Stiphout, R. G. P. M., Winchester, L. M., Wigfield, S.,
Harris, A. L., and Buffa, F. M. (2016) Genomic alterations underlie a pan-cancer
metabolic shift associated with tumour hypoxia. *Genome Biol.* **17**, 1–17

24. Mukodani, J., Ishikawa, Y., and Fukuzaki, H. (1990) Effects of hypoxia on sterol
synthesis, acyl-CoA:cholesterol acyltransferase activity, and efflux of cholesterol in
cultured rabbit skin fibroblasts. *Arteriosclerosis*. **10**, 106–110

25. Parathath, S., Mick, S. L., Feig, J. E., Joaquin, V., Grauer, L., Habiel, D. M., Gassmann,
M., Gardner, L. B., and Fisher, E. A. (2011) Hypoxia is present in murine atherosclerotic
plaques and has multiple adverse effects on macrophage lipid metabolism. *Circ. Res.*
109, 1141–1152

26. Wu, X., Niculite, C. M., Preda, M. B., Rossi, A., Tebaldi, T., Butoi, E., White, M. K.,
Tudoran, O. M., Petrusca, D. N., Jannasch, A. S., Bone, W. P., Zong, X., Fang, F.,
Burlacu, A., Paulsen, M. T., Hancock, B. A., Sandusky, G. E., Mitra, S., Fishel, M. L.,
Buechlein, A., Ivan, C., Oikonomopoulos, S., Gorospe, M., Mosley, A., Radovich, M.,
Davé, U. P., Ragoussis, J., Nephew, K. P., Mari, B., McIntyre, A., Konig, H., Ljungman,
M., Cousminer, D. L., Macchi, P., and Ivan, M. (2020) Regulation of cellular sterol
homeostasis by the oxygen responsive noncoding RNA lincNORS. *Nat. Commun.* **11**,
4755

27. Nguyen, A. D., McDonald, J. G., Bruick, R. K., and DeBose-Boyd, R. A. (2007)
Hypoxia stimulates degradation of 3-hydroxy-3-methylglutaryl-coenzyme A reductase
through accumulation of lanosterol and hypoxia-inducible factor-mediated induction of
insigs. *J. Biol. Chem.* **282**, 27436–27446

28. McKeown, S. R. (2014) Defining normoxia, physoxia and hypoxia in tumours-
implications for treatment response. *Br. J. Radiol.* **87**, 20130676

29. Dolt, K. S., Karar, J., Mishra, M. K., Salim, J., Kumar, R., Grover, S. K., and Qadar
Pasha, M. A. (2007) Transcriptional downregulation of sterol metabolism genes in
murine liver exposed to acute hypobaric hypoxia. *Biochem. Biophys. Res. Commun.* **354**,

148–153

30.   Cao, R., Zhao, X., Li, S., Zhou, H., Chen, W., Ren, L., Zhou, X., Zhang, H., and Shi, R.
1053   (2014) Hypoxia induces dysregulation of lipid metabolism in HepG2 cells via activation
1054   of HIF-2α. *Cell. Physiol. Biochem.* **34**, 1427–1441

1055   31.   Bellot, G., Garcia-Medina, R., Gounon, P., Chiche, J., Roux, D., Pouysségur, J., and
1056   Mazure, N. M. (2009) Hypoxia-induced autophagy is mediated through hypoxia-
1057   inducible factor induction of BNIP3 and BNIP3L via their BH3 domains. *Mol. Cell.*
1058   *Biol.* **29**, 2570–2581

1059   32.   Hornbeck, P. V., Zhang, B., Murray, B., Kornhauser, J. M., Latham, V., and Skrzypek,
1060   E. (2015) PhosphoSitePlus, 2014: mutations, PTMs and recalibrations. *Nucleic Acids*
1061   *Res.* **43**, D512–D520

1062   33.   Chua, N. K., Hart-Smith, G., and Brown, A. J. (2019) Non-canonical ubiquitination of
1063   the cholesterol-regulated degron of squalene monooxygenase. *J. Biol. Chem.* **294**, 8134–
1064   8147

1065   34.   Ivan, M., Kondo, K., Yang, H., Kim, W., Valiando, J., Ohh, M., Salic, A., Asara, J. M.,
1066   Lane, W. S., and Kaelin, J. (2001) HIFα targeted for VHL-mediated destruction by
1067   proline hydroxylation: implications for O2 sensing. *Science.* **292**, 464–468

1068   35.   Cockman, M. E., Lippl, K., Tian, Y. M., Pegg, H. B., Figg, W. D., Abboud, M. I., Heilig,
1069   R., Fischer, R., Myllyharju, J., Schofield, C. J., and Ratcliffe, P. J. (2019) Lack of
1070   activity of recombinant HIF prolyl hydroxylases (PHDs) on reported non-HIF
1071   substrates. *Elife.* 10.7554/ELIFE.46490

1072   36.   Zelcer, N., Sharpe, L. J., Loregger, A., Kristiana, I., Cook, E. C., Phan, L., Stevenson,
1073   J., and Brown, A. J. (2014) The E3 ubiquitin ligase MARCH6 degrades squalene
1074   monooxygenase and affects 3-hydroxy-3-methyl-glutaryl coenzyme A reductase and the
1075   cholesterol synthesis pathway. *Mol. Cell. Biol.* **34**, 1262–1270

1076   37.   Sharpe, L. J., Howe, V., Scott, N. A., Luu, W., Phan, L., Berk, J. M., Hochstrasser, M.,
1077   and Brown, A. J. (2019) Cholesterol increases protein levels of the E3 ligase MARCH6,
1078   and thereby stimulates protein degradation. *J. Biol. Chem.* **294**, 2436–2448

1079   38.   Hassink, G., Kikkert, M., Voorden, S. Van, Lee, S.-J., Spaapen, R., Laar, T. Van,
1080   Coleman, C. S., Bartee, E., Uh, K. F., Chau, V., and Wiertz, E. (2005) TEB4 is a C4HC3
1081   RING finger-containing ubiquitin ligase of the endoplasmic reticulum. *Biochem. J.* **388**,
1082   647–655

1083   39.   Zattas, D., Berk, J. M., Kreft, S. G., and Hochstrasser, M. (2016) A conserved C-terminal
1084   element in the yeast Doa10 and human MARCH6 ubiquitin ligases required for selective
1085   substrate degradation. *J. Biol. Chem.* **291**, 12105–12118

1086   40.   Lando, D., Peet, D. J., Gorman, J. J., Whelan, D. A., Whitelaw, M. L., and Bruick, R.
1087   K. (2002) FIH-1 is an asparaginyl hydroxylase enzyme that regulates the transcriptional
1088   activity of hypoxia-inducible factor. *Genes Dev.* **16**, 1466–1471

1089   41.   Scholz, C. C., Rodriguez, J., Pickel, C., Burr, S., Fabrizio, J. A., Nolan, K. A.,
1090   Spielmann, P., Cavadas, M. A. S., Crifo, B., Halligan, D. N., Nathan, J. A., Peet, D. J.,
1091   Wenger, R. H., Von Kriegsheim, A., Cummins, E. P., and Taylor, C. T. (2016) FIH

Regulates Cellular Metabolism through Hydroxylation of the Deubiquitinase OTUB1.
PLOS Biol. **14**, e1002347

42. McDonough, M. A., McNeill, L. A., Tilliet, M., Papamicaël, C. A., Chen, Q. Y., Banerji,
B., Hewitson, K. S., and Schofield, C. J. (2005) Selective inhibition of factor inhibiting
hypoxia-inducible factor. *J. Am. Chem. Soc.* **127**, 7680–7681

43. Hauß, T., Dante, S., Dencher, N. A., and Haines, T. H. (2002) Squalane is in the
midplane of the lipid bilayer: implications for its function as a proton permeability
barrier. *Biochim. Biophys. Acta - Bioenerg.* **1556**, 149–154

44. Padyana, A. K., Gross, S., Jin, L., Cianchetta, G., Narayanaswamy, R., Wang, F., Wang,
R., Fang, C., Lv, X., Biller, S. A., Dang, L., Mahoney, C. E., Nagaraja, N., Pirman, D.,
Sui, Z., Popovici-Muller, J., and Smolen, G. A. (2019) Structure and inhibition
mechanism of the catalytic domain of human squalene epoxidase. *Nat. Commun.* **10**, 97

45. Chua, N. K., Coates, H. W., and Brown, A. J. (2020) Squalene monooxygenase: a
journey to the heart of cholesterol synthesis. *Prog. Lipid Res.* **79**, 101033

46. Hong, Z., Liu, T., Wan, L., Fa, P., Kumar, P., Cao, Y., Prasad, C. B., Qiu, Z., Liu, J.,
Wang, H., Li, Z., Wang, Q.-E., Guo, P., Guo, D., Yilmaz, A. S., Lu, L., Papandreou, I.,
Jacob, N. K., Yan, C., Zhang, X., She, Q.-B., Ma, Z., and Zhang, J. (2022) Targeting
squalene epoxidase interrupts homologous recombination via the ER stress response and
promotes radiotherapy efficacy. *Cancer Res.* **82**, 1298–1312

47. Mahoney, C. E., Pirman, D., Chubukov, V., Sleger, T., Hayes, S., Fan, Z. P., Allen, E.
L., Chen, Y., Huang, L., Liu, M., Zhang, Y., McDonald, G., Narayanaswamy, R., Choe,
S., Chen, Y., Gross, S., Cianchetta, G., Padyana, A. K., Murray, S., Liu, W., Marks, K.
M., Murtie, J., Dorsch, M., Jin, S., Nagaraja, N., Biller, S. A., Roddy, T., Popovici-
Muller, J., and Smolen, G. A. (2019) A chemical biology screen identifies a vulnerability
of neuroendocrine cancer cells to SQLE inhibition. *Nat. Commun.* **10**, 96

48. Bensaad, K., Favaro, E., Lewis, C. A., Peck, B., Lord, S., Collins, J. M., Pinnick, K. E.,
Wigfield, S., Buffa, F. M., Li, J. L., Zhang, Q., Wakelam, M. J. O., Karpe, F., Schulze,
A., and Harris, A. L. (2014) Fatty acid uptake and lipid storage induced by HIF-1α
contribute to cell growth and survival after hypoxia-reoxygenation. *Cell Rep.* **9**, 349–
365

49. Hodges, B. D. M., and Wu, C. C. (2010) Proteomic insights into an expanded cellular
role for cytoplasmic lipid droplets. *J. Lipid Res.* **51**, 262–273

50. Pataki, C. I., Rodrigues, J., Zhang, L., Qian, J., Efron, B., Hastie, T., Elias, J. E., Levitt,
M., and Kopito, R. R. (2018) Proteomic analysis of monolayer-integrated proteins on
lipid droplets identifies amphipathic interfacial α-helical membrane anchors. *Proc. Natl.*
Acad. Sci. **115**, E8172–E8180

51. Bersuker, K., Peterson, C. W. H. W. H., To, M., Sahl, S. J. J., Savikhin, V., Grossman,
E. A. A., Nomura, D. K. K., and Olzmann, J. A. A. (2018) A proximity labeling strategy
provides insights into the composition and dynamics of lipid droplet proteomes. *Dev.*
Cell. **44**, 97–112

52. Du, X., Zhou, L., Aw, Y. C., Mak, H. Y., Xu, Y., Rae, J., Wang, W., Zadoorian, A.,
Hancock, S. E., Osborne, B., Chen, X., Wu, J. W., Turner, N., Parton, R. G., Li, P., and

Yang, H. (2020) ORP5 localizes to ER-lipid droplet contacts and regulates the level of
PI(4)P on lipid droplets. *J. Cell Biol.* **219**, e201905162

53.   Kalogirou, C., Linxweiler, J., Schmucker, P., Snaebjornsson, M. T., Schmitz, W., Wach,
S., Krebs, M., Hartmann, E., Puhr, M., Müller, A., Spahn, M., Seitz, A. K., Frank, T.,
Marouf, H., Büchel, G., Eckstein, M., Kübler, H., Eilers, M., Saar, M., Junker, K.,
Röhrig, F., Kneitz, B., Rosenfeldt, M. T., and Schulze, A. (2021) MiR-205-driven
downregulation of cholesterol biosynthesis through SQLE-inhibition identifies
therapeutic vulnerability in aggressive prostate cancer. *Nat. Commun.* **12**, 5066

54.   Bhaskaran, K., Douglas, I., Forbes, H., dos-Santos-Silva, I., Leon, D. A., and Smeeth,
L. (2014) Body-mass index and risk of 22 specific cancers: a population-based cohort
study of 5·24 million UK adults. *Lancet.* **384**, 755–765

55.   Onstad, M. A., Schmandt, R. E., and Lu, K. H. (2016) Addressing the role of obesity in
endometrial cancer risk, prevention, and treatment. *J. Clin. Oncol.* **34**, 4225–4230

56.   Berstein, L. M., Tchernobrovkina, A. E., Gamajunova, V. B., Kovalevskij, A. J.,
Vasilyev, D. A., Chepik, O. F., Turkevitch, E. A., Tsyrlina, E. V., Maximov, S. J.,
Ashrafian, L. A., and Thijssen, J. H. H. (2003) Tumor estrogen content and clinico-
morphological and endocrine features of endometrial cancer. *J. Cancer Res. Clin. Oncol.*
129, 245–249

57.   D'Agostino, R., and Pearson, E. S. (1973) Tests for departure from normality: empirical
results for the distributions of b2 and √b1. *Biometrika.* **60**, 613–622

58.   Kucharzewska, P., Christianson, H. C., and Belting, M. (2015) Global profiling of
metabolic adaptation to hypoxic stress in human glioblastoma cells. *PLoS One.* **10**,
e0116740

59.   Zhu, J., Jiang, X., and Chehab, F. F. (2014) FoxO4 interacts with the sterol regulatory
factor SREBP2 and the hypoxia inducible factor HIF2α at the CYP51 promoter. *J. Lipid*
Res. **55**, 431–442

60.   Liu, L., Cash, T. P., Jones, R. G., Keith, B., Thompson, C. B., and Simon, M. C. (2006)
Hypoxia-Induced Energy Stress Regulates mRNA Translation and Cell Growth. *Mol.*
Cell. **21**, 521–531

61.   Wong, J., Quinn, C. M., Gelissen, I. C., and Brown, A. J. (2008) Endogenous 24(S),25-
epoxycholesterol fine-tunes acute control of cellular cholesterol homeostasis. *J. Biol.*
Chem. **283**, 700–707

62.   Scott, N. A., Sharpe, L. J., Capell-Hattam, I. M., Gullo, S. J., Luu, W., and Brown, A. J.
(2020) The cholesterol synthesis enzyme lanosterol 14α-demethylase is post-
translationally regulated by the E3 ubiquitin ligase MARCH6. *Biochem. J.* **477**, 541–
555

63.   Nguyen, K. T., Lee, C.-S., Mun, S.-H., Nhung, T. T., Park, S. K., and Hwang, C.-S.
(2019) N-terminal acetylation and the N-end rule pathway control degradation of the
lipid droplet protein PLIN2. *J. Biol. Chem.* **294**, 379–388

64.   Zavacki, A. M., Arrojo E Drigo, R., Freitas, B. C. G., Chung, M., Harney, J. W., Egri,
P., Wittmann, G., Fekete, C., Gereben, B., and Bianco, A. C. (2009) The E3 ubiquitin

ligase TEB4 mediates degradation of type 2 iodothyronine deiodinase. *Mol. Cell. Biol.* **29**, 5339–47

65. Simonides, W. S., Mulcahey, M. A., Redout, E. M., Muller, A., Zuidwijk, M. J., Visser, T. J., Wassen, F. W. J. S., Crescenzi, A., Da-Silva, W. S., Harney, J., Engel, F. B., Obregon, M. J., Larsen, P. R., Bianco, A. C., and Huang, S. A. (2008) Hypoxia-inducible factor induces local thyroid hormone inactivation during hypoxic-ischemic disease in rats. *J. Clin. Invest.* **118**, 975–983

66. Bi, M., Naczki, C., Koritzinsky, M., Fels, D., Blais, J., Hu, N., Harding, H., Novoa, I., Varia, M., Raleigh, J., Scheuner, D., Kaufman, R. J., Bell, J., Ron, D., Wouters, B. G., and Koumenis, C. (2005) ER stress-regulated translation increases tolerance to extreme hypoxia and promotes tumor growth. *EMBO J.* **24**, 3470–3481

67. Howe, V., Chua, N. K., Stevenson, J., and Brown, A. J. (2015) The regulatory domain of squalene monooxygenase contains a re-entrant loop and senses cholesterol via a conformational change. *J. Biol. Chem.* **290**, 27533–27544

68. Olzmann, J. A., and Carvalho, P. (2019) Dynamics and functions of lipid droplets. *Nat. Rev. Mol. Cell Biol.* **20**, 137–155

69. Yue, S., Li, J., Lee, S. Y., Lee, H. J., Shao, T., Song, B., Cheng, L., Masterson, T. A., Liu, X., Ratliff, T. L., and Cheng, J.-X. X. (2014) Cholesteryl ester accumulation induced by PTEN loss and PI3K/AKT activation underlies human prostate cancer aggressiveness. *Cell Metab.* **19**, 393–406

70. Müllner, H., Zweytick, D., Leber, R., Turnowsky, F., and Daum, G. (2004) Targeting of proteins involved in sterol biosynthesis to lipid particles of the yeast Saccharomyces cerevisiae. *Biochim. Biophys. Acta - Biomembr.* **1663**, 9–13

71. Leber, R., Landl, K., Zinser, E., Ahorn, H., Spök, A., Kohlwein, S. D., Turnowsky, F., and Daum, G. (1998) Dual localization of squalene epoxidase, Erg1p, in yeast reflects a relationship between the endoplasmic reticulum and lipid particles. *Mol. Biol. Cell.* **9**, 375–386

72. Jacquier, N., Choudhary, V., Mari, M., Toulmay, A., Reggiori, F., and Schneiter, R. (2011) Lipid droplets are functionally connected to the endoplasmic reticulum in Saccharomyces cerevisiae. *J. Cell Sci.* **124**, 2424–2437

73. Wilfling, F., Wang, H., Haas, J. T., Krahmer, N., Gould, T. J., Uchida, A., Cheng, J. X., Graham, M., Christiano, R., Fröhlich, F., Liu, X., Buhman, K. K., Coleman, R. A., Bewersdorf, J., Farese, R. V., and Walther, T. C. (2013) Triacylglycerol synthesis enzymes mediate lipid droplet growth by relocalizing from the ER to lipid droplets. *Dev. Cell.* **24**, 384–399

74. Milla, P., Athenstaedt, K., Viola, F., Oliaro-Bosso, S., Kohlwein, S. D., Daum, G., and Balliano, G. (2002) Yeast oxidosqualene cyclase (Erg7p) is a major component of lipid particles. *J. Biol. Chem.* **277**, 2406–2412

75. M'barek, K. Ben, Ajjaji, D., Chorlay, A., Vanni, S., Forêt, L., and Thiam, A. R. (2017) ER membrane phospholipids and surface tension control cellular lipid droplet formation. *Dev. Cell.* **41**, 591–604

76.　Liu, Y., Fang, L., and Liu, W. (2021) High SQLE expression and gene amplification
correlates with poor prognosis in head and neck squamous cell carcinoma. *Cancer*
Manag. Res. **13**, 4709–4723

77.　Miettinen, T. A. (1982) Diurnal variation of cholesterol precursors squalene and methyl
sterols in human plasma lipoproteins. *J. Lipid Res.* **23**, 466–473

78.　Liu, G. C. K., Ahrens, E. H. J., Schreibman, P. H., and Crouse, J. R. (1976) Measurement
of squalene in human tissues and plasma: validation and application. *J. Lipid Res.* **17**,
38–45

79.　Goldstein, J. L., Basu, S. K., and Brown, M. S. (1983) Receptor-mediated endocytosis
of low-density lipoprotein in cultured cells. *Methods Enzymol.* **98**, 241–260

80.　Ran, F. A., Hsu, P. D., Wright, J., Agarwala, V., Scott, D. A., and Zhang, F. (2013)
Genome engineering using the CRISPR-Cas9 system. *Nat. Protoc.* **8**, 2281–2308

81.　Stevenson, J., Krycer, J. R., Phan, L., and Brown, A. J. (2013) A practical comparison
of ligation-independent cloning techniques. *PLoS One.* **8**, e83888

82.　Schmittgen, T. D., and Livak, K. J. (2008) Analyzing real-time PCR data by the
comparative CT method. *Nat. Protoc.* **3**, 1101–1108

83.　Labun, K., Montague, T. G., Krause, M., Torres Cleuren, Y. N., Tjeldnes, H., and Valen,
E. (2019) CHOPCHOP v3: expanding the CRISPR web toolbox beyond genome editing.
Nucleic Acids Res. **47**, W171–W174

84.　Kaakoush, N. O., Olzomer, E. M., Kosasih, M., Martin, A. R., Fargah, F., Lambie, N.,
Susic, D., Hoehn, K. L., Farrell, R., and Byrne, F. L. (2022) Differences in the active
endometrial microbiota across body weight and cancer in humans and mice. *Cancers*
(Basel). **14**, 2141

85.　Benjamini, Y., and Hochberg, Y. (1995) Controlling the false discovery rate: a practical
and powerful approach to multiple testing. *J. R. Stat. Soc. Ser. B*. **57**, 289–300

86.　Coates, H. W. H. W., Chua, N. K. N. K., and Brown, A. J. A. J. (2019) Consulting
prostate cancer cohort data uncovers transcriptional control: regulation of the MARCH6
gene. *Biochim. Biophys. Acta - Mol. Cell Biol. Lipids*. **1864**, 1656–1668

87.　Wong, J., Quinn, C. M., and Brown, A. J. (2007) Synthesis of the oxysterol, 24(S), 25-
epoxycholesterol, parallels cholesterol production and may protect against cellular
accumulation of newly-synthesized cholesterol. *Lipids Health Dis.* **6**, 10

1247

Supplementary tables

1248

Supplementary Table S1. siRNA and plasmids used for transfection

1249

siRNA	Description
SIC001	MISSION® universal negative control #1
SASI_Hs01_00105239	Targets human *MARCHF6* (NM_005885)
SASI_Hs01_00332063	Targets human *HIF1A* (NM_001530)
SASI_Hs01_00019152	Targets human *EPAS1* (*HIF2A*) (NM_001430)
SASI_Hs01_00016727	Targets human *HIF1AN* (*FIH-1*) (NM_017902)
Plasmid	**Description**
pCMV-(HA)$_3$-SM-V5	pcDNA3.1/V5-His TOPO vector containing the coding sequence of human squalene monooxygenase (SM; NM_003129.4) fused with three N-terminal HA epitope tags and C-terminal V5 and 6×His epitope tags, under the transcriptional control of a constitutive cytomegalovirus (CMV) promoter. Generated previously [9].
pCMV-(HA)$_3$-SM-V5 K82/90/100R	pCMV-(HA)$_3$-SM-V5 containing SM K82R, K90R and K100R substitutions, which blunt SM truncation [9]. Generated previously [9].
pCMV-(HA)$_3$-SM-V5 K290R	pCMV-(HA)$_3$-SM-V5 containing a K290R substitution, which disrupts a known ubiquitination site [32]. Generated during this study.
pCMV-(HA)$_3$-SM-V5 SM-N100 C/S/T>A	pCMV-(HA)3-SM-V5 containing SM T3A, T9A, T11A, S43A, C46A, S59A, S61A, S67A, S71A, S83A and S87A substitutions, which blunt cholesterol-induced degradation of SM [33]. Generated previously [9].
pCMV-(HA)$_3$-SM-V5 Y195F	pCMV-(HA)$_3$-SM-V5 containing an SM Y195F substitution that causes loss of catalytic activity [44]. Generated previously [9].
pCMV-SM(ΔN65)-V5	pCMV-(HA)$_3$-SM-V5 containing a deletion of the N-terminal (HA)$_3$ tag and first 65 residues of SM, which are lost upon truncation [9]. Generated previously [9].
pCMV-SM(ΔN65)-V5 Y195F	pCMV-SM(ΔN65)-V5 containing an SM Y195F substitution that causes loss of catalytic activity [44]. Generated during this study.
pCMV-GFP-ORP5	pEGFP-C1 vector containing the coding sequence of oxysterol binding protein-related protein-5 (ORP5) fused with an N-terminal GFP, under the transcriptional control of a constitutive CMV promoter. Generated previously [52].
pCMV-(HA)$_3$-SM-V5 Δ81–120	pCMV-(HA)$_3$-SM-V5 containing a deletion of SM residues 81–120, which are required for truncation [9]. Generated previously [9].
pCMV-SM-N100-GFP-V5	pcDNA3.1/V5-His TOPO vector containing the coding sequence of the first one hundred amino acids of human SM (SM-N100; NM_003129.4) fused with green fluorescent protein (GFP) and C-terminal V5 and 6×His epitope tags, under the transcriptional control of a constitutive CMV promoter. Generated previously [4].

1250

1251 **Supplementary Table S2. Primers used for qRT-PCR and DNA cloning**

1252 Non-annealing nucleotides for restriction enzyme cloning and DNA substitution are indicated
1253 in lowercase.

qRT-PCR primer pair		Primer sequence (5′–3′)	Source
VEGF	Forward Reverse	CCTGGTGGACATCTTCCAGG CTGTAGGAAGCTCATCTCTCC	[86]
CA9	Forward Reverse	GTGCCTATGAGCAGTTGCTGTC AAGTAGCGGCTGAAGTCAGAGG	Origene
RPL11	Forward Reverse	AGAGTGGAGACAGACTGACGCG CGGATGCCAAAGGATCTGACAG	Origene
GAPDH	Forward Reverse	GTCTCCTCTGACTTCAACAGCG ACCACCCTGTTGCTGTAGCCAA	Origene
ACTB	Forward Reverse	CACCATTGGCAATGAGCGGTTC AGGTCTTTGCGGATGTCCACGT	Origene
HIF1A	Forward Reverse	CAGTAACCAACCTCAGTGTGG CAGATGATCAGAGTCCAAAGC	Dr Laura Sharpe (Brown laboratory)
EPAS1 (*HIF2A*)	Forward Reverse	CTGTGTCTGAGAAGAGTAACTTCC TTGCCATAGGCTGAGGACTCCT	Origene
SQLE	Forward Reverse	GCTTCCTTCCTCCTTCATCAGTG GCAACAGTCATTCCTCCACCA	[9]
HMGCR	Forward Reverse	TTGGTGATGGGAGCTTGCTGTG AGTCACAAGCACGTGGAAGACG	[87]
HIF1AN (*FIH-1*)	Forward Reverse	CCTCTAACCTGCTGCTCATTGG GTAGAGGCACTCGAACTGATCC	Origene
DNA cloning primer pair		**Primer sequence (5′–3′)**	**Method**
SQLE gRNA 1 (promoter)	Forward Reverse	caccgAATGGAAACGTTCCGACCCG aaacCGGGTCGGAACGTTTCCATTc	*BbsI* cloning into PX598 vector (as in [80])
SQLE gRNA 2 (exon 1)	Forward Reverse	aaacAGTTCGCCCTCTTCTCGGATc caccgATCCGAGAAGAGGGCGAACT	
SM K290R	Forward Reverse	GATGGGCTTTTCTCCAgGTTCAGGAAAAGCCTG GCGATGCAATTTCCTCATTT	OEC (overlap extension cloning) [81]
SM Y195F	Forward Reverse	CCAGGTTGTAAATGGTTtCATGATTCATGATCAGG GCGATGCAATTTCCTCATTT	

1254