Allosteric Inhibition of the T Cell Receptor by a Designed Membrane Ligand

Yujie Ye [a], Shumpei Morita [b], Jay T. Groves [b] and Francisco N. Barrera [a, *]

[a], Department of Biochemistry & Cellular and Molecular Biology, University of Tennessee, Knoxville, TN 37996, U. S. A.

[b], Department of Chemistry, University of California-Berkeley, Berkeley, CA 94720, U. S. A.

* corresponding author: fbarrera@utk.edu

Abstract

The T cell receptor (TCR) is a complex molecular machine that directs the activation of T cells, allowing the immune system to fight pathogens and cancer cells. Despite decades of investigation, the molecular mechanism of TCR activation is still controversial. One of the leading activation hypotheses is the allosteric model. This model posits that binding of pMHC at the extracellular domain triggers a dynamic change in the transmembrane (TM) domain of the TCR subunits, which leads to signaling at the cytoplasmic side. We sought to test this hypothesis by creating a TM ligand for TCR. Previously we described a method to create a soluble peptide capable of inserting into membranes and bind to the TM domain of the receptor tyrosine kinase EphA2 (Alves *et al., eLife* 2018). Here we show that the approach is generalizable to complex membrane receptors, by designing a membrane ligand for TCR. We observed that the designed peptide caused a reduction of Lck phosphorylation of TCR at the CD3ζ subunit. As a result, in the presence of this Peptide Inhibitor of TCR (PITCR), the proximal signaling cascade downstream of TCR activation was significantly dampened in T cells. Co-localization and co-immunoprecipitation results in DIBMA native nanodiscs confirmed that PITCR was able to bind to the TCR. We propose that PITCR binds into a crevice present between the TM helices of the CD3ζ and CD3ε(δ) subunits. Our results additionally indicate that PITCR disrupts the allosteric changes in the compactness of the TM bundle that occur upon TCR activation, lending support to the allosteric TCR activation model. The TCR inhibition achieved by PITCR might be useful to treat inflammatory and autoimmune diseases and to prevent organ transplant rejection, as in these conditions aberrant activation of TCR contributes to disease.

Introduction

T cells are central players in the adaptive immune response. Different types of T cells recognize the presence of pathogenic organisms and cancer cells and orchestrate diverse immune activities intended to kill the damaging cells (Courtney et al., 2017; Ganti et al., 2020). The T cell receptor (TCR) is a protein complex present at the membrane of T cells that allows to detect the presence of foreign molecules. The TCR engages with antigen-presenting cells (APC), where peptide fragments are displayed at the major histocompatibility complex (pMHC) (Chakraborty & Weiss, 2014). Recognition of pMHC by the TCR triggers an intricate signaling cascade that activates the T cell response (Courtney et al., 2018; Kuhns & Davis, 2008). In αβ T cells, pMHC

binding occurs at the TCRαβ subunits. The TCR complex additionally contains four types of

CD3 subunits: ε, γ, δ and ζ, forming the TCR-CD3 complex (referred herein as TCR). The

dominant stoichiometry of TCR is composed of TCRαβ−CD3εγ−CD3εδ−CD3ζζ (Call et al., 2002;

Dong et al., 2019; Mariuzza et al., 2020). The CD3 subunits relay the information of the pMHC

binding event across the membrane, initiating TCR proximal signaling. TCR downstream

signaling cascade starts by phosphorylation of ITAMs (immunoreceptor tyrosine-based

activation motifs) present in all CD3 subunits, particularly at ζ, which leads to a transient

increase in calcium levels in the cytoplasm (Au-Yeung et al., 2018; Lo et al., 2018).

Despite decades of effort, there is not yet a clear understanding of how TCR is activated after

recognition of pMHC (Mariuzza et al., 2020). A number of different activation modes have been

proposed, namely the aggregation/clustering model, the segregation model, the

mechanosensing model, and the allosteric model. In this latter functional hypothesis, the signal

resulting from binding of pMHC to the extracellular region of the TCRαβ is dynamically

transmitted across the transmembrane (TM) region into the ITAMs. While there is growing

evidence supporting the allosteric model, as reviewed in (Schamel et al., 2019) , the molecular

mechanism of the allosteric triggering of TCR-CD3 is not clear. Recent reports provide a

plausible mechanistic framework on how allosteric changes affect the TM helical bundle: TCR

activation involves a quaternary relaxation of the TM helical bundle, whereby in the active state

there is a loosening in the interaction between the TM helices of the TCRβ and CD3ζ (Lanz et

al., 2021; Prakaash et al., 2021).

Here, we have used rational design to develop a peptide (PITCR) to target the TM region of

TCR. PITCR comprises the TM domain of the ζ subunit (Call et al., 2006) modified by the

addition of acidic residues to convert it to a conditional TM sequence. The PITCR peptide is

used to test the allosteric relaxation model, as its binding to the TM region of TCR can be

reasonably expected to alter the conformation and/or dynamics/packing of the helical bundle.

We observed that PITCR robustly inhibited the activation of the TCR. The results obtained in

this work support the allosteric relaxation activation model and provide new mechanistic insights

into TCR activation.

Results

PITCR decreases phosphorylation of the ζ chain upon TCR activation.

We recently reported an approach to transform the isolated TM domains of human receptors

into peptides that function as conditional TM sequences: they are highly soluble in water, while

they have the ability to insert into the membrane in the TM orientation that allows the peptide to

interact laterally with their natural binding partners (Alves et al., 2018). We applied this protocol

to the CD3ζ TM, to generate the PITCR peptide. Biophysical experiments in synthetic lipid

vesicles showed that the design for PITCR was successful, as it was water-soluble and was

able to insert into membranes (Figure 1-Figure supplementary 1).

The TCR at the surface of human Jurkat T cells is activated upon binding of the monoclonal

antibody (mAb) OKT3, which has been widely applied to study T cell signaling (Lo et al., 2018;

Lo et al., 2019). TCR activation is initiated by phosphorylation of tyrosine residues at the ITAMs

of the ζ chain by Lck (lymphocyte-specific protein tyrosine kinase) (Figure 1A) (Courtney et al.,

2017). To investigate whether PITCR affects TCR activation, we treated Jurkat cells with PITCR

before stimulation with OKT3. The immunoblot results revealed that PITCR reduced

phosphorylation of the ζ chain at residues Y142 and Y83 after TCR activation **(Figure 1B-C and**

Figure 1. PITCR reduces phosphorylation of the ζ chain in response to OKT3. (A) Cartoon that illustrates TCR proximal downstream signaling. The plasma membrane is shown as a horizontal bar, and phosphorylation sites are shown as yellow dots. ECD: extracellular domain; TMD: transmembrane domain; ICD: intracellular domain of TCR. **(B)** Jurkat cells were treated with PITCR, followed by stimulation with OKT3. Lysates were analyzed by immunoblot to detect TCR phosphorylation of ζ (pY142 and pY83). Total ζ levels were assessed and no change was observed. Data are representative at least five independent experiments. **(C)** Quantification of phosphorylation at both tyrosine residues in the presence of OKT3, normalized to data in the absence of PITCR (based on data from Figure 1-Figure supplementary 2). Error bars are the SD. *p* values were calculated using a two-tailed Mann-Whitney test.

Figure 1 - Figure supplementary 2). These results suggest that PITCR binding reduces

activation of the TCR.

Phosphorylation of TCR proximal signaling molecules is downregulated by PITCR.

Since PITCR inhibited TCR phosphorylation after activation, we sought to explore the effect of

PITCR on TCR downstream signaling. TCR activation induces the recruitment of Zap70 (ζ chain

– associated protein kinase 70) to the phosphorylated TCR, where it is itself phosphorylated by

Lck **(Figure 1A)**. The activated Zap70 then phosphorylates LAT (linker for activation of T cells)

and SLP76 (SH2 domain containing leukocyte protein of 76 kDa), and as a result PLCγ1

(phospholipase C-γ1) is recruited and phosphorylated (Courtney et al., 2018; Lo et al., 2018; Lo

Figure 2. PITCR reduces phosphorylation of TCR proximal signaling proteins after activation. (A) Immunoblot analysis of lysates to detect phosphorylation of Zap70 (pY319 and pY493), LAT (pY132 and pY191), SLP76 (pY128), and PLCγ1 (pY783). Total protein levels of Zap70, LAT and the housekeeping protein β-actin were assessed, revealing no changes. Data are representative of at least five independent experiments. **(B)** Quantification of phosphorylation in the presence of OKT3, normalized to data in the absence of PITCR (based on data from Figure 2-Figure supplementary 1). Error bars are the SD. *p* values were calculated using a two-tailed Mann-Whitney test.

& Weiss, 2021). In agreement with the hypothesis that PITCR inhibits TCR activation, in the

presence of peptide we observed a statistically significant decrease in the phosphorylation of

Zap70, LAT, SLP76 and PLCγ1 (**Figure 2 and Figure 2 - Figure supplementary 1**). However,

PITCR did not affect phosphorylation of Lck (**Figure 2-Figure supplementary 2**). Since basal

phosphorylation of Zap70, LAT, SLP76 and PLCγ1 was observed in the absence of OKT3

(**Figure 2A**), we could determine that the effect of PITCR is specific to stimulation of the TCR.

Figure 3. PITCR reduces the TCR intracellular calcium response. Jurkat cells were stained with the fluorescent dye indo-1 AM, followed by treatment with PITCR **(A)**, pHLIP as a negative control **(B),** or the variant PITCRG41P **(C)** and stimulated with OKT3. Ionomycin was applied as a positive calcium control. The indo-1 ratio was calculated from fluorescence at 405 nm (calcium-bound) divided by 475 nm (calcium-free). Data are representative of three independent experiments. Each independent experiment includes at least four technical replicates. Error bars are the SEM. **(D)** Quantification of the maximum Indo-1 increase after OKT3 activation, normalized to no peptide treatment. Error bars are the SD. *p* values were calculated from a Kruskal-Wallis test with Dunn's multiple comparisons test.

Our data therefore indicate that PITCR causes a robust inhibition of the proximal signaling

cascade triggered by TCR activation.

PITCR reduces intracellular calcium response.

After TCR activation, the active PLCγ1 hydrolyzes phosphatidyl inositol 4,5-bisphosphate (PIP_2)

to generate inositol trisphosphate (IP_3) and diacylglycerol (DAG). Free IP_3 diffuses across the

cytoplasm and binds to the IP3 receptor at the endoplasmic reticulum (ER), causing the release

of calcium from ER storage (Figure 1A) (Courtney et al., 2018; Lewis, 2001; Trebak & Kinet,

2019). Based in our previous results, we reasoned that PITCR should inhibit the cytoplasmic

calcium influx that follows TCR activation. We tested this idea with a kinetic analysis of

intracellular calcium release using the calcium indicator Indo-1 (Lo et al., 2018).

Consistent with our expectation, we observed that the calcium response after OKT3 stimulation

was attenuated in the PITCR treatment group compared with control conditions (**Figure 3 A and**

D, Figure 3 - Figure supplementary 1). We used as a negative control pHLIP (Scott et al.,

2019; Scott et al., 2017), a different conditional TM peptide that is not expected to interact with

membrane proteins (Alves et al., 2018). The dynamic calcium curve of pHLIP-treated group was

within the error of the control curve (**Figure 3 B and D**). To further test specificity of PITCR, we

performed a mutation (replacing Gly41 for a Pro) in PITCR that is expected to form a helical kink

and disrupt the TM helix. Control biophysical experiments indicated that the G41P mutation did

not prevent the peptide to act as a conditional TM (**Figure 1 - Figure supplementary 1**). We

observed that PITCRG41P was unable to inhibit the calcium influx in response to OKT3

treatment (**Figure 3 C and D**). These data indicate that PITCR specifically impaired the calcium

response that occurs in Jurkat cells after TCR activation, in agreement with the observed

decrease in phosphorylation of PLCγ1 (**Figure 2A**).

Inhibition of TCR engagement with antigen presenting cells.

While the OKT3 mAb efficiently stimulates TCR signaling, we sought to test the effect of TCR in

a more physiologically relevant TCR activation, consisting of T cell stimulation by binding to

pMHC in antigen presenting cells (APCs) (Lo et al., 2018). Similar to human primary cytotoxic

CD8[+] T cells, OT1[+]-TCR CD8[+] Jurkat T cells (J.OT1.CD8) can recognize the ovalbumin (OVA)

peptide presented by H-2K[b]-MHC I on T2 APC cells (T2K[b]) (**Figure 4 A and B**). TCR

engagement results in increased levels of CD69, a T cell activation marker (Lo et al., 2018; Lo

et al., 2019; Wolpert et al., 1997).We treated J.OT1.CD8 cells with PITCR followed by

incubation with T2K[b] cells pre-treated with a range of OVA concentrations and measured the

CD69 expression using flow cytometry (**Figure 4 C and D**). As expected, in the presence of

high OVA concentrations we observed CD69 upregulation (**Figure 4 C**). Our data showed that

PITCR caused a significant reduction of CD69 levels (**Figure 4 C and D**). To further examine

the specificity of PITCR, we again used pHLIP as a negative control, and we observed that

pHLIP did not elicit significant changes. Our data therefore indicate that PITCR specifically

impaired CD69 upregulation in J.OT1.CD8 cells in response to OVA stimulation. These results

show that PITCR also achieves robust inhibition when TCR is activated by binding to pMHC in

APC.

Figure 4. PITCR reduces CD69 expression after T cell activation by APC. (A) Cartoon showing T cell interaction with APC. **(B)** A live cell microscopy image that depicts engineered Jurkat-OT1$^+$ TCR-CD8$^+$ T cells interacting with T2K^b APC pre-incubated with the peptide antigen OVA. **(C)** Jurkat-OT1$^+$ TCR-CD8$^\pm$ T cells were treated with PITCR or pHLIP (as a negative control), and then incubated with T2K^b cells at different concentrations of OVA, followed by CD69 flow cytometry analysis. The upregulation of CD69 is representative of four independent experiments. Each independent experiment includes two technical replicates. Error bars are the SD. **(D)** Quantification of CD69-positive cells at [OVA] = 1 μM for PITCR (red) and negative control pHLIP (green). Each dot pair represents one independent experiment. p values were calculated from two-tailed paired t-test.

PITCR interacts with the TCR

Our results clearly show that PITCR inhibits TCR activation. We sought next to determine if this was a specific effect that resulted from a direct interaction between the peptide and TCR. First, we performed co-localization experiments in Jurkat cells. To this end, we fluorescently labeled PITCR with dylight680 (PITCR680). We employed super-resolution confocal microscopy with lightning deconvolution to investigate PITCR680 co-localization with TCR, as detected with an anti-CD3 ε antibody. We observed that PITCR680 localized to some areas of the Jurkat cells, corresponding to the plasma membrane and intracellular organelles, probably endosomes (**Figure 5 A**). In these two regions we observed co-localization between PITCR680 (magenta) and TCR (CD3ε,green) (**Figure 5 A**). To better assess co-localization, we plotted graphic profile curves on regions of interest (ROI), which revealed clear overlap in some areas. To quantify the degree of co-localization, we calculated the Mander's M1 coefficient, which showed a value of ~0.8 (**Figure 5 D**). This result reveals that around 80% of PITCR680 signal overlaps with the TCR We also observed robust co-localization using the Pearson correlation coefficient ($r = $ ~0.4) (**Figure 5 – Figure supplementary 2**) (Costes et al., 2004). To further explore whether TCR activation influences co-localization, we stimulated PITCR680-treated Jurkat cells with OKT3.

Figure 5. PITCR colocalizes with TCR. (A) PITCR680 and CD3ε co-localization was studied in the presence and absence of activation with OKT3. Scale bars = 10 μm. Representative areas with co-localization at the plasma membrane (*top*) and cytoplasm (*bottom*) were zoomed-in, where scale bars are 0.5 μm and 1 μm, respectively. Confocal images are representative of three independent experiments. **(B)** and **(C)** show graphic profile curves (dotted yellow lines) plotted across the zoom-in ROI in + OKT3 and – OKT3, respectively. Magenta lines denote PITCR, and green lines CD3ε. Overlap indicates colocalization. **(D)** Quantification of co-localization by Mander's coefficient (M1), corresponding to the fraction of PITCR that overlaps with CD3ε. Error bars indicate SD. *p* value was calculated from two tailed unpaired *t*-test.

We observed similar results, suggesting that PITCR is able to bind to TCR before it is activated

by OKT3.

We investigated if co-localization of PITCR and TCR also occurred in primary T cells. We

performed surface-selective imaging by total internal reflection microscopy in primary murine

CD4[+] T cells expressing AND TCR. Immune synapse formation was imaged (Biswas & Groves,

2019; Mossman et al., 2005; Yu et al., 2012), as T cells were stimulated by contact with

supported bilayers functionalized with agonist pMHC (MCC peptide labeled with Atto647N) and

the adhesion molecule ICAM-1 (Lin et al., 2019) (McAffee et al., 2021). TCR-pMHC complexes

were selectively distinguished from free pMHC using an elongated image exposure time

strategy (Lin et al., 2019; O'Donoghue et al., 2013; Pielak et al., 2017). We observed the c-

SMAC (central supramolecular activation cluster) structure at the core of the immune synapse

(Figure 6), as previously reported (Bromley et al., 2001). PITCR conjugated to AZDye 555

(PITCR555) could be detected in intracellular structures (**SI Movie**), consistent with confocal

microscopy results. Additionally, a distinct population plasma-membrane-bound peptide could

be also observed (**Figure 6**). In these cases, PITCR exhibited c-SMAC localization together with

TCR-pMHC complexes (**SI Movie**). Although the image resolution was insufficient to definitively

confirm molecular binding between PITCR and TCR-pMHC complex, their co-localization is

Figure 6: Colocalization of PITCR and the TCR-pMHC complex in primary murine CD4[+] T cells. **(A)** Images of plasma-membrane-localized PITCR555 and TCR-pMHC complex in a representative T cell adhering to supported lipid bilayer functionalized with pMHC (19-23 molecules/μm^2, 9% labeled with Atto-647N) and ICAM-1 (~20 molecules/μm^2). TCR-pMHC complex was selectively visualized with a long exposure time (500 ms). PITCR exhibited localization at c-SMAC together with the TCR-pMHC complex. **(B)** Vehicle control showed no signal at the PITCR channel.

suggestive of interaction. In aggregate, the co-localization experiments support binding of

PITCR to TCR in Jurkat and primary T cells.

We sought to determine if the observed co-localization indeed resulted from binding between

PITCR and TCR. We developed an assay to maintain the TCR complex of Jurkat cells in a

native lipid environment, consisting of using the polymer diisobutylene maleic acid (DIBMA) to

form native nanodiscs. On these samples we performed a co-immunoprecipitation (Co-IP)

experiment using an anti-CD3 ε antibody (UCHT1). We observed bands of TCR β, CD3 ε and

CD3 ζ in the anti-CD3 ε Co-IP lysates (**Figure 7**), indicating that TCR had been successfully

immunoprecipitated after solubilization with DIBMA. When cells were incubated with PITCR680,

Figure 7. PITCR interacts with TCR. Jurkat cells were treated with PITCR-680. TCR nanodiscs were immunoprecipitated with the mAb UCHT1, or with IgG as a negative control. Fluorescent detection of PITCR-680 after Co-IP, or run in the gel directly as a positive control (*side panel*). Below are shown immunoblot analysis of Co-IP samples and whole lysates to probe CD3 ε, TCR β and CD3 ζ. β-actin was a loading control. Data are representative of at least three independent experiments.

we observed in the Co-IP samples a fluorescent band of molecular weight (~5 kDa) similar to

that of PITCR680 (4.8 kDa) (**Figure 7**). To further explore whether TCR activation would affect

binding of PITCR680 to the complex, we applied OKT3 as previously described. We observed

the ~5 kDa fluorescent band as well. These results indicate that PITCR680 interacts with the

TCR irrespective of activation by OKT3, in agreement with the co-localization results in Jurkat

cells (**Figure 5**).

PITCR weakens the interaction of the ζ subunit with the rest of the complex after TCR

activation.

We next sought to understand the mechanism by which PITCR partially inhibits TCR activation.

It has been recently proposed that TCR activation involves a change in robustness of the TM

helical bundle. This allosteric change can be detected by immunoprecipitation after

solubilization in DDM, as a loose complex is less resistant to this detergent (Lanz et al., 2021;

Prakaash et al., 2021). We optimized this assay for our experimental conditions and

investigated the interaction between the ζ chain and rest of the TCR complex. We first

examined the immunoprecipitated levels of CD3 ε, CD3 ζ and TCR β in the presence of DDM to

Figure 8. PITCR weakens the interaction of the ζ chain with the rest of the complex after TCR activation. (A) Immunoblot analysis of immunoprecipitated samples and whole lysate samples solubilized with DDM. Data are representative of at least three independent experiments. **(B)** Quantification of ζ / β2 and ζ / ε after OKT3 stimulation. **(C)** ζ / β2 and ζ / ε in PITCR-treated OKT3 samples, normalized to OKT3 stimulation. The β2 subunit was studied since it is the most abundant β subunit at the plasma membrane. Error bars are SD. *p* values were calculated from two-tailed Mann-Whitney test.

evaluate the specificity and efficacy of the use of the anti-CD3 ε antibody for our immunoprecipitatation assay. We observed the targeted bands in the immunoblot results for anti-CD3 ε-IP, and no bands in the negative control IgG-IP (**Figure 8A**). These results suggest that our approach results in the IP of the TCR complex. Once we validated our method, we tested the effect of OKT3 stimulation and PITCR treatment. We observed that OKT3 increased the levels of CD3ζ when compared to both TCRβ2 and CD3ε (ζ / β2 and ζ / ε) (**Figure 8**). These results suggested a more robust ζζ interaction with the rest of complex in response to OKT3 stimulation. Based on this result, we reasoned that PITCR could act by reversing the changes in quaternary robustness that occur in the membrane region upon TCR activation. In agreement with our hypothesis, we observed that the ζ / β2 ratio was decreased when cells were incubated with PITCR and OKT3. However, ζ / ε did not change (**Figure 8**). Taken together, these results suggested that PITCR disrupts the allosteric changes in TM quaternary robustness that occur upon TCR activation.

Discussion

For this study we developed a novel conditional transmembrane peptide to target the T cell receptor. The design of PITCR involved strategically introducing glutamic acid residues into the transmembrane sequence of the human CD3 ζ chain, as previously described for TYPE7, a rationally designed TM ligand for the EphA2 receptor (Alves et al., 2018). We additionally introduced small and polar residues (**Figure 1-Figure supplementary 1 A**). PITCR selectively inserted into synthetic lipid vesicles at acidic pH (**Figure 1-Figure supplementary 1**). However, we observed that TCR activation in Jurkat cells was severely disrupted by acidic pH (data not shown). We therefore performed experiments at physiological pH, where we observed that PITCR efficiently targeted cells (**Figure 5 and Figure 6**). This observation is not unexpected, since the acidity-responsive TYPE7 also targeted cellular membranes at neutral pH, because the presence in the membrane of its target receptor shifts the pH-responsiveness to cause membrane insertion at pH 7.4. We suggest that a similar situation might occur for TCR. However, we speculate that PITCR will more effectively inhibit T cells that reside and survive in acidic environments. The solubility of PITCR is a useful property to facilitate delivery to tissues. Furthermore, the pH-responsiveness of PITCR could potentially be used for targeted therapies in pathologies that are characterized by acidic extracellular environments, including inflammatory (Andreev et al., 2007) and autoimmune diseases (Marunaka, 2015), and solid tumors (Cheng et al., 2015). PITCR could also be used to overcome a critical limitation of

allogeneic CAR T therapy, since TCR inhibition is required to prevent graft-versus-host disease
resulting from side effects of this therapy. The use of PITCR would additionally overcome the
risk resulting from genetic manipulation transfection with of allogeneic T cells before injection
into patients (Michaux et al., 2022).

PITCR caused robust inhibition throughout the signaling cascade that is triggered when TCR is
activated, from phosphorylation of the ζ chain to calcium influx. Upon TCR ligation, immunoblot
analysis showed that PITCR inhibited phosphorylation at multiple sites in Zap70, LAT, SLP76
and PLCγ1 (**Figure 2**), but not of Lck) (**Figure 2-Figure supplementary 2**). The observed
specific inhibition of TCR-downstream signaling suggests that PITCR is unlikely to
inhibit/activate a broad range of kinases or phosphatases. Rather, PITCR is likely to specifically
inhibit TCR triggering, while maintaining Lck association with coreceptors and its
phosphorylation (Ashouri et al., 2022; Guy et al., 2022).

What is the molecular mechanism of PITCR inhibition of TCR? A recent molecular dynamics
simulation study of TCR (Prakaash et al., 2021) offers clues on a possible binding site for
PITCR. The model reveals a hydrophobic gap between the TM domain of the ζ and ε(δ)
subunits that can accommodate PITCR (Figure 9). Interestingly, the cryo-EM structure of TCR
reveals a binding interface between the TM of ζ and ε(δ) (Dong et al., 2019). PITCR maintains
all the residues that form the helical face of ζ that interacts with ε(δ). Therefore, we propose that
PITCR binds into the membrane gap between ζ and ε(δ), where it establishes native

Figure 9. Cartoon model of PITCR interaction with TCR. Schematic view looking from the
extracellular side of the TM bundle of TCR, with PITCR is shown in magenta. TM helices are shown
as cylinders or circles. The disulfide bond between the ζ subunits is not shown for clarity.

interactions with the ε(δ) subunit. We found that when we replaced glycine at position 41 with

proline, the peptide no longer inhibited calcium influx (**Figure 3**). We speculate that proline

induced a local helical kink the TM helix peptide that prevented binding to the TM gap, or a

productive interaction between PITCR and the ε(δ) subunit of TCR.

The results of the immunoprecipitation experiments in the presence of DDM (Figure 8) showed

that upon TCR activation with OKT3, ζζ strengthened its association with the β subunit. We

observed that PITCR caused the opposite effect. This result shows that PITCR acts by

reversing the allosteric changes in TM compactness induced by OKT3 activation. However,

there was an interesting difference in the interaction with the ε subunit, which increased with

OKT3 activation but was not altered by PITCR. This observation suggests that not all the TM

interactions are equally important with regards to contributing to TCR activation. Our results

additionally suggest that the interface between the ζ and β subunits is a target for

pharmacological inhibition of the TCR.

We based the DDM immunoprecipitation assay on a previous report (Lanz et al., 2021), but our

protocol contains significant differences. We performed the immunoprecipitation for the native

receptor, while the previous protocol involved IP of an HA-tagged TCR, which was also

activated by different means to ours. Probably as a result of these differences, our DDM

immunoprecipitation result showed an opposite effect into how TCR activation affected the

association of TCRαβ with ζζ (Brazin et al., 2018; Lanz et al., 2021). Nevertheless, the two

experimental lines of evidence still agree in showing changes in the quaternary stability of the

TCR upon activation, and support that this effect could be an important element of the allosteric

activation of the TCR. Overall, our data indicate that binding of PITCR to the transmembrane

region of TCR results in opposite allosteric changes to those that occur upon TCR activation,

and we propose this effect reduces signal transduction into the intracellular region.

Even though the activation of TCR is an intricate process (Chai, 2020; Dong et al., 2019;

Mariuzza et al., 2020; Reinherz, 2019), significant progress has been made to understand the

conformational changes it entails. For example, in response to TCR engagement, the

juxtamembrane domains of the ζζ homodimer have been experimentally reported to switch from

a divaricated to a parallel conformation (Lee et al., 2015). TCR activation additionally involves a

conformational change of the ITAMs of CD3ε that releases the interaction of basic residues in

the intracellular domain of CD3 ζ (Zhang et al., 2011) and ε (Xu et al., 2008) giving access to

Lck for phosphorylation. We suggest a plausible mechanism where PITCR insertion into the gap

between ζζ homodimer and the rest of TCR-CD3 complex may additionally hinder the activating

conformational change in ζζ. This hypothesis might further explain the observed decrease of ζ /

β2 in the PITCR-treated Jurkat cells upon TCR activation.

The human adaptive immune system contains numerous types of T cells. T cells present a

broad repertoire of TCR (Davis & Bjorkman, 1988). For example, in human peripheral blood

samples, 10^4 varieties of TCRβs can be found (Kidman et al., 2020), while more than 10^{15}

combinations of TCRαβ are could theoretically be formed (Davis & Bjorkman, 1988) (Wong et

al., 2007) (Freeman et al., 2009). The TCR repertoire plays a critical role in the adaptive

immune response, but it also brings challenges to achieve immunosuppression to treat

inflammatory and autoimmune diseases. Our data show that PITCR interacts with different

types of TCRs: it inhibited TCR signaling in Jurkat-WT (**Figure 1, Figure 2 and Figure 3**), and

colocalized with the cSMAC structure formed by AND-TCR in primary CD4+ T cells (**Figure 6**).

Furthermore, PITCR reduced CD69 upregulation in OT1-TCR Jurkat cells (**Figure 4**). These

results suggest that PITCR may possess the ability to interact with a broad range of TCR types.

We hypothesized that this would be possible because the PITCR design is based on targeting

the TM region of the TCR, a sequence that is largely conserved.

Taken together, our data show that rational design of a membrane ligand inhibits TCR

activation. PITCR has potential clinical value to treat autoimmune and inflammatory diseases or

to avoid transplant rejection. The strategy used to design PITCR can be applied to develop

targeted ligands for other receptors causative of disease.

Materials and Methods

Cell lines

Human male leukemic Jurkat T cells (Clone E6-1, TIB-152) were obtained from the American

Type Culture Collection (Manassas, United States). Jurkat.OT1-TCRα-GFP-TCRβ.hCD8+

(J.OT1.hCD8+) cells and T2-K^b cells were kindly provided by Arthur Weiss (UCSF). Cell lines

were maintained in RPMI 1640 (Gibco 11875119) supplemented with 10 % fetal bovine serum

(Gibco 10437028), 1 % L-Glutamine (Gibco 25030081) and 1 % penicillin & streptomycin (Gibco

15140122) in a 37 °C & 5 % CO_2 humid tissue culture incubator (Panasonic Healthcare, Wood

Dale, United States). The identity of Jurkat cells was authenticated using ATCC services.

Mycoplasma contamination was ruled out by PCR (Abcam 289834).

Peptide synthesis

327   Peptides were synthesized by Thermo Fisher Scientific (Waltham, United States) and were

328   confirmed by matrix-assisted laser desorption ionization-time-of-flight (MALDI-TOF) mass

329   spectrometry and reverse phase high performance liquid chromatography (HPLC). Purities of

330   the peptides are over 95 %. PITCR sequence: DPKLSYLLDGILFGYGVELTALFLEVGFSESAD.

Intracellular calcium assay

Jurkat wild-type (WT) cells were washed twice with PBS and then stained with the calcium

sensor dye Indo-1 AM (Invitrogen I1223), at 37 °C and 5 % CO_2 for 30 mins as described (Lo et

al., 2018; Lo et al., 2019). The final concentration of Indo-1 AM was 1 µM. Stained Jurkat cells

were washed twice with PBS and then were transferred to a 96 well flat bottom black plate.

PITCR was added and incubated at 37 °C and 5 % CO_2 for 20 mins. Next, the plate was

transferred to a prewarmed (37 °C) and 5 % CO_2 Cytation V plate reader (BioTeK, Winooski,

United States) and incubated for another 10 mins. The final concentration of PITCR in the each

well was 10 µM. The baseline for each well was recorded for the first 100 seconds, followed by

auto-injection of anti-CD3 (OKT3 clone, Tonbo-70-0037). The final concentration of anti-CD3

was 1 µM. Ionomycin (Invitrogen I24222) was used as a positive control to prove that Jurkat

cells respond to calcium influx. The fluorescent signal collected from Jurkat-WT cells without

staining was subtracted from the signal collected from Indo-1 AM stained cells since Jurkat-WT

cells have auto-fluorescent signals.

SDS-PAGE Immunoblot analysis of proximal signal molecules of TCR-CD3, ζ-Y83 and ζ-Y142

Jurkat-WT cells were washed twice with PBS and treated with PITCR at 37 °C and 5 % CO_2 for

30 mins, followed by stimulation with anti-CD3 antibody (OKT3 clone) for 5 mins. The final cell

density was 5 x 10^6 cells/ml and the final concentration of PITCR was 10 µM. The final

concentration of anti-CD3 was 1 µM. Cells were lysed in 1 % NP-40 lysis buffer (50 mM Tris-Cl

pH 7.4, 150 mM sodium chloride, 2 mM PMSF, 5 mM EDTA, 0.25 % sodium deoxycholate with

proteinase inhibitors (Thermo Scientific A32955) and phosphatase inhibitors (Sigma-Aldrich

P0044) for 30 minutes on ice, followed by centrifugation of 16.2 x 10^3 g, 30 mins, 4 °C.

Supernatants were collected and detergent-compatible protein assay (Bio-Rad 5000112) was

performed to quantify the protein concentration of each sample. Equal amounts of protein

samples were run in 10 %, 12 % or 15 % SDS-PAGE gels and transferred to 0.45 µm nitro

cellulose membranes. Membranes were blocked with 3 % bovine serum albumin (BSA)

dissolved in TBS, followed by overnight incubation with primary antibodies at 4 °C. IRDye

800CW or IRDye 680LT secondary antibodies were used to incubate the blots at the 2nd day,

followed by detection with an Odyssey Infrared Scanner (Li-Cor Biosciences, Lincoln, United

States). All primary antibodies were diluted in 5 % BSA dissolved in 0.1 % TBST except specific

mentions and all secondary antibodies were diluted in 5 % non-fat milk dissolved in 0.1 % TBST

unless mentioned otherwise.

Co-localization assay and analysis

All steps were performed at room temperature (RT), unless noted otherwise. Jurkat-WT cells

were washed once with PBS, resuspended in RMPI1640 phenol-red free media, and treated

with dylight680 labeled PITCR (PITCR680) at 37 °C and 5 % CO_2 for 15 mins, followed by

stimulation in presence or absence of anti-CD3 (OKT3 clone) for 5 mins. The final concentration

of anti-CD3 was 1 µM. The final cell density was 5 x 10^6 cells/ml and the final concentration of

PITCR680 was 10 µM. PITCR680 treated cells were washed twice with cold PBS and

resuspended in cold RPMI1640 phenol-red free media. 100 µL of cells were transferred to each

chamber of µ-Slide 8 Well ibiTreat (ibidi80826) and rested for 10 mins. Next, RPMI1640 phenol-

red free media was removed gently. 150 µL cold fixing buffer (1.998 % - formaldehyde and 0.2

% - glutaraldehyde in filtered PBS) was added and incubated for 10 mins. Fixed PITCR680

treated Jurkat cells were washed twice with cold DPBS/Modified (HycloneSH30264.01). 150 µL

permeabilization buffer (0.1 % Triton X100 in PBS) was incubated with the sample for another

15 mins. Each chamber was washed with blocking buffer (5 % goat serum in 0.01 % PBST)

twice. The sample was blocked 1 hr, followed by washing once with antibody dilution buffer (1 %

BSA in 0.01 % PBST). The samples were incubated with 1:200 diluted anti-CD3ε (UCHT1

clone, sc-1179) primary antibody in the wet box at4 °C, overnight. On the 2nd day, each

chamber was washed twice with cold DPBS/Modified (HycloneSH30264.01). 1:200 diluted

secondary antibody (Invitrogen A32723) was incubated with samples in a foil covered wet box

for 1 hr, followed by washing twice with cold DPBS/Modified (HycloneSH30264.01). 1:1000

diluted DAPI (Thermo Scientific 62248) was stained with samples for 2 mins, followed by

washing once with cold DPBS/Modified. The samples were mounted with Vectashield (H-1000),

followed by sealing the chambers with parafilm until imaging.

Samples were imaged using a Leica SP8 White Light Laser Confocal Microscope (Leica,

Wetzlar, Germany) equipped with a 63X oil immersion objective, zoom 5 through LAS X

software. Z-stack scanning was applied, followed by a lightning deconvolution analysis. Each

image was chosen from one time point at each Z-stack section. Graphic profile curves of the

Region of Interest in Zoom-in images were analyzed using Image J RGB Profile Plot Plugin. The

Mander's M1 and Pearson's *r* coefficients were calculated using Image J Just Another

Colocalization Plugin (JACoP). **CD69 activation flow cytometry assay**

2×10^6 cells/ml T2-K^b cells were washed twice with PBS and treated with a series of diluted

ovalbumin (OVA) derived peptide (SIINFEKL) at 37 °C and 5 % CO_2 for 1 hr. 5 x 10^6 cells/ml

J.OT1.h$CD8^+$ cells were washed twice with PBS and treated with PITCR at 37 °C and 5 % CO_2

for 30 mins. PITCR-treated J.OT1.h$CD8^+$ cells were added to OVA-treated T2-K^b cells and the

ratio of these two cells was 1:1. Final concentration of PITCR was 10 µM. The incubation time

was 3 hrs. Next, anti-CD69 - Allophycocyanin (Biolegend 310910) and Isotype -

Allophycocyanin (Biolegend 400122) were applied to stain the cells, respectively followed by

LSRII Flow Cytometer (BD Bioscience, Franklin Lakes, United States) analysis. Data was

quantified using FlowJo_v10.8.0 software.

Co-Immunoprecipitation of TCR-CD3 complex and immunoblot analysis

Jurkat-WT T cells were washed twice with PBS and treated with Dylight680 labeled PITCR at 37

°C and 5 % CO_2 for 30 mins, followed by stimulation with 1 µM anti-CD3 antibody (OKT3 clone).

Cells were lysed in 2.5 % Diisobutylene Maleic Acid (DIBMA, AnatraceBMA101) lysis buffer (20

mM Tris-Cl pH 8.0, 137 mM NaCl, 2 mM EDTA, 1 mM PMSF, 5 mM iodoacetamide, 1 mM NaF,

proteinase inhibitors (Promega G6521) and phosphatase inhibitors (Sigma-Aldrich P0044 and

P5726) at warm room (37 °C) for 2 hrs followed by rotating at least 12 hrs in the cold room (4

°C). Lysates were ultracentrifuged 10,000 g, 4 °C, 45 mins to get rid of debris, and the

supernatants were collected and ultracentrifuged again 10,000 g, 4 °C, 1hr. The protein

concentrations were quantified using a detergent compatible protein assay. 40 µl whole lysates

were saved to detect the TCR-CD3 complex. 1 % BSA blocked protein G agarose (Thermo

Scientific 20398) and anti-CD3ε (UCHT1 clone, sc-1179) were added to the rest of lysates.

Samples were gently rotated at cold room (4 °C) for 16 hrs. Samples were centrifuged 5,000 g,

4 °C, 3mins, followed by washing twice with cold wash buffer (20 mM Tris-Cl pH 8.0, 137 mM

NaCl, 2 mM EDTA, 1 mM PMSF, 5 mM iodoacetamide, 1 mM NaF) and rinsing once with cold

wash buffer. SDS sample buffer was applied to elute the captured protein followed by incubation

at 95 °C for 5 mins.

Equal amount of whole lysate samples and captured protein samples were loaded in 15 % SDS-

PAGE glycine gels and transferred to 0.45 µm or 0.2 µm nitro cellulose membranes. 3 % BSA

was used to block the membranes, followed by incubation with primary antibodies: anti-TCRβ,

anti-CD3ε and anti-CD3ζ (6B10.2 clone) overnight at 4 °C. In the whole lysates' group, the

housekeeping protein β-actin was also probed. PITCR was directly detected using channel 700

nm of an Odyssey Infrared Scanner (Li-Cor Biosciences, Lincoln, United States). The 2[nd] day,

same approaches were performed as described in SDS-PAGE Immunoblot analysis of proximal

signal molecules of TCR-CD3, ζ-Y83 and ζ-Y142.

Liposome preparation

1-palmitoyl-2-oleolyl-glycero-3-phosphocholine (POPC) and 1-palmitoyl-2-oleoyl-sn-glycero-3-

phospho-L-serine (POPS) were purchased from Avanti Polar Lipids, Alabaster, AL. POPC and

POPS were dissolved in cold chloroform and stocks prepared at 32.89 mM. POPS and POPC

were mixed in a round-bottom test tube, dried together under a stream of argon gas, and placed

in a vacuum overnight. The dried lipid film was rehydrated in 10 mM sodium phosphate buffer

(pH 7.4), followed by extrusion with a Mini-Extruder (Avanti Polar Lipids, Alabaster, AL) through

a 0.1 µm Nuclepore Track-Etch membrane (Whatman, Maidstone, United Kingdom). The final

large unilamellar vesicles (LUVs) stock concentration was 4 mM. LUVs contain 10% POPS and

90% POPC.

Circular dichroism (CD)

CD experiments were performed as described previously (Nguyen et al., 2015) . Briefly, the

secondary structure of PITCR in aqueous solution (10 mM sodium phosphate, pH 7.4) and

liposomes (10 % POPS and 90 % POPC LUVs at pH 5.0 and pH 7.4, respectively) was

determined in a Jasco J-815 spectropolarimeter at RT. The Circular Dichroism (CD) spectra

were measured from 195 nm to 260 nm in a 2 mm path length quartz cuvette. The peptide to

lipid molar ratio was 1: 200 and the final concentration of the peptide was 5 µM. To obtain the

desired pH, the experimental samples were adjusted by adding either 100 mM sodium

phosphate (pH 7.4) or 100 mM sodium acetate (pH 4.0). The appropriate liposome or buffer

backgrounds were subtracted. Molar ellipticity was calculated with the following equation: $[\theta] =$

$\theta/[10lc(N-1)]$, where θ is the measured ellipticity in millidegree, l is the cell path length, c is

the protein concentration and N is the number of amino acids (*N*= 33).

pK $_{CD}$ determination assay

The apparent pK $_{CD}$ is defined as a pH midpoint, where half of the peptide changes its

secondary structure in presence of liposomes (Scott et al., 2017). The liposome preparation (10

% POPS and 90% POPC LUVs) was followed as described in liposome preparation. To reach a

series of different desired pH values, the experimental samples were adjusted with either 100

mM sodium phosphate or 100 mM sodium acetate. The final pH was measured by a 2.5-mm

bulb pH electrode (Microelectrodes, Bedford, NH) after recording CD spectrum. The CD spectra

were recorded from 220 nm to 262 nm. The appropriate liposome blanks were subtracted. The

ellipticity values at 222 nm were subtracted from that at 262 nm. The subtracted ellipticity were

plotted for a range of pH values. The pK $_{CD}$ was fitted in the following equation: $Signal =$

$\frac{[(F_a+S_a \times pH)+(F_b+S_b \times pH) \times 10^{[m \times (pH-pK)]}]}{[1+10^{[m \times (pH-pK)]}]}$, where F_a is the acidic baseline, F_b is the basic baseline, m

is the slope of the transition, and pK is the midpoint of the curve.

Peptide conjugation

Cysteine was added to the N-terminal of PITCR, termed NEC-PITCR (sequence:

ECDPKLSYLLDGILFGYGVELTALFLEVGFSESAD). NEC-PITCR was labeled with dylight680

maleimide (Thermo Scientific-46618) and AZDye 555 maleimide (Fluoroprobes-1168-1). Both

dyes labeled peptides were purified using reverse phase HPLC purification to remove

unconjugated dye. The molecular weight was confirmed by MALDI-TOF. After that, samples

were aliquoted, lyophilized and stored at -80 °C.

MALDI-TOF mass spectrometry

PITCR associated peptides were dissolved in 1 mM sodium phosphate buffer (pH 7.4, filtered).

The matrix α-Cyano-4-hydroxycinamic acid (TCI C1768), was dissolved in 75 % HPLC-level

acetonitrile coupled with 0.1 % TFA and sonicated 15 minutes, RT. The dissolved samples were

mixed with the dissolved matrix. After that, the mixed matrix samples were loaded onto the MSP

target plate (Bruker, Billerica, United States) drop by drop and dried using filtered air. The

Bruker Microflex MALDI-TOF mass spectrometer (Bruker, Billerica, United States) was

calibrated with ProteoMass MALDI-MS calibration standards (Sigma-Aldrich I6279-5X1VL,

I6154-5X1VL, C8857-5X1VL and P2613-5X1VL). All PITCR associated peptides were

measured in a negative mode. The pHLIP was measured in a positive mode. Data were

analyzed using FlexAnalysis software (Bruker, Billerica, United States) and graphs were plotted

using Origin 9.1 (research lab) software.

HPLC

All peptides were dissolved in 1 mM sodium phosphate buffer (pH 7.4, filtered). Dissolved

peptides were injected into an Agilent 1200 series HPLC system (Agilent Technologies, Santa

Clara, United States). A semi-preparative Agilent Zorbax 300 SB-C18 column (P.N. 880995-

202) was used to purify the PITCR associated peptides. A stable-bond analytical Agilent Zorbax

300 SB-C18 column (P.N. 880995-902) was used to identify the purity of each peptide. The

running procedure used a gradient (solvent A: 0.05% TFA HPLC-level water plus solvent B:

0.05% TFA HPLC-level acetonitrile) from 5% B to 100% B. PITCR associated peptides were

eluted around 80% B.

Immunoprecipation and immunoblot analysis

Jurkat-WT T cells were washed twice with PBS and treated with PITCR at 37 °C and 5 % CO_2

for 30 mins, followed by stimulation with 1 µM anti-CD3 antibody (OKT3 clone). Cells were lysed

in cold 0.5% Dodecyl-β-D-maltopyranoside (DDM, VWR-97063-172) lysis buffer (20 mM Tris-Cl

pH 8.0, 10 mM NaF, 166.67 mM NaCl, 20 mM iodoacetamide, Benzonase endonuclease 50

U/ml (Sigma-1016970001), proteinase inhibitors (Thermo Scientific A32955) and phosphatase

inhibitors (Sigma-Aldrich P0044 and P5726)) for 30 minutes on ice, followed by centrifugation of

16.2 x 10^3 g, 15 mins, 4 °C. Supernatants were collected and detergent compatible protein

assay was performed to quantify the protein concentration of each sample. 30-50 µl whole

lysates were saved to detect the TCR complex. Protein G agarose and anti-CD3ε (UCHT1

clone) were used as described in the methodology of co-IP of TCR. Samples were continuously

rotated at 4 °C for 4 hrs. Other steps follow the methodology of co-IP of TCR, except the wash

buffer (20 mM Tris-Cl pH 8.0, 10 mM NaF, 166.67 mM NaCl, 20 mM iodoacetamide, Benzonase

endonuclease 50 U/ml, Sigma-1016970001) and the elution condition (70 °C for 10 mins).

Immunoblot analysis were performed as described in the methodology of co-IP of TCR.

Antibodies

Antibodies	Sources	Catalogue #	Dilutions
Zap70 (pY319)	Cell Signaling Technology	2717	1:1000 (WB)
Zap70 (pY493)	Cell Signaling Technology	2704	1:1000 (WB)
Zap70 (total)	Cell Signaling Technology	3165	1:1000 (WB)
LAT (pY191)	Cell Signaling Technology	3584 (discontinued)	1:1000 (WB)
LAT (pY132)	Abcam	ab4476	1:2000 (WB)
LAT total	Cell Signaling Technology	45533	1:1000 (WB)
Lck (pY394)	R & D systems	755103	1:2000 (WB)
Lck (pY505)	Cell Signaling Technology	2751	1:1000 (WB)
Lck (total)	Cell Signaling Technology	2984	1:1000 (WB)

β-actin	Abcam	ab6276	1:5000 (WB)
CD3 ε (OKT3)	Tonbo biosciences	70-0037	1:50 (stimulation)
CD3 ε (UCHT1)	Santa Cruz biotechnology	sc-1179	1: 200 (IF), IP (see methodology)
CD3 ε (CD3-12)	Cell Signaling Technology	4443	1:1000 (WB)
ζ (6B10.2)	Santa Cruz biotechnology	sc-1239	1:500 (WB)
ζ (pY142)	BD Biosciences	558402	1:1000 (WB)
ζ (pY83)	Abcam	ab68236	1:1000 (WB)
TCR β	Cell Signaling Technology	77046	1:2000 (WB)
SLP76 (pY128)	BD Biosciences	558367	1:2000 (WB)
PLCγ1 (pY783)	Cell Signaling Technology	2821	1:1000 (WB)
IgG	Cell Signaling Technology	5415	IP (see methodology)
CD69-APC (FN50)	Biolegend	50-166-584	1:100 (Flowcytometry)
IgG-APC	Biolegend	50-168-838	Flowcytometry
Goat anti-Mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, Alexa Fluor™ Plus 488	ThermoFisher Scientific (Invitrogen)	A32723	1: 200 (IF)
IRDye® 800CW Goat anti-Rabbit IgG Secondary Antibody	LI-COR Bioscience	926-32211	1:5000 (WB)
IRDye® 800CW Goat anti-Mouse IgG Secondary Antibody	LI-COR Bioscience	926-32210	1:5000 (WB)
IRDye® 800CW Goat anti-Rat IgG Secondary Antibody	LI-COR Bioscience	925-32219	1:5000 (WB)

TIRF imaging of AND-TCR T cells

AND-TCR T cells were incubated with PITCR-AZDye555 by mixing 50 μL of 100 μM solution in 10 mM sodium phosphate buffer pH 7.4 with 450 μL of 2.5 M cells/mL T cells in RVC medium with IL-2 (final concentration of 10 μM PITCR-AZDye555, 2.2 M cells/mL, 37 °C, 30 min), rinsed by imaging buffer, then applied to the imaging chamber with SLB functionalized with ICAM-1 (~20 molecules/μm2) and pMHC (19-23 molecules/μm2, MCC peptide 9.1% labeled with Atto647N) at 37 °C. The real-time images of just-adhering cells with RICM, TIRF at 561 nm excitation (50 ms exposure), and TIRF at 640 nm excitation (500 ms exposure) channels were recorded at the average frame rate of 1 frame per 4 seconds. After 15-30 min, the snapshots of the cells forming immune synapses were recorded with the same three channels. The experiment was performed with 5 cells (real-time) and about 50-100 cells (snapshots) from 1 mouse. The cells with dominant plasma-membrane-bound PITCR signal could be found only in snapshot measurements due to low population. The vehicle control was performed by treating the cells with phosphate buffer instead of PITCR-AZDye555 solution using the cells from the same mouse.

Reagents

T cell culture medium: DMEM (Gibco, Thermo Fisher) with 10% FBS, 1 mM sodium pyruvate, 2 mM L-glutamine, 1x Corning nonessential amino acids (Fisher Scientific), 1x Corning MEM vitamin solution (Fisher Scientific), 0.67 mM L-arginine, 0.27 mM L-asparagine, 14 μM folic acid, 1x Corning Penicillin/Streptomycin (100 IU, 0.1 mg/mL respectively) (Fisher Scientific), 50 μM β-mercaptoethanol.

Imaging buffer for TIRF experiment: 20 mM HEPES, 137 mM NaCl, 5 mM KCl, 1 mM $MgCl_2$, 1.8 mM $CaCl_2$, 0.1% w/v D-glucose, 0.1% w/v BSA, pH 7.4.

AND-TCR T cell culture

Primary AND-TCR T cells were prepared and cultured basically as previously described (Smith et al., 2011).T cells from the lymph nodes and spleens were harvested from (B10.Cg-Tg(TcrAND)53Hed/J) × (B10.BR-H2k2 H2-T18a/SgSnJ) hybrid mice (Jackson Laboratory) and kept in T cell culture medium (day 1). The cells were activated by 2 μM MCC peptide at day 1, then cultured with IL-2 after day 2. Cells were used at day 5 or 6 for imaging. All animal work was performed with prior approval by Lawrence Berkeley National Laboratory Animal Welfare and Research Committee under the approved protocol 17702.

pMHC and ICAM-1 preparation

ICAM-1 extracellular domain with His10 tag and MHC class II I-Ek with two His6 tags were

expressed and purified as previously described (Nye & Groves, 2008).

Peptides for pMHC were prepared and loaded to MHC molecule as previously described

(O'Donoghue et al., 2013)MCC peptide (ANERADLIAYLKQATK) and MCC-GGSC

(ANERADLIAYLKQATKGGSC) were synthesized on campus (D. King, Howard Hughes Medical

Institute Mass Spectrometry Laboratory at University of California, Berkeley) or commercially

(Elim Biopharmaceuticals, Hayward, CA). MCC-GGSC was labeled with Atto647N-maleimide

(ATTO-TEC), purified by C18 column reversed phase HPLC, and identified by MALDI-TOF

mass spectrometry.

Excess amount of MCC and MCC-GGSC-Atto647N were separately loaded on MHC molecules

in loading buffer (PBS acidified with citric acid to pH 4.5, 1% BSA) over night at 37°C. Then

mixed at a 10:1 molar ratio to achieve 9.1% labeling efficiency. The mixture was diluted-

concentrated with TBS and 10 kDa MWCO filters (Spin-X UF, Corning) for two times to remove

excess peptides, then used for bilayer functionalization.

Supported lipid bilayer preparation

Small unilamellar vesicles with 98 mol% 1,2-dioleoyl-sn-glycero-3-phosphocholine (DOPC,

Avanti Polar Lipids) and 2 mol% 1,2-dioleoyl-sn-glycero-3-[(N-(5-amino-1-

carboxypentyl)iminodiacetic acid)succinyl] nickel salt (DGS-NTA, Avanti Polar Lipids) was

prepared by sonicating 0.5 mg/mL lipid suspension in water followed by centrifugation (21,000

g, 20 min, 4°C). Then, supported lipid bilayer (SLB) was prepared upon 25 mm #1.5 glass

coverslip set into Attofluor cell chamber (Invitrogen, Thermo Fisher). Coverslips were cleaned

by sonication in 1:1 water:2-propanol then etched with piranha solution (1:3 mixture of 30%

H_2O_2 and sulfuric acid), rinsed by water and set into clean chambers. SLB was formed by

adding 1:1 mixture of SUV solution and TBS into chambers and incubating for more than 30

min. SLB was rinsed by TBS then incubated with 10 mM $NiCl_2$ in TBS for 5 min. Chambers were

then incubated in imaging buffer for more than 30 min for blocking defects by BSA, then used

for functionalization. ICAM-1 (~10 nM) and pMHC (concentration adjusted by determined

densities) were added to chambers and incubated for 30 min, then rinsed by imaging buffer.

The ICAM-1 density is estimated to be ~20 molecule/μm^2 based on a previously reported

estimate (Lin et al., 2019). pMHC labeled by Atto647N was imaged by TIRF to determine the

molecular density. The density was determined by extrapolating the calibration curve of density-

intensity relationship from lower densities where the molecular density can be directly

determined by single molecule localization (below 0.5 molecules/μm^2) using TrackMate

(Tinevez et al., 2017).

TIRF microscopy and image processing

TIRF microscopy was performed on a motorized inverted microscope (Nikon Eclipse Ti-E;

Technical Instruments, Burlingame, CA) with Lumen Dynamics X-Cite® 120LED Fluorescence

Illumination System (Excelitas Technologies, Waltham, MA, USA) and a motorized stage (MS-

2000; Applied Scientific Instrumentation, Eugene, OR). A laser launch with 561-, and 640-nm

diode lasers (Coherent OBIS, Santa Clara, CA) was aligned into a custom-built fiber launch

(Solamere Technology Group Inc., Salt Lake City, UT). For TIRF imaging, laser excitation was

illuminated through a four bands beam splitter (ZT405/488/561/640rpc) to the objective lens (NA

1.49, 100×, oil immersion, Apochromat TIRF, Nikon), then filtered through an emission filter

(ET600/50M or ET700/75M). For RICM, LED excitation was illuminated through D546/10x

excitation filter and 50/50 beam splitter. Emission was captured on an EM-CCD (iXon Ultra 897;

Andor Inc., South Windsor, CT). All optical filters were purchased from Chroma Technology

Corp. (Bellows Falls, VT). The sample and objective lens were kept at 37 °C with temperature

controller system (CU-109, Live Cell Instrument, Republic of Korea). The equipment was

controlled using the software MicroManager (Edelstein et al., 2010) Laser power and exposure

time was set to 2 mW, 50 ms for 561 nm excitation and 1 mW, 500 ms for 640 nm excitation.

The pixel size was 0.16 μm square and the field of view was 81.92 μm square (512×512 pixels).

Cell footprint was determined from RICM images with the following procedures. RICM image

was gaussian-blurred (sigma: 2 pixels), manually background-subtracted, and converted to

absolute values pixel-wise. The obtained intensity images were segmented by semi-automatic

way: the inner region and the encompassing region of the cell of interest were manually

selected. Image was thresholded by the intensity in the inner region multiplied by an arbitrary

factor of 0.5. The obtained segment was filtered within the encompassing region, then cleaned

by binary opening (kernel: 3×3 pixels square). The regions smaller than 200 pixels were deleted

and the remaining regions (multiple regions were allowed to exist if any) were used as the cell

footprint.

The background signal was measured using the chamber containing only imaging buffer and

subtracted from TIRF images. The inhomogeneity of the TIRF illumination were corrected using

the images from the solution of rhodamine B (561 nm excitation, from Sigma Aldrich) and 3,3'-

diethylthiadicarbocyanine iodide (640 nm excitation, from Sigma Aldrich).

Statistical analysis

All statistical analyses of experiments were performed using GraphPad Prism 9.4.0. P values are provided as exact values. 95% confidence level was used to determine statistically significance in all experiments and ns stands for not significant. All statistical correspond to biological replicates only and all n values reflect biological replicates. Detailed statistical analyses were illustrated in each result.

Acknowledgements

This work was supported by grant R35GM140846 (to F.N.B.), and by a Faculty-Graduate student award to YYJ (University of Tennessee). We thank Art Weiss (UCSF) for generous advice to YYJ and for providing J.OT1.CD8 and T2Kb cells, and to L. Teyton (Scripps Research) and M. Davis (Stanford University) for providing the MHC and ICAM-1 bacmids. We also appreciate the advice provided to YYJ by Barry Bruce (University of Tennessee), Matthew Call (Walter and Eliza Hall Institute of Medical Research), and Peiqing Sun (Wake Forest Baptist Medical Center). We are also thankful for the technical advice of Tim Sparer and Trevor Hancock (University of Tennessee), Jaydeep Kolape (AMIC, University of Tennessee) and Ed Wright (BRF, University of Tennessee). We appreciate the members of the Barrera lab Jen Schuster, Jordan Pyron and Boomer Russell for insights on the manuscript.

Competing Interests

The authors declare that not competing or non-competing interest exist.

References

Alves, D. S., Westerfield, J. M., Shi, X., Nguyen, V. P., Stefanski, K. M., Booth, K. R., Kim, S., Morrell-Falvey, J., Wang, B. C., Abel, S. M., Smith, A. W., & Barrera, F. N. (2018). A novel pH-dependent membrane peptide that binds to EphA2 and inhibits cell migration. *Elife*, *7*. https://doi.org/10.7554/eLife.36645

Andreev, O. A., Dupuy, A. D., Segala, M., Sandugu, S., Serra, D. A., Chichester, C. O., Engelman, D. M., & Reshetnyak, Y. K. (2007). Mechanism and uses of a membrane peptide that targets tumors and other acidic tissues in vivo. *Proc Natl Acad Sci U S A*, *104*(19), 7893-7898. https://doi.org/10.1073/pnas.0702439104

Ashouri, J. F., Lo, W. L., Nguyen, T. T. T., Shen, L., & Weiss, A. (2022). TZAP70, too little, too much can lead to autoimmunity. *Immunol Rev*, *307*(1), 145-160. https://doi.org/10.1111/imr.13058

Au-Yeung, B. B., Shah, N. H., Shen, L., & Weiss, A. (2018). ZAP-70 in Signaling, Biology, and Disease. *Annu*
Rev Immunol, *36*, 127-156. https://doi.org/10.1146/annurev-immunol-042617-053335
Biswas, K. H., & Groves, J. T. (2019). Hybrid Live Cell-Supported Membrane Interfaces for Signaling
Studies. *Annu Rev Biophys*, *48*, 537-562. https://doi.org/10.1146/annurev-biophys-070317-
033330
Brazin, K. N., Mallis, R. J., Boeszoermenyi, A., Feng, Y., Yoshizawa, A., Reche, P. A., Kaur, P., Bi, K., Hussey,
R. E., Duke-Cohan, J. S., Song, L., Wagner, G., Arthanari, H., Lang, M. J., & Reinherz, E. L. (2018).
The T Cell Antigen Receptor alpha Transmembrane Domain Coordinates Triggering through
Regulation of Bilayer Immersion and CD3 Subunit Associations. *Immunity*, *49*(5), 829-841 e826.
https://doi.org/10.1016/j.immuni.2018.09.007
Bromley, S. K., Burack, W. R., Johnson, K. G., Somersalo, K., Sims, T. N., Sumen, C., Davis, M. M., Shaw, A.
S., Allen, P. M., & Dustin, M. L. (2001). The immunological synapse. *Annu Rev Immunol*, *19*, 375-
396. https://doi.org/10.1146/annurev.immunol.19.1.375
Call, M. E., Pyrdol, J., Wiedmann, M., & Wucherpfennig, K. W. (2002). The organizing principle in the
formation of the T cell receptor-CD3 complex. *Cell*, *111*(7), 967-979.
https://doi.org/10.1016/s0092-8674(02)01194-7
Call, M. E., Schnell, J. R., Xu, C., Lutz, R. A., Chou, J. J., & Wucherpfennig, K. W. (2006). The structure of
the zetazeta transmembrane dimer reveals features essential for its assembly with the T cell
receptor. *Cell*, *127*(2), 355-368. https://doi.org/10.1016/j.cell.2006.08.044
Chai, J. (2020). Atomic-resolution view of complete TCR-CD3 revealed. *Protein Cell*, *11*(3), 158-160.
https://doi.org/10.1007/s13238-019-00677-7
Chakraborty, A. K., & Weiss, A. (2014). Insights into the initiation of TCR signaling. *Nat Immunol*, *15*(9),
798-807. https://doi.org/10.1038/ni.2940
Cheng, C. J., Bahal, R., Babar, I. A., Pincus, Z., Barrera, F., Liu, C., Svoronos, A., Braddock, D. T., Glazer, P.
M., Engelman, D. M., Saltzman, W. M., & Slack, F. J. (2015). MicroRNA silencing for cancer
therapy targeted to the tumour microenvironment. *Nature*, *518*(7537), 107-110.
https://doi.org/10.1038/nature13905
Costes, S. V., Daelemans, D., Cho, E. H., Dobbin, Z., Pavlakis, G., & Lockett, S. (2004). Automatic and
quantitative measurement of protein-protein colocalization in live cells. *Biophys J*, *86*(6), 3993-
4003. https://doi.org/10.1529/biophysj.103.038422
Courtney, A. H., Amacher, J. F., Kadlecek, T. A., Mollenauer, M. N., Au-Yeung, B. B., Kuriyan, J., & Weiss,
A. (2017). A Phosphosite within the SH2 Domain of Lck Regulates Its Activation by CD45. *Mol*
Cell, *67*(3), 498-511 e496. https://doi.org/10.1016/j.molcel.2017.06.024
Courtney, A. H., Lo, W. L., & Weiss, A. (2018). TCR Signaling: Mechanisms of Initiation and Propagation.
Trends Biochem Sci, *43*(2), 108-123. https://doi.org/10.1016/j.tibs.2017.11.008
Davis, M. M., & Bjorkman, P. J. (1988). T-cell antigen receptor genes and T-cell recognition. *Nature*,
334(6181), 395-402. https://doi.org/10.1038/334395a0
Dong, Zheng, L., Lin, J., Zhang, B., Zhu, Y., Li, N., Xie, S., Wang, Y., Gao, N., & Huang, Z. (2019). Structural
basis of assembly of the human T cell receptor-CD3 complex. *Nature*, *573*(7775), 546-552.
https://doi.org/10.1038/s41586-019-1537-0
Edelstein, A., Amodaj, N., Hoover, K., Vale, R., & Stuurman, N. (2010). Computer control of microscopes
using microManager. *Curr Protoc Mol Biol*, *Chapter 14*, Unit14 20.
https://doi.org/10.1002/0471142727.mb1420s92
Freeman, J. D., Warren, R. L., Webb, J. R., Nelson, B. H., & Holt, R. A. (2009). Profiling the T-cell receptor
beta-chain repertoire by massively parallel sequencing. *Genome Res*, *19*(10), 1817-1824.
https://doi.org/10.1101/gr.092924.109

Ganti, R. S., Lo, W. L., McAffee, D. B., Groves, J. T., Weiss, A., & Chakraborty, A. K. (2020). How the T cell
signaling network processes information to discriminate between self and agonist ligands. *Proc*
Natl Acad Sci U S A, *117*(42), 26020-26030. https://doi.org/10.1073/pnas.2008303117
Guy, C., Mitrea, D. M., Chou, P. C., Temirov, J., Vignali, K. M., Liu, X., Zhang, H., Kriwacki, R., Bruchez, M.
P., Watkins, S. C., Workman, C. J., & Vignali, D. A. A. (2022). LAG3 associates with TCR-CD3
complexes and suppresses signaling by driving co-receptor-Lck dissociation. *Nat Immunol*, *23*(5),
757-767. https://doi.org/10.1038/s41590-022-01176-4
Kidman, J., Principe, N., Watson, M., Lassmann, T., Holt, R. A., Nowak, A. K., Lesterhuis, W. J., Lake, R. A.,
& Chee, J. (2020). Characteristics of TCR Repertoire Associated With Successful Immune
Checkpoint Therapy Responses. *Front Immunol*, *11*, 587014.
https://doi.org/10.3389/fimmu.2020.587014
Kuhns, M. S., & Davis, M. M. (2008). The safety on the TCR trigger. *Cell*, *135*(4), 594-596.
https://doi.org/10.1016/j.cell.2008.10.033
Lanz, A. L., Masi, G., Porciello, N., Cohnen, A., Cipria, D., Prakaash, D., Balint, S., Raggiaschi, R., Galgano,
D., Cole, D. K., Lepore, M., Dushek, O., Dustin, M. L., Sansom, M. S. P., Kalli, A. C., & Acuto, O.
(2021). Allosteric activation of T cell antigen receptor signaling by quaternary structure
relaxation. *Cell Rep*, *36*(6), 109531. https://doi.org/10.1016/j.celrep.2021.109531
Lee, M. S., Glassman, C. R., Deshpande, N. R., Badgandi, H. B., Parrish, H. L., Uttamapinant, C., Stawski, P.
S., Ting, A. Y., & Kuhns, M. S. (2015). A Mechanical Switch Couples T Cell Receptor Triggering to
the Cytoplasmic Juxtamembrane Regions of CD3zetazeta. *Immunity*, *43*(2), 227-239.
https://doi.org/10.1016/j.immuni.2015.06.018
Lewis, R. S. (2001). Calcium Signaling Mechanisms in T Lymphocytes. *Annual Review of Immunology*,
19(1), 497-521. https://doi.org/10.1146/annurev.immunol.19.1.497
Lin, J. J. Y., Low-Nam, S. T., Alfieri, K. N., McAffee, D. B., Fay, N. C., & Groves, J. T. (2019). Mapping the
stochastic sequence of individual ligand-receptor binding events to cellular activation: T cells act
on the rare events. *Sci Signal*, *12*(564). https://doi.org/10.1126/scisignal.aat8715
Lo, W. L., Shah, N. H., Ahsan, N., Horkova, V., Stepanek, O., Salomon, A. R., Kuriyan, J., & Weiss, A.
(2018). Lck promotes Zap70-dependent LAT phosphorylation by bridging Zap70 to LAT. *Nat*
Immunol, *19*(7), 733-741. https://doi.org/10.1038/s41590-018-0131-1
Lo, W. L., Shah, N. H., Rubin, S. A., Zhang, W., Horkova, V., Fallahee, I. R., Stepanek, O., Zon, L. I., Kuriyan,
J., & Weiss, A. (2019). Slow phosphorylation of a tyrosine residue in LAT optimizes T cell ligand
discrimination. *Nat Immunol*, *20*(11), 1481-1493. https://doi.org/10.1038/s41590-019-0502-2
Lo, W. L., & Weiss, A. (2021). Adapting T Cell Receptor Ligand Discrimination Capability via LAT. *Front*
Immunol, *12*, 673196. https://doi.org/10.3389/fimmu.2021.673196
Mariuzza, R. A., Agnihotri, P., & Orban, J. (2020). The structural basis of T-cell receptor (TCR) activation:
An enduring enigma. *J Biol Chem*, *295*(4), 914-925. https://doi.org/10.1074/jbc.REV119.009411
Marunaka, Y. (2015). Roles of interstitial fluid pH in diabetes mellitus: Glycolysis and mitochondrial
function. *World J Diabetes*, *6*(1), 125-135. https://doi.org/10.4239/wjd.v6.i1.125
McAffee, D. B., O'Dair, M. K., Lin, J. J., Low-Nam, S. T., Wilhelm, K. B., Kim, S., Morita, S., & Groves, J. T.
(2021). Discrete LAT condensates encode antigen information from single pMHC:TCR binding
events. *bioRxiv*, 2021.2012.2016.472676. https://doi.org/10.1101/2021.12.16.472676
Michaux, A., Mauen, S., Breman, E., Dheur, M. S., Twyffels, L., Saerens, L., Jacques-Hespel, C., Gauthy, E.,
Agaugue, S., Gilham, D. E., & Sotiropoulou, P. A. (2022). Clinical Grade Manufacture of CYAD-
101, a NKG2D-based, First in Class, Non-Gene-edited Allogeneic CAR T-Cell Therapy. *J*
Immunother, *45*(3), 150-161. https://doi.org/10.1097/CJI.0000000000000413
Mossman, K. D., Campi, G., Groves, J. T., & Dustin, M. L. (2005). Altered TCR signaling from geometrically
repatterned immunological synapses. *Science*, *310*(5751), 1191-1193.
https://doi.org/10.1126/science.1119238

Nguyen, V. P., Alves, D. S., Scott, H. L., Davis, F. L., & Barrera, F. N. (2015). A Novel Soluble Peptide with
733   pH-Responsive Membrane Insertion. *Biochemistry*, *54*(43), 6567-6575.
734   https://doi.org/10.1021/acs.biochem.5b00856

735   Nye, J. A., & Groves, J. T. (2008). Kinetic control of histidine-tagged protein surface density on supported
736   lipid bilayers. *Langmuir*, *24*(8), 4145-4149. https://doi.org/10.1021/la703788h

737   O'Donoghue, G. P., Pielak, R. M., Smoligovets, A. A., Lin, J. J., & Groves, J. T. (2013). Direct single
738   molecule measurement of TCR triggering by agonist pMHC in living primary T cells. *Elife*, *2*,
739   e00778. https://doi.org/10.7554/eLife.00778

740   Pielak, R. M., O'Donoghue, G. P., Lin, J. J., Alfieri, K. N., Fay, N. C., Low-Nam, S. T., & Groves, J. T. (2017).
741   Early T cell receptor signals globally modulate ligand:receptor affinities during antigen
742   discrimination. *Proc Natl Acad Sci U S A*, *114*(46), 12190-12195.
743   https://doi.org/10.1073/pnas.1613140114

744   Prakaash, D., Cook, G. P., Acuto, O., & Kalli, A. C. (2021). Multi-scale simulations of the T cell receptor
745   reveal its lipid interactions, dynamics and the arrangement of its cytoplasmic region. *PLoS*
746   *Comput Biol*, *17*(7), e1009232. https://doi.org/10.1371/journal.pcbi.1009232

747   Reinherz, E. L. (2019). The structure of a T-cell mechanosensor. *Nature*, *573*(7775), 502-504.
748   https://doi.org/10.1038/d41586-019-02646-w

749   Schamel, W. W., Alarcon, B., & Minguet, S. (2019). The TCR is an allosterically regulated macromolecular
750   machinery changing its conformation while working. *Immunol Rev*, *291*(1), 8-25.
751   https://doi.org/10.1111/imr.12788

752   Scott, H. L., Heberle, F. A., Katsaras, J., & Barrera, F. N. (2019). Phosphatidylserine Asymmetry Promotes
753   the Membrane Insertion of a Transmembrane Helix. *Biophys J*, *116*(8), 1495-1506.
754   https://doi.org/10.1016/j.bpj.2019.03.003

755   Scott, H. L., Westerfield, J. M., & Barrera, F. N. (2017). Determination of the Membrane Translocation pK
756   of the pH-Low Insertion Peptide. *Biophys J*, *113*(4), 869-879.
757   https://doi.org/10.1016/j.bpj.2017.06.065

758   Smith, A. W., Smoligovets, A. A., & Groves, J. T. (2011). Patterned two-photon photoactivation
759   illuminates spatial reorganization in live cells. *J Phys Chem A*, *115*(16), 3867-3875.
760   https://doi.org/10.1021/jp108295s

761   Tinevez, J. Y., Perry, N., Schindelin, J., Hoopes, G. M., Reynolds, G. D., Laplantine, E., Bednarek, S. Y.,
762   Shorte, S. L., & Eliceiri, K. W. (2017). TrackMate: An open and extensible platform for single-
763   particle tracking. *Methods*, *115*, 80-90. https://doi.org/10.1016/j.ymeth.2016.09.016

764   Trebak, M., & Kinet, J. P. (2019). Calcium signalling in T cells. *Nat Rev Immunol*, *19*(3), 154-169.
765   https://doi.org/10.1038/s41577-018-0110-7

766   Wolpert, E. Z., Petersson, M., Chambers, B. J., Sandberg, J. K., Kiessling, R., Ljunggren, H. G., & Karre, K.
767   (1997). Generation of CD8+ T cells specific for transporter associated with antigen processing
768   deficient cells. *Proc Natl Acad Sci U S A*, *94*(21), 11496-11501.
769   https://doi.org/10.1073/pnas.94.21.11496

770   Wong, J., Obst, R., Correia-Neves, M., Losyev, G., Mathis, D., & Benoist, C. (2007). Adaptation of TCR
771   repertoires to self-peptides in regulatory and nonregulatory CD4+ T cells. *J Immunol*, *178*(11),
772   7032-7041. https://doi.org/10.4049/jimmunol.178.11.7032

773   Xu, C., Gagnon, E., Call, M. E., Schnell, J. R., Schwieters, C. D., Carman, C. V., Chou, J. J., &
774   Wucherpfennig, K. W. (2008). Regulation of T cell receptor activation by dynamic membrane
775   binding of the CD3epsilon cytoplasmic tyrosine-based motif. *Cell*, *135*(4), 702-713.
776   https://doi.org/10.1016/j.cell.2008.09.044

777   Yu, Y., Fay, N. C., Smoligovets, A. A., Wu, H. J., & Groves, J. T. (2012). Myosin IIA modulates T cell
778   receptor transport and CasL phosphorylation during early immunological synapse formation.
779   *PLoS One*, *7*(2), e30704. https://doi.org/10.1371/journal.pone.0030704

780   Zhang, H., Cordoba, S. P., Dushek, O., & van der Merwe, P. A. (2011). Basic residues in the T-cell receptor
781   zeta cytoplasmic domain mediate membrane association and modulate signaling. *Proc Natl Acad*
782   *Sci U S A*, *108*(48), 19323-19328. https://doi.org/10.1073/pnas.1108052108