Using developmental rules to align microevolution with macroevolution

ABSTRACT

Macroevolutionary biologists have classically rejected the notion that higher level patterns of divergence arise through microevolutionary processes acting within populations. For morphology, this consensus partly derives from the inability of quantitative genetics models to correctly predict the behavior of evolutionary processes at the scale of millions of years. Developmental studies (evo-devo) have been proposed to reconcile micro and macroevolution. However, there has been little progress in establishing a formal framework to apply evo-devo models of phenotypic diversification. Here, we reframe this issue by asking if using evo-devo models to quantify biological variation can improve the explanatory power of comparative models, thus helping us bridge the gap between micro- and macroevolution. We test this prediction by evaluating the evolution of primate lower molars in a comprehensive dataset densely sampled across living and extinct taxa. Our results suggest that biologically-informed morphospaces alongside quantitative genetics models allow a seamless transition between the micro and macro scales, while biologically uninformed spaces do not. We show that the adaptive landscape for primate teeth is corridor-like, with changes in morphology within the corridor being nearly neutral. Overall, our framework provides a basis for integrating evo-devo into the modern synthesis, allowing an operational way to evaluate the ultimate causes of macroevolution.

Macroevolution | **Microevolution** | **Evo-Devo** | **Inhibitory Cascade Model** | **Morphospace**

"Macroevolution" is the field of study that aims to understand how the diversification of life occurred on our planet over large time scales[1]. Like any other historical science, it seeks to make sense of patterns over time ingrained in the fossil record and phylogenetic trees by referencing well-understood processes known from direct observations and experimentation[2]. In the case of evolutionary biology, this knowledge comes mainly from fields such as ecology and genetics, which tend to map evolutionary phenomena that take place during shorter time scales. For this reason, these studies are sometimes called "microevolution" and are designed to understand how population-level phenomena can produce evolutionary change. However, despite the presumed direct relationship between micro- and macro levels, quantitative studies have struggled to explain most macroevolutionary patterns in terms of microevolutionary processes[3–7]. Nevertheless, empirical results have consistently shown that the availability of genetic variation correlates strongly with rates of evolution[8–12], suggesting some effect of lower level microevolutionary processes at the macroevolutionary scale. The debate whether we can bridge the gap between these two scales is still open, with some arguing for their essential irreconcilability[3,13,14] and others advocating for reconciliation within the context of the modern synthesis[5,8–10,15–22].

One long-standing suggestion for bridging the gap between micro- and macroevolution has been through the study of developmental biology and ontogeny (*i.e.* evo-devo)[10,18,23–27]. This suggestion, however, has been challenging to implement. In a microevolutionary context, development can often be reasonably assumed to be a smooth genotype-to-phenotype (GP) map, *i.e.* a genotypic variation translates to phenotypic variation in a linear way, with traits being influenced by multiple genes of smaller effect. Such a smooth GP map would, in turn, allow the modeling of evolution and adaptation of the adult phenotype using a quantitative genetic framework, exactly because these classes of models entail this simplified, linear GP mapping[17–19,28]. On larger time scales, however, genetic architectures can change, selection can fluctuate, and development can be reorganized, generating non-linearities between genotypic and phenotypic divergence, even if the GP was originally smooth. The emergence of said non-linearities would then impede the straightforward extrapolation of microevolutionary processes over million-year timescales[18,26,29–31]. Therefore, in the absence of in-depth knowledge of development and the GP map, it is likely that macroevolutionary studies will find heterogeneity and discontinuities, even if the underlying genetic changes at the microevolutionary scale are relatively smooth and continuous[18]. Alternatively, if we can use developmental models as the basis of the quantification of morphology, we might smooth out some of these non-linearities, maximizing our ability to seamlessly connect micro and macroevolutionary scales[9,10,18,29,32–35].

Here, we propose a new framework for the investigation of morphological evolution over macroevolutionary time that explicitly models evolution at this scale as a consequence of underlying microevolutionary processes (Fig. 1). To deal with potential non-linearities that might arise over long time scales, we suggest the construction of developmentally-informed spaces (Fig. 1Bi), which coupled with quantitative genetics modeling (Fig. 1E-G) and comparative methods (Fig. 1D) can facilitate a conceptual bridge between micro and macro-scales. Under our proposed framework, we are able to directly compare microevolutionary-inspired models (henceforth called "microevolutionary models") to non-microevolutionary-inspired ones

that account for a wider variety of rate and state-heterogenous evolutionary processes (henceforth called "macroevolutionary models"). We test this workflow to investigate the evolution of Primate molars, which is an ideal model system for this kind of investigation. First, there is a simple yet powerful evo-devo model that describes the development and evolution of the mammalian molars, the inhibitory cascade model (ICM). The ICM models teeth proportions (*i.e.* molar row shape) as the result of a balance between inhibition and activation factors and predicts a positive relationship between the ratios of the areas of the last two molars in relation to the first one (m2/m1 and m3/m1,[34]). Second, there are several studies characterizing aspects of additive genetic variation in molars for Primates[36–39], as well as large-scale life history and demographic information for the group[40,41], parameters that are essential to model microevolutionary processes such as drift and selection (Fig. 1E-G). Third, tooth enamel is the most mineralized substance in vertebrate tissues, making teeth especially resistant to taphonomic processes and abundant in the fossil record (Fig. 1C). The use of a dense fossil record allows us to bridge some phylogenetic gaps between extant species, ensuring that heterogeneities along the tree are more likely due to differences in process rather than incomplete sampling. This extensive availability of a comprehensive combination of paleontological and neontological data enables unprecedented power to evaluate evolutionary dynamics through deep-time using data-hungry phylogenetic comparative methods[42–44]. We apply our framework to an expansive dataset of both extant (232 taxa) and extinct (248 taxa) species summarized from more than 250 different sources (see Supplementary Material), integrated with a newly published comprehensive phylogenetic context[45]. To address our hypotheses, we use a model-fitting approach based on information theory (Bayesian Information Criteria) and parsimony (minimizing model complexity/parameter number). We expect that variables devised to quantify developmental processes (ICM variables) will privilege microevolutionary models, while data embedded in biologically "naïve" spaces (those with no direct correspondence to any developmental model) will privilege complex macroevolutionary models.

Results

We used three morphospaces to test our hypothesis that evo-devo-informed quantification would provide a better bridge between micro and macroevolution (Fig S1). The first is based on the linear distances taken directly from the teeth ("distance space") and the second is based on the occlusal areas of each molar ("area space"). These two spaces are considered naïve because they make no assumptions about underlying developmental processes. As an evo-devo inspire space, we used the ICM description of the molar development[34], and constructed a morphospace based on the relation between the relative occlusal area of m2 and m3 in relation to m1 (m2/m1 and m3/m1, respectively). The result of model-based clustering showed a tendency of the ICM space to find less groups than the naïve spaces, suggesting the former is less patchy than the latter (Fig. 2).

Figure 1. Integrative evolutionary modeling framework employed in the present study. Morphological data are obtained from specimens from direct measurements or the literature (Fig. 1A). Data is then quantified and used to construct morphospaces (Fig. 1B). The morphospace construction step can either be informed by evo-devo models of trait ontogenesis (Fig. 1Bi) or be "naïve" in relation to them Fig. 1Bii). Data quantified in a certain way, together with a phylogeny (Fig. 1C) is used in an evolutionary modeling process (Fig. 1D), in which two main types of models are confronted: macroevolutionary (Fig. 1Di) and microevolutionary (Fig. 1Dii). Macroevolutionary models here are not only phenomenological (*i.e.* their parameters lack any precise underlying meaning in terms of microevolutionary processes) but also allow model heterogeneity along the tree (multi-regime models). Microevolutionary models, on the other hand, incorporates information from quantitative genetics (Fig. 1E), life history (Fig. 1F) and demographics (Fig. 1G) to model the expected rate of evolutionary change under drift.

The use of biologically naïve morphospaces favored evolutionary model complexity. Specifically, the best model for these spaces was a multi-regime multivariate BM model, which allows for rate heterogeneity along the tree (Table 1, S1,S2). For molar occlusal areas, the best model had three main regimes (Fig 3). The first regime covers most fossil groups (thus named "ancestral regime"), including Plesiadapiformes, stem-Haplorhini, part of stem-Simiiformes and Tarsiidae. The second regime refers to Strepsirrhini, both crown and stem, and the third refers to crown Simiiformes (monkeys and apes, including humans). This three-regime model was also considered a better fit than any global model (microevolutionary inspired or not) for the morphospace defined by linear distances, even though it was not the best solution found (see Supplementary Information, Table S2). In both morphospaces, the ancestral regime accumulated more variance over time than any derived regime, suggesting a weaker constraint on the former (Fig S8). It may be tempting to assign interpretations that are either biological (e.g. higher divergence rates after the K-Pg extinction event resulting from ecological opportunity) or statistical in nature (e.g. increased phylogenetic uncertainty of fossil placement resulting in upwardly biased rates[46]). However, we find that such patterns do not appear universally across morphospaces.

By contrast, the investigation of evo-devo-inspired variables based on the ICM paints a strikingly different picture (Fig. 2C). Instead of privileging more complex and heterogeneous models, the ICM morphospace privileges a single global OU process instead of one with multiple regimes (Table 1). An inspection of point estimates of rates of evolution is consistent with the identification of a single regime for this space, as estimated rates are homogeneous throughout the tree and between groups (Fig. 5,S8). Both macroevolutionary and microevolutionary versions of this model (OU and $OU_{\Sigma \propto G}$, respectively) had essentially

Figure 2. Principal component analysis of the full-sample covariance matrix for the linear distances (A), areas (B) and ICM ratios (C) morphospaces. Dots represent species averages, and colors represents groups identified in the Gaussian mixture model. Axes are not depicted to scale for convenience, so distances in the graph should not be considered representative of the metric of the underlying space.

the same BICs, suggesting that their information content is effectively the same (Table 1). However, the inspection of the confidence intervals for the macroevolutionary OU model reveals that the 95% intervals for its parameters overlap with the values implied by the microevolutionary model (Table S3). Furthermore, the microevolutionary model assumes an evolutionary rate matrix (Σ), which is proportional to the additive genetic covariance matrix of the traits (**G**-matrix; equations 1,2). This implies that, instead of optimizing values for each entry of Σ, this model only fits one proportionality parameter (κ), which makes it simpler and more parsimonious than the macroevolutionary one. Furthermore, parameter estimates of κ are within the range of values consistent with genetic drift around a single stationary optimum[40,41]. This suggests that the OU$_{\Sigma \propto G}$ model can be interpreted in terms of microevolutionary processes not only in terms of patterns but also in terms of the magnitude of variation. So, we choose the microevolutionary OU$_{\Sigma \propto G}$ as our preferred model not only because it reports the best BIC, but also because of its simplicity and biological interpretability.

Following a strict microevolutionary interpretation of our preferred model, the variation introduced by drift is aligned with the distribution of phenotypes on the ICM morphospace (Fig. 4A), suggesting that the similarity between intra and interspecific patterns of trait variation (see[11]) is consistent with drift. This is further reinforced by the investigation of node-specific rates of evolution, which shows a huge overlap with rates expected under genetic drift (Fig 5). However, drift alone would generate more variation than the total observed disparity during the period in which primates have evolved (Fig. S7A), suggesting that stabilizing selection played a crucial role in shaping the pattern of evolution in the group as well.

The investigation of the adaptive landscape implied by the best model shows that stabilizing selection is also aligned with the interspecific distribution of phenotypes (Fig. 4B). An examination of the half-lives ($t_{1/2}$) in different directions of this adaptive landscape shows that $t_{1/2}$ are higher along the activation-inhibition gradient direction of the ICM and lower in directions that would lead to deviations from the ICM (Fig. S5). These results indicate that the macroevolution of primate teeth is being shaped by a strong stabilizing selection against deviation from the ICM pattern while allowing evolution to occur along the activation-inhibition gradient, in a corridor-like manner.

Table 1. Comparison of models of primate lower row evolution fit through Maximum Likelihood.

Traits[a]	Model[b]	N_p [c]	logLik [d]	BIC [e]
Linear distances	BM	27	2585.03	-5003.38
	OU	54	2684.57	-5035.75
	$BM_{\Sigma \propto P}$	7	2108.57	-4173.93
	$OU_{\Sigma \propto P}$	34	2174.45	-4138.99
	$BM_{\Sigma \propto G}$	7	1480.63	-2918.04
	$OU_{\Sigma \propto G}$	34	1510.57	-2811.23
	Three-regime BM [f]	69	2823.03	**-5484.50** [g]
Areas	BM	9	415.80	-776.03
	OU	18	413.98	-716.83
	$BM_{\Sigma \propto P}$	4	92.39	-160.09
	$OU_{\Sigma \propto P}$	13	275.16	-470.06
	$BM_{\Sigma \propto G}$	4	74.41	-124.13
	$OU_{\Sigma \propto G}$	13	258.02	-435.78
	Three-regime BM [f]	21	471.42	**-813.19** [g]
ICM	BM	5	589.16	-1147.46
	OU^D	9	604.56	<u>-1153.55</u> [h]
	$BM_{\Sigma \propto P}$	3	538.31	-1058.10
	$OU_{\Sigma \propto P}$	8	584.68	-1119.98
	$BM_{\Sigma \propto G}$	3	575.69	-1132.86
	$OU^D_{\Sigma \propto G}$	7	598.54	**-1153.86** [g]
	Three-regime OU [f]	26	597.35	-1034.18

[a]Morphospace used to quantify molar form variation; either a biologically naïve space (linear distances or areas) or an evo-devo inspired space (ICM)

[b]Model type. Either global BM or OU models or a mixed model, which allows model and parameter heterogeneity. BM and OU can also incorporate the microevolutionary assumption that the evolutionary rate matrix (Σ) is proportional to G or P ($\Sigma \propto G$ or P models). Uppercase "D" indicates OU models with a diagonal H

[c]Number of model parameters.

[d]Log-likelihood of the model

[e]Bayesian Information Criterion used for model comparison

[f]Results for the mixed model for Linear distances and ICM are based on the best regime combination found for the areas morphospace

[g]Bold represents the best models

[h]Underline represents the ones with BIC 2 units away from the best model.

Discussion

Previous work has usually highlighted that larger-scale morphological evolution tends to conform to the expectation of microevolutionary models qualitatively but rarely (if ever) in terms of magnitudes of change[6]. In other words, while macroevolution seems to follow directions with more genetic variation, as expected due to neutral change[6, 11, 12, 21, 47, 48], rates of evolution tend to fall below those expected under genetic drift[6–8, 49]. This paradox has been used to argue for a fundamental mismatch between micro and macroevolution, as simplistic quantitative genetics models seem unlikely to represent million-year evolutionary processes[7, 50, 51]. Here, we constrained the proportionality parameter for our preferred microevolutionary-inspired model to be within realistic values for primates (equation 2). This constraint results in an estimated rate matrix that is compatible with drift around a stationary adaptive peak not only in patterns of trait association but also in magnitude. The key modeling choice that led to this conclusion was the quantification of developmentally-informed traits, which smoothed out transitions between microevolutionary and macroevolutionary data–defining and identifying a neutral subspace aligned with a conserved developmental process. A defining feature of the ICM space is its lack of discontinuities along the diversity of Primate molars (Fig. 2), which likely reduces the need for heterogeneous rates. Furthermore, by focusing on relative shape changes, which are governed by the balance of inhibitory and activation factors, this space limits the influence of other factors, such as static allometry, possibly allowing a closer match between genetic and macroevolutionary variation.

The differences we observe among morphometric representations might be partly due to how different spaces codify

size variation. Both naïve spaces contain size information, while the ICM variables do not. By using m1 size as a scaling factor, the ICM variables still include information regarding allometric variation (technically, they are unscaled versions of the Mosimann shape ratios, see[52]). Both area and distance spaces are log-scaled, meaning they fit a power-law allometric model of variation[53]. Therefore, a higher heterogeneity in size variation in the naïve spaces might privilege more complex models, while the same is not true for ICM variables. While this suggests size-correction can smooth out much of the heterogeneity in this case, this appears to derive from the fact that using shape-ratios can provide a means to quantify localized ontogenetic effects[54]. Nevertheless, without actual knowledge of developmental systems, it is hard to know beforehand that shape-ratios will necessarily lead to better conformity between micro and macroevolutionary scales. In fact, depending on the system, raw measurements and shape ratios might produce similar results[8]. Thus, studying a well-understood system such as molar development allows us to piece apart the possible role of ontogenetic models in helping us connect micro to macro scales.

Figure 3. Primate phylogenetic tree painted by the best regime combination found on the mixed model search for the individual molar areas. For areas and linear distances, the best model overall is a multi-regime BM, while for ICM ratios the best mixed model is a multi-regime OU (Table 1). However, for ICM ratios, single-regime microevolutionary models outperform all mixed models.

Previous work in primates has suggested that some traits have evolved with rates that are consistent with those expected under drift[49,55], including some dental features[56,57]. These works have largely been focused on hominin species, which could bias interpretations regarding the dental evolution of Primates. Our results partly agree with these results and extend them to the origin of the group (Fig. 5). While this could suggest that drift guided over 70 myrs of dental evolution in primates, our model-fitting approach tells otherwise. Within the set of microevolutionary-inspired models, OU models outperformed the BM models (Table 1), which suggests a crucial role of stabilizing selection in shaping macroevolutionary patterns. Considering the amount of variation introduced by drift every Myr (Fig 4A), a purely neutral process would result in overdispersion of tip values and higher phylogenetic signals (Fig S7). Instead, the patterns of stabilizing selection seem to be essential in shaping the ICM pattern by both constraining variation that deviates from the ICM pattern and facilitating evolution along the activation-inhibition gradient (Fig S5,S6).

Figure 4. Graphical representation of the best selected model ($OU^D_{\Sigma \propto G}$) for the ICM variables based on molar ratios m2/m1 and m3/m1. Dots are species averages, and horizontal and vertical lines depict ±1.96 standard deviations. Ellipses are covariance matrices representing the following parameters of the best model: A. Stochastic rate matrix Σ attributed to the amount of genetic drift introduced in the system every 1 myr. B. Individual adaptive landscape Ω based on model estimates for the rate of adaptation towards the optima (*H*). See Materials and Methods for explanations for these parameters. Multiple ellipses were calculated from parameter value combinations that are sampled along the multi-dimensional likelihood contour 2 log-likelihoods away from the peak.

Even though these two results might seem contradictory –rates of evolution consistent with drift and best model including stabilizing selection– we foresee at least two possibilities of how they both might be true: one has to do with the topography of the inferred adaptive landscape, and the other with the estimates of the evolutionary parameters. Regarding the adaptive landscape, the shape of the landscape implied by the preferred model is almost corridor-like (Fig 4B). If this corridor is relatively smooth internally (no great selection differentials within its limits), this would mean that species are free to explore this landscape neutrally, within the bounds of the corridor. Additionally, because G is aligned with the corridor as well (Fig 4A), this means most neutral changes will happen in accordance with the landscape, and will not produce great stabilizing selection. The other possibility is based on the precision of the rate-parameter estimates. Even though the OU model was the preferred one, estimated rates of evolution for BM models are remarkably similar to the ones of the full model S6–S9). Considering that node-specific rates of evolution are usually calculated under the assumption of a BM mode[58], this could mean that a dense fossil sample in a comprehensive phylogenetic framework might allow for a good estimation of rates of evolution, even under model violation. Irrespective of which is true (or even if both are), the observation that most evolutionary rates are compatible with drift is a pattern rarely seen for macroevolutionary data[6–8,49].

Together, these results point to the interplay of genetic variation, selection, and development leading to a homogeneous macroevolutionary process within a defined subspace. It has been argued that selection can mold genetic patterns of trait association and variation[28,59,60], specifically by altering developmental pathways and genetic interactions[61,62]. Conversely, development has also been argued to impose direct selective pressures (*i.e.* internal selection) by reducing the viability of non-conforming phenotypes[28,63], which could, in turn, trickle down to the organization of genetic variation. While in the present case we can observe this triple alignment between genetics, ontogeny and selection, its origins are harder to decipher. The ICM was originally described in rodents and later verified in many other mammalian groups[64,65], suggesting it is the ancestral condition for molar development in the group. In this case, ontogeny is viewed as the organizing factor behind both selective patterns and the organization of genetic variation[11]. Furthermore, this explains the near-neutral quality of primate dental evolution, as conformity to the developmental process would be the main selective pressure on relative tooth sizes[66].

Figure 5. Lande's Generalized Genetic Distance (LGGD) used to measure the node-specific rate of evolution throughout Primate divergence and diversification. Dashed lines represent the expected rates under genetic drift. For each node, a distribution of values were calculated by integrating over the variation in heritability, effective population size and generation time. Dots represent the median values and vertical lines at the 95% confidence interval.

However, some mammalian groups have been shown to deviate from the predictions of the ICM to different degrees, suggesting that the ontogenetic process itself could be malleable[64, 65, 67–69]. Indeed, it has been argued that molar tooth eruption timing in Primates is shaped by biomechanical demands at different ontogenetic stages[70], revealing a possible mechanism through which external selection could shape development, and indirectly, the morphology of the molar row.

Conclusions

To what degree microevolution can be extended to macroevolution is a central question in evolutionary biology[4]. While there is little doubt that the fundamental causes at both levels are the same (*e.g.* selection, drift, mutation), efforts to model the connection have generally failed beyond the qualitative alignment of patterns. When it comes to morphological evolution, the consensus has been overwhelmingly to reject any straightforward connection between both levels, specifically because of the fact that empirical evolutionary rates are orders of magnitude inferior to the ones expected by genetic drift[6, 7]. The results presented here reject this consensus, as we show that microevolutionary models can fit well to the data, as long as we properly choose the proper morphometric representation. Even the relatively simple task of characterizing the multivariate dimensions of three molars poses a large number of choices for measurement[10, 34, 71, 91]. Our results suggest that phenotypic quantification based on evo-devo models maximally narrows the gap between both levels of analysis–and allow for the discovery of the underlying subspaces that both qualitatively and quantitatively align macroevolutionary patterns with microevolutionary processes.

While primate molar seems unique in both the presence of a well-constrained ontogenetic model and abundance of data, other systems might also fit the requirements for the methods described here. The existence of evolutionary stable developmental pathways and modules suggests a long history of similarly stable selective pressures[24, 61, 72, 73]. This makes developmental modules good systems to investigate adaptive landscapes in deep-time[31, 54, 74]. Furthermore, assuming that these pathways are shaped by natural selection to optimize the generation of adaptive variation[59–61, 72], they are a likely place to identify simple connections between micro and macroevolutionary scales[31, 54]. So, other evolutionary stable systems are the likely candidates to verify the connection between scales of organization. Good examples are modules built upon serially homologous structures, like phalanges[75] and vertebrae[76]. For more complex structures formed by the interaction of multiple tissues, it might be harder to devise simple models that sufficiently describe the system ontogeny and variation. However, works that focus on the mammalian skull, and that used individualized bone measurements have had a good track-record of modeling multivariate evolution of these structures under microevolutionary models[12, 21, 55, 77, 78], going even further than the simple alignment between variation and evolutionary rates[8, 49, 55]. Since vertebrate skull bones are elements with deep individualized

history, measuring them individually might represent a good first approximation of the multiple morphogenic fields that interact to form the complete structure. This perspective contrasts with the regular practice of constructing morphospaces as comprehensive, phenomenological, and statistical descriptors of biological form without a clear connection to underlying biological processes[18,35,79,80]. However, given that different morphometric methods seem to point to similar overall patterns of trait variation[81], finding the correct quantification protocol might be a matter of proper scaling of morphometric variables than a radical departure from classically established measurement practices.

Lastly, our investigation also provides a new framework in which developmental biology can be more fully incorporated into macroevolutionary modeling. It has long been considered that developmental biology was left out of the evolutionary synthesis[25,26,54], and indeed such data are rarely incorporated into comparative analyses. Recent efforts have had different degrees of success, with many pointing out how complexities of the ontogenetic systems can lead to core violations of the modern synthesis[25,29,30,54,82]. By reframing the question of microevolutionary model adequacy into a problem of quantification of biological phenomena[79,80], we show how evo-devo is essential for a fully unified view in the context of the evolutionary synthesis.

Methods

Sample and morphometrics

We used the standard mesiodistal length (MD) and buccolingual breadth (BL) as basic descriptors of each molar. MD and BL were obtained for each tooth of the lower molar row (m1, m2 and m3). We obtained raw measurements from available datasets in the literature (n=6142)[42–44] and from newly measured museum specimens using a caliper (n=150). For rare species, we took measurements from images that were either published or provided to us (n=266). All photos used had a scale and were digitized using the Fiji software[83]. See the supplementary material for a full list of references and sources used for data collection. In total, we compiled a sample of 6558 individuals distributed among 480 species, divided between 232 extant and 248 extinct species. To evaluate the evolution of these traits on a phylogenetic framework, we used the most comprehensive phylogeny available that included both living and fossil primate species[45]. Our sample covered all genera and 52.98 % of the species diversity included in[45], spanning the full 75 mya of the group's evolution.

We constructed three distinct morphospaces to quantify molar variation (Fig. S1). For our biologically naïve representation of tooth form, we used a "distance space" based on linear distances obtained from each tooth and an "area space" based on each tooth's occlusal area. Occlusal molar area was approximated using a crown index (BLxMD)[64,65]. For our evo-devo-informed space, we used the ratios of the second and third molars in relation to the third (m2/m1 and m3/m1, respectively), as defined by the inhibitory cascade model of molar development[34]. We call this last morphospace the "ICM space". To evaluate morphospace patchiness we performed a clustering based on parameterized finite Gaussian mixture models[84]. This method test for a series of nested models, where groups are modeled according to different covariance structures.

Measurement error was accounted for by calculating the standard deviation of each measurement for each species. When a species had a sample of n=1, we assigned a standard deviation equal to the pooled within-group standard deviation. This implies a very high measurement error for species known from single specimens, such as the case of many fossils. The degree of genetic association between traits was approximated both by the intraspecific pooled phenotypic covariance matrix P, and an independently derived additive genetic covariance matrix G obtained from a pedigreed *Papio hamadryas* baboon population[36,37].

Phylogenetic comparative methods

To model morphological evolution, we used a maximum-likelihood (ML) model selection approach, which fits different Brownian-motion (BM) and Ornstein-Uhlenbeck (OU) models under a mixed Gaussian phylogenetic models (MGPM) framework[85]. Under this framework, the evolution of a p-dimensional multivariate trait is modeled as an OU process as follows:

$$dx(t) = -H(x(t) - \theta(t))dt + \Sigma_x dW(t) \tag{1}$$

where H is the $p \times p$ selective rate matrix, $x(t)$ is a p vector of trait values at time t, $\theta(t)$ is a p vector of each trait evolutionary optima at time t, Σ_x is the Cholesky factor of the $p \times p$ stochastic rate matrix Σ (sometimes called evolutionary rate matrix) and $W(t)$ denotes the p-dimensional standard Wiener process.

Under a strict quantitative genetics interpretation[86], the diagonal of H contains the rate of adaptation to the optima of each trait (α_p) and the off-diagonal measures the shape of co-selection among traits. Conversely, the diagonal of Σ contains the rate of evolution due to drift, with its off-diagonal elements containing the amount of coevolution due to genetic covariation. If H is a matrix of zeros, the model collapses into a multivariate BM model.

Under this microevolutionary perspective, Σ is not an entirely free parameter. Instead, if Σ is the genetic drift parameter, then it has to be proportional to the additive genetic covariance matrix G of those traits[87] as follows

$$\Sigma = G \frac{t_g}{Ne} \tag{2}$$

where t_g is the time in generations and Ne is the effective population size. Because t_g and Ne, and even the size of G are hard to estimate at evolutionary time scales, some have argued for treating t_g/Ne as a nuisance parameter, reducing the investigation of drift at the macroevolutionary scale to a simple evaluation of the proportionality between Σ and G[12,21,50]. Consistent with these suggestions, here we implement a series of proportionality models, or κ-models[11], in which Σ is set to be equal to a target matrix times a scaling factor κ. We used both the intraspecific pooled phenotypic covariance matrix P and G as target matrices. These models are implemented in the package PCMkappa (https://github.com/FabioLugar/PCMkappa).

Because the proportionality models are more tightly connected to a microevolutionary interpretation of the OU model, we call them "microevolutionary models". Full models (models where all parameters are estimated freely) are called "macroevolutionary models" because they do not have explicit microevolutionary assumptions.

We fitted two macroevolutionary BM and OU models and two microevolutionary models, using either P or G as a target matrix, for both BM and OU, totaling six global models (full BM and OU, $BM_{\Sigma \propto P}$, $BM_{\Sigma \propto G}$, $OU_{\Sigma \propto P}$, $OU_{\Sigma \propto G}$) for each morphospace. For the OU models, we investigated the confidence intervals of the parameters (see Supplementary Material) to evaluate if the model could be further reduced. Specifically, if the confidence interval of the off-diagonal elements of H overlapped with 0, another model was fit, setting H to be a diagonal matrix[85].

In addition, we performed a mixed Gaussian phylogenetic model search, which searches for the combination of regimes, models and model parameters that best fit the data[85]. For both the mixed model search and model comparison, we used the Bayesian Information Criterion (BIC), which minimizes parameter inflation due to large samples and is most appropriate for our model-selection question[88] (i.e. asymptotically identifying the data-generating process as opposed to minimizing trait prediction error). For the mixed gaussian models, we only fit full BM and OU models, and no κ-model due to software restrictions. Therefore, the mixed models are also considered macroevolutionary models.

To ensure that the κ-models were compatible with microevolutionary processes, we constrained the κ parameter to be within the range of expected values under drift, as expressed in equation 2. Because Σ is given in the tree (myr) scale, we found approximations for t_g and of Ne for Primates to infer the expected scaling factor κ. t_g was estimated as $t_g = 1myr/g_t$, where g_t is the average generation time in years obtained from[40]. Because we lack good estimates of g_t for fossil species, we used the phylogenetic average $\pm$ the standard deviation (SD) throughout the phylogeny. This was done by trimming the dataset to only the species with g_t data and obtaining the ancestral value and SD at the base through ML[89]. For Ne we used 20,000-1,000,000 as the range of possible values consistent with the genomics estimates for multiple primate species and hypothetical common ancestors[41]. While g_t and Ne are expected to vary over the tree, we assumed that the effect of this variation would be at least partially canceled out by the fact that these two quantities are generally inversely related to each other.

To evaluate the fitted model mechanistically under quantitative genetics theory, we generalized the equation for the adaptive landscape[86,90] to the multivariate case as

$$\Omega = H^{-1/2}GH^{-1/2} - P \tag{3}$$

Rates of evolution

Rates of evolution were used to evaluate if the evolutionary change conforms to the expectation of genetic drift. To calculate rates of evolution, we employed Lande's generalized genetic distance (LGGD,[87])

$$LGGD = \frac{N_e}{t_g} \Delta z^t \mathbf{G}^{-1} \Delta z \tag{4}$$

where Δz is the phenotypic divergence calculated as the time-standardized phylogentic independent contrasts (PIC) for each node[8,58]. We produced a distribution of values for each node by altering values of G, N_e and t_g between the range defined above. Confidence intervals for the null-hypothesis of drift were generated from simulations based on equation 2[8]. Values that fall within the bounds of the null-distribution are thought to conform to the expectation under genetic drift. Values that fall above or below are thought to be indicative of directional or stabilizing selection, respectively.

References

1. Dobzhansky, T. *Genetics and the Origin of Species*. 11 (Columbia university press, 1982).

2. de Santis, M. D. Misconceptions about historical sciences in evolutionary biology. *Evol. biology* **48**, 94–99, DOI: 10.1007/s11692-020-09526-6 (2021).

3. Stanley, S. M. A theory of evolution above the species level. *Proc. Natl. Acad. Sci. United States Am.* **72**, 646–650, DOI: 10.1073/pnas.72.2.646 (1975).

4. Gould, S. J. *The structure of evolutionary theory* (Harvard University Press, 2002).

5. Uyeda, J. C., Hansen, T. F., Arnold, S. J. & Pienaar, J. The million-year wait for macroevolutionary bursts. *Proc. Natl. Acad. Sci. United States Am.* **108**, 15908–15913, DOI: 10.1073/pnas.1014503108 (2011).

6. Houle, D., Bolstad, G. H., van der Linde, K. & Hansen, T. F. Mutation predicts 40 million years of fly wing evolution. *Nature* **548**, 447–450, DOI: 10.1038/nature23473 (2017).

7. Lynch, M. The rate of morphological evolution in mammals from the standpoint of the neutral expectation. *The Am. Nat.* **136**, 727–741, DOI: 10.1086/285128 (1990).

8. Machado, F. A., Marroig, G. & Hubbe, A. The pre-eminent role of directional selection in generating extreme morphological change in glyptodonts (cingulata; xenarthra). *Proc. Royal Soc. B: Biol. Sci.* **289**, DOI: 10.1098/rspb.2021.2521 (2022).

9. Hlusko, L. J. Integrating the genotype and phenotype in hominid paleontology. *Proc. Natl. Acad. Sci. United States Am.* **101**, 2653–2657, DOI: 10.1073/pnas.0307678101 (2004).

10. Hlusko, L. J., Schmitt, C. A., Monson, T. A., Brasil, M. F. & Mahaney, M. C. The integration of quantitative genetics, paleontology, and neontology reveals genetic underpinnings of primate dental evolution. *Proc. Natl. Acad. Sci. United States Am.* **113**, 9262–9267, DOI: 10.1073/pnas.1605901113 (2016).

11. Mongle, C. *et al.* Developmental processes mediate alignment of the micro- and macroevolution of primate molars. *Evolution* (In revision).

12. Marroig, G. & Cheverud, J. M. Did natural selection or genetic drift produce the cranial diversification of neotropical monkeys? *The Am. Nat.* **163**, 417–428, DOI: 10.1086/381693 (2004).

13. Erwin, D. H. Macroevolution is more than repeated rounds of microevolution. *Evol. & Dev.* **2**, 78–84, DOI: 10.1046/j.1525-142x.2000.00045.x (2000).

14. Hautmann, M. What is macroevolution? *Palaeontology* DOI: 10.1111/pala.12465 (2019).

15. Lande, R. The dynamics of peak shifts and the pattern of morphological evolution. *Paleobiology* **12**, 343–354, DOI: 10.1017/S0094837300003092 (1986).

16. Charlesworth, B., Lande, R. & Slatkin, M. A neo-darwinian commentary on macroevolution. *Evolution* **36**, 474, DOI: 10.2307/2408095 (1982).

17. Arnold, S. J., Pfrender, M. E. & Jones, A. G. The adaptive landscape as a conceptual bridge between micro- and macroevolution. *Genetica* **112-113**, 9–32, DOI: 10.1007/978-94-010-0585-2_2 (2001).

18. Polly, P. D. Developmental dynamics and g-matrices: Can morphometric spaces be used to model phenotypic evolution? *Evol. biology* **35**, 83–96, DOI: 10.1007/s11692-008-9020-0 (2008).

19. Hansen, T. F. Macroevolutionary quantitative genetics? a comment on polly (2008). *Evol. Biol.* **35**, 182–185 (2008).

20. Melo, D., Porto, A., Cheverud, J. M. & Marroig, G. Modularity: genes, development and evolution. *Annu. review ecology, evolution, systematics* **47**, 463–486, DOI: 10.1146/annurev-ecolsys-121415-032409 (2016).

21. Machado, F. A. Selection and constraints in the ecomorphological adaptive evolution of the skull of living canidae (carnivora, mammalia). *The Am. Nat.* **196**, 197–215, DOI: 10.1086/709610 (2020).

22. Laland, K. N. *et al.* The extended evolutionary synthesis: its structure, assumptions and predictions. *Proc. royal society B: biological sciences* **282**, 20151019 (2015).

23. Thompson, D. W. & Thompson, D. W. *On growth and form*, vol. 2 (Cambridge university press Cambridge, 1942).

24. Waddington, C. H. *The strategy of the genes* (Routledge, 1957).

25. Alberch, P., Gould, S. J., Oster, G. F. & Wake, D. B. Size and shape in ontogeny and phylogeny. *Paleobiology* **5**, 296–317, DOI: 10.1017/S0094837300006588 (1979).

26. Alberch, P. Ontogenesis and morphological diversification. *Am. zoologist* **20**, 653–667, DOI: 10.1093/icb/20.4.653 (1980).

27. Love, A. C. Evolutionary morphology, innovation, and the synthesis of evolutionary and developmental biology. *Biol. Philos.* **18**, 309–345 (2003).

28. Cheverud, J. M. Quantitative genetics and developmental constraints on evolution by selection. *J. Theor. Biol.* **110**, 155–171, DOI: 10.1016/s0022-5193(84)80050-8 (1984).

29. Salazar-Ciudad, I. & Jernvall, J. A computational model of teeth and the developmental origins of morphological variation. *Nature* **464**, 583–586, DOI: 10.1038/nature08838 (2010).

30. Linde-Medina, M. & Diogo, R. Do correlation patterns reflect the role of development in morphological evolution? *Evol. biology* **41**, 494–502, DOI: 10.1007/s11692-014-9275-6 (2014).

31. Hether, T. D. & Hohenlohe, P. A. Genetic regulatory network motifs constrain adaptation through curvature in the landscape of mutational (co)variance. *Evolution* **68**, 950–964, DOI: 10.1111/evo.12313 (2014).

32. Pearson, K. & Davin, A. G. On the biometric constants of the human skull. *Biometrika* **16**, 328, DOI: 10.2307/2331712 (1924).

33. Raup, D. M. Geometric analysis of shell coiling: general problems. *J. paleontology* 1178–1190 (1966).

34. Kavanagh, K. D., Evans, A. R. & Jernvall, J. Predicting evolutionary patterns of mammalian teeth from development. *Nature* **449**, 427–432, DOI: 10.1038/nature06153 (2007).

35. Alba, V., Carthew, J. E., Carthew, R. W. & Mani, M. Global constraints within the developmental program of the drosophila wing. *eLife* **10**, DOI: 10.7554/{eLife}.66750 (2021).

36. Hlusko, L. J., Maas, M.-L. & Mahaney, M. C. Statistical genetics of molar cusp patterning in pedigreed baboons: implications for primate dental development and evolution. *J. Exp. Zool. Part B, Mol. Dev. Evol.* **302**, 268–283, DOI: 10.1002/jez.b.21 (2004).

37. Hlusko, L. J., Sage, R. D. & Mahaney, M. C. Modularity in the mammalian dentition: mice and monkeys share a common dental genetic architecture. *J. Exp. Zool. Part B, Mol. Dev. Evol.* **316**, 21–49, DOI: 10.1002/jez.b.21378 (2011).

38. Hardin, A. M. Genetic correlations in the dental dimensions of saguinus fuscicollis. *Am. J. Phys. Anthropol.* **169**, 557–566 (2019).

39. Hardin, A. M. Genetic correlations in the rhesus macaque dentition. *J. Hum. Evol.* **148**, 102873, DOI: 10.1016/j.jhevol.2020.102873 (2020).

40. Pacifici, M. *et al.* Generation length for mammals. *Nat. Conserv.* **5**, 89 (2013).

41. Brevet, M. & Lartillot, N. Reconstructing the history of variation in effective population size along phylogenies. *Genome Biol. Evol.* **13**, DOI: 10.1093/gbe/evab150 (2021).

42. Plavcan, J. M. *Sexual dimorphism in the dentition of extant anthropoid primates*. Ph.D. thesis, Duke University (1990).

43. Godfrey, L. R., Samonds, K. E., Jungers, W. L. & Sutherland, M. R. Teeth, brains, and primate life histories. *Am. J. Phys. Anthropol. The Off. Publ. Am. Assoc. Phys. Anthropol.* **114**, 192–214 (2001).

44. Delson, E., Harcourt-Smith, W. E., Frost, S. R. & Norris, C. A. Databases, data access, and data sharing in paleoanthropology: first steps (2007).

45. Wisniewski, A. L., Lloyd, G. T. & Slater, G. J. Extant species fail to estimate ancestral geographical ranges at older nodes in primate phylogeny. *Proc. Royal Soc. B* **289**, 20212535 (2022).

46. Liam, L. J., Harmon, L. J. & Collar, D. C. Phylogenetic signal, evolutionary process, and rate. *Syst. biology* **57**, 591–601 (2008).

47. Schluter, D. Adaptive radiation along genetic lines of least resistance. *Evolution* **50**, 1766, DOI: 10.2307/2410734 (1996).

48. McGlothlin, J. W. *et al.* Adaptive radiation along a deeply conserved genetic line of least resistance in anolis lizards. *Evol. letters* **2**, 310–322, DOI: 10.1002/evl3.72 (2018).

49. Schroeder, L. & von Cramon-Taubadel, N. The evolution of hominoid cranial diversity: A quantitative genetic approach. *Evolution* **71**, 2634–2649 (2017).

50. Ackermann, R. R. & Cheverud, J. M. Discerning evolutionary processes in patterns of tamarin (genus saguinus) craniofacial variation. *Am. J. Phys. Anthropol.* **117**, 260–271, DOI: 10.1002/ajpa.10038 (2002).

51. Hansen, T. F. Stabilizing selection and the comparative analysis of adaptation. *Evolution* **51**, 1341, DOI: 10.2307/2411186 (1997).

52. Mosimann, J. E. Size allometry: Size and shape variables with characterizations of the lognormal and generalized gamma distributions. *J. Am. Stat. Assoc.* **65**, 930, DOI: 10.2307/2284599 (1970).

53. Jolicoeur, P. The multivariate generalization of the allometry equation. *Biometrics* **19**, 497–499 (1963).

54. Gould, S. J. *Ontogeny and phylogeny* (Belknap press, 1977).

55. Weaver, T. D., Roseman, C. C. & Stringer, C. B. Close correspondence between quantitative- and molecular-genetic divergence times for neandertals and modern humans. *Proc. Natl. Acad. Sci. United States Am.* **105**, 4645–4649, DOI: 10.1073/pnas.0709079105 (2008).

56. Gómez-Robles, A. Dental evolutionary rates and its implications for the neanderthal-modern human divergence. *Sci. Adv.* **5**, eaaw1268, DOI: 10.1126/sciadv.aaw1268 (2019).

57. Monson, T. A., Fecker, D. & Scherrer, M. Neutral evolution of human enamel-dentine junction morphology. *Proc. Natl. Acad. Sci. United States Am.* **117**, 26183–26189, DOI: 10.1073/pnas.2008037117 (2020).

58. Felsenstein, J. Phylogenies and quantitative characters. *Annu. review ecology systematics* **19**, 445–471, DOI: 10.1146/annurev.es.19.110188.002305 (1988).

59. Melo, D. & Marroig, G. Directional selection can drive the evolution of modularity in complex traits. *Proc. Natl. Acad. Sci. United States Am.* **112**, 470–475, DOI: 10.1073/pnas.1322632112 (2015).

60. Jones, A. G., Bürger, R., Arnold, S. J., Hohenlohe, P. A. & Uyeda, J. C. The effects of stochastic and episodic movement of the optimum on the evolution of the g-matrix and the response of the trait mean to selection. *J. Evol. Biol.* **25**, 2210–2231, DOI: 10.1111/j.1420-9101.2012.02598.x (2012).

61. Riedl, R. *Order in living organisms: a systems analysis of evolution* (John Wiley & Sons, 1978).

62. Watson, R. A., Wagner, G. P., Pavlicev, M., Weinreich, D. M. & Mills, R. The evolution of phenotypic correlations and "developmental memory". *Evolution* **68**, 1124–1138, DOI: 10.1111/evo.12337 (2014).

63. Cheverud, J. The evolution of genetic correlation and developmental constraints. In *Population genetics and evolution*, 94–101 (Springer, 1988).

64. Polly, P. D. Development with a bite. *Nature* **449**, 413–414 (2007).

65. Halliday, T. J. D. & Goswami, A. Testing the inhibitory cascade model in mesozoic and cenozoic mammaliaforms. *BMC Evol. Biol.* **13**, 79, DOI: 10.1186/1471-2148-13-79 (2013).

66. Monson, T. A. *et al.* Evidence of strong stabilizing effects on the evolution of boreoeutherian (mammalia) dental proportions. *Ecol. Evol.* DOI: 10.1002/ece3.5309 (2019).

67. Asahara, M. Unique inhibitory cascade pattern of molars in canids contributing to their potential to evolutionary plasticity of diet. *Ecol. Evol.* **3**, 278–285, DOI: 10.1002/ece3.436 (2013).

68. Varela, L., Tambusso, P. S. & Fariña, R. A. Unexpected inhibitory cascade in the molariforms of sloths (folivora, xenarthra): a case study in xenarthrans honouring gerhard storch's open-mindedness. *Frühförderung interdisziplinär* **76**, 1–16, DOI: 10.37520/fi.2020.002 (2020).

69. Sadier, A. *et al.* Growth rate as a modulator of tooth patterning during adaptive radiations. *BioRxiv* DOI: 10.1101/2021.12.05.471324 (2021).

70. Glowacka, H. & Schwartz, G. T. A biomechanical perspective on molar emergence and primate life history. *Sci. Adv.* **7**, eabj0335, DOI: 10.1126/sciadv.abj0335 (2021).

71. Monson, T. A. *et al.* Keeping 21st century paleontology grounded: Quantitative genetic analyses and ancestral state reconstruction re-emphasize the essentiality of fossils. *Biology* **11**, DOI: 10.3390/biology11081218 (2022).

72. Riedl, R. A systems-analytical approach to macro-evolutionary phenomena. *The Q. Rev. Biol.* **52**, 351–370 (1977).

73. Houle, D. & Rossoni, D. M. Complexity, evolvability, and the process of adaptation. *Annu. Rev. Ecol. Evol. Syst.* **53** (2022).

74. Olson, E. C. & Miller, R. L. *Morphological integration* (University of Chicago Press, Chicago and Longon, 1958).

75. Kavanagh, K. D. *et al.* Developmental bias in the evolution of phalanges. *Proc. Natl. Acad. Sci. United States Am.* **110**, 18190–18195, DOI: 10.1073/pnas.1315213110 (2013).

76. Carraco, G., Martins-Jesus, A. P. & Andrade, R. P. The vertebrate embryo clock: Common players dancing to a different beat. *Front. cell developmental biology* **10**, 944016, DOI: 10.3389/fcell.2022.944016 (2022).

77. Schroeder, L., Roseman, C. C., Cheverud, J. M. & Ackermann, R. R. Characterizing the evolutionary path(s) to early homo. *Plos One* **9**, e114307, DOI: 10.1371/journal.pone.0114307 (2014).

78. Schroeder, L., Elton, S. & Ackermann, R. R. Skull variation in afro-eurasian monkeys results from both adaptive and non-adaptive evolutionary processes. *Sci. Reports* **12**, 12516, DOI: 10.1038/s41598-022-16734-x (2022).

79. Bookstein, F. L. Measurement, explanation, and biology: Lessons from a long century. *Biol. theory* **4**, 6–20, DOI: 10.1162/biot.2009.4.1.6 (2009).

80. Houle, D., Pélabon, C., Wagner, G. P. & Hansen, T. F. Measurement and meaning in biology. *The Q. Rev. Biol.* **86**, 3–34, DOI: 10.1086/658408 (2011).

81. Machado, F. A., Hubbe, A., Melo, D., Porto, A. & Marroig, G. Measuring the magnitude of morphological integration: The effect of differences in morphometric representations and the inclusion of size. *Evolution* **73**, 2518–2528, DOI: 10.1111/evo.13864 (2019).

82. Mitteroecker, P. The developmental basis of variational modularity: Insights from quantitative genetics, morphometrics, and developmental biology. *Evol. biology* **36**, 377–385, DOI: 10.1007/s11692-009-9075-6 (2009).

83. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat. methods* **9**, 676–682 (2012).

84. Scrucca, L., Fop, M., Murphy, T. B. & Raftery, A. E. mclust 5: clustering, classification and density estimation using Gaussian finite mixture models. *The R J.* **8**, 289–317 (2016).

85. Mitov, V., Bartoszek, K. & Stadler, T. Automatic generation of evolutionary hypotheses using mixed gaussian phylogenetic models. *Proc. Natl. Acad. Sci.* **116**, 16921–16926 (2019).

86. Lande, R. Natural selection and random genetic drift in phenotypic evolution. *Evolution* **30**, 314, DOI: 10.2307/2407703 (1976).

87. Lande, R. Quantitative genetic analysis of multivariate evolution, applied to brain:body size allometry. *Evolution* **33**, 402–416, DOI: 10.1111/j.1558-5646.1979.tb04694.x (1979).

88. Dennis, B., Ponciano, J. M., Taper, M. L. & Lele, S. R. Errors in statistical inference under model misspecification: evidence, hypothesis testing, and AIC. *Front. ecology evolution* **7**, DOI: 10.3389/fevo.2019.00372 (2019).

89. Schluter, D., Price, T., Mooers, A. Ø. & Ludwig, D. Likelihood of ancestor states in adaptive radiation. *Evolution* **51**, 1699–1711, DOI: 10.1111/j.1558-5646.1997.tb05095.x (1997).

90. Hansen, T. F. & Martins, E. P. Translating between microevolutionary process and macroevolutionary patterns: the correlation structure of interspecific data. *Evolution* **50**, 1404–1417, DOI: 10.1111/j.1558-5646.1996.tb03914.x (1996).

91. Marroig, G. & Cheverud, J. Size as a line of least resistance II: direct selection on size or correlated response due to constraints? *Evolution* **64**, 1470–1488, DOI: 10.1111/j.1558-5646.2009.00920.x (2010).

Supplementary Materials

Contents

A Morphometrics

A.1 Morphospaces

We used three morphospaces to test our hypothesis that evo-devo-informed quantification would provide a better bridge between micro and macroevolution (Fig S1). The first is based on the linear distances taken directly from the teeth, constituting a 6-dimensional space ("distance space"). The second morphospace is based on the occlusal areas of each molar, which were calculated using a rectangular approximation[64,65], producing a 3-dimensional space ("area space"). These two spaces are considered naïve because they make no assumptions about underlying developmental processes. As an evo-devo inspire space, we used the ICM description of the molar development[34], and constructed a morphospace based on the relation between the relative occlusal area of m2 and m3 in relation to m1 (m2/m1 and m3/m1, respectively), resulting in a 2-dimensional space ("ICM space"). Variables were log-transformed in both the trait and area datasets, but not in the ICM space.

Figure S1. Schematic representation of variables used to construct morphospaces. The trait-space was built on the mesiodistal length (MD, vertical) and buccolingual breadth (BL, horizontal) taken from each molar. The area-space was built by estimating the occlusal areas of each molar as the $A = MDxBL$. The ICM-space was built by calculating the relative area size for m2 and m3 in relation to m1.

To evaluate morphospace patchiness we performed a clustering based on parameterized finite Gaussian mixture models[84]. This method test for a series of nested models, where groups are modeled according to different covariance structures that can be either spherical (all variances equal, no covariances) or ellipsoidal (different variances and non-zero covariances), and can share or not aspects of their covariance matrices, like size, shape or orientation. The permutation of these aspects produces a total of 14 total models, which are fitted with an Expectation-Maximization algorithm, and compared through BIC[84]. In the present case, the best models were the ones in which covariance matrices were ellipsoidal with the same shape, but with either the same orientation and different volumes (VEE) or the same volume and different orientations (EEV). Specifically, both naïve spaces showed a preference to the VVE model, whereas the ICM space showed a preference for the EEV model (Fig. S2). Furthermore, both naïve spaces showed a tendency for finding more groups, with the maximum BIC associated with eight clusters for linear variables and 5 for areas, while the ICM space preferred only two clusters (Fig. S2).

Figure S2. BIC values for the Gaussian mixture models for the linear distances, areas and ICM variables. EEV- Elipsoidal model with the same shape, same volume and different orientations VEE- Elipsoidal model with the same shape, different volumes volume and same orientation.

To visualize these results, we performed a Principal Component Analysis (PCA) on the variance-covariance matrix for the full sample on each space. We then inspected the two leading PCs to evaluate the space discontinuity. Because the ICM space is only made up of two variables, the leading two PCs represent all variation on the sample. The inspection of the leading PCs, and the distribution of the groups found through the clustering analysis show that the naïve spaces are more patchy than the ICM space (fig. 2).

To visualize these results, we performed a Principal Component Analysis (PCA) on the variance-covariance matrix for the full sample on each space. We then inspected the two leading PCs to evaluate the space discontinuity. Because the ICM space is only made up of two variables, the leading two PCs represent all variation on the sample. The inspection of the leading PCs, and the distribution of the groups found through the clustering analysis show that the naïve spaces are more patchy than the ICM space (fig. 2).

A.2 G-matrix

To model the evolution of these traits under quantitative genetics models, we need an estimate of the additive genetic covariance matrix **G** for our molar traits. One common practice in comparative analysis is to use the pooled intraspecific phenotypic covariance matrix **P** as an approximation of **G**[8,12,21,91]. This is justified on the bases of the high similarity between **P** and **G** for morphological traits[11,91]. However, given that **P**s contain also non-genetic information, we also used **G** estimated from a pedigreed population of baboons as a direct model of the Primate **G**[36].

Because the *G* provided by[36] was estimated for the raw, untransformed BL and MD measures, we had to transform it in order to match our three morphospaces. We did this through a Monte-Carlo approach, in which we used the published *G* and

population means to generate $n = 300$ samples of additive genetic effects. This number of samples was chosen to be compatible with effective sample sizes for the pedigreed population, and not over-represent the accuracy of the estimate[36]. For the linear distance space, we took the generated genetic effects and log-transformed them. For the area space, we log-transformed the product of the effects for each tooth BL and MD. For the ICM space we took the product of the effects for each tooth BL and MD and divided the ones relative to m2 and m3 by m1. We then calculated the maximum-likelihood covariance matrix for the resulting effects for each space. This procedure was done 10,000 times, and the mean covariance matrix was taken as a point estimate of G for the lower-dimensionality morphospaces (area and ICM spaces).

To test the validity of this approach we compared the simulation results to the analytic approximation for covariance matrices of products of random variables. Specifically, we investigated if the covariance matrix obtained for the product of the additive effect for each tooth BL and MD (area space) are similar to what would be expected analytically as

$$\sigma_{xy}^2 = \mu_x^2 \sigma_y^2 + \mu_y^2 \sigma_x^2 + (\sigma_{xy})^2 + 2\mu_x \mu_y \sigma_{xy} + \sigma_x^2 \sigma_y^2 \tag{S1}$$

$$\sigma_{xy,uv}^2 = \mu_x \mu_u \sigma_{y,v} + \mu_x \mu_v \sigma_{y,u} + \mu_y \mu_u \sigma_{x,v} + \mu_y \mu_v \sigma_{y,u} + \sigma_{xu} \sigma_{yv} \sigma_{yu} \tag{S2}$$

where μ are means, σ^2 variances and σ are covariances for the random variables x, y, u and v[91]. Variable pairs x:y and u:v are BL and MD variable pairs for each tooth. From these equations we were able to construct the covariance matrix for the area space on a cm^2 scale. This procedure was done only for this non-log area space as a proof-of-concept and to limit the number of assumptions necessary to derived log-scale and ratio spaces.

To compare analytical and Monte-Carlo estimates of G we first mean-scaled both matrices as follows:

$$\Sigma_\mu = \Sigma \oslash \bar{z}\bar{z}^t \tag{S3}$$

were Σ is the original covariance matrix, $\oslash$ is the element-wise product and $\bar{z}$ is a vector of means[19]. We then calculated the absolute differences from each Monte-Carlo sample of these standardized matrices. Differences in the variances on these matrices are equal to the difference in coefficient of variation between matrices, and thus provide a dimensionless scale of comparison.

The results show that matrices are extremely similar, with a slight bias for covariances being higher on the Monte Carlo samples (fig. S3). Despite this, all absolute differences are very small (< 0.002), suggesting that differences are negligible. A matrix correlation analysis reinforced this interpretation, showing values > 0.97 for all samples. Together, these results suggest that the Monte Carlo estimates are a reliable approximation for the covariance on lower dimensionalities.

Figure S3. Differences between the Monte Carlo sampling approach for generating covariances for areas and the analytical approximation. Horizontal lines within violins highlights the 95% interval for each matrix cell.

B Model comparison

We performed model comparison under the maximum likelihood framework of the Phylogenetic mixed Gaussian models[85],[92]. This framework has some desirable features. First, it incorporates both measurement errors and missing data during the model fitting procedure. The latter is especially important in our case due to the abundance of fossil taxa in our sample and the presence of species lacking m3s, such as callitrichines and the fossil *Xenothrix mcgregori*. Second, it takes into account process heterogeneity along the tree, allowing regimes to be under different kinds of models and parameter combinations. Lastly, the parameters of the fitted model can be interpreted under quantitative genetics theory in terms of genetic drift and stabilizing selection[51].

To test our hypothesis, we fit global models that included some microevolutionary assumptions (κ-models) and some that did not. Evidence supporting κ-models could then be seen as evidence for a microevolutionary interpretation of the data. For each morphometric space, we fitted regular BM and OU models, as well as two κ versions of these models. For these models, the rate matrix Σ was set to be proportional to either P or G. We also performed mixed model heuristic searches, which try to find the best regime combination for a given tree, allowing for regimes to be under different model types (BM or OU). For each space, we conducted 10 searches and recorded the result with the lowest BIC. Lastly, in addition to the heuristic result, we tested mixed models with pre-determinate regime shifts based on Fig. 3 which is the best result for the area morphospace. This regime was chosen as a common point of reference to compare the effect of model and parameter heterogeneity on the fitting process.

B.1 Distance space

For the distance space only 6 out of 10 heuristic searches converged. Nevertheless, they all produced better BIC values than any global model (table S1). The best model (Search 5) was a multi-rate BM model, with two regimes, one for Simiiformes and one for the remaining of the tree (Fig.S4A). This model deviates from the best model chosen to represent the rate heterogeneity on the other spaces by essentially fusing the ancestral and the Strepsirrhini regimes into one (Fig. 3). In terms of model parameters, this model performed similarly to the tree-regime one in the sense that the ancestral regime had higher rates of evolution than the Simiiformes one (see below fig 5).

Despite the best model being different than the one described in the main text, Search 4 produced a regime combination that was essentially identical to the one for areas (Fig. S4B). Even though the BIC was worse than the best model, this run performed better than any of the global models and is therefore adequate to highlight how this morphospace privileges more heterogeneous models instead of global ones.

Figure S4. Regimes for different runs of the heuristic search. A- Best model (Search 5). B- Model compatible with the best model for areas (Fig. 3)

B.2 Area space

For the area space, 9 out of the 10 heuristic searches converged. The best mixed model (Search 7) and the best overall model was a multi-rate BM model with three regimes, as described in the main text (Fig. 3).

B.3 ICM space

For the ICM space, all 10 heuristic searches converged. From these, four runs produced the same result as the global OU model (table S3), suggesting an absence of model shifts for this space. The global fits show that both OU and BM models are equally good fits to the data and that the addition of the microevolutionary assumption of proportionality between Σ and G for the global OU model produces a slightly superior model ($OU_{\Sigma\propto G}$). Although all three models seem to provide a good fit to the data, the investigation of confidence intervals for parameters of the OU models shows that the confidence interval of the off-diagonal element of H overlaps with zero (tables S6,S7), suggesting that this parameter can be excluded from the model (see below).

A model that omits the off-diagonal elements of H is implemented as a default model in the PCMfit package[85]. We chose not to include those models initially because our data is rich enough to estimate parameters from even fairly complex models. Instead, we chose to fit the most complex model and perform *post-hoc* model simplification. Using a κ-OU model with a diagonal H produces a model with vastly superior BIC ($OU_{\Sigma\propto G}^{d}$). The removal of the off-diagonal elements of H did not change significantly the parameter estimates (table S9,S8). Lastly, although OU^d and $OU_{\Sigma\propto G}^{d}$ had similar BICs, the confidence interval for all common parameters showed great overlap, suggesting that both models are equivalent. Furthermore, the $OU_{\Sigma\propto G}^{d}$ model had tighter confidence intervals, fewer parameters and a slightly larger BIC. For these reasons, we chose this as the best model for our data.

Table S1. Comparison of models of the linear distance morphospace fit through Maximum Likelihood.

Model [a]	N_p [b]	logLik [c]	BIC [d]
BM	27	2585.03	-5112.72
OU	54	2684.57	-5247.16
$BM_{\Sigma\propto P}$	7	2108.57	-4202.91
$OU_{\Sigma\propto P}$	34	2174.45	-4275.55
$BM_{\Sigma\propto G}$	7	1480.63	-2947.02
$OU_{\Sigma\propto G}$	34	1510.57	-2947.79
Search 1	78	2788.91	-5391.08
Search 2	51	2796.77	-5479.14
Search 3	97	2868.67	-5493.57
Search 4	74	2823.03	-5470.65
Search 5	51	2878.96	**-5643.53**
Search 6	78	2820.77	-5454.80

[a]Model type; either global BM or OU (full or κ models) or a search run of mixed model.

[b]Number of model parameters.

[c]Log-likelihood of the model.

[d]Bayesian Information Criterion used for model comparison. Bold represents the best models, while underline represents the ones with BIC 2 units away from the best model.

Table S2. Comparison of models of the area morphospace fit through Maximum Likelihood.

Model [a]	N_p [b]	logLik [c]	BIC [d]
BM	9	415.80	-813.21
OU	18	413.98	-790.47
$BM_{\Sigma \propto P}$	4	92.39	-176.70
$OU_{\Sigma \propto P}$	13	275.16	-523.54
$BM_{\Sigma \propto G}$	4	74.41	-140.74
$OU_{\Sigma \propto G}$	13	258.02	-489.26
Search 1	10	413.61	-806.75
Search 2	10	414.98	-809.48
Search 3	19	420.25	-800.85
Search 4	10	415.85	-811.23
Search 5	18	434.43	-831.37
Search 6	26	469.34	-883.58
Search 7	26	471.42	**-887.74**
Search 8	10	415.17	-809.87
Search 9	27	448.71	-840.08

[a]Model type; either global BM or OU (full or κ models) or a search run of mixed model.
[b]Number of model parameters.
[c]Log-likelihood of the model.
[d]Bayesian Information Criterion used for model comparison. Bold represents the best models, while underline represents the ones with BIC 2 units away from the best model.

Table S3. Comparison of models of the ICM morphospace fit through Maximum Likelihood.

Model [a]	N_p [b]	logLik [c]	BIC [d]
BM	5	589.16	-1147.46
OU	10	604.53	-1147.33
OU^d	9	604.56	<u>-1153.55</u>
$BM_{\Sigma \propto P}$	3	538.31	-1058.10
$OU_{\Sigma \propto P}$	8	584.68	-1119.98
$BM_{\Sigma \propto G}$	3	575.69	-1132.86
$OU_{\Sigma \propto G}$	8	598.96	-1148.53
$OU^d_{\Sigma \propto G}$	7	598.54	**-1153.86**
Three-regime OU	26	597.35	-1034.18
Search 1	10	604.56	-1147.37
Search 2	15	618.60	-1144.59
Search 3	15	618.56	-1144.52
Search 4	15	617.78	-1142.95
Search 5	10	604.55	-1147.37
Search 6	15	617.80	-1142.99
Search 7	10	604.56	-1147.37
Search 8	6	589.16	-1141.28
Search 9	10	604.55	-1147.36
Search 10	15	617.80	-1143.00

[a]Model type; either global BM or OU (full or κ models) or a search run of mixed model.
[b]Number of model parameters.
[c]Log-likelihood of the model.
[d]Bayesian Information Criterion used for model comparison. Bold represents the best models, while underline represents the ones with BIC 2 units away from the best model.

C Confidence Intervals

Confidence intervals for model parameters and derived statistics were obtained by exploring the likelihood surface and getting parameter combinations from models that were 2 log-likelihood units away from the peak. This was done using the "dentist" package (https://github.com/bomeara/dentist), which allows the exploration of the likelihood surface by "denting" of the surface- setting the region around the peak to have extremely low likelihoods and sampling on the resulting contour around the dented peak. We used 10,000 steps to sample the likelihood surface and obtain confidence intervals simultaneously for all parameters for models fit on the ICM morphospace. If maximum and minimum values sampled by the "denting" approach matched the confidence intervals, then the sampling did not encompass the true confidence interval, and the sampling range had to be expanded. Below we report both the confidence interval as well the extreme values examined to show that adequate confidence intervals are being reported.

For the OU and $\text{OU}_{\Sigma \propto G}$ models, the off-diagonal elements of the H matrix ($H_{1,2}$) have confidence intervals that overlap with 0, suggesting that this parameter can be omitted from the model (table S6, S7). The removal of the $H_{1,2}$ from the model did not change substantially parameter estimates (table S8, S9), nor did it reduce the likelihoods of the models, improving BIC scores (table S3). All models showed a great overlap in the common parameters (table S6–S9). This is also true for the reconstructed Σ matrix for the *kappa* models. Specifically, for $\text{OU}^d_{\Sigma \propto G}$, Σ confidence intervals are $\Sigma_{1,1} = 0.0243 - 0.0276$, $\Sigma_{1,2} = 0.0324 - 0.0368$ and $\Sigma_{2,2} = 0.0666 - 0.0756$, placing them within the expected for the $\text{OU}_{\Sigma \propto G}$ model.

Table S4. Confidence interval for the BM model. Lower and upper CI- parameters that are 2 log-likelihood units away from the ML estimate. Lowest and Highest examined values- Extreme values examined by the "denting" approach.

	best	lower CI	upper CI	lowest examined	highest examined
$X0_1{}^a$	1.086	0.993	1.184	0.801	1.200
$X0_2{}^a$	1.184	1.003	1.364	0.801	1.599
$\Sigma_{1,1}{}^b$	0.018	0.016	0.020	0.010	0.499
$\Sigma_{1,2}{}^b$	0.030	0.026	0.032	0.010	0.498
$\Sigma_{2,2}{}^b$	0.071	0.065	0.077	0.010	0.497

[a]Ancestral state at the root
[b]Entries of the rate matrix.

Table S5. Confidence interval for the $\text{BM}_{\Sigma \propto G}$ model. Lower and upper CI- parameters that are 2 log-likelihood units away from the ML estimate. Lowest and Highest examined values- Extreme values examined by the "denting" approach.

	best	lower CI	upper CI	lowest examined	highest examined
$X0_1{}^a$	1.090	0.983	1.195	0.800	1.200
$X0_2{}^b$	1.182	1.005	1.348	0.800	1.593
$\kappa{}^b$	0.361	0.317	0.413	0.102	0.899

[a]Ancestral state at the root
[b]Proportionality constant between the target matrix (**G**-matrix) and Σ.

Table S6. Confidence interval for the OU model. Lower and upper CI- parameters that are 2 log-likelihood units away from the ML estimate. Lowest and Highest examined values- Extreme values examined by the "denting" approach.

	best	lower CI	upper CI	lowest examined	highest examined
$X0_1{}^a$	1.132	1.008	1.200	0.801	1.200
$X0_2{}^a$	1.306	1.075	1.466	0.804	1.597
$H_{1,1}{}^b$	0.034	0.025	0.043	0.010	0.498
$H_{1,2}{}^b$	-0.214	-0.485	0.473	-0.490	0.498
$H_{2,2}{}^b$	0.031	0.021	0.040	0.010	0.498
$\theta_1{}^c$	1.064	1.012	1.115	0.808	1.198
$\theta_2{}^c$	1.066	0.971	1.175	0.801	1.593
$\Sigma_{1,1}{}^d$	0.022	0.020	0.025	0.010	0.495
$\Sigma_{1,2}{}^d$	0.034	0.030	0.036	0.010	0.500
$\Sigma_{2,2}{}^d$	0.080	0.074	0.086	0.010	0.498

[a]Ancestral state at the root
[b]Entries of the **H**-matrix
[c]Multivariate optima
[d]Entries of the rate matrix

Table S7. Confidence interval for the $OU_{\Sigma \propto G}$ model. Lower and upper CI- parameters that are 2 log-likelihood units away from the ML estimate. Lowest and Highest examined values- Extreme values examined by the "denting" approach.

	best	lower CI	upper CI	lowest examined	highest examined
$X0_1{}^a$	1.097	0.966	1.192	0.800	1.199
$X0_2{}^a$	1.232	1.031	1.376	0.802	1.599
$\kappa{}^b$	0.486	0.416	0.553	0.101	0.894
$H_{1,1}{}^c$	0.049	0.037	0.057	0.010	0.499
$H_{1,2}{}^c$	0.226	-0.352	0.469	-0.497	0.500
$H_{2,2}{}^c$	0.024	0.013	0.035	0.010	0.499
$\theta_1{}^d$	1.066	1.027	1.106	0.800	1.198
$\theta_2{}^d$	1.058	0.885	1.175	0.802	1.597

[a]Ancestral state at the root
[b]Proportionality constant between the target matrix (**G**-matrix) and Σ.
[c]Entries of the **H**-matrix
[d]Multivariate optima

Table S8. Confidence interval for the OU^d model. Lower and upper CI- parameters that are 2 log-likelihood units away from the ML estimate. Lowest and Highest examined values- Extreme values examined by the "denting" approach.

	best	lower CI	upper CI	lowest examined	highest examined
$X0_1{}^a$	1.102	1.012	1.192	0.801	1.199
$X0_2{}^a$	1.248	1.068	1.428	0.800	1.592
$H_{1,1}{}^b$	0.036	0.025	0.043	0.010	0.499
$H_{2,2}{}^b$	0.031	0.019	0.039	0.010	0.499
$\theta_1{}^c$	1.066	1.016	1.112	0.801	1.200
$\theta_2{}^c$	1.073	0.969	1.211	0.808	1.585
$\Sigma_{1,1}{}^d$	0.022	0.020	0.025	0.010	0.498
$\Sigma_{1,2}{}^c$	0.033	0.030	0.036	0.010	0.499
$\Sigma_{2,2}{}^c$	0.079	0.073	0.084	0.010	0.499

[a]Ancestral state at the root
[b]Entries of the **H**-matrix
[c]Multivariate optima
[d]Entries of the rate matrix

Table S9. Confidence interval for the $OU^d_{\Sigma \propto G}$ model. Lower and upper CI- parameters that are 2 log-likelihood units away from the ML estimate. Lowest and Highest examined values- Extreme values examined by the "denting" approach.

	best	lower CI	upper CI	lowest examined	highest examined
$X0_1{}^a$	1.107	0.963	1.190	0.800	1.200
$X0_2{}^a$	1.232	1.036	1.393	0.800	1.598
κ^b	0.484	0.416	0.536	0.101	0.900
$H_{1,1}{}^c$	0.044	0.034	0.050	0.010	0.498
$H_{1,2}{}^c$	0.025	0.013	0.037	0.010	0.495
$\theta_1{}^d$	1.066	1.029	1.109	0.800	1.200
$\theta_2{}^d$	1.063	0.917	1.209	0.802	1.594

[a]Ancestral state at the root
[b]Proportionality constant between the target matrix (**G**-matrix) and Σ.
[c]Entries of the **H**-matrix
[d]Multivariate optima

D Phylogenetic Half-lives

The investigation of the adaptive landscape implied by the best model shows a corridor-like topography (Fig. 4), suggesting that selection is relaxed along the activation-inhibition axis and intensified against deviations from the ICM. To further verify this without assuming a microevolutionary model, we calculated the phylogenetic half-lives ($t_{1/2}$) along the activation-inhibition gradient and the deviations from the ICM. $t_{1/2}$ measures the time it takes for the phenotype to move halfway in the direction of the optimum and is calculated as

$$t_{1/2} = ln(2)/\alpha \tag{S4}$$

with α being the rate of adaptation for each trait given by the diagonal of the H matrix[51]. Because H is given as a function of the original traits (m2/m1 and m3/m1) we have to perform a rotation of the original matrix into the eigenvectors of the adaptive landscape as

$$H^r = V^t H V \tag{S5}$$

were V are the eigenvectors of Ω (see equation 3). $t_{1/2}$ were calculated integrating over the confidence limits of parameters. In addition to the rotated space, we also calculated $t_{1/2}$ on the original space. This was done to evaluate an additional hypothesis of the ICM process that m3s, because they develop last, are under less intense stabilizing selection[11].

The results for molar ratios are in line with this prediction, as the half-life for m3/m1 is considerably larger than for m2/m1, suggesting a stronger selection on the latter than in the former (fig S5). For the components of the ICM, results are compatible with the expected for a corridor-like adaptive landscape, with half-lives along the activation-inhibition gradient being higher than on the one for deviations from the ICM (fig S5). This suggests that selection against deviations from ICM is far stronger than the ones along the activation-inhibition gradient, allowing the near-neutral evolution within the adaptive corridor. An inspection of the evolutionary trait-grams of these variables reinforces this idea, as deviations from the ICM are more tightly confined around the optima, and evolution along the activation-inhibition gradient seems greater in amplitude (fig S6).

Figure S5. Phylogenetic half life for ICM ratios (m2/m1 and m3/m1) and components of the ICM model (Activation-Inhibition gradient and deviations from the ICM).

E Disparity and phylogenetic signal

To illustrate the effect of stabilizing selection on both overall disparities in the sample and phylogenetic signal, we employed a simulation approach. We generated tip data using the phylogenetic tree and the best evolutionary model in two situations. In one, we used the whole model to produce data under an OU model; in another, we set the H matrix to be 0, producing a BM model with the same rate parameters as the OU model. For each model, we generated 1,000 datasets, and for each of these datasets, we extracted the overall disparity and the phylogenetic signal. The disparity was calculated simply as the sum of variances in the tip data, as a measure of overall morphospace occupancy[93]. For Phylogenetic signal, we employed the multivariate version of Blomberg's K[94,95]. Results from the simulations show the expected pattern, with the BM model showing higher disparity and phylogenetic signal than the OU model (Fig. S7). Because these simulations are based on our preferred microevolutionary model ($OU^d_{\Sigma \propto G}$), the BM represents what would take place under a pure genetic drift process, while the OU model represents the action of stabilizing selection in modulating the action of drift. So, despite the fact that most of the divergence within the group is compatible with drift (Figure 5), drift alone would produce a wider range of phenotypic values (Fig. S7A, left panel), thus requiring the action of stabilizing selection to not only constrain the total amount of divergence, but also to shape the pattern distribution of phenotypes (Fig. S7A, right panel).

Figure S6. Traitgrams of the components of the ICM for primates. Black lines represents the phylogeny mapped to measured trait values (black points) and golden lines and dots represents each trait optimum.

F Regime-specific disparity

To compare the disparity-generating potential of different regimes of the mixed-model depicted in Fig. 3, we employed a simulation approach. Specifically, we took each model for each regime and simulated phenotypic evolution according to that model on a star-phylogeny of equal size to the full phylogeny. This was done to standardize differences in tree structure, species sample size and model differences (BM or OU) between regimes. Simulations were performed 100 times, and for each run we computed the disparity of the simulated tip data. This was done for the three-regime model for each morphospace. Higher and lower values of disparity indicate a less or more constrained evolution, respectively. Results show that, for all spaces, the ancestral regime is less constrained than the Strepsirrhini and Simiiform regimes (Fig. S8). For areas and distances, the difference between the ancestral and derived regimes disparity is greater than for the ICM morphospace, with Simiiform showing a higher disparity than Strepsirrhini.

Figure S7. Simulated disparity (A) and phylogenetic signal (B) by assuming the best model (OU) or a brownian motion (BM) model with the same rate parameters as the best OU model. Ellipses represent the covariance matrix of the simulated tip values, and thus do not represent any evolutionary parameter (*e.g.* Σ, H, Ω), but the empirical phenotypic distributions. Dots are observed species averages.

G Supplementary references

References

91. Roseman, C. C. & Delezene, L. K. The inhibitory cascade model is not a good predictor of molar size covariation. *Evol. Biol.* **46**, 229–238 (2019).

92. Mitov, V., Bartoszek, K., Asimomitis, G. & Stadler, T. Fast likelihood calculation for multivariate gaussian phylogenetic models with shifts. *Theor. Popul. Biol.* **131**, 66–78, DOI: 10.1016/j.tpb.2019.11.005 (2020).

93. Hopkins, M. J. & Gerber, S. Morphological disparity. *Evol. developmental biology: A reference guide* 965–976 (2021).

94. Blomberg, S. P., Garland, T. & Ives, A. R. Testing for phylogenetic signal in comparative data: behavioral traits are more labile. *Evolution* **57**, 717–745, DOI: 10.1111/j.0014-3820.2003.tb00285.x (2003).

95. Adams, D. C. A generalized k statistic for estimating phylogenetic signal from shape and other high-dimensional multivariate data. *Syst. Biol.* **63**, 685–697, DOI: 10.1093/sysbio/syu030 (2014).

Figure S8. Simulated regime-specific disparities for Three-regime model on morphospace.

H Data source references

References

97. Laura Abbazzi, Flaviano Fanfani, Marco P Ferretti, Lorenzo Rook, L Cattani, Federico Masini, Francesco Mallegni, Fabio Negrino, and Carlo Tozzi. New human remains of archaic homo sapiens and lower palaeolithic industries from visogliano (duino aurisina, trieste, italy). *Journal of Archaeological Science*, 27(12):1173–1186, 2000.

98. David M Alba. Fossil apes from the vallès-penedès basin. *Evolutionary Anthropology: Issues, News, and Reviews*, 21(6):254–269, 2012.

99. David M Alba, Sergio Almécija, Isaac Casanovas-Vilar, Josep M Méndez, and Salvador Moyà-Solà. A partial skeleton of the fossil great ape hispanopithecus laietanus from can feu and the mosaic evolution of crown-hominoid positional behaviors. *PLoS One*, 7(6):e39617, 2012.

100. David M Alba, Isaac Casanovas-Vilar, Sergio Almécija, Josep M Robles, Júlia Arias-Martorell, and Salvador Moyà-Solà. New dental remains of hispanopithecus laietanus (primates: Hominidae) from can llobateres 1 and the taxonomy of late miocene hominoids from the vallès-penedès basin (ne iberian peninsula). *Journal of Human Evolution*, 63(1):231–246, 2012.

101. David M Alba, Josep Fortuny, Miriam Perez de Los Rios, Clément Zanolli, Sergio Almécija, Isaac Casanovas-Vilar, Josep M Robles, and Salvador Moyà-Solà. New dental remains of anoiapithecus and the first appearance datum of hominoids in the iberian peninsula. *Journal of Human Evolution*, 65(5):573–584, 2013.

102. David M Alba, Salvador Moyà-Solà, Assumpció Malgosa, Isaac Casanovas-Vilar, Josep M Robles, Sergio Almécija, Jordi Galindo, Cheyenn Rotgers, and Juan Vicente Bertó Mengual. A new species of pliopithecus gervais, 1849 (primates: Pliopithecidae) from the middle miocene (mn8) of abocador de can mata (els hostalets de pierola, catalonia, spain). *American Journal of Physical Anthropology: The Official Publication of the American Association of Physical Anthropologists*, 141(1):52–75, 2010.

103. David M Alba, Salvador Moyà-Solà, Josep M Robles, and Jordi Galindo. Brief communication: The oldest pliopithecid record in the iberian peninsula based on new material from the vallès-penedès basin. *American Journal of Physical Anthropology*, 147(1):135–140, 2012.

104. Peter Andrews. Two new fossil primates from the lower miocene of kenya. *Nature*, 228(5271):537–540, 1970.

105. Peter Andrews. New species of dryopithecus from kenya. *Nature*, 249:188–190, 1974.

106. Mark RL Anthony and Richard F Kay. Tooth form and diet in ateline and alouattine primates: reflections on the comparative method. *American Journal of Science*, 293(A):356, 1993.

107. Berhane Asfaw, Tim White, Owen Lovejoy, Bruce Latimer, Scott Simpson, and Gen Suwa. Australopithecus garhi: a new species of early hominid from ethiopia. *Science*, 284(5414):629–635, 1999.

108. Amy L Atwater and E Christopher Kirk. New middle eocene omomyines (primates, haplorhini) from san diego county, california. *Journal of Human Evolution*, 124:7–24, 2018.

109. Amy Lynn Atwater et al. *New Middle Eocene Omomyines (Primates, Haplorhini) from the Friars Formation of San Diego County, Southern California*. PhD thesis, 2017.

110. Anne-Marie Bacon, Fabrice Demeter, Mathieu Schuster, Vu The Long, Nguyen Kim Thuy, Pierre-Olivier Antoine, Sevket Sen, Ha Huu Nga, and Nguyen Mai Huong. The pleistocene ma u'oi cave, northern vietnam: palaeontology, sedimentology and palaeoenvironments. *Geobios*, 37(3):305–314, 2004.

111. Sunil Bajpai, Richard F Kay, Blythe A Williams, Debasis P Das, Vivesh V Kapur, and BN Tiwari. The oldest asian record of anthropoidea. *Proceedings of the National Academy of Sciences*, 105(32):11093–11098, 2008.

112. John C Barry, Louis L Jacobs, and Jay Kelley. An early middle miocene catarrhine from pakistan with comments on the dispersal of catarrhines into eurasia. *Journal of Human Evolution*, 15(6):501–508, 1986.

113. K Christopher Beard, Matthew F Jones, Nicholas A Thurber, and Oscar Sanisidro. Systematics and paleobiology of chiromyoides (mammalia, plesiadapidae) from the upper paleocene of western north america and western europe. *Journal of Vertebrate Paleontology*, 39(6):e1730389, 2019.

114. K Christopher Beard, Laurent Marivaux, Yaowalak Chaimanee, Jean-Jacques Jaeger, Bernard Marandat, Paul Tafforeau, Aung Naing Soe, Soe Thura Tun, and Aung Aung Kyaw. A new primate from the eocene pondaung formation of myanmar and the monophyly of burmese amphipithecids. *Proceedings of the Royal Society B: Biological Sciences*, 276(1671):3285–3294, 2009.

115. K Christopher Beard, Grégoire Métais, Faruk Ocakoğlu, and Alexis Licht. An omomyid primate from the pontide microcontinent of north-central anatolia: Implications for sweepstakes dispersal of terrestrial mammals during the eocene. *Geobios*, 66:143–152, 2021.

116. KC Beard. A new species of carpocristes (mammalia: Primatomorpha) from the middle tiffanian of the bison basin, wyoming, with notes on carpolestid phylogeny. *Annals of Carnegie Museum*, 69(3):195–208, 2000.

117. David R Begun. Dryopithecus crusafonti sp. nov., a new miocene hominoid species from can ponsic (northeastern spain). *American Journal of Physical Anthropology*, 87(3):291–309, 1992.

118. Lee R Berger and John E Parkington. A new pleistocene hominid-bearing locality at hoedjiespunt, south africa. *American Journal of Physical Anthropology*, 98(4):601–609, 1995.

119. Raymond L Bernor, Lawrence J Flynn, Terry Harrison, S Taseer Hussain, and Jay Kelley. Dionysopithecus from southern pakistan and the biochronology and biogeography of early eurasian catarrhines. *Journal of Human Evolution*, 17(3):339–358, 1988.

120. Jonathan I Bloch, Doug M Boyer, Philip D Gingerich, and Gregg F Gunnell. New primitive paromomyid from the clarkforkian of wyoming and dental eruption in plesiadapiformes. *Journal of Vertebrate Paleontology*, 22(2):366–379, 2002.

121. Jonathan I Bloch and Philip D Gingerich. Carpolestes simpsoni, new species (mammalia, proprimates) from the late paleocene of the clark's fork basin, wyoming. 1998.

122. Bennett Blumenberg and Andrew T Lloyd. Australopithecus and the origin of the genus homo: Aspects of biometry and systematics with accompanying catalog of tooth metric data. *BioSystems*, 16(2):127–167, 1983.

123. Thomas M Bown and Kenneth D Rose. Patterns of dental evolution in early eocene anaptomorphine primates (omomyidae) from the bighorn basin, wyoming. *Memoir (The Paleontological Society)*, pages 1–162, 1987.

124. Thomas M Bown and Kenneth D Rose. Evolutionary relationships of a new genus and three new species of omomyid primates (willwood formation, lower eocene, bighorn basin, wyoming). *Journal of Human Evolution*, 20(6):465–480, 1991.

125. Doug M Boyer, Loïc Costeur, and Yaron Lipman. Earliest record of platychoerops (primates, plesiadapidae), a new species from mouras quarry, mont de berru, france. *American Journal of Physical Anthropology*, 149(3):329–346, 2012.

126. Doug M Boyer, Stephanie A Maiolino, Patricia A Holroyd, Paul E Morse, and Jonathan I Bloch. Oldest evidence for grooming claws in euprimates. *Journal of Human Evolution*, 122:1–22, 2018.

127. Doug M Boyer, Craig S Scott, and Richard C Fox. New craniodental material of pronothodectes gaoi fox (mammalia,"plesiadapiformes") and relationships among members of plesiadapidae. *American Journal of Physical Anthropology*, 147(4):511–550, 2012.

128. Doug M Boyer, Erik R Seiffert, Justin T Gladman, and Jonathan I Bloch. Evolution and allometry of calcaneal elongation in living and extinct primates. *PloS one*, 8(7):e67792, 2013.

129. Doug M Boyer, Erik R Seiffert, and Elwyn L Simons. Astragalar morphology of afradapis, a large adapiform primate from the earliest late eocene of egypt. *American Journal of Physical Anthropology*, 143(3):383–402, 2010.

130. C Loring Brace. Krapina,"classic" neanderthals, and the evolution of the european face. *Journal of Human Evolution*, 8(5):527–550, 1979.

131. Michel Brunet, Franck Guy, David Pilbeam, Daniel E Lieberman, Andossa Likius, Hassane T Mackaye, Marcia S Ponce de Leon, Christoph PE Zollikofer, and Patrick Vignaud. New material of the earliest hominid from the upper miocene of chad. *Nature*, 434(7034):752–755, 2005.

132. Michel Brunet, Franck Guy, David Pilbeam, Hassane Taisso Mackaye, Andossa Likius, Djimdoumalbaye Ahounta, Alain Beauvilain, Cécile Blondel, Hervé Bocherens, Jean-Renaud Boisserie, et al. A new hominid from the upper miocene of chad, central africa. *Nature*, 418(6894):145–151, 2002.

133. Gregory A Buckley. A new species of purgatorius (mammalia; primatomorpha) from the lower paleocene bear formation, crazy mountains basin, south-central montana. *Journal of Paleontology*, 71(1):149–155, 1997.

134. Benjamin J Burger and James G Honey. Plesiadapidae (mammalia, primates) from the late paleocene fort union formation of the piceance creek basin, colorado. *Journal of Vertebrate Paleontology*, 28(3):816–825, 2008.

135. Benjamin John Burger. Skull of the eocene primate omomys carteri from western north america. *Paleontological Contributions*, 2010(2):1–19, 2010.

136. Yaowalak Chaimanee. Siamopithecus eocaenus, anthropoid primate from the late eocene of krabi, thailand. *Anthropoid origins: new visions*, pages 341–368, 2004.

137. Yaowalak Chaimanee, Olivier Chavasseau, K Christopher Beard, Aung Aung Kyaw, Aung Naing Soe, Chit Sein, Vincent Lazzari, Laurent Marivaux, Bernard Marandat, Myat Swe, et al. Late middle eocene primate from myanmar and the initial anthropoid colonization of africa. *Proceedings of the National Academy of Sciences*, 109(26):10293–10297, 2012.

138. Yaowalak Chaimanee, Vincent Lazzari, Mouloud Benammi, Adélaïde Euriat, and Jean-Jacques Jaeger. A new small pliopithecoid primate from the middle miocene of thailand. *Journal of Human Evolution*, 88:15–24, 2015.

139. Yaowalak Chaimanee, Renaud Lebrun, Chotima Yamee, and Jean-Jacques Jaeger. A new middle miocene tarsier from thailand and the reconstruction of its orbital morphology using a geometric–morphometric method. *Proceedings of the Royal Society B: Biological Sciences*, 278(1714):1956–1963, 2011.

140. Stephen GB Chester and Jonathan I Bloch. Systematics of paleogene micromomyidae (euarchonta, primates) from north america. *Journal of Human Evolution*, 65(2):109–142, 2013.

141. Russell Ciochon. *New interpretations of ape and human ancestry*. Springer Science & Business Media, 2012.

142. William A Clemens. Purgatorius, an early paromomyid primate (mammalia). *Science*, 184(4139):903–905, 1974.

143. William A Clemens, GREGORY P Wilson, and LB Albright. Early torrejonian mammalian local faunas from northeastern montana, usa. *Museum of Northern Arizona Bulletin*, 65:111–158, 2009.

144. Siobhán B Cooke, Alfred L Rosenberger, and Samuel Turvey. An extinct monkey from haiti and the origins of the greater antillean primates. *Proceedings of the National Academy of Sciences*, 108(7):2699–2704, 2011.

145. Susanne Cote, Nasser Malit, and Isaiah Nengo. Additional mandibles of rangwapithecus gordoni, an early miocene catarrhine from the tinderet localities of western kenya. *American Journal of Physical Anthropology*, 153(3):341–352, 2014.

146. Frank P Cuozzo, Emilienne Rasoazanabary, Laurie R Godfrey, Michelle L Sauther, Ibrahim Antho Youssouf, and Marni M LaFleur. Biological variation in a large sample of mouse lemurs from amboasary, madagascar: Implications for interpreting variation in primate biology and paleobiology. *Journal of Human Evolution*, 64(1):1–20, 2013.

147. Frank Patrick Cuozzo. *Craniodental variation in dwarf galagos (Galagoides) and mouse lemurs (Microcebus) and its implications for the paleotaxonomy of small-bodied Eocene primates*. University of Colorado at Boulder, 2000.

148. Eric De Bast, Cyril Gagnaison, and Thierry Smith. Plesiadapid mammals from the latest paleocene of france offer new insights on the evolution of plesiadapis during the paleocene-eocene transition. *Journal of Vertebrate Paleontology*, 38(3):e1460602, 2018.

149. Louis de Bonis and George D Koufos. The face and the mandible of ouranopithecus macedoniensis: description of new specimens and comparisons. *Journal of Human Evolution*, 24(6):469–491, 1993.

150. Louis de Bonis, George D Koufos, Franck Guy, Stéphane Peigné, and Ioanna Sylvestrou. Nouveaux restes du primate hominoïde ouranopithecus dans les dépôts du miocène supérieur de macédoine (grèce). *Comptes Rendus de l'Académie des Sciences-Series IIA-Earth and Planetary Science*, 327(2):141–146, 1998.

151. Alexander R Dutchak. *Mammalian faunal change during the early Eocene climatic optimum (Wasatchian and Bridgerian) at Raven Ridge in the Northeastern Uinta Basin, Colorado and Utah*. University of Colorado at Boulder, 2010.

152. Simon Engelberger. *Annotated catalogue of primate type specimens in the mammal collection of the Museum of Natural History Vienna*. PhD thesis, University of Vienna, 2010.

153. Joan Femenias-Gual, Judit Marigó, Raef Minwer-Barakat, and Salvador Moyà-Solà. New dental and postcranial material of agerinia smithorum (primates, adapiformes) from the type locality casa retjo-1 (early eocene, iberian peninsula). *Journal of Human Evolution*, 113:127–136, 2017.

154. Joan Femenias-Gual, Raef Minwer-Barakat, Judit Marigó, and Salvador Moyà-Solà. Agerinia smithorum sp. nov., a new early eocene primate from the iberian peninsula. *American Journal of Physical Anthropology*, 161(1):116–124, 2016.

155. John G Fleagle, Dennis W Powers, Glenn C Conroy, and John Paul Watters. New fossil platyrrhines from santa cruz province, argentina. *Folia primatologica*, 48(1-2):65–77, 1987.

156. John G Fleagle and Elwyn L Simons. Micropithecus clarki, a small ape from the miocene of uganda. *American Journal of Physical Anthropology*, 49(4):427–440, 1978.

157. John J Flynn, André R Wyss, Reynaldo Charrier, and Carl C Swisher. An early miocene anthropoid skull from the chilean andes. *Nature*, 373(6515):603–607, 1995.

158. Richard C Fox. Pronothodectes gaoi n. sp. from the late paleocene of alberta, canada, and the early evolution of the plesiadapidae (mammalia, primates). *Journal of Paleontology*, 64(4):637–647, 1990.

159. Richard C Fox. Saxonella (plesiadapiformes:? primates) in north america: S. naylori, sp. nov., from the late paleocene of alberta, canada. *Journal of Vertebrate Paleontology*, 11(3):334–349, 1991.

160. Richard C Fox. The dentition and relationships of carpodaptes cygneus (russell)(carpolestidae, plesiadapiformes, mammalia), from the late paleocene of alberta, canada. *Journal of Paleontology*, 76(5):864–881, 2002.

161. Jens L Franzen, Philip D Gingerich, Jörg Habersetzer, Jørn H Hurum, Wighart Von Koenigswald, and B Holly Smith. Complete primate skeleton from the middle eocene of messel in germany: morphology and paleobiology. *PLoS one*, 4(5):e5723, 2009.

162. Leo Gabunia and Abesalom Vekua. A plio-pleistocene hominid from dmanisi, east georgia, caucasus. *Nature*, 373(6514):509–512, 1995.

163. Charles Lewis Gazin. Paleocene mammalian faunas of the bison basin in south-central wyoming. *Smithsonian Miscellaneous Collections*, 1956.

164. Charles Lewis Gazin. A review of the middle and upper eocene primates of north america. *Smithsonian Miscellaneous Collections*, 1958.

165. CL Gazin. A new primate from the torrejon middle paleocene of the san juan basin, new mexico. *Proceedings of the Biological Society of Washington*, 81:629–34, 1968.

166. Insaf Gençturk, Berna Alpagut, and Peter Andrews. Interproximal wear facets and tooth associations in the paşalar hominoid sample. *Journal of human evolution*, 54(4):480–493, 2008.

167. Christopher C Gilbert, Emily D Goble, and Andrew Hill. Miocene cercopithecoidea from the tugen hills, kenya. *Journal of Human Evolution*, 59(5):465–483, 2010.

168. Christopher C Gilbert, Biren A Patel, N Premjit Singh, Christopher J Campisano, John G Fleagle, Kathleen L Rust, and Rajeev Patnaik. New sivaladapid primate from lower siwalik deposits surrounding ramnagar (jammu and kashmir state), india. *Journal of human evolution*, 102:21–41, 2017.

169. Philip D Gingerich. Dental variation in early eocene teilhardina belgica, with notes on the anterior dentition of some early tarsiiformes. *Folia Primatologica*, 28(2):144–153, 1977.

170. Philip D Gingerich. New species of eocene primates and the phytogeny of european adapidae. *Folia Primatologica*, 28(1):60–80, 1977.

171. Philip D Gingerich. The stuttgart collection of oligocene primates from the fayum province of egypt. *Paläontologische Zeitschrift*, 52:82–92, 1978.

172. Philip D Gingerich, Demberelyin Dashzeveg, and Donald E Russell. Dentition and systematic relationships of altanius orlovi (mammalia, primates) from the early eocene of mongolia. *Geobios*, 24(5):637–646, 1991.

173. Philip D Gingerich and Richard A Haskin. Dentition of early eocene pelycodus jarrovii (mammalia, primates) and the generic attribution of species formerly referred to pelycodus. 1981.

174. Philip D Gingerich, B Holly Smith, and Karen Rosenberg. Allometric scaling in the dentition of primates and prediction of body weight from tooth size in fossils. *American Journal of Physical Anthropology*, 58(1):81–100, 1982.

175. Philip Dean Gingerich. *Cranial anatomy and evolution of early Tertiary Plesiadapidae (Mammalia, Primates)*. Yale University, 1974.

176. Marc Godinot. A summary of adapiform systematics and phylogeny. *Folia primatologica*, 69(7):218–249, 1998.

177. Marc Godinot, Donald E Russell, and Pierre Louis. Oldest known nannopithex (primates, omomyiformes) from the early eocene of france. *Folia Primatologica*, 58(1):32–40, 1992.

178. Ana Gracia-Téllez, Juan-Luis Arsuaga, Ignacio Martínez, Laura Martín-Francés, María Martinón-Torres, José-María Bermúdez de Castro, Alejandro Bonmatí, and Jaime Lira. Orofacial pathology in homo heidelbergensis: The case of skull 5 from the sima de los huesos site (atapuerca, spain). *Quaternary international*, 295:83–93, 2013.

179. William K Gregory, Milo Hellman, and G Edward Lewis. *Fossil anthropoids of the Yale-Cambridge India expedition of 1935*. Carnegie institution of Washington, 1938.

180. Gregg F Gunnell. Evolutionary history of microsyopoidea (mammalia,? primates) and the relationship between plesiadapiformes and primates. 1989.

181. Gregg F Gunnell. Omomyid primates (tarsiiformes) from the bridger formation, middle eocene, southern green river basin, wyoming. *Journal of Human Evolution*, 28(2):147–187, 1995.

182. Gregg F Gunnell and Philip D Gingerich. A new species of niptomomys (microsyopidae) from the early eocene of wyoming. *Folia Primatologica*, 36(1-2):128–137, 1981.

183. Gregg F Gunnell and Ellen R Miller. Origin of anthropoidea: dental evidence and recognition of early anthropoids in the fossil record, with comments on the asian anthropoid radiation. *American Journal of Physical Anthropology: The Official Publication of the American Association of Physical Anthropologists*, 114(3):177–191, 2001.

184. Yohannes Haile-Selassie, Luis Gibert, Stephanie M Melillo, Timothy M Ryan, Mulugeta Alene, Alan Deino, Naomi E Levin, Gary Scott, and Beverly Z Saylor. New species from ethiopia further expands middle pliocene hominin diversity. *Nature*, 521(7553):483–488, 2015.

185. T Harrison. New finds of small fossil apes from the miocene locality at koru in kenya. *Journal of human Evolution*, 10(2):129–137, 1981.

186. Terry Harrison. New fossil anthropoids from the middle miocene of east africa and their bearing on the origin of the oreopithecidae. *American Journal of Physical Anthropology*, 71(3):265–284, 1986.

187. Terry Harrison. A taxonomic revision of the small catarrhine primates from the early miocene of east africa. *Folia Primatologica*, 50(1-2):59–108, 1988.

188. Terry Harrison. A new species of micropithecus from the middle miocene of kenya. *Journal of Human Evolution*, 18(6):537–557, 1989.

189. Terry Harrison. A reassessment of the taxonomic and phylogenetic affinities of the fossil catarrhines from fort ternan, kenya. *Primates*, 33:501–522, 1992.

190. Terry Harrison. Dendropithecoidea, proconsuloidea, and hominoidea. *Cenozoic mammals of Africa*, pages 429–469, 2010.

191. Terry Harrison, Jan Van der Made, and Francesc Ribot. A new middle miocene pliopithecid from sant quirze, northern spain. *Journal of Human Evolution*, 42(4):371–377, 2002.

192. Terry Harrison and Gu Yumin. Taxonomy and phylogenetic relationships of early miocene catarrhines from sihong, china. *Journal of Human Evolution*, 37(2):225–277, 1999.

193. Winfried Henke, Ian Tattersall, et al. Handbook of paleoanthropology (vol iii). 2015.

194. Andrew Hill, Kay Behrensmeyer, Barbara Brown, Alan Deino, Michael Rose, Jeffrey Saunders, Steven Ward, and Alisa Winkler. Kipsaramon: a lower miocene hominoid site in the tugen hills, baringo district, kenya. *Journal of Human Evolution*, 20(1):67–75, 1991.

195. Andrew Hill, Isaiah Odhiambo Nengo, and James B Rossie. A rangwapithecus gordoni mandible from the early miocene site of songhor, kenya. *Journal of Human Evolution*, 65(5):490–500, 2013.

196. LESLEA J Hlusko. A new late miocene species of paracolobus and other cercopithecoidea (mammalia: Primates) fossils from lemudong'o, kenya. *Kirtlandia*, 56(72):e85, 2007.

197. Leslea J Hlusko, Joshua P Carlson, Debbie Guatelli-Steinberg, Kristin L Krueger, Ben Mersey, Peter S Ungar, and Alban Defleur. Neanderthal teeth from moula-guercy, ardèche, france. *American Journal of Physical Anthropology*, 151(3):477–491, 2013.

198. Jeremy J Hooker and David L Harrison. A new clade of omomyid primates from the european paleogene. *Journal of Vertebrate Paleontology*, 28(3):826–840, 2008.

199. JJ Hooker. A new microchoerine omomyid (primates, mammalia) from the english early eocene and its palaeobiogeographical implications. *Palaeontology*, 50(3):739–756, 2007.

200. JJ Hooker, DE Russell, and Al Phélizon. A new family of plesiadapiformes (mammalia) from the old world lower paleogene. *Palaeontology*, 42(3):377–407, 1999.

201. Inés Horovitz and Ross DE MacPhee. The quaternary cuban platyrrhineparalouatta varonaiand the origin of antillean monkeys. *Journal of Human Evolution*, 36(1):33–68, 1999.

202. Jean-Jacques Hublin, Abdelouahed Ben-Ncer, Shara E Bailey, Sarah E Freidline, Simon Neubauer, Matthew M Skinner, Inga Bergmann, Adeline Le Cabec, Stefano Benazzi, Katerina Harvati, et al. New fossils from jebel irhoud, morocco and the pan-african origin of homo sapiens. *Nature*, 546(7657):289–292, 2017.

203. Mykolas Imbrasas. *Performance of 2D geometric morphometric analysis in hominin taxonomy and its application to the mandibular molar sample from Lomekwi, Kenya*. PhD thesis, University of Kent (United Kingdom), 2018.

204. Louis L Jacobs. Miocene lorisid primates from the pakistan siwaliks. *Nature*, 289(5798):585–587, 1981.

205. J-J Jaeger, Tin Thein, M Benammi, Y Chaimanee, Aung Naing Soe, Thit Lwin, Than Tun, San Wai, and S Ducrocq. A new primate from the middle eocene of myanmar and the asian early origin of anthropoids. *Science*, 286(5439):528–530, 1999.

206. Jean-Jacques Jaeger, K Christopher Beard, Yaowalak Chaimanee, Mustafa Salem, Mouloud Benammi, Osama Hlal, Pauline Coster, Awad A Bilal, Philippe Duringer, Mathieu Schuster, et al. Late middle eocene epoch of libya yields earliest known radiation of african anthropoids. *Nature*, 467(7319):1095–1098, 2010.

207. Jean-Jacques Jaeger, Olivier Chavasseau, Vincent Lazzari, Aung Naing Soe, Chit Sein, Anne Le Maître, Hla Shwe, and Yaowalak Chaimanee. New eocene primate from myanmar shares dental characters with african eocene crown anthropoids. *Nature Communications*, 10(1):3531, 2019.

208. RF Kay. A new small platyrrhine and the phyletic position of callitrichinae. *Vertebrate Paleontology in the Neotropics: the Miocene fauna of La Venta, Colombia*, pages 435–458, 1997.

209. Richard F Kay. Sivapithecus simonsi, a new species of miocene hominoid, with comments on the phylogenetic status of the ramapithecinae. *International Journal of Primatology*, 3:113–173, 1982.

210. Richard F Kay. Large fossil platyrrhines from the rio acre local fauna, late miocene, western amazonia. *J. Hum. Evol.*, 25:319–327, 1993.

211. Richard F Kay. "giant" tamarin from the miocene of colombia. *American Journal of Physical Anthropology*, 95(3):333–353, 1994.

212. Richard F Kay. A new primate from the early miocene of gran barranca, chubut province, argentina: Paleoecological implications. *The Paleontology of Gran Barranca: Evolution and Environmental Change through the Middle Cenozoic of Patagonia. Cambridge University Press, Cambridge*, pages 220–239, 2010.

213. Richard F Kay and Mario Alberto Cozzuol. New platyrrhine monkeys from the solimões formation (late miocene, acre state, brazil). *Journal of Human Evolution*, 50(6):673–686, 2006.

214. Richard F Kay, JG Fleagle, TRT Mitchell, Matthew Colbert, Tom Bown, and Dennis W Powers. The anatomy of dolichocebus gaimanensis, a stem platyrrhine monkey from argentina. *Journal of Human Evolution*, 54(3):323–382, 2008.

215. Richard F Kay, John G Fleagle, and Elwyn L Simons. A revision of the oligocene apes of the fayum province, egypt. *American Journal of Physical Anthropology*, 55(3):293–322, 1981.

216. Richard F Kay, Lauren A Gonzales, Wout Salenbien, Jean-Noël Martinez, Siobhán B Cooke, Luis Angel Valdivia, Catherine Rigsby, and Paul A Baker. Parvimico materdei gen. et sp. nov.: A new platyrrhine from the early miocene of the amazon basin, peru. *Journal of human evolution*, 134:102628, 2019.

217. Richard F Kay, Richard H Madden, and Javier Guerrero Díaz. Nuevos hallazgos de monos en el mioceno de colombia. *Ameghiniana*, 25(3):203–212, 1988.

218. Richard F Kay, Richard H Madden, J Michael Plavcan, Richard L Cifelli, and Javier Guerrero Díaz. Stirtonia victoriae, a new species of miocene colombian primate. *Journal of Human Evolution*, 16(2):173–196, 1987.

219. Richard F Kay, Jonathan MG Perry, MD Malinzak, Kari L Allen, E Christopher Kirk, J Michael Plavcan, and John G Fleagle. The paleobiology of santacrucian primates. *Early Miocene Paleobiology in Patagonia: High-Latitude Paleocommunities of the Santa Cruz Formation*, pages 306–330, 2012.

220. Richard F Kay and Elwyn L Simons. The ecology of oligocene african anthropoidea. *International Journal of Primatology*, 1:21–37, 1980.

221. Richard F Kay and Peter S Ungar. Dental evidence for diet in some miocene catarrhines with comments on the effects of phylogeny on the interpretation of adaptation. *Function, phylogeny, and fossils: Miocene hominoid evolution and adaptations*, pages 131–151, 1997.

222. Richard F Kay and Blythe A Williams. Dental evidence for anthropoid origins. *Anthropoid origins*, pages 361–445, 1994.

223. Richard Frederick Kay and Jonathan Marcus G Perry. New primates from the río santa cruz and río bote (early-middle miocene), santa cruz province, argentina. *Publicación Electrónica de la Asociación Paleontológica Argentina*, 19(2):230–238, 2019.

224. Jay Kelley. A new large species of sivapithecus from the siwaliks of pakistan. *Journal of Human Evolution*, 17(3):305–324, 1988.

225. Jay Kelley, Steve Ward, Barbara Brown, Andrew Hill, and Dana L Duren. Dental remains of equatorius africanus from kipsaramon, tugen hills, baringo district, kenya. *Journal of human evolution*, 42(1-2):39–62, 2002.

226. Thomas S Kelly and Paul C Murphey. Mammals from the earliest uintan (middle eocene) turtle bluff member, bridger formation, southwestern wyoming, usa, part 1: Primates and rodentia. *Palaeontologia Electronica*, 19:1–55, 2016.

227. E Christopher Kirk and Elwyn L Simons. Diets of fossil primates from the fayum depression of egypt: a quantitative analysis of molar shearing. *Journal of Human Evolution*, 40(3):203–229, 2001.

228. E Christopher Kirk and Blythe A Williams. New adapiform primate of old world affinities from the devil's graveyard formation of texas. *Journal of Human Evolution*, 61(2):156–168, 2011.

229. László Kordos and David R Begun. A new reconstruction of rud 77, a partial cranium of dryopithecus brancoi from rudabánya, hungary. *American Journal of Physical Anthropology: The Official Publication of the American Association of Physical Anthropologists*, 103(2):277–294, 1997.

230. George D Koufos. Mandible of ouranopithecus macedoniensis (hominidae, primates) from a new late miocene locality of macedonia (greece). *American Journal of Physical Anthropology*, 91(2):225–234, 1993.

231. George D Koufos. The first female maxilla of the hominoid ouranopithecus macedoniensis from the late miocene of macedonia, greece. *Journal of Human Evolution*, 29(4):385–399, 1995.

232. David W Krause and Philip D Gingerich. Mammalian fauna from douglass quarry, earliest tiffanian (late paleocene) of the eastern crazy mountain basin, montana. 1983.

233. Yutaka Kunimatsu. A revision of the hypodigm of nyanzapithecus vancouveringi. *African Study Monographs*, 13(4):231–235, 1992.

234. Yutaka Kunimatsu, Benjavun Ratanasthien, Hideo Nakaya, Haruo Saegusa, and Shinji Nagaoka. Earliest miocene hominoid from southeast asia. *American Journal of Physical Anthropology: The Official Publication of the American Association of Physical Anthropologists*, 124(2):99–108, 2004.

235. Yutaka Kunimatsu, Benjavun Ratanasthien, Hideo Nakaya, Haruo Saegusa, Shinji Nagaoka, Yûsuke Suganuma, Akira Fukuchi, and Bantita Udomkan. An additional specimen of a large-bodied miocene hominoid from chiang muan, northern thailand. *Primates*, 46:65–70, 2005.

236. Emily K Lannoye. *A new middle Paleocene mammalian fauna from the Fort Union Formation, Great Divide Basin, Wyoming*. PhD thesis, University of Colorado at Boulder, 2015.

237. LSB Leakey. Lower dentition of kenyapithecus africanus. *Nature*, 217(5131):827–830, 1968.

238. Meave G Leakey, Fred Spoor, Frank H Brown, Patrick N Gathogo, Christopher Kiarie, Louise N Leakey, and Ian McDougall. New hominin genus from eastern africa shows diverse middle pliocene lineages. In *Human Evolution Source Book*, pages 118–127. Routledge, 2016.

239. MG Leakey, Peter S Ungar, and A Walker. A new genus of large primate from the late oligocene of lothidok, turkana district, kenya. *Journal of Human Evolution*, 28(6):519–531, 1995.

240. RE Leakey, Meave G Leakey, and AC Walker. Morphology of turkanapithecus kalakolensis from kenya. *American Journal of Physical Anthropology*, 76(3):277–288, 1988.

241. Jason A Lillegraven. Primates from later eocene rocks of southern california. *Journal of Mammalogy*, 61(2):181–204, 1980.

242. Wu Liu and Liang Zheng. Comparisons of tooth size and morphology between the late miocene hominoids from lufeng and yuanmou, china, and their implications. *Anthropological Science*, 113(1):73–77, 2005.

243. Donald L Lofgren, James G Honey, Malcolm C McKenna, Robert L Zondervan, Erin E Smith, X Wang, and LG Barnes. Paleocene primates from the goler formation of the mojave desert in california. *Geology and Vertebrate Paleontology of Western and Southern North America, Contributions in Honor of David P. Whistler: National History Museum of Los Angeles County Science Series*, 41:11–28, 2008.

244. Sergi López-Torres. *The Paromomyidae (Primates, Mammalia): Systematics, Evolution, and Ecology*. PhD thesis, University of Toronto (Canada), 2017.

245. Sergi López-Torres and Mary T Silcox. The european paromomyidae (primates, mammalia): taxonomy, phylogeny, and biogeographic implications. *Journal of Paleontology*, 92(5):920–937, 2018.

246. Sergi López-Torres, Mary T Silcox, and Patricia A Holroyd. New omomyoids (euprimates, mammalia) from the late uintan of southern california, usa, and the question of the extinction of the paromomyidae (plesiadapiformes, primates). *Palaeontologia Electronica*, 21(3):1–28, 2018.

247. Teresa Elise MacDonald. Late paleocene (tiffanian) mammal-bearing localities in superposition, from near drumheller, alberta. 1996.

248. DG MacInnes. Notes on the east african miocene primates. *Journal of the East Africa and Uganda Natural History Society*, 17(3-4):141–181, 1943.

249. Ross DE MacPhee and Inés Horovitz. New craniodental remains of the quaternary jamaican monkey xenothrix mcgregori (xenotrichini, callicebinae, pitheciidae), with a reconsideration of the aotus hypothesis1. *American Museum Novitates*, 2004(3434):1–51, 2004.

250. Cary T Madden and G Edward Lewis. Indopithecus giganteus distinct from sivapithecus indicus. *Primates*, 21:572–576, 1980.

251. Anne Dambricourt Malassé, Pu Zhang, and Patricia Wils. A new molar from the middle pleistocene hominid assemblage of yanhuidong, tongzi, south china. *Acta Anthropologica Sinica*, 37:1–17, 2018.

252. Judit Marigó, Raef Minwer-Barakat, and Salvador Moyà-Solà. Nievesia sossisensis, a new anchomomyin (adapiformes, primates) from the early late eocene of the southern pyrenees (catalonia, spain). *Journal of Human Evolution*, 64(6):473–485, 2013.

253. Judit Marigó, I Susanna, R Minwer-Barakat, J Madurell-Malapeira, S Moyà-Solà, I Casanovas-Vilar, JM Robles, and DM Alba. The primate fossil record in the iberian peninsula. *Journal of Iberian Geology*, 40(1):179–211, 2014.

254. Laurent Marivaux, Sylvain Adnet, Ali J Altamirano-Sierra, Myriam Boivin, François Pujos, Anusha Ramdarshan, Rodolfo Salas-Gismondi, Julia V Tejada-Lara, and Pierre-Olivier Antoine. Neotropics provide insights into the emergence of new world monkeys: New dental evidence from the late oligocene of peruvian amazonia. *Journal of Human Evolution*, 97:159–175, 2016.

255. Laurent Marivaux, Walter Aguirre-Diaz, Aldo Benites-Palomino, Guillaume Billet, Myriam Boivin, Francois Pujos, Rodolfo Salas-Gismondi, Julia V Tejada-Lara, Rafael M Varas-Malca, and Pierre-Olivier Antoine. New record of neosaimiri (cebidae, platyrrhini) from the late middle miocene of peruvian amazonia. *Journal of human evolution*, 146:102835, 2020.

256. Laurent Marivaux, Pierre-Olivier Antoine, Syed Rafiqul Hassan Baqri, Mouloud Benammi, Yaowalak Chaimanee, Jean-Yves Crochet, Dario De Franceschi, Nayyer Iqbal, Jean-Jacques Jaeger, Grégoire Métais, et al. Anthropoid primates from the oligocene of pakistan (bugti hills): data on early anthropoid evolution and biogeography. *Proceedings of the National Academy of Sciences*, 102(24):8436–8441, 2005.

257. Evgeny N Maschenko and Masanura Takai. Primates of the genus altanius (mammalia, primates) from the lower eocene of tsagan-khushu, southern mongolia. *Russ. J. Theriol*, 9(2):61–69, 2010.

258. Mark A Mason. *New fossil primates from the Uintan (Eocene) of southern California*. Museum of Paleontology, University of California, 1990.

259. Hirofumi Matsumura, Nguyen Lan Cuong, Nguyen Kim Thuy, and Tomoko Anezaki. Dental morphology of the early hoabinian, the neolithic da but and the metal age dong son civilized peoples in vietnam. *Zeitschrift für Morphologie und anthropologie*, pages 59–73, 2001.

260. Spencer G Mattingly, K Christopher Beard, Pauline MC Coster, Mustafa J Salem, Yaowalak Chaimanee, and Jean-Jacques Jaeger. A new parapithecine (primates: Anthropoidea) from the early oligocene of libya supports parallel evolution of large body size among parapithecids. *Journal of Human Evolution*, 153:102957, 2021.

261. Monte L McCrossin. New species of bushbaby from the middle miocene of maboko island, kenya. *American Journal of Physical Anthropology*, 89(2):215–233, 1992.

262. D Jeffery Meldrum and Richard F Kay. Nuciruptor rubricae, a new pitheciin seed predator from the miocene of colombia. *American Journal of Physical Anthropology: The Official Publication of the American Association of Physical Anthropologists*, 102(3):407–427, 1997.

263. Ulrike Maria Menz. *Evolution und Funktionsvariabilität von bunodonten Molaren bei Primaten*. PhD thesis, Dissertation, Frankfurt am Main, Johann Wolfgang Goethe-Universität, 2017, 2016.

264. ER Miller, BR Benefit, ML McCrossin, JM Plavcan, MG Leakey, AN El-Barkooky, MA Hamdan, MK Abdel Gawad, SM Hassan, and EL Simons. Systematics of early and middle miocene old world monkeys. *Journal of Human Evolution*, 57(3):195–211, 2009.

265. Raef Minwer-Barakat, Judit Marigó, Damien Becker, and Loïc Costeur. A new primate assemblage from la verrerie de roches (middle eocene, switzerland). *Journal of human evolution*, 113:137–154, 2017.

266. Raef Minwer-Barakat, Judit Marigo, Joan Femenias-Gual, and Salvador Moya-Sola. New material of pseudoloris parvulus (microchoerinae, omomyidae, primates) from the late eocene of sossís (northeastern spain) and its implications for the evolution of pseudoloris. *Journal of Human Evolution*, 83:74–90, 2015.

267. William J Morris. An eocene fauna from the cathedral bluffs tongue of the washakie basin, wyoming. *Journal of Paleontology*, pages 195–203, 1954.

268. Paul E Morse, Stephen GB Chester, Doug M Boyer, Thierry Smith, Richard Smith, Paul Gigase, and Jonathan I Bloch. New fossils, systematics, and biogeography of the oldest known crown primate teilhardina from the earliest eocene of asia, europe, and north america. *Journal of Human Evolution*, 128:103–131, 2019.

269. Salvador Moyà-Solà, Meike Köhler, David M Alba, Isaac Casanovas-Vilar, Jordi Galindo, Josep M Robles, Lluís Cabrera, Miguel Garcés, Sergi Almécija, and Elisabet Beamud. First partial face and upper dentition of the middle miocene hominoid dryopithecus fontani from abocador de can mata (vallès-penedès basin, catalonia, ne spain): taxonomic and phylogenetic implications. *American Journal of Physical Anthropology: The Official Publication of the American Association of Physical Anthropologists*, 139(2):126–145, 2009.

270. Kathleen M Muldoon and Gregg F Gunnell. Omomyid primates (tarsiiformes) from the early middle eocene at south pass, greater green river basin, wyoming. *Journal of human evolution*, 43(4):479–511, 2002.

271. Paul C Murphey and Rachel H Dunn. Hemiacodon engardae, a new species of omomyid primate from the earliest uintan turtle bluff member of the bridger formation, southwestern wyoming, usa. *Journal of Human Evolution*, 57(2):123–130, 2009.

272. Michael E Nelson. Middle eocene primates (mammalia) from southwestern wyoming. *The Southwestern Naturalist*, pages 487–493, 1977.

273. Isaiah Odhiambo Nengo and Tood C Rae. New hominoid fossils from the early miocene site of songhor, kenya. *Journal of Human Evolution*, 23(5):423–429, 1992.

274. Allison Marie Neumann. *A new Early Eocene (Lostcabinian) mammal assemblage from the Main Body of the Wasatch Formation, Northern Green River Basin; The Pinnacles, Sweetwater Co., Wyoming*. University of Alberta (Canada), 2013.

275. Xijun Ni, Yaoming Hu, Yuanqing Wang, and CHUANKUI LI. A clue to the asian origin of euprimates. *Anthropological Science*, 113(1):3–9, 2005.

276. Xijun Ni, Qiang Li, Lüzhou Li, and K Christopher Beard. Oligocene primates from china reveal divergence between african and asian primate evolution. *Science*, 352(6286):673–677, 2016.

277. Xijun Ni, Jin Meng, K Christopher Beard, Daniel L Gebo, Yuanqing Wang, and Chuankui Li. A new tarkadectine primate from the eocene of inner mongolia, china: phylogenetic and biogeographic implications. *Proceedings of the Royal Society B: Biological Sciences*, 277(1679):247–256, 2010.

278. Nelson M Novo and John G Fleagle. New specimens of platyrrhine primates from patagonia (pinturas formation, early miocene). *Ameghiniana*, 52(3):367–372, 2015.

279. Nelson M Novo, Marcelo F Tejedor, María E Pérez, and J Marcelo Krause. New primate locality from the early miocene of patagonia, argentina. *American Journal of Physical Anthropology*, 164(4):861–867, 2017.

280. Harvard University. Museum of Comparative Zoology. *Bulletin of the Museum of Comparative Zoology*, volume 48. The Museum, 1905.

281. Biren A Patel and Ari Grossman. Dental metric comparisons of morotopithecus and afropithecus: Implications for the validity of the genus morotopithecus. *Journal of Human Evolution*, 51(5):506–512, 2006.

282. Rajeev Patnaik, David Cameron, JC Sharma, and Jacob Hogarth. Extinction of siwalik fossil apes: a review based on a new fossil tooth and on palaeoecological and palaeoclimatological evidence. *Anthropological Science*, 113(1):65–72, 2005.

283. Jonathan MG Perry, Richard F Kay, Sergio F Vizcaíno, and M Susana Bargo. Oldest known cranium of a juvenile new world monkey (early miocene, patagonia, argentina): Implications for the taxonomy and the molar eruption pattern of early platyrrhines. *Journal of Human Evolution*, 74:67–81, 2014.

284. Martin Pickford. A new look at kenyapithecus based on recent discoveries in western kenya. *Journal of Human Evolution*, 14(2):113–143, 1985.

285. Martin Pickford and Yutaka Kunimatsu. Catarrhines from the middle miocene (ca. 14.5 ma) of kipsaraman, tugen hills, kenya. *Anthropological Science*, 113(2):189–224, 2005.

286. Martin Pickford, Brigitte Senut, D Gommery, and E Musiime. La distinción entre ugandapithecus y proconsul. *Estudios geológicos*, 65(2):183–241, 2009.

287. David Pilbeam. New hominoid skull material from the miocene of pakistan. *Nature*, 295(5846):232–234, 1982.

288. David R Pilbeam. Tertiary pongidae of east africa: evolutionary relationships and taxonomy. *Bulletin of the Peabody Museum of Natural History*, 31:1–185, 1969.

289. D Tab Rasmussen, Myron Shekelle, Stephen L Walsh, and Bradford O Riney. The dentition of dyseolemur, and comments on the use of the anterior teeth in primate systematics. *Journal of Human Evolution*, 29(4):301–320, 1995.

290. Peter Robinson. Fossil mammalia of the huerfano formation, eocene, of colorado. 1966.

291. Peter Robinson. Diversity starts early: notharctid primates from the sandcouleean (early eocene) of the powder river basin, wyoming, usa. *Historical Biology*, 30(1-2):189–203, 2018.

292. Kenneth D Rose. Anterior dentition and relationships of the early eocene omomyids arapahovius advena and teilhardina demissa, sp. nov. *Journal of Human Evolution*, 28(3):231–244, 1995.

293. Kenneth D Rose and Thomas M Bown. Additional fossil evidence on the differentiation of the earliest euprimates. *Proceedings of the National Academy of Sciences*, 88(1):98–101, 1991.

294. Kenneth D Rose, Stephen GB Chester, Rachel H Dunn, Doug M Boyer, and Jonathan I Bloch. New fossils of the oldest north american euprimate teilhardina brandti (omomyidae) from the paleocene–eocene thermal maximum. *American Journal of Physical Anthropology*, 146(2):281–305, 2011.

295. Kenneth D Rose, Amy E Chew, Rachel H Dunn, Mary J Kraus, Henry C Fricke, and Shawn P Zack. Earliest eocene mammalian fauna from the paleocene-eocene thermal maximum at sand creek divide, southern bighorn basin, wyoming. 2012.

296. Kenneth D Rose, Ross DE MacPhee, and John P Alexander. Skull of early eocene cantius abditus (primates: Adapiformes) and its phylogenetic implications, with a reevaluation of "hesperolemur" actius. *American Journal of Physical Anthropology: The Official Publication of the American Association of Physical Anthropologists*, 109(4):523–539, 1999.

297. Kenneth David Rose. *The Carpolestidae, early tertiary primates from North America*. Museum of Comparative Zoology, Harvard University, 1975.

298. Alfred L Rosenberger. The face of strigorhysis: implications of another tarsier-like, large-eyed eocene north american tarsiiform primate. *The Anatomical Record: Advances in Integrative Anatomy and Evolutionary Biology*, 294(5):797–812, 2011.

299. Alfred L Rosenberger, Zachary S Klukkert, Siobhán B Cooke, and Renato Rímoli. Rethinking antillothrix: the mandible and its implications. *American Journal of Primatology*, 75(8):825–836, 2013.

300. James B Rossie and Andrew Hill. A new species of simiolus from the middle miocene of the tugen hills, kenya. *Journal of human evolution*, 125:50–58, 2018.

301. James B Rossie and Laura MacLatchy. A new pliopithecoid genus from the early miocene of uganda. *Journal of Human Evolution*, 50(5):568–586, 2006.

302. Damián Ruiz-Ramoni, Ascanio D Rincón, Andrés Solórzano, and Salvador Moya-Sola. The first fossil platyrrhini (primates: anthropoidea) from venezuela: a capuchin monkey from the plio-pleistocene of el breal de orocual. *Journal of Human Evolution*, 105:127–131, 2017.

303. Joshua X Samuels, L Barry Albright, and Theodore J Fremd. The last fossil primate in north america, new material of the enigmatic ekgmowechashala from the arikareean of oregon. *American Journal of Physical Anthropology*, 158(1):43–54, 2015.

304. Anek R Sankhyan, Jay Kelley, and Terry Harrison. A highly derived pliopithecoid from the late miocene of haritalyangar, india. *Journal of Human Evolution*, 105:1–12, 2017.

305. Ross Secord. The tiffanian land-mammal age (middle and late paleocene) in the northern bighorn basin, wyoming. 2008.

306. Erik R Seiffert. Early primate evolution in afro-arabia. *Evolutionary Anthropology: Issues, News, and Reviews*, 21(6):239–253, 2012.

307. Erik R Seiffert, Jonathan MG Perry, Elwyn L Simons, and Doug M Boyer. Convergent evolution of anthropoid-like adaptations in eocene adapiform primates. *Nature*, 461(7267):1118–1121, 2009.

308. Erik R Seiffert, Elwyn L Simons, and Yousry Attia. Fossil evidence for an ancient divergence of lorises and galagos. *Nature*, 422(6930):421–424, 2003.

309. Erik R Seiffert, Elwyn L Simons, Doug M Boyer, Jonathan MG Perry, Timothy M Ryan, and Hesham M Sallam. A fossil primate of uncertain affinities from the earliest late eocene of egypt. *Proceedings of the National Academy of Sciences*, 107(21):9712–9717, 2010.

310. Erik R Seiffert, Elwyn L Simons, William C Clyde, James B Rossie, Yousry Attia, Thomas M Bown, Prithijit Chatrath, and Mark E Mathison. Basal anthropoids from egypt and the antiquity of africa's higher primate radiation. *Science*, 310(5746):300–304, 2005.

311. Erik R Seiffert, Elwyn L Simons, John G Fleagle, and Marc Godinot. Paleogene anthropoids. *Cenozoic mammals of Africa. Berkeley: University of California Press. p*, pages 369–391, 2010.

312. Erik R Seiffert, Elwyn L Simons, Timothy M Ryan, and Yousry Attia. Additional remains of wadilemur elegans, a primitive stem galagid from the late eocene of egypt. *Proceedings of the National Academy of Sciences*, 102(32):11396–11401, 2005.

313. Erik R Seiffert, Marcelo F Tejedor, John G Fleagle, Nelson M Novo, Fanny M Cornejo, Mariano Bond, Dorien De Vries, and Kenneth E Campbell Jr. A parapithecid stem anthropoid of african origin in the paleogene of south america. *Science*, 368(6487):194–197, 2020.

314. Takeshi Setoguchi and Alfred L Rosenberger. A fossil owl monkey from la venta, colombia. *Nature*, 326(6114):692–694, 1987.

315. Takeshi Setoguchi, Nobuo Shigehara, Alfred L Rosenberger, and Alberto Cadena G. Primate fauna from the miocene la venta, in the tatacoa desert, department of huila, colombia. *Caldasia*, pages 761–773, 1986.

316. Mary T Silcox, Gregg F Gunnell, and Jonathan I Bloch. Cranial anatomy of microsyops annectens (microsyopidae, euarchonta, mammalia) from the middle eocene of northwestern wyoming. *Journal of Paleontology*, 94(5):979–1006, 2020.

317. Mary T Silcox, David W Krause, Mary C Maas, and Richard C Fox. New specimens of elphidotarsius russelli (mammalia,? primates, carpolestidae) and a revision of plesiadapoid relationships. *Journal of Vertebrate Paleontology*, 21(1):132–152, 2001.

318. EL Simons, DT Rasmussen, and DL Gebo. A new species of propliopithecus from the fayum, egypt. *American Journal of Physical Anthropology*, 73(2):139–147, 1987.

319. Elwyn Simons. Egyptian oligocene primates: a review. *American Journal of Physical Anthropology*, 38(S21):199–238, 1995.

320. Elwyn L Simons. The dentition of ourayia: its bearing on relationships of omomyid prosimians. *Postilla*, 54:1–20, 1961.

321. Elwyn L Simons. Miocene monkey (prohylobates) from northern egypt. *Nature*, 223(5207):687–689, 1969.

322. Elwyn L Simons. Parapithecus grangeri (parapithecidae, old world higher primates): new species from the oligocene of egypt and the initial differentiation of ceropithecoidea. *Postilla*, (166), 1974.

323. Elwyn L Simons. Description of two genera and species of late eocene anthropoidea from egypt. *Proceedings of the National Academy of Sciences*, 86(24):9956–9960, 1989.

324. Elwyn L Simons. Diversity in the early tertiary anthropoidean radiation in africa. *Proceedings of the National Academy of Sciences*, 89(22):10743–10747, 1992.

325. Elwyn L Simons. Crania of apidium: primitive anthropoidean (primates, parapithecidae) from the egyptian oligocene. american museum novitates; no. 3124. 1995.

326. Elwyn L Simons. Preliminary description of the cranium of proteopithecus sylviae, an egyptian late eocene anthropoidean primate. *Proceedings of the National Academy of Sciences*, 94(26):14970–14975, 1997.

327. Elwyn L Simons and Thomas M Bown. Afrotarsius chatrathi, first tarsiiform primate (? tarsiidae) from africa. *Nature*, 313(6002):475–477, 1985.

328. Elwyn L Simons and Richard F Kay. Qatrania, new basal anthropoid primate from the fayum, oligocene of egypt. *Nature*, 304(5927):624–626, 1983.

329. Elwyn L Simons and Richard F Kay. New material of qatrania from egypt with comments on the phylogenetic position of the parapithecidae (primates, anthropoidea). *American Journal of Primatology*, 15(4):337–347, 1988.

330. Elwyn L Simons and D Tab Rasmussen. The generic classification of fayum anthropoidea. *International journal of primatology*, 12:163–178, 1991.

331. Elwyn L Simons and D Tab Rasmussen. A remarkable cranium of plesiopithecus teras (primates, prosimii) from the eocene of egypt. *Proceedings of the National Academy of Sciences*, 91(21):9946–9950, 1994.

332. Elwyn L Simons, Erik R Seiffert, Prithijit S Chatrath, and Yousry Attia. Earliest record of a parapithecid anthropoid from the jebel qatrani formation, northern egypt. *Folia Primatologica*, 72(6):316–331, 2001.

333. Elwyn L Simons, Erik R Seiffert, Timothy M Ryan, and Yousry Attia. A remarkable female cranium of the early oligocene anthropoid aegyptopithecus zeuxis (catarrhini, propliopithecidae). *Proceedings of the National Academy of Sciences*, 104(21):8731–8736, 2007.

334. George Gaylord Simpson. The fort union of the crazy mountain field, montana, and its mammalian faunas. *Bulletin of the United States National Museum*, 1937.

335. George Gaylord Simpson, Walter Granger, et al. The tiffany fauna, upper paleocene. 3, primates, carnivora, condylarthra, and amblypoda. american museum novitates; no. 817. 1935.

336. Matthew M Skinner, Meave G Leakey, Louise N Leakey, Fredrick K Manthi, and Fred Spoor. Hominin dental remains from the pliocene localities at lomekwi, kenya (1982–2009). *Journal of Human Evolution*, 145:102820, 2020.

337. Thierry Smith, Kenneth D Rose, and Philip D Gingerich. Rapid asia–europe–north america geographic dispersal of earliest eocene primate teilhardina during the paleocene–eocene thermal maximum. *Proceedings of the National Academy of Sciences*, 103(30):11223–11227, 2006.

338. Salvador Moyà Solà and Meike Köhler. New partial cranium of dryopithecus lartet, 1863 (hominoidea, primates) from the upper miocene of can llobateres, barcelona, spain. *Journal of Human Evolution*, 29(2):101–139, 1995.

339. Nancy J Stevens, Erik R Seiffert, Patrick M O'Connor, Eric M Roberts, Mark D Schmitz, Cornelia Krause, Eric Gorscak, Sifa Ngasala, Tobin L Hieronymus, and Joseph Temu. Palaeontological evidence for an oligocene divergence between old world monkeys and apes. *Nature*, 497(7451):611–614, 2013.

340. Chester Stock. Microsyopsinae and hyopsodontidae in the sespe upper eocene, california. *Proceedings of the National Academy of Sciences*, 20(6):349–354, 1934.

341. Chester Stock. A tarsiid primate and a mixodectid from the poway eocene, california. *Proceedings of the National Academy of Sciences*, 24(7):288–293, 1938.

342. GEN Suwa, Berhane Asfaw, Yohannes Haile-Selassie, TIM White, Shigehiro Katoh, Giday WoldeGabriel, William K Hart, Hideo Nakaya, and Yonas Beyene. Early pleistocene homo erectus fossils from konso, southern ethiopia. *Anthropological Science*, 115(2):133–151, 2007.

343. Frederick S Szalay. Systematics of the omomyidae (tarsiiformes, primates): taxonomy, phylogeny, and adaptations. bulletin of the amnh; v. 156, article 3. (3), 1976.

344. Masanaru Takai. New specimens of neosaimiri fieldsi from la venta, colombia: a middle miocene ancestor of the living squirrel monkeys. *Journal of Human Evolution*, 27(4):329–360, 1994.

345. Masanaru Takai and Federico Anaya. New specimens of the oldest fossil platyrrhine, branisella boliviana, from salla, bolivia. *American Journal of Physical Anthropology: The Official Publication of the American Association of Physical Anthropologists*, 99(2):301–317, 1996.

346. Masanaru Takai, Federico Anaya, Nobuo Shigehara, and Takeshi Setoguchi. New fossil materials of the earliest new world monkey, branisella boliviana, and the problem of platyrrhine origins. *American Journal of Physical Anthropology: The Official Publication of the American Association of Physical Anthropologists*, 111(2):263–281, 2000.

347. Masanaru Takai, Federico Anaya, Hisashi Suzuki, Nobuo Shigehara, and Takeshi Setoguchi. A new platyrrhine from the middle miocene of la venta, colombia, and the phyletic position of callicebinae. *Anthropological Science*, 109(4):289–307, 2001.

348. Masanaru Takai, Takeshi Nishimura, Nobuo Shigehara, and Takesh Setoguch. Meaning of the canine sexual dimorphism in fossil owl monkey, aotus dindensis from the middle miocene of la venta, colombia. In *Comparative Dental Morphology*, volume 13, pages 55–59. Karger Publishers, 2009.

349. Marcelo Fabian Tejedor and Alfred L Rosenberger. A neotype for homunculus patagonicus ameghino, 1891, and a new interpretation of the taxon. 2008.

350. Matthew A Tornow. Systematic analysis of the eocene primate family omomyidae using gnathic and postcranial data. *Bulletin of the Peabody Museum of Natural History*, 49(1):43–129, 2008.

351. Erik Trinkaus. Dental remains from the shanidar adult neanderthals. *Journal of Human evolution*, 7(5):369–382, 1978.

352. Gerrit D Van den Bergh, Yousuke Kaifu, Iwan Kurniawan, Reiko T Kono, Adam Brumm, Erick Setiyabudi, Fachroel Aziz, and Michael J Morwood. Homo floresiensis-like fossils from the early middle pleistocene of flores. *Nature*, 534(7606):245–248, 2016.

353. Wei Wang. New discoveries of gigantopithecus blacki teeth from chuifeng cave in the bubing basin, guangxi, south china. *Journal of Human Evolution*, 57(3):229–240, 2009.

354. Michael William Webb. Late paleocene mammals from near drayton valley, alberta. 1996.

355. Robert M West. Fossil mammals from the lower buck hill group, eocene of trans-pecos texas: Marsupicarnivora, primates, taeniodonta, condylarthra, bunodont artiodactyla, and dinocerata. Technical report, Texas Memorial Museum, The University of Texas at Austin, 1982.

356. James W Westgate. Uintan land mammals (excluding rodents) from an estuarine facies of the laredo formation (middle eocene, claiborne group) of webb county, texas. *Journal of Paleontology*, 64(3):454–468, 1990.

357. Brandon C Wheeler. Community ecology of the middle miocene primates of la venta, colombia: the relationship between ecological diversity, divergence time, and phylogenetic richness. *Primates*, 51:131–138, 2010.

358. Tim D White, Gen Suwa, and Berhane Asfaw. Australopithecus ramidus, a new species of early hominid from aramis, ethiopia. *Nature*, 371(6495):306–312, 1994.

359. Tim D White, Gen Suwa, Scott Simpson, and Berhane Asfaw. Jaws and teeth of australopithecus afarensis from maka, middle awash, ethiopia. *American Journal of Physical Anthropology: The Official Publication of the American Association of Physical Anthropologists*, 111(1):45–68, 2000.

360. Blythe A Williams and Herbert H Covert. New early eocene anaptomorphine primate (omomyidae) from the washakie basin, wyoming, with comments on the phylogeny and paleobiology of anaptomorphines. *American Journal of Physical Anthropology*, 93(3):323–340, 1994.

361. Blythe A Williams and E Christopher Kirk. New uintan primates from texas and their implications for north american patterns of species richness during the eocene. *Journal of Human Evolution*, 55(6):927–941, 2008.

362. XLIE Xiang-xu and E Delson. A new species of dryopithecus from gansu, china. *Kexue tongbao (English edition)*, 34(3):223–229, 1989.

363. Zhang Yinyun. Variability and evolutionary trends in tooth size of gigantopithecus blacki. *American Journal of Physical Anthropology*, 59(1):21–32, 1982.

364. Pan Yuerong, Diane M Waddle, and John G Fleagle. Sexual dimorphism in laccopithecus robustus, a late miocene hominoid from china. *American Journal of Physical Anthropology*, 79(2):137–158, 1989.

365. Yahdi Zaim, Russell L Ciochon, Joshua M Polanski, Frederick E Grine, E Arthur Bettis III, Yan Rizal, Robert G Franciscus, Roy R Larick, Matthew Heizler, K Lindsay Eaves, et al. New 1.5 million-year-old homo erectus maxilla from sangiran (central java, indonesia). *Journal of Human Evolution*, 61(4):363–376, 2011.

366. Iyad S Zalmout, William J Sanders, Laura M MacLatchy, Gregg F Gunnell, Yahya A Al-Mufarreh, Mohammad A Ali, Abdul-Azziz H Nasser, Abdu M Al-Masari, Salih A Al-Sobhi, Ayman O Nadhra, et al. New oligocene primate from saudi arabia and the divergence of apes and old world monkeys. *Nature*, 466(7304):360–364, 2010.

367. Yingqi Zhang and Terry Harrison. Gigantopithecus blacki: a giant ape from the pleistocene of asia revisited. *American Journal of Physical Anthropology*, 162:153–177, 2017.

368. Yingqi Zhang, Changzhu Jin, Yanjun Cai, Reiko Kono, Wei Wang, Yuan Wang, Min Zhu, and Yaling Yan. New 400e320 ka gigantopithecus blacki remains from hejiang cave, chongzuo city, guangxi, south china. *Quaternary International*, 30(1):e11, 2013.

369. LX Zhao and LZ Zhang. New fossil evidence and diet analysis of gigantopithecus blacki and its distribution and extinction in south china. *Quaternary International*, 286:69–74, 2013.

370. Jelle S Zijlstra, Lawrence J Flynn, and Wilma Wessels. The westernmost tarsier: A new genus and species from the miocene of pakistan. *Journal of human evolution*, 65(5):544–550, 2013.