Title: NBR1 -mediated selective autophagy of ARF7 modulates root branching

Running title: Autophagy modulates root branching

Ebstrup E[1+], Ansbøl J[1+], Chevalier J[1],Clemmens P[1], Rodriguez E[1*]

[1]Innate immunity and cellular reprogramming group. Functional Genomic section, Department of
Biology, University of Copenhagen, Denmark.

[+]These authors contributed equally to this work.

*Correspondence to: Eleazar.rodriguez@bio.ku.dk

Abstract

Auxin orchestrates growth and developmental regulation in plants. For instance, auxin dictates root

architecture via the Auxin Response Factor (ARF) family of transcription factors, which control lateral

root (LR) formation. In *Arabidopsis*, ARF7 is essential for the correct definition of LR's prebranch sites

(PBS) through gene expression oscillation and plays a pivotal role during LR initiation. Despite ARF7's

importance to this process, there is a surprising lack of knowledge about how ARF7's turnover is

regulated and how this contributes to root architecture patterning. Here, we show that ARF7

accumulates in autophagy mutants and is degraded through NBR1-dependent selective autophagy.

Moreover, our data indicates that the previously reported rhythmic changes to ARF7 abundance in

the root's oscillation zone its observed in the rest of the plant and this is modulated via autophagy.

Surprisingly, autophagy deficiency did not seem to cause major changes in PBS establishment but

ultimately reduces LR formation. In support of these findings, our microscopy data shows that ARF7

and autophagy markers mainly co-localize in mature tissue and the process can be triggered by

auxin. In conclusion we report a novel role for autophagy during plant development, namely by

enacting auxin-induced selective degradation of ARF7 to optimize root branching.

Introduction

The plant root system plays a fundamental role as a point of contact with the plant's surroundings and allows adaptation to multiple stresses. Contrary to primary roots, lateral root (LR) development is post embryonic, which means that LR architecture is heavily influenced by growth conditions (Lv *et al*, 2021). As LR growth is modular, perception of the correct developmental conditions authorizes the repetition of LR modules along the primary root, maximizing the root's surface area. LRs originate from a cell layer called pericycle which surrounds the vasculature, and occurs sequentially in 4 steps: positioning, initiation, outgrowth, and emergence. During the positioning step, the plant ensures the spatiotemporal distribution, priming and activation of LR founder cells (LRFC) (Du & Scheres, 2018). LR spacing is controlled by an oscillatory root clock which dictates gene expression and cycles every 4-6 hours (Moreno-Risueño *et al*, 2010; Wachsman *et al*, 2020). This oscillatory clock is regulated, among others, by the phytohormone auxin and can be characterized through monitoring of reporter markers under the control of DR5, a synthetic promoter containing tandem repeats of an auxin responsive TGTCTC element (Ulmasov *et al*, 1997). DR5 expression analyses lead to definition of 2 contrasting oscillation phases: in-phase (concurrent with DR5 maximum signal) and anti-phase (DR5 minimum signal) which reciprocally repress each other (Moreno-Risueño *et al*, 2010). One of the genes displaying anti-phase expression is AUXIN RESPONSIVE FACTOR 7 (*ARF7*), a key transcription factor (TF) which, among others, regulates LR formation (Okushima *et al*, 2007). Importantly, analysis of *ARF7* loss of function mutant (Moreno-Risueño *et al*, 2010; Perianez-Rodriguez *et al*, 2021) indicates that this TF's oscillation is essential for the correct establishment of LR prebranch sites (PBS).

While positioning and initiation are broadly regulated by different mechanisms, ARF7 and its 2 direct targets LATERAL ORGAN BOUNDARIES DOMAIN (LBD) 16 and LBD33 are among the few genes that participate in both processes (Moreno-Risueño *et al*, 2010); this attests to ARF7's pivotal role in organizing root architecture. During the initiation step, the auxin signalling node IAA14-ARF7-ARF19 controls LBD16 activation in the LRFC, which is necessary to achieve asymmetry and initiate cell division (Goh *et al*, 2019). In addition to the importance of *ARF7* transcriptional oscillation for LR formation, it has also been shown that the subcellular localization of ARF7 is important for its function: nuclear localization correlates with auxin responsive tissue while in less responsive areas, activating ARFs concentrate to form cytoplasmic condensates with liquid-like and solid-like proprieties (Powers *et al*, 2019). These body of evidence demonstrate the regulatory complexity coordinating ARFs-dependent responses, as an intricate web of processes tightly regulate ARFs levels and compartmentalization. Yet surprisingly, little is known about how ARF7 proteostasis is managed and how it impacts the cellular decision making behind LR formation authorization.

Macroautophagy (herein autophagy) is an evolutionary conserved catabolic process which partakes
in cellular recycling of macromolecules and organelles (Levine & Klionsky, 2004). Autophagy is
characterized by the formation of an isolation membrane enclosing a portion of cytoplasm to form a
double-membraned vesicle termed the autophagosome. Originally characterized during starvation
stress, autophagy was viewed as a random process due to the observation of what seemed to be an
indiscriminate engulfment of portions of the cytoplasm. Further studies unveiled the existence of
several selective cargo adaptors which allow for targeted degradation of cellular constituents (*e.g.,*
(Svenning *et al*, 2011; Nolan *et al*, 2017; Stephani *et al*, 2020; Wu *et al*, 2021), meaning that
autophagy can be highly selective and participate in targeted remodelling of the cell. Perhaps one of
the most characterized cargo receptors in plants is NEIGHBOUR OF BRCA1 (NBR1, Svenning *et al*,
2011), a hybrid of mammalian *NBR1* and *p62*, possessing a PB1 domain necessary for
homopolymerization and biomolecular condensate formation, and an UBA domain for ubiquitin
binding. It is through the latter that NBR1 binds ubiquitinated cargo, which are then targeted for
autophagic degradation via NBR1's interaction with the autophagosome decorating proteins ATG8s
(*e.g.,* Zhou et al, 2013, 2014; Hafrén et al, 2017; Ji et al, 2020; Tarnowski et al, 2020). The ability to
function selectively allows autophagy to participate in a variety of stress responses, namely by
promoting the cellular reprogramming necessary to enact appropriated responses (Rodriguez *et al*,
2020). Without surprise, autophagic deficiency has been shown to cause myriads of developmental
defects; interference with core autophagic machinery component often causes lethality during early
developmental stages, severe developmental defects, and shortened life span in metazoans (*e.g.,*
Scott *et al*, 2004; Liu *et al*, 2015; Yoshii *et al*, 2016). In contrast, plants are able to tolerate disruption
of autophagy activity without major penalties: autophagy deficient mutants display subtle growth
phenotypes like reduced biomass, inflorescence numbers and premature senescence (Bassham *et al*,
2006), but importantly, are able to complete their life cycle and generate fertile progeny. It thus
seems that the phenotypic "struggles" of plant *atg* mutants become more apparent under stress, as
can be attested by the many publications linking autophagy with stress responses (Liu *et al*, 2009;
Hofius *et al*, 2009; Vanhee *et al*, 2011; Zhou *et al*, 2013b; Munch *et al*, 2014; Chen *et al*, 2015; Nolan
et al, 2017; Sedaghatmehr *et al*, 2019; Rodriguez *et al*, 2020). For instance, autophagy has been
shown to be involved in primary root growth and LR formation during nutrient starvation (Deb *et al*,
2014; Huang *et al*, 2019), which is consistent with the cross-kingdom descriptions of the starvation-
alleviating properties of autophagy (Scott *et al*, 2004; Desideri *et al*, 2012; Huang *et al*, 2015; Wu *et*
al, 2021). In contrast, the role of autophagy in regulating root architecture under normal
developmental conditions is less understood.

In this work, we explored the contribution of autophagy in mediating ARF7's dependent responses.
By using a combination of cell imaging, biochemical, genetics, molecular biology and proteomics
approaches, we demonstrate that ARF7's turnover is processed via NBR1-mediated selective
autophagy. Importantly, we demonstrate that ARF7's abundance rhythmically oscillates in the whole
plant and autophagy is necessary for proper establishment of ARF7's rhythm. Surprisingly, our data
indicates that autophagy mutants display normal PBS establishment but have reduced LR formation,
and together with ARF7 and ATG8 co-localization experiments, suggest that ARF7 might be regulated
by different mechanisms in a tissue specific manner. Additively, we show that short auxin treatment
is sufficient to trigger ARF7 to localize into cytoplasmic foci, regardless of the root zone, co-
localization of these cytoplasmic foci with ATG8a mainly occurs in the maturation zone. With these
findings, we reveal an important regulatory role for selective autophagy during auxin responses
through the turnover of the crucial auxin signalling regulator ARF7 and consequent fine-tuning of
root architecture.

Results

Autophagy modulates ARFs *turnover*

We have recently reported that autophagy participates in stem cell reprogramming and
differentiation in the distantly related species *P. patens* and *A. thaliana* (Rodriguez *et al*, 2020; Kanne
et al, 2021). Because *Arabidopsis* stem cell formation executes a program similar to LR formation
(Sugimoto *et al*, 2010), we wondered if key regulators of these processes might be modulated via
autophagy. As seen in Fig 1, ARF7 accumulates in different autophagy mutants and in the KO of the
autophagic cargo adaptor NBR1 (Fig 1A), when compared to Col-0. To address if the proteasome also
contributes to ARF7 proteostasis, we treated seedlings of those genotypes with the proteasome
inhibitor MG132 (Fig 1A). Col-0 samples treated with MG132 showed ARF7 accumulation to levels
comparable with those of untreated *atg* mutants. Moreover, simultaneous inhibition of both
degradation pathways leads to further accumulation of ARF7 which indicate that ARF7 proteostasis
is dependent on autophagy and the proteasome and it's in agreement with data by (Jing *et al*, 2022)
showing proteasomal degradation of ARF7. To rule out that elevated transcript levels in *atg* mutants
could be behind ARF7 accumulation in those lines, we performed qPCR and observed that *ARF7*'s
relative expression was around 30% lower in *atg* mutants than Col-0 (Fig 1B).

To further our claim that autophagy modulates ARF7 turnover, we introduced the translational
reporter *pARF7::gARF7-Venus* herein called *ARF7-Venus* (Orosa-Puente *et al*, 2018) into *atg2-1*

background. We then quantified ARF7-Venus levels by western blot analyses of protein extracts
from *ARF7-Venus* and *ARF7-Venus x atg2-1*¸ with this showing significantly higher levels of ARF7-
Venus when autophagy is dysfunctional (Fig 1C). Using the same lines, we also analysed ARF7-Venus
subcellular fluorescence intensity in the root's meristematic and maturation zones by confocal
microscopy (Fig 1 D-I). In agreement with our western blots, we detected significantly higher ARF7-
Venus fluorescence intensity in the *atg2-1* background for all tissues and subcellular localization
except for the meristem's nuclei (Fig1 I). Because ARF7 often works together with ARF19 during
regulation of root architecture, we also enquired about autophagy's role during ARF19 turnover.
Western blot probing of *pARF19::ARF19-GFP/nph4-1 arf19-1* seedlings (Okushima *et al*, 2007)
treated with or without the vacuolar protease inhibitors Pepstatin A and E-64D for 24h revealed
substantial accumulation of ARF19-GFP when vacuolar proteases were inhibited(Fig EV1) . Together,
our results indicate that autophagy participates in the turnover of these essential LR formation
regulators.

Figure 1: ARF7 is degraded via autophagy. A) ARF7 western blots of protein extracts from Col-0, *atg5-1*, *atg2-1* and *nbr1-c1*
with or without addition of the proteasome inhibitor MG132. B) *ARF7* relative expression in given genotypes. *ARF7*
expression was normalized to *ACTIN2* and is normalized to Col-0 (set to 1). The experiment was repeated 3 times
(biological replicates) with similar results. C) GFP western blot of protein extracts from Col-0 or *atg2-1* expressing *ARF7-*
Venus. Values below each band represent the relative intensity ratio between ARF7 (A) or ARF7-Venus (C) and RuBisCO
large subunit (Rbc L) as stained with Ponceau S used as loading control. Ratios were normalized to untreated Col-0 NT (MS+
solvent) which was arbitrarily set to 1. Values provided are the average and SD of 5 (A) or 7 (C) biological replicates,
asterisk depict statistical significance from Col-0 NT according to a Holm-Sidak test (A) or two-tailed t-test (C) (* 0.05; **
0.01; **** 0.0001). D-E) Confocal images of root maturation zone (top panel) and meristematic zone (bottom panel) in
Col-0 (D) or *atg2-1* (E) expressing *ARF7-Venus*. (F-I) Fluorescence intensity quantification of ARF7-Venus subcellular
localization in given lines. The values presented for ARF7-Venus fluorescence intensity in *atg2-1* background in the
maturation zone (F-G) or meristematic zone (H-I) were normalized to the values in Col-0 background (which was set to 1).
The outline of the violin plots represents the probability of the kernel density. Dotted lines represent interquartile ranges
(IQR), with the thick horizontal line representing the median; whiskers extend to the highest and lowest data point. Results
were obtained from three independent experiments with at least 11 plants per condition being analysed in total. At least

40 nuclei and cytoplasmic regions were measured per genotype/tissue. Asterisks mark statistical significance to NT
according to a Mann-Whitney U-test (****P < 0.0001).

Fig EV1: Autophagy inhibition promotes ARF19 accumulation. GFP western blot of 14-day-old *pARF19::ARF19-GFP/nph4-1*
arf19-1 seedlings grown in MS or MS supplemented with vacuolar protease inhibitors E-64D and Pepstatin A for 24h.
Values below each band represent the ratio between ARF19-GFP and the loading control (LC) normalized to NT (MS+
solvent), which was set to 1. Image is representative of 3 biological replicates; asterisk depict statistical significance from
NT according to a Man—Whitney U-test (P≤0.01)

ARF7 *interacts with* NBR1 *and* ATG8a *in planta*

To further substantiate our claim that autophagy participates in the targeted degradation of ARF7
we performed protein-protein interaction studies. For this purpose, we used *Arabidopsis* stable
transgenic lines expressing ARF7-Venus and performed coimmunoprecipitation (Co-IP) experiments.
The experiment revealed that NBR1 (Fig 2A) but not GOT1-YFP, co-immunoprecipitated with ARF7-
Venus; this is supported by ATG8a's enrichment in the same conditions (Fig 2B). Moreover, a web-
based analysis (Jacomin *et al*, 2016) identified several potential ATG8-interaction motifs (AIM) in
ARF7, supporting our findings. As a cautionary note, we were unable to detect ARF7-Venus in the
input of both Co-IPs which can likely be explained by the fact that ARF7-Venus is under the control of
its native promoter and thus lowly expressed. We also performed multi-colour Bimolecular
fluorescence complementation (mcBiFC) analysis with a system allowing simultaneous interaction
studies of several proteins (Gehl *et al*, 2009) to transiently express Venus[N]-ARF7 with SCFP[N]-ATG8a
and either SCFP[C]-NBR1 or SCFP[C]-TSPO, (TRYPTOPHAN-RICH SENSORY PROTEIN, an autophagic cargo
adaptor for aquaporins, Hachez *et al*, 2014) in *Nicotiana benthamiana* leaves. As expected, SCFP[C]-
NBR1 and SCFP[N]-ATG8a were able to reconstitute CFP to produce distinct fluorescence signal
localized in the nuclei and cytoplasm (Fig 2C). Moreover, Venus[N]-ARF7 and SCFP[C]-NBR1 produced
GFP signal, which was localized to the cytoplasm and nuclei, in many instances co-localizing with
NBR1-ATG8a CFP signal (Fig 2C). In contrast, SCFP[C]-TSPO did not produce GFP signal when co-
expressed with Venus[N]-ARF7. In combination, these results indicate that ARF7 interacts with

autophagic components and cumulative with data from Fig 1, indicate that this TF is degraded via

NBR1-mediated selective autophagy.

Figure 2 ARF7 associates with autophagic components NBR1 and ATG8a. **A, B)** ARF7 and ATG8/NBR1 association was

tested by Coimmunoprecipitation (Co-IP) assay. Protein extracted from *Arabidopsis* lines expressing *pARF7::gARF7-Venus*

or free *YFP* were immunoprecipitated with anti-GFP antibody and subsequently probed with antibodies raised against

NBR1 or ATG8a. **C)** Bimolecular fluorescence complementation experiments shows oligomerization of ARF7, NBR1 and

ATG8a in *N. Benthamiana*. Images were taken 48 hpi. cCFP-TSPO was used as control. Insets display close-ups of examples

of reconstitution of GFP and CFP signal, which are also indicated by arrows. Etched boxes in GFP and CFP channels indicate

the place of the insets in those channels. Scale bar: 10 μM

ARF7 co-localization with autophagosomes follows a daily rhythm and is triggered by auxin.

Having established that autophagy regulates ARF7 abundance, it would be expected that this TF co-

localize with autophagosome-like structures. To assess this, we crossed ARF7-Venus to mCherry-

ATG8a and mCherry-NBR1 reporter lines (Svenning *et al*, 2011) and used this for live cell imaging. As

expected, both autophagy-related proteins localized to cytoplasmic foci, in many instances co-

localizing with ARF7-Venus (Fig 3A). Moreover, mCherry-ATG8a also co-localized with ARF7-Venus in

the nucleus (Fig 3A). It should be noted that the co-localization of these autophagic proteins with

ARF7 in cytoplasmic foci is mainly observable in the root's maturation zone; in younger tissue ARF7

primarily localizes in the nucleus but some instances of cytoplasmic co-localization are also

observable (Fig 3A & B). Our data is in agreement with previous reports demonstrating that ARF7

partitioning is dependent on developmental stage of the tissue (Orosa-Puente *et al*, 2018; Powers *et*

al, 2019).

ARF7's nucleo-cytopartitioning has been ascribed to liquid-liquid phase separation (LLPS) (Powers *et*

al, 2019); likewise, autophagosome formation sites and protein cargo triage (Kirkin *et al*, 2009;

Zhang *et al*, 2009; Fujioka *et al*, 2020) are also dependent on LLPS. To address the nature of these

cytoplasmic ARF7 & ATG8 foci, we incubated seedlings expressing mCherry-ATG8 and ARF7-Venus

with the LLPS inhibitor 1,6-hexanediol (Wang *et al*, 2021; Xie *et al*, 2021) and tracked the stability of

those foci over time. Most of the foci in which ARF7-Venus and mCherry-ATG8a co-localized were

sensitive to 1,6-hexanediol treatment (Fig EV2), suggesting that these structures have liquid-like

properties. Interestingly, some of these co-localizing events were insensitive to 1,6-hexanediol

treatment, even after prolonged exposure (*i.e.,* more than 30 min), indicating that those assembles

have solid-like properties. Our data is both in agreement with findings for ARF7 (Powers *et al*, 2019)

and studies in mammalians showing that NBR1 (and p62) is involved in condensate formation during

autophagic degradation of substrates (Turco *et al*, 2021).

Figure EV2 ARF7 and ATG8 condensates have both liquid and solid properties. *pARF7::gARF7-Venus* x *mCherry ATG8a* plants were treated with the LLPS inhibitor 1,6-Hexanediol and imaged at given time points. Representative pictures were taken in the maturation zone of 10–14-day old *Arabidopsis* seedlings. Leftmost Merge column represent the merger of YFP and mCherry channels, while the rightmost also includes the T-PMT channel (Brightfield). Scale bar 10 µM.

While our data demonstrates that autophagy plays a clear role in controlling ARF7's abundance, the factors/conditions leading to ARF7 turnover are unclear. To tackle this, we decided to treat *ARF7 - Venus x mCherry ATG8a* seedlings with auxin or cytokinin treatment; our rationale being two-fold: a) auxin (NAA) and cytokinin (Kinetin) often act in contrast to each other during many developmental processes (Du & Scheres, 2018); b) we have recently shown that plants rapidly engage autophagy to remodel the proteome upon perception of wide range of treatments, among them auxin and cytokinin (Rodriguez *et al*, 2020). We started by imaging the maturation zone, as ARF7 cytoplasmic condensates are mostly observed in this region. We noticed that the co-localization of ARF7-Venus with mCherry-ATG8a varied during the time of the day, *i.e.,* there is a significant increase (45% to 85%, P<0.05) in ARF7 and ATG8a co-localization when comparing seedlings imaged during Zeitgeber time (ZT, time after the lights are switched on) ZT0 vs ZT6, respectively. This is consistent with previous reports (Perianez-Rodriguez *et al*, 2021) showing that ARF7 expression follows a periodic fluctuation of 4-6h in the oscillation zone. Given this, we decided to use ZT0 as the timepoint for subsequent treatments. Next, we incubated seedlings with vacuolar protease inhibitors (Pepstatin A and E-64d) which led to a dramatic increase in ARF7-Venus co-localization with mCherry-ATG8a (Fig 3C & D). This both supports our findings with *ARF7-Venus* x *atg2-1* (Figure 1) and also the proposition that ARF7 condensate formation can be linked to autophagic degradation. Similarly, NAA treatment promoted a significant increase of ARF7-Venus co-localization with mCherry-ATG8 positive foci in the cytoplasm (Fig 3 C &D) while Kinetin did not lead to increased ARF7-Venus co-localization with mCherry-ATG8a (P>0.05). Altogether, our results suggest that ARF7 cytoplasmic localization might be part of a proteome remodelling strategy which is mediated, at least in part, via autophagy.

Figure 3 ARF7 co-localization with autophagosomes displays rhythmic variation and is enhanced by NAA treatment. **A)** Co-localization of ARF7-Venus with mCherry-ATG8a (nuclei and/or cytoplasmic foci) or mCherry-NBR1 (cytoplasmic foci) in young (meristem/elongation) or maturation zone. Representative pictures were taken in the differentiation zone of 10–14-day old *Arabidopsis* seedlings. Scale bar 10 μM. **B)** Quantification of the co-localization of ARF7-Venus with mCherry-ATG8a or NBR1 under conditions depicted in **A)**; samples followed by the same letters are not statistically different (Dunn's test, P≤0.0001). **C)** ARF7-Venus co-localization with autophagosomes under selected treatments: No Treatment (NT), E64d + Pepstatin A (vacuolar protease inhibitors), NAA (1μM NAA for 30 min) and Kinetin (1μM Kinetin for 30 min). Scale bar 10 μM. **D)** Quantification of the co-localization of ARF7-Venus with mCherry-ATG8a under conditions depicted in **C)**. The outline of the violin plots represents the probability of the kernel density. Dotted lines represent interquartile ranges (IQR), with the thick horizontal line representing the median; whiskers extend to the highest and lowest data point. Results were obtained from three independent experiments with at least 8 plants per condition. Asterisks mark statistical significance to NT (MS+ solvents) according to a Wilcoxon-rank test (**0.01; **** 0.0001).

Autophagy regulates rhythmic changes to ARF7 *protein abundance but is not required for PBS*

establishment

Intrigued by our results showing that co-localization between ARF7 and ATG8a follows rhythmic

variation and knowing that *ARF7* expression also shows a similar pattern in the oscillation zone (OZ,

Perianez-Rodriguez *et al*, 2021), we decided to explore the impact of autophagy in the natural

variation of ARF7's abundance. Therefore, we performed western blots against ARF7 in Col-0 and

atg mutants' seedlings during a 12h period. As seen in Fig 4A, ARF7 levels increased from ZT0 to ZT9,

followed by progressive decline up to ZT12, which is in agreement with the abovementioned pattern

for *ARF7* transcript oscillation in the OZ. Moreover, these results likely explain the variation in co-

localization of ARF7-Venus with autophagosomes at different timepoints (Fig 3): condensation is

dependent on target concentration (Mcswiggen *et al*, 2019) and therefore more likely observable

under high ARF7 levels (ZT6). Our findings also illustrate that ARF7 abundance display rhythmic

changes in the whole plant, which is consistent ARF7's expression profile (*e.g.,* Rosado *et al*, 2012;

Lee *et al*, 2019) and with ARF7's role during hypocotyl gravi- and phototropism (Wang *et al*, 2020).

In marked contrast, the rhythmic variation in ARF7's abundance was lost (P≤0.05) when autophagy

was disrupted (Fig 4B-D). Moreover, the transcript levels of *LBD16*, which is directly regulated by

ARF7, also lost its rhythmic fluctuation in *atg* mutants (Fig 4E). Interestingly, *LBD16* levels are

substantially higher in *atg2-1* than Col-0, which could be related to the overall higher levels of ARF7

seen in *atg* mutants. Taken together, these results demonstrate the importance of autophagy in

regulating ARF7-dependent rhythm.

Figure 4: ARF7 abundance displays whole-plant rhythmic variation and is modulated by autophagy. **A-D)** ARF7 western
blots from proteins extracted from Col-0 (**A**); *atg2-1* (**B**); *atg5-1* (**C**) or *nbr1-c1* (**D**) at given Zeitgeber time (ZT0-ZT12).
Values below each band represent the relative intensity ratio between ARF7 and RuBisCO large subunit (Rbc L) as stained
with Ponceau S used as loading control. Ratios were normalized to untreated Col-0 (NT) which was arbitrarily set to 1.
Values given are the mean ± SD of 3 independent experiments, Col-0 was statistically significant (P≤0.05) from all
autophagy mutant, according to a Paired-T-test. **E)** Relative gene expression oscillation of *LBD16* in Col-0 and *atg2-1* at
given timepoints. *LBD16* expression was normalized to *ACTIN2* and set to 1 for each genotype at timepoint 0h. **F)**
Prebranch site (PBS) in *DR5::GFP* and *DR5::GFP x atg5-1*. PBS of 5-day post germination, vertically grown seedlings were
scored with a stereo microscope. The outline of the violin plots represents the probability of the kernel density. Dotted
lines represent interquartile ranges (IQR), with the thick horizontal line representing the median; whiskers extend to the
highest and lowest data point. Results were obtained from 4 independent experiments with a total of at least 80 plants per
genotype. Mann-Whitney was used for the statistical test. **G)** ARF7-Venus and mCherry-ATG8a co-localization in root young
tissue treated with MS + solvent (top panel) or MS+NAA (bottom panel) for 30 min. **H)** Quantification of the co-localization
of ARF7-Venus with mCherry-ATG8a under conditions depicted in **G)**. The outline of the violin plots represents the
probability of the kernel density. Dotted lines represent interquartile ranges (IQR), with the thick horizontal line
representing the median; whiskers extend to the highest and lowest data point. Results were obtained from three
independent experiments with at least 8 plants per condition. Asterisks mark statistical significance to NT according to a
Mann-Whitney test (**** 0.001).

ARF7 oscillation has been established as an essential step during PBS determination as *arf7-1*
mutants display abnormal PBS patterning (Perianez-Rodriguez *et al*, 2021). Because ARF7 abundance
does not seem to change when autophagy is impaired, we questioned whether PBS establishment
could be affected in *atg* mutants. To track this, we compared PBS numbers in Col-0 and *atg5-1* lines
expressing *DR5::GFP* with a stereo microscope. As seen in Fig 4F, there was no major differences in
PBS count (P>0.05) between both genotypes, being that the total number of PBS we report are
comparable to those available in the literature for Col-0 (Moreno-Risueño *et al*, 2010). Taken
together, our results suggest that while autophagy is necessary for ARF7's rhythmic fluctuation,
deficiencies in the degradation process do not produce major interference with PBS establishment.

Autophagy deficiency negatively impacts auxin mediated proteome remodelling

Intriguingly, our results could suggest that ARF7 oscillation might not be necessary for PBS
establishment, which disagrees with previous reports (Perianez-Rodriguez *et al*, 2021). Several
explanations could justify these seemingly contradictory differences. Firstly, *ARF7*'s oscillation
relevance during PBS establishment was inferred from analysing *arf7-1*(Perianez-Rodriguez *et al*,
2021) in which there is no ARF7 expression. On the other hand, in *atg* mutants, ARF7 is expressed
but does not oscillate due to defective proteostasis (Fig 1-4). It could thus be possible that while
ARF7 is undoubtedly important for correct PBS establishment, the oscillation of this protein might
not be essential for the process. Alternatively, *ARF7's* levels in the OZ zone could be regulated by
different mechanisms than in the rest of the plant; this would be consistent with the
aforementioned nucleo-cytopartitioning of ARF7 (Fig 3, Powers *et al*, 2019). To analyse this, we
decided to investigate auxin's effect on ARF7 and ATG8a co-localization but now focusing on
younger tissue. Like seen for the maturation zone (Fig 3C-D), NAA caused ARF7-Venus to massively
localize in cytoplasmic foci (Fig 4G); indicating that auxin promotes ARF7 condensations. Likewise,
we also observed a massive increase in mCherry-ATG8 foci (Fig 4G), which is consistent with our
previous reports showing that quick bursts of auxin lead to rapid increase in autophagic activity
(Rodriguez *et al*, 2020). Surprisingly, these dramatic increases of ARF7 and ATG8a cytoplasmic foci
only produced a modest, albeit significant, increase in co-localization (Fig 4G & H), which contrasts
with our maturation zone data (Fig 3). Given that a) autophagy has a well characterized role in
aggregate degradation (*e.g.*, Lim *et al*, 2015); b) the root meristem has been shown to display zone
specific chaperonin expression to prevent protein aggregation (Llamas *et al*, 2021); c) ARF7
condensates have liquid and solid properties (Fig EV2, Powers et al 2019); d) ARF7 condensates are
more prominent in older tissue (Fig3, Powers *et al*, 2019); our results here likely reveal a mechanism

by which auxin treatment leads to a rapid activation of autophagy to modulate ARF7 levels though

this mainly occurs in older tissue, where aggregation is more prone to occur.

In sum, these experiments reveal that while the oscillation of ARF7 abundance seems to be a whole

plant phenomenon, the means by which this regulation occurs might be zone specific. Likewise, this

apparent zone-specific regulation of ARF7 turnover also reconciles our findings with those of

Perianez-Rodriguez et al, 2021, namely the relevance of *ARF7* oscillation to PBS establishment and

the normal PBS establishment in *atg* mutants.

Given that NAA induced rapid ARF7 localization to cytoplasmic foci, we wondered if this would have

any effect in ARF7 abundance. Hence, we checked ARF7 levels by western blot with protein extracts

from Col-0 and *atg2-1* seedlings treated with NAA for 1h at ZT0 (Fig EV3). Our data revealed that

NAA produced a mild but significant decrease (P≤0.01) in ARF7 levels in Col-0, which is consistent

with the NAA-induced increase co-localization between ARF7 and autophagic components (Fig 3 &

4). In combination, our results (Fig 3-4, EV3) support a mechanism by which auxin promotes ARF7's

condensation to modulate ARF7 levels, preventing excessive nuclear ARF7 and consequent

exaggerated responses. In contrast to previous reports (e.g., Wang *et al*, 2005; Perianez-Rodriguez *et*

al, 2021), our data suggests that auxin treatment induces changes to ARF7 abundance; which has

also been reported at the transcript level after 2h of IAA treatment (Biswas *et al*, 2019). These

differences might be easily conciliated due to different experimental set ups among reports: longer

auxin treatments (3h-16h) and different tissues (*i.e.,* root's OZ or protoplast) were used in those

reports (Perianez-Rodriguez *et al*, 2021; Wang *et al*, 2005) as opposed to the experimental setup

used here. This is supported by our data, which indicates that tissue specific mechanisms and time of

the day appear to be important factors during ARF7 turnover and by predicted buffering

mechanisms of ARF7 signalling (Perianez-Rodriguez *et al*, 2021). Concerning *atg2-1* seedlings, NAA

treatment did not lead to significant changes in ARF7 levels (Fig EV3) which might not be surprising

given that ARF7 levels are already elevated when autophagy is impaired and that other

compensatory mechanisms (*e.g.*, proteasome) could be upregulated to compensate for defective

autophagy.

Figure EV3 NAA-induced changes to ARF7s abundance is lost in *atg* mutant. ARF7 western blot from proteins extracted
from Col-0 or *atg2-1* treated with MS + solvent or MS supplemented with NAA (1µM, 60 min), at Zeitgeber time 0 (ZT0).
Values below each band represent the relative intensity ratio between ARF7 (**A**) and RuBisCO large subunit (Rbc L) as
stained with Ponceau S used as loading control. Ratios were normalized to each genotype's NT sample (MS+ solvent),
which was set to 1. Values are mean ± SD of 5 biological replicates. Statistical analysis was done using a Holm-Sidak test (*
0.05; *** 0.001).

Accumulation of ARF7 in the cytoplasm, which we have shown is impacted by autophagy (Fig 1 D-I;
Fig3), has been linked to decreased auxin-dependent responses (Powers *et al*, 2019). Moreover,
atg2-1 is unable to execute ARF7 turnover upon rapid auxin treatment (Fig EV3) and we wondered if
this could be indicative of more broad implications, namely altered proteome remodelling during
rapid auxin enacted responses. To evaluate this, we gave a short auxin treatment (30 min) to Col-0
and *atg2-1* seedlings and performed Mass Spectometry (MS). Despite the rapid nature of the
treatment with NAA, proteome remodelling was engaged in both genotypes (Fig 5A), and as
expected, more proteins were downregulated in Col-0 than *atg2-1* (167 vs 94, Fig 5B). This
difference in auxin-induced downregulated proteins between the genotypes is consistent with the
loss of a major proteostasis pathway in the *atg2-1* background. There were also differences, albeit
smaller, in what concerns auxin-induced protein accumulation, 146 for Col-0 vs 114 for *atg2-1*.
Surprisingly, only around 5% of the proteins with differential regulation upon NAA treatment were
similarly down (or up) regulated in both genotypes. These results indicate that while *atg2-1*
responds to NAA, the nature of this response is substantially different to that of Col-0, both
supporting and expanding our previous findings (Rodriguez et al 2020) showing defective NAA-
induced reprogramming in *atg* mutants.

Autophagy facilitates LR formation

As our results suggest that autophagy's contribution to ARF7 turnover might be more prominent in
mature tissue, we wondered if accumulation of ARF7 in *atg* mutants could have an impact in other
ARF7-dependent phenotypes. Besides PBS, ARF7 regulatory role during LR initiation is also well
established (*e.g.,* Okushima *et al*, 2007; Goh *et al*, 2019); consequently, we examined *atg* mutants'
ability to produce LRs by transplanting 5-day old seedlings to vertical plates and scored LRs after 9
days (Fig 5 C-D). As shown in Fig 5, *atg* mutants have a significantly lower LR density when compared
to Col-0. Because we have seen that lack of autophagy leads to condensation of ARF7 (Fig 3) which
has been linked with decreased auxin responses (Powers *et al*, 2019) and because NAA treatment
did not produce major changes to ARF7 abundance in *atg2-1* seedlings (Fig EV3), we questioned if

atg mutants might have dampened auxin responses. To ascertain this, we germinated Col-0 and *atg*

mutants and after 5 days, transferred them to vertical MS plates with or without NAA (0.1 µM) and

counted LRs 9 days after germination. As seen in Fig 5 C-D, NAA treatment caused an increase in LR

density for all genotypes, but it did not rescue the differences observed between Col-0 and *atg*

mutants. These results are in agreement with our previous findings (Rodriguez *et al*, 2020) regarding

atg mutants' reduced ability to respond effectively to callus inducing media (rich in auxin) to form

callus, a process resembling LR initiation.

Figure 5 Autophagy deficient mutants display impaired proteome remodelling upon auxin treatment and show decreased

LR formation capacity. **A)** Volcano plot displaying pairwise protein content changes in Col-0 or *atg2-1*, 30 min after Mock

(MS+ EtOH) or NAA treatment (MS+ NAA) with a F.D.R. <0.05 using an unpaired t-test. **B)** Venn diagrams showing the

number of differentially expressed proteins overlapping between the datasets (F.D.R. < 0.05). **C-D)** LR density

quantification in Col-0, *atg2-1* and *atg5-1* grown in MS plates supplemented with Solvent (MS) or 0.1 µM NAA. The outline

of the violin plots represents the probability of the kernel density. Dotted lines represent interquartile ranges (IQR), with

the thick horizontal line representing the median; whiskers extend to the highest and lowest data point but no more than

±1.5 times the IQR from the box. Results were obtained from three independent experiments with at least 40 plants per

condition, asterisks mark statistical significance to Col-0 according to the t-test (*0.05; **** 0.0001). Scale bar: 1 cm

Discussion

Here, we show that autophagy modulates ARF7 and ARF19 turnover and how through the selective

cargo adaptor NBR1 (Fig 1 & 2). We and others (Jing *et al*, 2022) have shown that MG132 treatment

also causes ARF7 to accumulate, which suggests proteasomal activity also regulates this TF turnover.

However, care should be taken when inferring proteasomal contribution based solely on MG132-

derived data as this inhibitor produces several off-target effects like transcriptional upregulation

(Wang et al, 2008; Cho et al, 2013), interference with lysosomal proteases (Garrison & Bangs, 2020)

and autophagosome assembly (Ji et al, 2016). Nonetheless, as we see a synergistic effect of MG132

and autophagy inhibition in ARF7 accumulation (Fig 1), it is likely that both degradation pathways are

involved in regulating ARFs turnover. The recruitment of both degradation pathways to promote

ARF7 turnover is in line with the necessity to keep ARF7 output under tight control (Moreno-Risueño

et al, 2010; Perianez-Rodriguez et al, 2021) and with the fact that excessive activating ARFs lead to

dramatic developmental retardation due to exaggerated auxin program execution (Powers et al,

2019). Interestingly, we find that while atg mutants over accumulate ARF7, autophagy blockage

leads to increased ARF7 cytoplasmic condensates, which have been linked to low responsiveness to

auxin. Consequently, despite having higher ARF7 levels than Col-0, atg mutants show diminished

auxin response (Fig EV3, Fig5), LR formation (Fig 5) and callus formation (Rodriguez et al, 2020), a

process reminiscent of LR initiation. How does accumulation of ARFs equate to diminished auxin

responses? It is possible that autophagy inhibition leads to ARF7 accumulation which may induce

supersaturation (Mcswiggen et al, 2019) and cause ARF7 condensation in the cytoplasm (Fig 1 & 3).

Recent reports (Powers et al, 2019; Jing et al, 2022) have linked ARFs condensation with decreased

auxin responsiveness, therefore, the disruption of protein degradation pathways we report here

causes ARF7 levels to increase, promoting cytoplasmic aggregation and consequent reduced auxin

responsiveness. While our data provides an indication that the mobilization of ARF7 to cytoplasmic

foci might be part of a mechanisms aimed at adjusting ARF7 abundance and auxin responsiveness,

more experiments are needed to fully understand this process. For instance, (Jing et al, 2022)

recently showed that the loss of function of AFF1 (F-box protein) displays reduced auxin

responsiveness due to cytoplasmic hyper condensation and consequent nuclear depletion of ARFs.

Despite these dramatic effects, the aff1 mutants only display mild developmental phenotypes and

surprisingly seems able to produce LRs when supplemented with 2-4D; this contrasts with arf7 x

arf19 mutant's severe LR formation defects, even when cultured with auxin (Okushima et al, 2007).

Moreover, we report that ARF7 cytoplasmic foci do not co-localize with autophagosomes at the

same extent in young vs mature tissues, even after inducing ARF7 cytoplasmic condensation with a

short NAA treatment (Fig 3). A caveat to this interpretation is that in young tissue, NAA-induced

condensates might have greater liquid-like properties, hence the lower co-location with autophagic

components. Given that condensate solidification is a progressive process, it is possible with time,

ARF7 condensates in young tissue would increase solidity and increase their co-localization with
ATG8a/NBR1

In sum, while we provide robust evidence indicating that ARF7 cytoplasmic condensation is part of a
triage mechanism for its autophagic degradation, further clarification is needed to completely
uncover how this process is regulated in different root zones.

In contrast to PBS establishment, which is not grossly affected in *atg5-1* (Fig 4), LR formation seems
to be impaired in *atg* mutants (Fig 5). In the context of LR formation, there is evidence of autophagic
participation in a wide range of processes like cell death regulation of endodermal cells above LRPs
(Escamez *et al*, 2020) and potentially also during PIN's degradation (Kleine-Vehn *et al*, 2008).
Moreover, autophagy modulates proteome turnover induced by different hormones like auxin (Fig
5; Rodriguez et al 2020), cytokinin (Acheampong *et al*, 2020; Rodriguez *et al*, 2020) and ABA
(Rodriguez *et al*, 2020); all of which are known to influence LR formation. For these reasons,
deficiencies in such a transversal process like autophagy complicate result interpretation and
ascribing phenotypes to a specific function of autophagy. For instance, ABA is known to interfere
with LRs formation (de Smet *et al*, 2003) and consequently, misregulation of ABA signalling in *atg*
mutants (Rodriguez *et al*, 2020) could be a source of defective LR formation. However, we have
previously seen that *atg* mutants have difficulties forming callus (Rodriguez *et al*, 2020) a process
resembling LR initiation and because ABA does not affect callogenesis (Sugimoto *et al*, 2010), it is
plausible that LR deficiencies in *atg* mutants occur at an earlier stage than ABA-induced effects (LR
emergence). Nonetheless, because our results unequivocally show that autophagy specifically
impacts the turnover of the important LR regulators ARF7 and ARF19, it is undeniable that at the
very least, defective ARFs turnover in *atg* mutants must contribute to the decreased auxin responses
and LR formation we report here.

In conclusion, we have uncovered a novel role for selective autophagy during post embryonic organ
development through the turnover of the key developmental regulators ARF7 and ARF19 and
provided further insight into the intricate mechanisms behind ARFs fine-tuning and how this affects
root architecture.

Acknowledgements

This work was funded by a Danish Research Agency grant to E.R. (DFF1- 1032-00249B). We would

like to thank Tom Guilfoyle (anti-ARF7 antibody), Koki Yoshimoto (anti-ATG8a antibody), Malcolm

Bennett (*pARF7::gARF7-Venus*), Yasin Dagdas (*DR5::GFP* and *atg5-1 DR5::GFP*), Liwen Jiang (*NBR1-*

C1) and Hidehiro Fukaki (*pARF19::ARF19-GFP nph4-1 arf19-1*) for sharing material/reagents. Thanks

are due to Miguel Moreno-Risueño and Nuria Coll for insightful discussion. We would also like to

thank Suksawad Vongvisuttikun for technical assistance, the 2022 classes from the courses "Plant

Molecular Biology" & "Experimental Higher Model Organisms" (University of Copenhagen, Denmark)

for performing the ARF19-GFP & ARF7-Venus western blots, respectively. Confocal microscopy

imaging was performed using equipment from the Center for Advanced Bioimaging (CAB) Denmark,

University of Copenhagen. Mass spectrometry analyses were performed by the Proteomics Research

Infrastructure (PRI) at the University of Copenhagen (UCPH), supported by the Novo Nordisk

Foundation (NNF) (grant agreement number NNF19SA0059305).

Author contributions

Conceptualization: ER; Investigation: ER, EE, JA, JC, PC. Formal analyses: ER, EE, JA. Writing: ER, EE, JA

Visualization: ER, JA; Supervision: ER; Funding acquisition ER. All authors read the manuscript and

agreed with the findings reported.

Methods and Materials

Plant material and phenotypic characterization

Arabidopsis seedlings were grown on solid, half-strength MS media (0.8% agar, 1% sucrose at 5.7 pH) and kept at 21 C° with a photoperiod of 16 h (120 µE/m^2/s). Seeds were sterilized with 1.3% bleach followed by 70% alcohol and then rinsed three times with sterilized water. The following lines were used throughout this study; Col-0, *atg2-1* (SALK_076727), *atg5-1* (SAIL_129_B07), *nbr1-c1* (Ji *et al*, 2020), mCherry ATG8a & mCherry NBR1 (Svenning *et al*, 2011), pARF7::gARF7 Venus (Orosa-Puente *et al*, 2018), pARF19::ARF19-GFP/*nph4-1 arf19-1* (Okushima *et al*, 2007), DR5::GFP and DR5::GFP *atg5-1* (Gift from Yasin Dagdas, GMI Vienna, Austria).

For the time course experiments, seedlings were acclimated in liquid half-strength MS media for at least 2 days before sampling. Sampling was done just before the chamber light was turned on ZT0, and then at indicated times. For LR and PBS scoring, seedlings were germinated on half-strength MS media and set to grow vertically for the indicated time, using a Leica M165 FC stereomicroscope. PBSs were counted after 7 days using GFP filter to visualize DR5 driven, GFP signal. Per genotype, at least 70 plants were analyzed derived from 6 independent experiments. LR's were counted at day 9 as described by (Dubrovsky & Forde, 2012), and only LR's that had at least reached stage IV were included. Per genotype/condition at least 40 plants were counted derived 6 independent experiment. All genotypes were blinded during counting in both assays to prevent potential bias.

Chemical treatments

For confocal microscopy and western blot assays, seedlings were acclimated in liquid half-strength MS media for 2 days acclimation before imaging. For hormone treatment effect on the co-localization of ARF7 with ATG8 & NBR1, samples were either treated with NAA (1 µM), Kinetin (1 µM), or MS for the given time. Chemical inhibition of autophagy was done by pre-treating seedlings with the vacuolar protease inhibitors E-64d (1:3000, stock 10 µM) and Pepstatin A (1 µM) 16 h prior imaging or western blotting. For inhibition of condensates in the mature regions of the root, roots were imaged just before and directly after treatment with condensate inhibitor 1,6-hexanediol (10%) as described by(Wang *et al*, 2021). Images were taken at 5 min interval for 30 min. Confocal imaging of the seedlings was taken with LSM700 Zeiss confocal microscope, at 20x or 63x water objective. Images were analyzed using Zen 2012 (Zeiss).

For protein analysis of ARF7 in *atg* mutants upon proteasomal blockage, samples were treated with or without MG132 (100 µM) for 16 h before harvest.

Protein extraction

Proteins were extracted using 3xSDS buffer (30% glycerol, 3% SDS, 94 nM Tris, pH 6.8, Bromophenol
blue, Complete ultra-tablets EDTA-free protease inhibitor cocktail (Roche), 50mM DTT), then
samples were centrifugated for 20 min at 4°C 14000 rpm and the supernatant was moved into a new
tube. Samples were denatured at 95°C for 5 min before loaded it for SDS-PAGE.

SDS-PAGE and western blotting

Protein samples were separated in a 4-20% pre-casted gel (Bio-Rad) and transferred overnight onto
nitrocellulose membrane (GE Healthcare). Membranes were blocked for 1 h with 5% skim milk
(Merck) in TBST [50 mM Tris-HCL, pH 7.5 150 nM NaCl, 0.1% Tween-20 (Sigma)] and incubated 2 h at
room temperature or overnight (4°C) with primary antibodies: anti-ARF7 (Wang et al, 2005), anti-
NBR1 (AS14 2805, Agrisera; 1:5000); anti-GFP (TP401 AMSBio, 1:1000); anti-ATG8a (Yoshimoto *et al*,
2004) Membranes were washed, followed by incubation in anti-rabbit HRP conjugate (Promega;
1:5000). Home-made chemiluminescent substrate (Mruk & Cheng, 2011) was applied before
detection. Bands were quantified using ImageJ and normalized to loading controls.

Sample preparation for proteomic analysis

Tissue extracts (100 µg) were diluted in digestion buffer (0.5% SDC in 50 mM TEAB), heat-treated for
5 min at 95 °C, and prepared by a modified filter-aided sample preparation (FASP) protocol (65). In
brief, tissue extracts were transferred to 0.5 ml (tilted) spin filters (Amicon), centrifuged at 14,000g
for 15 min, and reduced and alkylated by addition of digestion buffer containing 1:50 (v:v) tris(2-
carboxyethyl)phosphine (0.5 M, Sigma) and 1:10 (v:v) 2-chloroacetamide (Sigma) for 30 min at 37 °C.
Samples were digested in fresh digestion buffer containing 1 µg Typsin/LysC mix (Promega) and
0.01% ProteaseMAX (Promega) O/N at 37 °C. Digested samples were desalted using stage-tips
containing styrene divinylbenzene reversed-phase sulfonate material (SDB-RPS; 3 M).

The mesenteric artery-based library was generated using a pooled digested and stage-tipped sample
from the 12-week-old SHRs and WKYs that was fractionated. A high-pH reverse-phase peptide (HpH)
fractionation kit (Pierce, Thermo Scientific) was used to create the 15 fractionations.

Data acquisition by liquid chromatography–mass spectrometry (LC-MS)

Peptides were separated on 50 cm columns packed with ReproSil-Pur C18-AQ 1.9 µm resin (Dr
Maisch GmbH). Liquid chromatography was performed on an EASY-nLC 1200 ultra-high-pressure
system coupled through a nanoelectrospray source to an Exploris 480 mass spectrometer (Thermo
Fisher Scientific). Peptides were loaded in buffer A (0.1% formic acid) and separated applying a

nonlinear gradient of 5 to 65% buffer B (0.1% formic acid, 80% acetonitrile) at a flow rate of 300

nl/min over 100 min. Spray voltage was set to 2400 V. Data acquisition switched between a full scan

(120,000 resolution, 45 ms max. injection time, AGC target 300%) and 49 DIA scans with isolation

width of 13.7 m/z and windows overlap of 1 m/z spanning a precursor mass range of 361 to 1033

m/z (15,000 resolution, 22 ms max. injection time, AGC target 1000%). Normalized collision energy

was set to 27. All statistical analysis of LFQ derived protein expression data was performed using the

automated analysis pipeline of the Clinical Knowledge Graph (Santos *et al*, 2022).

Data availability

565   The mass spectrometry data from this publication has been deposited to the PRIDE archive (Perez-

566   Riverol et al 2022; https://www.ebi.ac.uk/pride/archive/) and given the identifier [PXD042834]

Vector construction and Plant Transformation for BiFC

To create pDEST-ATG8a-SCYNE, pDEST-NBR1-SCYCE, pDEST-ARF7-VYNE and pDEST-TSPO-SCYCE, we

PCR amplified (CloneAMP HIFI PCR premix, TAKARA) the coding sequences of *ATG8a*, *NBR1*, *ARF7*

and TSPO flanked recognition sequence for Gateway (Thermofischer) or In-Fusion (Clontech) cloning.

ATG8a, *NBR1 and ARF7's* PCR products were cloned into pENTR (Life Technologies) and

subsequentially transferred (LR clonase, Invitrogen) into the Gateway-compatible BiFC vectors (Gehl

C *et al*, 2009) pDEST-^{GW} SCYNE, pDEST-^{GW}SCYCE, pDEST-^{GW}VYNE. *TSPO* PCR amplicon was directly

cloned into pDEST-^{GW}SCYCE using In-Fusion cloning. *Agrobacterium* strain GV3101 containing

constructs were grown in Yeast-extract peptone media supplied with the respective antibiotics

overnight. Cultures were spun and resuspended in 10mM MgCl$_2$, 10 mM MES-K and 100 µM

acetosyringone to the density of 600nm = 0.15-0.2 per construct and mixed 1:1 and left at room

temperature for 2 h, followed by infiltration in 3 weeks old *N. Benthamiana*. Pictures were taken 48

hpi using a Zeiss LSM 700 confocal microscope.

Immunoprecipitation

Immunoprecipitation assay was described previously (Zuo *et al*, 2022). In brief, 1 g of 14-day old

seedlings were grinded with mortar and pestle and 2 ml of IP extraction buffer (50mM Tris-HCl pH

7.5, 150mM NaCl, 5% glycerol 1mM EDTA, 0.1% NP40, 10mM DTT, protease inhibitor (mini complete

ultra, EDTA-free) was added, followed by centrifugation at 4℃ 13000 rpm, 20 min. The supernatant

was incubated with 20 µl GFP trap beads (Chromotek) rotating for 2-4-hours at 4℃. The beads were

washed 3-4 times with washing buffer (20 mM Tris-HCl pH 7.5, 0.15 M NaCl, 0.1% NP40). 30 µL of

3xSDS was added to the beads and heat up 5 min at 95 ℃ before proceeding for SDS-PAGE.

RNA extraction

RNA was extracted as described previously (Zuo *et al*, 2022) using TRIzol reagent (Thermo Scientific),
treated with DNase (Thermo Scientific) and was reverse transcribed into cDNA using Reverse Aid kit
(Thermo Scientific) as described by the manufacturer. qPCR assays were executed on the
Quantstudio 1 (Applied Biosystems) system using SYBR Green master mix (ThermoFisher), *ACT2* was
used as internal control. All primers used for qPCR analysis are listed in Supplementary table 1. The
experiment was repeated three times.

Supplementary table 1

Cloning primers	
ARF7 Forward primer	5'-CACCATGAAAGCTCCTTCATCAA-3'
ARF7 Reverse primer	5'-TCACCGGTTAAACGAAGTGGC-3'
NBR1 Forward primer	5'-CACCATGGAGTCTACTGCTA-3'
NBR1 Reverse primer	5'-TCAAGCCTCCTTCTCCCCT-3'
TSPO Forward primer	5'-GCAGGCTCCGCGGCCATGGATTCTCAGGACATCAGA-3'
TSPO Reverse primer	5'-AGCTGGGTCGGCGCGTCACGCGACTGCAAGCTTTAC-3'
ATG8a Forward	5'-CACCATGATCTTTGCTTGCTTGA-3'
ATG8a Reverse	5'-TCATCAAGCAACGGTAAGAGAT-3'
qPCR primers	
ACT2 Forward primer	5'-CTTGTTCCAGCCCTCGTTTGTG-3'
ACT2 Reverse primer	5'-CCTTGGAGATCCACATCTGCTG-3'
ARF7 Forward primer	5'-CAAGGTCACAGTGAGCAAGTCG -3'
ARF7 Reverse primer	5'-TGTGGAGCATGCATATGAGCTTGG -3'
LBD16 Forward primer	5'-TCCATGATCGATGTGAAGCTGTCG-3'
LBD16 Reverse primer	5'-TGTGATTGCAAGAAAGCCACCTG-3'

References

Bassham DC, Laporte M, Marty F, Moriyasu Y, Ohsumi Y, Olsen LJ & Yoshimoto K (2006) Autophagy in development and stress responses of plants. *Autophagy* 2: 2–11

Biswas MS, Fukaki H, Mori IC, Nakahara K & Ichi Mano J' (2019) Reactive oxygen species and reactive carbonyl species constitute a feed-forward loop in auxin signaling for lateral root formation. *The Plant Journal* 100: 536–548

Chen L, Liao B, Qi H, Xie LJ, Huang L, Tan WJ, Zhai N, Yuan LB, Zhou Y, Yu LJ, *et al* (2015) Autophagy contributes to regulation of the hypoxia response during submergence in Arabidopsis thaliana. *Autophagy* 11: 2233–2246

Cho KS, Kwon KJ, Jeon SJ, Joo SH, Kim KC, Cheong JH, Bahn GH, Kim HY, Han SH, Shin CY, *et al* (2013) Transcriptional Upregulation of Plasminogen Activator Inhibitor-1 in Rat Primary Astrocytes by a Proteasomal Inhibitor MG132. *Biomol Ther (Seoul)* 21: 107

Deb S, Sankaranarayanan S, Wewala G, Widdup E & Samuel MA (2014) The S-Domain Receptor Kinase Arabidopsis Receptor Kinase2 and the U Box/Armadillo Repeat-Containing E3 Ubiquitin Ligase9 Module Mediates Lateral Root Development under Phosphate Starvation in Arabidopsis. *Plant Physiol* 165: 1647–1656

Desideri E, Filomeni G & Ciriolo MR (2012) Glutathione participates in the modulation of starvation-induced autophagy in carcinoma cells. *Autophagy* 8: 1769–1781

Du Y & Scheres B (2018) Lateral root formation and the multiple roles of auxin. *J Exp Bot* 69: 155–167

Dubrovsky JG & Forde BG (2012) Quantitative analysis of lateral root development: Pitfalls and how to avoid them. *Plant Cell* 24: 4–14

Escamez S, André D, Sztojka B, Bollhöner B, Hall H, Berthet B, Voß U, Lers A, Maizel A, Andersson M, *et al* (2020) Cell Death in Cells Overlying Lateral Root Primordia Facilitates Organ Growth in Arabidopsis. *Current Biology* 30: 455-464.e7

Fujioka Y, Alam JM, Noshiro D, Mouri K, Ando T, Okada Y, May AI, Knorr RL, Suzuki K, Ohsumi Y, *et al* (2020) Phase separation organizes the site of autophagosome formation. *Nature* 578: 301–305

Garrison P & Bangs JD (2020) p97 Inhibitor CB-5083 Blocks ERAD in Trypanosoma brucei. *Mol Biochem Parasitol* 239

Gehl C, Waadt R, Kudla J, Mendel R & Hansch R (2009) New GATEWAY vectors for high throughput analyses of protein-protein interactions by bimolecular fluorescence complementation. *Mol Plant*: 1051–1058

Goh T, Toyokura K, Yamaguchi N, Okamoto Y, Uehara T, Kaneko S, Takebayashi Y, Kasahara H, Ikeyama Y, Okushima Y, *et al* (2019) Lateral root initiation requires the sequential induction of transcription factors LBD16 and PUCHI in Arabidopsis thaliana. *New Phytologist* 224: 749–760

Hachez C, Veljanovski V, Reinhardt H, Guillaumot D, Vanhee C, Chaumont F & Batoko H (2014) The Arabidopsis abiotic stress-induced TSPO-Related protein reduces cell-surface

expression of the aquaporin PIP2;7 through protein-protein interactions and autophagic
degradation. *Plant Cell* 26: 4974–4990

Hafrén A, Macia J-L, Love AJ, Milner JJ, Drucker M & Hofius D (2017) Selective autophagy limits
cauliflower mosaic virus infection by NBR1-mediated targeting of viral capsid protein and
particles. *Proceedings of the National Academy of Sciences* 114: E2026–E2035

Hofius D, Schultz-Larsen T, Joensen J, Tsitsigiannis DI, Petersen NHTT, Mattsson O, Jørgensen
LB, Jones JDGG, Mundy J & Petersen M (2009) Autophagic Components Contribute to
Hypersensitive Cell Death in Arabidopsis. *Cell* 137: 773–783

Huang H, Kawamata T, Horie T, Tsugawa H, Nakayama Y, Ohsumi Y & Fukusaki E (2015) Bulk
RNA degradation by nitrogen starvation-induced autophagy in yeast. *EMBO J* 34: 154

Huang L, Yu LJ, Zhang X, Fan B, Wang FZ, Dai YS, Qi H, Zhou Y, Xie LJ & Xiao S (2019) Autophagy
regulates glucose-mediated root meristem activity by modulating ROS production in
Arabidopsis. *Autophagy* 15: 407–422

Jacomin AC, Samavedam S, Promponas V & Nezis IP (2016) iLIR database: A web resource for
LIR motif-containing proteins in eukaryotes. *Autophagy* 12: 1945–1953

Ji C, Zhou J, Guo R, Lin Y, Kung C-H, Hu S, Ng WY, Zhuang X & Jiang L (2020) AtNBR1 Is a
Selective Autophagic Receptor for AtExo70E2 in Arabidopsis. *Plant Physiol* 184: 777–791

Ji MM, Lee JM, Mon H, Xu J, Tatsuke T & Kusakabe T (2016) Proteasome inhibitor MG132
impairs autophagic flux through compromising formation of autophagosomes in Bombyx
cells. *Biochem Biophys Res Commun* 479: 690–696

Jing H, Korasick DA, Emenecker RJ, Morffy N, Wilkinson EG, Powers SK & Strader LC (2022)
Regulation of AUXIN RESPONSE FACTOR condensation and nucleo-cytoplasmic
partitioning. *Nat Commun* 13: 4015

Kanne J V., Ishikawa M, Bressendorff S, Ansbøl J, Hasebe M, Rodriguez E & Petersen M (2021)
Overexpression of ATG8/LC3 enhances wound-induced somatic reprogramming in
Physcomitrium patens. *Autophagy* 18: 1463–1466

Kirkin V, Lamark T, Sou YS, Bjørkøy G, Nunn JL, Bruun JA, Shvets E, McEwan DG, Clausen TH,
Wild P, *et al* (2009) A Role for NBR1 in Autophagosomal Degradation of Ubiquitinated
Substrates. *Mol Cell* 33: 505–516

Kleine-Vehn J, Leitner J, Zwiewka M, Sauer M, Abas L, Luschnig C & Friml J (2008) Differential
degradation of PIN2 auxin efflux carrier by retromer-dependent vacuolar targeting.
Proceedings of the National Academy of Sciences of the USA 105: 17812–17817

Lee HW, Cho C, Pandey SK, Park Y, Kim M-J & Kim J (2019) LBD16 and LBD18 acting
downstream of ARF7 and ARF19 are involved in adventitious root formation in
Arabidopsis. *BMC Plant Biology* 19: 1–11

Levine B & Klionsky DJ (2004) Development by Self-Digestion. *Dev Cell* 6: 463–477

Lim J, Lachenmayer ML, Wu S, Liu W & Kundu M (2015) Proteotoxic Stress Induces
Phosphorylation of p62 / SQSTM1 by ULK1 to Regulate Selective Autophagic Clearance of
Protein Aggregates. 1–28

Liu EY, Xu N, O'Prey J, Lao LY, Joshi S, Long JS, O'Prey M, Croft DR, Beaumatin F, Baudot AD, *et*
al (2015) Loss of autophagy causes a synthetic lethal deficiency in DNA repair.
Proceedings of the National Academy of Sciences 112: 773–778

Liu Y, Xiong Y & Bassham DC (2009) Autophagy is required for tolerance of drought and salt
stress in plants. *Autophagy* 5: 954–963

Llamas E, Torres-Montilla S, Lee HJ, Barja MV, Schlimgen E, Dunken N, Wagle P, Werr W,
Zuccaro A, Rodríguez-Concepción M, *et al* (2021) The intrinsic chaperone network of
Arabidopsis stem cells confers protection against proteotoxic stress. *Aging Cell* 20: 1–12

Lv B, Wei K, Hu K, Tian T, Zhang F, Yu Z, Zhang D, Su Y, Sang Y, Zhang X, *et al* (2021) MPK14-
mediated auxin signaling controls lateral root development via ERF13-regulated very-
long-chain fatty acid biosynthesis. *Mol Plant* 14: 285–297

Mcswiggen DT, Mir M, Darzacq X & Tjian R (2019) Evaluating phase separation in live cells:
diagnosis, caveats, and functional consequences. *Genes Dev* 33: 1619–1634

Moreno-Risueño MA, Van Norman JM, Moreno A, Zhang J, Ahnert SE & Benfey PN (2010)
Oscillating Gene Expression Determines Competence for Periodic Arabidopsis Root
Branching. *Science (1979)* 329: 1306–1311

Mruk DD & Cheng CY (2011) Enhanced chemiluminescence (ECL) for routine immunoblotting.
Spermatogenesis 1: 121–122

Munch D, Rodriguez E, Bressendorff S, Park OK, Hofius D & Petersen M (2014) Autophagy
deficiency leads to accumulation of ubiquitinated proteins, ER stress, and cell death in
Arabidopsis. *Autophagy* 10: 1579–1587

Nolan TM, Brennan B, Yang M, Chen J, Zhang M, Li Z, Wang X, Bassham DC, Walley J & Yin Y
(2017) Selective Autophagy of BES1 Mediated by DSK2 Balances Plant Growth and
Survival. *Dev Cell* 41: 33-46.e7

Okushima Y, Fukaki H, Onoda M, Theologis A & Tasaka M (2007) ARF7 and ARF19 regulate
lateral root formation via direct activation of LBD/ASL genes in Arabidopsis. *Plant Cell* 19:
118–130

Orosa-Puente B, Leftley N, von Wangenheim D, Banda J, Srivastava AK, Hill K, Truskina J,
Bhosale R, Morris E, Srivastava M, *et al* (2018) Root branching toward water involves
posttranslational modification of transcription factor ARF7. *Science* 362: 1407–1410

Perez-Riverol Y, Bai J, Bandla C, Hewapathirana S, García-Seisdedos D, Kamatchinathan S,
Kundu D, Prakash A, Frericks-Zipper A, Eisenacher M, Walzer M, Wang S, Brazma A,
Vizcaíno JA (2022). The PRIDE database resources in 2022: A Hub for mass spectrometry-
based proteomics evidences. Nucleic Acids Res 50(D1):D543-D552.

Perianez-Rodriguez J, Rodriguez M, Marconi M, Bustillo-Avendaño E, Wachsman G, Sanchez-
Corrionero A, De Gernier H, Cabrera J, Perez-Garcia P, Gude I, *et al* (2021) An auxin-
regulable oscillatory circuit drives the root clock in Arabidopsis. *Sci Adv* 7

Powers SK, Holehouse AS, Korasick DA, Schreiber KH, Clark NM, Jing H, Emenecker R, Han S,
Tycksen E, Hwang I, *et al* (2019) Nucleo-cytoplasmic Partitioning of ARF Proteins Controls
Auxin Responses in Arabidopsis thaliana. *Mol Cell* 76: 177–190

Rodriguez E, Chevalier J, Olsen J, Ansbøl J, Kapousidou V, Zuo Z, Svenning S, Loefke C, Koemeda
S, Drozdowskyj PS, *et al* (2020) Autophagy mediates temporary reprogramming and
dedifferentiation in plant somatic cells. *EMBO J* 39: e103315

Rosado A, Li R, van de Ven W, Hsu E & Raikhel N V. (2012) Arabidopsis ribosomal proteins
control developmental programs through translational regulation of auxin response
factors. *Proceedings of the National Academy of Sciences* 109: 19537–19544

Santos A, Colaço AR, Nielsen AB, Niu L, Strauss M, Geyer PE, Coscia F, Albrechtsen NJW, Mundt
F, Jensen LJ, *et al* (2022) A knowledge graph to interpret clinical proteomics data. *Nat
Biotechnol* 40: 692–702

Scott RC, Schuldiner O & Neufeld TP (2004) Role and Regulation of Starvation-Induced
Autophagy in the Drosophila Fat Body. *Dev Cell* 7: 167–178

Sedaghatmehr M, Thirumalaikumar VP, Kamranfar I, Marmagne A, Masclaux-Daubresse C &
Balazadeh S (2019) A regulatory role of autophagy for resetting the memory of heat stress
in plants. *Plant Cell Environ* 42: 1054–1064

Stephani M, Picchianti L, Gajic A, Beveridge R, Skarwan E, Hernandez VS de M, Mohseni A,
Clavel M, Zeng Y, Naumann C, *et al* (2020) A cross-kingdom conserved ER-phagy receptor
maintains endoplasmic reticulum homeostasis during stress. *Elife* 9: 1–105

Sugimoto K, Jiao Y & Meyerowitz EM (2010) Arabidopsis Regeneration from Multiple Tissues
Occurs via a Root Development Pathway. *Dev Cell* 18: 463–471

Svenning S, Lamark T, Krause K & Johansen T (2011) Plant NBR1 is a selective autophagy
substrate and a functional hybrid of the mammalian autophagic adapters NBR1 and
p62/SQSTM1. *Autophagy* 7: 993–1010

Tarnowski L, Rodriguez MC, Brzywczy J, Piecho-Kabacik M, Krčkova Z, Martinec J, Wawrzynska
A & Sirko A (2020) A selective autophagy cargo receptor NBR1 modulates abscisic acid
signalling in Arabidopsis thaliana. *Sci Rep* 10: 7778

Turco E, Savova A, Gere F, Ferrari L, Romanov J, Schuschnig M, Martens S & Savova A⬚ (2021)
Reconstitution defines the roles of p62, NBR1 and TAX1BP1 in ubiquitin condensate
formation and autophagy initiation. *Nat Commun* 12: 5212

Ulmasov T, Hagen G & Guilfoyle TJ (1997) ARF1, a transcription factor that binds to auxin
response elements. *Science (1979)* 276: 1865–1868

Vanhee C, Zapotoczny G, Masquelier D, Ghislain M & Batoko H (2011) The Arabidopsis
Multistress Regulator TSPO Is a Heme Binding Membrane Protein and a Potential
Scavenger of Porphyrins via an Autophagy-Dependent Degradation Mechanism. *Plant Cell*
23: 785–805

Wachsman G, Zhang J, Moreno-Risueño MA, Anderson CT & Benfey PN (2020) Cell wall
remodeling and vesicle trafficking mediate the root clock in *Arabidopsis*. *Science (1979)*
370: 819–823

Wang HQ, Liu HM, Zhang HY, Guan Y & Du ZX (2008) Transcriptional upregulation of BAG3
upon proteasome inhibition. *Biochem Biophys Res Commun* 365: 381–385

Wang S, Tiwari SB, Hagen G & Guilfoyle TJ (2005) AUXIN RESPONSE FACTOR7 Restores the
Expression of Auxin-Responsive Genes in Mutant Arabidopsis Leaf Mesophyll Protoplasts.
Plant Cell 17: 1979–1993

Wang X, Jiang B, Gu L⬚, Gu L, Chen Y, Mora M, Zhu M, Noory E, Wang Q & Lin C (2021) A
photoregulatory mechanism of the circadian clock in Arabidopsis. *Nat Plants* 7: 1397–
1408

Wang X, Yu R, Wang J, Lin Z, Han X, Deng Z, Fan L, He H, Deng XW & Chen H (2020) The
Asymmetric Expression of SAUR Genes Mediated by ARF7/19 Promotes the Gravitropism
and Phototropism of Plant Hypocotyls. *Cell Rep* 31: 107529

Wu J, Michaeli S, Picchianti L, Dagdas Y, Galili G & Peled-Zehavi H (2021) ATI1 (ATG8-interacting
protein 1) and ATI2 define a plant starvation-induced reticulophagy pathway and serve as
MSBP1/MAPR5 cargo receptors. *Autophagy* 17: 3375–3388

Xie D, Chen M, Niu J, Wang L, Li Y, Fang X, Li P & Qi Y (2021) Phase separation of SERRATE
drives dicing body assembly and promotes miRNA processing in Arabidopsis. *Nat Cell Biol*
23: 32–39

Yoshii SR, Kuma A, Akashi T, Hara T, Yamamoto A, Kurikawa Y, Itakura E, Tsukamoto S, Shitara
H, Eishi Y, *et al* (2016) Systemic Analysis of Atg5-Null Mice Rescued from Neonatal
Lethality by Transgenic ATG5 Expression in Neurons. *Dev Cell* 39: 116–130

Zhang Y, Yan L, Zhou Z, Yang P, Tian E, Zhang K, Zhao Y, Li Z, Song B, Han J, *et al* (2009) SEPA-1
Mediates the Specific Recognition and Degradation of P Granule Components by
Autophagy in C. elegans. *Cell* 136: 308–321

Zhou J, Wang J, Cheng Y, Chi YJ, Fan B, Yu JQ & Chen Z (2013a) NBR1-Mediated Selective
Autophagy Targets Insoluble Ubiquitinated Protein Aggregates in Plant Stress Responses.
PLoS Genet 9: 1004477

Zhou J, Wang J, Cheng Y, Chi YJ, Fan B, Yu JQ & Chen Z (2013b) NBR1-Mediated Selective
Autophagy Targets Insoluble Ubiquitinated Protein Aggregates in Plant Stress Responses.
PLoS Genet 9: 1004477

Zhou J, Zhang Y, Qi J, Chi Y, Fan B, Yu JQ & Chen Z (2014) E3 Ubiquitin Ligase CHIP and NBR1-
Mediated Selective Autophagy Protect Additively against Proteotoxicity in Plant Stress
Responses. *PLoS Genet* 10: e1004116

Zuo Z, Roux M, Rodriguez E & Petersen M (2022) mRNA Decapping Factors LSM1 and PAT
Paralogs Are Involved in Turnip Mosaic Virus Viral Infection. *Molecular Plant-Microbe*
Interactions 5: 125–129

A

	Col-0		*atg5-1*		*atg2-1*		*nbr1-c1*	
	NT	MG132	NT	MG132	NT	MG132	NT	MG132
Rel. Int.	1.0	1.9**	1.7*	2.8****	1.9*	2.6****	2.3**	3.2****
STD	0.0	0.4	0.2	1.1	0.6	0.6	1.0	1.0

α-ARF7

Rbc L

B

Relative *ARF7* expression

C

α-GFP

Rbc L

Rel. Int.	1.0	7.2***
STD	0.0	3.1

D pARF7::gARF7 Venus

Venus | Inset | Bright-Field

E pARF7::gARF7 Venus *atg-2-1*

Venus | Inset | Bright-Field

F Cytosol signal — Mature zone

Fluorescence (AU)

ARF7-Venus | ARF7-Venus atg2 (****)

G Cytosol signal — Mature zone

Fluorescence (AU)

ARF7-Venus | ARF7-Venus atg2 (****)

H Nuclear signal — Meristem

Fluorescence (AU)

ARF7-Venus | ARF7-Venus atg2 (****)

I Nuclear signal — Meristem

Fluorescence (AU)

ARF7-Venus | ARF7-Venus atg2 (ns)

Figure 2

Figure 4

A

Mock vs. auxin treated

Col-0

atg2-1

-log10(P - value)

Log2(Fold Change)

B

Downregulated proteins

Col-0   atg2-1

167   13   94

Upregulated proteins

Col-0   atg2-1

146   12   114

C

LR Density (LR mm⁻¹)

MS
0.1 µM NAA

Col-0   atg2-1   atg5-1   Col-0   atg2-1   atg5-1

D

Col-0   atg2-1   atg5-1

NT

0.1 µM NAA