Deep-sea bacteriophages facilitate host utilization of polysaccharides

Chong Wang[1,2,4], Rikuan Zheng[1,2,4], Tianhang Zhang[1,2,3,4], Chaomin Sun[1,2,3,4*]

[1]CAS and Shandong Province Key Laboratory of Experimental Marine Biology & Center of Deep Sea Research, Institute of Oceanology, Chinese Academy of Sciences, Qingdao, China.

[2]Laboratory for Marine Biology and Biotechnology, Qingdao National Laboratory for Marine Science and Technology, Qingdao, China.

[3]College of Earth Science, University of Chinese Academy of Sciences, Beijing, China.

[4]Center of Ocean Mega-Science, Chinese Academy of Sciences, Qingdao, China.

[*]Corresponding author

Chaomin Sun

Tel.: +86 532 82898857; fax: +86 532 82898857.

E-mail address: sunchaomin2020@126.com; sunchaomin@qdio.ac.cn

Key words: deep-sea virus, polysaccharides, bacteriophages, auxiliary metabolic genes, carbon cycle

Running title: Bacteriophages assist host polysaccharide metabolism

Abstract

Bacteriophages are ubiquitous in nature and play key roles in various ecosystems. They impact microbial community composition and reprogram the metabolism of diverse host cells using auxiliary metabolic genes (AMGs). Whether bacteriophages can reprogram host polysaccharide metabolism through AMGs remains unclear, however. Here, we found for the first time that polysaccharides induce the production of different types of bacteriophages in two deep-sea *Lentisphaerae* strains (WC36 and zth2). Through physiological assays, genomic analysis and transcriptomics assays, we found that these bacteriophages might assist their hosts in metabolizing polysaccharides through AMGs. Moreover, the isolated bacteriophages could effectively assist a marine bacterium (*Pseudomonas stutzeri* 273) in metabolizing and utilizing polysaccharide to promote its growth. These findings shed light on the importance of atypical and poorly understood virus-host interactions and bring us closer to understanding the potential role of deep-sea viruses in marine ecosystems.

Introduction

Viruses are the most abundant and genetically diverse biological entities on this planet (Suttle, 2007), and potentially play a role in shaping microbial abundances, community structure, and the evolutionary trajectory of core metabolisms (Sullivan et al., 2006; Samson et al., 2013; Zimmerman et al., 2020). The effect of bacteriophages (viruses that infect bacteria) on their specific hosts and within the microbial community is largely determined by their lifestyle. Traditionally, most bacteriophage life cycles are classified as being either lytic or lysogenic (Du Toit, 2017). During the lytic cycle, phages hijack the host's metabolic machinery for replication of their own genome and the production of progeny particles that are released through lysis. By contrast, during lysogeny, phages integrate their genomes into the host chromosome (or exist in the extrachromosomal form) and enter a dormant state, with the potential to re-enter a lytic cycle and release progeny by environmental changes at a later stage. Currently, more and more attention has been paid to chronic life cycles where bacterial growth continues despite phage reproduction (Hoffmann Berling and Maze, 1964). During a chronic cycle, progeny phage particles are released from host cells via extrusion or budding without killing the host (Putzrath and Maniloff, 1977; Russel, 1991; Marvin et al., 2014), instead of lysis of the host cell. Chronic infections are common among eukaryotic viruses (Godkin and Smith, 2017), however, only a handful of chronic viruses have been described for prokaryotes (Roux et al., 2019; Alarcón-Schumacher et al., 2022). The best-studied examples of chronic infection are those from filamentous ssDNA (single-stranded DNA) phages that to date have primarily been identified in the order *Tubulavirales*, which currently involves three families (*Inoviridae*, *Paulinoviridae* and *Plectroviridae*) (Ackermann, 2007, 2009; Knezevic et al., 2021; Chevallereau et al., 2022). One of the most distinctive features of inoviruses is their ability to establish a chronic infection whereby their genomes may either integrate into the host chromosome or reside within the cell in a non-active form, and inoviruses particles are continuously released without killing the host. In addition, tailless phages enclosed in lipid membrane are also released from hosts

during chronic life cycles (Liu et al., 2022). Recent studies indicate that chronic

cycles are more widespread in nature than previously thought (Mäntynen et al., 2021),

and the abundance of inoviruses in ecosystem may have been far underestimated

(Roux et al., 2019). These findings have suggested that plenty of novel phages in the

biosphere may be produced through chronic life cycles, but whether chronic cycles

are associated with specific environments or ecological conditions remains to be

thoroughly explored (Chevallereau et al., 2022).

Marine ecosystems always influence the operating conditions for life on earth via

microbial interaction networks (Falkowski et al., 1998; Faust and Raes, 2012;

Wigington et al., 2016), which are modulated by viruses through impacting their

lifespan, gene flow and metabolic outputs (Suttle, 2005). A majority of these viruses

are bacteriophages, which exist widely in oceans (Breitbart, 2012). In addition to core

viral genes encoding viral structural proteins (Brum et al., 2016), bacteriophages also

encode various auxiliary metabolic genes (AMGs) (Brum and Sullivan, 2015) that

provide supplemental support to key steps of host metabolism. For instance, AMGs of

marine bacteriophages have been predicted to be involved in photosynthesis (Mann et

al., 2003), nitrogen- (Ahlgren et al., 2019; Gazitúa et al., 2021), sulfur-

(Anantharaman et al., 2014; Roux et al., 2016), and phosphorus cycling (Zeng and

Chisholm, 2012), nucleotide metabolism (Sullivan et al., 2005; Dwivedi et al., 2013;

Enav et al., 2014) and almost all central carbon metabolisms in host cells (Hurwitz et

al., 2013). However, AMGs of chronic phages have not been reported. Thus, it is

worth exploring whether chronic phages could carry AMGs and assist host

metabolism during chronic infection.

The deep sea harbors abundant and undiscovered viruses, which potentially

control the metabolism of microbial hosts and influence biogeochemical cycling (Li et

al., 2021). However, the vast majority of deep-sea bacteria cannot be cultivated,

which greatly limits the host range of deep-sea bacteriophages, resulting in most

unknown bacteriophages could not be isolated in the laboratory. Thus, it is of great

significance to further identify unknown phages in the deep sea and explore their

relationship with microbial hosts and even marine ecosystems. Here, we report that

polysaccharides can induce deep-sea *Lentisphaerae* bacteria (difficult-to-cultivate microorganisms) to release some bacteriophages. These bacteriophages might assist host metabolism to degrade polysaccharide and thereby promote host growth via AMGs.

Results and Discussion

Polysaccharides promote the growth of deep-sea *Lentisphaerae* strain WC36 and stimulate the expression of phage-associated genes

As a primary carbon source, polysaccharides are a ubiquitous energy source for microorganisms in both terrestrial and marine ecosystems (Zheng et al., 2021a). In our previous work, we successfully isolated a novel *Bacteroidetes* species through a polysaccharide degradation-driven strategy from the deep-sea cold seep (Zheng et al., 2021a). Of note, using the same approach and deep-sea sample replicates, we also cultured a bacterial strain WC36 that was identified as a member of the phylum *Lentisphaerae* by 16S rRNA gene sequence-based phylogenetic analysis (Fig. 1A). As expected, growth assays showed that two kinds of polysaccharides including laminarin and starch could promote strain WC36 growth (Fig. 1B). In particular, supplementation of 10 g/L laminarin and 10 g/L starch in rich medium resulted in ~7-fold and ~4-fold growth promotion respectively compared to cultivation in rich medium (Fig. 1B).

To explore the reasons behind this significant growth promotion by polysaccharide supplementation, we performed a transcriptomic analysis of strain WC36 cultured in rich medium supplemented either with or without 10 g/L laminarin for 5 days and 10 days. In addition to the up-regulation of genes related to glycan transport and degradation, when 10 g/L laminarin was added in the rich medium, the most upregulated genes were phage-associated (e. g. phage integrase, phage portal protein) (Fig. 1C and Supplementary Table 1). Consistently, RT-qPCR results (Fig. 1D) confirmed upregulation of some genes encoding phage-associated proteins, as shown in Fig. 1C. Therefore, we speculate that the metabolism of polysaccharides in strain WC36 might be closely connected with the phage production.

Fig. 1. Polysaccharides promote the deep-sea *Lentisphaerae* strain WC36 growth and
stimulate the expression of phage-associated genes. (A) Maximum likelihood phylogenetic tree
of 16S rRNA gene sequences from strain WC36, strain zth2, and some
Planctomycetes-Verrucomicrobia-Chlamydia (PVC) group bacteria. *Bacillus cereus* ATCC 14579
was used as the outgroup. Bootstrap values (%) > 80 are indicated at the base of each node with
the gray dots (expressed as percentages of 1000 replications). (B) Growth assays of strain WC36
cultivated in rich medium either without supplementation, with 5 g/L or 10 g/L laminarin or with 5
g/L or 10 g/L starch. (C) Transcriptomics-based heat map showing all up-regulated genes
encoding phage-associated proteins in strain WC36 cultured in rich medium supplemented with 10
g/L laminarin. "Rich" indicates strain WC36 cultivated in rich medium; "Lam" indicates strain
WC36 cultivated in rich medium supplemented with 10 g/L laminarin. (D) RT-qPCR detection of
the expression of some genes encoding phage-associated proteins shown in panel C. The heat map
was generated by Heml 1.0.3.3. The numbers in panel C represent multiple differences in gene
expression (by taking $\log_2$ values).

Polysaccharides induce the production of bacteriophages in *Lentisphaerae* strain WC36

To test our hypothesis that polysaccharides might induce strain WC36 to release bacteriophages, we grew strain WC36 in rich medium supplemented with or without laminarin or starch and isolated bacteriophages from the cell suspension supernatant according to established standard protocols (Lin et al., 2012). Based on the growth curve of strain WC36, we found that the strictly anaerobic WC36 grew much slower than normal aerobic bacteria. Therefore, different to the typical sampling time (24 h) for bacteriophage isolation from other bacteria (Jiang and Paul, 1996; Weinbauer et al., 1999), we selected a longer sampling time (10 days) to extract bacteriophages. TEM observation showed that many different shapes of phage-like structures indeed existed in the supernatant of WC36 cells cultured in rich medium supplemented with laminarin (Fig. 2A, panels II-IV) or starch (Fig. 2A, panels V-VIII). In contrast, we did not observe any phage-like structures in the supernatant of WC36 cells cultured in rich medium (Fig. 2A, panel I) or in different media without culturing bacterial cells (Supplementary Fig. 1A, B). These results suggest that polysaccharides indeed stimulate the production of various bacteriophages from strain WC36. Correspondingly, in the presence of laminarin, different forms of bacteriophages were observed by TEM, with filamentous ones dominant (Fig. 2A). The length of these filamentous phages was about ~0.4 μm to ~8.0 μm (Fig. 2A, panel II), and the size of the hexagonal phages was less than 100 nm (Fig. 2A, panel IV). In the presence of starch, in addition to filamentous and hexagonal phages (Fig. 2A, panels V, VI, and VIII), we also observed a kind of icosahedral *Microviridae*-like phage with a diameter of 25-30 nm (Fig. 2A, panel VII).

Given that we found bacteriophages in cultures of strain WC36, we next sought to explore whether bacteriophages adhering to bacterial cells could be observed. To this end, we checked the morphology of strain WC36 in the absence or presence of polysaccharides. As expected, with TEM we observed many filamentous phage-like structures around strain WC36 cells cultivated in rich medium supplemented with laminarin (Fig. 2B, panel II) and starch (Fig. 2B, panels III-IV). In contrast, we could

not find any phage-like structures around strain WC36 cells cultivated either in rich medium (Fig. 2B, panel I), in rich medium supplemented with 10 g/L laminarin (Supplementary Fig. 2A) or in 10 g/L starch without culturing bacterial cells (Supplementary Fig. 2B). Moreover, based on TEM observation of ultrathin sections of strain WC36 cultured in medium supplemented with polysaccharide, we even observed filamentous phages that were being released from or entering into host cells via a kind of extrusion or budding structure around bacterial cells (Fig. 2C, panels II-IV), suggesting that filamentous bacteriophages might be released chronically as reported previously (Chevallereau et al., 2022; Liu et al., 2022).

Fig. 2. Polysaccharides induce the production of bacteriophages in *Lentisphaerae* strain WC36. (A) TEM observation of phages extracted from the supernatant of WC36 cells cultured in rich medium supplemented with or without polysaccharide. Panel I shows the absence of phages in the supernatant of WC36 cells cultivated in rich medium. Panels II-IV show the morphology of phages present in the supernatant of WC36 cells cultivated in rich medium supplemented with 10 g/L laminarin. Yellow hollow arrows indicate the typical granular structure of filamentous phages,

and green arrows indicate phage-like particles with other shapes. Panels V-VIII show the morphology of phages present in the supernatant of strain WC36 cells cultivated in rich medium supplemented with 10 g/L starch. Typical filamentous phages are indicated with yellow hollow arrows, and two other kinds of phage-like particles with different shapes are indicated by orange and green arrows, respectively. (B) TEM observation of strain WC36 cultured in rich medium supplemented with or without 10 g/L laminarin or 10 g/L starch. Panel I shows representative morphology of strain WC36 cultivated in rich medium. Panel II shows the morphology of strain WC36 cultivated in rich medium supplemented with 10 g/L laminarin. Panels III-IV show the morphology of strain WC36 cultivated in rich medium supplemented with 10 g/L starch. Red arrows indicate filamentous phages associated with bacterial cells. (C) TEM of an ultrathin section of strain WC36 cultured in rich medium supplemented with or without 10 g/L laminarin or 10 g/L starch. Panel I shows an ultrathin section of strain WC36 cultivated in rich medium; Panels II-III show ultrathin sections of strain WC36 cultivated in rich medium supplemented with 10 g/L laminarin; Panel IV shows an observation of the ultrathin section of strain WC36 cultivated in rich medium supplemented with 10 g/L starch. Red arrows indicate filamentous phages being released from or entering bacterial cells.

To further verify the release style of bacteriophages from strain WC36 in the presence of polysaccharide, we prolonged the incubation time of this strain up to 30 days in medium supplemented with 5 g/L or 10 g/L laminarin. Strain WC36 showed stable growth after it entered the stationary phase (Fig. 3A). The bacterial cells maintained their cell shape intact regardless of the presence of laminarin, indicating their healthy state after for a long time (Supplementary Fig. 3A, panels I-III). Apparently, the replication and release of bacteriophages in strain WC36 did not kill the host cell, consistent with the key feature of chronic bacteriophages (Howard-Varona et al., 2017). With an increase of culture time (5, 10, or 30 days) and laminarin concentration (5 or 10 g/L), the number of released bacteriophages increased (Supplementary Fig. 3A, panels IV-VI; Supplementary Fig. 4, panels IV-IX). Specifically, the average number per microliter of filamentous phages and hexagonal phages extracted from the supernatant of strain WC36 cultured in rich medium supplemented with 10 g/L laminarin for 5, 10, 30 days (around 9.7, 29, 65.3 and 9, 30, 46.7) was higher than those cultured in rich medium supplemented with 5 g/L laminarin for 5, 10, 30 days (around 4.3, 13.7, 35.3 and 4, 11.3, 17.7), respectively (Fig. 3B, C).

Fig. 3. Bacteriophages induced from *Lentisphaerae* strain WC36 by polysaccharides are released via chronic infections. (A) Growth curve of strain WC36 cultivated in either rich medium alone or rich medium supplemented with 5 g/L or 10 g/L laminarin for 30 days. The number of filamentous phages (B) and hexagonal phages (C) extracted from 1 μL of the supernatant of strain WC36 cell cultured in rich medium supplemented with or without 5 g/L or 10 g/L laminarin (for 5, 10, and 30 days). The average numbers are shown at the top of the bar charts.

In addition, in the presence of laminarin we observed many extrusions or buddings around bacterial cells (Supplementary Fig. 3B, panels II-III), which are typical cellular structures in host cells used for chronic bacteriophage release (Marvin et al., 2014; Krupovic and Consortium, 2018). In contrast, phage-like structures with extrusions or buddings were not found in cells of strain WC36 cultivated in rich medium (Supplementary Fig. 3B, panel I). Hence, these results collectively suggest that filamentous phages induced from deep-sea strain WC36 replicate in a chronic manner. However, the morphology and process of hexagonal phages that entry and exit into the WC36 cells were not observed.

Polysaccharides promote the growth of deep-sea *Lentisphaerae* strain zth2 and induce bacteriophage production

To explore whether the production of bacteriophages induced by polysaccharide is an individual case, we further checked growth promotion and bacteriophage induction by polysaccharide in another cultured deep-sea *Lentisphaerae* - strain zth2 - which was isolated by the same polysaccharide degradation-driven strategy as that used for strain WC36. 16S rRNA gene sequence similarity calculation using the NCBI server indicated that the closest relative of strain zth2 was strain WC36 (90.4%), which was lower than the threshold sequence identity (94.5%) for distinct genera (Yarza et al., 2014). Based on a maximum likelihood tree and similarity of 16S rRNA gene sequences, we propose that strain zth2 and WC36 are the type strains of two different novel genera, both belonging to the *Lentisphaeraceae* family, *Lentisphaerales* order, *Lentisphaeria* class. Consistently, laminarin indeed promoted the growth of strain zth2 (Fig. 4A). Analysis of the differential transcriptome and RTq-PCR of strain zth2 cultured in either unsupplemented medium or with laminarin showed that many genes encoding critical factors associated with the phage life cycle were up-regulated (ranging from 4 to 200 fold) (Fig. 4B, C and Supplementary Table 2), suggesting that phages might also be involved in the utilization of laminarin by strain zth2, the same as in strain WC36.

To further confirm the existence of bacteriophages in the supernatant of strain zth2 cell suspension cultured in medium supplemented with polysaccharide, we performed bacteriophage isolation from the supernatant of strain zth2 cultures followed by TEM. Indeed, many filamentous-, hexagonal- and icosahedral phages were clearly observed in the culture of strain zth2 grown in the presence of laminarin (Fig. 4D, panels I-III) and starch (Fig. 4D, panels IV-V), but not in rich medium short of polysaccharide (Fig. 4D, panel VI). This suggests that bacteriophages were induced by supplementation of polysaccharide to strain zth2.

Fig. 4. Polysaccharides promote the growth of deep-sea *Lentisphaerae* strain zth2 and induce

the production of bacteriophages. (A) Growth assays of strain zth2 cultivated in rich medium

supplemented with or without 3 g/L laminarin for four days. (B) Transcriptomics-based heat map

showing all up-regulated genes encoding phage-associated proteins in strain zth2 cultured in rich

medium supplemented with 3 g/L laminarin. (C) RT-qPCR detection of the expressions of some

genes encoding phage-associated proteins shown in panel B. The numbers in panel C represent

multiple differences in gene expression (by taking $\log_2$ values). (D) TEM observation of phages

extracted from the supernatant of a cell suspension of strain zth2 cultured in the rich medium

supplemented with or without polysaccharides. Panels I-III show the morphology of phages in the

cell supernatant of strain zth2 cultivated in rich medium supplemented with 3 g/L laminarin.

Typical filamentous phage is indicated with yellow hollow arrows, and two other kinds of

phage-like particles with different shapes are indicated by orange and green arrows, respectively.

Panels IV-V show the morphology of phages present in the cell supernatant of strain zth2

cultivated in rich medium supplemented with 3 g/L starch. Typical filamentous phages are

indicated with red arrows, and other phage-like particles with different shapes are indicated by

green arrows. Panel VI shows that no phages were observed in the cell supernatant of strain zth2

cultivated in rich medium. "Rich" indicates strain zth2 cultivated in rich medium; "Lam" indicates

strain zth2 cultivated in rich medium supplemented with 3 g/L laminarin.

In addition, transcriptome and RT-qPCR results showed that genes encoding

bacterial secretion system-related proteins were up-regulated in strains WC36

(Supplementary Fig. 5A, B) and zth2 (Supplementary Fig. 5C, D) cultured in medium

supplemented with laminarin. It has been reported that filamentous phages utilise host

cell secretion systems for their own egress from bacterial cells (Davis et al., 2000;

Bille et al., 2005). Therefore, the filamentous phages induced from strains WC36 and

zth2 might also get in and out of host cells using bacterial secretion systems. Taken

together, we conclude that laminarin effectively promotes the growth of deep-sea

Lentisphaerae strains WC36 and zth2, and induces the production of bacteriophages.

Distinct from previous reports that bacteriophages are induced under traditional

conditions such as low temperature (Wang et al., 2007), mitomycin C (Mazaheri

Nezhad Fard et al., 2010), and ultraviolet irradiation (McKay and Baldwin, 1973),

here, we report that polysaccharides induce the production of bacteriophages.

Bacteriophages reprogram the polysaccharide metabolism of *Lentisphaerae*

strains WC36 and zth2

In natural ecosystems, viral infections indirectly influence microbial metabolic fluxes,

metabolic reprogramming, and energy homeostasis of host cells (Howard-Varona et

al., 2020). For example, cyanoviruses possess AMGs encoding core photosynthetic

reaction centers (Sullivan et al., 2006) which are highly expressed during infection,

thus stimulating photosynthesis and enhancing viral abundance (Lindell et al., 2007).

In this study, we clearly show that polysaccharide effectively promotes bacterial

growth and simultaneously induces the production of bacteriophages, suggesting a

close relationship between bacteriophages and host polysaccharide metabolism.

Therefore, we sought to ask whether bacteriophages could reprogram polysaccharide

metabolism in deep-sea bacteria. For this purpose, we sequenced the genomes of the

bacteriophages induced by polysaccharides in *Lentisphaerae* strains WC36 and zth2.

Eventually, two assembled genomes (Phage-WC36-1, 8.0 kb; Phage-WC36-2, 28.3 kb)

were obtained from the bacteriophages of strain WC36 (Fig. 5A, B and

Supplementary Tables 3, 4). Meanwhile, two assembled genomes (Phage-zth2-1, 8.0

kb; Phage-zth2-2, 40.4 kb) were obtained from the phages of strain zth2 (Fig. 5A, C

and Supplementary Tables 3, 5). Subsequently, we carefully compared the

bacteriophage genomes with those of the corresponding hosts (strains WC36 and zth2) using Galaxy Version 2.6.0 (https://galaxy.pasteur.fr/) (Afgan et al., 2018) with the NCBI BLASTN method and used BWA-mem software for read mapping from host whole genome sequencing (WGS) to these bacteriophages. These analyses both showed that the bacteriophage genomes are completely outside of the host chromosomes, as reported previously (Chevallereau et al., 2022). Through comparison of genome sequence (Supplementary Fig. 6), IMG database analysis, and phylogenetic analysis (Supplementary Fig. 7), we confirmed that Phage-WC36-1 was the same as Phage-zth2-1 (inoviruses, belonging to the family *Inoviridae*, order *Tubulavirales*, class *Faserviricetes*), indicating that inoviruses might be common in the *Lentisphaerae* phylum. Inovirus members were the dominant population in the active group of viruses from deep-sea sediments, and encode some proteins contributing to host phenotypes and ultimately alter the fitness and other behaviors of their hosts (Engelhardt et al., 2015; Mai-Prochnow et al., 2015). A maximum likelihood tree showed that both Phage-WC36-2 and Phage-zth2-2 belonged to the class *Caudoviricetes* (Supplementary Fig. 8). In addition to genes encoding phage-associated proteins within the assembled bacteriophage genomes, there are also many genes encoding proteins associated with polysaccharide transport and degradation. These polysaccharide metabolizing proteins include glycoside hydrolase family 18 protein (GH18, HMM e-value 1.5e-14) (Fig. 5A), TonB-system energizer ExbB (Koebnik, 2005), O-polysaccharide acetyltransferase (Lunin et al., 2020), glucosaminidase, glycoside hydrolase family 127 protein (best BLASTP hit e-value 3e-25) (Fig. 5B) and transglycosylase (Fig. 5C). These CAZymes (GH18 and GH127) are specifically responsible for degrading polysaccharides (Cantarel et al., 2009). Energizer ExbB is an important part of the TonB-system, which is often present in *Bacteroides* and responsible for transport of oligosaccharides from the outer membrane into the periplasm (Foley et al., 2016). It is absent in the strain WC36 and zth2 genomes, however. Therefore, these AMGs encoding TonB-system energizer ExbB and glycoside hydrolases might reprogram the polysaccharide metabolism of host cells (strains WC36 and zth2) by providing supplemental support to key

358 metabolic processes. Overall, bacteriophages induced by polysaccharides contain

359 specific AMGs associated with polysaccharide degradation, which may play a key

360 role in the degradation and utilization of polysaccharide by host cells.

Fig. 5. Genomic organization of bacteriophages released from *Lentisphaerae* strains WC36 and zth2 cultured in rich medium supplemented with polysaccharide. (A) Schematic of the genomic composition of Phage-WC36-1 and Phage-zth2-1. (B) A diagram showing the genomic composition of Phage-WC36-2. (C) The genomic composition of Phage-zth2-2. Arrows represent different ORFs and the direction of transcription. The main putative gene products of the phages are shown, and the numbers in brackets indicate the numbers of amino acids within each ORF. Hypothetical proteins are indicated with gray arrows, structural modules are indicated by green arrows, the replication module is indicated by blue-grey arrows, the regulatory module is indicated by golden arrows, the assembly and secretion modules are indicated by blue arrows and auxiliary metabolic genes (AMGs) encoding proteins associated with polysaccharide transport and degradation are indicated by red arrows. The size of the phage genomes is shown behind each gene cluster.

Bacteriophages released from *Lentisphaerae* strain WC36 promote the growth and polysaccharide metabolism of a marine bacterium, *Pseudomonas stutzeri* 273

Given that the bacteriophages released from *Lentisphaerae* strain WC36 (named Phages-WC36 in this study) could reprogram bacterial polysaccharide metabolism and thereby promote host growth, we next sought to determine the capability of these

bacteriophages to reprogram polysaccharide metabolism and promote growth in other marine bacteria. To this end, we chose a marine bacterium - *Pseudomonas stutzeri* 273 (Wu et al., 2016) - and tested the potential growth promotion driven by Phages-WC36. Growth assays showed that Phages-WC36 could promote strain 273 growth by assisting its metabolism and utilization of starch (Fig. 6A). In particular, supplementation with Phages-WC36 and 5 g/L starch in basal medium resulted in 6~10-fold growth promotion compared to basal medium without Phages-WC36 or starch (Fig. 6A). In addition, we observed many filamentous phage and hexagonal phage structures in the supernatant of strain 273 cultivated in basal medium supplemented with 5 g/L starch and Phages-WC36 for 24 h. In contrast, we could not find any phage-like structures in three other types of media. Specifically, the average number of filamentous phages and hexagonal phages extracted from the supernatant of strain 273 cultured in basal medium supplemented with 5 g/L starch and Phages-WC36 (~44.7 and ~29, respectively) was higher than that in three other media types (where it was close to zero) (Fig. 6B). To test whether Phages-WC36 could infect strain 273, the strain 273 cultivated in basal medium supplemented with 5 g/L starch and Phages-WC36 was inoculated in basal medium and basal medium supplemented with 5 g/L starch for three consecutive times (Fig. 6C). After 3 passages of serial cultivation in basal medium supplemented with 5 g/L starch, we found that filamentous phages (~45.3, Fig. 6D) and hexagonal phages (~21.7, Fig. 6E) were also present in basal medium supplemented with starch, but not in the basal medium, which may mean that Phages-WC36 could infect strain 273 and starch is an important inducer. Consistently, genes encoding bacterial type II/IV/VI secretion system-related proteins were up-regulated in strain 273 cultured in basal medium supplemented with 5 g/L starch and 20 μl/mL Phages-WC36 in comparison to basal medium with 5 g/L starch (Supplementary Fig. 9). Upregulation of these genes may help filamentous phages get in and out of host cells (Davis et al., 2000; Bille et al., 2005). Moreover, the expression of genes encoding the TonB-dependent receptor and CAZymes was upregulated in strain 273 cultured in basal medium supplemented with starch and Phages-WC36 in comparison to basal medium with starch (Supplementary

409 Fig. 10A and Supplementary Table 6) or Phages-WC36 (Supplementary Fig. 10B and

410 Supplementary Table 7). These results suggest that Phages-WC36 proliferate in strain

411 273 and promote growth by assisting in starch metabolism and utilization.

412 Consequently, we propose that Phages-WC36 might infect other hosts and assist them

413 in degradation and utilization of polysaccharides. This provides stronger evidence that

414 the relationship between Phages-WC36 and their hosts is not a fight to the death

415 between enemies, but in contrast a mutualistic relationship between partners

416 (Shkoporov et al., 2022). Most deep-sea microbes are anaerobic, which limits

417 investigation of bacteriophages in their original host. *P. stutzeri* 273 is a typical

418 aerobic bacterium, and thus our study provides an effective way to replicate deep-sea

419 bacteriophages and perform functional analysis in aerobic heterologous hosts.

Fig. 6. Bacteriophages released from *Lentisphaerae* strain WC36 promote the growth and polysaccharide metabolism of a marine bacterium *Pseudomonas stutzeri* 273. (A) Growth curve and status of *P. stutzeri* 273 cultivated in basal medium, basal medium supplemented with 20 µl/mL Phages-WC36, basal medium supplemented with 5 g/L starch, basal medium supplemented with 5 g/L starch and 20 µl/mL Phages-WC36. (B) Number of filamentous phages and hexagonal phages extracted from 1 µL of *P. stutzeri* 273 cell supernatant after culture in four different types of media as in panel A. The average numbers are shown at the top of the bar charts. (C) Continuous culture model of Phages-WC36 infected strain 273. The number of filamentous phages (D) and hexagonal phages (E) extracted from 1 µL of *P. stutzeri* 273 cell supernatant after three successive cultures. The average numbers are shown at the top of the bar charts.

In summary, we demonstrate for the first time that polysaccharides induce the

production of bacteriophages in *Lentisphaerae* strains WC36 and zth2, and these

bacteriophages contain AMGs encoding TonB-system energizer ExbB and CAZymes

that assist host metabolism and utilization of polysaccharide, thereby promoting host

growth. We also demonstrate that Phages-WC36 could infect and assist a marine

strain 273 to utilize polysaccharides, further expanding our understanding of the

significant role of deep-sea bacteriophages. Moreover, the study of deep-sea

bacteriophages has the potential to provide insight into the molecular evolution of

virus-host interactions in prokaryotes.

Materials and methods

Samples, media, and cultivation conditions

Deep-sea samples used for bacteria isolation in this study were collected from a

typical cold seep (E119º17'07.322", N22º06'58.598") at a depth of 1146 m by *RV*

KEXUE in July of 2018. Inorganic medium (including 1.0 g/L NH_4Cl, 1.0 g/L

$NaHCO_3$, 1.0 g/L CH_3COONa, 0.5 g/L KH_2PO_4, 0.2 g/L $MgSO_4·7H_2O$, 0.7 g/L

cysteine hydrochloride, 500 µl/L 0.1 % (w/v) resazurin, 1 L filtered seawater, pH 7.0)

supplemented with 1.0 g/L of different polysaccharides (laminarin, fucoidan, starch,

and alginate) under a 100% N_2 atmosphere was used to enrich microbes at 28 °C for

one month (Zheng et al., 2021a). The detailed isolation and purification steps were

performed as described previously (Zheng et al., 2021b). After pure cultures of

deep-sea bacteria were obtained, we cultured them in a rich medium (containing 1.0

g/L peptone, 5.0 g/L yeast extract, 1.0 g/L NH_4Cl, 1.0 g/L $NaHCO_3$, 1.0 g/L

CH_3COONa, 0.5 g/L KH_2PO_4, 0.2 g/L $MgSO_4·7H_2O$, 0.7 g/L cysteine hydrochloride,

500 µl/L 0.1 % (w/v) resazurin, 1 L filtered seawater, pH 7.0) or rich medium

supplemented with different polysaccharides for the future research.

Phylogenetic analysis

The 16S rRNA gene tree was constructed based on full-length 16S rRNA gene

sequences by the maximum likelihood method. The full-length 16S rRNA gene

sequences of *Lentisphaerae* strains WC36 and zth2 were obtained from their genomes, and the 16S rRNA gene sequences of other related taxa used for phylogenetic analysis were obtained from NCBI (www.ncbi.nlm.nih.gov/). The maximum likelihood phylogenetic trees of Phage-WC36-1 were constructed based on the zona occludens toxin (Zot) and single-stranded DNA-binding protein. The maximum likelihood phylogenetic tree of Phage-WC36-2 and Phage-zth2-2 was constructed based on the terminase large subunit protein (*terL*). These proteins used to construct the phylogenetic trees were all obtained from the NCBI databases. All the sequences were aligned by MAFFT version 7 (Katoh et al., 2019) and manually corrected. The phylogenetic trees were constructed using the W-IQ-TREE web server (http://iqtree.cibiv.univie.ac.at) (Trifinopoulos et al., 2016). Finally, we used the online tool Interactive Tree of Life (iTOL v5) (Letunic and Bork, 2021) to edit the tree.

Growth assay and transcriptomic analysis of *Lentisphaerae* strains WC36 and zth2 cultured in medium supplemented with laminarin

To assess the effect of polysaccharide on the growth of strain WC36, 100 µl of freshly incubated cells were inoculated in a 15 mL Hungate tube containing 10 mL rich medium supplemented with or without 5 g/L or 10 g/L laminarin (or starch). To assess the effect of laminarin on the growth of strain zth2, 100 µL of freshly incubated cells were inoculated in a 15 mL Hungate tube containing 10 mL rich medium supplemented with or without 3 g/L laminarin. Each condition was performed in three replicates. All the Hungate tubes were incubated at 28 °C. The bacterial growth status was monitored by measuring the OD_{600} value via a microplate reader (Infinite M1000 Pro; Tecan, Mannedorf, Switzerland) every day until cell growth reached the stationary phase.

For transcriptomic analysis, a cell suspension of strain WC36 was harvested after culture in 1.5 L rich medium either without supplementation or with 10 g/L laminarin for 5 days and 10 days. A cell suspension of strain zth2 was harvested after culture in rich medium either without supplementation or with 3 g/L laminarin for 4 days.

Thereafter, these collected samples were used for transcriptomic analysis by
Novogene (Tianjin, China). The detailed procedures of the transcriptomic assays
including library preparation, clustering, and sequencing, and data analyses are
described in the supplementary information.

Bacteriophages isolation

Phage isolation was as described previously with some modifications (Yamamoto et
al., 1970; Tseng et al., 1990; Kim and Blaschek, 1991). To induce the production of
bacteriophages, strain WC36 was inoculated in rich medium supplemented with or
without 10 g/L laminarin or 10 g/L starch, and strain zth2 was cultured in rich
medium supplemented with or without 3 g/L laminarin or 3 g/L starch. After 10 days
incubation, 60 mL of the different cultures was respectively collected and cells were
removed by centrifugation at $8,000 \times g$, 4 °C for 20 minutes three times. The
supernatant was filtered through a 0.22 μm millipore filter (Pall Supor, New York,
America) and subsequently 1 M NaCl was added to lyse the bacteria and separate
phage particles from these bacterial fragments. Then the supernatant was filtered
through a 0.22 μm millipore filter and collected by centrifugation at $8,000 \times g$, 4 °C
for 20 minutes, and phage particles were immediately precipitated with 100 g/L
polyethylene glycol (PEG8000) at 4 °C for 6 hours, and collected by centrifugation at
$10,000 \times g$, 4 °C for 20 minutes. The resulting phage particles were suspended in 2
mL SM buffer (0.01% gelatin, 50 mM Tris-HCl, 100 mM NaCl, and 10 mM $MgSO_4$),
then the suspension was extracted three times with an equal volume of chloroform
(Lin et al., 2012) and collected by centrifugation at $4,000 \times g$, 4 °C for 20 minutes.
Finally, clean phage particles were obtained.

Transmission electron microscopy (TEM)

To observe the morphology of bacteriophages, 10 μL phage virions were allowed to
adsorb onto a copper grid for 20 minutes, and then stained with phosphotungstic acid
for 30 seconds. Next, micrographs were taken with TEM (HT7700, Hitachi, Japan)
with a JEOL JEM 12000 EX (equipped with a field emission gun) at 100 kV.

To observe the morphology of strain WC36, cell suspension in rich medium

supplemented with or without polysaccharide was centrifuged at 5,000 × *g* for 10

minutes to obtain cell pellets. Subsequently, one part of the cell collection was

adsorbed to the copper grids for 20 minutes, then washed with 10 mM phosphate

buffer solution (PBS, pH 7.4) for 10 minutes and dried at room temperature for TEM

observation. Another part was first preserved in 2.5% glutaraldehyde for 24 hours at

4 °C, and then washed three times with PBS and dehydrated in ethanol solutions of

30%, 50%, 70%, 90%, and 100% for 10 minutes each time. Samples were further

fixed in 1% osmium tetroxide for 2 hours and embedded in plastic resin (Zechmann

and Zellnig, 2009; Fortunato et al., 2016). Thereafter, an ultramicrotome (Leica EM

UC7, Germany) was used to prepare ultrathin sections (50 nm) of cells, and the

obtained sections were stained with uranyl acetate and lead citrate (Graham and

Orenstein, 2007; Panphut et al., 2011). Finally, all samples were observed with the

JEOL JEM-1200 electron microscope as described above. In parallel, the filamentous

phage' number in each condition was respectively calculated.

Genome sequencing of bacteriophages

To sequence the genomes of bacteriophages, phage genomic DNA was extracted from

different purified phage particles. Briefly, 1 μg/mL DNase I and RNase A were added

to the concentrated phage solution for nucleic acid digestion overnight at 37 °C. The

digestion treatment was inactivated at 80 °C for 15 minutes, followed by extraction

with a Viral DNA Kit (Omega Bio-tek, USA) according to the manufacturer's

instructions. Then, genome sequencing was performed by Biozeron Biological

Technology Co. Ltd (Shanghai, China). The detailed process of library construction,

sequencing, genome assembly and annotation is described in the supplementary

information. In order to detect whether these bacteriophage genomes were located

within or outside of the host chromosomes, we used the new alignment algorithm

BWA-MEM (version 0.7.15) (Li, 2013) with default parameters to perform read

mapping of host WGS to these phages. In addition, we also evaluated the assembly

graph underlying the host consensus assemblies. Clean reads were mapped to the

bacterial complete genome sequences by Bowtie 2 (version 2.5.0) (Langmead and

Salzberg, 2012), BWA (version 0.7.8) (Li and Durbin, 2009), and SAMTOOLS

(version 0.1.18) (Li et al., 2009) with the default parameters.

Growth assay of *Pseudomonas stutzeri* 273 cultured in basal medium
supplemented with starch and Phages-WC36

549
550

In order to verify whether the bacteriophages induced from strain WC36 were able to

assist host metabolism and utilization of polysaccharides, we used *Pseudomonas*

stutzeri 273 isolated from deep-sea sediment (Wu et al., 2016). Specifically, 50 μL

freshly incubated cells (*P. stutzeri* 273) were inoculated in 5 mL basal medium (10 g/L

NaCl, 1 L sterilized distilled water), basal medium supplemented with 20 µl/L

Phages-WC36, basal medium supplemented with 5 g/L starch, basal medium

supplemented with 5 g/L starch and 20 µl/L Phages-WC36 for 24 hours at 28 °C,

respectively. The Phages-WC36 used in this growth assay were purified multiple

times and eventually suspended in SM buffer. Each condition was performed with

three replicates. Bacterial growth was monitored by measuring the OD_{600} value via a

microplate reader (Infinite M1000 Pro; Tecan, Mannedorf, Switzerland) every 4 hours

until cell growth reached the stationary phase. In addition, phages were extracted from

the supernatants of the above four media and their morphology was observed by TEM.

In parallel, filamentous phage and hexagonal phage numbers in each condition was

respectively calculated by randomly selecting TEM photos.

For transcriptomic analysis, we used cell suspensions of *P. stutzeri* 273 cultured

in basal medium supplemented with 20 µl/L Phages-WC36, basal medium

supplemented with 5 g/L starch, basal medium supplemented with 5 g/L starch and 20

µl/L Phages-WC36 for 24 hours at 28 °C. These cultures were harvested and sent to

Novogene (Tianjin, China) for transcriptome sequencing analysis. The detailed

procedures are described in the supplementary information.

Data availability

573 The complete genome sequences of strains WC36 and zth2 presented in this study

have been deposited in the GenBank database with accession numbers CP085689 and CP071032, respectively. The whole 16S rRNA gene sequences of strains WC36 and zth2 have been deposited in the GenBank database with accession numbers OK614042 and MW729759, respectively. The genome sequences of Phage-WC36-1 (Phage-zth2-1), Phage-WC36-2, and Phage-zth2-2 have been deposited in the GenBank database with accession numbers ON698679, OL791266, and OL791268, respectively. The raw sequencing reads from the transcriptomics analysis have been deposited to the NCBI Short Read Archive (accession numbers: PRJNA946146, PRJNA756144, and PRJNA865570).

Acknowledgements

This work was supported by the Science and Technology Innovation Project of Laoshan Laboratory (Grant No. LSKJ202203103; 2022QNLM030004-3), the NSFC Innovative Group Grant (No. 42221005), Shandong Provincial Natural Science Foundation (ZR2021ZD28), National Natural Science Foundation of China (Grant No. 92051107), Strategic Priority Research Program of the Chinese Academy of Sciences (Grant No. XDA22050301), China Ocean Mineral Resources R&D Association Grant (Grant No. DY135-B2-14), Key Collaborative Research Program of the Alliance of International Science Organizations (Grant No. ANSO-CR-KP-2022-08), China Ocean Mineral Resources R&D Association Grant (Grant No. DY295-B2-14), Key deployment projects of Center of Ocean Mega-Science of the Chinese Academy of Sciences (Grant No. COMS2020Q04), and Taishan Scholars Program for Chaomin Sun. The authors are grateful to the captain and crew of the R/V *KEXUE*, as well as the *FAXIAN* ROV team for assistance with sample collection.

Author contributions

CW, RZ, and CS conceived and designed the study; CW and RZ conducted most of the experiments; TZ helped to isolate the *Lentisphaerae* strain zth2; CW and CS led the writing of the manuscript; all authors contributed to and reviewed the manuscript.

Conflict of interest

The authors declare no competing interests.

References

Ackermann, H.W. (2007) **5500 Phages examined in the electron microscope** *Arch Virol* **152**:227-243

Ackermann, H.W. (2009) **Phage classification and characterization** *Methods Mol Biol* **501**:127-140

Afgan, E., Baker, D., Batut, B., van den Beek, M., Bouvier, D., Cech, M., Chilton, J., Clements, D., Coraor, N., Grüning, B.A., et al. (2018) **The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update** *Nucleic Acids Res* **46**:W537-W544

Ahlgren, N.A., Fuchsman, C.A., Rocap, G., and Fuhrman, J.A. (2019) **Discovery of several novel, widespread, and ecologically distinct marine *Thaumarchaeota* viruses that encode *amoC* nitrification genes** *ISME J* **13**:618-631

Alarcón-Schumacher, T., Naor, A., Gophna, U., and Erdmann, S. (2022) **Isolation of a virus causing a chronic infection in the archaeal model organism *Haloferax volcanii* reveals antiviral activities of a provirus** *Proc Natl Acad Sci USA* **119**:e2205037119

Anantharaman, K., Duhaime, M.B., Breier, J.A., Wendt, K.A., Toner, B.M., and Dick, G.J. (2014) **Sulfur oxidation genes in diverse deep-sea viruses** *Science* **344**:757-760

Bille, E., Zahar, J.R., Perrin, A., Morelle, S., Kriz, P., Jolley, K.A., Maiden, M.C., Dervin, C., Nassif, X., and Tinsley, C.R. (2005) **A chromosomally integrated bacteriophage in invasive meningococci** *J Exp Med* **201**:1905-1913

Breitbart, M. (2012) **Marine viruses: truth or dare** *Ann Rev Mar Sci* **4**:425-448

Brum, J.R., and Sullivan, M.B. (2015) **Rising to the challenge: accelerated pace of discovery transforms marine virology** *Nat Rev Microbiol* **13**:147-159

Brum, J.R., Ignacio-Espinoza, J.C., Kim, E.H., Trubl, G., Jones, R.M., Roux, S., VerBerkmoes, N.C., Rich, V.I., and Sullivan, M.B. (2016) **Illuminating structural proteins in viral "dark matter" with metaproteomics** *Proc Natl Acad Sci USA* **113**:2436-2441

Cantarel, B.L., Coutinho, P.M., Rancurel, C., Bernard, T., Lombard, V., and Henrissat, B. (2009) **The Carbohydrate-Active EnZymes database (CAZy): an expert resource for Glycogenomics** *Nucleic Acids Res* **37**:D233-238

Chevallereau, A., Pons, B.J., van Houte, S., and Westra, E.R. (2022) **Interactions between bacterial and phage communities in natural environments** *Nat Rev Microbiol* **20**:49-62

Davis, B.M., Lawson, E.H., Sandkvist, M., Ali, A., Sozhamannan, S., and Waldor, M.K. (2000) **Convergence of the secretory pathways for cholera toxin and the filamentous phage, CTXphi** *Science (New York, NY)* **288**:333-335

Du Toit, A. (2017) **Viral infection: The language of phages** *Nat Rev Microbiol*
15:134-135

Dwivedi, B., Xue, B., Lundin, D., Edwards, R.A., and Breitbart, M. (2013) **A**
bioinformatic analysis of ribonucleotide reductase genes in phage genomes and
metagenomes *BMC Evol Biol* **13**:33

Enav, H., Mandel-Gutfreund, Y., and Béjà, O. (2014) **Comparative metagenomic**
analyses reveal viral-induced shifts of host metabolism towards nucleotide
biosynthesis *Microbiome* **2**:9

Engelhardt, T., Orsi, W.D., and Jørgensen, B.B. (2015) **Viral activities and life**
cycles in deep subseafloor sediments *Environ Microbiol Rep* **7**:868-873

Falkowski, P.G., Barber, R.T., and Smetacek, V. (1998) **Biogeochemical controls**
and feedbacks on ocean primary production *Science* **281**:200-206

Faust, K., and Raes, J. (2012) **Microbial interactions: from networks to models** *Nat*
Rev Microbiol **10**:538-550

Foley, M.H., Cockburn, D.W., and Koropatkin, N.M. (2016) **The Sus operon: a**
model system for starch uptake by the human gut *Bacteroidetes* *Cell Mol Life Sci*
73:2603-2617

Fortunato, T.M., Beltrami, C., Emanueli, C., De Bank, P.A., and Pula, G. (2016)
Platelet lysate gel and endothelial progenitors stimulate microvascular network
formation in vitro: tissue engineering implications *Sci Rep* **6**:25326

Gazitúa, M.C., Vik, D.R., Roux, S., Gregory, A.C., Bolduc, B., Widner, B.,
Mulholland, M.R., Hallam, S.J., Ulloa, O., and Sullivan, M.B. (2021) **Potential**
virus-mediated nitrogen cycling in oxygen-depleted oceanic waters *ISME J*
15:981-998

Godkin, A., and Smith, K.A. (2017) **Chronic infections with viruses or parasites:**
breaking bad to make good *Immunology* **150**:389-396

Graham, L., and Orenstein, J.M. (2007) **Processing tissue and cells for transmission**
electron microscopy in diagnostic pathology and research *Nat Protoc* **2**:2439-2450

Hoffmann Berling, H., and Maze, R. (1964) **Release of male-specific**
bacteriophages from surviving host bacteria *Virology* **22**:305-313

Howard-Varona, C., Hargreaves, K.R., Abedon, S.T., and Sullivan, M.B. (2017)
Lysogeny in nature: mechanisms, impact and ecology of temperate phages *ISME*
J **11**:1511-1520

Howard-Varona, C., Lindback, M.M., Bastien, G.E., Solonenko, N., Zayed, A.A.,
Jang, H., Andreopoulos, B., Brewer, H.M., del Rio, T.G., Adkins, J.N., et al. (2020)
Phage-specific metabolic reprogramming of virocells *ISME J* **14**:881-895

Hurwitz, B.L., Hallam, S.J., and Sullivan, M.B. (2013) **Metabolic reprogramming**
by viruses in the sunlit and dark ocean *Genome Biol* **14**:R123

Jiang, S.C., and Paul, J.H. (1996) **Occurrence of lysogenic bacteria in marine**
microbial communities as determined by prophage induction *Mar Ecol Prog Ser*
142:27-38

Katoh, K., Rozewicki, J., and Yamada, K.D. (2019) **MAFFT online service:**
multiple sequence alignment, interactive sequence choice and visualization *Brief*
Bioinform **20**:1160-1166

Kim, A.Y., and Blaschek, H.P. (1991) **Isolation and characterization of a**
filamentous viruslike particle from *Clostridium acetobutylicum* **NCIB 6444** *J*
Bacteriol **173**:530-535

Knezevic, P., Adriaenssens, E.M., and Ictv Report, C. (2021) **ICTV Virus**
Taxonomy Profile: Inoviridae *J Gen Virol* **102**:

Koebnik, R. (2005) **TonB-dependent trans-envelope signalling: the exception or**
the rule? *Trends Microbiol* **13**:343-347

Krupovic, M., and Consortium, I.R. (2018) **ICTV virus taxonomy profile:**
Plasmaviridae J Gen Virol **99**:617-618

Langmead, B., and Salzberg, S.L. (2012) **Fast gapped-read alignment with Bowtie**
2 *Nat Methods* **9**:357-359

Letunic, I., and Bork, P. (2021) **Interactive Tree Of Life (iTOL) v5: an online tool**
for phylogenetic tree display and annotation *Nucleic Acids Res* **49**:W293-W296

Li, H. (2013) **Aligning sequence reads, clone sequences and assembly contigs with**
BWA-MEM 1303:

Li, H., and Durbin, R. (2009) **Fast and accurate short read alignment with**
Burrows-Wheeler transform *Bioinformatics* **25**:1754-1760

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G.,
Abecasis, G., and Durbin, R. (2009) **The Sequence Alignment/Map format and**
SAMtools *Bioinformatics* **25**:2078-2079

Li, Z., Pan, D., Wei, G., Pi, W., Zhang, C., Wang, J.H., Peng, Y., Zhang, L., Wang,
Y., Hubert, C.R.J., et al. (2021) **Deep sea sediments associated with cold seeps are**
a subsurface reservoir of viral diversity *ISME J* **15**:2366-2378

Lin, Y.R., Chiu, C.W., Chang, F.Y., and Lin, C.S. (2012) **Characterization of a new**
phage, termed phi A318, which is specific for *Vibrio alginolyticus Arch Virol*
157:917-926

Lindell, D., Jaffe, J.D., Coleman, M.L., Futschik, M.E., Axmann, I.M., Rector, T.,
Kettler, G., Sullivan, M.B., Steen, R., Hess, W.R., et al. (2007) **Genome-wide**
expression dynamics of a marine virus and host reveal features of co-evolution
Nature **449**:83-86

Liu, Y., Alexeeva, S., Bachmann, H., Guerra Martínez, J.A., Yeremenko, N., Abee,
T., and Smid, E.J. (2022) **Chronic release of tailless phage particles from**
Lactococcus lactis Appl Environ Microbiol **88**:e0148321

Lunin, V.V., Wang, H.-T., Bharadwaj, V.S., Alahuhta, M., Peña, M.J., Yang, J.-Y.,
Archer-Hartmann, S.A., Azadi, P., Himmel, M.E., Moremen, K.W., et al. (2020)
Molecular mechanism of polysaccharide acetylation by the arabidopsis xylan
O-acetyltransferase XOAT1 *Plant Cell* **32**:2367-2382

Mäntynen, S., Laanto, E., Oksanen, H.M., Poranen, M.M., and Díaz-Muñoz, S.L.
(2021) **Black box of phage-bacterium interactions: exploring alternative phage**
infection strategies *Open Biol* **11**:210188

Mai-Prochnow, A., Hui, J.G., Kjelleberg, S., Rakonjac, J., McDougald, D., and Rice,
S.A. (2015) **'Big things in small packages: the genetics of filamentous phage and**
effects on fitness of their host' *FEMS Microbiol Rev* **39**:465-487

Mann, N.H., Cook, A., Millard, A., Bailey, S., and Clokie, M. (2003) **Marine**
ecosystems: bacterial photosynthesis genes in a virus *Nature* **424**:741

Marvin, D.A., Symmons, M.F., and Straus, S.K. (2014) **Structure and assembly of**
filamentous bacteriophages *Prog Biophys Mol Bio* **114**:80-122

Mazaheri Nezhad Fard, R., Barton, M.D., and Heuzenroeder, M.W. (2010) **Novel**
bacteriophages in *Enterococcus* **spp** *Curr Microbiol* **60**:400-406

McKay, L.L., and Baldwin, K.A. (1973) **Induction of prophage in** *Streptococcus*
lactis **C2 by ultraviolet irradiation** *Appl Microbiol* **25**:682-684

Panphut, W., Senapin, S., Sriurairatana, S., Withyachumnarnkul, B., and Flegel, T.W.
(2011) **A novel integrase-containing element may interact with Laem-Singh virus**
(LSNV) to cause slow growth in giant tiger shrimp *BMC Vet Res* **7**:18

Putzrath, R.M., and Maniloff, J. (1977) **Growth of an enveloped mycoplasmavirus**
and establishment of a carrier state *J Virol* **22**:308-314

Roux, S., Krupovic, M., Daly, R.A., Borges, A.L., Nayfach, S., Schulz, F., Sharrar,
A., Matheus Carnevali, P.B., Cheng, J.F., Ivanova, N.N., et al. (2019) **Cryptic**
inoviruses revealed as pervasive in bacteria and archaea across Earth's biomes
Nat Microbiol **4**:1895-1906

Roux, S., Brum, J.R., Dutilh, B.E., Sunagawa, S., Duhaime, M.B., Loy, A., Poulos,
B.T., Solonenko, N., Lara, E., Poulain, J., et al. (2016) **Ecogenomics and potential**
biogeochemical impacts of globally abundant ocean viruses *Nature* **537**:689-693

Russel, M. (1991) **Filamentous phage assembly** *Mol Microbiol* **5**:1607-1613

Samson, J.E., Magadán, A.H., Sabri, M., and Moineau, S. (2013) **Revenge of the**
phages: defeating bacterial defences *Nat Rev Microbiol* **11**:675-687

Shkoporov, A.N., Turkington, C.J., and Hill, C. (2022) **Mutualistic interplay**
between bacteriophages and bacteria in the human gut *Nat Rev Microbiol*

Sullivan, M.B., Coleman, M.L., Weigele, P., Rohwer, F., and Chisholm, S.W. (2005)
Three *Prochlorococcus* **cyanophage genomes: signature features and ecological**
interpretations *PLoS Biol* **3**:e144

Sullivan, M.B., Lindell, D., Lee, J.A., Thompson, L.R., Bielawski, J.P., and
Chisholm, S.W. (2006) **Prevalence and evolution of core photosystem II genes in**
marine cyanobacterial viruses and their hosts *PLoS Biol* **4**:e234

Suttle, C.A. (2005) **Viruses in the sea** *Nature* **437**:356-361

Suttle, C.A. (2007) **Marine viruses--major players in the global ecosystem** *Nat Rev*
Microbiol **5**:801-812

Trifinopoulos, J., Nguyen, L.T., von Haeseler, A., and Minh, B.Q. (2016)
W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis
Nucleic Acids Res **44**:W232-W235

Tseng, Y.H., Lo, M.C., Lin, K.C., Pan, C.C., and Chang, R.Y. (1990)
Characterization of filamentous bacteriophage phi Lf from *Xanthomonas*
campestris **pv. campestris** *J Gen Virol* **71(Pt8)**:1881-1884

Wang, F., Wang, F., Li, Q., and Xiao, X. (2007) **A novel filamentous phage from**
the deep-sea bacterium *Shewanella piezotolerans* **WP3 is induced at low**
temperature *J Bacteriol* **189**:7151-7153

Weinbauer, M.G., Wilhelm, S.W., and Suttle, C.A. (1999) **Sunlight-lnduced DNA damage and resistance in natural viral communities** *Aquat Microb Ecol* **17**:111-120

Wigington, C.H., Sonderegger, D., Brussaard, C.P., Buchan, A., Finke, J.F., Fuhrman, J.A., Lennon, J.T., Middelboe, M., Suttle, C.A., Stock, C., et al. (2016) **Re-examination of the relationship between marine virus and microbial cell abundances** *Nat Microbiol* **1**:15024

Wu, S., Liu, G., Jin, W., Xiu, P., and Sun, C. (2016) **Antibiofilm and anti-Infection of a marine bacterial exopolysaccharide against *Pseudomonas aeruginosa* Front Microbiol** **7**:102

Yamamoto, K.R., Alberts, B.M., Benzinger, R., Lawhorne, L., and Treiber, G. (1970) **Rapid bacteriophage sedimentation in the presence of polyethylene glycol and its application to large-scale virus purification** *Virology* **40**:734-744

Yarza, P., Yilmaz, P., Pruesse, E., Glöckner, F.O., Ludwig, W., Schleifer, K.H., Whitman, W.B., Euzéby, J., Amann, R., and Rosselló-Móra, R. (2014) **Uniting the classification of cultured and uncultured bacteria and archaea using 16S rRNA gene sequences** *Nat Rev Microbiol* **12**:635-645

Zechmann, B., and Zellnig, G. (2009) **Rapid diagnosis of plant virus diseases by transmission electron microscopy** *J Virol Methods* **162**:163-169

Zeng, Q., and Chisholm, S.W. (2012) **Marine viruses exploit their host's two-component regulatory system in response to resource limitation** *Curr Biol* **22**:124-128

Zheng, R., Cai, R., Liu, R., Liu, G., and Sun, C. (2021a) *Maribellus comscasis* **sp. nov., a novel deep-sea *Bacteroidetes* bacterium, possessing a prominent capability of degrading cellulose** *Environ Microbiol* **23**:4561-4575

Zheng, R., Liu, R., Shan, Y., Cai, R., Liu, G., and Sun, C. (2021b) **Characterization of the first cultured free-living representative of *Candidatus* Izemoplasma uncovers its unique biology** *ISME J* **15**:2676-2691

Zimmerman, A.E., Howard-Varona, C., Needham, D.M., John, S.G., Worden, A.Z., Sullivan, M.B., Waldbauer, J.R., and Coleman, M.L. (2020) **Metabolic and biogeochemical consequences of viral infection in aquatic ecosystems** *Nat Rev Microbiol* **18**:21-34