Polysaccharides induce deep-sea *Lentisphaerae* strains to release chronic bacteriophages

Chong Wang[1,2,4], Rikuan Zheng[1,2,4], Tianhang Zhang[1,2,3,4], Chaomin Sun[1,2,3,4*]

[1]CAS and Shandong Province Key Laboratory of Experimental Marine Biology & Center of Deep Sea Research, Institute of Oceanology, Chinese Academy of Sciences, Qingdao, China.

[2]Laboratory for Marine Biology and Biotechnology, Qingdao National Laboratory for Marine Science and Technology, Qingdao, China.

[3]College of Earth Science, University of Chinese Academy of Sciences, Beijing, China.

[4]Center of Ocean Mega-Science, Chinese Academy of Sciences, Qingdao, China.

[*]Corresponding author

Chaomin Sun

Tel.: +86 532 82898857; fax: +86 532 82898857.

E-mail address: sunchaomin@qdio.ac.cn

Key words: deep-sea virus, polysaccharides, chronic bacteriophages, auxiliary metabolic genes, virus-host interaction

Running title: Induction of chronic bacteriophages by polysaccharides

Abstract

Viruses are ubiquitous in nature and play key roles in various ecosystems. Notably, some viruses (e.g. bacteriophage) exhibit alternative life cycles, such as chronic infections without cell lysis. However, the impact of chronic infections and their interactions with the host organisms remains largely unknown. Here, we found for the first time that polysaccharides induced the production of multiple temperate phages infecting two deep-sea *Lentisphaerae* strains (WC36 and zth2). Through physiological assays, genomic analysis, and transcriptomics assays, we found these bacteriophages were released via a chronic style without host cell lysis, which might reprogram host polysaccharide metabolism through the potential auxiliary metabolic genes (AMGs). The findings presented here, together with recent discoveries made on the reprogramming of host energy-generating metabolisms by chronic bacteriophages, shed light on the poorly explored marine virus-host interaction and bring us closer to understanding the potential role of chronic viruses in marine ecosystems.

Introduction

Viruses are the most abundant and genetically diverse biological entities on this planet (Suttle, 2007), and potentially play a role in shaping microbial abundances, community structure, and the evolutionary trajectory of core metabolisms (Sullivan et al., 2006; Samson et al., 2013; Zimmerman et al., 2020). The effect of bacteriophages (viruses that infect bacteria) on their specific hosts and within the microbial community is largely determined by their lifestyle. Traditionally, most bacteriophage life cycles are classified as being either lytic or lysogenic (Du Toit, 2017). During the lytic cycle, phages hijack the host's metabolic machinery for replication of their own genome and the production of progeny particles that are released through lysis. By contrast, during lysogeny, phages integrate their genomes into the host chromosome (or exist in the extrachromosomal form) and enter a dormant state, with the potential to re-enter a lytic cycle and release progeny by environmental changes at a later stage. Currently, more and more attention has been paid to chronic life cycles where bacterial growth continues despite phage reproduction (Hoffmann Berling and Maze, 1964), which was different from the lysogenic life cycle that could possibly lyse the host under some specific conditions. During a chronic cycle, progeny phage particles are released from host cells via extrusion or budding without killing the host (Putzrath and Maniloff, 1977; Russel, 1991; Marvin et al., 2014), instead of lysis of the host cell. Chronic infections are common among eukaryotic viruses (Godkin and Smith, 2017), however, to our best knowledge, only few phages have been described for prokaryotes in the pure isolates up to date (Roux et al., 2019; Alarcón-Schumacher et al., 2022; Liu et al., 2022). The best-studied examples of chronic infection are those from filamentous ssDNA (single-stranded DNA) phages that to date have primarily been identified in the order *Tubulavirales*, which currently involves three families (*Inoviridae*, *Paulinoviridae* and *Plectroviridae*) (Ackermann, 2007, 2009; Knezevic et al., 2021; Chevallereau et al., 2022). One of the most distinctive features of inoviruses is their ability to establish a chronic infection whereby their genomes may either integrate into the host chromosome or reside within the cell in a non-active form, and

inoviruses particles are continuously released without killing the host. In addition,

tailless phages enclosed in lipid membrane are also released from hosts during chronic

life cycles (Liu et al., 2022). Recent studies indicate that chronic cycles are more

widespread in nature than previously thought (Mäntynen et al., 2021), and the

abundance of inoviruses in ecosystem may have been far underestimated (Roux et al.,

2019). Therefore, a large percentage of phages in nature are proposed to replicate

through chronic life cycles, but whether chronic cycles are associated with specific

environments or ecological conditions remains to be thoroughly explored

(Chevallereau et al., 2022).

Marine ecosystems always influence the operating conditions for life on earth via

microbial interaction networks (Falkowski et al., 1998; Faust and Raes, 2012;

Wigington et al., 2016), which are modulated by viruses through impacting their

lifespan, gene flow and metabolic outputs (Suttle, 2005). A majority of these viruses

are bacteriophages, which exist widely in oceans and affect the life activities of

microbes (Breitbart, 2012; Roux et al., 2016; Gregory et al., 2019; Dominguez-Huerta

et al., 2022). In addition to core viral genes encoding viral structural proteins (Brum et

al., 2016), bacteriophages also encode various auxiliary metabolic genes (AMGs)

(Brum and Sullivan, 2015) that provide supplemental support to key steps of host

metabolism. For instance, AMGs of marine bacteriophages have been predicted to be

involved in photosynthesis (Mann et al., 2003), nitrogen cycling (Ahlgren et al., 2019;

Gazitúa et al., 2021), sulfur cycling (Anantharaman et al., 2014; Roux et al., 2016),

phosphorus cycling (Zeng and Chisholm, 2012), nucleotide metabolism (Sullivan et

al., 2005; Dwivedi et al., 2013; Enav et al., 2014), and almost all central carbon

metabolisms in host cells (Hurwitz et al., 2013). However, AMGs of chronic phages

have not been reported. Thus, it is worth exploring whether chronic phages could

carry AMGs and assist host metabolism during chronic infection.

The deep sea harbors abundant and undiscovered viruses, which potentially

control the metabolism of microbial hosts and influence biogeochemical cycling (Li et

al., 2021). However, due to the vast majority of deep-sea microbes cannot be

cultivated in the laboratory, most bacteriophages could not be isolated. Thus, it is of

great significance to further identify unknown phages in the deep sea and explore their relationship with microbial hosts and even marine ecosystems. Here, we report that polysaccharides can induce deep-sea *Lentisphaerae* bacteria (difficult-to-cultivate microorganisms) to release some chronic bacteriophages. These chronic bacteriophages might assist host polysaccharides metabolism via corresponding potential AMGs.

Results and Discussion

Polysaccharides promote the growth of deep-sea *Lentisphaerae* strain WC36 and stimulate the expression of phage-associated genes

As a primary carbon source, polysaccharides are a ubiquitous energy source for microorganisms in both terrestrial and marine ecosystems (Zheng et al., 2021a). In our previous work, we successfully isolated a novel *Bacteroidetes* species through a polysaccharide degradation-driven strategy from the deep-sea cold seep (Zheng et al., 2021a). Of note, using the same approach and deep-sea sample replicates, we also cultured a bacterial strain WC36 that was identified as a member of the phylum *Lentisphaerae* by 16S rRNA gene sequence-based phylogenetic analysis (Fig. 1A). As expected, growth assays showed that two kinds of polysaccharides including laminarin and starch could promote strain WC36 growth (Fig. 1B). In particular, supplementation of 10 g/L laminarin and 10 g/L starch in rich medium resulted in ~7-fold and ~4-fold growth promotion respectively compared to cultivation in rich medium (Fig. 1B).

To explore the reasons behind this significant growth promotion by polysaccharide supplementation, we performed a transcriptomic analysis of strain WC36 cultured in rich medium supplemented either with or without 10 g/L laminarin for 5 days and 10 days. In addition to the up-regulation of genes related to glycan transport and degradation, when 10 g/L laminarin was added in the rich medium, the most upregulated genes were phage-associated (e. g. phage integrase, phage portal protein) (Fig. 1C and Supplementary Table 1), which were expressed at the background level in the rich medium alone. Consistently, RT-qPCR results (Fig. 1D)

141 confirmed upregulation of some genes encoding phage-associated proteins, as shown

142 in Fig. 1C. Therefore, we speculate that the metabolism of polysaccharides in strain

143 WC36 might be closely connected with the phage production.

Polysaccharides induce the production of bacteriophages in *Lentisphaerae* strain

WC36

To test our hypothesis that polysaccharides might induce strain WC36 to release

bacteriophages, we grew strain WC36 in rich medium supplemented with or without

laminarin or starch and isolated bacteriophages from the cell suspension supernatant

according to established standard protocols (Lin et al., 2012). Based on the growth

curve of strain WC36, we found that the growth rate of strictly anaerobic strain WC36

was relatively slow. Therefore, different to the typical sampling time (24 h) for

bacteriophage isolation from other bacteria (Jiang and Paul, 1996; Weinbauer et al.,

1999), we selected a longer sampling time (10 days) to extract bacteriophages. TEM

observation showed that many different shapes of phage-like structures indeed existed

in the supernatant of WC36 cells cultured in rich medium supplemented with

laminarin (Fig. 2A, panels II-IV) or starch (Fig. 2A, panels V-VIII). In contrast, we

did not observe any phage-like structures in the supernatant of WC36 cells cultured in

rich medium (Fig. 2A, panel I). We also tested and confirmed that there were not any

phage-like structures in rich medium supplemented with 10 g/L laminarin alone

(Supplementary Fig. 1A) or in 10 g/L starch alone (Supplementary Fig. 1B), ruling

out the possibility of phage contamination from the polysaccharides (laminarin/

starch). These results suggest that polysaccharides indeed stimulate the production of

various bacteriophages from strain WC36. Correspondingly, in the presence of

laminarin, different forms of bacteriophages were observed by TEM, with filamentous

ones dominant (Fig. 2A). The length of these filamentous phages was about ~0.4 μm

to ~8.0 μm (Fig. 2A, panel II), and the size of the hexagonal phages was less than 100

nm (Fig. 2A, panel IV). In the presence of starch, in addition to filamentous and

hexagonal phages (Fig. 2A, panels V, VI, and VIII), we also observed a kind of

icosahedral *Microviridae*-like phage with a diameter of 25-30 nm (Fig. 2A, panel

VII).

Given that we found bacteriophages in cultures of strain WC36, we next sought

to explore whether bacteriophages adhering to bacterial cells could be observed. To

this end, we checked the morphology of strain WC36 in the absence or presence of

polysaccharides. As expected, with TEM we observed many filamentous phage-like

structures around strain WC36 cells cultivated in rich medium supplemented with

laminarin (Fig. 2B, panel II) and starch (Fig. 2B, panels III-IV). In contrast, we could

not find any phage-like structures around strain WC36 cells cultivated in rich medium

(Fig. 2B, panel I). Meanwhile, we also checked the polysaccharides (laminarin/ starch)

in rich medium directly by TEM and did not find any phage-like structures

(Supplementary Fig. 2). Moreover, based on TEM observation of ultrathin sections of

strain WC36 cultured in medium supplemented with polysaccharide, we even

observed filamentous phages that were being released from or entering into host cells

via a kind of extrusion or budding structure around bacterial cells (Fig. 2C, panels

II-IV; Supplementary Fig. 3B, panels II-III), which were typical cellular structures in

host cells used for chronic bacteriophage release (Marvin et al., 2014; Krupovic and

Consortium, 2018). In contrast, phage-like structures with extrusions or buddings

were not found in cells of strain WC36 cultivated in rich medium (Fig. 2C, panel I;

Supplementary Fig. 3B, panel I). Hence, these results collectively suggest that

filamentous bacteriophages induced from deep-sea strain WC36 replicate in a chronic

manner as reported previously (Chevallereau et al., 2022; Liu et al., 2022). However,

the entry and exit of the hexagonal phages into the WC36 cells were not observed.

To further verify the release style of bacteriophages from strain WC36 in the

presence of polysaccharide, we prolonged the incubation time of this strain up to 30

days in medium supplemented with 5 g/L or 10 g/L laminarin. Strain WC36 showed

stable growth after it entered the stationary phase (Fig. 3A). Regardless of whether the

laminarin was present, the bacterial cells kept their cell shape intact, indicating they

were still healthy after 30 days (Supplementary Fig. 3A, panels I-III). Apparently, the

replication and release of bacteriophages in strain WC36 did not kill the host cell,

consistent with the key feature of chronic bacteriophages (Howard-Varona et al.,

2017). With an increase of culture time (5, 10, or 30 days) and laminarin

concentration (5 or 10 g/L), the number of released bacteriophages increased

(Supplementary Fig. 3A, panels IV-VI; Supplementary Fig. 4, panels IV-IX).

Specifically, the average number per microliter of filamentous phages (9.7, 29 or 65.3)

extracted from the supernatant of strain WC36 cultured in rich medium supplemented

with 10 g/L laminarin for 5, 10 or 30 days was higher than that cultured in rich

medium supplemented with 5 g/L laminarin (4.3, 13.7 or 35.3) (Fig. 3B). The average

number per microliter of hexagonal phages (9, 30, 46.7) extracted from the

supernatant of strain WC36 cultured in rich medium supplemented with 10 g/L

laminarin for 5, 10 or 30 days was higher than that cultured in rich medium

supplemented with 5 g/L laminarin (4, 11.3 or 17.7) (Fig. 3C).

Polysaccharides promote the growth of deep-sea *Lentisphaerae* strain zth2 and

induce bacteriophage production

To explore whether the production of bacteriophages induced by polysaccharide is an

individual case, we further checked the effect of polysaccharides on another cultured

deep-sea *Lentisphaerae* strain zth2. Strain zth2 was isolated by the same

polysaccharide degradation-driven strategy as that used for strain WC36 (Zhang et al.,

2022). 16S rRNA gene sequence similarity calculation using the NCBI server

indicated that the closest relative of strain zth2 was strain WC36 (90.4%), which was

lower than the threshold sequence identity (94.5%) for distinct genera (Yarza et al.,

2014). Based on a maximum likelihood tree and similarity of 16S rRNA gene

sequences, we propose that strain zth2 and WC36 are the type strains of two different

novel genera, both belonging to the *Lentisphaeraceae* family, *Lentisphaerales* order,

Lentisphaeria class. Consistently, laminarin indeed promoted the growth of strain zth2

(Fig. 4A). Analysis of the differential transcriptome and RT-qPCR of strain zth2

cultured in either unsupplemented medium or with laminarin showed that many genes

encoding critical factors associated with the phage life cycle were up-regulated

(ranging from 4 to 200 fold) (Fig. 4B, C and Supplementary Table 2), suggesting that

phages might also be involved in the utilization of laminarin by strain zth2, the same

as in strain WC36.

To further confirm the existence of bacteriophages in the supernatant of strain

zth2 cell suspension cultured in medium supplemented with polysaccharide, we

performed bacteriophage isolation from the supernatant of strain zth2 cultures

followed by TEM. Indeed, many filamentous-, hexagonal- and icosahedral phages

were clearly observed in the culture of strain zth2 grown in the presence of laminarin

(Fig. 4D, panels I-III) and starch (Fig. 4D, panels IV-V), but not in rich medium short

of polysaccharide (Fig. 4D, panel VI). This suggests that bacteriophages were induced

by supplementation of polysaccharide to strain zth2.

In addition, transcriptome and RT-qPCR results showed that genes encoding

bacterial secretion system-related proteins were up-regulated in strains WC36

(Supplementary Fig. 5A, B) and zth2 (Supplementary Fig. 5C, D) cultured in medium

supplemented with laminarin. It has been reported that filamentous phages utilize host

cell secretion systems for their own egress from bacterial cells (Davis et al., 2000;

Bille et al., 2005). Therefore, the filamentous phages induced from strains WC36 and

zth2 might also get in and out of host cells using bacterial secretion systems. Taken

together, we conclude that laminarin effectively promotes the growth of deep-sea

Lentisphaerae strains WC36 and zth2, and induces the production of bacteriophages.

Distinct from previous reports that bacteriophages are induced under traditional

conditions such as low temperature (Wang et al., 2007), mitomycin C (Mazaheri

Nezhad Fard et al., 2010), and ultraviolet irradiation (McKay and Baldwin, 1973),

here, we report for the first time that polysaccharides induce the production of

bacteriophages.

Bacteriophages reprogram the polysaccharide metabolism of *Lentisphaerae*

strains WC36 and zth2

In this study, we clearly show that polysaccharide effectively promotes bacterial

growth and simultaneously induces the production of bacteriophages, suggesting a

close relationship between bacteriophages and host polysaccharide metabolism.

Therefore, we sought to ask whether bacteriophages could reprogram polysaccharide

metabolism in deep-sea bacteria. For this purpose, we sequenced the genomes of the bacteriophages induced by polysaccharides in *Lentisphaerae* strains WC36 and zth2. Eventually, two incomplete assembled genomes (Phage-WC36-1, 6.3 kb; Phage-WC36-2, 28.3 kb) were obtained from the bacteriophages of strain WC36 (Fig. 5A, B and Supplementary Tables 3, 4). Meanwhile, two incomplete assembled genomes (Phage-zth2-1, 6.3 kb; Phage-zth2-2, 40.4 kb) were obtained from the phages of strain zth2 (Fig. 5A, C and Supplementary Tables 3, 5). Subsequently, we carefully compared the bacteriophage genomes with those of the corresponding hosts (strains WC36 and zth2) using Galaxy Version 2.6.0 (https://galaxy.pasteur.fr/) (Afgan et al., 2018) with the NCBI BLASTN method and used BWA-mem software for read mapping from host whole genome sequencing (WGS) to these bacteriophages. These analyses both showed that the bacteriophage genomes are completely outside of the host chromosomes. Through comparison of genome sequence (Supplementary Fig. 6), IMG database analysis, and phylogenetic analysis (Supplementary Fig. 7), we confirmed that Phage-WC36-1 was the same as Phage-zth2-1 (inoviruses, belonging to the family *Inoviridae*, order *Tubulavirales*, class *Faserviricetes*), indicating that inoviruses might be common in the *Lentisphaerae* phylum. Inovirus members were the dominant population in the active group of viruses from deep-sea sediments, and encode some proteins contributing to host phenotypes and ultimately alter the fitness and other behaviors of their hosts (Engelhardt et al., 2015; Mai-Prochnow et al., 2015). A maximum likelihood tree showed that both Phage-WC36-2 and Phage-zth2-2 belonged to the class *Caudoviricetes* (Supplementary Fig. 8). In addition to genes encoding phage-associated proteins within the assembled bacteriophage genomes, there are also many genes encoding proteins associated with polysaccharide transport and degradation, which were potential AMGs (Shaffer et al., 2020; Pratama et al., 2021). These polysaccharide metabolizing proteins include TonB-system energizer ExbB (Koebnik, 2005), O-polysaccharide acetyltransferase (Lunin et al., 2020), glucosaminidase, glycoside hydrolase family 127 protein (best BLASTP hit e-value 3e-25), and transglycosylase (Fig. 5). GH127 is one of the CAZymes, which are specifically responsible for degrading polysaccharides (Cantarel et al., 2009).

Energizer ExbB is an important part of the TonB-system, which is often present in

Bacteroides and responsible for transport of oligosaccharides from the outer

membrane into the periplasm (Foley et al., 2016). However, it is absent in the strain

WC36 and zth2 genomes. In natural ecosystems, viral infections indirectly influence

microbial metabolic fluxes, metabolic reprogramming, and energy homeostasis of

host cells (Howard-Varona et al., 2020). For example, cyanoviruses possess AMGs

encoding core photosynthetic reaction centers (Sullivan et al., 2006) which are highly

expressed during infection, thus stimulating photosynthesis and enhancing viral

abundance (Lindell et al., 2007). Therefore, these potential AMGs encoding

TonB-system energizer ExbB and glycoside hydrolases might reprogram the

polysaccharide metabolism of host cells (strains WC36 and zth2) by providing

supplemental support to key metabolic processes. The above results provide stronger

evidence that the relationship between Phages-WC36 and their host is not a fight to

the death between enemies, but in contrast a mutualistic relationship between partners

(Shkoporov et al., 2022). Overall, bacteriophages induced by polysaccharides contain

potential AMGs associated with polysaccharide degradation, which may play a key

role in the degradation and utilization of polysaccharide by host cells.

Moreover, it was recently demonstrated that selfish bacteria, which were

common throughout the water column of the ocean, could bind, partially hydrolyze,

and transport polysaccharides into the periplasmic space without loss of hydrolysis

products (Reintjes et al., 2017; Giljan et al., 2023). Based on our results, we

hypothesized that these chronic phages might also enter the host through this

"selfishness" mechanism while assisting the host in metabolizing polysaccharides,

thus not lysing the host. On the other hand, these chronic phages might hijack this

"selfishness" mechanism to improve their infectivity and entry, rather than helping

their hosts to grow and proliferate, so they could reap the benefits of simply having

more hosts to infect. In the future, we need to construct a genetic operating system of

the strictly anaerobic host strain WC36 to detailedly reveal the relationship between

chronic phage and host.

Materials and methods

Samples, media, and cultivation conditions

Deep-sea samples used for bacteria isolation in this study were collected from a typical cold seep (E119º17'07.322", N22º06'58.598") at a depth of 1146 m by *RV KEXUE* in July of 2018. Inorganic medium (including 1.0 g/L NH_4Cl, 1.0 g/L $NaHCO_3$, 1.0 g/L CH_3COONa, 0.5 g/L KH_2PO_4, 0.2 g/L $MgSO_4 \cdot 7H_2O$, 0.7 g/L cysteine hydrochloride, 500 µl/L 0.1 % (w/v) resazurin, 1 L filtered seawater, pH 7.0) supplemented with 1.0 g/L of different polysaccharides (laminarin, fucoidan, starch, and alginate) under a 100% N_2 atmosphere was used to enrich microbes at 28 °C for one month (Zheng et al., 2021a). The detailed isolation and purification steps were performed as described previously (Zheng et al., 2021b). After pure cultures of deep-sea bacteria were obtained, we cultured them in a rich medium (containing 1.0 g/L peptone, 5.0 g/L yeast extract, 1.0 g/L NH_4Cl, 1.0 g/L $NaHCO_3$, 1.0 g/L CH_3COONa, 0.5 g/L KH_2PO_4, 0.2 g/L $MgSO_4 \cdot 7H_2O$, 0.7 g/L cysteine hydrochloride, 500 µl/L 0.1 % (w/v) resazurin, 1 L filtered seawater, pH 7.0) or rich medium supplemented with different polysaccharides for the future research.

Genome sequencing, annotation, and analysis of strains WC36 and zth2

For genomic sequencing, strains WC36 and zth2 were grown in the liquid rich medium supplemented with 5 g/L laminarin and starch and harvested after one week of incubation at 28 °C. Genomic DNA was isolated by using the PowerSoil DNA isolation kit (Mo Bio Laboratories Inc., Carlsbad, CA). Thereafter, the genome sequencing was carried out with both the Illumina NovaSeq PE150 (San Diego, USA) and Nanopore PromethION platform (Oxford, UK) at the Beijing Novogene Bioinformatics Technology Co., Ltd. A complete description of the library construction, sequencing, and assembly was performed as previously described (Zheng et al., 2021b). We used seven databases to predict gene functions, including Pfam (Protein Families Database, http://pfam.xfam.org/), GO (Gene Ontology, http://geneontology.org/) (Ashburner et al., 2000), KEGG (Kyoto Encyclopedia of Genes and Genomes, http://www.genome.jp/kegg/) (Kanehisa et al., 2004), COG

(Clusters of Orthologous Groups, http://www.ncbi.nlm.nih.gov/COG/) (Galperin et al., 2015), NR (Non-Redundant Protein Database databases), TCDB (Transporter Classification Database), and Swiss-Prot (http://www.ebi.ac.uk/uniprot/) (Bairoch and Apweiler, 2000). A whole genome Blast search (E-value less than 1e-5, minimal alignment length percentage larger than 40%) was performed against above seven databases.

Phylogenetic analysis

The 16S rRNA gene tree was constructed based on full-length 16S rRNA gene sequences by the maximum likelihood method. The full-length 16S rRNA gene sequences of *Lentisphaerae* strains WC36 and zth2 were obtained from their genomes, and the 16S rRNA gene sequences of other related taxa used for phylogenetic analysis were obtained from NCBI (www.ncbi.nlm.nih.gov/). All the accession numbers were listed in the Supplementary Table 6. The maximum likelihood phylogenetic trees of Phage-WC36-1 were constructed based on the zona occludens toxin (Zot) and single-stranded DNA-binding protein (Zeng et al., 2021; Evseev et al., 2023). The maximum likelihood phylogenetic tree of Phage-WC36-2 and Phage-zth2-2 was constructed based on the terminase large subunit protein (*terL*). These proteins used to construct the phylogenetic trees were all obtained from the NCBI databases. All the sequences were aligned by MAFFT version 7 (Katoh et al., 2019) and manually corrected. The phylogenetic trees were constructed using the W-IQ-TREE web server (http://iqtree.cibiv.univie.ac.at) with the "GTR+F+I+G4" model (Trifinopoulos et al., 2016). Finally, we used the online tool Interactive Tree of Life (iTOL v5) (Letunic and Bork, 2021) to edit the tree.

Growth assay and transcriptomic analysis of *Lentisphaerae* strains WC36 and zth2 cultured in medium supplemented with laminarin

To assess the effect of polysaccharide on the growth of strain WC36, 100 μl of freshly incubated cells were inoculated in a 15 mL Hungate tube containing 10 mL rich medium supplemented with or without 5 g/L or 10 g/L laminarin (or starch). To assess

the effect of laminarin on the growth of strain zth2, 100 µL of freshly incubated cells

were inoculated in a 15 mL Hungate tube containing 10 mL rich medium

supplemented with or without 3 g/L laminarin. Each condition was performed in three

replicates. All the Hungate tubes were incubated at 28 °C. The bacterial growth status

was monitored by measuring the OD_{600} value via a microplate reader (Infinite M1000

Pro; Tecan, Mannedorf, Switzerland) every day until cell growth reached the

stationary phase.

For transcriptomic analysis, a cell suspension of strain WC36 was harvested after

culture in 1.5 L rich medium either without supplementation or with 10 g/L laminarin

for 5 days and 10 days. A cell suspension of strain zth2 was harvested after culture in

rich medium either without supplementation or with 3 g/L laminarin for 4 days.

Thereafter, these collected samples were used for transcriptomic analysis by

Novogene (Tianjin, China). The detailed procedures of the transcriptomic assays

including library preparation, clustering, and sequencing, and data analyses are

described in the supplementary information.

Bacteriophages isolation

Phage isolation was as described previously with some modifications (Yamamoto et

al., 1970; Tseng et al., 1990; Kim and Blaschek, 1991). To induce the production of

bacteriophages, strain WC36 was inoculated in rich medium supplemented with or

without 10 g/L laminarin or 10 g/L starch, and strain zth2 was cultured in rich

medium supplemented with or without 3 g/L laminarin or 3 g/L starch. After 10 days

incubation, 60 mL of the different cultures was respectively collected and cells were

removed by centrifugation at 8,000 $\times$ *g*, 4 °C for 20 minutes three times. The

supernatant was filtered through a 0.22 µm millipore filter (Pall Supor, New York,

America) and subsequently 1 M NaCl was added to lyse the bacteria and separate

phage particles from these bacterial fragments. Then the supernatant was filtered

through a 0.22 µm millipore filter and collected by centrifugation at 8,000 $\times$ *g*, 4 °C

for 20 minutes, and phage particles were immediately precipitated with 100 g/L

polyethylene glycol (PEG8000) at 4 °C for 6 hours, and collected by centrifugation at

10,000 × *g*, 4 °C for 20 minutes. The resulting phage particles were suspended in 2

mL SM buffer (0.01% gelatin, 50 mM Tris-HCl, 100 mM NaCl, and 10 mM MgSO$_4$),

then the suspension was extracted three times with an equal volume of chloroform

(Lin et al., 2012) and collected by centrifugation at 4,000 × *g*, 4 °C for 20 minutes.

Finally, clean phage particles were obtained.

Transmission electron microscopy (TEM)

To observe the morphology of bacteriophages, 10 μL phage virions were allowed to

adsorb onto a copper grid for 20 minutes, and then stained with phosphotungstic acid

for 30 seconds. Next, micrographs were taken with TEM (HT7700, Hitachi, Japan)

with a JEOL JEM 12000 EX (equipped with a field emission gun) at 100 kV.

To observe the morphology of strain WC36, cell suspension in rich medium

supplemented with or without polysaccharide was centrifuged at 5,000 × *g* for 10

minutes to obtain cell pellets. Subsequently, one part of the cell collection was

adsorbed to the copper grids for 20 minutes, then washed with 10 mM phosphate

buffer solution (PBS, pH 7.4) for 10 minutes and dried at room temperature for TEM

observation. Another part was first preserved in 2.5% glutaraldehyde for 24 hours at

4 °C, and then washed three times with PBS and dehydrated in ethanol solutions of

30%, 50%, 70%, 90%, and 100% for 10 minutes each time. Samples were further

fixed in 1% osmium tetroxide for 2 hours and embedded in plastic resin (Zechmann

and Zellnig, 2009; Fortunato et al., 2016). Thereafter, an ultramicrotome (Leica EM

UC7, Germany) was used to prepare ultrathin sections (50 nm) of cells, and the

obtained sections were stained with uranyl acetate and lead citrate (Graham and

Orenstein, 2007; Panphut et al., 2011). Finally, all samples were observed with the

JEOL JEM-1200 electron microscope as described above. In parallel, the phage'

number in each condition was respectively calculated by ten random TEM images.

Genome sequencing of bacteriophages

To sequence the genomes of bacteriophages, phage genomic DNA was extracted from

different purified phage particles. Briefly, 1 μg/mL DNase I and RNase A were added

to the concentrated phage solution for nucleic acid digestion overnight at 37 °C. The digestion treatment was inactivated at 80 °C for 15 minutes, followed by extraction with a Viral DNA Kit (Omega Bio-tek, USA) according to the manufacturer's instructions. Then, genome sequencing was performed by Biozeron Biological Technology Co. Ltd (Shanghai, China). The detailed process of library construction, sequencing, genome assembly and annotation is described in the supplementary information. In order to detect whether these bacteriophage genomes were located within or outside of the host chromosomes, we used the new alignment algorithm BWA-MEM (version 0.7.15) (Li, 2013) with default parameters to perform read mapping of host WGS to these phages. In addition, we also evaluated the assembly graph underlying the host consensus assemblies. Clean reads were mapped to the bacterial complete genome sequences by Bowtie 2 (version 2.5.0) (Langmead and Salzberg, 2012), BWA (version 0.7.8) (Li and Durbin, 2009), and SAMTOOLS (version 0.1.18) (Li et al., 2009) with the default parameters.

Data availability

The complete genome sequences of strains WC36 and zth2 presented in this study have been deposited in the GenBank database with accession numbers CP085689 and CP071032, respectively. The whole 16S rRNA gene sequences of strains WC36 and zth2 have been deposited in the GenBank database with accession numbers OK614042 and MW729759, respectively. The genome sequences of Phage-WC36-1 (Phage-zth2-1), Phage-WC36-2, and Phage-zth2-2 have been deposited in the GenBank database with accession numbers PP701473, OL791266, and OL791268, respectively. The raw sequencing reads from the transcriptomics analysis have been deposited to the NCBI Short Read Archive (accession numbers: PRJNA946146, PRJNA756144, and PRJNA865570).

Acknowledgements

This work was funded by the Science and Technology Innovation Project of Laoshan Laboratory (Grant nos. 2022QNLM030004-3 and LSKJ202203103), the NSFC

Innovative Group Grant (No. 42221005), Major Research Plan of the National Natural Science Foundation (Grant no. 92351301), Shandong Provincial Natural Science Foundation (Grant nos. ZR2021ZD28 and ZR2023QD010), Strategic Priority Research Program of the Chinese Academy of Sciences (Grant no. XDA22050301), Key Collaborative Research Program of the Alliance of International Science Organizations (Grant no. ANSO-CR-KP-2022-08), and the Taishan Scholars Program (Grant nos. tstp20230637 and tsqn202312264).

Author contributions

CW, RZ, and CS conceived and designed the study; CW and RZ conducted most of the experiments; TZ helped to isolate the *Lentisphaerae* strain zth2; CW and CS led the writing of the manuscript; all authors contributed to and reviewed the manuscript.

Conflict of interest

The authors declare no competing interests.

References

Ackermann, H.W. (2007) **5500 Phages examined in the electron microscope** *Arch Virol* **152**:227-243

Ackermann, H.W. (2009) **Phage classification and characterization** *Methods Mol Biol* **501**:127-140

Afgan, E., Baker, D., Batut, B., van den Beek, M., Bouvier, D., Cech, M., Chilton, J., Clements, D., Coraor, N., Grüning, B.A., et al. (2018) **The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update** *Nucleic Acids Res* **46**:W537-W544

Ahlgren, N.A., Fuchsman, C.A., Rocap, G., and Fuhrman, J.A. (2019) **Discovery of several novel, widespread, and ecologically distinct marine *Thaumarchaeota* viruses that encode *amoC* nitrification genes** *ISME J* **13**:618-631

Alarcón-Schumacher, T., Naor, A., Gophna, U., and Erdmann, S. (2022) **Isolation of a virus causing a chronic infection in the archaeal model organism *Haloferax volcanii* reveals antiviral activities of a provirus** *Proc Natl Acad Sci USA* **119**:e2205037119

Anantharaman, K., Duhaime, M.B., Breier, J.A., Wendt, K.A., Toner, B.M., and Dick, G.J. (2014) **Sulfur oxidation genes in diverse deep-sea viruses** *Science* **344**:757-760

Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis,
A.P., Dolinski, K., Dwight, S.S., Eppig, J.T., et al. (2000) **Gene ontology: tool for**
the unification of biology. The Gene Ontology Consortium *Nat Genet* **25**:25-29

Bairoch, A., and Apweiler, R. (2000) **The SWISS-PROT protein sequence**
database and its supplement TrEMBL in 2000 *Nucleic Acids Res* **28**:45-48

Bille, E., Zahar, J.R., Perrin, A., Morelle, S., Kriz, P., Jolley, K.A., Maiden, M.C.,
Dervin, C., Nassif, X., and Tinsley, C.R. (2005) **A chromosomally integrated**
bacteriophage in invasive meningococci *J Exp Med* **201**:1905-1913

Breitbart, M. (2012) **Marine viruses: truth or dare** *Ann Rev Mar Sci* **4**:425-448

Brum, J.R., and Sullivan, M.B. (2015) **Rising to the challenge: accelerated pace of**
discovery transforms marine virology *Nat Rev Microbiol* **13**:147-159

Brum, J.R., Ignacio-Espinoza, J.C., Kim, E.H., Trubl, G., Jones, R.M., Roux, S.,
VerBerkmoes, N.C., Rich, V.I., and Sullivan, M.B. (2016) **Illuminating structural**
proteins in viral "dark matter" with metaproteomics *Proc Natl Acad Sci USA*
113:2436-2441

Cantarel, B.L., Coutinho, P.M., Rancurel, C., Bernard, T., Lombard, V., and
Henrissat, B. (2009) **The Carbohydrate-Active EnZymes database (CAZy): an**
expert resource for Glycogenomics *Nucleic Acids Res* **37**:D233-238

Chevallereau, A., Pons, B.J., van Houte, S., and Westra, E.R. (2022) **Interactions**
between bacterial and phage communities in natural environments *Nat Rev*
Microbiol **20**:49-62

Davis, B.M., Lawson, E.H., Sandkvist, M., Ali, A., Sozhamannan, S., and Waldor,
M.K. (2000) **Convergence of the secretory pathways for cholera toxin and the**
filamentous phage, CTXphi *Science* **288**:333-335

Dominguez-Huerta, G., Zayed, A.A., Wainaina, J.M., Guo, J., Tian, F., Pratama,
A.A., Bolduc, B., Mohssen, M., Zablocki, O., Pelletier, E., et al. (2022) **Diversity and**
ecological footprint of Global Ocean RNA viruses *Science* **376**:1202-1208

Du Toit, A. (2017) **Viral infection: The language of phages** *Nat Rev Microbiol*
15:134-135

Dwivedi, B., Xue, B., Lundin, D., Edwards, R.A., and Breitbart, M. (2013) **A**
bioinformatic analysis of ribonucleotide reductase genes in phage genomes and
metagenomes *BMC Evol Biol* **13**:33

Enav, H., Mandel-Gutfreund, Y., and Béjà, O. (2014) **Comparative metagenomic**
analyses reveal viral-induced shifts of host metabolism towards nucleotide
biosynthesis *Microbiome* **2**:9

Engelhardt, T., Orsi, W.D., and Jørgensen, B.B. (2015) **Viral activities and life**
cycles in deep subseafloor sediments *Environ Microbiol Rep* **7**:868-873

Evseev, P., Bocharova, J., Shagin, D., and Chebotar, I. (2023) **Analysis of**
Pseudomonas aeruginosa **isolates from patients with cystic fibrosis revealed novel**
groups of filamentous bacteriophages *Viruses* **15**:2215

Falkowski, P.G., Barber, R.T., and Smetacek, V. (1998) **Biogeochemical controls**
and feedbacks on ocean primary production *Science* **281**:200-206

Faust, K., and Raes, J. (2012) **Microbial interactions: from networks to models** *Nat*
Rev Microbiol **10**:538-550

Foley, M.H., Cockburn, D.W., and Koropatkin, N.M. (2016) **The Sus operon: a model system for starch uptake by the human gut** *Bacteroidetes* *Cell Mol Life Sci* **73**:2603-2617

Fortunato, T.M., Beltrami, C., Emanueli, C., De Bank, P.A., and Pula, G. (2016) **Platelet lysate gel and endothelial progenitors stimulate microvascular network formation in vitro: tissue engineering implications** *Sci Rep* **6**:25326

Galperin, M.Y., Makarova, K.S., Wolf, Y.I., and Koonin, E.V. (2015) **Expanded microbial genome coverage and improved protein family annotation in the COG database** *Nucleic Acids Res* **43**:D261-269

Gazitúa, M.C., Vik, D.R., Roux, S., Gregory, A.C., Bolduc, B., Widner, B., Mulholland, M.R., Hallam, S.J., Ulloa, O., and Sullivan, M.B. (2021) **Potential virus-mediated nitrogen cycling in oxygen-depleted oceanic waters** *ISME J* **15**:981-998

Giljan, G., Brown, S., Lloyd, C.C., Ghobrial, S., Amann, R., and Arnosti, C. (2023) **Selfish bacteria are active throughout the water column of the ocean** *ISME Commun* **3**:11

Godkin, A., and Smith, K.A. (2017) **Chronic infections with viruses or parasites: breaking bad to make good** *Immunology* **150**:389-396

Graham, L., and Orenstein, J.M. (2007) **Processing tissue and cells for transmission electron microscopy in diagnostic pathology and research** *Nat Protoc* **2**:2439-2450

Gregory, A.C., Zayed, A.A., Conceição-Neto, N., Temperton, B., Bolduc, B., Alberti, A., Ardyna, M., Arkhipova, K., Carmichael, M., Cruaud, C., et al. (2019) **Marine DNA Viral Macro- and Microdiversity from Pole to Pole** *Cell* **177**:1109-1123.e1114

Hoffmann Berling, H., and Maze, R. (1964) **Release of male-specific bacteriophages from surviving host bacteria** *Virology* **22**:305-313

Howard-Varona, C., Hargreaves, K.R., Abedon, S.T., and Sullivan, M.B. (2017) **Lysogeny in nature: mechanisms, impact and ecology of temperate phages** *ISME J* **11**:1511-1520

Howard-Varona, C., Lindback, M.M., Bastien, G.E., Solonenko, N., Zayed, A.A., Jang, H., Andreopoulos, B., Brewer, H.M., del Rio, T.G., Adkins, J.N., et al. (2020) **Phage-specific metabolic reprogramming of virocells** *ISME J* **14**:881-895

Hurwitz, B.L., Hallam, S.J., and Sullivan, M.B. (2013) **Metabolic reprogramming by viruses in the sunlit and dark ocean** *Genome Biol* **14**:R123

Jiang, S.C., and Paul, J.H. (1996) **Occurrence of lysogenic bacteria in marine microbial communities as determined by prophage induction** *Mar Ecol Prog Ser* **142**:27-38

Kanehisa, M., Goto, S., Kawashima, S., Okuno, Y., and Hattori, M. (2004) **The KEGG resource for deciphering the genome** *Nucleic Acids Res* **32**:D277-280

Katoh, K., Rozewicki, J., and Yamada, K.D. (2019) **MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization** *Brief Bioinform* **20**:1160-1166

Kim, A.Y., and Blaschek, H.P. (1991) **Isolation and characterization of a filamentous viruslike particle from *Clostridium acetobutylicum* NCIB 6444** *J Bacteriol* **173**:530-535

Knezevic, P., Adriaenssens, E.M., and Ictv Report, C. (2021) **ICTV Virus Taxonomy Profile: Inoviridae** *J Gen Virol* **102**:

Koebnik, R. (2005) **TonB-dependent trans-envelope signalling: the exception or the rule?** *Trends Microbiol* **13**:343-347

Krupovic, M., and Consortium, I.R. (2018) **ICTV virus taxonomy profile: *Plasmaviridae*** *J Gen Virol* **99**:617-618

Langmead, B., and Salzberg, S.L. (2012) **Fast gapped-read alignment with Bowtie 2** *Nat Methods* **9**:357-359

Letunic, I., and Bork, P. (2021) **Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree display and annotation** *Nucleic Acids Res* **49**:W293-W296

Li, H. (2013) **Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM** *ArXiv* preprint arXiv:1303.3997

Li, H., and Durbin, R. (2009) **Fast and accurate short read alignment with Burrows-Wheeler transform** *Bioinformatics* **25**:1754-1760

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G., and Durbin, R. (2009) **The Sequence Alignment/Map format and SAMtools** *Bioinformatics* **25**:2078-2079

Li, Z., Pan, D., Wei, G., Pi, W., Zhang, C., Wang, J.H., Peng, Y., Zhang, L., Wang, Y., Hubert, C.R.J., et al. (2021) **Deep sea sediments associated with cold seeps are a subsurface reservoir of viral diversity** *ISME J* **15**:2366-2378

Lin, Y.R., Chiu, C.W., Chang, F.Y., and Lin, C.S. (2012) **Characterization of a new phage, termed phi A318, which is specific for *Vibrio alginolyticus*** *Arch Virol* **157**:917-926

Lindell, D., Jaffe, J.D., Coleman, M.L., Futschik, M.E., Axmann, I.M., Rector, T., Kettler, G., Sullivan, M.B., Steen, R., Hess, W.R., et al. (2007) **Genome-wide expression dynamics of a marine virus and host reveal features of co-evolution** *Nature* **449**:83-86

Liu, Y., Alexeeva, S., Bachmann, H., Guerra Martínez, J.A., Yeremenko, N., Abee, T., and Smid, E.J. (2022) **Chronic release of tailless phage particles from *Lactococcus lactis*** *Appl Environ Microbiol* **88**:e0148321

Lunin, V.V., Wang, H.-T., Bharadwaj, V.S., Alahuhta, M., Peña, M.J., Yang, J.-Y., Archer-Hartmann, S.A., Azadi, P., Himmel, M.E., Moremen, K.W., et al. (2020) **Molecular mechanism of polysaccharide acetylation by the arabidopsis xylan O-acetyltransferase XOAT1** *Plant Cell* **32**:2367-2382

Mäntynen, S., Laanto, E., Oksanen, H.M., Poranen, M.M., and Díaz-Muñoz, S.L. (2021) **Black box of phage-bacterium interactions: exploring alternative phage infection strategies** *Open Biol* **11**:210188

Mai-Prochnow, A., Hui, J.G., Kjelleberg, S., Rakonjac, J., McDougald, D., and Rice, S.A. (2015) **'Big things in small packages: the genetics of filamentous phage and effects on fitness of their host'** *FEMS Microbiol Rev* **39**:465-487

Mann, N.H., Cook, A., Millard, A., Bailey, S., and Clokie, M. (2003) **Marine ecosystems: bacterial photosynthesis genes in a virus** *Nature* **424**:741

Marvin, D.A., Symmons, M.F., and Straus, S.K. (2014) **Structure and assembly of filamentous bacteriophages** *Prog Biophys Mol Bio* **114**:80-122

Mazaheri Nezhad Fard, R., Barton, M.D., and Heuzenroeder, M.W. (2010) **Novel bacteriophages in *Enterococcus* spp** *Curr Microbiol* **60**:400-406

McKay, L.L., and Baldwin, K.A. (1973) **Induction of prophage in *Streptococcus lactis* C2 by ultraviolet irradiation** *Appl Microbiol* **25**:682-684

Panphut, W., Senapin, S., Sriurairatana, S., Withyachumnarnkul, B., and Flegel, T.W. (2011) **A novel integrase-containing element may interact with Laem-Singh virus (LSNV) to cause slow growth in giant tiger shrimp** *BMC Vet Res* **7**:18

Pratama, A.A., Bolduc, B., Zayed, A.A., Zhong, Z.P., Guo, J., Vik, D.R., Gazitúa, M.C., Wainaina, J.M., Roux, S., and Sullivan, M.B. (2021) **Expanding standards in viromics: in silico evaluation of dsDNA viral genome identification, classification, and auxiliary metabolic gene curation** *PeerJ* **9**:e11447

Putzrath, R.M., and Maniloff, J. (1977) **Growth of an enveloped mycoplasmavirus and establishment of a carrier state** *J Virol* **22**:308-314

Reintjes, G., Arnosti, C., Fuchs, B.M., and Amann, R. (2017) **An alternative polysaccharide uptake mechanism of marine bacteria** *ISME J* **11**:1640-1650

Roux, S., Krupovic, M., Daly, R.A., Borges, A.L., Nayfach, S., Schulz, F., Sharrar, A., Matheus Carnevali, P.B., Cheng, J.F., Ivanova, N.N., et al. (2019) **Cryptic inoviruses revealed as pervasive in bacteria and archaea across Earth's biomes** *Nat Microbiol* **4**:1895-1906

Roux, S., Brum, J.R., Dutilh, B.E., Sunagawa, S., Duhaime, M.B., Loy, A., Poulos, B.T., Solonenko, N., Lara, E., Poulain, J., et al. (2016) **Ecogenomics and potential biogeochemical impacts of globally abundant ocean viruses** *Nature* **537**:689-693

Russel, M. (1991) **Filamentous phage assembly** *Mol Microbiol* **5**:1607-1613

Samson, J.E., Magadán, A.H., Sabri, M., and Moineau, S. (2013) **Revenge of the phages: defeating bacterial defences** *Nat Rev Microbiol* **11**:675-687

Shaffer, M., Borton, M.A., McGivern, B.B., Zayed, A.A., La Rosa, S.L., Solden, L.M., Liu, P., Narrowe, A.B., Rodríguez-Ramos, J., Bolduc, B., et al. (2020) **DRAM for distilling microbial metabolism to automate the curation of microbiome function** *Nucleic Acids Res* **48**:8883-8900

Shkoporov, A.N., Turkington, C.J., and Hill, C. (2022) **Mutualistic interplay between bacteriophages and bacteria in the human gut** *Nat Rev Microbiol*

Sullivan, M.B., Coleman, M.L., Weigele, P., Rohwer, F., and Chisholm, S.W. (2005) **Three *Prochlorococcus* cyanophage genomes: signature features and ecological interpretations** *PLoS Biol* **3**:e144

Sullivan, M.B., Lindell, D., Lee, J.A., Thompson, L.R., Bielawski, J.P., and Chisholm, S.W. (2006) **Prevalence and evolution of core photosystem II genes in marine cyanobacterial viruses and their hosts** *PLoS Biol* **4**:e234

Suttle, C.A. (2005) **Viruses in the sea** *Nature* **437**:356-361

Suttle, C.A. (2007) **Marine viruses--major players in the global ecosystem** *Nat Rev Microbiol* **5**:801-812

Trifinopoulos, J., Nguyen, L.T., von Haeseler, A., and Minh, B.Q. (2016)
W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis
Nucleic Acids Res **44**:W232-W235

Tseng, Y.H., Lo, M.C., Lin, K.C., Pan, C.C., and Chang, R.Y. (1990)
Characterization of filamentous bacteriophage phi Lf from *Xanthomonas*
campestris **pv. campestris** *J Gen Virol* **71(Pt8)**:1881-1884

Wang, F., Wang, F., Li, Q., and Xiao, X. (2007) **A novel filamentous phage from**
the deep-sea bacterium *Shewanella piezotolerans* **WP3 is induced at low**
temperature *J Bacteriol* **189**:7151-7153

Weinbauer, M.G., Wilhelm, S.W., and Suttle, C.A. (1999) **Sunlight-lnduced DNA**
damage and resistance in natural viral communities *Aquat Microb Ecol*
17:111-120

Wigington, C.H., Sonderegger, D., Brussaard, C.P., Buchan, A., Finke, J.F., Fuhrman,
J.A., Lennon, J.T., Middelboe, M., Suttle, C.A., Stock, C., et al. (2016)
Re-examination of the relationship between marine virus and microbial cell
abundances *Nat Microbiol* **1**:15024

Yamamoto, K.R., Alberts, B.M., Benzinger, R., Lawhorne, L., and Treiber, G. (1970)
Rapid bacteriophage sedimentation in the presence of polyethylene glycol and its
application to large-scale virus purification *Virology* **40**:734-744

Yarza, P., Yilmaz, P., Pruesse, E., Glöckner, F.O., Ludwig, W., Schleifer, K.H.,
Whitman, W.B., Euzéby, J., Amann, R., and Rosselló-Móra, R. (2014) **Uniting the**
classification of cultured and uncultured bacteria and archaea using 16S rRNA
gene sequences *Nat Rev Microbiol* **12**:635-645

Zechmann, B., and Zellnig, G. (2009) **Rapid diagnosis of plant virus diseases by**
transmission electron microscopy *J Virol Methods* **162**:163-169

Zeng, J., Wang, Y., Zhang, J., Yang, S., and Zhang, W. (2021) **Multiple novel**
filamentous phages detected in the cloacal swab samples of birds using viral
metagenomics approach *Virol J* **18**:240

Zeng, Q., and Chisholm, S.W. (2012) **Marine viruses exploit their host's**
two-component regulatory system in response to resource limitation *Curr Biol*
22:124-128

Zhang, T., Zheng, R., Liu, R., Li, R., and Sun, C. (2022) **Cultivation and functional**
characterization of a deep-sea *Lentisphaerae* **representative reveals its unique**
physiology and ecology *Front Mar Sci* **9**:848136

Zheng, R., Cai, R., Liu, R., Liu, G., and Sun, C. (2021a) *Maribellus comscasis* **sp.**
nov., a novel deep-sea *Bacteroidetes* **bacterium, possessing a prominent capability**
of degrading cellulose *Environ Microbiol* **23**:4561-4575

Zheng, R., Liu, R., Shan, Y., Cai, R., Liu, G., and Sun, C. (2021b) **Characterization**
of the first cultured free-living representative of *Candidatus* **Izemoplasma**
uncovers its unique biology *ISME J* **15**:2676-2691

Zimmerman, A.E., Howard-Varona, C., Needham, D.M., John, S.G., Worden, A.Z.,
Sullivan, M.B., Waldbauer, J.R., and Coleman, M.L. (2020) **Metabolic and**
biogeochemical consequences of viral infection in aquatic ecosystems *Nat Rev*
Microbiol **18**:21-34

Figure Legends

Fig. 1. Polysaccharides promote the deep-sea *Lentisphaerae* strain WC36 growth and stimulate the expression of phage-associated genes. (A) Maximum likelihood phylogenetic tree of 16S rRNA gene sequences from strain WC36, strain zth2, and some *Planctomycetes-Verrucomicrobia-Chlamydia* (PVC) group bacteria. *Bacillus cereus* ATCC 14579 was used as the outgroup. Bootstrap values (%) > 80 are indicated at the base of each node with the gray dots (expressed as percentages of 1000 replications). The accession number is displayed behind corresponding strain. (B) Growth assays of strain WC36 cultivated in rich medium either without supplementation, with 5 g/L or 10 g/L laminarin or with 5 g/L or 10 g/L starch. (C) Transcriptomics-based heat map showing all up-regulated genes encoding phage-associated proteins in strain WC36 cultured in rich medium supplemented with 10 g/L laminarin. "Rich" indicates strain WC36 cultivated in rich medium; "Lam" indicates strain WC36 cultivated in rich medium supplemented with 10 g/L laminarin. (D) RT-qPCR detection of the expression of some genes encoding phage-associated proteins shown in panel C. The heat map was generated by Heml 1.0.3.3. The numbers in panel C represent multiple differences in gene expression (by taking $\log_2$ values).

Fig. 2. Polysaccharides induce the production of bacteriophages in *Lentisphaerae* strain WC36. (A) TEM observation of phages extracted from the supernatant of WC36 cells cultured in rich medium supplemented with or without polysaccharide. Panel I shows the absence of phages in the supernatant of WC36 cells cultivated in rich medium. Panels II-IV show the morphology of phages present in the supernatant of WC36 cells cultivated in rich medium supplemented with 10 g/L laminarin. Yellow hollow arrows indicate the typical granular structure of filamentous phages, and green arrows indicate phage-like particles with other shapes. Panels V-VIII show the morphology of phages present in the supernatant of strain WC36 cells cultivated in rich medium supplemented with 10 g/L starch. Typical filamentous phages are indicated with yellow hollow arrows, and two other kinds of phage-like particles with different shapes are indicated by orange and green arrows, respectively. (B) TEM

observation of strain WC36 cultured in rich medium supplemented with or without 10 g/L laminarin or 10 g/L starch. Panel I shows representative morphology of strain WC36 cultivated in rich medium. Panel II shows the morphology of strain WC36 cultivated in rich medium supplemented with 10 g/L laminarin. Panels III-IV show the morphology of strain WC36 cultivated in rich medium supplemented with 10 g/L starch. Red arrows indicate filamentous phages associated with bacterial cells. (C) TEM of an ultrathin section of strain WC36 cultured in rich medium supplemented with or without 10 g/L laminarin or 10 g/L starch. Panel I shows an ultrathin section of strain WC36 cultivated in rich medium; Panels II-III show ultrathin sections of strain WC36 cultivated in rich medium supplemented with 10 g/L laminarin; Panel IV shows an observation of the ultrathin section of strain WC36 cultivated in rich medium supplemented with 10 g/L starch. Red arrows indicate filamentous phages being released from or entering bacterial cells.

Fig. 3. Bacteriophages induced from *Lentisphaerae* strain WC36 by polysaccharides are released via chronic manners. (A) Growth curve of strain WC36 cultivated in either rich medium alone or rich medium supplemented with 5 g/L or 10 g/L laminarin for 30 days. The number of filamentous phages (B) and hexagonal phages (C) extracted from 1 μL of the supernatant of strain WC36 cell cultured in rich medium supplemented with or without 5 g/L or 10 g/L laminarin (for 5, 10, and 30 days). The average numbers are shown at the top of the bar charts.

Fig. 4. Polysaccharides promote the growth of deep-sea *Lentisphaerae* strain zth2 and induce the production of bacteriophages. (A) Growth assays of strain zth2 cultivated in rich medium supplemented with or without 3 g/L laminarin for four days. (B) Transcriptomics-based heat map showing all up-regulated genes encoding phage-associated proteins in strain zth2 cultured in rich medium supplemented with 3 g/L laminarin. (C) RT-qPCR detection of the expressions of some genes encoding phage-associated proteins shown in panel B. The numbers in panel C represent multiple differences in gene expression (by taking $\log_2$ values). (D) TEM observation of phages extracted from the supernatant of a cell suspension of strain zth2 cultured in the rich medium supplemented with or without polysaccharides. Panels I-III show the morphology of phages in the cell supernatant of strain zth2 cultivated in rich medium

supplemented with 3 g/L laminarin. Typical filamentous phage is indicated with
yellow hollow arrows, and two other kinds of phage-like particles with different
shapes are indicated by orange and green arrows, respectively. Panels IV-V show the
morphology of phages present in the cell supernatant of strain zth2 cultivated in rich
medium supplemented with 3 g/L starch. Typical filamentous phages are indicated
with red arrows, and other phage-like particles with different shapes are indicated by
green arrows. Panel VI shows that no phages were observed in the cell supernatant of
strain zth2 cultivated in rich medium. "Rich" indicates strain zth2 cultivated in rich
medium; "Lam" indicates strain zth2 cultivated in rich medium supplemented with 3
g/L laminarin.

Fig. 5. Genomic organization of bacteriophages released from *Lentisphaerae*
strains WC36 and zth2 cultured in rich medium supplemented with
polysaccharide. (A) Schematic of the genomic composition of Phage-WC36-1 and
Phage-zth2-1. (B) A diagram showing the genomic composition of Phage-WC36-2. (C)
The genomic composition of Phage-zth2-2. Arrows represent different ORFs and the
direction of transcription. The main putative gene products of the phages are shown,
and the numbers in brackets indicate the numbers of amino acids within each ORF.
Hypothetical proteins are indicated with gray arrows, structural modules are indicated
by green arrows, the replication module is indicated by blue-grey arrows, the
regulatory module is indicated by golden arrows, the assembly and secretion modules
are indicated by blue arrows and potential AMGs encoding proteins associated with
polysaccharide transport and degradation are indicated by red arrows. The size of the
phage genomes is shown behind each gene cluster.

A

Rich
Rich+3 g/L laminarin

B

(Log₂ fold change)

Locus tag	Product
3.99	JYG24_06675	Phage terminase large subunit
3.35	JYG24_11755	Phage tail protein
4.27	JYG24_11775	Phage major tail tube protein
3.48	JYG24_11780	Phage tail sheath protein
9.30	JYG24_11790	Phage tail protein
7.96	JYG24_11795	Phage tail protein I
3.67	JYG24_11825	Phage baseplate assembly protein V
3.68	JYG24_11845	Phage major capsid protein
5.81	JYG24_11850	Phage portal protein
6.77	JYG24_11930	Phage antirepressor

Lam vs Rich

C

Rich+Lam

Rich+Sta

Rich