Paternal methamphetamine exposure induces higher sensitivity to methamphetamine in male offspring through driving ADRB1 on CaMKII-positive neurons in mPFC

Yanyan Zheng[a,1], Dekang Liu[a,1], Zhaosu Li[a], Hao Guo[a], Wenwen Chen[a], Tao Hu[a], Yuanyuan Zhang[a], Xiang Li[a], Ziheng Zhao[a], Qinglong Cai[a], Feifei Ge[a], Yu Fan[a,*], Xiaowei Guan[a,*]

[a]Department of Human Anatomy and Histoembryology, Nanjing University of Chinese Medicine, Nanjing 210023, China

[1]These authors are equally contribute to this work

Running title: Paternal METH and offspring ADRB1

*Correspondence to: Prof. X Guan (guanxw918@njucm.edu.cn), Dr. Y Fan (yfan@njucm.edu.cn)

Phone number: +86-25-85811387

Highlights

- Paternal methamphetamine (METH) exposure induces higher sensitivity to METH in male F1 during adulthood, accompanied with higher ADRB1 level in mPFC.

- METH use in F1 ($METH^{F1}$) evokes more CaMKII-positive neurons in mPFC of METH-sired than saline-sired male F1.

- Inhibiting ADRB1 function or knocking-down ADRB1 level on CaMKII-positive neurons ($ADRB1^{CaMKII}$) efficiently suppresses $METH^{F1}$-evoked mPFC activation, and ultimately rescues transgenerational susceptibility to addiction in male F1.

Abstract

Paternal abuse of drugs, such as methamphetamine (METH), elevates the risk of developing addiction in subsequent generations, however, its underlying molecular mechanism remains poorly understood. Male adult mice (F0) were exposed to METH for 30 days, followed by mating with naïve female mice to create the first-generation mice (F1). When growing to adulthood, F1 were subjected to conditioned place preference (CPP) test. Subthreshold dose of METH, insufficient to induce CPP normally, were used in F1 (METH^{F1}). Selective antagonist (betaxolol) for β1-adrenergic receptor (ADRB1) or its knocking-down virus were administrated into mPFC to regulate ADRB1 function and expression on CaMKII-positive neurons. METH-sired male F1 acquired METH^{F1}-induced CPP, indicating that paternal METH exposure induce higher sensitivity to METH in male F1. Compared with saline (SAL)-sired male F1, CaMKII-positive neuronal activity was normal without METH^{F1}, but strongly evoked after METH^{F1} treatment in METH-sired male F1 during adulthood. METH-sired male F1 had higher ADRB1 levels without METH^{F1}, which was kept at higher levels after METH^{F1} treatment in mPFC. Either inhibiting ADRB1 function with betaxolol, or knocking-down ADRB1 level on CaMKII-positive neurons (ADRB1^{CaMKII}) with virus transfection efficiently suppressed METH^{F1}-evoked mPFC acyivation, and ultimately blocked METH^{F1}-induced CPP in METH-sired male F1. In the process, the p-ERK1/2 and ΔFosB may be potential subsequent signals of mPFC ADRB1^{CaMKII}. The mPFC ADRB1^{CaMKII} mediates paternal METH exposure-induced higher sensitivity to drug addiction in male offspring, raising a promising pharmacological target for predicting or treating transgenerational addiction.

Keywords: paternal METH exposure, male offspring, higher sensitivity to METH, mPFC activation, ADRB1^{CaMKII}

Abbreviations

ADRB1, β1-adrenergic receptor; CPP, conditioned place preference; CREB, cAMP response element-binding protein; F1, first-generation; GPCRs, G protein coupled receptors; METH, methamphetamine; mPFC, medial prefrontal cortex; SAL, saline; PD, postnatal day; PFC, prefrontal cortex

1 Introduction

Paternal psychostimulants intake elevates the risk of developing addiction in subsequent generations [1, 2], which remains major health concern. Methamphetamine (METH) is a commonly abused psychostimulant worldwide. One recent study found that METH-sired rats exhibit more sensitized behaviors in response to acute drug use [3], indicating that paternal METH use may enhance susceptibility to drugs in offspring. The medial prefrontal cortex (mPFC) is critically implicated in the processes of METH addiction [4, 5] and susceptibility to drugs [6-8]. Human imaging studies show that prenatal METH exposure significantly affect the volume, cortical thickness and connection of offspring brains, especially the prefrontal cortex (PFC) [7, 9, 10]. Previously, we found that mPFC were blunted to stimulants in METH-sired mice during adulthood [11]. However, the intrinsic molecular mechanism of mPFC underlying transgenerational effects of paternal METH abuse on offspring remains much to be clarified.

Adrenergic innervation on mPFC glutamatergic (CaMKII-positive) neurons plays essential roles in neurophysiology and neuropathology. The β1-adrenergic receptor (ADRB1) is highly expressed on mPFC CaMKII-positive neurons (ADRB1^{CaMKII}), which has been reported to participate in cognition [12, 13], reward [14] , arousal [15], motivation [16], and stress [17]. Generally, activating ADRB1 triggers neuronal excitability within PFC [18]. In animal models of cocaine-induced conditioned place preference (CPP), a model to evaluate reward and cue-related stimulations of drugs, ADRB1 administration efficiently blocks the reinstatement behaviors to cocaine in mice [19]. Prenatal events exposure, malnutrition for example,

impairs the long-term potentiation in PFC of offspring, which leads to memory

deficits [20-22], but could be fully recovered by ADRB1 activation [23]. Our

previous transcriptomic analysis screened out a higher levels of *Adrb1* gene in

mPFC of METH-sired male F1 mice (not published). These findings suggest

that ADRB1 might involve in drug-related transgenerational neurotoxicity, such

as mPFC plasticity and vulnerability of addiction.

The β-adrenergic innervation could regulate subsequent signals though G

protein-dependent or/and β-arrestin-dependent pathway. As one of classical G

protein coupled receptors (GPCRs), ADRB1 has been well-studied to be

coupled to Gs protein which lead to an increase in protein kinase A

(PKA)-dependent signaling pathway including cAMP, PKA, and cAMP

response element-binding protein (CREB) [12, 24]. Recent reports have

revealed that several ligands called "biased ligands", such as β-adrenergic

receptors, elicit G protein-independent and β-arrestin-dependent pathway [25,

26], which may be served as negative feedback of the G protein-dependent

pathway [27]. Huang et al. [28] find that β-Arrestin-biased β-adrenergic

signaling promotes extinction learning of cocaine reward memory. At present,

the price signaling pathway of ADRB1 in drug-preferred behavior remains

much to be clarified.

In the present study, male adult mice (F0) were exposed to METH for 30

days, followed by mating with naïve female mice to create the first-generation

mice (F1). When growing up to adult age, F1 mice were subjected to

conditioned place preference (CPP) test to evaluate sensitivity to METH.

Subthreshold dose of METH, that is insufficient to induce CPP normally, were

used in F1 mice (METH^{F1}). The role of ADRB1^{CaMKII} in transgenerational

susceptibility of addiction were explored in METH-sired F1 mice.

2 Materials and methods

2.1 Animals

Male C57BL/6 wild type (WT) mice weighing 22-25 g were used. All animals were housed at constant humidity (40~60%) and temperature (24 ± 2ºC) with a 12-hour light/dark cycle and allowed free access to food and water. The animals were handled for three days prior to onset of experiments. All procedures were carried out in accordance with the National Institutes of Health Guide for the Care and Use of Laboratory Animals of China and approved by the Institutional Animal Care and Use Committee (IACUC) at Nanjing University of Chinese Medicine.

2.2 Drug treatment and tissue collection

Male C57BL/6 WT mice were assigned to administration of METH (5 mg/kg, dissolved in saline, i.p.) or saline (0.2 mL, i.p.) once daily at 10 a.m. for 30 consecutive days. At 24 h after the last injection, the F0 mice were mated with three naïve female C57BL/6 WT mice for 4 days to create METH-sired or saline-sired first-generation (F1) offspring mice.

On postnatal day 21 (PD21), male and female F1 mice were weaned and kept (4 male mice per cage) in homecages. On PD60, the mPFC tissue were collected from some male F1 mice. From PD70, some male and female F1 mice were subjected to METH-induced CPP training and test. In this study, to examine the sensitivity to METH, subthreshold dose of METH (0.25 mg/kg) were used to induce CPP in male F1 mice. Betaxolol, a specific β1-adrenergic receptor (ADRB1) antagonist (Cat# S2091, Sellekchem, USA) was diluted to

0.6 µg/µl in sterile saline. Brains of male F1 mice were collected on PD60 and

PD73. The brains of PD60 mice were collected without any treatment to

represent baseline condition. The brains of PD73 mice were collected within

60 min after CPP test.

Some male naïve mice were only subjected to saline or METH treatment.

On Day 1 (D1) and D2, mice were injected with saline (0.2 mL, i.p.) in the

morning, and saline (0.2 mL, i.p.) or METH (0.25 mg/kg, i.p.) in the afternoon

once daily at homecage. Their mPFC tissue were collected on D3 within 60

min of the last injection.

2.3 CPP

In the present study, the 0.25 mg / kg dose of METH administration (METH^{F1}),

being insufficient to produce CPP in naïve mice (subthreshold dose), was used

to induce CPP in male F1 mice and female F1 mice. The CPP test was

performed in the TopScan3D CPP apparatus (CleverSys, VA, USA), which is

constructed of two distinct chambers (15 × 15 × 23 cm each) separated by a

removable guillotine door. The CPP procedure consisted of three phases

including the pre-conditioning test (Pre-test, PD70), conditioning (CPP training,

PD71-72) and post-conditioning test (CPP test, PD73). During CPP training,

both METH-sired and SAL-sired F1 mice were injected with saline (0.2 mL,

i.p.) in the morning or METH (0.25 mg/kg, i.p.) in the afternoon once daily.

After each injection, the mice were confined to one conditioning chamber

(SAL-paired chamber or METH-paired chamber) for 45 min, and then returned

to their homecages. During the Pre-test and Test, mice freely access two

chambers for 15 min. The CPP score is the time spent in METH-paired

chamber minus that in SAL-paired chamber, and the ΔCPP score is the CPP

score of Test minus the CPP score of Pre-test.

2.4 Immunofluorescence

Briefly, the brains were perfused with 4% paraformaldehyde (PFA), and

coronal brain sections (30 μm) were cut on a cryostat (Leica, Germany). The

sections were incubated in 0.3% (v/v) Triton X-100 for 0.5 h, blocked with 5%

donkey serum for 1.5 h at room temperature, and incubated overnight at 4°C

with the following primary antibodies: rabbit anti-c-Fos (1:3000,

RRID:AB_2247211, Cell Signaling Technology, USA), mouse anti-NeuN (1:800,

RRID:AB_2313673, Millipore, USA), mouse anti-CaMKII α (1:100,

RRID:AB_626789, Santa Cruz Biotechnology, USA), mouse anti-CaMKII α

(1:50, RRID:AB_2721906,Cell Signaling Technology , USA), Rabbit

anti-ADRB1 (1:100, RRID:AB_10885544, Bioss, China), followed by the

corresponding fluorophore-conjugated secondary antibodies for 1.5 h at room

temperature. The following secondary antibodies were used here, including

Alexa Fluor 555-labeled donkey anti-rabbit secondary antibody (1:500, RRID:

AB_2762834, Thermo Fisher Scientific, USA), Alexa Fluor 488-labeled donkey

anti-mouse secondary antibody (1:500, RRID: AB_141607, Thermo Fisher

Scientific, USA). In detail of Figure 1k-l, brain slices were attached to slides

and placed in an oven at 60°C for 40 min, and then immersed in 4% PFA

solution for 15 min in a refrigerator at 4°C for fixation. Then graded

concentration alcohol dehydration (50%-70%-100%-100%) was performed

sequentially for 5 min each time. Next, antigen repair was performed by

immersing the slides with brain slices attached in 10 mM PH=6.0 trisodium

citrate solution and heating in a water bath at 82°C for 30 min. Normal staining

steps were then followed. Fluorescence signals were visualized using a Leica

DM6B upright digital research microscope (Leica, Germany) and Leica TCS

SP8 confocal microscope (Leica, Germany).

2.5 Western blot

Total protein was extracted from mPFC of individual mouse using RIPA lysis

buffer (Beijing ComWin Biotech Co., China), Phosphatase Inhibitor (Beijing

ComWin Biotech Co., China), and Protease Inhibitor (Beijing ComWin Biotech

Co., China) according to the manufacturer's instructions. Protein samples (15

or 20 μg) was separated by 10% SDS–PAGE, 12% SDS–PAGE and

electrophoretically transferred onto PVDF membranes. The transferred

membranes were blocked with 5% non-fat dry milk and 0.1% Tween 20 in

TBST buffer for 2 h at room temperature, then subsequently incubated with the

following primary antibodies: Rabbit anti-c-Fos (1:1000, RRID:AB_224721,

Cell Signaling Technology, USA), Rabbit anti-GAPDH (1:10000,

RRID:AB_2651132, Bioworld Technology, USA), Rabbit anti-CaMKII α (1:1000,

RRID:AB_305050, Abcam, USA), Rabbit anti-ADRB1 (1:1000,

RRID:AB_10885544, Bioss, China), Rabbit anti-ARRB2 (1:1000, BA3767,

Boster, China), Rabbit anti-ERK1/2 (1:1000, BM4326, Boster, China), Rabbit

anti-phospho-ERK1/2 (1:1000, BM4156, Boster, China), Rabbit anti-ΔFosB

(1:1000, bsm-52071R, Bioss, China) , Rabbit anti-PKA (1:1000, bs-0520R,

Bioss, China), Rabbit anti-CREB (1:1000, RRID:AB_2800317, Cell Signaling

Technology, USA), Rabbit anti-phospho-CREB (1:1000, RRID:AB_2561044,

Cell Signaling Technology, USA). The next day, the membranes were washed

in Tris-buffered saline with Tween 20 (TBST) and incubated with horseradish

peroxidase (HRP)-conjugated secondary antibody goat anti-rabbit or goat anti

mouse (1:5000, Beijing ComWin Biotech Co., China) at room temperature for 1

h. The blots were visualized by the eECL Western Blot kit (Beijing ComWin

Biotech Co., China, CW0049S) or ELC Kit (Vazyme^{TM}) and the signal was

visualized by imaging system (Tanon-5200, Shanghai, China). The blots were

washed with stripping buffer (Beyotime Institute of Biotechnology, China) to be

reprobed with other antibodies. In this study, GAPDH was used as the loading

control. The relative level of each protein expression was normalized to

GAPDH. Values for target protein levels were calculated using Image J

software (NIH, USA).

2.6 RNA extraction and quantitative real-time PCR

The mPFC were dissected immediately on ice. Total RNA was extracted from

the mPFC using FastPure Cell/Tissue Total RNA Isolation Kit (Vazyme, China),

according to the manufacturer's instructions. All samples were tested for

concentration and purified using a 4200 TapeStation (Agilent Technologies,

Santa Clara, USA). After RNA extraction, the amount of RNA was normalized

across samples, and cDNA was created using HiScript II Q RT SuperMix for

qPCR (+gDNA wiper) (Vazyme, China). *Gapdh* was used as the internal

control. The primers used in the study were as follows: *Adrb1* (forward,

CTCATCGTGGTGGGTAACGTG; reverse, ACACACAGCACATCTACCGAA).

c-Fos   (forward,   CGGGTTTCAACGCCGACTA;   reverse,

TTGGCACTAGAGACGGACAGA).   *Gapdh*   (forward,

AGGTCGGTGTGAACGGATTTG;   reverse,

TGTAGACCATGTAGTTGAGGTCA). The *Adrb1* and *c-Fos* mRNA levels were

normalized to *Gapdh* mRNA levels. The relative mRNA level was calculated by

the comparative CT method ($2^{-\Delta\Delta Ct}$).

2.7 Local administration of betaxolol in mPFC

The cannula-double (O.D. 0.41 mm - 27G, C.C 0.8 mm, Mates with M 3.5, Cat 62063, RWD, China) were planted into mPFC (AP + 2.7 mm, ML ± 0.4 mm, DV − 1.5 mm) of male F1 mice on PD63 before CPP training. After surgery, mice were maintained at home cage about 1 week. From PD70, male F1 mice were subjected to METH-induced CPP training. In Experiment 1 and S1, a volume of 500 nL of betaxolol (specific antagonist for ADRB1) were bilaterally injected into mPFC at a rate of 100 nL/min at 15 min prior to CPP test on PD73. In Experiment 2 and S2, a volume of 500 nL of betaxolol bilaterally into the mPFC at a rate of 100 nL/min at 15 min prior to CPP training on PD71 and PD72.

2.8 Specific knockdown of ADRB1 on mPFC CaMKII-positive neurons

On PD55, mice were anesthetized with 2% isoflurane in oxygen, and were fixed in a stereotactic frame (RWD, Shenzhen, China). A heating pad was used to maintain the core body temperature of the mice at 36°C. The coordinates of mPFC were defined as AP + 2.05 mm, ML ± 0.3 mm, DV − 1.9 mm. A volume of 400 nL virus of *CaMKIIap-EGFP-MIR155 (MCS)-SV40PolyA* (KD group) (Titer, 2.41F + 13 v.g/mL; Cat, GIDV0286541, GeneChem, Shanghai, China) or *CaMKIIap-EGFP-SV40PolyA* (Ctrl group) (Titer, 4.01E + 12 v.g/mL; Cat, AAV9/CON540, GeneChem, Shanghai, China) were injected bilaterally into mPFC at a rate of 80 nL/min. After surgery, mice were kept at homecage about 3 weeks.

2.9 Dendritic spine analysis

The dendritic spines of mPFC were photographed using the z-stack scanning function of Leica TCS SP8 confocal microscope (Leica, Germany). The Spine

quantification were performed under LAS X software (Leica, Germany). The

number of spines is counting only in the second or the third branches of

dendrites from the body of CaMKII-positive neurons. Density of spines were

scored in about 50 µm dendritic segments in length.

2.10 Statistical analysis

Statistical analysis was carried out using GraphPad Prism 8.0.2 software. All

data are presented as the Mean ± SD. CPP score are analyzed by two-way

analysis of variance (ANOVA) with Sidak's multiple comparisons test, and

other data are analyzed by unpaired t-tests. Statistical significance was set as

$P < 0.05$.

3 Results

3.1 METH-sired male F1 mice acquires METH^{F1}-induced CPP in adulthood

In the present study, the 0.25 mg / kg dose of METH administration (METH^{F1}),

being insufficient to produce CPP in naïve mice (subthreshold dose), was used

to induce CPP in male F1 mice (Figure 1a) and female F1 mice (Figure S1a).

In male F1 mice (Figure 1b-d), METH^{F1} failed to induce CPP in SAL-sired

male F1 (n = 8 mice, t = 0.9474, P = 0.5872, Figure 1b), but increased CPP

score of METH-sired male F1 (n = 10 mice, t = 6.587, P < 0.0001, Figure 1b)

when compared with the corresponding Pre-test data. The ΔCPP score (t =

3.685, P = 0.002, Figure 1c) in METH-sired male F1 was higher than that in

SAL-sired male F1.

In female F1 mice (Figure S1b-d), METH^{F1} failed to induce CPP in both

SAL-sired female F1 (n = 8 mice, t = 1.398, P = 0.3295, Figure S1b) and

METH-sired female F1 (n = 10 mice, t = 1.979, P = 0.1264, Figure S1b) when

compared with the corresponding Pre-test data. There was no significance on

ΔCPP score (t = 0.277, P = 0.7854, Figure S1c) between METH-sired and

SAL-sired female F1.

These results indicate that paternal METH exposure induce higher

sensitivity to METH in male, but not female offspring. As such, only male F1

mice were used to the subsequent experiments.

3.2 METH^{F1} activates more CaMKII-positive neurons and induces higher

ADRB1 levels in mPFC of METH-sired male F1 mice

The mPFC tissue were collected from male F1 mice on PD60 and PD73, which

represent the baseline condition and responsive condition to METH^{F1},

respectively. The c-Fos, CaMKII and NeuN are used as markers to label

neuronal activation, glutamatergic neuron and total neuron in mPFC,

respectively.

On PD60, there was no significance in the percentage of c-Fos-positive &

CaMKII-positive neurons in subregions of mPFC including cingulate cortex

(Cg1, t = 0.595, P = 0.5839), prelimbic cortex (PrL, t = 0.0371, P = 0.9722) and

infralimbic cortex (IL, t = 0.4451, P = 0.6793) between METH-sired male F1 (n

= 3 mice) and SAL-sired male F1 (n = 3 mice, Figure 1e). While on PD73

following METH^{F1}, the percentage of c-Fos-positive & CaMKII-positive neurons

were at higher levels in Cg1 (t = 3.447, P = 0.0137), PrL (t = 3.539, P = 0.0076)

and IL (t = 3.160, P = 0.0134) of METH-sired male F1 (n = 4-5 mice) than that

of SAL-sired male F1 (n = 4-5 mice, Figure 1f). On PD73, the neurons (labeled

as NeuN-positive neurons) were more activated in Cg1 (t = 3.031, P = 0.0163),

PrL (t = 4.227, P = 0.0022) and IL (t = 4.098, P = 0.0027, Figure S2a), and the

dendritic spine density of CaMKII-positive neurons was increased (t = 4.960, P

= 0.0026, Figure S2b) in mPFC of METH-sired male F1 (n = 4-6 mice) than of

SAL-sired male F1 (n = 4-6 mice). As one of important kinase, CaMKII can

also be used to reflect the activities of neurons. On PD60, the baseline levels

of mPFC c-Fos (t = 3.496, P = 0.0058) and CaMKII protein (t = 2.457, P =

0.0338) were obviously lower in METH-sired male F1 (n = 6 mice) than that in

SAL-sired male F1 (n = 6 mice, Figure 1g). On PD73 following METH^{F1}, there

was similar levels of c-Fos protein (t = 0.5061, P = 0.6238) but a higher levels

of CaMKII protein (t = 3.350, P = 0.0074) in METH-sired male F1 (n = 6 mice)

than in SAL-sired male F1 (n = 6 mice, Figure 1h). These results indicate that

paternal METH exposure did not change or even reduce the mPFC activity, but

obviously   increased   mPFC   response   to   METH^{F1},   especially   the

CaMKII-positive neurons in it.

On PD60, the baseline levels of mPFC *Adrb1* gene (t = 3.084, P = 0.0368,

Figure 1i) and ADRB1 protein (t = 2.400, P = 0.0373, Figure 1j) were higher in

METH-sired male F1 (qPCR: n = 3 mice; protein: n = 6 mice) than that in

SAL-sired male F1 (qPCR: n = 3 mice; protein: n = 6 mice). On PD73 following

METH^{F1}, mPFC *Adrb1* gene (t = 4.621, P = 0.0099, Figure 1i) and ADRB1

protein (t = 3.489, P = 0.0175, Figure 1j) were persistently at higher levels in

METH-sired male mice (qPCR: n = 3 mice; protein: n = 3 mice) than that in

SAL-sired male F1 (qPCR: n = 3 mice; protein: n = 4 mice).

Further, ADRB1 levels on neurons of mPFC and its subregions (Figure 1k-l)

were examined by immunostaining. On PD60, the mean gray value of ADRB1

on neurons was much higher in the Cg1 (t = 4.649, P = 0.0097) and PrL (t =

4.311, P = 0.0125), and a high trend in IL (t = 2.265, P = 0.0862) of METH-sired

male F1 (n = 3 mice) than that in SAL-sired male F1 (n = 3 mice). On PD73

following METH^{F1}, the mean gray value of ADRB1 on neurons was

continuously higher in PrL (t = 6.304, P = 0.0032), and a higher trend in Cg1 (t

= 2.382, P = 0.0758), IL (t = 2.635, P = 0.0579) of METH-sired male F1 (n = 3

mice), when compared to that in SAL-sired male F1 (n = 3 mice).

To observe the effects of METH^{F1} treatment on levels of c-Fos and ADRB1

on the mPFC under normal conditions, male naïve mice were subjected to

subthreshold dose of METH treatment (Figure S3a). Neither *c-Fos* gene (t =

0.4522, P = 0.6745, Figure S3b) and *Adrb1* (t = 0.9778, P = 0.3835, Figure

S3b), nor c-Fos protein (t = 1.115, P = 0.3274, Figure S3c) and ADRB1 protein

(t = 1.723, P = 0.1600, Figure S3c) in mPFC were changed by subthreshold

dose of METH administration in naïve mice (n = 3 mice per group).

With these results, we infer that paternal METH exposure-increased ADRB1

on mPFC neurons might contribute to the higher sensitivity to METH in male

F1 mice.

3.3 Blocking ADRB1 suppresses METH^{F1}-induced CPP in METH-sired

male F1 mice

Here, betaxolol was bilaterally injected into mPFC (Figure S4a) to block the

activation of ADRB1. In Experiment 1 (Figure 2a), a single dose of betaxolol

were injected into mPFC at 15 min prior to METH^{F1}-induced CPP test.

METH-sired male F1 did not exhibit METH-preferred behavior any more (n = 8

mice, t = 1.666, P = 0.2219), and METH-sired male F1 with vehicle treatment

exhibited METH^{F1}-induced CPP continuously (n = 8 mice, t = 3.518, P =

0.0068). The ΔCPP score (t = 3.666, P = 0.0025) in vehicle-injected

METH-sired male F1 was much higher than that in betaxolol-injected

METH-sired male F1 (Figure 2b-d).

In Experiment 2 (Figure 2e), betaxolol were injected into mPFC during the

whole period of CPP training. Betaxolol-treated METH-sired male F1 mice did

not acquire METH^{F1}-induced CPP (n = 8 mice, t = 0.2852, P = 0.9514), but

vehicle-treated METH-sired male F1 mice still exhibit METH^{F1}-induced CPP (n

= 8 mice, t = 6.570, P < 0.0001). The ΔCPP score (t = 4.847, P = 0.0003) in

vehicle-injected METH-sired male F1 was much higher than that in

betaxolol-injected METH-sired male F1 (Figure 2f-h).

Here, we also examine the effects of betaxolol on METH^{F1}-induced CPP in

SAL-sired male F1 mice. In Experiment S1 (Figure S4b-e), similar to the above

results, METH^{F1} could not induce CPP in SAL-sired male F1 (n = 5 mice, t =

0.1464, P = 0.9872). Further, a single dose of betaxolol administration at 15

min prior to CPP did not affect their CPP score as well (n = 6 mice, t = 0.1320,

P = 0.9896). There was no difference in the ΔCPP score (t = 0.01917, P =

0.9851) between vehicle-treated and betaxolol-treated SAL-sired male F1. In

Experiment S2 (Figure S4f-i), similarly, METH^{F1} treatment could not induce

CPP in SAL-sired male F1 (n = 6 mice, t = 0.5901, P = 0.8136). The betaxolol

injection during CPP training did not significantly affect their CPP scores as

well (n = 6 mice, t = 0.4447, P = 0.8884). There was no difference in the ΔCPP

score (t = 0.4406, P = 0.6689) between vehicle-treated and betaxolol-treated

SAL-sired male F1.

These results indicate that mPFC ADRB1 mediate both formation (CPP

training) and expression (CPP test) of METH^{F1}-induced CPP in METH-sired F1

mice, and contribute to paternal METH exposure-increased sensitivity to

METH in male F1 mice.

3.4 Knocking-down ADRB1^{CaMKII} blocks METH-induced CPP in

METH-sired male F1 mice

Here, adeno-associated virus (AAV) were used to knocking-down ADRB1 on

CaMKII-positive neurons (ADRB1^{CaMKII}) within mPFC (Figure 3a). As shown in

Figure 3b, AAV of *CaMKIIap-EGFP-MIR155(MCS)-SV40PolyA* (KD group) or

AAV of *CaMKIIap-EGFP-SV40PolyA* (Ctrl group) were bilaterally infused into

mPFC on P55, which could specifically transfect CaMKII-positive neurons. On

P70, about 60%~80% CaMKII-positive neurons were transfected with virus

(Figure 3c). The levels of *Adrb1* gene ($t = 7.361$, $P = 0.0018$, Figure 3d),

ADRB1 protein ($t = 4.286$, $P = 0.0128$, Figure 3e), and ADRB1^{CaMKII} expression

($t = 4.237$, $P = 0.0133$, Figure 3f) were all reduced in mPFC of male F1 by KD

virus (n = 3 mice), when compared with Ctrl virus (n = 3 mice), indicating good

efficiency of KD virus.

As shown Figure 3g-i, Ctrl virus-treated METH-sired male F1 mice still

acquired METH^{F1}-induced CPP in (n = 12 mice, $t = 8.228$, $P < 0.0001$), but KD

virus-treated METH-sired male F1 mice did not exhibit METH-preferred

behaviors and even produced METH-aversion behaviors (n = 10 mice, $t =$

3.253, $P = 0.008$), when compared with their corresponding Pre-test score.

The ΔCPP score of KD group was much lower than that in Ctrl group ($t = 7.95$,

$P < 0.0001$).

These results confirmed that mPFC ADRB1^{CaMKII} mediate paternal METH

exposure-induced higher sensitivity to METH in male F1 mice.

3.5 Knocking-down ADRB1^{CaMKII} reduces mPFC activation in METH-sired

male F1 mice

Compared to Ctrl group (protein: n = 3 mice; immunostaining: n = 6 mice;

dendritic spine density: n = 6 mice), knocking-down mPFC ADRB1^{CaMKII}

obviously decreased levels of c-Fos protein (t = 5.372, P = 0.0058, Figure 4a)

and immunostaining (t = 3.136, P = 0.0106, Figure 4b), and dendritic spine

density of CaMKII-positive neurons (t = 3.891, P = 0.003, Figure 4c) in mPFC

of METH-sired male F1 (protein: n = 3 mice; immunostaining: n = 6 mice;

dendritic spine density: n = 6 mice).

The potential subsequent molecules of β-adrenergic signaling pathways

especially β-arrestin2 and PKA, were examined in mPFC. Compared to Ctrl

virus group (n = 3-6 mice), protein levels of β-arrestin2 (t = 0.4597, P = 0.6555)

and PKA (t = 0.6055, P = 0.5775, Figure 4d) were not changed, the p-ERK1/2

(t = 8.202, P < 0.0001), ratio of p-ERK1/2 to ERK1/2 (t = 2.486, P = 0.0347)

and ΔFosB (t = 2.944, P = 0.0147) were decreased, and ERK1/2 (t = 0.9153, P

= 0.3839), CREB (t = 0.5454, P = 0.6145), p-CREB (t = 0.8085, P = 0.4641)

and ratio of p-CREB to CREB (t = 0.6042, P = 0.5783) were not changed in

mPFC by knocking-down ADRB1 (n = 3-6 mice, Figure 4e).

These results indicate that ADRB1^{CaMKII} positively activate mPFC

CaMKII-positive neurons, p-ERK1/2 and ΔFosB are the subsequent signals for

ADRB1 in the process.

4 Discussion

The current study found that paternal METH exposure induces higher ADRB1

levels on mPFC neurons. The subthreshold dose of METH (METH^{F1}), which is

insufficient to produce CPP normally, efficiently induced METH-related CPP in

METH-sired male F1, accompanied with more activated CaMKII-positive

neurons and continuous higher ADRB1 in their mPFC. Either inhibiting ADRB1

function or knocking-down ADRB1 expression on CaMKII-positive neurons

(ADRB1^{CaMKII}) efficiently reduced METH^{F1}-evoked mPFC activation and

blocked METH[F1]-induced CPP in METH-sired male F1. These results indicate

that paternal METH exposure induce a higher sensitivity to METH in male

offspring potentially through driving mPFC ADRB1[CaMKII] (Figure 5).

METH is commonly abused psychostimulant among men of reproductive

age. Consistent with previous studies, we found that METH[F1], which is

insufficient to induce CPP in naïve mice, effectively induced METH-preferred

behaviors in METH-sired male F1 and not in SAL-sired male F1, indicating an

enhanced sensitivities to drugs in F1 of sires exposed to METH. The mPFC

has been implicated in the vulnerability to drug abuse [6-8]. Currently, little

literature uncover the association of paternal METH intake with offspring

mPFC alteration. *In utero* exposure to cocaine had been reported to increase

the induction of long-term potentiation (LTP) and neuronal excitability at

pyramidal neurons in mPFC of offspring [29], an increase in dendritic spines of

II/III layer of pyramidal neurons in mPFC of offspring [30] , and a reduction of

the sensitivity of locomotor activity to cocaine in offspring rats [29, 31]. By

contrast, Yan et al. [32] reported prenatal cocaine exposure attenuated

baseline levels of mature BDNF protein in offspring brain, and Cantacorps et al.

[33] found prenatal alcohol exposure increased vulnerability to cocaine

addiction in adult mice via a greater glutamate receptor in mPFC. Similarly,

human imaging studies demonstrated that prenatal cocaine

exposed-adolescents exhibited less capability of increasing their PFC

activation in response to memory load [34], and infants with prenatal cocaine

use had a lesser gray matter in prefrontal brain regions [35, 36]. In addition,

adolescents with prenatal cocaine exposure had reduced activity in PFC when

exposing to favorite-food cues [37]. Although there are conflicting results

among these studies, they suggest that plasticity of frontal brain play essential roles in prenatal drugs intaking-elicited heritable susceptibility to addiction in offspring. Here, we found that paternal METH exposure seemed to reduce mPFC activity in male F1, as indicated by lower levels of c-Fos and CaMKII in mPFC. However, METH[F1] treatment, which could not activate mPFC in naïve mice, reversely triggered mPFC activation in METH-sired male F1, implying potential intrinsic molecules in mPFC that may contribute to paternal METH exposure-increased sensitivity to METH in male F1 mice.

ADRB1 is highly expressed on glutamatergic neurons of mPFC. In general, endogenous activation of ADRB1 excites neurons via eliciting the depolarization on the membrane potential in mPFC [38]. An early study found that prenatal cocaine exposure increased cortical ADRB1 about 68% in pup rats at 30 days of age, a time when these rats showed hyperactivity [39]. Unfortunately, few studies further explored the role of this increase of cortical ADRB1 in addiction or related psychiatric disorders of offspring. Administration of betaxolol, a selective ADRB1 antagonist, blocked reinstatement of cocaine CPP [19]. Further, blocking ADRB1 impaired reconsolidation and reinstatement of cocaine self-administration [40]. Here, we found that paternal METH exposure increased ADRB1 on mPFC neurons. We believe that this higher ADRB1 is kind of compensatory effect for brain injury in offspring of sires exposed to drugs, which could strengthen the sensitivity of neurons to internal or external stimuli. Following METH[F1] treatment, we observed a continuous higher ADRB1 level in METH-sired male F1. Blocking ADRB1 or knocking-down ADRB1 on CaMKII-positive neurons (ADRB1[CaMKII]) efficiently reduced METH[F1]-evoked mPFC activation, and ultimately blocked

METH^{F1}-induced CPP in METH-sired male F1. There results indicate that

ADRB1^{CaMKII} mediate mPFC activity, which is one key intrinsic molecule of

mPFC for the paternal METH exposure-induced higher sensitivity in male

offspring.

ADRB1 activates neurons by eliciting its downstream protein kinases and

signal pathways (e.g. PKA, β-arrestin2), or mediating the function of nearby

channels (e.g. TREK-2-like channels) or receptors (e.g. GluR1) on the

membrane of cells. The critical role of ADRB1/cyclic-AMP dependent PKA in

synaptic plasticity of the frontal cortex are well-studied [41]. Recently, Huang et

al. found that ADRB1/β-arrestin/ERK signaling in entorhinal cortex, acting as a

non-classical G protein-coupled pathway, mediated memory reconsolidation

[28]. Their lab later reported that overexpression of β-arrestin2 in PFC

promoted extinction learning of cocaine-induced CPP, which was blocked by

propranolol, a nonbiased β-adrenergic receptor antagonist [42]. Thus, we

detected the molecules related to ADRB1, especially those of

β-arrestin2-dependent and PKA-dependent signal proteins. Unexpected, the

levels of β-arrestin2 and PKA failed to change in mPFC after knocking-down

ADRB1. While, we found that knocking-down ADRB1 effectively reduce the

levels of p-ERK1/2 and ΔFosB in mPFC. Son et al. [43] has been reported that

acupuncture exert its therapeutic effects on METH-induced addiction via

suppression of METH-increased p-ERK1/2 and ΔFosB in PFC. The p-ERK1/2

signaling cascades in PFC has been thought of important contributor for the

development of METH addiction [44]. ΔFosB, a transcription factor, is

increased in PFC by drugs including METH [43] and Δ9-tetrahydrocannabinol

(THC) [45], which leads to a long-lasting enhanced behavioral response.

Further, ΔFosB is preferentially induced by cocaine in adolescent animals,

which engaged in the higher vulnerability to addiction [46]. Unlike c-Fos, Arc

and ERK, ΔFosB maintains adaptive changes in brain for a long time following

drugs of abuse [47], which is involved in the long-term pathological plasticity of

brain and persistent addictive behavior by drugs [48]. Together with our

present study, these findings revealed that p-ERK1/2 and ΔFosB may be

subsequent signals for ADRB1 mediating heritable susceptibility to addiction in

offspring.

Collectively, our findings reveal a novel molecular mechanism underlying

paternal METH exposure-induced higher sensitivity to addiction in male

offspring, and mPFC ADRB1^{CaMKII} is a promising pharmacological target for

predicting or treating transgenerational susceptibility to addiction, especially

caused by paternal drug exposure.

Acknowledgments

This work is supported by National Natural Science Foundation of China

(82271531 and 82071495), Natural Science Foundation of Jiangsu Province,

China (BK20201398) and Natural Science Foundation of the Higher Education

Institutions of Jiangsu Province, China (21KJB360007).

Author contribution

Zheng Y and Liu D have equal contribution to the manuscript. Zheng Y, Liu D

and Li Z performed behavioral tests, morphological tests. Guo H, Chen W, Liu

D performed molecular experiments. Hu T, Zhang Y, Li X, Fan Y carried out the

virus-related experiment. Fan Y and Liu D performed data analysis. Zhao Z,

Cai Q and Ge F assist the data analysis. Guan X and Fan Y wrote the

manuscript. Liu D help to support the proposal. Guan X developed the overall

concept.

Conflict of interests

The authors declare that they have no competing interests.

Data availability statement

The data that support the findings of this study are available from the

corresponding author upon reasonable request.

Reference

[1] R.H.J. Verstegen, G. Wang, K.P.S. Langenberg-Ververgaert, L.Y. Ren, I. Nulman, Paternal exposure to recreational drugs before conception and its effect on live-born offspring: A scoping review, Birth defects research 112(13) (2020) 970-988.

[2] L.R. Goldberg, T.J. Gould, Multigenerational and transgenerational effects of paternal exposure to drugs of abuse on behavioral and neural function, The European journal of neuroscience 50(3) (2019) 2453-2466.

[3] L. Mihalčíková, A. Ochozková, R. Šlamberová, Does paternal methamphetamine exposure affect the behavior of rat offspring during development and in adulthood?, Physiol Res 70(S3) (2021) S419-S430.

[4] R. Cordie, L.M. McFadden, Optogenetic inhibition of the medial prefrontal cortex reduces methamphetamine-primed reinstatement in male and female rats, Behavioural pharmacology 30(6) (2019) 506-513.

[5] A. Parsegian, R.E. See, Dysregulation of Dopamine and Glutamate Release in the Prefrontal Cortex and Nucleus Accumbens Following Methamphetamine Self-Administration and During Reinstatement in Rats, Neuropsychopharmacology : official publication of the American College of Neuropsychopharmacology 39(4) (2014) 811-822.

[6] J.L. Perry, J.E. Joseph, Y. Jiang, R.S. Zimmerman, T.H. Kelly, M. Darna, P. Huettl, L.P. Dwoskin, M.T. Bardo, Prefrontal cortex and drug abuse vulnerability: translation to prevention and treatment interventions, Brain Res Rev 65(2) (2011) 124-149.

[7] H. Li, J.-A. Chen, Q.-Z. Ding, G.-Y. Lu, N. Wu, R.-B. Su, F. Li, J. Li, Behavioral sensitization induced by methamphetamine causes differential alterations in gene expression and histone acetylation of the prefrontal cortex in rats, BMC neuroscience 22(1) (2021) 24-24.

[8] R.Z. Goldstein, N.D. Volkow, Dysfunction of the prefrontal cortex in addiction: neuroimaging findings and clinical implications, Nature reviews. Neuroscience 12(11) (2011) 652-669.

[9] B. González, S. Jayanthi, N. Gomez, O.V. Torres, M.H. Sosa, A. Bernardi, F.J. Urbano, E. García-Rill, J.-L. Cadet, V. Bisagno, Repeated methamphetamine and modafinil induce differential cognitive effects and specific histone acetylation and DNA methylation profiles in the mouse medial prefrontal cortex, Progress in neuro-psychopharmacology & biological psychiatry 82 (2018) 1-11.

[10] M. Haddar, K. Uno, K. Azuma, S.-I. Muramatsu, A. Nitta, Inhibitory effects of Shati/Nat8l overexpression in the medial prefrontal cortex on methamphetamine-induced conditioned place preference in mice, Addiction biology 25(3) (2020) e12749.

[11] Y. Fan, Z. Li, Y. Zheng, X. Wei, Z. Zhang, Q. Cai, D. Liu, F. Ge, X. Guan, Sex-specific neurobehavioural outcomes and brain stimulation pattern in adult offspring paternally exposed to methamphetamine, Addiction biology 27(3) (2022) e13175.

[12] L. Coutellier, P.M. Ardestani, M. Shamloo, β1-adrenergic receptor activation enhances memory in Alzheimer's disease model, Ann Clin Transl Neurol 1(5) (2014) 348-360.

[13] A.M. Goodman, B.M. Langner, N. Jackson, C. Alex, L.L. McMahon, Heightened Hippocampal β-Adrenergic Receptor Function Drives Synaptic Potentiation and Supports Learning and Memory in the TgF344-AD Rat Model during Prodromal Alzheimer's Disease, The Journal of neuroscience : the official journal of the Society for Neuroscience 41(26) (2021) 5747-5761.

[14] C.A. Rudoy, A.-R.S. Reyes, E.J. Van Bockstaele, Evidence for beta1-adrenergic receptor
involvement in amygdalar corticotropin-releasing factor gene expression: implications for
cocaine withdrawal, Neuropsychopharmacology : official publication of the American College
of Neuropsychopharmacology 34(5) (2009) 1135-1148.
[15] Q. Sun, J. Huang, D.L. Yang, X.N. Cao, W.L. Zhou, Activation of β-adrenergic receptors
during sexual arousal facilitates vaginal lubrication by regulating vaginal epithelial Cl(-)
secretion, The journal of sexual medicine 11(8) (2014) 1936-48.
[16] Z.-B. You, E. Galaj, F. Alén, B. Wang, G.-H. Bi, A.R. Moore, T. Buck, M. Crissman, S. Pari,
Z.-X. Xi, L. Leggio, R.A. Wise, E.L. Gardner, Involvement of the ghrelin system in the
maintenance and reinstatement of cocaine-motivated behaviors: a role of adrenergic action at
peripheral β1 receptors, Neuropsychopharmacology : official publication of the American
College of Neuropsychopharmacology 47(8) (2022) 1449-1460.
[17] S. Sugama, T. Takenouchi, M. Hashimoto, H. Ohata, Y. Takenaka, Y. Kakinuma,
Stress-induced microglial activation occurs through β-adrenergic receptor: noradrenaline as a
key neurotransmitter in microglial activation, Journal of neuroinflammation 16(1) (2019) 266.
[18] X.-H. Ji, X.-H. Cao, C.-L. Zhang, Z.-J. Feng, X.-H. Zhang, L. Ma, B.-M. Li, Pre- and
Postsynaptic β-Adrenergic Activation Enhances Excitatory Synaptic Transmission in Layer
V/VI Pyramidal Neurons of the Medial Prefrontal Cortex of Rats, Cerebral Cortex 18(7) (2007)
1506-1520.
[19] O. Vranjkovic, S. Hang, D.A. Baker, J.R. Mantsch, β-adrenergic receptor mediation of
stress-induced reinstatement of extinguished cocaine-induced conditioned place preference in
mice: roles for β1 and β2 adrenergic receptors, The Journal of pharmacology and
experimental therapeutics 342(2) (2012) 541-51.
[20] A. Hernández, H. Burgos, M. Mondaca, R. Barra, H. Núñez, H. Pérez, R. Soto-Moyano, W.
Sierralta, V. Fernández, R. Olivares, L. Valladares, Effect of prenatal protein malnutrition on
long-term potentiation and BDNF protein expression in the rat entorhinal cortex after
neocortical and hippocampal tetanization, Neural plasticity 2008 (2008) 646919.
[21] B.L. Thompson, P. Levitt, G.D. Stanwood, Prenatal exposure to drugs: effects on brain
development and implications for policy and education, Nature reviews. Neuroscience 10(4)
(2009) 303-312.
[22] C.-C. Huang, Y.-C. Liang, K.-S. Hsu, Prenatal cocaine exposure enhances long-term
potentiation induction in rat medial prefrontal cortex, International Journal of
Neuropsychopharmacology 14(4) (2011) 431-443.
[23] T. Kalpachidou, A. Raftogianni, P. Melissa, A.M. Kollia, F. Stylianopoulou, A. Stamatakis,
Effects of a Neonatal Experience Involving Reward Through Maternal Contact on the
Noradrenergic System of the Rat Prefrontal Cortex, Cerebral cortex (New York, N.Y. : 1991)
26(9) (2016) 3866-3877.
[24] J. Meitzen, J.I. Luoma, C.M. Stern, P.G. Mermelstein, β1-Adrenergic receptors activate
two distinct signaling pathways in striatal neurons, Journal of neurochemistry 116(6) (2011)
984-95.
[25] S.K. Shenoy, M.T. Drake, C.D. Nelson, D.A. Houtz, K. Xiao, S. Madabushi, E. Reiter, R.T.
Premont, O. Lichtarge, R.J. Lefkowitz, beta-arrestin-dependent, G protein-independent
ERK1/2 activation by the beta2 adrenergic receptor, The Journal of biological chemistry 281(2)
(2006) 1261-73.

[26] J. Wang, C. Gareri, H.A. Rockman, G-Protein-Coupled Receptors in Heart Disease,
Circulation research 123(6) (2018) 716-735.
[27] D.G. Tilley, G protein-dependent and G protein-independent signaling pathways and their
impact on cardiac function, Circulation research 109(2) (2011) 217-30.
[28] B. Huang, Y. Li, D. Cheng, G. He, X. Liu, L. Ma, β-Arrestin-biased β-adrenergic signaling
promotes extinction learning of cocaine reward memory, Science Signaling 11(512) (2018).
[29] H. Lu, B. Lim, M.-m. Poo, Cocaine exposure in utero alters synaptic plasticity in the medial
prefrontal cortex of postnatal rats, The Journal of neuroscience : the official journal of the
Society for Neuroscience 29(40) (2009) 12664-12674.
[30] B.A. Morrow, T. Hajszan, C. Leranth, J.D. Elsworth, R.H. Roth, Prenatal exposure to
cocaine is associated with increased number of spine synapses in rat prelimbic cortex,
Synapse 61(10) (2007) 862-5.
[31] C. Crozatier, R.M. Guerriero, F. Mathieu, B. Giros, M. Nosten-Bertrand, B.E. Kosofsky,
Altered cocaine-induced behavioral sensitization in adult mice exposed to cocaine in utero,
Brain research. Developmental brain research 147(1-2) (2003) 97-105.
[32] Q.S. Yan, S.Z. Zheng, S.E. Yan, Prenatal cocaine exposure decreases brain-derived
neurotrophic factor proteins in the rat brain, Brain research 1009(1-2) (2004) 228-33.
[33] L. Cantacorps, S. Montagud-Romero, M. Luján, O. Valverde, Prenatal and postnatal
alcohol exposure increases vulnerability to cocaine addiction in adult mice, British journal of
pharmacology 177(5) (2020) 1090-1105.
[34] Z. Li, P. Santhanam, C.D. Coles, M. Ellen Lynch, S. Hamann, S. Peltier, X. Hu, Prenatal
cocaine exposure alters functional activation in the ventral prefrontal cortex and its structural
connectivity with the amygdala, Psychiatry research 213(1) (2013) 47-55.
[35] K. Rando, T.M. Chaplin, M.N. Potenza, L. Mayes, R. Sinha, Prenatal cocaine exposure
and gray matter volume in adolescent boys and girls: relationship to substance use initiation,
Biological psychiatry 74(7) (2013) 482-489.
[36] K. Grewen, M. Burchinal, C. Vachet, S. Gouttard, J.H. Gilmore, W. Lin, J. Johns, M. Elam,
G. Gerig, Prenatal cocaine effects on brain structure in early infancy, NeuroImage 101 (2014)
114-123.
[37] S.W. Yip, C.M. Lacadie, R. Sinha, L.C. Mayes, M.N. Potenza, Prenatal cocaine exposure,
illicit-substance use and stress and craving processes during adolescence, Drug and alcohol
dependence 158 (2016) 76-85.
[38] K. Grzelka, P. Kurowski, M. Gawlak, P. Szulczyk, Noradrenaline Modulates the Membrane
Potential and Holding Current of Medial Prefrontal Cortex Pyramidal Neurons via
β(1)-Adrenergic Receptors and HCN Channels, Front Cell Neurosci 11 (2017) 341-341.
[39] M.G. Henderson, M.M. McConnaughey, B.A. McMillen, Long-term consequences of
prenatal exposure to cocaine or related drugs: effects on rat brain monoaminergic receptors,
Brain research bulletin 26(6) (1991) 941-5.
[40] H. Zhu, Y. Zhou, Z. Liu, X. Chen, Y. Li, X. Liu, L. Ma, β1-Adrenoceptor in the Central
Amygdala Is Required for Unconditioned Stimulus-Induced Drug Memory Reconsolidation,
The international journal of neuropsychopharmacology 21(3) (2018) 267-280.
[41] O. Flores, H. Pérez, L. Valladares, C. Morgan, A. Gatica, H. Burgos, R. Olivares, A.
Hernández, Hidden prenatal malnutrition in the rat: role of β_1-adrenoceptors on synaptic
plasticity in the frontal cortex, Journal of neurochemistry 119(2) (2011) 314-23.

[42] X. Liu, L. Ma, H.H. Li, B. Huang, Y.X. Li, Y.Z. Tao, L. Ma, β-Arrestin-biased signaling
mediates memory reconsolidation, Proceedings of the National Academy of Sciences of the
United States of America 112(14) (2015) 4483-4488.
[43] J.S. Son, Y.C. Jeong, Y.B. Kwon, Regulatory Effect of Bee Venom on
Methamphetamine-Induced Cellular Activities in Prefrontal Cortex and Nucleus Accumbens in
Mice, Biological and Pharmaceutical Bulletin 38(1) (2015) 48-52.
[44] W.L. Sun, P.M. Quizon, J. Zhu, Molecular Mechanism: ERK Signaling, Drug Addiction, and
Behavioral Effects, Progress in molecular biology and translational science 137 (2016) 1-40.
[45] M.F. Lazenka, M. Kang, D.D. De, D.E. Selley, L.J. Sim-Selley, Δ9-Tetrahydrocannabinol
Experience Influences ΔFosB and Downstream Gene Expression in Prefrontal Cortex,
Cannabis and Cannabinoid Research 2(1) (2017) 224-234.
[46] M.E. Ehrlich, J. Sommer, E. Canas, E.M. Unterwald, Periadolescent mice show enhanced
DeltaFosB upregulation in response to cocaine and amphetamine, The Journal of
neuroscience : the official journal of the Society for Neuroscience 22(21) (2002) 9155-9.
[47] L.I. Perrotti, R.R. Weaver, B. Robison, W. Renthal, I. Maze, S. Yazdani, R.G. Elmore, D.J.
Knapp, D.E. Selley, B.R. Martin, L. Sim-Selley, R.K. Bachtell, D.W. Self, E.J. Nestler, Distinct
patterns of DeltaFosB induction in brain by drugs of abuse, Synapse 62(5) (2008) 358-69.
[48] E.J. Nestler, M. Barrot, D.W. Self, DeltaFosB: a sustained molecular switch for addiction,
Proceedings of the National Academy of Sciences of the United States of America 98(20)
(2001) 11042-6.
689

Figure legends

Figure 1. The CPP, mPFC activity and ADRB1 level in male F1. a,

Experimental design and timeline. **b**, CPP score by male F1. **c**, ΔCPP score by

male F1. **d,** Traveled traces in CPP apparatus by male F1. **e-f**, The percentage

of c-Fos-positive cells in Cg1, PrL and IL on PD60 and on PD73. CaMKII and

c-Fos were used as markers for the pyramidal neurons and activated neurons,

respectively. Scale bar, 500 μm /100 μm /20 μm. **g-h**, Levels of c-Fos and

CaMKII protein on PD60 and PD73. **i**, Levels of *Adrb1* mRNA on PD60 and

PD73. **j**, Levels of ADRB1 protein on PD60 and PD73. **k-l**, Mean gray value of

ADRB1 on mPFC neurons on PD60 and PD73. Scale bar, 500 μm /100 μm /20

μm. Cg1, cingulate cortex; PrL, prelimbic cortex; IL, infralimbic cortex.

SAL-sired male F1, saline-sired male F1 mice; METH-sired male F1,

methamphetamine-sired male F1 mice. N.S., $P > 0.05$. [#], $P < 0.05$ vs baseline

(Pre-test CPP score). *, $P < 0.05$, **, $P < 0.01$ vs SAL-sired male F1.

Figure 2. Effects of betaxolol on METH[F1]-induced CPP in METH-sired

male F1. a, Timeline of Experiment 1. **b**, CPP score. **c**, ΔCPP score. **d,**

Traveled traces in the CPP apparatus by METH-sired male F1. **e**, Timeline of

Experiment 2. **f**, CPP score. **g**, ΔCPP score. **h,** Traveled traces in the CPP

apparatus by METH-sired male F1. F1-METH-Ts-Vcl, vehicle bilaterally into

the mPFC of METH-sired male F1 at 15 min prior to CPP test;

F1-METH-Ts-Bxol, betaxolol bilaterally into the mPFC of METH-sired male F1

at 15 min prior to CPP test; F1-METH-Tr-Vcl, vehicle bilaterally into the mPFC

of METH-sired male F1 during CPP training; F1-METH-Tr-Bxol, betaxolol

bilaterally into the mPFC of METH-sired male F1 during CPP training. N.S., *P*

> 0.05. [##], $P < 0.01$ vs Pre-test CPP score. **, $P < 0.01$ vs F1-METH-Ts-Vcl or

F1-METH-Tr-Vcl.

Figure 3. Effects of knocking-down ADRB1[CaMKII] on METH[F1]-induced CPP

in METH-sired male F1. a, Experimental design and timeline. **b**, Schematic

diagram of viral injection and representative images of viral transfection in

mPFC (middle pannel). Scale bar, 500 µm /20 µm. **c**, Percentage of viral

transfected neurons in the total pyramidal neurons of mPFC. Scale bar, 500

µm /100 µm. **d**, Levels of *Adrb1* mRNA in mPFC. **e**, Levels of ADRB1 protein in

mPFC. **f**, Mean gray value of ADRB1 on virus transfected neurons of mPFC.

Scale bar, 500 µm /100 µm. **g**, CPP score. **h**, .ΔCPP score. **i**, Traveled traces

in the CPP apparatus by METH-sired male mice. F1-METH-Ctrl, METH-sired

male F1 mice injected with Ctrl virus. F1-METH-KD, METH-sired male F1 mice

injected with KD virus. N.S., $P > 0.05$. [##], $P < 0.01$ vs pre-test CPP score. *, $P <$

0.05, **, $P < 0.01$ vs F1-METH-Ctrl.

Figure 4. Effects of knocking-down ADRB1[CaMKII] on mPFC activity and

subsequent signals in METH-sired male F1. a, Levels of c-Fos protein. **b**,

The c-Fos immunostaining. Scale bar, 500 µm /100 µm. **c**, Density of dendritic

spine. Scale bar, 50 µm /10 µm. **d**, Levels of β-arestin2 and PKA protein. **e**,

Levels of ERK1/2, p-ERK1/2, ERK1/2/p-ERK1/2, CREB, p-CREB, p-CREB/

CREB and ΔFosB protein. F1-METH-Ctrl, METH-sired male F1 mice injected

with Ctrl virus. F1-METH-KD, METH-sired male F1 mice injected with KD virus.

N.S., $P > 0.05$. *, $P < 0.05$, **, $P < 0.01$, vs F1-METH-Ctrl.

Figure 5. Schematic summary of this study. Paternal methamphetamine

(METH) exposure induces higher ADRB1 levels on mPFC neurons in male F1

mice during adulthood. Subthreshold dose of METH treatment in F1 (METH^{F1}),

which is not sufficient to induce CPP, efficiently produces METH-related CPP

in METH-sired male F1 mice, accompanied with more activated mPFC

CaMKII-positive neurons than that in saline-sired male F1 mice. Locally

inhibiting ADRB1 function or specifically knocking-down ADRB1 on

CaMKII-positive neurons (ADRB1^{CaMKII}) efficiently reduces METH^{F1}-evoked

mPFC activation, and ultimately blocks METH^{F1}-induced CPP in METH-sired

male F1 mice. The p-EKR1/2 and ΔFosB might be potential subsequent

signals of ADRB1 in the process.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5