Cell-surface protein YwfG of *Lactococcus lactis* binds to α-1,2-linked mannose

Short title: Lactococcal cell-surface protein binds to α-1,2-linked mannose

Wataru Tsuchiya[1], Zui Fujimoto[1], Noritoshi Inagaki[1], Hiroyuki Nakagawa[1], Miwa Tanaka[2],

Hiromi Kimoto-Nira[3,#a], Toshimasa Yamazaki[1], Chise Suzuki[4,#a,*]

[1] Research Center for Advanced Analysis, National Agriculture and Food Research

Organization (NARO), Tsukuba, Ibaraki, Japan

[2] Asahi Quality & Innovations, Ltd, Moriya, Ibaraki, Japan

[3] Institute of Food Research, NARO, Tsukuba, Ibaraki, Japan

[4] College of Bioresource Science, Nihon University, Fujisawa, Kanagawa, Japan

[#a] Previous affiliation: Institute of Livestock and Grassland Science, NARO, Tsukuba, Ibaraki,

Japan

* Corresponding author

E-mail: suzuki.chise@nihon-u.ac.jp

Abstract

Lactococcus lactis strains are used as starter cultures in the production of fermented dairy and vegetable foods, but the species also occurs in other niches such as plant material. *Lactococcus lactis* subsp. *lactis* G50 (G50) is a plant-derived strain and potential candidate probiotics. Western blotting of cell-wall proteins using antibodies generated against whole G50 cells detected a 120-kDa protein. MALDI-TOF MS analysis identified it as YwfG, a Leu-Pro-any-Thr-Gly cell-wall-anchor-domain–containing protein. Based on a predicted domain structure, a recombinant YwfG variant covering the N-terminal half (aa 28–511) of YwfG ($YwfG_{28-511}$) was crystallized and the crystal structure was determined. The structure consisted of an L-type lectin domain, a mucin-binding protein domain, and a mucus-binding protein repeat. Recombinant YwfG variants containing combinations of these domains ($YwfG_{28-270}$, $YwfG_{28-311}$, $YwfG_{28-511}$, MubR4) were prepared and their interactions with monosaccharides were examined by isothermal titration calorimetry; the only interaction observed was between $YwfG_{28-270}$, which contained the L-type lectin domain, and D-mannose. Among four mannobioses, α-1,2-mannobiose had the highest affinity for $YwfG_{28-270}$ (dissociation constant = 34 µM). $YwfG_{28-270}$ also interacted with yeast mannoproteins and yeast mannan. Soaking of the crystals of $YwfG_{28-511}$ with mannose or α-1,2-mannobiose revealed that both sugars bound to the L-type lectin domain in a similar manner, although the presence of the mucin-binding protein domain and the mucus-binding protein repeat within the recombinant protein inhibited the interaction between the L-type lectin domain and mannose residues. Three of the YwfG variants (except MubR4) induced aggregation of yeast cells. Strain G50 also induced aggregation of yeast cells, which was abolished by deletion of *ywfG* from G50, suggesting that surface YwfG contributes to the interaction with yeast cells. These findings provide new structural and functional insights into the interaction between *L. lactis* and its ecological niche via binding of the cell-surface protein YwfG with mannose.

Introduction

Gram-positive bacteria interact with their environment and other cells via proteins
expressed on their cell surface. An important subset of these cell-surface proteins are the
adhesins that mediate adhesion to other cells or environmental surfaces. For example, in the
initial stages of colonization, pathogenic bacteria use their cell-surface proteins to bind to
host cells [1].

In *Staphylococcus aureus*, mucus-binding proteins are anchored to the cell surface by a
sortase A–mediated system that recognizes the motif LPXTG (Leu-Pro-any-Thr-Gly) [2]. In
this system, the mucus-binding proteins are cleaved between the Thr and Gly of the LPXTG
motif by sortase A and then undergo transpeptidation and covalent attachment to
peptidoglycan in the bacterial cell wall. Sortase-anchored surface proteins fulfill key
functions during the infectious process and thus sortase A is essential for host colonization
and for the pathogenesis of invasive diseases. The roles of sortases and proteins containing
the LPXTG motifs are well documented in pathogens, but even in nonpathogenic lactic acid
bacteria a mechanism that mediates the relationship between cell surface proteins and the
host that is analogous to that in pathogenic bacteria has been identified [3, 4].

Lactobacilli are Gram-positive bacteria in order *Lactobacillales*, strains of which are
found in the human intestine and used in the production of fermented foods. The strains that
are used as probiotics are mostly of intestinal origin and are reported to adhere to the
intestinal mucus layer [5]. Understanding the interaction between lactobacilli and mucins is
critical to elucidate the survival strategies of lactobacilli in a constantly changing intestinal
environment, as well as to elucidate the mechanisms of their beneficial effects in the
intestinal tract. Cell-surface proteins likely play a key role in these strategies and mechanisms

by interacting with host receptors, such as the glycans in mucins [3, 5]. Lactobacilli express

several adhesins, including moonlighting proteins (multi-functional cytoplasmic proteins)

that exert adhesive functions when expressed on the cell surface [6]. Several species of

lactobacilli are known to possess surface-layer (S-layer) proteins [7] that are involved in the

adhesion of the cells to the host epithelium or to extracellular matrix components and in

modulating the host immune response [3]. For example, knockout of S-layer protein A in

Lactobacillus acidophilus significantly reduces binding of the mutant to DC-SIGN, a

dendritic cell (DC)–specific receptor, and S-layer protein A is involved in the modulation of

DC and T-cell functions [8]. Whereas certain strains of lactobacilli bind to epithelial cells via

mucin in order to reside in the intestinal tract, other strains bind yeast cells used in

fermentation processes [9]. In some cases, symbiosis between the yeast and lactobacilli

improves fermentation [10, 11].

Lactococcus is another genus in order *Lactobacillales*. Strains belonging to *Lactococcus*

lactis and *Lactococcus cremoris* are used as fermentation starters in the production of cheese

and other dairy and vegetable fermented products. Originally, it was thought that lactococci

did not reach the intestinal tract alive, but Klijn et al. [12] showed that a fraction of viable

cells of a strain of *L. lactis* was able to survive passage through the human gastrointestinal

tract. Recently, several lactococcal strains have been reported to have beneficial health effects

such as improvement of skin status [13] and activation of human plasmacytoid DCs [14].

There are few studies on lactococcal surface proteins that interact with intestinal epithelium

or DCs.

Lactococcus lactis is used extensively in the industrial production of fermented dairy

products. However, *L. lactis* strains have also been isolated from plant material [15, 16].

Lactococcus lactis subsp. *lactis* G50 (G50) was isolated from Napier grass and was

subsequently found to have immunostimulatory activity [17]. *In vitro*, intact G50, but not

heat-treated G50, can induce tumor necrosis factor α expression in the murine macrophage

cell line J774.1, suggesting that a heat-sensitive molecule is involved in this induction [18].

In ovomucoid-sensitized BALB/c mice, oral administration of intact G50 cells reduces total

IgE production [17]. Together, these findings indicate that G50 may have potential uses as a

probiotic.

Analysis of the G50 genome sequence [19] may help reveal the cell-surface proteins that

contribute to the activity of G50. Here, we detected a cell-surface protein by reaction of *L.*

lactis strains with anti-G50 antibody. We used matrix-assisted laser

desorption/ionization–time-of-flight mass spectrometry (MALDI-TOF MS) and X-ray

crystallography to investigate the structure and interactions of this protein.

Materials and Methods

Strains

The lactococcal strains used in this study are listed in Table 1. The strains were grown at

30°C for 18 h in M17 (BD Difco, Sparks, MD, USA) broth containing 0.5% (wt/vol) glucose

(GM17) or chemically defined medium [20] in which lactose was replaced with 2% glucose

(CDMG) [21]. *Escherichia coli* BL21(DE3) (Merck, Darmstadt, Germany) was used for the

expression of recombinant proteins. Three yeast strains were also used: *Saccharomyces*

cerevisiae X2180-1A, *S. cerevisiae* AB9 [MAT a/α *gpi10/gpi10 ura3/URA3 leu2/LEU2*], and

S. cerevisiae AB9-2 [MAT a/α *gpi10/gpi10 mnn2/mnn2 ura3/URA3 leu2/LEU2*] [22]. YPD

containing 1% yeast extract, 2% peptone, and 2% glucose was used for culturing the yeast

strains. A synthetic defined medium (SD-medium) (yeast nitrogen base without amino acids

containing 0.17% ammonium sulfate [BD Difco], 2% glucose, 0.13% casamino acids [BD

Difco]) was used for yeast culture for mannoprotein preparation.

Table 1. Lactococcal strains used in this study

Strain	Species	YwfG*	Origin [reference]
G50	*Lactococcus lactis* subsp. *lactis*	+	Napier grass [17]
5A	*L. lactis* subsp. *lactis*	-	*ΔywfG* of G50, this study
12A	*L. lactis* subsp. *lactis*	-	*ΔywfG* of G50, this study
S63	*L. lactis* subsp. *lactis*	-	Kimchi [18]
O29	*Lactococcus cremoris*	+	Kefir [15]
H46	*L. lactis* subsp. *lactis*	+	Kimchi [15]
H54	*L. lactis* subsp. *lactis*	+	Kimchi [15]
DRC1	*L. lactis* subsp. *lactis* bv. diacetylactis	+	NIRD [23]
N7	*L. lactis* subsp. *lactis* bv. diacetylactis	+/-	Commercial cheese starter [23, 24]
CVT8w	*L. lactis* subsp. *lactis* bv. diacetylactis	+	Commercial cheese starter [25]
CVT3w	*L. lactis* subsp. *lactis* bv. diacetylactis	+	Commercial cheese starter [25]
H59	*L. lactis* subsp. *lactis* bv. diacetylactis	+	Kimchi [15]
NIAI 527	*L. lactis* subsp. *lactis*	-	Laboratory collection [23, 24]
NIAI 712	*L. cremoris*	-	Laboratory collection [23, 24]
C13	*L. lactis* subsp. *lactis*	-	Cheese [15]
O31	*L. lactis* subsp. *lactis*	-	Kefir [15]
1256	*L. lactis* subsp. *lactis*	-	Raw milk [15]
S01	*L. lactis* subsp. *lactis*	-	Fermented milk [15]
P79	*L. lactis* subsp. *lactis*	-	Fermented vegetables [15, 18]
342	*L. lactis* subsp. *lactis*	-	Cheese [15, 18]

119 *, The presence (+) or absence (-) of 120-kDa band corresponding to YwfG was examined by western

120 blotting.

Preparation of cell-wall proteins, surface-exposed proteins, and

cell lysates

124   Cell-wall proteins were extracted from the cell-wall fraction by using lysozyme, as

125 reported by Meyrand et al. [26]. Briefly, *L. lactis* cells were cultured to an optical density

126 (OD_{600}) of 1.2 in 6 mL of GM17 medium at 30°C and centrifuged at $5,000 \times g$ for 10 min at

127 4°C. The cell pellet was washed once with 50 mM Tris-HCl (pH 7.0) and resuspended in 1

mL of 50 mM Tris-HCl (pH 7.0) containing protease inhibitors (cOmplete, Roche Applied

Science, Mannheim, Germany). The cells were disrupted with glass beads using a FastPrep

system (Qbiogene, Montreal, Quebec, Canada) set at 6.0 m s^{-1} for 40 s and centrifuged at low

speed (1,000 $\times$ g) for 2 min at 4°C to remove the glass beads and any unbroken bacteria. The

supernatant was centrifuged at 16,000 $\times$ g for 15 min at 4°C. The pellet containing the

bacterial cell envelopes was digested with lysozyme (1 mg L^{-1}) and mutanolysin (200 U

mL^{-1}) in 50 mM Tris-HCl (pH 7.0) containing 5 mM MgCl$_2$ for 3 h at 37°C, centrifuged at

10,000 $\times$ g for 10 min at 4°C, and the supernatants were analyzed by western blotting.

Surface-exposed proteins were prepared according to the shaving method [26]. Bacterial

cells were digested by trypsin in the presence of 1.2 M sucrose and 1 mM CaCl$_2$, centrifuged

at 20,000 $\times$ g for 10 min, and supernatants were used for western blotting.

To prepare whole cell lysates, bacterial cultures (5 mL) were centrifuged, pellets were

washed twice with phosphate buffer saline (PBS), and suspended in 300 μL of PBS

containing 0.1% SDS, protease inhibitor cocktail (cOmplete), and Lysing matrix B (MP

Biomedicals, Santa Ana, USA). Cells were disrupted with glass beads using a FastPrep

system as described above, centrifuged at 13,000 $\times$ g for 10 min, and the supernatants were

used for western blotting.

SDS-PAGE and western blotting

SDS-PAGE was performed in a 5%–20% linear gradient gel (Atto, Tokyo, Japan).

Proteins were visualized by Coomassie Brilliant Blue staining (CBB) or western blotting.

After SDS-PAGE, proteins were electrotransferred to a polyvinylidene difluoride membrane

(Millipore, Burlington, MA, USA) using a semi-dry transfer blotter (Atto). Rabbit polyclonal

antiserum to G50 was generated against intact G50 cells and purified by using a recombinant

protein A column (rProtein A Sepharose FF, Cytiva, Tokyo, Japan). The membrane was

incubated with anti-G50 antiserum (1:3,000 dilution) and horseradish peroxidase–linked

anti-rabbit immunoglobulin (Bio-Rad Laboratories Inc., Hercules, CA, USA; 1:5,000

dilution), and the blot was developed with an ECL Prime western blotting kit (Cytiva).

Protein identification by MALDI-TOF MS and database searching

CBB-stained bands corresponding to the bands detected by the anti-G50 antiserum were

excised from the gel and treated by in-gel digestion with a proteomics-grade trypsin

(Sigma-Aldrich, St Louis, MO, USA). The digested samples were purified by using ZipTip

C18 (Merck). The purified peptides were mixed with an equal volume of

α-cyano-4-hydroxycinnamic acid (4-CHCA) (Shimadzu GLC Ltd., Tokyo, Japan) matrix

solution (prepared as 10 mg/mL in 50% acetonitrile containing 0.1% TFA), and an aliquot of

the mixture (0.5 μL) was spotted onto an ABI 4800 MALDI plate (SCIEX, Framingham, MA,

USA). The mass spectrometer, a MALDI-TOF/TOF mass spectrometer (4800 Plus TOF/TOF

analyzer, SCIEX) was tuned and calibrated using a calibration standards mixture (Peptide

Calibration Standard II, Bruker Daltonics GmbH & Co. KG, Bremen, Germany) prior to the

measurements. After the spot had been allowed to dry at room temperature, MS and MS/MS

spectra were obtained. Proteins were identified using a Mascot search engine (ver. 2.4.0.;

Matrix Science, London, UK) and the NCBI database.

Protein sequence and structure analysis

Protein sequence analysis and domain fold prediction were performed on the Pfam

(http://pfam.xfam.org) and the Phyre2 (http://www.sbg.bio.ic.ac.uk/phyre2)servers. Protein

structures were compared on the Dali server (http://ekhidna2.biocenter.helsinki.fi/dali/)[27].

Preparation of YwfG variants

Total bacterial DNA was isolated by using a DNeasy Blood & Tissue Kit (Qiagen, Hilden,

Germany). Primers were designed using the coding sequence (CDS) of locus tag

LLG50_11005 of the G50 genome sequence (accession number CP025500.1) (Table S1).

DNA fragments encoding variants of YwfG (residues 28–270, 28–336, 28–511, and

860–1034) were amplified by polymerase chain reaction using the primer sets Fw28–Rv270,

Fw28–Rv336, Fw28–Rv511, and Fw860–Rv1034, respectively. The resulting fragments were

inserted in NdeI/BamHI-digested pET28b (Merck) by In-Fusion cloning reactions (In-Fusion

HD Cloning Kit, Takara Bio, Kusatsu, Japan) with an N-terminal His$_6$-tag. The resulting

constructs were transformed into *E. coli* BL21(DE3). Recombinant YwfG variants were

expressed at 16°C overnight in the presence of 0.2 mM isopropyl β-D-thiogalactoside and

purified by affinity chromatography using a nickel-chelating column (HisTrap HP, Cytiva),

followed by gel filtration using a HiLoad 26/600 Superdex 75 pg column (Cytiva) and cation

exchange chromatography using a Mono S 10/100 GL column (Cytiva). The obtained

recombinant YwfG variants were named YwfG$_{28-270}$, YwfG$_{28-336}$, YwfG$_{28-511}$ (numbers

indicate amino acid residues), and MubR4 (residues 860–1034).

Preparation of yeast mannoproteins

gpi10 mutants of *S. cerevisiae* secrete mannoproteins into the culture medium [22].

Mannoproteins of AB9 strain (wild-type) and of AB9-2 strain (*mnn2* mutant lacking

α-1,2-mannosyltransferase) were prepared. Yeast cells were cultured in the SD-medium for 6

days at 30°C. The cultures were centrifuged at $12,000 \times g$ for 10 min at 4°C to obtain a

cell-free supernatant. The mannoproteins were concentrated using a 10,000

201   molecular-weight-cut-off ultrafiltration unit (Centriprep, Merk Millipore, Cork, Ireland),

202   precipitated by adding 2 volumes of ethanol, and dissolved in 20 mM Tris-HCl (pH 7.5).

Isothermal titration calorimetry (ITC) analysis

A MicroCal iTC200 system (Malvern Panalytical, Almelo, Netherlands) was used to

measure enthalpy changes associated with interactions between the YwfG variants and the

following yeast cell-wall components: the monosaccharides D-glucose, D-galactose,

D-mannose, D-fucose, N-acetyl-D-glucosamine (GlcNAc), and

2-(acetylamino)-2-deoxy-D-galactose (GalNAc) (Sigma-Aldrich); α-1,2-, α-1,3-, α-1,4-, and

α-1,6-mannobiose (Dextra Laboratories, Reading, UK); mannan from *S. cerevisiae*

(Sigma-Aldrich); and yeast mannoproteins.

A solution of YwfG variant (100 μM $YwfG_{28-270}$, $YwfG_{28-336}$, $YwfG_{28-511}$, or MubR4 in

20 mM Tris-HCl [pH 7.5]) was placed in a 200-μL calorimeter cell, and test solution (10 mM

for monosaccharides, 2 mM for monosaccharides and mannobioses, 10 mg mL^{-1} for yeast

mannan, OD_{280} 1.0 for mannoproteins) was loaded into the injection syringe. The test

solution was titrated into the sample cell as a sequence of 20 injections of 2-μL aliquots each.

All experiments were performed at 25°C.

Crystallization, data collection, and structure refinement

220   Purified $YwfG_{28-511}$ was concentrated to 12.0 mg mL^{-1} and crystallized by the sitting-drop

221   vapor diffusion method at 20°C using a precipitant solution consisting of 0.1 M sodium

222   acetate (pH 5.0–5.6), 1.3 M Li_2SO_4, and 2% (w/v) polyethylene glycol 3350 (Hampton

223   Research, Aliso Viejo, CA, USA). We used 50 μL of the reservoir solution and a drop

consisting of 0.5 µL of the protein solution and 0.5 µL of the reservoir solution; plate-type

crystals appeared within 2–3 weeks.

　　Diffraction data were collected at two beamlines: BL-5A housed at the Photon Factory

and BL-NW12 housed at the Photon Factory Advanced Ring (both at the High Energy

Accelerator Research Organization, Tsukuba, Japan). Diffraction data were collected using a

Pilatus3 S6M or Pilatus3 S2M detector (Dectris, Philadelphia, PA, USA). Crystals were

collected in a 0.5-mm nylon cryoloop (0.5 mm, Hampton Research), soaked in a precipitant

drop containing 30% (v/v) glycerol, and cryocooled in a nitrogen gas stream to 95 K. Crystals

in complex with D-mannose or α-1,2-mannobiose were obtained by adding 0.5 µL 10%

D-mannose or α-1,2-mannobiose solution to the crystal drop a few minutes prior to the

diffraction experiment. The data were integrated using the XDS program package [28] and

scaled using the SCALA program [29] within the CCP4 program suite [30].

　　The ligand-free structure of $YwfG_{28-511}$ was determined by the molecular replacement

method using the Molrep program [31] with the lectin domain (residues 258–492) of SraP

(PDB code 4m01 [32]) as the search model. Sugar complex structures were built using the

ligand-free structure as the starting model. Manual model building and molecular refinement

were performed using Coot [33] and Refmac5 [34]. Data collection and refinement statistics

are shown in Table 2. The stereochemistry of the models was analyzed using the MolProbity

program [35] and structural drawings were prepared using the PyMOL

(https://www.pymol.org/pymol.html) or Cuemol2 program (http://cuemol.osdn.jp/en/).

Table 2. Data collection and refinement statistics of the YwfG fragment structures

	Native	D-mannose complex	α-1,2-mannobiose complex
PDB code	7YL4	7YL5	7YL6
Data collection			
Space group	*I*222	*I*222	*I*222

Cell parameters (Å)	$a = 43.2$	$a = 43.3$	$a = 43.5$
	$b = 272.9$	$b = 273.2$	$b = 276.6$
	$c = 320.3$	$c = 322.8$	$c = 321.8$
X-ray source	PF-AR BL-NW12	PF BL-5A	PF-AR BL-NW12
Wavelength (Å)	1.00000	1.00000	1.00000
Resolution (Å)	48.6–2.50 (2.64–2.50)	48.7–3.00 (3.16–3.00)	43.06–2.95 (3.11–2.95)
No. of reflections	871 880 (127 582)	507 098 (73 143)	546 645 (84 290)
Unique reflections	66 894 (9 601)	39 549 (5 675)	42 029 (6 070)
Completeness (%)	100.0 (100.0)	100.0 (100.0)	99.9 (100.0)
Multiplicity	13.0 (13.3)	12.8 (12.9)	13.0 (13.9)
R-merge	0.093 (0.810)	0.145 (1.053)	0.203 (1.187)
Average I/σ	21.3 (3.4)	16.0 (2.7)	14.0 (2.5)
Refinement			
Resolution (Å)	45.5–2.50 (2.57–2.50)	45.6–3.00 (3.08–3.00)	41.5–2.95 (3.03–2.95)
R-factor	0.246 (0.385)	0.227 (0.365)	0.211 (0.361)
R-free	0.280 (0.395)	0.296 (0.392)	0.268 (0.387)
No. of water molecules	74	33	41
Average B-value (Å^2)	65.1	79.9	65.2
r.m.s.d. from ideals			
lengths (Å)	0.011	0.021	0.014
angles (°)	1.75	2.28	2.03
Ramachandran plot (%) (favored/allowed/disallowed)	94.7 / 5.0 / 0.3	93.7 / 5.5 / 0.8	95.4 / 4.3 / 0.3

*Values in parentheses refer to the highest-resolution shell. r.m.s.d., root-mean-square deviation.

Interaction of yeast cells and YwfG variants and lactococcal cells

Saccharomyces cerevisiae X2180-1A was cultured in YPD medium to an OD$_{600}$ of 0.6. In

microplate wells, 50 μL yeast culture, 10 μL each recombinant YwfG protein (0.5 μM or

1 μM as indicated in Fig 7), and 40 μL of PBS, D-mannose (0.5 mM), methyl α-D-mannoside

(0.5 mM), or each mannobiose (1 mM) were mixed and left to settle for 1–2 h, and

aggregation was observed.

Lactococcal cells were cultured in CDMG and suspended in PBS to an OD_{600} of 1.0. Then,

60 μL of each yeast and the lactococcal cells were added to 80 μL PBS and left to settle for

1–2 h, and aggregation was observed.

Deletion of *ywfG* gene in *L. lactis* G50

The *ywfG* of *L. lactis* G50 was deleted using the pG^+host-based allelic exchange method

[36-38]. Construction of the plasmid (pCS339) used for *ywfG* deletion is described in S1 Fig.

L. lactis G50 was treated with lithium acetate and dithiothreitol [39] and used for

electroporation of pCS339. Chromosomal *ywfG* deletion *(ΔywfG)* strains were produced

according to the method of Oxaran et al. [37]. Briefly, erythromycin-resistant colonies grown

at 30°C were incubated in the presence of erythromycin at 37°C, which is a non-permissive

temperature for pG^+host replication in strains with pCS339 integrated into the chromosome.

Cells were incubated in the absence of erythromycin at 30°C to induce plasmid excision, and

then erythromycin-sensitive strains were selected at 37°C. *ywfG* deletion was confirmed by

PCR and western blotting (S1 Fig).

Results

Identification of cell-surface proteins that reacted with anti-G50 antibody

273   Five lactococcal strains showing high immunostimulatory activity in our previous study

274   (G50, S63, O29, H46, and H54) [18] were selected. Cell-wall proteins were prepared from

275   the cell-wall fraction by using lysozyme and analyzed by western blotting by using an

276   antibody generated against live G50 cells (Fig 1). A 37-kDa band was detected in all strains,

277   and a strong 120-kDa band was detected in G50, O29, H46, and H54, but not in S63.

Fig 1. Western blot and Coomassie Brilliant Blue staining of lactococcal cell-wall proteins. (A) Western blot of cell-wall proteins of five lactococcal strains using anti-G50 antibody. (B) Coomassie Brilliant Blue staining of the same samples. The black triangle indicates the thin band of the 120-kDa protein.

In addition, a surface-exposed protein fraction was prepared from five strains (G50, S63, O29, P79, and 342) by trypsin cleavage of cell surface–exposed proteins in the presence of 1.2 M sucrose. A 120-kDa band was detected in G50 and O29 (S2A Fig). Cell lysates of 13 strains were also analyzed by western blotting (S2B Fig). The results suggested that YwfG is located both at the cell surface and in the cytoplasm, so not all of the expressed YwfG is covalently bound to the cell wall. Among the 18 lactococcal strains examined, the 120-kDa band was detected in 9 strains (Table 1).

The CBB-stained 120-kDa and 37-kDa bands from the fraction of cell-wall proteins were excised, subjected to in-gel trypsin digestion, and analyzed by MALDI-TOF MS and a Mascot search. The obtained peptide fragments from the 120-kDa protein were all aligned to the translated sequence of a CDS (LLG50_11005, GenBank: AUS70730.2) of the G50 genome (S3 Fig). This protein consists of 1107 residues and contains mucus-binding protein repeats and the LPXTG motif. The peptide fragments from the 120-kDa protein from strains O29, H46, and H54 were almost the same as those from strain G50. Interestingly, a C-terminal peptide sequence DSKPTDVLPSTGDSQK containing LPSTG was detected in the 120-kDa proteins of strains G50, O29, H46, and H54, suggesting that this LPXTG-motif is not used for anchoring the protein to the cell wall.

Amino acid sequence analysis using the Phyre2 and Pfam servers indicated that the 120-kDa protein contained a leguminous-type (L-type) lectin domain (Lec), a mucin-binding

protein domain (Mbp), and four mucus-binding protein repeats (MubRs), followed by a

C-terminal LPXTG motif (Fig 2). From its similarity to the LPXTG

cell-wall-anchor-domain–containing protein of *L. lactis* subsp. *lactis* Il1403 (GenBank:

AYV53882.1) [40], we named the protein YwfG, although the N-terminal 180 amino acid

residues of YwfG of G50, including the lectin domain, are missing from that of Il1403.

Fig 2. Schematic drawing of YwfG and constructions of variants. (Top) Predicted domain

organization of YwfG. (Bottom) Recombinant variants constructed in the present study.

   A Mascot search suggested that the 37-kDa protein was a phosphoglycerate kinase

(GenBank: AUS68744.1). Because phosphoglycerate kinases have been reported to be

cell-surface adhesins in *Ligilactobacillus agilis* [41] and Streptococci [42], the present

lactococcal phosphoglycerate kinase could possibly function as a moonlighting protein on the

cell surface.

Interactions of the YwfG variants with monosaccharides and mannobioses

   The expression of the whole YwfG protein has been attempted but expression and

stability were so low that the quantity and purity of the protein could not be achieved for use

in experiments. Then, four recombinant YwfG variants (YwfG$_{28–270}$, Lec domain; YwfG$_{28–336}$,

Lec and Mbp; YwfG$_{28–511}$, Lec, Mbp, and MubR1; and MubR4) (Fig 2) were expressed and

purified. The binding of the four variants with six monosaccharides (D-glucose, D-galactose,

D-mannose, D-fucose, GlcNAc, and GalNAc) and methyl α- D-mannoside was examined

using ITC. The only combination of variant and monosaccharide that showed an interaction

was YwfG$_{28–270}$ with D-mannose; that is, the Lec domain alone could bind to D-mannose (S4

Fig).

The binding of the three variants to four mannobioses (α-1,2-, α-1,3-, α-1,4-, and

α-1,6-mannobiose) was also examined using ITC. The thermodynamic parameters for the

interactions observed in the ITC assay are shown in Table 3. No binding of YwfG$_{28–336}$, or

YwfG$_{28–511}$ to any mannobiose was observed. YwfG$_{28–270}$ bound to all of the mannobioses,

but the binding capacity of α-1,2-mannobiose was the greatest; the dissociation constant of

α-1,2-mannobiose (K_D = 34 μM) was much lower than that of D-mannose (K_D = 500 μM).

Table 3. Thermodynamic parameters from isothermal titration calorimetry assay

Substrate	N (sites)	K_D (μM)	ΔH (cal/mol)	ΔS (cal/mol/deg)
D-mannose*	(1.00)	500 ± 37	−2098 ± 83.59	8.07
α-1,2-mannobiose	1.19 ± 0.0104	34.4 ± 0.97	−5236 ± 62.07	2.86
α-1,3-mannobiose	0.757 ± 0.148	172 ± 21	−6073 ± 1387	−3.15
α-1,4-mannobiose	1.82 ± 0.171	328 ± 91	−2206 ± 309.7	9.55
α-1,6-mannobiose	1.21 ± 0.0839	210 ± 13	−4235 ± 372.8	2.62

337   * Thermodynamic parameters determined with fixed stoichiometry (N = 1).

338   N, stoichiometry; K_D, dissociation constant; ΔH, enthalpy change; ΔS, entropy change.

Crystal structure of YwfG$_{28–511}$

Next, we elucidated the crystal structure of recombinant YwfG$_{28–511}$. We solved it by the

molecular replacement method at 2.50-Å resolution (Table 2). The crystal belonged to space

group *I*222 with two molecules (I and II) per asymmetric unit. Molecule I was modeled with

continuous residues from 40–511, whereas only residues 41–417 and 446–465 were modeled

in molecule II. The electron density map of N-terminal residues 28–39 could not be modeled

in either molecule. YwfG$_{28–511}$ fragment had an overall elongated shape with an overall

diameter of 40 Å and length of 180 Å, comprised 35 β-sheets and 4 α-helices, and was

formed by head-to-tail tandem connection of the three domains, Lec, Mbp, and MubR1 (Fig

3A, B). Compared with molecule I, molecule II had a more bent structure with a larger angle

between domains (Fig 3C).

Fig 3. Overall structure of YwfG$_{28–511}$. (A) X-ray crystal structure and (B) topology diagram

of YwfG$_{28–511}$ molecule I. The Lec, Mbp, Sub-N, Sub-C domains, and calcium ion are colored

blue, green, orange, red, and black, respectively. (C) Superimposition of the crystal structures

of YwfG$_{28–511}$ molecules I (green) and II (red).

Starting from the N-terminus, the first domain (residues 41–268) had a jelly roll–type

β-sandwich structure comprising 15 β-strands and 3 short α-helices. The core structure

consisted of an antiparallel β-sheet fold formed from two sheets: sheet-I (β1, β3, β5, β10-12,

β14, and β15) and sheet-II (β2, β4, β6-β9, and β13). A calcium-binding site was observed

between sheet-II and the loop connecting the β7 and β8 strands (loop 7–8). One Ca^{2+} ion was

coordinated by Asp152 (Oδ1 and Oδ2) belonging to β7; Tyr154 (O), Asn156 (Oδ1), and

Asp166 (Oδ1) located in loop 7–8; and one water molecule. Structural similarity analysis

using the Dali server [27] revealed that this domain had the highest similarity to *S. aureus*

SraP lectin domain (PDB code 4m01), with a structural alignment Z-score of 35.5 and

root-mean-square deviation (r.m.s.d.) of 1.4 Å over 226 aligned residues (Fig 4A).

Fig 4. Comparison of YwfG domain structures. (A) Superimposed ribbon models of the

Lec domain of YwfG (green) and *Staphylococcus aureus* SraP (light blue, PDB code 4m01).

(B) Superimposition of the Mbp domain structures of YwfG (green) and *L. monocytogenes*

Imo0835 (yellow, PDB code 2kvz; pink, PDB code 2kt7). Superimposition of the (C) Sub-N

and (D) Sub-C structures of YwfG MubR1 on the B1 and B2 subdomains, respectively, of *L.*

reuteri MubRV (cyan, PDB code 4mt5) and MubR5 (magenta, PDB code 3i57).

The second domain (residues 271–336) comprised six β-strands folded in a β-sandwich

composed of a mixed β-sheet formed from β16, β17, β19, and β21 and an antiparallel β-sheet

consisting of β18 and β20. This structure is similar to the Mbp domains of *Listeria*

monocytogenes mucin-binding and putative peptidoglycan-binding protein Imo0835 (PDB

codes 2kt7 and 2kvz, respectively) with a structural alignment Z-score of 7.7–8.2 and r.m.s.d.

of 1.6–1.9 Å over 64–66 Cα atoms (Fig 4B).

The third domain (residues 337–511) had a structure in which two subdomains were

connected by a three-stranded β-sheet. The N-terminal subdomain (Sub-N, residues 337–409)

had a structure called a ubiquitin-like β-grasp fold, similar to that found in proteins in the

immunoglobulin-binding superfamily, formed from six β-strands (β22–β27) and one α-helix

(α4). The C-terminal subdomain (Sub-C, residues 410–511) consisted of a second

domain-like fold formed from six β-strands (β28, β29, β32–β35) and a twisted antiparallel

β-sheet (β28, β30, β31) located between the subdomains. The modeling of Sub-C in molecule

(II) was incomplete because the electron density map was unclear. Structural comparison

analysis revealed that the third YwfG domain showed high similarity with MubR folds of a

mucus-binding protein from *Limosilactobacillus reuteri*, Mub-RV (PDB code 4mt5) [4] and

Mub-R5 (PDB code 3i57) [43], with a Z-score of 12.3–14.5 and r.m.s.d. of 3.3–6.3 Å over

155–165 aligned residues. The Sub-N and Sub-C domains of YwfG were very similar to the

B1 and B2 domains of the *L. reuteri* MubRs, respectively, although the β-sheet of the third

domain of YwfG is more bent (Fig 4C, D).

Substrate recognition mechanism

A structural analysis of the substrate-binding complex was performed to elucidate the

substrate recognition mechanism of YwfG for the glycoconjugates with which an interaction

was confirmed by the ITC assay. To obtain crystals of substrate-bound YwfG, ligand-free

crystals were soaked in a solution containing D-mannose or α-1,2-mannobiose. The structures

of the D-mannose– and α-1,2-mannobiose–YwfG$_{28–511}$ complexes were determined at 3.00 Å

and 2.95 Å, respectively. In each complex, a single substrate molecule was bound to the

recognition site, which was located in the groove-like region between loop 7–8 and loop

13–14 on the YwfG$_{28–270}$ surface and partially overlapped with the calcium-binding site.

In the Man–YwfG$_{28–511}$ complex, the recognition of D-mannose by the Lec domain was

the result of a hydrogen bond network involving six amino acid residues (Fig 5A). The

D-mannose-OH6 group was recognized by the main-chain N of Gly250 and Phe252 and the

main-chain O of Phe252 in loop 13–14. The main-chain amide of Ala251 also formed a

hydrogen bond with D-mannose-O5. The D-mannose-OH4 group interacted with the

sidechain of Asn156 (Nδ2) coordinating with the calcium ion and sidechain of Asp117 (Oδ1)

in the β6 strand. Furthermore, the D-mannose-OH3 group interacted indirectly with the

main-chain N of Gly136 in loop 6–7 via a water molecule. In addition, the aromatic sidechain

of Tyr154 formed a CH-π stacking interaction with the hydrophobic face of D-mannose.

Fig 5. Substrate recognition by the YwfG Lec domain. (A) D-mannose and (B)

α-1,2-mannobiose binding mechanism. The sugar substrate molecules and the YwfG residues

that interact with these are indicated by the yellow and green ball-and-stick models,

respectively. Calcium ions are shown as blue-white spheres.

The crystal structure of the α-1,2-mannobiose–YwfG$_{28–511}$ complex showed a wider range

of substrate interactions than was seen for the recognition of mannose (Fig 5B). The OH4

group of the non-reducing terminal mannose residue (mannose-1) formed a hydrogen bond

with the sidechain of the Ser81 (Oγ) in loop 3–4. The reducing terminal mannose residue

(mannose-2) was recognized in a manner similar to that of the monosaccharide mannose. The

mannose-2-OH1 group was not involved in the recognition mechanism and was oriented

toward the open space outside the ligand-binding pocket.

428 Interaction of YwfG variants with yeast mannoproteins

Yeast *gpi10* mutants secrete mannoproteins into the culture media [22]. Here, secreted

mannoproteins were concentrated from the culture filtrate by ultrafiltration and ethanol

precipitation. Interaction of $YwfG_{28–270}$ with mannoproteins was examined using ITC.

Thermal energetic changes, which indicate an interaction, were observed for mannoproteins

obtained from AB9 and AB9-2 and for the commercial yeast mannan. Yeast mannoproteins

are highly glycosylated cell-wall proteins consisting of 20% protein and 80% D-mannose.

Mannoprotein from AB9 is wild-type high-mannose glycan, whereas that from AB9-2 is

high-mannose glycan lacking α-1,2-linked mannose. Consistent with the observed binding

capacity to α-1,2-mannobiose (Table 2), the interactions of $YwfG_{28–270}$ with mannoprotein

from AB9 and yeast mannan were more intense than that with the mannoprotein from AB9-2

(Fig 6).

Fig 6. Interaction of $YwfG_{28–270}$ with yeast mannan proteins. Isothermal titration of

$YwfG_{28–270}$ against (A) AB9 (wild type) mannoprotein, (B) AB9-2 (*mnm2* mutant)

mannoprotein, (C) SD-medium, and (D) yeast mannan.

445 Interaction of YwfG variants with yeast cells

The interactions of the four recombinant variants (YwfG$_{28-270}$, YwfG$_{28-336}$, YwfG$_{28-511}$,

and MubR4) with yeast X2180-1A, which expresses high-mannose glycans on its cell surface,

were investigated (Fig 7). When X2180-1A and each recombinant variant (1 µM) were added

to a well of a round-bottomed titer plate, the cells were distributed uniformly at the bottom of

the wells in the presence of YwfG$_{28-270}$, YwfG$_{28-336}$, YwfG$_{28-511}$, but not in the presence of

MubR4 (Fig 7A). Non-aggregated yeast cells settle toward the center of round-bottomed

vessels and wells, but when aggregated, they assume a uniform distribution across the bottom

of the vessel or well. Such a uniform distribution or "creeping" phenotype [44] is observed in

glycosylation-deficient mutants [45] and in mating yeast cells. Arras et al. [44] have

described that yeast cells 'creep' up the sides of the culture vessel when cells of opposite

mating types are mixed, and they have shown that this 'creeping' phenotype is a consequence

of cell aggregation. The aggregation induced by YwfG$_{28-336}$ but not by YwfG$_{28-270}$ or

YwfG$_{28-511}$ was inhibited by the addition of 1 mM D-mannose or α methyl- D-mannoside (Fig

7B). Similarly, when 1 mM mannobiose was added, the aggregation induced by YwfG$_{28-336}$

and YwfG$_{28-511}$ but not by YwfG$_{28-270}$ was clearly inhibited in the presence of

α-1,2-mannobiose and was weakly inhibited in the presence of α-1,3-, α-1,4-, or

α-1,6-mannobiose (Fig 7C). Aggregation was observed in the presence of *L. lactis* G50, but

this aggregation was inhibited by the addition of 1 mM methyl α-D-mannoside. Furthermore,

aggregation was not observed in the presence of *ΔywfG* strains (Fig 7D), indicating that

cell-surface YwfG contributes to the interaction with yeast cells.

Fig 7. Interaction of yeast cells with YwfG variants and *L. lactis*. (A) Yeast cells in the

presence of 1 µM of each variant (270 [YwfG$_{28-270}$], 366 [YwfG$_{28-366}$], 511 [YwfG$_{28-511}$], MR

[MubR4]) or PBS. (B) Yeast cells in the presence of 0.5 µM of each variant and 0.5 mM

D-mannose or 0.5 mM methyl α-D-mannoside (methyl-M). (C) Yeast cells in the presence of

1 µM of each variant and 1 mM of each mannobiose (α-1,2, α-1,3, α-1,4, or

α-1,6-mannobiose). (D) Yeast cells in the presence of *L. lactis* G50 or the *ywfG* deletion

strains (5A, 12A) in the presence or absence of methyl-M.

Discussion

Here, we found that YwfG is a cell-surface protein that reacts with a strain-specific

antibody against *L. lactis* G50. This protein is reportedly to be an SrtA substrate [40]. Sortase

cleaves between the Thr and Gly of this motif and covalently anchors the substrate protein to

the cell wall. We detected a peptide sequence including LPSTG (DSKPTDV<u>LPSTG</u>DSQK)

by MALDI-TOF MS not only in YwfG of strain G50 but also in that of the other strains

examined. Since YwfG was detected in the surface-exposed protein fraction, in cell lysates,

and in the cell-wall protein fraction, it is unclear whether YwfG with an uncleaved LPXTG

motif is located not only in the cytoplasm but also at the cell surface. The *ΔywfG* strains did

not cause yeast aggregation, suggesting that YwfG on the cell surface contributes to the

aggregation or interaction with mannose residues.

Structural analysis revealed that $YwfG_{28-511}$ is composed of three different domains: Lec,

Mbp, and MubR. These domains have structural features of many gram-positive bacterial cell

wall adhesins, suggesting that YwfG has a similar function [4, 43, 46, 47]. Adhesins are

considered to function most effectively when their substrate-binding sites protrude as far as

possible from the bacterial surface [4]. *Lactococcus lactis* G50 YwfG has a Lec domain at the

tip of an approximately 600–650-Å long column consisting of one Mbp and four MubRs

extending from the cell wall, suggesting that the structure is suitable for binding to distant

target glycans.

The Lec domains of YwfG and *S. aureus* SraP have similar structures, although their

substrate specificities differ markedly. The legume (L)-type lectin–like module of SraP

specifically binds to *N*-acetylneuraminic acid and may mediate adhesion of *S. aureus* to host

sialylated receptors and interaction of *S. aureus* with host epithelial cells [32]. In contrast, the

ITC analysis in the present study revealed that $YwfG_{28-270}$ exhibits high affinity for

D-mannose and mannobiose, especially α-1,2-mannobiose. Crystallographic analysis of the

YwfG–substrate complexes provided structural evidence of this substrate specificity. In the

recognition mechanism of α-1,2-mannobiose by YwfG, mannose-2 (the reducing terminal

mannose residue) was involved in multiple interactions, whereas only one hydrogen bond

was observed for mannose-1 (the non-reducing terminal mannose residue), indicating that

mannose-2 and mannose-1 are the major and auxiliary recognition sites, respectively.

Furthermore, the mannose-2-OH1 group is not involved in the interaction and is oriented

toward the outside of the substrate-binding pocket. Since the OH1 group contributes to the

polymerization of sugars, we hypothesize that the Lec domain of YwfG will also bind to

longer-chain sugars possessing the mannose-α-1,2-mannose structure. This difference in the

sugar-binding affinities of the Lec domains of lactococcal YwfG and the L-type lectin-like

modules of staphylococcal SraP may explain the different host specificities, with the former

interacting with plant substrates or yeast cells and the latter interacting with intestinal

epithelium.

Because YwfG was recognized by an anti-G50 antibody that was raised against intact

G50 cells, YwfG was recognized as an epitope and could interact with host immune cells. *L.*

reuteri mucus-binding protein repeats, Mub-R5 and Mub-RV, which have high structural

similarity to the MubR domain of YwfG, have been reported to bind not only to mucin

glycans but also to mammalian secretory Ig such as IgA, IgM, and IgG at the mucosal surface

of the digestive tract [43]. These interactions of the *L. reuteri* MubRs are considered to

contribute to the promotion of bacterial retention in the host mucosal epithelium and the

modulation of host immune responses [43]; MubR of YwfG may have similar functions.

Mannose-specific adhesin (Msa) of *Lactiplantibacillus plantarum* WCSF1 also contains a

MubR and an L-type lectin domain, as does YwfG, and its mannose-specific binding function

is considered an essential factor for the probiotic properties of *L. plantarum* WCSF1 [46].

The lectin domains of Msa and YwfG, which contribute to mannose-specific binding, have

only 37% amino acid identity. Mannose binding may be necessary for *L. lactis* to bind to

plant substrates or yeast cells. Similarly, it may play an important role in interactions with

mannose-containing receptors and other factors in the gastrointestinal tract. Further studies

are required to elucidate the role of *L. lactis* YwfG in the host–probiotic interaction and the

host response when bacterial adhesins adhere to intestinal epithelial cells or immune cells.

A Blastp search [48] showed that at least 55 lactococcal strains have YwfG homologs

with more that 90% identity and query cover. YwfG is composed of a complete or incomplete

N-terminal Lec domain and various numbers of MubRs. The plant-derived *L. lactis* KF147

has a CDS of YwfG (GenBank: ADA65975.1) with complete Lec, Mbp, and two MubR

domains. *Lactococcus cremoris* MG1363, a plasmid-free progeny of the dairy starter strain

NCDO712, has a CDS of YwfG (GenBank: CAL99029.1) with complete Lec, Mbp, and three

MubR domains. The complete set of amino acid residues involved in mannose and calcium

binding (D117, D152, D166, N156, Y154, and A251) are conserved in the Lec domains of

YwfG of strains KF147, MG1363, and G50. YwfG was not detected in the surface proteome

of *L. lactis* NZ9000, a derivative of MG1363 [49]. The authors digested cell-surface proteins

with trypsin in the presence of sucrose and DTT and analyzed the resulting peptides by

LC/MS, but because YwfG has no cysteine residue, it may be trypsin resistant even in the

presence of DTT. Indeed, we did observe the 120 kDa band in western blots of cell

surface–exposed proteins prepared by using trypsin (S2A Fig).

In contrast, *L. lactis* Il1403 has a CDS of YwfG (GenBank: AYV53882.1) lacking half

(120 amino acid residues) of the N-terminal Lec domain of YwfG. The preceding CDS

(GenBank: AYV53883.1) encodes 157 amino acids that are similar to the N-terminal 140

amino acid residues of YwfG of G50. There is a single-base deletion between 2259225 and

2259227 that fragments the complete YwfG sequence. It is unclear whether this is a

sequencing error or an actual mutation. The YwfG of Il1403 has been confirmed to be a

cell-wall-associated protein that is present in the wild-type strain and absent in the sortase A

mutant [40], suggesting that those proteins are likely substrates of sortase A.

Strains of *L. lactis* are widely used in the production of fermented vegetables and various

fermented dairy products, and these strains are frequently isolated from non-dairy niches,

including soil and plant substrates [50]. The plant-derived *L. lactis* KF147 has gene sets for

degrading complex plant polymers and is adapted to grow on substrates derived from plant

cell walls [51]. Some plant cell walls contain galactomannans and glucomannans, which are

polymers consisting primarily of mannose (α-D-mannosides). Degradation of mannans to

D-mannose can be followed by conversion to D-fructose by mannose isomerase. In the

present study, a 35-kb region (1462553 to 1497747) of the G50 genome showed more than

98% identity with 100% coverage with more than 40 *Lactococcus* strains including KF147

and Il1403. This region encodes various proteins involved in glycan degradation and

transport, such as fructokinase, β-glucosidase, endo-β-*N*-acetylglucosaminidase,

glucohydrolase, lacto-*N*-biosidase, sugar transporters, α-mannosidases, and xylose operon.

These proteins therefore may contribute to the degradation of mannans and/or plant substrates

and uptake of monosaccharides for the growth of the strains. Anchoring lectins that bind to

mannose-containing glycans distributed in plant cell walls to the cell-surface layer may be a

necessary strategy for degrading plant substrates.

Here, we experimentally clarified the structure and function of the lactococcal cell surface

protein YwfG. Categorized as a mucus-binding protein, this protein is homologous to the

Staphylococcus aureus SraP lectin, but unlike the lectins of the pathogenic bacteria, YwfG

bound mannose. These findings provide new structural and functional insights into the

interaction between *L. lactis* and its ecological niche, showing that these interactions are

distinct from the bacteria–host interactions of pathogenic bacteria.

Acknowledgements

576 The authors thank Ms. Tomoko Sato for her outstanding technical support in the

577 MALDI-TOF MS analysis and Dr. E. Maguin for supplying the plasmid pG$^+$host9.

References

1. Chan JM, Gori A, Nobbs AH, Heyderman RS. Streptococcal serine-rich repeat proteins in colonization and disease. Front Microbiol. 2020;11:593356. doi: 10.3389/fmicb.2020.593356.

2. Schneewind O, Missiakas D. Sortases, surface proteins, and their roles in *Staphylococcus aureus* disease and vaccine development. Microbiol Spectr. 2019;7(1). doi: 10.1128/microbiolspec.PSIB-0004-2018.

3. Muscariello L, De Siena B, Marasco R. *Lactobacillus* cell surface proteins involved in interaction with mucus and extracellular matrix components. Curr Microbiol. 2020;77(12):3831-3841. doi: 10.1007/s00284-020-02243-5.

4. Etzold S, Kober OI, Mackenzie DA, Tailford LE, Gunning AP, Walshaw J, et al. Structural basis for adaptation of lactobacilli to gastrointestinal mucus. Environ Microbiol. 2014;16(3):888-903. doi: 10.1111/1462-2920.12377.

5. Nishiyama K, Sugiyama M, Mukai T. Adhesion properties of lactic acid bacteria on intestinal mucin. Microorganisms. 2016;4(3). doi: 10.3390/microorganisms4030034.

6. Savijoki K, Nyman TA, Kainulainen V, Miettinen I, Siljamäki P, Fallarero A, et al. Growth mode and carbon source Impact the surfaceome dynamics of *Lactobacillus rhamnosus* GG. Front Microbiol. 2019;10:1272. doi: 10.3389/fmicb.2019.01272.

7. Hynönen U, Palva A. *Lactobacillus* surface layer proteins: structure, function and applications. Appl Microbiol Biotechnol. 2013;97(12):5225-5243. doi: 10.1007/s00253-013-4962-2.

8. Konstantinov SR, Smidt H, de Vos WM, Bruijns SC, Singh SK, Valence F, et al. S layer protein A of *Lactobacillus acidophilus* NCFM regulates immature dendritic cell and T cell functions. Proc Natl Acad Sci U S A. 2008;105(49):19474-19479. doi: 10.1073/pnas.0810305105.

9. Yamasaki-Yashiki S, Sawada H, Kino-Oka M, Katakura Y. Analysis of gene expression profiles of *Lactobacillus paracasei* induced by direct contact with *Saccharomyces cerevisiae* through recognition of yeast mannan. Biosci Microbiota Food Health. 2017;36(1):17-25. doi: 10.12938/bmfh.BMFH-2016-015.

10. Hirayama S, Furukawa S, Ogihara H, Morinaga Y. Yeast mannan structure necessary for co-aggregation with *Lactobacillus plantarum* ML11-11. Biochem Biophys Res Commun. 2012;419(4):652-655. doi: 10.1016/j.bbrc.2012.02.068.

11. Nejati F, Junne S, Neubauer P. A big world in small grain: a review of natural milk kefir starters. Microorganisms. 2020;8(2). doi: 10.3390/microorganisms8020192.

12. Klijn N, Weerkamp AH, de Vos WM. Genetic marking of *Lactococcus lactis* shows its survival in the human gastrointestinal tract. Appl Environ Microbiol. 1995;61(7):2771-2774. doi: 10.1128/aem.61.7.2771-2774.1995.

13. Kimoto-Nira H. New lactic acid bacteria for skin health via oral intake of heat-killed or live cells. Anim Sci J. 2018;89(6):835-842. doi: 10.1111/asj.13017.

14. Sugimura T, Jounai K, Ohshio K, Tanaka T, Suwa M, Fujiwara D. Immunomodulatory effect of *Lactococcus lactis* JCM5805 on human plasmacytoid dendritic cells. Clin Immunol. 2013;149(3):509-518. doi: 10.1016/j.clim.2013.10.007.

15. Nomura M, Kobayashi M, Narita T, Kimoto-Nira H, Okamoto T. Phenotypic and molecular characterization of *Lactococcus lactis* from milk and plants. J Appl Microbiol. 2006;101(2):396-405. doi: 10.1111/j.1365-2672.2006.02949.x.

16. Fallico V, McAuliffe O, Fitzgerald GF, Ross RP. Plasmids of raw milk cheese isolate *Lactococcus lactis* subsp. *lactis* biovar diacetylactis DPC3901 suggest a plant-based origin for the strain. Appl Environ Microbiol. 2011;77(18):6451-6462. doi: 10.1128/AEM.00661-11.

17.   Kimoto H, Mizumachi K, Okamoto T, Kurisaki J. New *Lactococcus* strain with
immunomodulatory activity: enhancement of Th1-type immune response. Microbiol
Immunol. 2004;48(2):75-82. doi: 10.1111/j.1348-0421.2004.tb03490.x.
18.   Suzuki C, Kimoto-Nira H, Kobayashi M, Nomura M, Sasaki K, Mizumachi K.
Immunomodulatory and cytotoxic effects of various *Lactococcus* strains on the murine
macrophage cell line J774.1. Int J Food Microbiol. 2008;123(1-2):159-165. doi:
10.1016/j.ijfoodmicro.2007.12.022.
19.   Nakano K, Minami M, Shinzato M, Shimoji M, Ashimine N, Shiroma A, et al.
Complete genome sequence of *Lactococcus lactis* subsp. *lactis* G50 with immunostimulating
activity, isolated from Napier grass. Genome Announc. 2018;6(8). doi:
10.1128/genomeA.00069-18.
20.   Otto R, Tenbrink B, Veldkamp H, Konings WN. The relation between growth rate
and electrochemical proton gradient of *Streptococcus cremoris*. FEMS Microbiology Letters.
1983;16(1):69-74.
21.   Suzuki C, Kobayashi M, Kimoto-Nira H. Novel exopolysaccharides produced by
Lactococcus lactis subsp. *lactis*, and the diversity of *epsE* genes in the exopolysaccharide
biosynthesis gene clusters. Biosci Biotechnol Biochem. 2013;77(10):2013-2018. doi:
10.1271/bbb.130322.
22.   Tanaka M, Odani T, Yuuki T, Ohtake Y, Shimma Y-I, Yoko-O T, et al. Yeast strain
releasing mannan protein and method of producing mannan protein WO2006025295A1 2005.
23.   Nomura M, Kobayashi M, Okamoto T. Rapid PCR-based method which can
determine both phenotype and genotype of *Lactococcus lactis* subspecies. Appl Environ
Microbiol. 2002;68(5):2209-2213. doi: 10.1128/AEM.68.5.2209-2213.2002.
24.   Nomura M, Kimoto H, Someya Y, Suzuki I. Novel characteristic for distinguishing
Lactococcus lactis subsp. *lactis* from subsp. *cremoris*. Int J Syst Bacteriol. 1999;49 Pt
1:163-166. doi: 10.1099/00207713-49-1-163.
25.   Kimoto H, Nomura M, Suzuki I. Growth energetics of *Lactococcus lactis* subsp.
lactis biovar diacetylactis in cometabolism of citrate and glucose. Int Dairy J.
1999;9(12):857-863.
26.   Meyrand M, Guillot A, Goin M, Furlan S, Armalyte J, Kulakauskas S, et al. Surface
proteome analysis of a natural isolate of *Lactococcus lactis* reveals the presence of pili able to
bind human intestinal epithelial cells. Mol Cell Proteomics. 2013;12(12):3935-3947. doi:
10.1074/mcp.M113.029066.
27.   Holm L, Rosenstrom P. Dali server: conservation mapping in 3D. Nucleic Acids Res.
2010;38(Web Server issue):W545-549. doi: 10.1093/nar/gkq366.
28.   Kabsch W. XDS. Acta Crystallogr D Biol Crystallogr. 2010;66(Pt 2):125-132. doi:
10.1107/S0907444909047337.
29.   Evans P. Scaling and assessment of data quality. Acta Crystallogr D Biol Crystallogr.
2006;62(Pt 1):72-82. doi: 10.1107/S0907444905036693.
30.   Winn MD, Ballard CC, Cowtan KD, Dodson EJ, Emsley P, Evans PR, et al.
Overview of the CCP4 suite and current developments. Acta Crystallogr D Biol Crystallogr.
2011;67(Pt 4):235-242. doi: 10.1107/S0907444910045749.
31.   Vagin A, Teplyakov A. Molecular replacement with MOLREP. Acta Crystallogr D
Biol Crystallogr. 2010;66(Pt 1):22-25. doi: 10.1107/S0907444909042589.
32.   Yang YH, Jiang YL, Zhang J, Wang L, Bai XH, Zhang SJ, et al. Structural insights
into SraP-mediated *Staphylococcus aureus* adhesion to host cells. PLoS Pathog.
2014;10(6):e1004169. doi: 10.1371/journal.ppat.1004169.
33.   Emsley P, Cowtan K. Coot: model-building tools for molecular graphics. Acta
Crystallogr D Biol Crystallogr. 2004;60(Pt 12 Pt 1):2126-2132. doi:
10.1107/S0907444904019158.

34.   Murshudov GN, Skubak P, Lebedev AA, Pannu NS, Steiner RA, Nicholls RA, et al.
REFMAC5 for the refinement of macromolecular crystal structures. Acta Crystallogr D Biol
Crystallogr. 2011;67(Pt 4):355-367. doi: 10.1107/S0907444911001314.
35.   Chen VB, Arendall WB, 3rd, Headd JJ, Keedy DA, Immormino RM, Kapral GJ, et
al. MolProbity: all-atom structure validation for macromolecular crystallography. Acta
Crystallogr D Biol Crystallogr. 2010;66(Pt 1):12-21. doi: 10.1107/S0907444909042073.
36.   Biswas I, Gruss A, Ehrlich SD, Maguin E. High-efficiency gene inactivation and
replacement system for Gram-positive bacteria. J Bacteriol. 1993;175(11):3628-3635. doi:
10.1128/jb.175.11.3628-3635.1993.
37.   Oxaran V, Ledue-Clier F, Dieye Y, Herry JM, Pechoux C, Meylheuc T, et al. Pilus
biogenesis in *Lactococcus lactis*: molecular characterization and role in aggregation and
biofilm formation. PLos One. 2012;7(12):e50989. doi: 10.1371/journal.pone.0050989.
38.   Maguin E, Prevost H, Ehrlich SD, Gruss A. Efficient insertional mutagenesis in
lactococci and other Gram-positive bacteria. J Bacteriol. 1996;178(3):931-935. doi:
10.1128/jb.178.3.931-935.1996.
39.   Papagianni M, Avramidis N, Filioussis G. High efficiency electrotransformation of
Lactococcus lactis spp. *lactis* cells pretreated with lithium acetate and dithiothreitol. BMC
Biotechnol. 2007;7:15. doi: 10.1186/1472-6750-7-15.
40.   Dieye Y, Oxaran V, Ledue-Clier F, Alkhalaf W, Buist G, Juillard V, et al.
Functionality of sortase A in *Lactococcus lactis*. Appl Environ Microbiol.
2010;76(21):7332-7337. doi: 10.1128/AEM.00928-10.
41.   Kinoshita H, Ohuchi S, Arakawa K, Watanabe M, Kitazawa H, Saito T. Isolation of
lactic acid bacteria bound to the porcine intestinal mucosa and an analysis of their
moonlighting adhesins. Biosci Microbiota Food Health. 2016;35(4):185-196. doi:
10.12938/bmfh.16-012.
42.   Amblee V, Jeffery CJ. Physical features of intracellular proteins that moonlight on
the cell surface. PLoS One. 2015;10(6):e0130575. doi: 10.1371/journal.pone.0130575.
43.   MacKenzie DA, Tailford LE, Hemmings AM, Juge N. Crystal structure of a
mucus-binding protein repeat reveals an unexpected functional immunoglobulin binding
activity. J Biol Chem. 2009;284(47):32444-32453. doi: 10.1074/jbc.M109.040907.
44.   Arras SDM, Hibbard TR, Mitsugi-McHattie L, Woods MA, Johnson CE, Munkacsi
A, et al. Creeping yeast: a simple, cheap and robust protocol for the identification of mating
type in *Saccharomyces cerevisiae*. FEMS Yeast Res. 2022;22(1). doi:
10.1093/femsyr/foac017.
45.   Suzuki C, Shimma YI. P-type ATPase *spf1* mutants show a novel resistance
mechanism for the killer toxin SMKT. Mol Microbiol. 1999;32(4):813-823. doi:
10.1046/j.1365-2958.1999.01400.x.
46.   Pretzer G, Snel J, Molenaar D, Wiersma A, Bron PA, Lambert J, et al.
Biodiversity-based identification and functional characterization of the mannose-specific
adhesin of *Lactobacillus plantarum*. J Bacteriol. 2005;187(17):6128-6136. doi:
10.1128/JB.187.17.6128-6136.2005.
47.   von Ossowski I, Satokari R, Reunanen J, Lebeer S, De Keersmaecker SC,
Vanderleyden J, et al. Functional characterization of a mucus-specific LPXTG surface
adhesin from probiotic *Lactobacillus rhamnosus* GG. Appl Environ Microbiol.
2011;77(13):4465-4472. doi: 10.1128/AEM.02497-10.
48.   Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al.
BLAST+: architecture and applications. BMC Bioinformatics. 2009;10:421. doi:
10.1186/1471-2105-10-421.
49.   Berlec A, Zadravec P, Jevnikar Z, Štrukelj B. Identification of candidate carrier
proteins for surface display on *Lactococcus lactis* by theoretical and experimental analyses of

the surface proteome. Appl Environ Microbiol. 2011;77(4):1292-1300. doi:
10.1128/AEM.02102-10.
50.   Yanagida F, Chen YS, Shinohara T. Searching for bacteriocin-producing lactic acid
bacteria in soil. J Gen Appl Microbiol. 2006;52(1):21-28. doi: 10.2323/jgam.52.21.
51.   Siezen RJ, Starrenburg MJ, Boekhorst J, Renckens B, Molenaar D, van Hylckama
Vlieg JE. Genome-scale genotype-phenotype matching of two *Lactococcus lactis* isolates
from plants identifies mechanisms of adaptation to the plant niche. Appl Environ Microbiol.
2008;74(2):424-436. doi: 10.1128/AEM.01850-07.
735
736

Supporting information

S1 Table Plasmids and primers used in this study.

S1 Fig A Map of the pCS339 plasmid for *ywfG* deletion. To construct this plasmid, a 984-bp upstream fragment (2240620–2241603) and a 991-bp downstream fragment (2236506–2237496) of the *ywfG* gene (LLG50_11005, complementary 2237254–2240577) were amplified by PCR; each fragment was ligated into the pGEM-T Easy plasmid, resulting in pCS336 and pCS337, respectively. The PstI fragment (967 bp) of pCS336 was ligated into the PstI site of pCS337, resulting in pCS338. SacII–digested pCS338 and pG+host9 were ligated, resulting in pCS339. **B**, Western blots of the cell-wall proteins of *L. lactis* G50, O29, H46, and *ΔywfG* strains.

S2 Fig A Western blots of surface-exposed proteins of lactococcal strains S63, G50, P79, 342, and O29. **B**, Western blots of cell lysates of 13 lactococcal strains.

S3 Fig Deduced amino acid sequence of LLG50_11005 and peptides obtained by MALDI-TOF MS analysis. Peptides obtained were aligned to the deduced sequence and are shown underlined and in bold. Peptides that were detected on both sides of the trypsin cleavage site are indicated by a triangle.

S4 Fig Interaction of YwfG$_{28-270}$ with monosaccharides. Isothermal titration of YwfG$_{28-270}$ against D-glucose, D-galactose, D-mannose, D-fucose, GlcNAc, GalNAc.

S1_raw_Fig 1 image. Two gels loaded with the same samples, one for CBB staining and one for western blotting, were simultaneously subjected to SDS-PAGE. Proteins were visualized by CBB staining (Left panel, Fig 1B) or western blotting visualizing by exposure with ECL regent (Right panel, Fig 1A). The middle panel is the PVDF membrane after exposure with ECL, visualizing the protein on which the ECL reagent bound.

S2_raw_Fig 7 image. Interaction of yeast cells and YwfG derivatives was observed in a

microtiter plate. Photographing yeast cells accumulated at the bottom of the wells requires

oblique lighting, and it is difficult to take a photograph all wells at once, so the photographs

were staggered and cut and pasted.

7YL4_D_1300030883_val-report-full_P1. A PDB Summary Validation Report for 7YL4,

the ligand-free structure of YwfG$_{28-511}$.

7YL5_D_1300030884_val-report-full_P1. A PDB Summary Validation Report for 7YL5,

the D-mannose complex structure of YwfG$_{28-511}$.

7YL6_D_1300030939_val-report-full_P1. A PDB Summary Validation Report for 7YL6,

the α-1,2-mannobiose complex structure of YwfG$_{28-511}$.

Fig 1

Fig 2

A

Lec

Mbp

180 Å

Sub-N

MubR1

Sub-C

B

Sheet-I   Sheet-II

β12   β10   β14   α2   β4   β6   β8

β3   β2

β15   α1

β11   β5   β1   β13   β7   β9

N

α3

β16   β18

β17   β20

β21

β19

β22   β24

β27   β26

β23   β25   α4

β31   β28

β30   β34   β32

β35

β29   β33

C

C

(I)

(II)

A

C

N

C

N

B

N

C

D

N

C

Fig 4

Fig 6

A

270 336 511 MR

YwfG variant 1 μM

PBS

B

PBS

YwfG variant 0.5 μM

+mannose

+methyl-M

270 336 511 MR

C

      - 270 336 511 MR

YwfG variant 1 μM

mannobiose

α-1,2 +

α-1,3 +

α-1,4 +

α-1,6 +

D

    +methyl-M

PBS

G50 wt

5A *ΔywfG*

12A *ΔywfG*

Fig 7