1 **Universal gut microbial relationships in the gut microbiome of wild baboons**
2

3 **Authors:** Kimberly E. Roche[1], Johannes R. Björk[2,3,4], Mauna R. Dasari[4,5], Laura Grieneisen[6,7],
4 David Jansen[4], Trevor J. Gould[6], Laurence R. Gesquiere[8], Luis B. Barreiro[9,10,11], Susan C.
5 Alberts[8,12,13], Ran Blekhman[6,14], Jack A. Gilbert[15], Jenny Tung[8,12,13,16], Sayan Mukherjee[1,17,18,19],
6 Elizabeth A. Archie[4]
7
Affiliations:
[1] Program in Computational Biology and Bioinformatics, Duke University, Durham, NC, USA
[2] University of Groningen and University Medical Center Groningen, Department of
Gastroenterology and Hepatology, Groningen, The Netherlands
[3] University of Groningen and University Medical Center Groningen, Department of Genetics,
Groningen, The Netherlands
[4] Department of Biological Sciences, University of Notre Dame, Notre Dame, IN, USA
[5] Department of Biological Sciences, University of Pittsburgh, Pittsburgh, PA, USA
[6] Department of Genetics, Cell Biology, and Development, University of Minnesota,
Minneapolis, MN, USA
[7] Department of Biology, University of British Columbia-Okanagan Campus, Kelowna, BC,
Canada
[8] Department of Biology, Duke University, Durham, NC, USA
[9] Committee on Genetics, Genomics, and Systems Biology, University of Chicago, Chicago, IL,
USA.
[10] Section of Genetic Medicine, Department of Medicine, University of Chicago, Chicago, IL,
USA.
[11] Committee on Immunology, University of Chicago, Chicago, IL, USA.
[12] Department of Evolutionary Anthropology, Duke University, Durham, NC, USA
[13] Duke University Population Research Institute, Duke University, Durham, NC, USA
[14] Department of Ecology, Evolution, and Behavior, University of Minnesota, Minneapolis, MN,
USA
[15] Department of Pediatrics and the Scripps Institution of Oceanography, University of
California, San Diego, San Diego, CA, USA
[16] Department of Primate Behavior and Evolution, Max Planck Institute for Evolutionary
Anthropology, 04103 Leipzig, Germany
[17] Departments of Statistical Science, Mathematics, Computer Science, and Bioinformatics &
Biostatistics, Duke University, Durham, NC, USA
[18] Center for Scalable Data Analytics and Artificial Intelligence, University of Leipzig, Leipzig
Germany
[19] Max Plank Institute for Mathematics in the Natural Sciences, Leipzig, Germany
39
40 * Correspondence to: earchie@nd.edu

Abstract

Ecological relationships between bacteria mediate the services that gut microbiomes provide to their hosts. Knowing the overall direction and strength of these relationships within hosts, and their generalizability across hosts, is essential to learn how microbial ecology scales up to affect microbiome assembly, dynamics, and host health. Here we gain insight into these patterns by inferring thousands of correlations in bacterial abundance between pairs of gut microbiome taxa from extensive time series data, consisting of 5,534 microbiome profiles from 56 wild baboon hosts over a 13-year period. We model these time series using a statistically robust, multinomial logistic-normal modeling framework and test the degree to which bacterial abundance correlations are consistent across hosts (i.e., "universal") or individualized to each host. We also compare these patterns to two publicly available human data sets. We find that baboon gut microbial relationships are largely universal: correlation patterns within each baboon host reflect a mixture of idiosyncratic and shared patterns, but the shared pattern dominates by almost 2-fold. Surprisingly, the strongest and most consistently correlated bacterial pairs across hosts were overwhelmingly positively correlated and typically belonged to the same family—a 3-fold enrichment compared to pairs drawn from the data set as a whole. The bias towards universal, positive bacterial correlations was also apparent in monthly samples from human infants, and bacterial families that had universal relationships in baboons also tended to be universal in human infants. Together, our results advance our understanding of the relationships that shape gut microbial ecosystems, with implications for microbiome personalization, community assembly and stability, and the feasibility of microbiome interventions to improve host health.

Introduction

Mammalian gut microbiomes are highly diverse, dynamic communities whose members exhibit the full spectrum of ecological relationships, from strong mutualisms like syntrophy and cross-feeding, to competition, parasitism, and predation [1-4]. These relationships mediate a variety of biological processes that have powerful effects on host health and fitness, including the metabolism of complex carbohydrates and toxins, and the synthesis of physiologically important compounds, like short-chain fatty acids, neurotransmitters, and vitamins [1-8]. Despite their importance, major gaps remain in our understanding of microbial relationships in the gut microbiome [1, 9, 10]. We typically do not know if the abundance of one microbe consistently predicts the abundance of other microbes in the same host community, nor do we understand whether these correlative relationships are consistent in strength or direction across hosts ([10-13]).

Knowing the overall direction and strength of these correlative relationships is important, not only because they partly reflect the ecological relationships that mediate gut microbial processes, but also because overall correlation patterns can affect gut microbiome assembly, stability, and productivity [14, 15]. For instance, sets of microbes that exhibit strong, positive relationships within hosts sometimes represent networks of cooperating taxa that promote each other's growth [5, 9, 16]. In turn, these strong, mutualistic interdependencies can create an ecological house of cards where microbes rise and fall together, hampering community assembly and stability [14, 17]. In addition, understanding the degree to which correlative relationships between microbes are the same or different in different hosts can shed light on whether hosts share similar, underlying microbial ecologies [9, 10, 18-20]. Filling this knowledge gap has consequences for the generalizability of microbiome assembly processes, stability, and the

ecosystem services that emerge from microbiome dynamics to affect host health [9, 10, 12, 14, 17, 21].

To date, there are several reasons to think that correlative relationships in the gut microbiome will not be consistent across hosts and will instead be individualized to each host. For instance, several common community and evolutionary processes—such as horizontal gene transfer, genotype by environment interactions, and priority effects—can lead microbiome taxa to fill different ecological roles in different hosts [3, 22-26]. Further, some microbes can adopt context-dependent metabolisms and ecological roles depending on their microbial neighbors or other aspects of the environment—all phenomena that could lead to personalized interspecies relationships in gut microbiota [27-30]. Finally, the common observation that gut microbial community compositions (i.e., the presence and abundance of taxa) are highly individualized is sometimes proposed to arise from host-specific microbial ecologies and relationships [22-26, 31-35].

However, to date, the handful of studies that have tested the generalizability of gut microbial relationships across hosts suggest that these relationships are not highly individualized and are instead largely consistent (i.e., "universal") across hosts (**Fig. 1A**; [10, 18-20, 36]). For instance, Bashan et al. [10] inferred "universal" gut microbial relationships in the human gut microbiome by applying dissimilarity-overlap analysis (DOA) to cross-sectional samples from several human data sets. DOA infers universal microbial relationships by testing whether pairs of hosts who share many of the same microbiome taxa also tend to have similar abundances of those taxa [10, 18-20, 36]. This approach relies on the assumption that, when two communities share many of the same species and have similar abundances of those species, they do so because of a shared, underlying set of between-species abundance relationships [10, 36]. While many studies using this approach find evidence that microbial relationships are "universal" [10, 18-20], DOA's assumptions have been questioned because some conditions can lead to the spurious detection of universality, including environmental gradients, the strength of stochastic processes, and the presence of many non-interactive species [10, 36, 37].

An obvious alternative is to measure microbial correlations directly from microbiome time series collected from several hosts [9, 38]. Unlike DOA, this approach should be able to pinpoint which microbiome taxa exhibit the most and least consistent relationships across hosts. However, measuring microbial correlations from longitudinal, multi-host microbiome time series has its own challenges: time series with adequately dense sampling are rare, and most such data sets exhibit temporal autocorrelation and irregular sampling [38]. Further, the most common, and still most feasible, way to collect microbiome community data—via high-throughput sequencing—generates noisy count data that usually can only be interpreted in terms of relative (not absolute) abundances [39, 40].

To overcome these hurdles, here we combine extensive time-series data on the stool-associated microbiota with a multinomial logistic-normal modeling framework (**Fig. 1;** n=5,534 samples from 56 baboons; 75 to 181 samples per baboon across 6 to 13.3 years, between 2000 and 2013; [41-43]). This framework uses 16S rRNA sequencing count data to learn a smoothly evolving Gaussian process. The baboon hosts were the subject of long-term research on individually recognized animals by the Amboseli Baboon Research Project in Kenya, which has been studying baboon ecology and behavior in the Amboseli ecosystem since 1971 [41]. The baboons range over the same habitat and experience similar diets and sources of microbial colonization, facilitating inference about the consistency of microbial correlations across hosts (**Fig. S1;** [42, 43]). Our modeling approach accounts for variation attributable to seasonal

changes in the animals' diets, proportionality in the count data, and irregularity in sampling to
produce per-individual, per-taxon trajectories of log-ratio abundances that we used to estimate
pairwise microbial correlations within each host.
We pursued four main objectives. First, we characterized the overall sign and strength of
pairwise correlations in bacterial abundance within each host. Second, we tested the degree to
which these correlation patterns are systematically consistent across hosts or individualized by
host (**Fig. 1A**). Third, we identified taxonomic, phylogenetic, environmental, and host-related
predictors of the direction and universality of bacterial correlations. Finally, we tested the
generalizability of our findings by comparing the patterns of universality in our data set to two
microbiome time series from humans [34, 44].
Our predictions for these analyses were influenced by ideas from community and
microbial ecology. First, because strong interdependencies can hamper community assembly and
destabilize community dynamics [14, 15, 17], we expected that most microbial correlations
would be weak with few strong positive relationships between microbes. Second, consistent with
studies that used DOA, we expected that microbial relationships would be more consistent across
hosts than individualized (see **Fig. 1A** for a visualization of this prediction). This result would
suggest that personalized microbiota—their compositions and dynamics—do not arise from host-
specific microbiome ecologies [10, 18-20]. Third, because closely related gut bacteria may have
similar functional properties, we expected to observe many positive correlations between those
that are close phylogenetic relatives. Alternatively, competitive exclusion may lead closely
related taxa to exhibit neutral or negative relationships. Fourth, because the environments
experienced by baboons in Amboseli are far more uniform than those experienced by typical
human study subjects [42, 43], we expected that the signature of "universality" in baboons would
be stronger than that observed in humans. We discuss the implications of these patterns for
individual microbiome community assembly and dynamics, and for understanding how
microbiome communities are structured across hosts—a key requirement for successful
intervention to improve host health [10, 11, 45].

Figure 1. Testing the generalizability of gut microbial correlations across hosts. (A)
Schematic illustrating our approach for testing the degree to which gut microbial abundance
correlations are consistent (i.e., "universal" [10]) across different baboon hosts. The left-hand set
of images show our expectations for consistent correlation patterns; the right-hand images show
our expectation for individualized correlation patterns. Colored circles next to each baboon
represent prevalent microbial taxa found in at least 20% of samples in each host (and excluding
putative duplicate 16S gene copies; see methods). In each host, we inferred centered log-ratio
(CLR) abundance trajectories for these taxa using a multinomial-logistic normal modeling
approach implemented in the R package 'fido' [46]. Cartoons of two such trajectories for the
orange and blue taxa are below each baboon. We used these trajectories to infer covariances
between each pair of taxa in all baboons (represented by covariance matrices). We then
converted these covariances to Pearson's correlations and compared bacterial correlation patterns
across all hosts, shown as heat maps (red cells are positively correlated taxa; blue cells reflect
negatively correlated taxa). **(B)** Irregular time series of fecal samples used to infer microbial
CLR abundance trajectories in 56 baboon hosts (n=5,534 total samples; 75-181 samples per
baboon across 6 to 13.3 years). Each point represents a fecal sample collected from a known
individual baboon (y-axis) on a given date (x-axis). Samples from the same baboon were
collected a median of 20 days apart (range=0 to 723 days; 25th percentile=7 days, 75th
percentile =49 days). **(C)** Relative abundances of the 8 most prevalent gut bacterial orders and

179 families over time (x-axis) for all 56 hosts (samples from females are labeled with an F; male
180 samples with an M). Microbiota were somewhat individualized to each host (**Fig. S2**; [42, 43]).
181
182

Results
Most bacterial correlations within individuals are weak and negative
We began by characterizing the overall sign, strength, and significance of pairwise
correlations in bacterial abundance within each host. To do so, we applied the approach outlined
in **Fig. 1A** to stool-associated time series from 56 baboons (**Fig. 1B**) and calculated Pearson's
correlations between all possible pairs of bacterial taxa for three taxonomic partitions of the data.
These partitions were: (1) all pairs of CLR-transformed amplicon sequence variants (ASVs)
found in >20% of samples in each host and were unlikely to represent a duplicate 16S rRNA
gene copy ([47]; see Methods; n=125 ASVs; **Fig. 2A; Table S1**); (2) all pairs of bacterial phyla
found in >20% of samples in each host (n=12 phyla; **Table S2; Fig. S3**); and (3) all pairs of taxa
agglomerated to the most granular possible family, order, or class found in >20% of samples in
each host (n=37 taxa; **Table S3; Fig. S3**). We assessed the false discovery rate for each
correlation with a threshold for significance of FDR ≤ 0.05, by comparing the nominal p-values
for each observed correlation to an empirical permutation-based null, obtained by shuffling
taxonomic identities within microbiome samples 10 times for each host and re-calculating the
Pearson correlation p-values obtained from the permutations (**Fig. 2B**). We also confirmed that
the resulting correlation patterns were insensitive to several modeling choices and were not
primarily driven by seasonal shifts in microbiome composition (see results below and the
Supplement).
Consistent with the expectation that most bacterial correlations in the gut microbiome are
weak [14, 17], only 17% of ASV-ASV correlations in the heat map in **Fig. 2A** were stronger than
expected by chance (FDR ≤ 0.05; **Fig. S4A**; 20% of phylum-phylum; 21% of family/order/class
correlations; **Fig. S3**). The strongest negatively correlated pair in **Fig. 2A** included two ASVs in
the family Lachnospiraceae that had a median correlation of -0.562 (+/- 0.118 s.d.) across all
baboon hosts (**Fig. 2C**; ASV25 and ASV107; **Tables S1** and **S4**). The strongest positively
correlated pair of ASVs included two members of the genus *Prevotella* that had a median
correlation of 0.801 (+/- 0.053 s.d.) across all baboons (**Fig. 2D**; ASV2 and ASV3; **Tables S1**
and **S4**). While these two ASVs were assigned to the same genus, their V4 16S DNA sequence
identity was 97.6%, indicating they are probably not duplicate 16S gene copies in the same taxa
[47] (**Table S4**).
In support of the idea that positive bacterial interdependencies are rare [14, 15, 17], only
8.8% of ASV pairs were significantly positively correlated within hosts over time, and the
overall bacterial correlation patterns were slightly skewed towards negative relationships—
especially for relationships between bacterial phyla. For instance, at the ASV-level, the median
correlation coefficient in **Fig. 2A** was -0.016, and 53% of these correlations were negative
(binomial test p < 0.0001). For family/order/class-level taxa, 55% of all correlations in were
negative (**Figs. S3A** and **S4A**; median family/order/class-level correlation=-0.031; binomial test
p < 0.0001). Correlations between phyla exhibited the strongest negative skew, with 64% of
phyla-phyla correlations having a negative sign (**Figs. S3B and S4A;** median phyla-level
correlation=-0.092; binomial test p < 0.0001). This bias towards negative relationships may
reflect the fact that different phyla exhibit substantial differences in metabolism and lifestyle and
likely respond to distinct environmental drivers.

226
227
Figure 2. Bacterial correlation patterns across hosts. The heat map in panel **(A)** shows
Pearson's correlation coefficients of CLR abundances between all pairs of ASVs (x-axis) in each
of the 56 baboon hosts (y-axis). Each pair of ASVs is represented on the x-axis, including all
pairwise combinations of 125 ASVs resulting in 7,750 ASV-ASV pairs in each host (434,000
total correlations across all 56 hosts). Columns are ordered by the mean correlation coefficient
between ASV-ASV pairs, from negative (blue) to positive (red). **(B)** Pairwise correlations
generated from random permutations of the data. Taxonomic identities were shuffled within
samples and pairwise ASV-ASV correlations were estimated to produce a null model of ASV-
ASV correlation patterns within and between hosts. Column order is the same as in Panel A.
Panels **(C)** and **(D)** show example trajectories of CLR abundances for two pairs of ASVs in the
same five hosts. Panel **(C)** shows a strongly negatively correlated pair (median r across all
hosts=-0.562; two ASVs in family Lachnospiraceae: ASV25 (orange) and ASV107 (blue);
Tables S1 and S4) and panel **(D)** shows one strongly positively correlated pair (median r across
all hosts=0.801; two ASVs in the genus Prevotella 9; ASV2 (orange) and ASV3 (blue); **Tables**
S1 and S4).

Within-host bacterial correlation patterns are largely consistent across baboons

Next, we tested the degree to which within-host ASV-ASV correlations were consistent across hosts. We began by plotting the absolute value of each ASV pair's median Pearson's correlation coefficient as a function of the consistency of their correlation sign (positive or negative) across the 56 hosts (**Figs. 3A and 3B**). These plots provide two main insights into the consistency of bacterial associations. First, in support of the idea that ASVs do not exhibit vastly different correlative relationships in different hosts, no ASV pairs were both strongly and inconsistently correlated across hosts (**Figs. 3A and 3B**; **Fig. S5A**). Instead, the ASV pairs that had inconsistent correlation signs across hosts always had weak and often non-significant median absolute correlation coefficients within hosts (**Figs. 3A and 3B**). Second, the pairs with the most consistent sign agreement across hosts also exhibited the largest median absolute correlation coefficients across hosts (**Figs. 3A and 3B**; Spearman's r=0.844, p<0.0001). Hence, pairs of ASVs that have the strongest relationships, and are therefore likely to play the most important roles in structuring gut microbiome dynamics, also tend to have the most consistent relationships in different hosts. Indeed, for the sets of positively or negatively correlated ASV-pairs that showed universal agreement in the sign of their correlation across all hosts (i.e., where x=1 in **Figs. 3A and 3B**), the median correlation coefficient is 0.398, compared to 0.113 for those with no sign consistency (x=0.5 in **Figs. 3A and 3B**). Note, that the correlation strength for a given pair of ASVs was only weakly predicted by bacterial abundance. When both members of the pair were relatively abundant, pairs tended to exhibit stronger median correlations (r=0.012, p<0.0001**; Fig. S6**). However, while this effect is significant, it explained <1% of the variance in median correlation strength.

Visual inspection of the patterns in **Figs. 2A**, **3A, and 3B** indicate that ASV-ASV correlations are largely consistent across baboons, as opposed to individualized to each baboon. To explicitly quantify the relative strength of shared versus individualized signatures in the heat map in **Fig. 2A**, we calculated the population mean pattern for the ASV-ASV correlation matrix, m. For each host, we then estimated the residual difference, e, between that individual's observed ASV-ASV correlation matrix, y, and the population mean matrix: $y - m$ (see **Fig. S7A** for a cartoon example). We reasoned that the observed correlation matrix for each host can be approximated by a mixture of contributions from the population mean matrix m and the host-specific residual matrix e. To identify the optimal mixture for each host (i.e., the mixture of consistent vs. individualized correlation patterns that best explained the observed data), we titrated the contribution (i.e., weight) of e from 0% to 100% (and correspondingly, the contribution of m from 100% to 0%) and identified the value that minimized the Frobenius distance between the simulated combination and the observed correlation matrix, y.

In support of prior observations of "universality" [10, 18-20], we found that, across hosts, the optimal mixture involved contributions from the shared correlation structure (i.e., m) of between 50% and 70% (median 65%) and a host-level contribution (i.e., from e) of between 50% and 30% (median 35%). Hence, population-level signatures contributed almost twice the weight as host-level signatures (a median population:host ratio of 1.86:1; **Fig. S7B**). As a result, ASV-level relationships tend to be more consistent across hosts than host-specific.

284
285
Figure 3. **None of the ASV pairs were strongly and inconsistently correlated across hosts,**
and the strongest and most consistently correlated ASVs are typically positively correlated.
Plots in (**A**) and (**B**) show the median correlation strength for each ASV-ASV pair across all 56
hosts as a function of the consistency in direction of that pair's correlation across hosts,
measured as the proportion of hosts that shared the majority correlation sign (positive or
negative; ASV pairs that were positively correlated in half of the 56 hosts have a consistency of
0.5; ASV pairs that were positively [or negatively] correlated in all hosts have a consistency of
1.0). Panel (**A**) presents this relationship for consensus positively correlated features and panel
(**B**) shows consensus negatively correlated features. The Spearman correlation between median
association strength and the proportion of shared sign for all correlated features is 0.844 (p <
0.0001). Multiplying the two axes in either panel (**A**) or (**B**) creates a "universality score", whose
distribution is shown in panel (**C**). This score reflects the strength and consistency of pairwise
microbial correlations across hosts and ranges from 0 to 1, where a score of 1 indicates ASV-
ASV pairs with perfect correlations of the same sign in all hosts. A vertical line indicates the

minimum significant universality score. **(D)** Correlation networks for the top 2.5% most strongly and consistently correlated ASV pairs across hosts (i.e., the top 2.5% highest universality scores; pairs with rank 1-194 in **Table S4**). Network edges are colored by the consensus sign of the correlation between that pair (black for pairs where most hosts had a positive correlation; gray for pairs where most hosts had a negative correlation). Node labels indicate the ASV identity in **Table S1** and colors represent bacterial families. **(E)** Significantly enriched bacterial families in the network in panel D (Fisher's Exact Test p < 0.01 all, FDR ≤ 0.05; see **Table S5** enrichment statistics for all families). **(F)** Significantly enriched same-family pairings in the network in panel D (**Table S5**). Note that for visualization, the estimated log2 odds ratio intervals have been truncated at 5; full estimates are given in **Table S5.**

The most consistent ASV-level correlations are positive and between phylogenetically related taxa

One advantage of our approach, compared to dissimilarity overlap analyses [10], is we can identify the bacterial pairs that exhibit the most consistent relationships across hosts. Hence, we next conducted several analyses to understand why some pairs of ASVs exhibit more consistent correlation patterns across hosts than others. To do so, we created a "universality" score that could be calculated for each ASV pair. The score multiplies the pair's median correlation coefficient across hosts (y-axis of **Fig. 3A, 3B**) with its correlation consistency across hosts (i.e., proportion of shared sign; x-axis of **Fig. 3A, 3B**). The resulting scores range from 0 to 1, where a score of 1 equates to perfect "universality" (i.e., all hosts have a correlation coefficient of 1 or all hosts have a correlation coefficient of -1). Applying this score to all pairs of ASVs reveals a right-skewed distribution, reflecting the fact that most bacterial correlations are weak, with inconsistent sign directions across hosts (**Fig. 3C**; **Fig. S4B**). However, 49% of these scores were higher than expected by chance (permutation test; FDR ≤ 0.05; **Fig. 3C**; **Fig. S4B**), reflecting a signal of universality in our data.

Interestingly, the ASV-pairs with the highest universality scores almost always exhibited net positive correlations across hosts, as opposed to net negative relationships, suggesting that the most universal relationships occur between pairs of ASVs that respond similarly to shared drivers or facilitate each other's growth. For example, among the ASV pairs in the top 1% of universality scores (n=78 pairs), 96.2% exhibited net positive correlations across all hosts, while only 5.6% (3 of 78 pairs) exhibited net negative correlations (**Table S4**). In the top 2.5% most universal ASV pairs (n=194), 78.4% had net positive correlations across all hosts (**Table S4**).

To visualize these highly consistent positive correlations, we plotted bacterial co-abundance networks connecting the top 2.5% most universal ASV pairs (**Fig. 3C**). A handful of ASVs were highly connected within this network. The most connected ASV was ASV107 (family Lachnospiraceae; **Table S1**; **Table S4**), which was connected to 20 other ASVs. Ten other ASVs were connected to more than 10 other ASVs, including six other members of Lachnospiraceae (ASV9, ASV25, ASV30, ASV106, ASV107, and ASV111), two members of Coriobacteria (ASV115 in the family Coriobacteriaceae and ASV30 in the genus *Slackia*), one member of Bifidobacteriaceae (ASV50), and one member of Prevotellaceae (ASV71). The ASVs involved in these top 2.5% pairs were enriched for the families Atopobiaceae, Bifidobacteriaceae, Coriobacteriaceae, Eggerthellaceae, Erysipelotrichaceae, and Lachnospiraceae (**Fig. 3E; Table S5**; all Fisher's Exact Test p-values < 0.01; FDR ≤ 0.05).

The network in **Fig. 3D** revealed clusters of positive connections, often between ASVs
assigned to the same family (**Fig. 3F**). In fact, same-family pairs were enriched by >3-fold in the
2.5% most universal taxon pairs (52 pairs observed vs. 19 expected, p < 0.0001). The cluster of
interconnected red nodes in **Fig. 3D** represents members of Lachnospiraceae, and
Lachnospiraceae-Lachnospiraceae pairings were 3.7 times more common in this network than
overall (30 pairs observed vs. 9 pairs expected **Fig. 3F**). Bifidobacteriaceae also tended to exhibit
within-family ASV pairings (**Fig. 3F**).
The observation that the most consistent correlations often occur among ASVs in the
same family raises another question: does the phylogenetic distance between a pair predict the
nature of their relationship? In support of the idea that closely related ASVs respond similarly to
the environment or facilitate each other's growth, we found a significant relationship between the
universality score of a given pair of ASVs and their phylogenetic distance (Pearson's r for
positively correlated pairs=-0.213; p < 0.0001; **Fig. 4A**). In contrast, negatively correlated ASV
pairs exhibited a weak positive relationship between phylogenetic distance and universality such
that closely related taxa tended to be less universal than more distantly related taxa (Pearson's
r=0.049; p=0.004; **Fig. 4B**). In other words, the strongest and most consistently *negatively*
correlated taxa tend to be only distantly related. Positively correlated, closely related pairs were
often members of the families Atopobiaceae, Eggerthellaceae, and Lachnospiraceae, especially
pairs where both members belonged to the family Lachnospiraceae (**Fig. 4C-D**; **Table S6).**
364

365
Figure 4. The most consistent ASV-level correlations are positive and often between close
evolutionary relatives. Pairwise universality scores are plotted as a function of phylogenetic
distance between the ASV-ASV pair for consensus positively correlated pairs in red (**A**) and
negatively correlated pairs in blue (**B**). Phylogenetic distance (x-axis) is binned into 0.1
increments; each point represents a given ASV pair, and box plots represent the median and

interquartile ranges for a given interval of phylogenetic distance. Phylogenetic distance is
negatively correlated with universality score in positive pairs (Pearson's correlation for
positively associated ASV pairs=-0.213, p-value < 0.0001), and positively correlated with
universality score in negatively associated pairs (Pearson's correlation for negatively associated
ASV pairs=0.049, p = 0.004). Panel **(C)** shows families for the ASV pairs enriched in the closest
related (distance < 0.2) and highly universal (score > 0.5) pairs. Panel **(D)** shows enriched
family-family pairings for the same subset of closely related and highly universal ASV pairs in
panel C. Note that for visualization, the estimated log2 odds ratio intervals have been truncated
at 5, which excludes 5 pairs with high uncertainty in the odds ratio; full estimates are given in
Table S6.
381
382
Genetic relatives, and hosts with similar microbiome compositions, have more similar
bacterial correlation patterns
We next asked whether host-level variables, including sex, social group membership,
genetic relationships, and baseline gut microbiome composition predict host differences in
patterns of bacterial correlation. Consistent with prior research [10], the strongest predictor of
distance in bacterial correlation patterns was distance in terms of baseline microbiome
composition. Indeed, a Mantel test correlating compositional distance of average microbial
profiles (as Aitchison distances between the per-host mean of centered log-ratio-transformed
samples) with distance in microbial correlation patterns between hosts (via Frobenius distance)
revealed that 34% of the variation in correlation patterns was explained by baseline microbiome
community composition (Mantel: r^2=0.343; p=0.001; **Fig. 5A**; **Table S7**).
Consistent with prior research in our population, which finds widespread heritability of
the abundance of individual gut microbiome taxa [43], we also found a weak but significant
relationship between host genetic distance and the distance in microbial correlation patterns
between hosts. Hosts who were more distantly related based on a multigenerational pedigree
have slightly less similar ASV-level correlation matrices, as measured by Frobenius distance
(**Fig. 5B**; **Table S7**; r^2=0.025; Mantel p-value=0.001). We found no evidence that members of
the same social group or sex exhibit especially similar microbial correlation patterns (social
group: F=1.994; p=0.106; sex: F=1.784; p=0.187; **Table S7**).

Figure 5. Baboons with more similar bacterial correlation patterns are more likely to have more similar baseline microbiome compositions and are more likely to be genetic relatives. In panel **(A)** each point is a pair of hosts; the y-axis shows the similarity of these hosts' bacterial correlation patterns (via Frobenius distance) as a function of their microbiome compositional similarity (via Aitchison distance; Mantel: r^2=0.343; p=0.001). Colors show samples from pairs of baboons living in the same social group and grey dots are pairs of animals living in different social groups; there is no detectable effect of social group on correlation pattern similarity. Panel **(B)** shows the same Frobenius distances as a function of host genetic dissimilarity (1 – the coefficient of genetic relatedness between hosts; R^2=0.025; p-value Mantel test 0.001). Colors reflect pairs of hosts living in the same social group, as in panel A.

Universality in Amboseli is not solely explained by seasonality or synchrony

Without experiments, we cannot disentangle whether our observed bacterial correlations are due to ecological interactions between bacterial species (e.g., mutualisms, direct or indirect competition etc.) or to shared responses to environmental gradients. While our modeling approach accounts for seasonal changes in the first three principal components of the baboons' diets, to identify other potential effects of season we re-estimated the ASV-ASV correlation matrix after removing an oscillating seasonal trend from the observed log-ratio abundance for each ASV (**Fig. S8**). Removing this trend had little effect on the ASV-ASV correlation matrix; the variance explained by the seasonal oscillation is small for all ASVs (median 1.1%, minimum=0%, maximum=6%) and the between-ASV correlation estimates were almost identical to those derived from our original model (Pearson's r=0.979, p<0.0001; **Fig. S8C**). We also tested whether pairs of ASVs with especially consistent between-host correlation patterns tend to show large seasonal changes in CLR abundance. To do so, we focused on 13 families that exhibit significant seasonal changes in CLR abundance, based on a previous analysis of the same data set [42]. While ASV pairs in which one member belongs to one of these significantly "seasonal" families are slightly more universal, this effect is small (difference of 0.026, p<0.0001 vs. pairs where 0 or 1 partner were "seasonal"; **Fig. S9**).

Because the high level of universality we observed was not well explained by season, we also tested whether universality is explained by synchronized dynamics. We reasoned that if one member of an ASV pair shows highly synchronized dynamics across different hosts, and the other member is also strongly synchronized across hosts, then universality could be an inevitable outcome of each member of the pair's strong synchrony. We quantified synchrony as the degree to which the observed dynamics of a single, focal ASV are consistent across hosts, such that high synchrony (near 1) implies that the timing and direction of shifts in log-ratio ASV abundance are identical across hosts in the population (see Methods; **Fig. S10**). Estimates of synchrony ranged from 0.019 to 0.477 (median=0.196). Interestingly, ASVs in the 13 "seasonal" families are not more likely to have high synchrony than other families (ANOVA, p=0.358; **Fig. S11**). However, the average synchrony of an ASV-ASV pair did predict that pair's universality score (r=0.264, p<0.0001): ASV pairs that are more synchronous on average are also more likely to show consistent correlations across hosts. These high synchrony, high universality pairs are highly enriched for Bifidobacteriaceae-Bifidobacteriaceae and Lachnospiraceae-Lachnospiraceae family pairs (**Fig. S12**).

Baboon microbiomes are not substantially more "universal" than human microbiomes

Finally, to investigate the generalizability and applicability of our observations in baboons, we turned to two publicly available gut microbial time-series data sets: daily samples from 34 adults over a 17-day span (483 total samples; hereafter "Johnson et al." [34]), and the DIABIMMUNE cohort that consists of 285 samples, collected monthly over 3 years, from 15 infants and toddlers living in Russian Karelia ([44]; at the time of writing, these cohorts were the only publicly available data sets we could find that included large numbers of repeated samples from the same subjects). Because baboons in Amboseli experience less heterogeneity in their environments and diets than humans [42, 43], we expected they would exhibit greater consistency in microbial correlations than either human cohort. Note that we compared each host cohort's universality at the family/order/class level because this taxonomic level offered the greatest comparative power (10.1% of families/orders/classes overlap between the cohorts compared to just 3.1% of genera and no ASVs).

Contrary to our expectations, we find comparable evidence of universality in baboons and the DIABIMMUNE infant/toddler cohort, but weak universality in Johnson et al. (**Figs. 6A-6D**). Bacterial families in the DIABIMMUNE cohort yielded universality scores slightly higher than those observed in Amboseli (25th percentile=0.132, median=0.206, 75th percentile=0.316 for DIABIMMUNE; 25th percentile=0.088, median=0.150, 75th percentile=0.234 for Amboseli), driven by relationships between families that were stronger on average than those estimated for baboons (median DIABIMMUNE family-family correlation strength=0.270; median Amboseli family-family correlation strength=0.181). The high level of consistency between both infants/toddlers and baboons in one wild population is surprising and may point to the similar sampling intervals for these cohorts. Both cohorts were sampled approximately monthly, while Johnson et al.'s subjects were sampled daily [17, 48]. Median correlation strengths and universality scores for the Johnson et al. [34] cohort were substantially lower (median correlation=0.099; 25th percentile universality=0.050, median=0.076, 75th percentile=0.111) than the DIABIMMUNE cohort or the baboons.

Despite considerable differences in the hosts, time scales, and designs of these studies, all three data sets exhibited a positive correlation between correlation strength and sign consistency for family pairs (**Fig. 6C**). This correlation was strongest in the Amboseli baboons (Spearman's

r=0.844; p<0.0001); weaker in the DIABIMMUNE cohort (r=0.686; p<0.0001) and weakest in
Johnson et al. [34] (r=0.644; p<0.0001). Further, the observation that the most universal family-
family associations skew positive in baboons was replicated in the infant data set, but not in
Johnson et al. [34]. All of the top 1% and top 2.5% most universal family pairs (6 of 6 and 16 of
16 pairs, respectively) are positively associated in the DIABIMMUNE cohort, compared to 86%
and 71% of these pairs in the Amboseli baboons.
Finally, we examined the relationship between universality scores for family pairs that
overlapped between Amboseli and DIABIMMUNE (n=45 pairs), and between Amboseli and
Johnson et al. [34] (**Fig. 6D**; n=21 pairs; only 10 family pairs overlapped between all three data
sets). For these overlapping pairs, scores in the Amboseli data predicted scores for the same
family-family pair in the DIABIMMUNE data set (r=0.449, p=0.023). The association between
scores in the Amboseli data and the Johnson et al. data was negative, but not statistically
significant (r=-0.222, p=0.071).

492
493
Figure 6. Patterns of universality in baboons are recapitulated in the DIABIMMUNE
study. Following **Fig. 2A**, Panels **(A)**, **(B)**, and **(C)** show the Pearson's correlation coefficients
of CLR abundances between all pairs of families (x-axis) in two time series data sets from
human subjects: **(A)** the Amboseli baboons, **(B)** the DIABIMMUNE cohort, consisting of 15
infants/toddlers sampled monthly over 3 years in Russian Karelia [44], and **(C)** the diet study of
Johnson et al. [34], including 34 adults sampled daily over 17 days. Following **Figs. 3A and B**,
panel **(D)** shows the median correlation strength of each family pair's correlation coefficient
across hosts as a function of the consistency in direction of that pair's correlation across hosts
(i.e., the proportion of hosts that shared the majority correlation sign, positive or negative).
Median correlation strength is low overall in Johnson et al. (median=0.099), whereas the

Amboseli baboon and DIABIMMUNE infant/toddler cohorts show similar relationships between
median correlation strength and the proportion shared correlation sign across hosts (Spearman's r
in Amboseli=0.844; Spearman's r in DIABIMMUNE=0.686). **(E)** Universality scores for
overlapping family pairs from the infant/toddler subjects of the DIABIMMUNE study and
baboons in the Amboseli study are significantly correlated (r=0.449, p=0.0226). Panel D shows
universality scores for overlapping gut bacterial family pairs in the Amboseli baboon and
DIABIMMUNE infant/toddler data sets (black outlines), as well as the Amboseli and Johnson et
al. data sets (gray outlines) on opposing axes. Color represents the taxonomic identities of the
family pairs.
513
514
Discussion
Do different hosts have different microbiome "ecologies"? Answering this question is
essential for predicting gut microbiome community assembly and dynamics, and for
understanding the degree to which the species interactions that govern these processes are shared
across hosts. Here, we overcome the constraints of previous cross-sectional analyses by
measuring bacterial correlations directly from longitudinal, multi-host microbiome time series.
Our results provide independent confirmation for prior studies that tested for universal gut
microbial relationships via dissimilarity overlap analyses (DOA; [10, 18-20, 36]). We confirm
that bacterial correlation patterns are largely shared across hosts in the same population, as
opposed to idiosyncratic to individual hosts, and that hosts with the most similar bacterial
correlation patterns are those with the most similar baseline microbiome compositions—a core
assumption of DOA. Because prior analyses of these microbiome time series find that each
baboon exhibits a highly personalized microbiome composition and dynamics [42], our findings
suggest that such compositional personalization, which is also common in humans [22-26, 31-
35], cannot be easily explained by personalized microbiome ecologies. Further, in terms of
microbiome therapeutics, our results suggest that widely applicable microbiome interventions
may be more attainable than personalized microbiome compositions would suggest.
By measuring bacterial correlations in multiple hosts, we were also able, for the first
time, to pinpoint which pairs of bacterial taxa exhibit the most consistent relationships across
hosts. Surprisingly, we found that the most universal bacterial pairs are almost always positively
(as opposed to negatively) correlated. Positive bacterial interactions have been the subject of
recent controversy [9, 15, 49]. Ecological theory predicts that strong positive interactions should
be rare in natural communities because species interdependencies can hamper community
assembly and stability [14, 17]. This theory is supported by experiments that directly measure the
effects of one bacterial species on another's growth [50-53] (but see [49]). Our results suggest
that positive bacterial correlations are indeed uncommon in intact, unmanipulated microbiomes:
significant positive relationships made up just 8.8% of all of the pairwise correlations we
observed. However, when they occur, they often contain taxa that belong to the same bacterial
families or are otherwise phylogenetically close, suggesting they may be members of the same
ecological guild and respond similarly to shared resources and other environmental drivers. This
pattern may partly explain the abundance of positively associated Lachnospiraceae pairs in our
data, a family in which positive, within-family interactions are known to contribute to
hydrolyzing starch and other complex carbohydrates, and ultimately the regulation of short chain
fatty acids (SCFAs) [54-56].

These observations—that bacterial correlation patterns are largely consistent across hosts,
and that the most consistent correlations are typically positive—were also apparent in one human
data set, despite differences in study design, host age, and time scale. Specifically, both the
Amboseli baboons and the DIABIMMUNE infant/toddler cohort from Russia [44] exhibit
comparable levels of universality. This outcome was surprising, given that baboons are expected
to experience less heterogeneity in their environments and diets than human children from birth
to age three years—even if those infants are from the same population (Russian Karelia). We
also found that the most universal bacterial families in baboons tended to be highly universal in
human infants/toddlers. Hence, some bacterial families may exhibit consistent microbial
relationships within hosts, across host populations, and across host species. Finally, a recent,
independent study also identified consistent bacterial correlation patterns across four different
populations of human hosts [9]. While this study lacked resolution at the level of individual
hosts, it did identify a highly conserved network of positively associated and closely related
microbes similar to those we identify in **Fig. 3**. The authors speculate that these conserved
associations may indicate strong partner fidelity or obligate partnerships.
We did, however, fail to detect universality in a second human data set reported in
Johnson et al. [34], in which subjects were sampled daily, rather than weekly or monthly. The
lack of universality in Johnson et al. [34] may be due to this difference in sampling time scale,
especially if daily abundances and correlations are noisier than covariances modeled over the
longer time scales in our study. In support, many fewer of the microbial correlations were
stronger than random chance in Johnson et al. as compared to the baboons or children in the
DIABIMMUNE cohort. The subjects in Johnson et al. [34] also consumed substantially different
diets from each other, perhaps more so than the children in the DIABIMMUNE cohort, and this
inter-host difference in diet may reduce the universality of microbial correlations.
In terms of understanding microbiome ecology, an essential caveat to our findings is that
the correlation patterns we observed could reflect either direct or indirect relationships, or
uncontrolled environmental gradients, and they cannot be mapped directly to standard categories
of pairwise ecological interactions, such as mutualism, commensalism, amensalism, exploitation,
or competition. Experimental approaches that directly measure the effects of one species on
another's growth *in vitro* are better suited to characterizing these relationships [49-53]. However,
even then, caution is required because a microbe's community and environmental context can
have important consequences for its metabolism, functional capacities, and relationships with
other microbes. We surmise that most of the correlation patterns we observed are not attributable
to environmental gradients because our signature of universality persisted, even when we
accounted for diet, oscillating seasonal drivers, and microbial synchrony between hosts. Hence,
some of correlations we observed may derive from microbial interactions themselves, rather than
shared environmental drivers creating shared dynamics.
Our finding that correlations between gut microbial taxa are largely consistent across
hosts is important, considering that many studies find highly personalized gut microbial
compositions and single-taxon dynamics [27-29]. Personalized compositions and dynamics in
the gut microbiome are commonly attributed to horizontal gene transfer and functional
redundancy, which may lead some bacteria to perform different functions and exhibit different
environmental responses in different hosts. Our results suggest these processes do not
substantially alter pairwise microbial associations in the gut, at least for highly prevalent taxa at
the level of ASVs and above, and on the time scales in our study (on the order of weeks and
months). Because ASVs encompass multiple species and strains, reflecting the functional

diversity of these taxa, their dynamics may be somewhat buffered against idiosyncrasies driven
by horizontal gene transfer and functional redundancy, which affect single strains more strongly
than whole species or genera. If so, personalized gut microbial compositions may emerge instead
from personalized assembly processes [57, 58], the fact that most microbial relationships are
weak, and the effects of rare, host-specific taxa (which were necessarily excluded from our
analyses). A logical next step would be to confirm the stability of the microbial correlations we
observed using culture-based approaches, which will help reveal the stability of these
correlations in vitro and whether they can be attributed to direct effects of one microbe on
another's growth.
604
Methods
Study population and microbiome profiles
The baboon hosts in this study were members of the Amboseli baboon population, which
has been studied by the Amboseli Baboon Research Project since 1971 [41]. The microbiome
compositional profiles are derived from V4 16S rRNA gene amplicon sequencing data that were
previously analyzed in [42, 43]. Our analyses use 5,534 of these profiles from 56 especially well-
sampled baboons, collected over a 13.3-year span between 2000 to 2013 (**Fig. 1B**). Each baboon
host in this data set was sampled at least 75 times (mean number of samples=99; range=75 to
181 samples; median number of days between samples within hosts=20 days; 25th percentile=7
days, 75th percentile =49 days). DNA was extracted from each sample using the MoBio and
QIAGEN PowerSoil kit with a bead-beating step. All samples were sequenced on an Illumina
HiSeq 2500, with a median read count of 48,827 reads per sample across all 5,534 samples
(range=982 to 459,315 reads per sample). Further details of sample collection, DNA extraction,
and sequencing can be found in [42, 43].
619
Filtering of low-abundance taxa
Data sets of per-sample taxonomic counts were produced at each of three taxonomic
levels, from finest to coarsest: ASV, taxonomic assignments finer than phyla, but above the
genus level (e.g., class, order, family), and phylum. At the intermediate and coarsest levels, taxa
were agglomerated using phyloseq's tax_glom() function [59] such that all sequence variants
sharing taxonomic identity at that level were collapsed into a single taxon (e.g. family
Bifidobacteraceae).
To reduce sparsity in the data set, remove 16S sequences that could represent gene
duplications, and focus only on taxa that were prevalent in all 56 hosts, we further filtered as
follows: (1) in each of the three taxonomically defined data sets (i.e. ASV, taxa assigned to
family/order/class, and phylum), we identified taxa present in a minimum of 20% of each host's
samples; (2) if a given ASV was >99% genetically similar to another ASV we removed the least
abundant of the pair to minimize the risk of including duplicate 16S rRNA gene copies from the
same taxa [47]; and (3) counts associated with all other taxa were combined into a dummy
category, hereafter referred to as "other." The "other" category therefore includes a combination
of rare and host-specific gut microbes. This category was retained in the data set (although not
analyzed directly) because "other" counts still inform the precision of the observed relative
abundances in our model. Characteristics of the filtered data at each taxonomic level are
provided in **Tables S1-S3**.
639
640

Modeling log-ratio dynamics

Estimates of taxon-taxon covariance were obtained from the *basset* model of the "fido" package in R [46]. Data for each host took the form of a $D \times N$ count matrix, where D gives the number of taxa and N the number of samples collected for a given host. The following model was fit to each host's count matrix (Y) where Y_i represents the counts associated with a single sample:

$$Y_i \sim \mathrm{Multinomial}(\pi_i)$$
$$\pi_i = \mathrm{ALR}^{-1}(\eta_i)$$
$$\eta \sim \mathrm{Normal}(\Lambda, \Sigma, I)$$
$$\Lambda \sim \mathrm{GP}(\Theta[X], \Sigma, \Gamma[X])$$
$$\Sigma \sim \mathrm{inv\text{-}Wishart}(\Xi, \nu)$$

The observed relative abundances are considered to be drawn from a multinomial distribution parameterized by a set of proportions (π) which have an analogous representation in the additive log-ratio. The dynamics of these log-ratio abundances (η) are described by what amounts to a state space model in the 3$^{\mathrm{rd}}$ and 4$^{\mathrm{th}}$ lines of the specification above, where a Gaussian process models the evolution of a "latent" state. The matrix Σ captures covariation in log-ratio abundances (the D rows of the observed count matrix). Sample-sample covariation arising from nearness in time (autocorrelation) is modeled by the kernel matrix Γ. Both the kernel matrix and the expected baseline log-ratio abundances (Θ) are parameterized by a set of time-varying covariates X which included the day of sampling (where the date of first sample is defined as zero) and the first three principal components of diet composition, calculated following [42, 43] as the diet all juveniles and females living in the host's social group in the 30 days prior to sample collection. All group members consume highly similar diets as they travel in a together across the habitat, encountering the same resources at the same time [42, 43]. These data are collected via random-order behavioral observations collected two to four times per week on adult females and juveniles in each social group. Parameterization of the kernel matrix is further described in the Supplement.

Posterior inference on this model is performed as described in [46] and yields estimates of the distributions of parameters necessary to reconstruct trajectories for all log-ratio taxa across sampling time. In particular, we extract the posterior estimates of one such parameter, Σ, the covariance of additive log-ratio (ALR) taxa, from the fitted models for each host. We convert these covariance matrices over ALR taxa to the centered log-ratio (CLR) form (a simple linear transformation of the matrix). We then normalize estimated CLR covariance matrices to Pearson correlation matrices in R using the built-in cov2cor() function.

Calculating universality scores for taxon-taxon pairs

We devised a universality score for each pair of taxa intended to capture the strength and consistency of taxon-taxon correlations across hosts. The majority direction is negative otherwise. This score identifies the sign of the taxon-taxon correlation (positive or negative) that is most common across the 56 hosts (i.e., occurs in >50% of the 56 hosts in the data set). The direction of this sign is the "majority correlation sign."

For a pair of taxa i, let n_i^{maj} be the number of hosts with CLR correlation over pair i with the majority correlation sign for that pair and let n be the total number of hosts. Let R be the

681 subset of estimated CLR correlations for pair i across hosts with the majority sign. The
682 universality score u_i for that taxon-taxon pair is then given by
683

$$u_i = \frac{n_i^{\text{maj}}}{n} \times \text{median}(R)$$

684
685   This score is the product of the median CLR correlation across hosts and the proportion
686 of hosts with the majority correlation sign, and is bounded between 0 and 1. Scores near 1
687 indicate strong universality and near-zero scores indicate weak universality. Strong universality
688 can only be achieved by taxon-taxon correlations that are both large in magnitude and highly
689 concordant across hosts.
690
Defining a cutoff for significant bacterial correlations and universality scores
We identified correlations stronger than expected under random simulations using
permutations of the data set to define empirical null distributions (**Fig. S4A**). Specifically, we
permuted the data by randomly shuffling taxon identity within each sample 10 times for each of
the 56 hosts. This procedure maintained relative abundance patterns within a sample but
scrambled the covariance patterns of relative abundances. The distributions of ASV-level CLR
correlations in the original and permuted data are shown in **Fig. S4A**. We identified "significant"
correlations as those below FDR ≤ 0.05 (Benjamini-Hochberg), testing against the permuted
data.
We applied an analogous permutation test to derive a null distribution for taxon-taxon
universality scores. In a single iteration of this permutation procedure, rows and columns of the
observed taxon-taxon correlation matrix for each host were shuffled, maintaining the distribution
over observed correlations at the host level but randomizing the identity of taxon pairs across
hosts. This procedure was repeated 100 times and universality scores were calculated from each
of these shuffled data sets to give a pseudo-null distribution of universality scores. The observed
and null distributions of universality scores at the ASV level are shown in **Fig. S4B**. We used
this empirical null distribution to identify universality scores significantly greater than expected
(FDR ≤ 0.05).
709
Estimating the ratio of population-level to host-level contributions to observed taxon-taxon
correlation patterns
We used simulations to estimate the degree of shared "signal" between hosts in terms of
taxon-taxon correlations. Each host's "observed correlations" were defined as the *basset*
estimated maximum a posteriori (MAP) estimates of centered log-ratio ASV correlations for that
host. We computed the *mean* correlations across the population using the function estcov() from
the shapes package in R [60] and estimated a host-specific contribution to the observed
correlations as the residual *difference* between per-host observed and these mean correlations.
That is,
719

observed host correlations = mean population correlations + host residual

720
721   For each host, we simulated a hypothetical set of composite taxon-taxon correlations as a
722 convex combination of mean and host residual:
723

$$\text{composite correlations} = (1-\alpha) \times \text{mean population correlations} + \alpha \times \text{host residual}$$

A cartoon example of this procedure is given in **Fig. S7A**. For example, one such
simulated set of taxon-taxon correlations might constitute a mixture of 90% host contribution
and 10% shared population-level "signal" (α=0.9). Alternatively, a small host-level contribution
might have α=0.1.
For each host, we iterated over increasing proportions of host-level contribution (from
0% to 100%), generating simulated composite correlation matrices according to the formula
above. We compared these simulated patterns to those observed for the same host, reasoning that
simulated correlation matrices that minimize the distance between the observed correlation
matrices and the simulated mixtures provide the best description of the underlying true mixture.
734

Estimating synchrony
Seasonal autoregressive models were fit independently to each CLR-transformed ASV
with arima() in R, using covariate matrices which included per-host intercepts and an oscillating
periodic trend to mimic wet-dry season oscillation. For each ASV, all hosts' samples were
combined into a single series, yielding per-ASV models of CLR dynamics. This procedure is
detailed in the Supplemental Methods. Residuals were extracted from these fitted models as
seasonally "de-trended" data and CLR correlation matrices across ASV pairs were estimated
directly from these adjusted data using cov() in R (**Fig. S9**).
"Synchrony" was estimated by sampling aligned microbiome compositional profiles
across hosts. We identified all samples collected from pairs of hosts within 1 calendar day. For
instance, a sample collected from host F01 on 2011-03-14 could pair with a sample from M04 on
2011-03-15. For all possible pairs of hosts, we selected one such aligned pair of samples,
yielding 1540 joint observations of gut microbiome composition. For each such paired sample,
one host was arbitrarily designated as host A and the other as host B. The "synchrony" of a given
taxon was estimated as the correlation of a taxon's model-inferred log-ratio abundance across the
set of samples from hosts labeled A and the set of samples from hosts labeled B. The cartoon in
Fig. S10 illustrates this sample pairing.
752

Enrichment analyses
We performed enrichment analyses for bacterial families and family pairs in several
settings. In each case we computed the frequency of ASVs belonging to a given family, or of
pairs belonging to a family pair, on a subset of the data. These were compared to the overall
frequencies of ASVs belonging to those families or pairs.
To determine the enrichment of families and family pairs in the most universal ASV pairs
(**Fig. 3E and 3F**), we calculated the frequencies of ASV families and pairs in the top 2.5% of
pairs by universality scores. Significant enrichment of families or pairs was identified using a
one-sided Fisher's Exact Test. Multiple test correction was applied as a Benjamini-Hochberg
adjustment to observed p-values.
Phylogenetic distances between ASV sequences were calculated with the dist.ml function
in the "phangorn" package in R [61] using default settings for amino acid substitution rates. In
Fig. 4C and 4D, low phylogenetic distance/high median correlation strength pairs were
identified as those with phylogenetic distances of less than 0.2 and median correlation strengths
of greater than 0.5. Again, significance of these was evaluated against overall frequencies of the
same families and pairs.

To determine enrichment of low synchrony/high universality or of high synchrony/high
universality families and pairs (shown in **Fig. S12A and 12B**), we defined the low
synchrony/high universality cohort as those ASV pairs with synchrony estimates of less than 0.3
and universality estimates greater than 0.4. We defined the high synchrony/high universality
cohort as those ASV pairs with synchrony greater than 0.3 and universality greater than 0.4. The
frequency of these subsets was evaluated against the overall frequencies of the same families and
pairs.
776
Evaluating explanatory factors
Variation in taxon-taxon correlation patterns explained by kinship and baseline
composition. To evaluate a possible explanatory effect of distances in terms of kinship or
baseline gut bacterial composition on distances in terms of taxon-taxon correlation patterns, we
applied Mantel tests. However, because population structure can lead to anticonservative p-
values [62], we also developed a second simulation-based procedure for evaluating the
significance of baseline composition, using a permutation procedure of our own design. Firstly,
baseline composition for each host was estimated by transforming all of a given host's samples
to the centered log-ratio representation after adding a small fraction (0.5) to remove zeros. The
vector of per-taxon averages of these CLR values was used as that host's "baseline" CLR
composition. The Euclidean distances between hosts in terms of these per-host baselines were
compared against distances in terms of correlation patterns to give an r^2 value.
In the case of the customized permutation test, this observed result was evaluated against
a pseudo-null distribution computed in the following way. The identity of each taxon in the
baseline composition was shuffled for each host independently. Euclidean distances across these
shuffled baselines were computed and an r^2 value calculated for these distances against the
observed distances computed from taxon-taxon correlation patterns. This procedure was repeated
1000 times to give a distribution of "random" r^2 values we used as an empirical null.
Variation in taxon-taxon correlation patterns explained by sex and social group. To test
whether host sex or social group membership predicted similarity in terms of correlation
patterns, we used an ANOVA-like strategy. We calculated the F-statistic, a ratio of between- to
within-group variation, on the observed correlation patterns (strictly, the vectorized CLR taxon-
taxon correlation matrices; Z in the equation below) and segmented samples into groups defined
by either sex or social group. The F-statistic was calculated as
801

$$F = \frac{\text{between-group variation}}{\text{within-group variation}} = \frac{\sum_{i=1}^{K} n_i (\bar{Z}_i - \bar{Z})^2 / K - 1}{\sum_{i=1}^{K} \sum_{j=1}^{n_i} (Z_{ij} - \bar{Z}_i)^2 / (N - K)}$$

802
803   and significance was evaluated via an F-distribution parameterized by the appropriate degrees of
804   freedom. Here K represents the number of groups (e.g. two, in the case of sex) and N, the total
805   number of hosts. The matrix $\bar{Z}_i$ consists of the mean taxon-taxon correlations for group i and $\bar{Z}$,
806   the population mean correlations.
807
Comparison to microbiome time series from human populations
We compared our findings to those generated from two human data sets: the
DIABIMMUNE project's infant/toddler cohort from Russian Karelia [44] and the adult diet-
microbiome association study of Johnson et al. [34]. In both cases, count tables were obtained
from the project's public website and subject identity and sampling schedules were available in

the associated metadata. We compared each host cohort's universality at the family/order/class
level because this taxonomic level offered the greatest comparative power (10.1% of
families/orders/classes overlap between the cohorts compared to just 3.1% of genera and no
ASVs). The *basset* model from the "fido" R package [46] was fit to each subject's data set using
model settings analogous to those employed on the Amboseli baboon series: first, only taxa with
non-zero counts in at least 20% of all subjects' series were retained; second, Gaussian process
kernel bandwidth settings were chosen in such a way as to encode an expectation of minimum
autocorrelation between samples at a distance in time of 90 days. We extracted centered log-ratio
estimates of taxa at the family level in the same manner as described previously for the Amboseli
data set.
823
824 **Data and code availability**
825 16S rRNA gene sequences are available on EBI-ENA (project 590 ERP119849) and Qiita (study
826 12949). Analyzed data and code is available on GitHub at:
827 https://github.com/kimberlyroche/rulesoflife
828
References

1. Faust K, Raes J. Microbial interactions: from networks to models. Nature Reviews Microbiology.
2012;10(8):538-50. doi: 10.1038/nrmicro2832. PubMed PMID: 22796884.
2. Foster KR, Bell T. Competition, Not Cooperation, Dominates Interactions among Culturable
Microbial Species. Current Biology. 2012;22(19):1845-50. doi: 10.1016/j.cub.2012.08.005. PubMed
PMID: WOS:000309792900031.
3. Dolinsek J, Goldschmidt F, Johnson DR. Synthetic microbial ecology and the dynamic interplay
between microbial genotypes. Fems Microbiology Reviews. 2016;40(6):961-79. doi:
10.1093/femsre/fuw024. PubMed PMID: WOS:000387995000010.
4. Seth EC, Taga ME. Nutrient cross-feeding in the microbial world. Front Microbiol. 2014;5:350. Epub
20140708. doi: 10.3389/fmicb.2014.00350. PubMed PMID: 25071756; PubMed Central PMCID:
PMCPMC4086397.
5. Backhed F, Ley RE, Sonnenburg JL, Peterson DA, Gordon JI. Host-bacterial mutualism in the human
intestine. Science. 2005;307(5717):1915-20. doi: Doi 10.1126/Science.1104816. PubMed PMID:
ISI:000227957300045.
6. Gould AL, Zhang V, Lamberti L, Jones EW, Obadia B, Korasidis N, et al. Microbiome interactions
shape host fitness. Proc Natl Acad Sci U S A. 2018. doi: 10.1073/pnas.1809349115. PubMed PMID:
30510004.
7. Pontrelli S, Szabo R, Pollak S, Schwartzman J, Ledezma-Tejeida D, Cordero OX, et al. Metabolic
cross-feeding structures the assembly of polysaccharide degrading communities. Sci Adv.
2022;8(8):eabk3076. Epub 20220223. doi: 10.1126/sciadv.abk3076. PubMed PMID: 35196097;
PubMed Central PMCID: PMCPMC8865766.
8. Degnan PH, Taga ME, Goodman AL. Vitamin B12 as a modulator of gut microbial ecology. Cell
Metab. 2014;20(5):769-78. doi: 10.1016/j.cmet.2014.10.002. PubMed PMID: 25440056; PubMed
Central PMCID: PMCPMC4260394.
9. Loftus M, Hassouneh SA, Yooseph S. Bacterial associations in the healthy human gut microbiome
across populations. Sci Rep. 2021;11(1):2828. Epub 20210202. doi: 10.1038/s41598-021-82449-0.
PubMed PMID: 33531651; PubMed Central PMCID: PMCPMC7854710.
10. Bashan A, Gibson TE, Friedman J, Carey VJ, Weiss ST, Hohmann EL, et al. Universality of human
microbial dynamics. Nature. 2016;534(7606):259-62. doi: 10.1038/nature18301. PubMed PMID:
27279224; PubMed Central PMCID: PMCPMC4902290.

11. Widder S, Allen RJ, Pfeiffer T, Curtis TP, Wiuf C, Sloan WT, et al. Challenges in microbial ecology:
building predictive understanding of community function and dynamics. ISME Journal.
2016;10(11):2557-68. doi: 10.1038/ismej.2016.45. PubMed PMID: 27022995.

12. Cao HT, Gibson TE, Bashan A, Liu YY. Inferring human microbial dynamics from temporal
metagenomics data: Pitfalls and lessons. Bioessays. 2017;39(2):1600188. doi:
10.1002/bies.201600188. PubMed PMID: 28000336.

13. Faust K, Raes J. Host-microbe interaction: Rules of the game for microbiota. Nature.
2016;534(7606):182-3. doi: 10.1038/534182a. PubMed PMID: 27279206.

14. Coyte KZ, Schluter J, Foster KR. The ecology of the microbiome: Networks, competition, and
stability. Science. 2015;350(6261):663-6. doi: 10.1126/science.aad2602. PubMed PMID: 26542567.

15. Palmer JD, Foster KR. Bacterial species rarely work together. Science. 2022;376(6593):581-2. Epub
20220505. doi: 10.1126/science.abn5093. PubMed PMID: 35511986.

16. Wu G, Zhao N, Zhang C, Lam YY, Zhao L. Guild-based analysis for understanding gut microbiome
in human health and diseases. Genome Med. 2021;13(1):22. Epub 2021/02/11. doi: 10.1186/s13073-
021-00840-y. PubMed PMID: 33563315; PubMed Central PMCID: PMCPMC7874449.

17. Coyte KZ, Rao C, Rakoff-Nahoum S, Foster KR. Ecological rules for the assembly of microbiome
communities. PLoS Biol. 2021;19(2):e3001116. Epub 20210219. doi: 10.1371/journal.pbio.3001116.
PubMed PMID: 33606675; PubMed Central PMCID: PMCPMC7946185.

18. Gao C, Montoya L, Xu L, Madera M, Hollingsworth J, Purdom E, et al. Fungal community assembly
in drought-stressed sorghum shows stochasticity, selection, and universal ecological dynamics. Nat
Commun. 2020;11(1):34. Epub 2020/01/09. doi: 10.1038/s41467-019-13913-9. PubMed PMID:
31911594; PubMed Central PMCID: PMCPMC6946711.

19. Vila JCC, Liu YY, Sanchez A. Dissimilarity-Overlap analysis of replicate enrichment communities.
ISME J. 2020;14(10):2505-13. Epub 20200618. doi: 10.1038/s41396-020-0702-7. PubMed PMID:
32555503; PubMed Central PMCID: PMCPMC7490387.

20. San-Juan-Vergara H, Zurek E, Ajami NJ, Mogollon C, Pena M, Portnoy I, et al. A Lachnospiraceae-
dominated bacterial signature in the fecal microbiota of HIV-infected individuals from Colombia,
South America. Sci Rep. 2018;8(1):4479. Epub 20180314. doi: 10.1038/s41598-018-22629-7.
PubMed PMID: 29540734; PubMed Central PMCID: PMCPMC5852036.

21. Gonze D, Coyte KZ, Lahti L, Faust K. Microbial communities as dynamical systems. Current
Opinion in Microbiology. 2018;44:41-9. doi: 10.1016/j.mib.2018.07.004. PubMed PMID:
WOS:000447581000007.

22. Franzosa EA, Huang K, Meadow JF, Gevers D, Lemon KP, Bohannan BJM, et al. Identifying
personal microbiomes using metagenomic codes. Proceedings of the National Academy of Sciences.
2015;112(22):E2930-E8. doi: 10.1073/pnas.1423854112. PubMed PMID: WOS:000355832200014.

23. Faith JJ, Guruge JL, Charbonneau M, Subramanian S, Seedorf H, Goodman AL, et al. The long-term
stability of the human gut microbiota. Science. 2013;341(6141):1237439. Epub 2013/07/06. doi:
10.1126/science.1237439. PubMed PMID: 23828941; PubMed Central PMCID: PMC3791589.

24. Bik EM, Costello EK, Switzer AD, Callahan BJ, Holmes SP, Wells RS, et al. Marine mammals
harbor unique microbiotas shaped by and yet distinct from the sea. Nat Commun. 2016;7:10516.
Epub 20160203. doi: 10.1038/ncomms10516. PubMed PMID: 26839246; PubMed Central PMCID:
PMCPMC4742810.

25. Caporaso JG, Lauber CL, Costello EK, Berg-Lyons D, Gonzalez A, Stombaugh J, et al. Moving
pictures of the human microbiome. Genome Biology. 2011;12(5):R50. doi: Artn R50
Doi 10.1186/Gb-2011-12-5-R50. PubMed PMID: ISI:000295732700014.

26. Costello EK, Lauber CL, Hamady M, Fierer N, Gordon JI, Knight R. Bacterial community variation
in human body habitats across space and time. Science. 2009;326(5960):1694-7. doi: Doi
10.1126/Science.1177486. PubMed PMID: ISI:000272839000053.

27. Louca S, Polz MF, Mazel F, Albright MBN, Huber JA, O'Connor MI, et al. Function and functional
redundancy in microbial systems. Nat Ecol Evol. 2018;2(6):936-43. Epub 2018/04/18. doi:
10.1038/s41559-018-0519-1. PubMed PMID: 29662222.

28. Rainey PB, Quistad SD. Toward a dynamical understanding of microbial communities. Philos Trans
R Soc Lond B Biol Sci. 2020;375(1798):20190248. Epub 2020/03/24. doi: 10.1098/rstb.2019.0248.
PubMed PMID: 32200735; PubMed Central PMCID: PMCPMC7133524.
29. Martiny JB, Jones SE, Lennon JT, Martiny AC. Microbiomes in light of traits: A phylogenetic
perspective. Science. 2015;350(6261):aac9323. doi: 10.1126/science.aac9323. PubMed PMID:
26542581.
30. Debray R, Herbert RA, Jaffe AL, Crits-Christoph A, Power ME, Koskella B. Priority effects in
microbiome assembly. Nat Rev Microbiol. 2022;20(2):109-21. Epub 20210827. doi: 10.1038/s41579-
021-00604-w. PubMed PMID: 34453137.
31. Risely A, Wilhelm K, Clutton-Brock T, Manser MB, Sommer S. Diurnal oscillations in gut bacterial
load and composition eclipse seasonal and lifetime dynamics in wild meerkats. Nat Commun.
2021;12(1):6017. Epub 20211014. doi: 10.1038/s41467-021-26298-5. PubMed PMID: 34650048;
PubMed Central PMCID: PMCPMC8516918.
32. Kolodny O, Weinberg M, Reshef L, Harten L, Hefetz A, Gophna U, et al. Coordinated change at the
colony level in fruit bat fur microbiomes through time. Nature Ecology & Evolution. 2019;3(1):116-
24. doi: 10.1038/s41559-018-0731-z. PubMed PMID: WOS:000453767000021.
33. Flores GE, Caporaso JG, Henley JB, Rideout JR, Domogala D, Chase J, et al. Temporal variability is
a personalized feature of the human microbiome. Genome Biology. 2014;15(12):531. doi: ARTN 531
10.1186/s13059-014-0531-y. PubMed PMID: WOS:000346609500011.
34. Johnson AJ, Vangay P, Al-Ghalith GA, Hillmann BM, Ward TL, Shields-Cutler RR, et al. Daily
Sampling Reveals Personalized Diet-Microbiome Associations in Humans. Cell Host & Microbe.
2019;25(6):789-802. Epub 2019/06/14. doi: 10.1016/j.chom.2019.05.005. PubMed PMID: 31194939.
35. Smits SA, Marcobal A, Higginbottom S, Sonnenburg JL, Kashyap PC. Individualized Responses of
Gut Microbiota to Dietary Intervention Modeled in Humanized Mice. mSystems. 2016.
36. Kalyuzhny M, Shnerb NM. Dissimilarity-overlap analysis of community dynamics: Opportunities
and pitfalls. Methods in Ecology and Evolution. 2017;8:1764-73.
37. Marsland R, 3rd, Cui W, Mehta P. A minimal model for microbial biodiversity can reproduce
experimentally observed ecological patterns. Sci Rep. 2020;10(1):3308. Epub 20200224. doi:
10.1038/s41598-020-60130-2. PubMed PMID: 32094388; PubMed Central PMCID:
PMCPMC7039880.
38. Faust K, Lahti L, Gonze D, de Vos WM, Raes J. Metagenomics meets time series analysis:
unraveling microbial community dynamics. Current Opinion in Microbiology. 2015;25:56-66. doi:
10.1016/j.mib.2015.04.004. PubMed PMID: 26005845.
39. Gloor GB, Macklaim JM, Pawlowsky-Glahn V, Egozcue JJ. Microbiome datasets are compositional:
and this is not optional. Front Microbiol. 2017;8:2224. Epub 2017/12/01. doi:
10.3389/fmicb.2017.02224. PubMed PMID: 29187837; PubMed Central PMCID:
PMCPMC5695134.
40. Quinn TP, Richarrson MF, Lovell D, Crowley TM. propr: An R-package for Identifying
Proportionally Abundant Features Using Compositional Data Analysis Scientific Reports.
2017;7:16252.
41. Alberts SC, Altmann J. The Amboseli Baboon Research Project: Themes of continuity and change.
In: Kappeler P, Watts DP, editors. Long-term field studies of primates: Springer Verlag; 2012. p. 261-
87.
42. Bjo□rk J, Dasari M, Grieneisen L, Gould TJ, Grenier JC, Yotova V, et al. Synchrony and
idiosyncrasy in the gut microbiome of wild primates. Nature Ecology & Evolution. 2022;6:955-64.
doi: https://www.biorxiv.org/content/10.1101/2021.11.24.469913v1.
43. Grieneisen L, Dasari M, Gould TJ, Bjo□rk JR, Grenier JC, Yotova V, et al. Gut microbiome
heritability is near-universal but environmentally contingent. Science. 2021;373:181-6.
44. Vatanen T, Kostic AD, d'Hennezel E, Siljander H, Franzosa EA, Yassour M, et al. Variation in
Microbiome LPS Immunogenicity Contributes to Autoimmunity in Humans. Cell. 2016;165(4):842-

53. Epub 20160428. doi: 10.1016/j.cell.2016.04.007. PubMed PMID: 27133167; PubMed Central
PMCID: PMCPMC4950857.
45. Cullen CM, Aneja KK, Beyhan S, Cho CE, Woloszynek S, Convertino M, et al. Emerging Priorities
for Microbiome Research. Front Microbiol. 2020;11:136. Epub 2020/03/07. doi:
10.3389/fmicb.2020.00136. PubMed PMID: 32140140; PubMed Central PMCID:
PMCPMC7042322.
46. Silverman JD, Roche K, Holmes ZC, David LA, Mukherjee S. Bayesian Multinomial Logistic
Normal Models through Marginally Latent Matrix-T Processes. Journal of Machine Learning
Research. 2022;23:1-42.
47. Vetrovsky T, Baldrian P. The variability of the 16S rRNA gene in bacterial genomes and its
consequences for bacterial community analyses. PLoS One. 2013;8(2):e57923. Epub 20130227. doi:
10.1371/journal.pone.0057923. PubMed PMID: 23460914; PubMed Central PMCID:
PMCPMC3583900.
48. Guittar J, Shade A, Litchman E. Trait-based community assembly and succession of the infant gut
microbiome. Nat Commun. 2019;10(1):512. Epub 2019/02/03. doi: 10.1038/s41467-019-08377-w.
PubMed PMID: 30710083; PubMed Central PMCID: PMCPMC6358638.
49. Kehe J, Ortiz A, Kulesa A, Gore J, Blainey PC, Friedman J. Positive interactions are common among
culturable bacteria. Science Advances. 2021;45:eabi7159.
50. Weiss AS, Burrichter AG, Durai Raj AC, von Strempel A, Meng C, Kleigrewe K, et al. In vitro
interaction network of a synthetic gut bacterial community. ISME J. 2022;16(4):1095-109. Epub
20211202. doi: 10.1038/s41396-021-01153-z. PubMed PMID: 34857933; PubMed Central PMCID:
PMCPMC8941000.
51. Ortiz A, Vega NM, Ratzke C, Gore J. Interspecies bacterial competition regulates community
assembly in the C. elegans intestine. ISME J. 2021;15(7):2131-45. Epub 20210215. doi:
10.1038/s41396-021-00910-4. PubMed PMID: 33589765; PubMed Central PMCID:
PMCPMC8245486.
52. Carlstrom CI, Field CM, Bortfeld-Miller M, Muller B, Sunagawa S, Vorholt JA. Synthetic microbiota
reveal priority effects and keystone strains in the Arabidopsis phyllosphere. Nat Ecol Evol.
2019;3(10):1445-54. Epub 20190926. doi: 10.1038/s41559-019-0994-z. PubMed PMID: 31558832;
PubMed Central PMCID: PMCPMC6774761.
53. Venturelli OS, Carr AC, Fisher G, Hsu RH, Lau R, Bowen BP, et al. Deciphering microbial
interactions in synthetic human gut microbiome communities. Mol Syst Biol. 2018;14(6):e8157.
Epub 20180621. doi: 10.15252/msb.20178157. PubMed PMID: 29930200; PubMed Central PMCID:
PMCPMC6011841.
54. Meehan CJ, Beiko RG. A phylogenomic view of ecological specialization in the Lachnospiraceae, a
family of digestive tract-associated bacteria. Genome Biol Evol. 2014;6(3):703-13. doi:
10.1093/gbe/evu050. PubMed PMID: 24625961; PubMed Central PMCID: PMCPMC3971600.
55. Vacca M, Celano G, Calabrese FM, Portincasa P, Gobbetti M, De Angelis M. The Controversial Role
of Human Gut Lachnospiraceae. Microorganisms. 2020;8(4). Epub 20200415. doi:
10.3390/microorganisms8040573. PubMed PMID: 32326636; PubMed Central PMCID:
PMCPMC7232163.
56. Stackebrandt E. The Family Lachnospiraceae. The Prokaryotes. 2014.
57. Costello EK, Stagaman K, Dethlefsen L, Bohannan BJ, Relman DA. The application of ecological
theory toward an understanding of the human microbiome. Science. 2012;336(6086):1255-62. Epub
2012/06/08. doi: science.1224203 [pii]
10.1126/science.1224203. PubMed PMID: 22674335.
58. Walter J, Ley R. The human gut microbiome: ecology and recent evolutionary changes. Annual
Review of Microbiology. 2011;65:411-29. Epub 2011/06/21. doi: 10.1146/annurev-micro-090110-
102830. PubMed PMID: 21682646.
59. McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis and graphics
of microbiome census data. PLoS One. 2013;8(4):e61217. Epub 20130422. doi:

10.1371/journal.pone.0061217. PubMed PMID: 23630581; PubMed Central PMCID:
PMCPMC3632530.
60. Yaqing Chen [aut c, Alvaro Gajardo [aut], Jianing Fan [aut], Zhong Q, Dubey P, Han K, et al. frechet:
Statistical Analysis for Random Objects and Non-Euclidean Data. Available: https://CRANR-
projectorg/package=frechet. 2020.
61. Schliep KP. phangorn: phylogenetic analysis in R. Bioinformatics. 2011;27(4):592-3. Epub
20101217. doi: 10.1093/bioinformatics/btq706. PubMed PMID: 21169378; PubMed Central PMCID:
PMCPMC3035803.
62. Guillot G, Rousset F. Dismantling the Mantel tests. In: Harmon L, editor. Methods in Ecology and
Evolution. 42013. p. 336–44.
1022