Physiological and behavioural resistance of malaria vectors in rural West-Africa : a data mining study to adress their fine-scale spatiotemporal heterogeneity, drivers, and predictability

Paul Taconet[1], Dieudonne Diloma Soma[2], Barnabas Zogo[3], Karine Mouline[1], Frederic Simard[1], Alphonsine Amanan Koffi[3], Roch Kounbobr Dabire[2], Cedric Pennetier[1,2] and Nicolas Moiroux[1]

[1]Institut de Recherche pour le Developpement, MIVEGEC, Univ. Montpellier, CNRS, IRD, 91 Avenue Agropolis, BP 64501, 34394, Montpellier, France
[2]Institut de Recherche en Sciences de La Sante (IRSS), Bobo-Dioulasso, Burkina Faso
[3]Institut Pierre Richet (IPR), Bouake, Cote d'Ivoire

ABSTRACT
Insecticide resistance and behavioral adaptation of malaria mosquitoes impact the efficacy of long-lasting
insecticide nets - currently the main malaria vector control tool. To develop and deploy complementary,
efficient and cost-effective control interventions, a good understanding of the drivers of these physiological
and behavioural traits is needed. In this data-mining work, we modeled a set of indicators of physiological
resistances to insecticide (prevalence of three target-site mutations) and biting behaviours (early- and
late-biting, exophagy) of anopheles mosquitoes in two rural areas of West-Africa, located in Burkina Faso
and Cote d'Ivoire. To this aim, we used mosquito field collections along with heterogeneous, multisource
and multi-scale environmental data. The objectives were i) to assess the small-scale spatial and temporal
heterogeneity of the indicators, ii) to better understand their drivers, and iii) to assess their spatio-temporal
predictability, at scales that are consistent with operational action. The explanatory variables covered a wide
range of potential environmental determinants of vector resistance to insecticide or feeding behaviour :
vector control, human availability and nocturnal behaviour, macro and micro-climatic conditions, landscape,
etc. The resulting models revealed many statistically significant associations, although their predictive powers
were overall weak. We interpreted and discussed these associations in light of several topics of interest, such
as : respective contribution of public health and agriculture in the development of physiological resistances,
biological costs associated with physiological resistances, biological mechanisms underlying biting behavior,
and impact of micro-climatic conditions on the time or place of biting. To our knowledge, our work is the first
studying insecticide resistance and feeding behaviour of malaria vectors at such fine spatial scale with such a
large dataset of both mosquito and environmental data.

21
22 **Keywords:** malaria, anopheles, insectide resistance, behavioural adaptation, exophagy, early-biting, late-biting, spatiotemporal distribu-
23 tion, statistical modeling, data mining

Introduction

Malaria remains a major public health concern in Africa, with 234 million cases and 593 000 death over the continent in 2021 (WHO, 2022). After years of steady reduction in the disease transmission mainly due to the scale-up of vector control (VC) interventions (in particular insecticide-based tools such as long lasting insecticide nets (LLIN) and indoor residual spraying (IRS)) (Bhatt et al., 2015), progress is now stalling since 2015 (WHO, 2022). Involved in such worrying trends are a combination of biological, environmental and socio-economical factors. The mosquito biology, with the buildup of adaptive changes in the mosquito vectors populations enabling them to avoid or circumvent the lethal effects of insecticides, is most likely playing a very important contribution (Killeen, 2014). These changes are framed as vector *resistance* to insecticides. As a consequence of the widespread use of insecticides (in agriculture and public health), vector resistance has arisen rapidly in malaria vectors in many areas of Africa and above (Durnez and Coosemans, 2013; Riveron et al., 2018); and as previously indicated, is now at such level that it compromises the effectiveness of the most efficient malaria control interventions (Gatton et al., 2013; Hemingway et al., 2016; Killeen, 2014; Sokhna et al., 2013). Complementary and locally-tailored VC strategies taking into account the great diversity of vectors resistance mechanisms (see below) are therefore needed to target these vectors contributing to residual malaria transmission (Corbel and N'Guessan, 2013; Durnez and Coosemans, 2013; Hemingway et al., 2016; Moiroux, 2012; Riveron et al., 2018; Sokhna et al., 2013; WHO, 2017).

Vector resistances to insecticide are usually split into two categories : *physiological* and *behavioural* resistance (Lockwood et al., 1984; Sokhna et al., 2013). Physiological resistance refers to biochemical and morphological mechanisms (e.g. target-site modifications, metabolic resistance, cuticular thickness) that enable the mosquito to withstand the effects of insecticide despite its contact with it (Davidson, 1957). Among the physiological resistances, the target-site mutations L1014F (*kdr-w*) (Martinez-Torres et al., 1998), L1014S (*kdr-e*) (Ranson et al., 2000), and G119S (*ace-1*) (Weill et al., 2004), conferring insecticide resistance to pyrethroids (*kdr-w* and *kdr-e*) and to carbamates and organophosphates (*ace-1*), have been extensively described. Behavioral resistance, on its side, refers to any modification of mosquito behavior that facilitates avoidance or circumvention of insecticides (Carrasco et al., 2019; Gatton et al., 2013; Riveron et al., 2018). Behavioral resistance of mosquitoes to insecticides can be qualitative (i.e. modifications that prevent or limit the contact with the insecticide) or quantitative (i.e. modifications that stop, limit or reduce insecticide action once contact has occurred, e.g. escaping, behavioural thermoregulation or curative self-medication) (Carrasco et al., 2019). Up-to-date, the behavioral resistance mechanisms described in the literature are mainly qualitative and consist in spatial, temporal, or trophic avoidance. In particular, in the anopheline populations, the following behavioral qualitative resistance mechanisms have been described after the scale-up of insecticide-based VC tools (Durnez and Coosemans, 2013) : i) increase of exophagic or exophilic behaviours (spatial avoidance), where mosquitoes shifted from biting or resting indoor to outdoor, ii) increase of early- or late-biting behaviours (temporal avoidance), where mosquitoes shifted from biting at night to earlier in the evening or later in the morning, iii) increase of zoophagic behaviours (trophic avoidance), where mosquitoes shifted from biting on humans to biting on animals.

To help develop and deploy complementary VC strategies that are efficient and cost-effective, a better understanding of the spatiotemporal distribution and drivers of both vector physiological resistance and feeding behaviour is needed at a local scale. We raise here a set of questions that, among others, must be explored further at local scale towards this aim :

> ***What is the respective contribution of public health and agriculture in the development of physiological resistances in Anopheles vectors ?*** The molecular and genetic basis of physiological resistance has been widely acknowledged: under the pressure of insecticides, mutations that enable the vectors to survive are naturally selected and then spread over the generations (Labbé et al., 2017; Martinez-Torres et al., 1998). The main

force that governs the development of a physiological mechanism of resistance in a population of insects is therefore the selective pressure induced by insecticide exposure. This pressure can be induced by the vector control tools, or by the runoff of pesticides used in agriculture (in many cases, the same as those used for impregnation of bed nets) into the malaria vectors breeding sites (Chandre et al., 1999; Hien et al., 2017; Reid and McKenzie, 2016; Yadouleton et al., 2011). Assessing the relative contribution of these two selection pressures on the development of resistant phenotypes is critical to further predict the relative impacts of public health and agriculture on the growth of physiological resistances and act consequently.

> *What are the biological mechanisms underlying behavioural resistances ?* Contrary to physiological resistance, the biological mechanisms underlying behavioral resistance are still poorly known (Carrasco et al., 2019; Durnez and Coosemans, 2013; Killeen, 2014; Main et al., 2016). In particular, a pending question, having important implications for vector control, is whether behavioural shifts reflect evolutionary adaptations in response to selection pressures from vector control tools, as for physiological resistances (*constitutive resistance*) or are manifestations of pre-existing phenotypic plasticity which is triggered when facing the insecticide or in response to environmental variation that reduces human host availability (*inducible resistance*). Inducible resistance imply that vectors rapidly revert to baseline behaviours when VC interventions are lifted, whereas constitutive resistance might progressively and durably erode the effectiveness of current VC tools. Understanding the biological mechanisms underlying behavioural resistances is therefore important to assess the long-term efficacy of insecticide-based VC interventions.

> *Are mosquito biting behaviours modulated by local-scale environmental conditions other than insecticide-related ones ?* As aforementioned, the overall rise of behavioral resistances is likely caused by the widespread of insecticide-based vector control interventions. However, local environmental conditions can modulate vector behaviours at the time of foraging activity. Local climatic conditions – e.g. wind, rain, temperature, humidity, luminosity - may for example affect the timing and location of vector biting, as it has been noted in some studies (Kirby and Lindsay, 2004; Kreppel et al., 2020; Ngowo et al., 2017). Mosquitoes with natural endophagic / endophilic preferences might, for example, bite or rest outside if temperature inside is too high or humidity too low, in order to decrease their risk of desiccation-related mortality (Kreppel et al., 2020; Ngowo et al., 2017). Land cover, as well, can affect biting rhythms. It has been noted for example that distance to breeding sites may influence nocturnal host-seeking behaviour, with vectors biting on average earlier in the night in households located close to the breeding sites (Njan Nloga et al., 1993; Snow and Gilles, 2002). Assessing whether and to which extent behavioural resistance traits are influenced by local environmental (climatic or landscape) settings may help design VC tools exploiting the vulnerabilities of vectors.

> *Are there associations between behavioural and physiological resistances ?* Physiological and behavioral resistances may likely coexist in mosquito populations, with the first possibly influencing the second. In fact, physiologically resistant mosquitoes may, theoretically, use the recognition of insecticide-based control tool as a proxy for host presence (framed as *behavioural exploitation* (Carrasco et al., 2019)). Several studies have actually showed that the kdr mutation can modify the host-seeking or biting behavior of *Anopheles* in presence of insecticide-treated net (Diop, Moiroux, et al., 2015; Diop, Chandre, et al., 2021; Porciani et al., 2017). Such behavioural exploitation could potentially lead to a better host recognition/localization and have a dramatic impact, with the control intervention having the opposite effect to the one expected. It is hence important to assess if and to which extent physiologically resistant mosquitoes exhibit different biting behaviours than their susceptible counterparts.

> *Which adaptative strategy (physiological or behavioural resistance) arises faster ?* Understanding the relative capacity of mosquitoes to develop physiological resistance and to shift their behaviour in response to vector control is necessary to highlight where and when mitigation efforts should be prioritized (Sanou et al.,

2021). After introduction / re-introduction of insecticide-based tools, if vectors rapidly shift their behaviour to feed outside or at times when people are not protected by an LLIN, interventions that target such mosquitoes should be quickly deployed. In contrast, the rapid emergence of physiological resistance in vectors who continue to feed indoors and at night indicates that switching to alternative insecticide classes in indoor-based interventions may have a greater impact. Additionally, for a given environment, assessing the relative rate of development of physiological and behavioral resistances is of direct epidemiological importance : it has been showed for example that under a scenario where LLIN and IRS are both heavily used, changes in the susceptibility to insecticide is likely to have a bigger epidemiological impact than changes in biting times (Sherrard-Smith et al., 2019).

> ***Are resistance rates heterogeneous at small spatiotemporal scales ?*** Mosquito presence and abundance has already been found heterogeneous in space and time at fine-scale, calling for locally-tailored (species-, season-, and village-specific) control interventions (Moiroux, Bio-Bangana, et al., 2013; Moiroux, Djènontin, et al., 2014; Taconet, Porciani, et al., 2021). However, little is known about the small-scale spatiotemporal heterogeneity of vector resistance. The potential drivers of the development or triggering of resistant phe-notypes (vector control use, land cover, micro-climate, human behaviour, etc.) are likely to vary at small spatiotemporal scales, and so may, at the end of the line, vector resistance. As for abundances, assessing the level of heterogeneity of resistance rates in space and time is important to assess the spatiotemporal scale at which management of vector resistance should be considered.

> ***To what extent can we explain and predict vector resistance and biting behaviour in space and time ?*** Assessing the levels of explainability and predictability of vector resistance and biting behaviour is important for both scientific and operational purposes. Towards this aim, generating statistical models linking vector resistances or biting behaviours to their potential drivers and assessing their explanatory and predictive powers can help (Shmueli, 2010; Shmueli and Koppius, 2010). High explanatory or predictive powers in the models might suggest that the conditions driving a vector to resist are well understood, and conversely, low explanatory powers might suggest that resistances are driven by factors either yet undiscovered or not included in the models. Additionally, assessing the predictability of resistances in vector populations in space and time is an important step towards mapping vector resistance at every place (e.g. village) and time (e.g. season) in the area, with such decision-support tools important to deploy the right intervention, at the right place and time (Taconet, Porciani, et al., 2021).

In this study, we used field mosquito collections and environmental data collected simultaneously in two rural areas of West-Africa to bring elements of answer to these questions for our areas. Guided by these questions, our overall objectives were i) to assess the fine-scale prevalence and spatiotemporal heterogeneity of physiological resistances and at-risk biting behaviours of malaria vectors in these areas and ii) to better understand their drivers. To do so, we modeled a set of indicators of physiological resistances and biting behaviours (namely exophagy, early-biting, late-biting, *kdr-w*, *kdr-e*, and *ace-1* target-site mutations) at the individual mosquito level using this fine-grained dataset and advanced statistical methods in an exploratory and data-driven approach. Patterns found in the data were interpreted and discussed in light of the topics aforementioned, of importance for the management of malaria residual transmission. We concluded with a set of recommendations to manage vector resistances in our study areas.

Methods

Entomological and environmental data

The data used in this work were collected in the frame of the REACT project (Soma, Zogo, Somé, et al., 2020; Zogo et al., 2019). In this projet, a total of fifty-five villages, distributed in two West-African rural areas (~ 2500 squared km each) located in the areas of Diébougou (southwestern Burkina Faso (BF)) and Korhogo (northern Ivory Coast (IC)) were selected according to the following criteria: accessibility during the rainy season, 200–500 inhabitants per village, and distance between two villages higher than two kilometers. After an exhaustive census of the population in these villages at the beginning of the project, entomological and human behaviours surveys were regularly conducted during 15 months (1.25 year) in the Diébougou area and 18 months (1.5 year) in the Korhogo area. Vector control interventions were implemented both as part of the project and of the national malaria control programs (see below). Figure 1 shows the study areas and the corresponding timelines for data collection and vector control interventions. Entomological data were collected in the field, and environmental data were collated from specific devices (see below) or created from heterogeneous field and satellite-based sources. Below is a description of the data used in our work.

Figure 1. Villages of the study and timeline for data collection + vector control

> *Anopheles collections*

Several rounds of mosquito collections (eight in the Korhogo (IC) area, seven in the Diébougou (BF) area) were conducted in each village. The periods of the surveys span the typical climatic conditions of these tropical areas (except the high rainy season - July to September) (see Additional file 1.A for the spatiotemporal trends of the meteorological conditions). Mosquitoes were collected using the Human Landing Catch (HLC) technique from 17:00 to 09:00 both indoors and outdoors at four sites per village (i.e. eight collection points) for one night during each survey. Malaria vectors were identified using morphological keys. All individuals belonging to

the *Anopheles funestus* group (in both study areas) and *Anopheles gambiae s.l.* complex (in BF) were identified to the species level using PCR. In IC, due to the very large numbers of *An. gambiae s.l.* vectors collected, a sub-sample only of these individuals (randomly selected in space and time) was identified to species. Finally, PCR assay were carried out on all the *An. gambiae s.s.* and *An. coluzzii* collected in BF to detect the L1014F (*kdr-w*), the L1014S (*kdr-e*) and the G119S (*ace-1*) target-site mutations. Detailed descriptions of the methods used to obtain these data are provided in Soma, Zogo, Somé, et al. (2020); Zogo et al. (2019). These data are published as a Darwin Core archive in the Global Biodiversity Information Facility (GBIF) and available at this URL : https://doi.org/10.15468/v8fvyn (DD Soma et al., 2023).

> ***Data on weather preceding mosquito collections and during mosquito collections***

Weather can impact the fitness or the activity of resistant genotypes (Kliot and Ghanim, 2012), as well as the biting behaviour of the mosquitoes (see Introduction). In this work, we recorded or retrieved weather conditions : (i) during mosquito collections (i.e. the HLC sessions), (ii) during the day of collection, and (iii) during the month preceding collection. Weather on the day of collection and during mosquito collection may impact the relative activity of each genotype and phenotypes associated with resistances. Weather during the month preceding the survey, on its side, can impact development and survival rates of both the current and parental generations of collected mosquitoes. Regarding our outputs (prevalence of behavioral phenotypes and target-site mutations - see next section), weather during the month preceding collection may therefore impact the fitness of the studied genotypes (for target-site mutations) or possible – and unknown - genotypes associated with studied behavioral phenotypes.

Micro-climatic conditions (temperature, relative humidity, luminosity and atmospheric pressure) were simultaneously recorded where mosquito collections were being conducted. Instruments used to record these data were : for temperature and relative humidity : Hygro Buttons 23 Data Loggers [Proges Plus DAL0084] (temporal resolution (TR): 15 minutes) ; for luminosity : HOBO Pendant® Temperature/Light 8K Data Logger (TR: 15 minutes) ; for atmospheric pressure : Extech SD700 Data Loggers (TR : 10 minutes). Hygro and Hobo loggers were positioned both inside and outside the houses where mosquito sampling was conducted, close to the sampling positions. The barometer was positioned at the center of the village. These field data were completed with satellite or modeled data available at coarser spatial resolutions : rainfall (spatial resolution (SR) : ~ 11 km, TR : 30 min, source : Global Precipitation Measurement (GPM) IMERG (GSFC, 2019), wind speed (SR : ~ 28 km,TR : 1h, source : ERA5 (Hersbach et al., 2020)), apparent magnitude of the Moon (SR : 0.001 degrees, TR : 1 day, source : Institute of celestial mechanics and ephemeris calculations).

Meteorological conditions on the day of collection and during the month preceding collection were extracted from satellite imagery. Namely, rainfall estimates were extracted from the GPM - IMERG daily Final products (Center, 2019). Diurnal and nocturnal temperatures were derived from the Moderate Resolution Imaging Spectroradiometer (MODIS) daily Land Surface Temperature (LST) Terra and Aqua products (Wan et al., 2015a,b). Rainfall and temperatures were averaged in a 2 km buffer zone around each HLC collection point to create meteorological variables on both the day of collection and the month preceding collection. Detailed descriptions of the methods used to collect and process these data are provided in Taconet, Porciani, et al. (2021).

> ***Data on host availability and human behaviour***

The nocturnal behaviour of humans (hours inside the dwellings, hours of use of LLINs) drives host availability for the mosquitoes and can therefore impact their behaviour. For instance, high LLIN use rate can drive mosquitoes to feed outside, at times when people are not protected, or on alternative sources of blood

(Durnez and Coosemans, 2013). Here, human population was counted in each village, through an exhaustive census conducted at the beginning of the project. Then, several human behavioural surveys (two in IC, three in BF) were carried out in each village (see Figure 1). For each survey and village, several households (mean = 14, SD = 2) were randomly selected, and for each household, one to three persons in each age class (0–5 years old, 6–17 years old and ≥ 18 years old) were selected. The head of the household was then asked, for each selected person, on the night preceding the survey : i) whether he/she used an LLIN or not, ii) the time at which he/she entered and exited his own house, and iii) the time at which he/she entered and exited his LLIN-protected sleeping space (where appropriate). Households for human behavioural surveys were independently selected from households for entomological surveys. The surveys were conducted after the distribution of the LLINs (see below), and span the typical climatic conditions of the areas. Detailed descriptions of the methods used to collect these data are provided in Soma, Zogo, Taconet, et al. (2021).

> ### *Landscape data*

Landscape can have an impact on mosquito foraging behaviour (e.g. the distance to breeding sites can impact biting rhythms) or physiological resistance (e.g. through pesticides used in crops) (see Introduction). Digital land cover maps were produced for each study area by carrying out a Geographic Object-Based Image Analysis (Hay and Castilla, 2008) using multisource very high and high resolution satellite-derived products. From these maps, several variables were derived : the percentage of landscape occupied respectively by cotton fields, by rice fields, and by the other crops (mainly leguminous crops, millet, sorghum) in a 2 km buffer size area around each collection point ; and the distance to the nearest stream (as a proxy for the distance to the breeding site). For cotton, the variable was binarized as presence / absence of cotton cultivated due to the small range of values. In addition, the geographical location of the households was recorded, and used to derive two indices : the degree of clustering of the households in each village, and the distance from each collection point to the edge of the village. Detailed descriptions of the methods used to collect, generate and process these data are provided in Taconet, Porciani, et al. (2021).

> ### *Vector control*

Repeated exposure to insecticides used in vector control interventions is undoubtedly one of the most important drivers of the development of resistance (see Introduction). In both Burkina Faso and Ivory Coast, LLINs have been universally distributed every 3-4 years since 2010 (PNLP, 2014a,b). In BF, a mass distribution of LLINs (PermaNet 2.0) was carried out by the National Malaria Control Program in July 2016 (i.e. 6 months before our first entomological survey). In IC, our team distributed LLINs in the villages of the project in June 2017 (i.e. height month after the first entomological survey and ten months before the last one). Complementary VC tools were implemented in some of the villages in the middle of the project - namely IRS, ivermectin to peri-domestic animals (IVM), intensive Information Education and Communication to the populations (IEC), and larval control (Larv.) as part of a randomized controlled trial aiming at assessing the benefits of new, complementary VC strategies (Soma, Zogo, Somé, et al., 2020; Zogo et al., 2019) (see Figure 1, and Additional file 1 available online at this URL (along with the other supplementary material) : https://doi.org/10.23708/VJEEMU (Taconet, DD Soma, Zogo, Mouline, Simard, Koffi Amanan, et al., 2023)).

Statistical analyses

Dependent and independant variables

Six vector potential resistance indicators were modelled :

- three indicators of physiological resistance to insecticide : *kdr-w* mutation, *kdr-e* mutation, *ace-1* mutation,

- three indicators of biting behaviours : early biting, late biting, exophagy. **Here, it is unknown whether these behaviours are real *behavioural resistances* as defined in the introduction (i.e. induced by the use of insecticides) or are natural behaviours (i.e. behaviours that exist naturally in vector populations). However, to harmonise the vocabulary used and facilitate the reading, in the remainder of this manuscript, we will qualify these three phenotypes, associated with resistant behaviours, as 'behavioural resistance'.**

Exophagy was defined as the probability for a host-seeking mosquito to bite outdoor (as opposed to indoor). Early biting was defined as the probability for a host-seeking mosquito to bite before 50 % of the LLIN users were declared to be under their bednet in the evening, and late biting was defined as the probability for a host-seeking mosquito to bite after 50 % of the LLIN users were declared to be out of their bednet in the morning (based on the closest - in space and time - human behaviour survey). *Kdr-w*, *kdr-e* and *ace-1* mutations were defined as the probabilities for an allele of a host-seeking mosquito to be mutated (as opposed to wild type). The statistical unit was therefore the mosquito for biting behaviour models and the allele for physiological resistance models. Dependent variables were all binary (0 = absence of resistance/mutation, 1 = presence of resistance/mutation) and models outcomes were probabilities for a mosquito (resp. allele) to be resistant (resp. mutated). Each indicator was modeled separately for each main species in each study area, as determinants of resistance might be species- or site-specific (i.e. mosquitoes might respond differently to environmental variations depending on the species and study area, due to potential local chromosomal forms, adaptation, etc.) (Durnez and Coosemans, 2013; Riveron et al., 2018). As three main species were found in BF and two in IC (see Results section), a total of twenty-one dependent variables were built (exophagy : 3 in BF and 2 in IC ; early biting : 3 in BF and 2 in IC ; late biting : 3 in BF and 2 in IC ; *kdr-w* : 2 in BF ; *kdr-e* : 2 in BF ; *ace-1* : 2 in BF). Based on literature (see Introduction) and available data, we then built independent variables representing potential determinants of each of these resistant phenotypes. These variables are provided in Table 1. To build these variables, the source data were possibly aggregated in space or time, at varying resolutions depending on the considered dependent variable. Replication data are available online at https://doi.org/10.23708/LV8GEW (Taconet, DD Soma, Zogo, Mouline, Simard, Koffi, et al., 2023).

Table 1. Dependent and independent variables for the resistance models. Each bullet point in the **Independant variables** columns represents an independent variable built and used for the considered model. These variables belonged to several families of potential drivers : vector control, vector resistance, host availability, micro-climate at the time of collection, meteorological conditions preceding collection, landscape, and vector abundance. Some independent variables, mentioned as BF only, were available only for the BF study area.

Resistance type	Model name	Dependent variable : Statistical unit & values	Independant variables							
			Vector control	Host availability	Vector resistance / behaviour	Micro-climatic conditions during collection	Meteorological conditions on the day of collection	Meteorological conditions on the month preceding collection	Landscape & crops	Others
Behavioural resistance	Exophagy	**Mosquito** **1 = exophagic** **0 = endophagic**	- Vector control intervention implemented - Time since distribution of LLIN (BF) / Time since first entomological survey (IC)	- Average LLIN use rate in the village on the season of collection - Human population in the village - Perc. of population indoor in the village on the hour of collection - Perc. of the population under an LLIN in the village on the hour of collection	- Kdr-w mutation alleles on the collected mosquito (BF only, An. gambiae s.s. and An. coluzzii) - Kdr-e mutation alleles on the collected mosquito (BF only, An. gambiae s.s. and An. coluzzii)	At the hour of collection : - Temperature indoor - Humidity indoor - Luminosity outdoor - Atmospheric pressure - Relative temperature difference b\w indoor and outdoor - Relative humidity difference b\w indoor and outdoor - Relative luminosity difference b\w indoor and outdoor - Presence / absence of rainfall - Wind speed outdoor - Apparent magnitude of the Moon	- Diurnal temperature - Cumulated rainfall	- Avg. diurnal temperature - Avg. nocturnal temperature - Cumulated rainfall	- Distance from the collection point to the edge of the village - Distance from the collection point to the nearest stream - Degree of clustering of the households in the village	
	Early biting	**Mosquito** **1 = early biter** **0 = all other**		- Average LLIN use rate in the village on the season of collection - Human population in the village	- same as above - Place of collection of the collected mosquito (indoor / outdoor)					
	Late biting	**Mosquito** **1 = late biter** **0 = all other**				During the night of collection : - Avg. temperature - Avg. humidity - Avg. luminosity - Avg. atmospheric pressure - Cumulated rainfall - Avg. wind speed outdoor				
Physiological resistance (BF area only)	Kdrw mutation Kdre mutation Ace1 mutation	**Allele** **1 = resistant** **0 = sensible**		- Average LLIN use rate in the village on the season of collection - Human population in the village - Perc. of population indoor in the village on the hour of collection - Perc. of the population under an LLIN in the village on the hour of collection	- Kdr-w mutation alleles on the collected mosquito - Kdr-w mutation alleles on the collected mosquito - Ace-1 mutation alleles on the collected mosquito	At the hour of collection : - Temperature - Humidity - Luminosity - Atmospheric pressure		- Avg. diurnal temperature - Avg. nocturnal temperature - Cumulated rainfall	In a 2 km buffer area around the collection points : - Presence / absence of cotton fields - Perc. landscape occupied by rice fields - Perc. landscape occupied by other crops	- Number of mosquitoes collected during the night of collection

Modeling workflow

A graphical representation of the modeling workflow (explained below) is available in Additional figure 2. A replication R script (starting from the section 'Multivariate modeling part 1 : Explanatory model') is available online at this URL : https://doi.org/10.23708/LV8GEW (Taconet, DD Soma, Zogo, Mouline, Simard, Koffi, et al., 2023).

Pre-processing. Before modeling, we excluded the dependent variables that had too few 'resistant' observations, according to the following criteria : 'resistant' class $\leq$ 50 observations & $\leq$ 3% of the total observations. Next, we implemented the modeling workflow described below for each remaining dependent variable.

Bivariate modeling. We first excluded the independent variables that were poorly associated with the dependent variable (criteria for exclusion : p-value > 0.2 of a bivariate Generalized Linear binomial Mixed-effect Model (GLMM) with nested random effects at the village and collection site level). Next, we filtered-out collinear variables (Pearson correlation coefficient > 0.7) based on empirical knowledge. With the set of remaining independent variables, two distinct multivariate models were built, with complementary objectives, as explained in the Box 1 below.

Multivariate modeling part 1 : Explanatory model. A binomial GLMM was fitted to the data. Nested random effects were introduced in the model at the village and collection place level. Variables were deleted recursively using an automatic backward variable selection procedure based on the reduction of the Akaike Information Criterion (AIC). Variables belonging to the 'vector control' (for all resistance models) and 'crops' (for physiological resistance models only) groups were forced in the multivariate models (i.e. they were not filtered-out in the variable selection procedure) because there are strong *a priori* assumptions associated with these variables. Such variable selection procedure therefore retained all the 'vector control' and 'crops' variables (whether significantly associated or not with the dependent variable), and the additional variables that decreased the AIC of the multivariate model.

Multivariate modeling part 2 : Predictive model. We additionally fitted a Random Forest (RF) model (Breiman, 2001a) to the data. The model hyperparameters were optimized using a random 5-combinations grid search (Chicco, 2017). Whenever the dependent variable was imbalanced (more than 1/3 imbalance ratio between the positive and negative class), data were up-sampled within the model resampling procedure to cope with well-known problems of machine-learning (ML) models regarding class imbalance (Tyagi and Mittal, 2020).

Assessment of effect sizes and significance of independent variables. To interpret the effect of each independent variable in the GLMM model, we plotted, for each independent variable retained in the final model, the predicted probabilities of resistance across available values of that variable (all other things being equals) (i.e. "partial dependence plot" (PDP) (Friedman and Popescu, 2008)). For reporting and discussion in the manuscript, we kept only variables that had a p-value < 0.05 (results containing the 'full' models are provided in supplementary material, see Results section). To uncover the possible complex relationships that the RF model had learned, we generated smoothed versions of PDPs for each independent variable. However, we restricted the generation of PDPs to the following cases : i) the Area Under the Receiver Operating Characteristics (AUC) (see below) of the model was > 0.6 (because model interpretation tools of ML models (e.g. PDPs) should be trusted only if the predictive power of the underlying model is good enough (Zhao and Hastie, 2021)) and ii) the range of predicted probabilities of resistance was > 0.05 (i.e. the independent variable, over its range of available values, changed the probability of resistance by at least 5 percentage points).

Assessment of models performance. The explanatory power of the GLMM was assessed by calculating the marginal coefficient of determination (R^2) (Nakagawa and Schielzeth, 2013) from the observed and in-sample

predicted values. Marginal R^2 is a goodness-of-fit metric that measures the overall variance explained by the fixed effects in the GLMM. R^2 values were interpreted according to the criteria defined by Cohen (2013) : $R^2 \in$ {0; 0.02{ : very weak ; $R^2 \in$ {0.02; 0.13{ : very weak ; $R^2 \in$ {0.02; 0.13{ : weak ; $R^2 \in$ {0.13; 0.26{ : moderate ; $R^2 \in$ {0.26; 1{ : substantial. The predictive power of the RF model was assessed by leave village - out cross-validation (CV), with the Area under the ROC Curve (AUC) chosen as the performance metric. This CV strategy consisted in recursively leaving-out the observations belonging to one village of collection (i.e. the validation fold), training the model with the observations coming from the other villages (i.e. the training fold), and predicting on the left-out set of observations. We hence measured the ability of the model to predict resistance status ('resistant' or 'non-resistant') on individual mosquitoes caught on new - unseen villages of collection. AUC values were interpreted according to the following criteria : AUC $\in$ {0.5; 0.6{ : very weak ; AUC $\in$ {0.6; 0.65{ : weak ; AUC $\in$ {0.65; 0.75{ : moderate ; AUC $\in$ {0.75; 1{ : substantial.

Box 1 : **Why modeling with both logistic regression and random forests ?**

"Data" models (Breiman, 2001b) - like linear of logistic regression - and "algorithmic" models (Breiman, 2001b) – so-called *machine learning* models like random forests or support vector machines - play complementary roles in scientific theory building and testing. On the one hand, data models are useful to test existing theories and to reach to "statistical" conclusions about causal relationships that exist at the theoretical level, e.g. : vector control significantly impacts vector resistance (or not). These models are hence useful in explanatory modeling, where the main objective is to apply statistical models to data for testing causal hypotheses about theoretical constructs (Shmueli, 2010). On the other hand, algorithmic models, thanks to their ability to inherently capture complex and hidden patterns contained in the data (e.g. interactions, non-linear associations), are useful for assessing the relevance of a theory and for developing new theories or improve existing ones, through predictive analytics (Shmueli and Koppius, 2010). Predictive analytics is the process of building a model aimed at making empirical predictions and then assessing its predictive power. Shmueli and Koppius (2010) describe six roles for predictive analytics in theory building and testing, including generation of new theory, improvement of existing theory, assessment of relevance of a theory, and assessment of level of predictability of a phenomena. In a "big data" context such as that of this study (large datasets, containing lots of observations and variables), predictive analytics is every time more used to support scientific theory development through these roles (Breiman, 2001b; Karpatne et al., 2017; Shmueli and Koppius, 2010).

In our study, we used GLMMs to i) test whether vector control significantly increases vector resistance, as could be expected, and ii) possibly find environmental factors that impact vector resistance in a statistically significant way. We used RFs to i) account for potential unhypothesized, complex relationships (e.g. interactions, non-linear associations) between independent and dependent variables in the statistical modeling process, ii) possibly reveal part of the complex relationships learned by the model (through PDPs), iii) assess distance between theory (potential / identified determinants of vector resistance) and practice (are they enough to predict resistance on unseen mosquitoes ? If not, why ?).

Software and libraries used

The softwares used in this work were exclusively free and open source. The R programming language (R Core Team, 2018) and the R-studio environment (RStudio Team, 2020) were used as the main programming tools. The QGIS software (QGIS Development Team, 2021) and the 'ggplot2' R package (Wickham, 2016) were used

to create respectively the map of the study area and the timeline for data collection. The 'glmmTMB' (Brooks et al., 2017) package was used for the bivariate modeling. The 'buildmer' package (Voeten, 2020) was used to fit the GLMM models with stepwise selection in the multivariate modeling. The 'caret' (Wing et al., 2018) and 'ranger' (Wright and Ziegler, 2017) packages were used were to fit the random forest models in the multivariate modeling. The 'MLmetrics' (Yan, 2016) and 'MuMIn' (Bartoń, 2020) packages were used to calculate respectively the AUC of the RFs and the marginal R^2 of the GLMMs. The 'jtools' (Long, 2020) and 'pdp' (Greenwell, 2017) packages were used to generate the partial dependence plots of respectively the GLMMs and the RFs. The 'broom.mixed' (Bolker and Robinson, 2020) package was used to extract the coefficients / odd ratios, confidence intervals and p-values of the multivariate GLMMs. The 'patchwork' (Pedersen, 2019) and 'gridExtra' (Auguie, 2017) packages were used to create various plot compositions. The 'tidyverse' meta-package (Wickham, 2017) was used throughout the entire analysis. A few additional R packages were used to create, tidy, and transform the data used in this work (see (Taconet, Porciani, et al., 2021)). The LibreOffice suite was used to create the tables and other plot compositions.

Results

Spatio-temporal heterogeneity of vector abundance

In the Korhogo area (IC), a total of 2048 human-nights of HLC was conducted. A sum of 57722 vectors belonging to the *Anopheles* genus was collected. The main species/complex found were *An. gambiae s.l.* and *An. funestus* (respectively 56267 (97% of all the *Anopheles* collected) and 714 (1%) individuals collected). Among the 56267 *An. gambiae s.l.* collected, 3922 (7%) were identified to species: 3726 (95% of the individual identified to species) were *An. gambiae s.s.* and 196 (5%) were *An. coluzzii.* Hence, in the rest of this article, we will consider the *An. gambiae s.l.* collected in the Korhogo area as *An. gambiae s.s.* In the Diébougou area (BF), a total of 1512 human-nights of HLC was conducted. A sum of 3056 vectors belonging to the *Anopheles* genus was collected. The main species found were *An. coluzzii*, *An. gambiae s.s.* and *An. funestus* (respectively 1321 (43% of all the *Anopheles* collected), 616 (20%) and 708 (23%) individuals collected). As expected, mosquito abundance was heterogeneous in time and space (except for *An. funestus* in IC, for which the vast majority (93 %) of the individuals was collected in the first entomological survey, and almost half of the individuals (42 %) were collected within one single village) (see additional file 1 and additional figure 3 for maps and charts of the spatiotemporal distribution of vector abundance).

Spatio-temporal heterogeneity of vector resistance

Table 2 and Figure 2 show, respectively, global and spatiotemporal descriptive statistics on the resistances of the main vector species collected in the two areas.

A | Temporal distribution

1 dot = 1 entomological survey (all villages combined)
ribbons = spatial variability for the considered survey
dot sizes = % of mosquitoes collected in the considered survey over all the surveys (the biggest, the more)
red dashed horizontal line = overall weighted mean (all surveys considered)

B | Spatial distribution

1 bar = 1 village (all entomological surveys combined)
error bars = temporal variability for the considered village
bar colors = % of mosquitoes collected in the considered village over all the villages (the brighter, the more)
red dashed horizontal line = overall weighted mean (all villages considered)

Figure 2. (Previous page) Spatio-temporal distributions of the resistances of the main vector species collected (panel A : temporal distribution, panel B : spatial distribution). *For behavioural resistances, the y-axis represents th percentage of resistant mosquitoes for the considered survey / village. For physiological resistances, the y-axis represents the allele frequency of the considered mutation for the considered survey / village. Confidence intervals (A : ribbons, B : lineranges) provide indicators of variability of the resistance indicator (A : mean ± standard deviation of the resistance indicator calculated at the village level for the considered entomological survey ; B : mean ± standard deviation of the resistance indicator calculated at the entomological survey level for the considered village). To avoid excessive consideration of small sample sizes, the total number of mosquito collected was represented graphically using the size of dots (A) or the color of the bars (B).*

Table 2. (below) Descriptive statistics for the resistances of the main vector species collected.

Resistance indicator	Study area	Species	n° collected	n° resistant	% resistant	Temporal confidence interval & range	Spatial confidence interval & range
Exophagy	Korhogo	An. gambiae s.l.	56267	31295	56 %	± 2 % (44 – 60)	± 7 %] (38 – 71)
		An. funestus	714	493	69 %	± 7 % (0 – 100)	± 16 % (0 – 100)
	Diébougou	An. coluzzii	1321	577	44 %	± 5 % (38 – 64)	± 9 % (0 – 100)
		An. gambiae s.s.	616	268	44 %	± 7 % (18 – 56)	± 12 % (0 – 75)
		An. funestus	708	250	35 %	± 6 % (19 – 40)	± 8 % (0 – 100)
Early biting	Korhogo	An. gambiae s.l.	56267	1670	3 %	± 1 % (1 – 6)	± 2 % (0 – 10)
		An. funestus	714	92	13 %	± 6 % (0 – 100)	± 12 % (0 – 100)
	Diébougou	An. coluzzii	1321	28	2 %	± 1 % (0 – 4)	± 2 % (0 – 75)
		An. gambiae s.s.	616	19	3 %	± 1 % (0 – 6)	± 3 % (0 – 14)
		An. funestus	708	9	1 %	± 1 % (0 – 2)	± 4 % (0 – 100)
Late biting	Korhogo	An. gambiae s.l.	56267	499	1 %	± 0 % (0 – 1)	± 1 % (0 – 9)
		An. funestus	714	4	1 %	± 1 % (0 – 12)	± 1 % (0 – 7)
	Diebougou	An. coluzzii	1321	46	3 %	± 3 % (0 – 14)	± 3 % (0 – 14)
		An. gambiae s.s.	616	8	1 %	± 3 % (0 – 11)	± 5 % (0 – 100)
		An. funestus	708	82	12 %	± 3 % (0 – 22)	± 10 % (0 – 100)
Kdr-w mutation		An. coluzzii	1321	NA	59 %	± 5 % (55 – 69)	± 8 % (12 – 100)
		An. gambiae s.s.	616	NA	90 %	± 8 % (59 – 100)	± 9 % (68 – 100)
Kdr-e mutation	Diébougou	An. coluzzii	1321	NA	17 %	± 8 % (0 – 43)	± 10 % (0 – 50)
		An. gambiae s.s.	616	NA	4 %	± 4 % (0 – 19)	± 4 % (0 – 17)
Ace-1 mutation		An. coluzzii	1321	NA	2 %	± 1 % (0 – 7)	± 1 % (0 – 6)
		An. gambiae s.s.	616	NA	21 %	± 6 % (11 – 50)	± 8 % (0 – 75)

The columns 'Temporal confidence interval and range' and 'Spatial confidence interval and range' provide indicators of the variability and range of resistance around the overall mean (percentage resistant) respectively in time (i.e. variability between the entomological surveys) and space (i.e. variability between the villages). Format of these columns: standard deviation (minimum – maximum). Computation of standard deviation : to take into account the uneven sample size between entomological surveys (resp. villages) (i.e. to avoid excessive consideration of small / very small sample size), standard deviations for temporal (resp. spatial) variability were extracted by first calculating the resistance indicator at the entomolgial survey (resp. village) level and then computing the weighted standard deviation (weights = number of mosquitoes collected in each entomolgial survey (resp. village)).

Exophagy rates. In the Korhogo area (IC), the overall exophagy rate (% of bites received outdoor) was 56 % for *An. gambiae s.l.* and 69 % for *An. funestus*. The exophagy rate of *An. gambiae s.l.* varied little, both amongst the entomological surveys and the villages (Temporal Standard Deviation (TSD) (see legend of Table 2 for definition) = ± 2 %, Spatial Standard Deviation (SSD) (see legend of Table 2 for definition) = ± 7 %). The exophagy rate of *An. funestus* was more heterogeneously distributed in time and space (TSD = ± 7 %, SSD = ± 16 %). In the Diebougou area (BF), the overall exophagy rate was 44 % for *An. coluzzii*, 44 % for *An. gambiae s.s.* and 35 % for *An. funestus*. For the three species, the exophagy rate varied quite sensibly among the en-

tomological surveys (TSD = ± 5%, ± 7%, ± 6% respectively) and the villages (SSD = ± 9%, ± 12%, ± 8% respectively).

Early and late biting rates. In the Korhogo area (IC), the early biting rate (i.e. % of bites received before 50% of the LLIN users were declared to be under their bednet at night) was 3% for *An. gambiae s.l.* and 13% for *An. funestus*. The early biting rate was overall stable among the surveys and villages for *An. gambiae s.l.* (TSD = ± 1%, SSD = ± 2%) and was more heterogeneously distributed for *An. funestus* (TSD = ± 6%, SSD = ± 12%). The late biting rate (i.e. % of bites received after 50% of the LLIN users were declared to be out of their bednet in the morning) was lower than the early biting rate : 1% for both *An. gambiae s.l.* and *An. funestus* (only 4 late-bites for *An. funestus*) and was overall stable among the surveys and villages (TSD = ± 0% and SSD = ± 1% for *An. gambiae s.l.*). In the Diébougou area (BF), the early biting rate was respectively 2%, 3% and 1% for *An. coluzzii*, *An. gambiae s.s.* and *An. funestus*. The early biting rate was overall stable among the surveys (TSD = ± 1% for the three species) and to some extent more heterogeneous among the villages (SSD = ± 2%, ± 3%, ± 4% respectively). The late biting rate was respectively 3%, 1% and 12% for *An. coluzzii*, *An. gambiae s.s.* and *An. funestus*. Late biting rates were more heterogeneously distributed than early biting rates, both among the surveys (TSD = ± 3% for the three species) and villages (SSD = ± 3%, ± 5%, ± 10% respectively).

Allele frequencies of kdr-e, kdr-w, ace-1 mutations. In the BF area, the allele frequency of the *kdr-w* mutation was 90% for *An. gambiae s.s.* and 59% for *An. coluzzii*. It varied to some extent among the surveys and villages (for *An. gambiae s.s.* : TSD = 8%, SSD = 9% ; for *An. coluzzii* : TSD = 5%, SSD = 8%). The allele frequency of the *kdr-e* mutation was 4% for *An. gambiae s.s.* and 17% for *An. coluzzii*. For *An. gambiae s.s.*, it remained low among the surveys and villages (TSD = SSD = 4%) and for *An. coluzzii*, it varied more sensibly (TSD = 8%, SSD = 10%). The allele frequency of the *ace-1* mutation was 21 % for *An. gambiae s.s.* and 2% for *An. coluzzii*. For *An. gambiae s.s*, it varied sensibly among the surveys and villages (TSD = 6%, SSD = 8%), and for *An. coluzzii* it was overall stably low (TSD = SSD = 1%).

Dependent variables excluded from the modeling process

Seven of the original twenty-one dependent variables were excluded before statistical modeling due to the very small size of their 'resistant' class (see Table 2) :

- early-biting in BF for the three species,
- late-biting in BF for *An. coluzzii* and *An. gambiae s.s.*,
- late-biting in IC for *An. funestus*,
- ace-1 in BF for *An. coluzzii*.

Associations between physiological resistance and environmental variables

For the remaining five models of physiological resistance in the Diébougou area (BF), Figure 3 shows the PDPs of the independent variables retained in the modeling workflow. For the GLMMs, numerical values of odd-ratios, 95% confidence intervals, and p-values are provided in Additional file 4.

Figure 3. (Previous page) Results of the statistical models of probability of physiological resistance in the malaria vectors. *For each model, the top plot shows the explanatory power (R^2) and predictive power (AUC) of respectively the GLMM and the RF model. The other plots show the predicted probabilities of collecting a resistant vector across available values of each independent variable, holding everything else in the model equal (yellow line : probability predicted by the GLMM model ; green line : probability predicted by the RF model) (short reminder : vector control and crops variables were forced-in, and the other variables were retained only if they improved the AIC of the model. In addition, for the GLMM models, the other variables were plotted only if their p-value was < 0.05. For the RF models, the predicted probability (i.e. green line) was plotted only if the AUC of the model was > 0.6 and the range of predicted probabilities of resistance for the considered variable was > 0.05). In these plots, the y-axis represents the probability for an allele to be resistant. The red horizontal dashed line represents the overall rate of resistance (see Table 2). The p-values of the GLMMs are indicated through the stars : * : p < 0.05, ** p < 0.01, *** p < 0.001. The grey squares distributed along the x-axis at the top and bottom of each plot represent the measured values available in the data (the darker the square, the more the number of observations) (NB : for atmospheric pressure, the values in the x-axis are centered around the mean)*

Associations with variables encoding vector control interventions. No statistically significant association
was found between the likelihood of collecting an *Anopheles* carrying any of the target-site mutations and
the type of VC intervention (LLIN + complementary tool compared to LLIN only) within the time frame of the
study. However, the likelihood of collecting a host-seeking *An. gambiae s.s.* or *An. coluzzii* carrying a resistant
kdr-e allele increased with the time since LLIN distribution, and as well with the % of users of LLINs in the
village population. Regarding the others target-site mutations (*kdr-w* or *ace-1*), the likelihood of collecting a
host-seeking *Anopheles* carrying them did not increase with the time since LLIN distribution.
456
457   ***Associations with variables encoding crops.*** No statistically significant association was found between the
458 likelihood of collecting a host-seeking *Anopheles* carrying any of the target-site mutations and the % of land-
459 scape occupied by crop fields (cotton, rice, or other crops) in a 2 km-wide buffer area around the collection point.
460
Associations with variables encoding micro-climate at the time (hour) of foraging activity. Positive asso-
ciations were found between the likelihood of collecting a host-seeking *An. coluzzii* carrying the *kdr-e* mutation
and atmospheric pressure, humidity and temperature at the time of collection, as well as that of collecting an
An. gambiae s.s. carrying the *kdr-e* mutation and atmospheric pressure at the time of collection. A negative
association was found between the likelihood of collecting a host-seeking *An. gambiae s.s.* carrying the *kdr-w*
mutation and humidity at the time of collection.
467
Associations with variables encoding meteorological conditions during the month preceding collection.
Negative associations were found between the likelihood of collecting a host-seeking : *An. coluzzii* carrying the
kdr-w mutation and cumulated rainfall, *An. gambiae s.s.* carrying the *kdr-w* mutation and both cum. rainfall and
mean diurnal temperatures, *An. coluzzii* carrying the *kdr-e* mutation and mean nocturnal temperatures, *An.*
gambiae s.s. carrying *ace-1* mutation and both mean diurnal and nocturnal temperatures during the month
preceding collection. A positive association was found between the likelihood of collecting a host-seeking *An.*
coluzzii carrying the *kdr-e* mutation and cumulated rainfall.

Associations between behavioural resistance and environmental variables

476 For the remaining nine models of behavioural resistance, Figure 4 shows the PDPs of the independent variables
477 retained in the modeling workflow. For the GLMMs, numerical values of odd-ratios, 95% confidence intervals
478 and p-values are provided in Additional file 4.

Figure 4. (Previous page) Results of the statistical models of probability of behavioural resistance in the malaria vectors. *For each model, the top plot shows the explanatory power (R^2) and predictive power (AUC) of respectively the GLMM and the RF model. The other plots show the predicted probabilities of collecting a resistant vector across available values of each independent variable, holding everything else in the model equal (yellow line : probability predicted by the GLMM model ; green line : probability predicted by the RF model) (short reminder : vector control variables were forced-in, and the other variables were retained only if they improved the AIC of the model. In addition, other variables were plotted only if their p-value was < 0.05. For the RF models, the predicted probability (i.e. green line) was plotted only if the AUC of the model was > 0.6 and the range of predicted probabilities of resistance for the considered variable was > 0.05). In these plots, the y-axis represents the probability for a mosquito to be resistant. The red horizontal dashed line represents the overall rate of resistance (see Table 2). The p-values of the GLMMs are indicated through the stars * : p < 0.05, ** p < 0.01, *** p < 0.001. The grey squares distributed along the x-axis at the top and bottom of each plot represent the measured values available in the data (the darker the square, the more the number of observations) (NB : for atmospheric pressure, the values in the x-axis are centered around the mean).*

Associations with variables encoding vector control interventions. No statistically significant association
was found between the likelihood of collecting an exophagic, early- or late- biting *Anopheles* and neither the
type of VC intervention (LLIN + complementary tool compared to LLIN only) nor the time since LLIN distribution
within the time frame of the study.

Associations with variables encoding host availability. In the Korhogo area (IC), the likelihood of exophagy
of *An. gambiae s.s.* slightly increased with the % of the population under an LLIN at the time of collection. The
likelihood of early-biting of *An. gambiae s.s.* increased with the % of users of LLINs in the village population. In
the Diébougou (BF) area, the likelihood of exophagy of *An. funestus* increased with the % of the population
under an LLIN at the time of collection.

Associations with variables encoding landscape. In the Korhogo area (IC), the likelihood of exophagy of
An. funestus increased with increasing distance to the edge of the village. The likelihood of early-biting of *An.*
gambiae s.s. decreased with increasing distance to the edge of the village. In the Diébougou (BF) area, the
likelihood of exophagy of *An. coluzzii* increased with increasing distance to the nearest stream.

Associations with variables encoding micro-climate at the time (hour) of foraging activity. In the Korhogo
area (IC), the likelihood of exophagy of *An. gambiae s.s.* decreased when humidity indoors increased and when
humidity got relatively higher indoors compared to outdoors. In addition, it increased when luminosity got
relatively higher indoors compared to outdoors. In the Diébougou area (BF), the likelihood of exophagy of *An.*
funestus increased when temperature or humidity got relatively higher indoors compared to outdoors.

Associations with variables encoding meteorological conditions on the day or night of collection. Positive
associations were found between the likelihood of : exophagy of *An. coluzzii* and rainfall (BF area), early-biting
of *An. gambiae s.s.* and temperature (IC area), late-biting of *An. gambiae s.s.* and both rainfall and temperature
(IC area), late-biting of *An. funestus* and temperature (BF area). A negative association was found between the
likelihood of exophagy of *An. gambiae s.s.* and rainfall (IC area).

Associations with variables encoding meteorological conditions during the month preceding collection.
Negative associations were found between the likelihood of : exophagy of *An. gambiae s.s.* and both cumulated
rainfall and mean diurnal temperatures (IC area), exophagy of *An. coluzzii* and mean nocturnal temperatures
(BF area), late biting of *An. gambiae s.s.* and mean nocturnal temperature (IC area). A positive association was
found between the likelihood of exophagy of *An. gambiae s.s.* and mean nocturnal temperatures (BF area).

Associations with variables encoding physiological resistances. As a reminder, the genotypes for the target-
site mutations of individual collected mosquitoes were introduced as independent variables in the behavioural
resistance models in the Diébougou area (BF). Here, these variables were not retained in the variable selection
procedure, i.e. no statistically significant association was found between any of the behavioural resistance
indicator and *kdr-w*, *kdr-e*, or *ace-1* mutations.

Explanatory and predictive power of the statistical models

521 Additional figure 5 provides boxplots of observed resistance status vs. predicted probabilities by each model.

522
523 ***Exophagy.*** For the models of exophagy, the explanatory power of the GLMM models was : 'very weak' for *An.*
524 *gambiae s.s.* in the Korhogo area (IC), 'moderate' for *An. funestus* in the Korhogo area (IC)', weak' for *An. funestus*,
525 *An. coluzzii* and *An. gambiae s.s.* in the Diébougou area (BF). The predictive power of the RF models of exophagy
526 was 'very weak' for all the species in the two study areas.

527
Early and late biting. For the models of early biting, the explanatory power of the GLMM models was 'weak'
for both *An. gambiae s.s.* and *An. funestus* in the Korhogo area (IC). For the models of late biting, the explanatory
power of the GLMM was 'weak' for *An. gambiae s.s.* in the Korhogo area (IC) and 'substantial' for *An. funestus* in
the Diébougou area (BF). The predictive power of the RF models of early and late biting was 'very weak' for all
species in the two study areas, except for the model of late biting of *An. gambiae s.s.* in the Korhogo area (IC)
for which it was 'weak'.

534
Kdr-w, kdr-e, ace-1. For the *kdr-w* mutation in the Diébougou area (BF), the explanatory power of the GLMM
models was 'weak' for *An. coluzzii* and 'substantial' for *An. gambiae s.s.* ; and the predictive power of the RF
models was 'weak' for *An. coluzzii* and 'moderate' for *An. gambiae s.s.* For the *kdr-e* mutation in the Diébougou
area (BF), the explanatory power of the GLMM models was 'substantial' for both *An. coluzzii* and *An. gambiae s.s.*
; and the predictive power of the RF models was 'moderate' for *An. coluzzii* and 'weak' for *An. gambiae s.s.* For
the *ace-1* mutation in the Diébougou area (BF), the explanatory power of the GLMM models was 'weak' for *An.*
gambiae s.s. ; and the predictive power of the RF model was 'very weak'.

Discussion

In this work, we studied indicators of physiological and behavioural resistances of several malaria vectors in
rural West-Africa at a fine spatial scale (approximately the extent of a health district), using longitudinal data
collected in two areas belonging to two different countries, respectively 27 and 28 villages per area, and across
1.25 to 1.5 year. The objectives were to describe the spatial and temporal heterogeneity of vector resistance,
and to better understand their drivers, at scales that are consistent with operational action. To our knowledge,
our work is the first studying the heterogeneity of vector resistance at such fine spatial scale with such a large
dataset of mosquito collection and potential drivers of resistance. In this discussion, we first use our results
to provide elements of answers to the questions raised in introduction of this article. We then discuss some
implications of the findings for the management of vector resistance in our areas.

Physiological resistances : potential drivers and spatiotemporal heterogeneity

553 The main drivers of physiological resistances are insecticides, used either in public health for vector control or
554 in agriculture (see Introduction). In this study, we found that the probability of collecting a host-seeking *An.*
555 *gambiae s.s.* or *An. coluzzii* in the Diébougou area carrying a *kdr-e* resistant allele significantly increased with
556 both the time since LLIN distribution and the % of LLIN users in the village population. In contrast, there was no
557 significant association between any of the target-site mutations and any of the crop-related variable. This could

indicate that **within the spatiotemporal frame of our study, the development of the kdr-e mutation in the vector population was more likely due to insecticides used in public health than pesticides used in agriculture**. In Burkina Faso, pesticides are widely used for cotton and sugar cane (Ouedraogo et al., 2011), but only in lesser proportions in market gardening and cereal production (maize and rice are the only cereals that are treated to a significant extent (MERSI et al., 2016)). Here, in the 2-km wide buffer zones around our collection points crops occupied up to 40 % of the total land, but were mainly made of leguminous crops, millet, sorghum, with cotton and rice being only marginally present. Hence, pesticides are likely not much used (field surveys regarding the use of pesticides by the farmers could confirm this hypothesis). This could explain the absence of association between target-site mutations and the crops-related variables. Noteworthy, the *kdr-w* and *ace-1* mutations did not increase significantly with the time since LLIN distribution. The absence of increase of the *kdr-w* mutation may be explained by its very high baseline allelic frequencies ; while that of the *ace-1* mutation may be explained by the type of insecticide used to impregnate the LLINs - deltamethrin, which does not select the ace-1 mutation).

The statistical models captured many associations between the likelihood of collecting a physiologically resistant *Anopheles* and the variables encoding weather, both during the month preceding collection and at the hour of collection. As stated previously, weather may impact the fitness or the activity of mosquitoes carrying resistant genotypes ; and may therefore *in fine* impact the probability of collecting a physiologically resistant mosquito. Here, the associations that were captured could hence traduce **biological costs associated with target-site mutations, both in terms of fitness and activity**, as found elsewhere for other mosquito species (Kliot and Ghanim, 2012). Regarding fitness, we found that the likelihood of collecting a host-seeking mosquito (*An. gambiae s.s.* or *An. coluzzii*) carrying a mutated allele, overall, decreased (to varying extents depending on the species and mutation) in the hot seasons (i.e. when diurnal or nocturnal temperatures during the month preceding collection increased). Carrying a kdr mutation might be associated with a reduced ability to seek out optimal temperatures, potentially resulting in a decreased longevity, fecundity, or ovarian development rates (Foster et al., 2003). Regarding activity, we observed that the likelihood of collecting a mosquito carrying a mutated allele (for the *kdr-e* mutation) decreased when atmospheric pressure, humidity, or temperature at the hour of collection got lower ; implying that mosquitoes carrying the *kdr-e* mutation could be less active in colder or drier conditions, or when atmospheric pressure is lower.

Lastly, we observed that the allelic frequencies of the target-site mutations, within each vector species and for each mutation, were overall quite stable across the villages and seasons within the spatiotemporal frame of the study. At larger spatial and temporal scales, physiological resistances were found more heterogeneous (Moyes et al., 2020). In our study, such homogeneity might be due to a relative homogeneity in space and time of the main determinants of physiological resistance (access and use of insecticide-based vector control interventions). The relative seasonal homogeneity might traduce that fitness costs, despite their existence, might be limited within the range of meteorological conditions in our area.

Behavioural resistances : potential drivers and spatiotemporal heterogeneity

An important and pending question is the genetic (constitutive) or plastic (inducible) nature of behavioual resistances (see Introduction). In this study, we found no statistically significant association between any of the indicators of behavioural resistance and neither the time since LLIN distribution nor the VC tool implemented. There was hence no evidence of growing frequencies of behavioural resistances (exophagy, early- and late-biting) in response to vector control within the 1.25 to 1.5 years of this study, i.e. **no clear indication of constitutive resistance**. Nonetheless, some of the associations captured by the models, as well as a comparison of the measured resistance rates with that of previous studies in the same countries, suggest that there may still be a genetic component in mosquito foraging behaviour. First, we found many statistically significant associations between the likelihood of collecting a behaviourally resistant *Anopheles* and the

meteorological conditions during the month preceding collection. This could mean that there might exist a fitness cost to behavioural resistance, associated with (i) maternal or paternal effects or (ii) development or survival at the larval stage (see Taconet, Porciani, et al. (2021)), that support (at least for the option (ii)) the hypothesis of genetic bases for some host-seeking behavioural phenotypes. Second, the exophagy rates of the main malaria vectors in both areas (on average, 41% in the Diébougou area and 56% in the Korhogo area) were substantially higher than those, overall, historically reported for these species ; in general in Africa (Huho et al., 2013; Sinka et al., 2010) and especially in Burkina Faso (Sherrard-Smith et al., 2019). Such high levels of outdoor biting, in comparison with past levels, suggest that behavioural adaptations may be ongoing in the study areas, most probably in response to the widespread and prolonged use of insecticide-based vector control tools. The hypothesis of a hereditary component in the behavior of malaria vectors (at least for the biting hour) is supported by a recent study which has observed, for *Anopheles arabiensis* in Tanzania, that F2 from early-biting F0 (grandmothers) were - to some extent - more likely to bite early than F2 from mid or late-biting F0 (Govella et al., 2021). In our study, the absence of significant association between the probability of behavioural resistances and insecticide-related variables might be due to the relatively short length of the study. In fact, mosquito behaviours are likely complex multigenic traits (Main et al., 2016) and might therefore respond slowly to selection (at least, slower than target-site mutations, which are linked to single genes and may hence respond rapidly and efficiently to selection). In a recent two-years long longitudinal study in another area of Burkina Faso, the authors also observed an absence of association between time and behavioural resistance (for early biting, exophagy and exophily) (Sanou et al., 2021) ; however, such associations were actually found in a four-years long study in Tanzania (for exophagy) (Kreppel et al., 2020). Long-term monitoring of vector behaviour, particularly in areas with a long history of use of insecticides in public health, is critical to better understand the biological mechanisms underlying behavioural resistances, to potentially assess their development rate, and more broadly to assess residual malaria transmission risk (Durnez and Coosemans, 2013; Kreppel et al., 2020; Sanou et al., 2021).

Weather can impact the fitness of possible genotypes associated with resistant behavioural phenotypes, but may also influence the activity of these phenotypes (see Introduction). Here, we found many associations between mosquito host-seeking behaviour and variables representing meteorological conditions on the day or at the hour of collection. For instance, the probability for an *An. gambiae s.s.* to be collected outdoor in the Korhogo area increased when the air indoor was dry, or when the air outdoor became relatively more humid than indoor. Likewise, in the Diébougou area, the probability for an *An. funestus* to be collected outdoor increased when the air outdoor became relatively cooler than indoor. These observations are consistent with the hypothesis of mosquitoes shifting from indoor to outdoor host-seeking in case of desiccation-related mortality risk indoors, as observed and discussed elsewhere (Kessler and Guerin, 2008; Kreppel et al., 2020; Ngowo et al., 2017). The meteorological conditions seemed to cause not only spatial, but also temporal shifts in host-seeking activity. For instance, we found that the probability of collecting a late-biting *An. gambiae s.s.* in the Korhogo area increased when the average noctural temperature increased. Several associations also suggest that some malaria vectors may modify their behaviour in response to environmental variation that reduces host availability, as hypothesized elsewhere (Durnez and Coosemans, 2013). For instance, the likelihood of collecting an *An. gambiae s.s.* (in the Korhogo area) or an *An. funestus* (in the BF area) outdoor increased at hours when people were protected by their LLINs. Likewise, the likelihood of collecting an early-biting *An. gambiae s.s.* in the Korhogo area increased when the % of LLIN users in the village increased. Altogether, all these associations suggest that in our study areas **mosquito foraging behaviour is driven – to a certain extent - by environmental conditions at the time of foraging activity**, i.e. that vectors likely modify their time or place of biting according to climatic conditions or host availability. The many associations that were captured here in field conditions could be further tested experimentally, to quantify their effect more precisely and validate the underlying biological hypothesis.

Although many significant associations between environmental parameters and foraging behaviours have been captured by the models, their explanatory and predictive powers were overall weak. A low explanatory power can indicate either i) that variations in the dependent variable (here, individual vector resistance) are only marginally caused by the independent variables or ii) that the statistical model does not capture properly the true nature of the underlying relationships between the studied effect and its drivers (Karpatne et al., 2017) (e.g. a linear regression cannot, by definition, capture non-linear relationships that might exist in nature). Here, we minimized the risk of omitting important, complex associations by using, in addition to the binomial regression model, a machine-learning model (namely a random forest) that is inherently able to capture complex patterns contained in the data (e.g. non-linear relationships, interactions) (Breiman, 2001a). Still, the models had low predictive powers. Altogether, these results indicate that very likely, despite the amount, granularity and diversity of potential factors measured and introduced in the models, most of the factors driving the individual host-seeking behaviours of the mosquitoes were not introduced in the models. Another possibility could be that some of our independent variables did not represent the actual "reality" in the field (e.g. the distance to the nearest stream is not necessary an ideal proxy for the distance to the breeding site). Nevertheless, since we used a wide range of variables encoding the environmental conditions at the time of foraging activity, we can hypothesize that within the spatiotemporal frame of the study, **mosquito foraging behaviour was only marginally driven by environmental variations**. This leaves the floor to other factors, like genetics (see above), learning, or randomness.

To test whether physiological resistance impacts the behaviour of host-seeking mosquitoes, we introduced in the behaviour resistance models of *An. coluzzii* and *An. gambiae s.s.* in the Diébougou area two variables encoding the genotypes for respectively the *kdr-w* and *kdr-e* mutations. No statistically significant association was found. In other words, **we could not find, in the field, a behavioural phenotype (among those studied, i.e. exophagy, early- and late-biting) associated with a genotype for one of the target-site mutations**. The only study, to our knowledge, having investigated the relationship between the kdr mutation and biting time or location in the field has also reported no statistically significant association between these two mechanisms of resistance to insecticide (Djènontin et al., 2021). Noteworthy, in our study, there was few variabilities in the genotypes of the collected mosquitoes (i.e. few homozygote susceptible mosquitoes captured, particularly for the *kdr-w* mutation), making it unfavorable to detect assocations between physiological and behavioural resistances. In the Korhogo area, such analysis could not be performed because physiological resistance was not available at the individual mosquito level.

Finally, we observed that the behavioural resistance rates for each vector species in each health district were, overall, relatively homogeneous across the villages and seasons within the spatiotemporal frame of the study (as for physiological resistances). This could mean that the overall dynamics of behavioural resistance occur at broader spatial and temporal scales than those of our study. At larger scales (i.e. among countries and across years in Africa), exophagy rates of *Anopheles* mosquitoes seem, actually, to be more variable (Sherrard-Smith et al., 2019).

Implications of the findings for the management of vector resistance in the study areas

Long-lasting insecticidal nets have undoubtedly played a major role in reducing malaria cases throughout Africa, thanks both to their barrier and killing effects. More locally, we highlighted the efficacy of their barrier role in the Diébougou area by showing that, for their users, they prevented more than 80% of *Anopheles* bite exposure in the area (Soma, Zogo, Taconet, et al., 2021). Despite these successes, our study adds to the growing body of evidence that the insecticides they are impregnated with are responsible for the rise of physiological resistances in the malaria vectors populations. We also highlighted that in response to an LLIN distribution, physiological resistance seems to grow quite rapidly in a susceptible population. Besides the

development of physiological resistance, comparison with historical data suggests that the vectors may also be progressively changing their feeding behaviour to avoid the effects of the insecticides - although there was no clear evidence of this in the strict context of this study. Such trends in vector resistance may have an important epidemiological impact (Sherrard-Smith et al., 2019). Altogether, these results show, if still necessary, that we urgently need to think more strategically about our use of insecticides in public health tools in our areas. Switching to alternative insecticides, rotating or mixing insecticides, using current or novel insecticides in vector control tools others than long-lasting nets, entirely removing the insecticides from the vector control toolbox, or fostering the use of insecticidal-free tools, are all actions that could be envisaged (Paaijmans and Huijben, 2020). Burkina Faso has, actually, distributed LLINs that mixes pyrethroid with Piperonyl butoxide (PBO) in its last universal LLIN distribution, in 2019.

Here, we observed that both behavioural and physiological resistances of mosquitoes were quite stable across the villages and seasons within the spatiotemporal frame of the study. This contrasts with their biting rates, which was found, in another study (Taconet, Porciani, et al., 2021), highly variable across the villages, seasons, and amongst the species. At small spatiotemporal scales, this calls for different strategies for respectively vector control (interventions aiming at reducing the human-vector contact) and resistance management (interventions aiming at reducing the development of physiological or behavioural resistance) at such spatiotemporal scales. While vector control plans should be very locally-tailored (species-, season-, and village-specific) (Taconet, Porciani, et al., 2021), resistance management strategies would probably not gain much in being adapted to the season or village within our areas. In other words, while resistance management plans are undoubtedly urgently needed, there is no compelling evidence – in the current state of the knowledge - that they should be tailored at very fine scales (village, season). Noteworthy, no entomological survey was performed during the high rainy season (July to September), at the known mosquito abundance and malaria transmission peaks. It would be worth collecting mosquitoes at this season to confirm the observed resistance rates.

Conclusion

Less than a decade after the first universal LLIN distribution, malaria vectors in two areas of rural West-Africa seem to be growingly adapting to avoid or circumvent the lethal effects of insecticides used in control interventions. After an LLIN distribution, rapid widespread of physiological resistance occurring in tandem with probable lower acting behavioral adaptations, are very likely contributing to the erosion of the insecticides impact. In an attempt to better understand the drivers of the intensity and spatio-temporal heterogeneity of physiological and behavioural resistance in malaria vectors, at the scale of a rural health district over a period of 1.5 years, we have mainly (i) shown that resistance (both physiological and behavioural) was quite homogeneous across the villages and seasons at theses scales, and (ii) hypothesized that at these spatiotemporal scales, vector resistance seemed to be only marginally driven by environmental factors other than those linked to insecticide use in current vector control. We believe that without waiting to understand precisely the underlying drivers, mechanisms, and rates of development of resistances, the malaria control community needs to think very strategically about the use and usefulness of current and novel insecticide-based control interventions.

Acknowledgements

We thank populations of the villages for their kind support and collaboration. We also thank all the field and laboratory staff for their strong commitment to the REACT project.

Abbreviations

AIC: Akaike Information Criterion; AUC: Area under the ROC Curve; BF: Burkina Faso; CV: Cross-Validation; GLMM: Generalized Linear binomial Mixed-effect Model; HLC: Human Landing Catch; IC: Ivory Coast; IRS: Indoor Residual Spraying; LLIN: Long Lasting Insecticide Nets; ML: Machine Learning; RF: Random Forest; SD : Standard Deviation; SR : Spatial Resolution; SSD: Spatial Standard Deviation; TSD: Temporal Standard Deviation; TR: Temporal Resolution; VC: Vector Control;

Data, scripts, code and supplementary information availability

Data and scripts are available online: https://doi.org/10.23708/LV8GEW (Taconet, DD Soma, Zogo, Mouline, Simard, Koffi, et al., 2023)

Supplementary information are available online: https://doi.org/10.23708/VJEEMU (Taconet, DD Soma, Zogo, Mouline, Simard, Koffi Amanan, et al., 2023)

Ethics approval and consent to participate

Ethical clearance for the study was granted by the National ethics committee (No. 063/MSHP/CNER-kp) in Côte d'Ivoire and by the Institutional Ethics Committee of the Institut de Recherche en Sciences de la Santé (No. A06/2016/CEIRES) in Bukina Faso. We received community agreement before the beginning of the study, and we obtained written informed consent from all the mosquito collectors and supervisors. Yellow fever vaccines were administered to all the field staff. Collectors were treated free of charge when they were diagnosed with malaria during the study period according to WHO recommendations. They were also free to withdraw from the study at any time without any consequences.

Conflicts of interest disclosure

The authors declare that they comply with the PCI rule of having no financial conflicts of interest in relation to the content of the article.

Funding

This work was part of the REACT project funded by the French Initiative 5%—Expertise France (no. 15SANIN213), and the ANORHYTHM project funded by the French National Research Agency (no. ANR-16-CE35-008). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. PT was supported by the French Institute of Research for Sustainable Development (IRD) through an international volunteer fellowship.

References

Auguie B (2017). *gridExtra: Miscellaneous Functions for "Grid" Graphics*.

Bartoń K (2020). *MuMIn: Multi-Model Inference*.

Bhatt S, DJ Weiss, E Cameron, D Bisanzio, B Mappin, U Dalrymple, KE Battle, CL Moyes, A Henry, PA Eckhoff, EA Wenger, O Briët, MA Penny, TA Smith, A Bennett, J Yukich, TP Eisele, JT Griffin, CA Fergus, M Lynch, F Lindgren, JM Cohen, CLJ Murray, DL Smith, SI Hay, RE Cibulskis, and PW Gething (2015). The effect of malaria control on Plasmodium falciparum in Africa between 2000 and 2015. en. *Nature* 526, 207–211. https://doi.org/10.1038/nature15535.

Bolker B and D Robinson (2020). *broom.mixed: Tidying Methods for Mixed Models*.

Breiman L (2001a). Random forests. *Machine learning* 45, 5–32.

— (2001b). Statistical Modeling: The Two Cultures. *Statistical Science* 16, 199–215.

Brooks ME, K Kristensen, KJv Benthem, A Magnusson, CW Berg, A Nielsen, HJ Skaug, M Maechler, and BM Bolker (2017). glmmTMB Balances Speed and Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. *The R Journal* 9, 378–400.

Carrasco D, T Lefèvre, N Moiroux, C Pennetier, F Chandre, and A Cohuet (2019). Behavioural adaptations of mosquito vectors to insecticide control. en. *Current Opinion in Insect Science* 34, 48–54. https://doi.org/10.1016/j.cois.2019.03.005.

Center NGESDAIS (2019). GPM IMERG Final Precipitation L3 1 day 0.1 degree x 0.1 degree V06. type: dataset. https://doi.org/10.5067/GPM/IMERGDF/DAY/06.

Chandre F, F Darriet, S Manguin, C Brengues, P Carnevale, and P Guillet (1999). Pyrethroid cross resistance spectrum among populations of Anopheles gambiae s.s. from Côte d'Ivoire. ENG. *Journal of the American Mosquito Control Association* 15, 53–59.

Chicco D (2017). Ten quick tips for machine learning in computational biology. en. *BioData Mining* 10. https://doi.org/10.1186/s13040-017-0155-3.

Cohen J (2013). *Statistical Power Analysis for the Behavioral Sciences*. en. 0th ed. Routledge. ISBN: 978-1-134-74270-7. https://doi.org/10.4324/9780203771587.

Corbel V and R N'Guessan (2013). Distribution, Mechanisms, Impact and Management of Insecticide Resistance in Malaria Vectors: A Pragmatic Review. en. *Anopheles mosquitoes - New insights into malaria vectors*. https://doi.org/10.5772/56117.

Davidson G (1957). Insecticide Resistance in Anopheles Sundaicus. en. *Nature* 180, 1333–1335. https://doi.org/10.1038/1801333a0.

Diop MM, N Moiroux, F Chandre, H Martin-Herrou, P Milesi, O Boussari, A Porciani, S Duchon, P Labbé, and C Pennetier (2015). Behavioral Cost & Overdominance in Anopheles gambiae. en. *PLOS ONE* 10. Ed. by Lazzari CR, e0121755. https://doi.org/10.1371/journal.pone.0121755.

Diop MM, F Chandre, M Rossignol, A Porciani, M Chateau, N Moiroux, and C Pennetier (2021). Sub-lethal insecticide exposure affects host biting efficiency of Kdr-resistant Anopheles gambiae. en. *Peer Community Journal* 1, e28. https://doi.org/10.24072/pcjournal.15.

Djènontin A, A Bouraima, C Soares, S Egbinola, and G Cottrell (2021). Human biting rhythm of Anopheles gambiae Giles, 1902 (Diptera: Culicidae) and sleeping behaviour of pregnant women in a lagoon area in Southern Benin. *BMC Research Notes* 14, 200. https://doi.org/10.1186/s13104-021-05615-7.

Durnez L and M Coosemans (2013). Residual Transmission of Malaria: An Old Issue for New Approaches. en. In: *Anopheles mosquitoes - New insights into malaria vectors*. Ed. by Manguin S. InTech. ISBN: 978-953-51-1188-7. https://doi.org/10.5772/55925.

Foster SP, S Young, MS Williamson, I Duce, I Denholm, and GJ Devine (2003). Analogous pleiotropic effects of insecticide resistance genotypes in peach–potato aphids and houseflies. en. *Heredity* 91, 98–106. https://doi.org/10.1038/sj.hdy.6800285.

Friedman JH and BE Popescu (2008). Predictive learning via rule ensembles. en. *The Annals of Applied Statistics* 2, 916–954. https://doi.org/10.1214/07-AOAS148.

Gatton ML, N Chitnis, T Churcher, MJ Donnelly, AC Ghani, HCJ Godfray, F Gould, I Hastings, J Marshall, H Ranson, M Rowland, J Shaman, and SW Lindsay (2013). THE IMPORTANCE OF MOSQUITO BEHAVIOURAL ADAPTATIONS TO MALARIA CONTROL IN AFRICA. en. *Evolution* 67, 1218–1230. https://doi.org/10.1111/evo.12063.

Govella NJ, PCD Johnson, GF Killeen, and HM Ferguson (2021). *Heritability and phenotypic plasticity of biting time behaviors in the major African malaria vector Anopheles arabiensis*. en. preprint. Evolutionary Biology. https://doi.org/10.1101/2021.05.17.444456.

Greenwell BM (2017). pdp: An R Package for Constructing Partial Dependence Plots. *The R Journal* 9, 421–436.

GSFC PPS(AN (2019). GPM IMERG Final Precipitation L3 Half Hourly 0.1 degree x 0.1 degree V06. Type: dataset. https://doi.org/10.5067/GPM/IMERG/3B-HH/06.

Hay GJ and G Castilla (2008). Geographic Object-Based Image Analysis (GEOBIA): A new name for a new discipline. en. In: *Object-Based Image Analysis*. Ed. by Blaschke T, Lang S, and Hay GJ. Berlin, Heidelberg: Springer Berlin Heidelberg, pp. 75–89. ISBN: 978-3-540-77057-2 978-3-540-77058-9. https://doi.org/10.1007/978-3-540-77058-9_4.

Hemingway J, H Ranson, A Magill, J Kolaczinski, C Fornadel, J Gimnig, M Coetzee, F Simard, DK Roch, CK Hinzoumbe, J Pickett, D Schellenberg, P Gething, M Hoppé, and N Hamon (2016). Averting a malaria disaster: will insecticide resistance derail malaria control? en. *The Lancet* 387, 1785–1788. https://doi.org/10.1016/S0140-6736(15)00417-1.

Hersbach H, B Bell, P Berrisford, S Hirahara, A Horányi, J Muñoz-Sabater, J Nicolas, C Peubey, R Radu, D Schepers, A Simmons, C Soci, S Abdalla, X Abellan, G Balsamo, P Bechtold, G Biavati, J Bidlot, M Bonavita, G Chiara, P Dahlgren, D Dee, M Diamantakis, R Dragani, J Flemming, R Forbes, M Fuentes, A Geer, L Haimberger, S Healy, RJ Hogan, E Hólm, M Janisková, S Keeley, P Laloyaux, P Lopez, C Lupu, G Radnoti, P Rosnay, I Rozum, F Vamborg, S Villaume, and J Thépaut (2020). The ERA5 global reanalysis. en. *Quarterly Journal of the Royal Meteorological Society* 146, 1999–2049. https://doi.org/10.1002/qj.3803.

Hien AS, DD Soma, O Hema, B Bayili, M Namountougou, O Gnankiné, T Baldet, A Diabaté, and KR Dabiré (2017). Evidence that agricultural use of pesticides selects pyrethroid resistance within Anopheles gambiae s.l. populations from cotton growing areas in Burkina Faso, West Africa. en. *PLOS ONE* 12. Ed. by Carvalho LH, e0173098. https://doi.org/10.1371/journal.pone.0173098.

Huho B, O Briët, A Seyoum, C Sikaala, N Bayoh, J Gimnig, F Okumu, D Diallo, S Abdulla, T Smith, and G Killeen (2013). Consistently high estimates for the proportion of human exposure to malaria vector populations occurring indoors in rural Africa. en. *International Journal of Epidemiology* 42, 235–247. https://doi.org/10.1093/ije/dys214.

Karpatne A, G Atluri, JH Faghmous, M Steinbach, A Banerjee, A Ganguly, S Shekhar, N Samatova, and V Kumar (2017). Theory-Guided Data Science: A New Paradigm for Scientific Discovery from Data. *IEEE Transactions on Knowledge and Data Engineering* 29, 2318–2331. https://doi.org/10.1109/TKDE.2017.2720168.

Kessler S and PM Guerin (2008). Responses of Anopheles gambiae, Anopheles stephensi, Aedes aegypti, and Culex pipiens mosquitoes (Diptera: Culicidae) to cool and humid refugium conditions. en. *Journal of Vector Ecology* 33, 145–149. https://doi.org/10.3376/1081-1710(2008)33[145:ROAGAS]2.0.CO;2.

Killeen GF (2014). Characterizing, controlling and eliminating residual malaria transmission. *Malaria Journal* 13, 330. https://doi.org/10.1186/1475-2875-13-330.

Kirby M and S Lindsay (2004). Responses of adult mosquitoes of two sibling species, *Anopheles arabiensis* and *A. gambiae* s.s. (Diptera: Culicidae), to high temperatures. en. *Bulletin of Entomological Research* 94, 441–448. https://doi.org/10.1079/BER2004316.

Kliot A and M Ghanim (2012). Fitness costs associated with insecticide resistance: Fitness cost and insecticide resistance. en. *Pest Management Science* 68, 1431–1437. https://doi.org/10.1002/ps.3395.

Kreppel KS, M Viana, BJ Main, PCD Johnson, NJ Govella, Y Lee, D Maliti, FC Meza, GC Lanzaro, and HM Ferguson (2020). Emergence of behavioural avoidance strategies of malaria vectors in areas of high LLIN coverage in Tanzania. en. *Scientific Reports* 10. https://doi.org/10.1038/s41598-020-71187-4.

Labbé P, JP David, H Alout, P Milesi, L Djogbénou, N Pasteur, and M Weill (2017). Evolution of Resistance to Insecticide in Disease Vectors. en. In: *Genetics and Evolution of Infectious Diseases*. Elsevier, pp. 313–339. ISBN: 978-0-12-799942-5. https://doi.org/10.1016/B978-0-12-799942-5.00014-7.

Lockwood JA, TC Sparks, and RN Story (1984). Evolution of Insect Resistance to Insecticides: A Reevaluation of the Roles of Physiology and Behavior. en. *Bulletin of the Entomological Society of America* 30, 41–51. https://doi.org/10.1093/besa/30.4.41.

Long JA (2020). *jtools: Analysis and Presentation of Social Scientific Data*.

Main BJ, Y Lee, HM Ferguson, KS Kreppel, A Kihonda, NJ Govella, TC Collier, AJ Cornel, E Eskin, EY Kang, CC Nieman, AM Weakley, and GC Lanzaro (2016). The Genetic Basis of Host Preference and Resting Behavior in the Major African Malaria Vector, Anopheles arabiensis. en. *PLOS Genetics* 12. Ed. by Zwiebel LJ, e1006303. https://doi.org/10.1371/journal.pgen.1006303.

Martinez-Torres D, F Chandre, MS Williamson, F Darriet, JB Berge, AL Devonshire, P Guillet, N Pasteur, and D Pauron (1998). Molecular characterization of pyrethroid knockdown resistance (kdr) in the major malaria vector Anopheles gambiae s.s. en. *Insect Molecular Biology* 7, 179–184. https://doi.org/10.1046/j.1365-2583.1998.72062.x.

MERSI, CNRST, and IRSS (2016). Utilisation des pesticides agricoles dans trois régions à l'ouest du Burkina Faso et évaluation de leur impact sur la santé et l'environnement: cas des Régions de la Boucle du Mouhoun, des Cascades et des Hauts-Bassins.

Moiroux N (2012). Modélisation du risque d'exposition aux moustiques vecteurs de Plasmodium spp. dans un contexte de lutte anti-vectorielle. 2012MON20177. PhD thesis.

Moiroux N, AS Bio-Bangana, A Djènontin, F Chandre, V Corbel, and H Guis (2013). Modelling the risk of being bitten by malaria vectors in a vector control area in southern Benin, west Africa. *Parasites & Vectors* 6, 71. https://doi.org/10.1186/1756-3305-6-71.

Moiroux N, A Djènontin, AS Bio-Bangana, F Chandre, V Corbel, and H Guis (2014). Spatio-temporal analysis of abundances of three malaria vector species in southern Benin using zero-truncated models. en. *Parasites & Vectors* 7, 103. https://doi.org/10.1186/1756-3305-7-103.

Moyes CL, DK Athinya, T Seethaler, KE Battle, M Sinka, MP Hadi, J Hemingway, M Coleman, and PA Hancock (2020). Evaluating insecticide resistance across African districts to aid malaria control decisions. en. *Proceedings of the National Academy of Sciences* 117, 22042–22050. https://doi.org/10.1073/pnas.2006781117.

Nakagawa S and H Schielzeth (2013). A general and simple method for obtaining R^2 from generalized linear mixed-effects models. en. *Methods in Ecology and Evolution* 4. Ed. by O'Hara RB, 133–142. https://doi.org/10.1111/j.2041-210x.2012.00261.x.

Ngowo H, E Kaindoa, J Matthiopoulos, H Ferguson, and F Okumu (2017). Variations in household microclimate affect outdoor-biting behaviour of malaria vectors. English. *Wellcome Open Research* 2, 102–102. https://doi.org/10.12688/wellcomeopenres.12928.1.

Njan Nloga A, V Robert, J Toto, and P Carnevale (1993). La durée du cycle gonotrophique d'Anopheles moucheti varie de trois à quatre jours en fonction de la proximité par rapport aux gites de ponte. FRE. *Bulletin de Liaison et de Documentation - OCEAC* 26, 69–72.

Ouedraogo M, A M., T Z., and P I. (2011). Pesticides in Burkina Faso: Overview of the Situation in a Sahelian African Country. en. In: *Pesticides in the Modern World - Pesticides Use and Management*. Ed. by Stoytcheva M. InTech. ISBN: 978-953-307-459-7. https://doi.org/10.5772/16507.

Paaijmans KP and S Huijben (2020). Taking the 'l' out of LLINs: using insecticides in vector control tools other than long-lasting nets to fight malaria. en. *Malaria Journal* 19. https://doi.org/10.1186/s12936-020-3151-x.

Pedersen TL (2019). *patchwork: The Composer of Plots*.

PNLP (2014a). Directives nationales pour la prise en charge du paludisme dans les formations sanitaires du Burkina Faso. Ministère de la Santé/Burkina Faso.

— (2014b). Programme National de Lutte Contre le Paludisme en Côte d'Ivoire. 2014. Plan stratégique national de lutte contre le paludisme 2012–2015 (période replanifiée 2014–2017). Approche stratifiée de mise à l'échelle des interventions de lutte contre le paludisme en Côte d'Ivoire et consolidation des acquis. Abidjan: Ministère de La Santé et l'Hygiène Publique. 149 p.

Porciani A, M Diop, N Moiroux, T Kadoke-Lambi, A Cohuet, F Chandre, L Dormont, and C Pennetier (2017). Influence of pyrethroïd-treated bed net on host seeking behavior of Anopheles gambiae s.s. carrying the kdr allele. en. *PLOS ONE* 12. Ed. by Favia G, e0164518. https://doi.org/10.1371/journal.pone.0164518.

QGIS Development Team (2021). *QGIS Geographic Information System*. QGIS Association.

R Core Team (2018). *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing.

Ranson H, B Jensen, JM Vulule, X Wang, J Hemingway, and FH Collins (2000). Identification of a point mutation in the voltage-gated sodium channel gene of Kenyan Anopheles gambiae associated with resistance to DDT and pyrethroids. en. *Insect Molecular Biology* 9, 491–497. https://doi.org/10.1046/j.1365-2583.2000.00209.x.

Reid MC and FE McKenzie (2016). The contribution of agricultural insecticide use to increasing insecticide resistance in African malaria vectors. en. *Malaria Journal* 15. https://doi.org/10.1186/s12936-016-1162-4.

Riveron JM, M Tchouakui, L Mugenzi, BD Menze, MC Chiang, and CS Wondji (2018). Insecticide Resistance in Malaria Vectors: An Update at a Global Scale. en. In: *Towards Malaria Elimination - A Leap Forward*. Ed. by Manguin S and Dev V. InTech. ISBN: 978-1-78923-550-0 978-1-78923-551-7. https://doi.org/10.5772/intechopen.78375.

RStudio Team (2020). *RStudio: Integrated Development Environment for R*. Boston, MA: RStudio, PBC.

Sanou A, L Nelli, WM Guelbéogo, F Cissé, M Tapsoba, P Ouédraogo, N Sagnon, H Ranson, J Matthiopoulos, and HM Ferguson (2021). Insecticide resistance and behavioural adaptation as a response to long-lasting insecticidal net deployment in malaria vectors in the Cascades region of Burkina Faso. en. *Scientific Reports* 11, 17569. https://doi.org/10.1038/s41598-021-96759-w.

Sherrard-Smith E, JE Skarp, AD Beale, C Fornadel, LC Norris, SJ Moore, S Mihreteab, JD Charlwood, S Bhatt, P Winskill, JT Griffin, and TS Churcher (2019). Mosquito feeding behavior and how it influences residual malaria transmission across Africa. en. *Proceedings of the National Academy of Sciences* 116, 15086–15095. https://doi.org/10.1073/pnas.1820646116.

Shmueli G (2010). To Explain or to Predict? en. *Statistical Science* 25, 289–310. https://doi.org/10.1214/10-STS330.

Shmueli G and O Koppius (2010). Predictive Analytics in Information Systems Research. en. *SSRN Electronic Journal*. https://doi.org/10.2139/ssrn.1606674.

Sinka ME, MJ Bangs, S Manguin, M Coetzee, CM Mbogo, J Hemingway, AP Patil, WH Temperley, PW Gething, CW Kabaria, RM Okara, T Van Boeckel, HCJ Godfray, RE Harbach, and SI Hay (2010). The dominant Anopheles vectors of human malaria in Africa, Europe and the Middle East: occurrence data, distribution maps and bionomic précis. en. *Parasites & Vectors* 3. https://doi.org/10.1186/1756-3305-3-117.

Snow RW and HM Gilles (2002). The epidemiology of malaria. *Essential malariology* 4.

Sokhna C, MO Ndiath, and C Rogier (2013). The changes in mosquito vector behaviour and the emerging resistance to insecticides will challenge the decline of malaria. en. *Clinical Microbiology and Infection* 19, 902–907. https://doi.org/10.1111/1469-0691.12314.

Soma DD, B Zogo, P Taconet, K Mouline, LPA Alou, RK Dabiré, AA Koffi, C Pennetier, and N Moiroux (2023). Anopheles collections in the health districts of Korhogo (Côte d'Ivoire) and Diébougou (Burkina Faso) (2016-2018). en. Type: dataset. https://doi.org/10.15468/V8FVYN.

Soma, B Zogo, P Taconet, A Somé, S Coulibaly, L Baba-Moussa, GA Ouédraogo, A Koffi, C Pennetier, KR Dabiré, and N Moiroux (2021). Quantifying and characterizing hourly human exposure to malaria vectors bites to address residual malaria transmission during dry and rainy seasons in rural Southwest Burkina Faso. en. *BMC Public Health* 21. https://doi.org/10.1186/s12889-021-10304-y.

Soma, B Zogo, A Somé, BN Tchiekoi, DFdS Hien, HS Pooda, S Coulibaly, JE Gnambani, A Ouari, K Mouline, A Dahounto, GA Ouédraogo, F Fournet, AA Koffi, C Pennetier, N Moiroux, and RK Dabiré (2020). Anopheles bionomics, insecticide resistance and malaria transmission in southwest Burkina Faso: A pre-intervention study. en. *PLOS ONE* 15, e0236920. https://doi.org/10.1371/journal.pone.0236920.

Taconet P, A Porciani, DD Soma, K Mouline, F Simard, AA Koffi, C Pennetier, RK Dabiré, M Mangeas, and N Moiroux (2021). Data-driven and interpretable machine-learning modeling to explore the fine-scale environmental determinants of malaria vectors biting rates in rural Burkina Faso. en. *Parasites & Vectors* 14. https://doi.org/10.1186/s13071-021-04851-x.

Taconet P, DD Soma, B Zogo, K Mouline, F Simard, AA Koffi, RK Dabiré, C Pennetier, and N Moiroux (2023). *Replication data and script for the manuscript "Physiological and behavioural resistance of malaria vectors*

in rural West-Africa : a data mining study to adress their fine-scale spatiotemporal heterogeneity, drivers, and predictability". Version VERSION PROVISOIRE. https://doi.org/10.23708/LV8GEW.

Taconet P, DD Soma, B Zogo, K Mouline, F Simard, A Koffi Amanan, RK Dabiré, C Pennetier, and N Moiroux (2023). *Supplementary information for the manuscript 'Physiological and behavioural resistance of malaria vectors in rural West-Africa : a data mining study to adress their fine-scale spatiotemporal heterogeneity, drivers, and predictability'*. Version VERSION PROVISOIRE. https://doi.org/10.23708/VJEEMU.

Tyagi S and S Mittal (2020). Sampling Approaches for Imbalanced Data Classification Problem in Machine Learning. en. In: *Proceedings of ICRIC 2019*. Ed. by Singh PK, Kar AK, Singh Y, Kolekar MH, and Tanwar S. Vol. 597. Cham: Springer International Publishing, pp. 209–221. ISBN: 978-3-030-29406-9 978-3-030-29407-6. https://doi.org/10.1007/978-3-030-29407-6_17.

Voeten CC (2020). *buildmer: Stepwise Elimination and Term Reordering for Mixed-Effects Regression*.

Wan Z, S Hook, and G Hulley (2015a). MOD11A1 MODIS/Terra Land Surface Temperature/Emissivity Daily L3 Global 1km SIN Grid V006. type: dataset. https://doi.org/10.5067/MODIS/MOD11A1.006.

— (2015b). MYD11A1 MODIS/Aqua Land Surface Temperature/Emissivity Daily L3 Global 1km SIN Grid V006. type: dataset. https://doi.org/10.5067/MODIS/MYD11A1.006.

Weill M, C Malcolm, F Chandre, K Mogensen, A Berthomieu, M Marquine, and M Raymond (2004). The unique mutation in ace-1 giving high insecticide resistance is easily detectable in mosquito vectors. en. *Insect Molecular Biology* 13, 1–7. https://doi.org/10.1111/j.1365-2583.2004.00452.x.

WHO (2017). WHO | Global vector control response 2017–2030.

— (2022). World malaria report 2022. en.

Wickham H (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York. ISBN: 978-3-319-24277-4.

— (2017). *tidyverse: Easily Install and Load the 'Tidyverse'*.

Wing MKCfJ, S Weston, A Williams, C Keefer, A Engelhardt, T Cooper, Z Mayer, B Kenkel, tRC Team, M Benesty, R Lescarbeau, A Ziem, L Scrucca, Y Tang, C Candan, and T Hunt (2018). *caret: Classification and Regression Training*.

Wright MN and A Ziegler (2017). ranger: A Fast Implementation of Random Forests for High Dimensional Data in C++ and R. *Journal of Statistical Software* 77, 1–17. https://doi.org/10.18637/jss.v077.i01.

Yadouleton A, T Martin, G Padonou, F Chandre, A Asidi, L Djogbenou, R Dabiré, R Aïkpon, M Boko, I Glitho, and M Akogbeto (2011). Cotton pest management practices and the selection of pyrethroid resistance in Anopheles gambiae population in Northern Benin. en. *Parasites & Vectors* 4. https://doi.org/10.1186/1756-3305-4-60.

Yan Y (2016). *MLmetrics: Machine Learning Evaluation Metrics*.

Zhao Q and T Hastie (2021). Causal Interpretations of Black-Box Models. en. *Journal of Business & Economic Statistics* 39, 272–281. https://doi.org/10.1080/07350015.2019.1624293.

Zogo B, DD Soma, BN Tchiekoi, A Somé, LP Ahoua Alou, AA Koffi, F Fournet, A Dahounto, B Coulibaly, S Kandé, RK Dabiré, L Baba-Moussa, N Moiroux, and C Pennetier (2019). *Anopheles* bionomics, insecticide resistance mechanisms, and malaria transmission in the Korhogo area, northern Côte d'Ivoire: a pre-intervention study. *Parasite* 26, 40. https://doi.org/10.1051/parasite/2019040.