1   **Title:** Fear and foraging interact with temperature and seasonality to drive diversity in fish life
2   histories
3
4   **Running Title:** Ecosystem size spectra drive life history diversity
5
6   **Authors:**

7   Kindsvater, Holly K.[1], Juan-Jordá, Maria-José[2,3], Dulvy, Nicholas K.[2], Horswill, Cat[4,5,6],
8   Matthiopoulos, Jason[7], Mangel, Marc[8,9]
9
10
1. Dept. of Fish and Wildlife Conservation, Virginia Polytechnic Institute and State University,
310 W. Campus Dr., Blacksburg, VA 24061, USA
2. Earth to Ocean Research Group, Dept. of Biological Sciences, Simon Fraser University,
Burnaby, BC V5A 1S6 Canada
3. AZTI Tecnalia, Herrera Kaia, Gipuzkoa, Spain
4. ZSL Institute of Zoology, Regent's Park, London, NW1 4RY, UK
5. Centre for Biodiversity and Environmental Research, Department of Genetics, Evolution and
Environment, University College London, Gower Street, London, WC1E 6BT, UK
6. Department of Zoology, University of Cambridge, Downing St, Cambridge CB2 3EJ, UK
7. Institute of Biodiversity, Animal Health and Comparative Medicine, Graham Kerr Building,
University of Glasgow, Glasgow, G12 8QQ, Scotland.
8. Theoretical Ecology Group, Dept. of Biology, University of Bergen, Bergen 5020, Norway
9. Institute of Marine Sciences and Dept. of Applied Mathematics and Statistics, University of
California, Santa Cruz, CA 95064, USA.

25
26
27   **Keywords –** Body size evolution, ecosystem size spectra, climate change, metabolic theory,
28   state-dependent models
29
30
31

Abstract

Understanding how fish growth and reproduction will adapt to changing environmental
conditions is a fundamental question in evolutionary ecology, but predicting the responses of
specific taxa is challenging. Analyses of the physiological effects of climate change upon life
histories and population demography rarely consider size-dependent foraging and risk of
predation. We embedded a state-dependent energetic model in an ecosystem size spectrum to
model prey availability (foraging) and risk of predation (fear) experienced by individual fish as
they grow. We found that asymptotic growth emerged from size-based foraging and reproductive
and mortality patterns in the context of ecosystem food web interactions, without *a priori*
assumptions about the onset of reproduction. While more productive ecosystems led to larger
maximum body sizes, the effects of temperature on body size were surprisingly small. To
validate our model, we ran it for abiotic scenarios corresponding to the three ecological lifestyles
of tunas, considering environments that included seasonal variation in temperature. We
successfully predicted realistic patterns of growth, reproduction, and mortality of all three tuna
species with distinct ecological lifestyles. Unexpectedly, we found that individuals grew larger
when they experienced higher productivity or colder temperatures for part of the year, due to
seasonal migrations. This increase in body size emerged because spawning opportunities were
seasonally constrained, and explains the evolution of gigantism in temperate tunas. Our approach
combines direct and indirect effects of productivity (foodscapes) and individual risk (fearscapes)
with metabolic processes and offers a promising approach to understand fish life-history
responses to changing ocean conditions.

Introduction

The body sizes and population biomasses of aquatic species are changing rapidly to human-induced environmental change (Oke et al. 2020), motivating the need for mechanistic models to predict growth and reproduction in response to changing environmental conditions (Cheung et al. 2008, Cheung et al. 2010, Blanchard et al. 2012, Fernandes et al. 2020). Beyond thermal physiology, the evolutionary consequences resulting from changing ecosystem productivity (foodscapes) and predator abundance (fearscapes) are difficult to disentangle from the effects of changes in prey abundance, predation risk or seasonality on growth and related life history traits (Daufresne et al. 2009, Audzijonyte and Richards 2018, Neubauer and Andersen 2019). Classic and recent advances predicting optimal growth and reproduction of harvested populations (Holt 1958; Beverton and Holt 1959; White et al. 2022) have been combined with size-structured community interactions to predict the evolution of life history traits in specific taxa (Shuter et al. 2016). This approach provides a promising foundation for predicting ecological and evolutionary responses to environmental and climate change. While specific taxa within an ecosystem will adapt according to contextual factors such as phylogenetic lineage, adaptive capacity, and environmental variability, mechanistic models incorporating the roles of foodscapes and fearscapes, seasonality, and other metabolic processes could explain adaptive variation among closely related taxa and refine our mechanistic understanding of species' responses to changing environmental and climate conditions (Free et al. 2019). By examining closely related tuna species (Scombridae) that have invaded different environmental niches, we can gain insight into how physiological adaptations to thermal conditions, productivity and their seasonality affect allocation to growth and reproduction. This can help move us beyond broad macro-ecological

patterns toward a more mechanistic understanding of the drivers of existing diversity and

improve predictions of species-specific responses to future change (Alvarez-Noriega et a. 2023).

In ectothermic species with indeterminate growth, including fish, food availability and risk of

predation both increase as individuals grow larger, and growth rates can be selected to be slower

or faster depending on the size-dependence of per-capita resource availability and predication

risk (Conover and Munch 2002; Walsh and Reznick 2009, Hulthén et al 2021). At the same time,

development at higher temperatures can lead to smaller body sizes, within and between

ectothermic species (Kingsolver and Huey 2008). Trade-offs among these life-history traits

determine individual body sizes and population demographic rates (Beverton and Holt 1959,

Gadgil and Bossert 1970, Kozlowski 1992, Kozlowski 1996), but existing theory struggles to

predict how size-dependent changes in prey availability and decreases in predation risk, along

with interacting effects of temperature and seasonality, affect the evolution of growth (Varpe

2017). There is longstanding interest in understanding the mechanisms leading to biphasic or

asymptotic growth patterns (Quince et al. 2008). The earliest growth models, including the von

Bertalanffy growth model, were based on hypothesized differences in the allometric scaling of

anabolism (resources taken in) and catabolism (resources spent) (von Bertalanffy 1960,

Audzijonyte et al. 2022). Such growth models are routinely fit to data on size-at-age of fishes,

but their mechanistic justification is necessarily simplistic; such models struggle to explain

spatial and temporal trends in body size within and among species of fish (Audzijonyte et al.

2020), suggesting that they do not sufficiently capture the relevant intrinsic physiological and

extrinsic ecological drivers of fish life histories.

The role of metabolic requirements and the trade-off between growth and reproduction in

shaping growth trajectories has been given increasing attention in recent decades (Jørgensen et

al. 2016, Wong et al. 2021, White et al 2022). Physiological processes can vary with the

environmental temperature experienced by organisms (Brown et al, 2004, Clarke and Johnston,

1999). The allometric scaling of metabolic costs in different temperature regimes (known as the

Metabolic Theory of Ecology [MTE]; Gillooly et al. 2001) has been used to predict individual

body size according to the benefits of growing large (to increase foraging success and avoid

predation), balanced against the costs of increased metabolic overhead and diverting resources

from reproduction to growth (Wong et al. 2021). More detailed dynamic models of energetic

allocation have linked variable environmental conditions to growth, reproduction, and longevity

(Kozłowski and Teriokhin 1999, Cichoń and Kozłowski, 2000, Lika and Kooijman 2003,

Jorgensen and Fiksen 2006, Audzijonyte et al. 2022). Understandably, the results of these

energetic budget models depend on specific assumptions regarding mass-based foraging success

and risk of predation. Generalizing these prior results requires a unified framework incorporating

metabolic demands, access to resources, and predation risk in environments of differing

temperatures, productivity, and seasonality.

We draw on ecosystem size-spectra theory to reduce the number of *ad hoc* assumptions required

about the scaling of ecological processes - foodscapes and fearscapes - that arise from

community interactions. We hypothesize this approach could yield more realistic predictions of

diversity in fish life histories, providing significant insights into the mechanisms that determine

adaptive responses to changing environmental conditions. In aquatic ecosystems, size spectra are

an emergent property of community interactions (Sheldon et al. 1977, Thygesen et al. 2005, Law

et al. 2009, Sprules and Barth 2016, Christensen and Andersen 2011, Andersen et al. 2016,

Andersen 2019). In a community size spectrum, flow of energy between trophic levels via

consumption and predation rates are characterized by individual mass, instead of species'

identity (Benoît and Rochet 2004, Blanchard et al. 2009, Andersen 2019). The key property of

size spectra is that an individual's relative position (mass) determines both its prey field

(foodscape – or the area under the size spectrum to the left of an individual's mass) and its risk

of predation (fearscape – the area to the right). Individuals are born small and grow through the

size spectrum over their lifetime, consuming prey that are a fraction of their own size.

Furthermore, size-spectrum theory assumes the lower limit of prey size preference depends on

the profitability of the prey, and any upper limit depends on the maximum gape size of the

predator. These interactions could equally apply to interactions within size-structured

populations of the same species (i.e., cannibalism). Therefore, predation and consumption rates

determined by different areas under a size-spectrum curve could be used to simultaneously

characterize the mass-specific caloric resource availability and risk of predation experienced by

an individual as it grows (Benoît and Rochet 2004, Giacomini et al. 2013, Shuter et al. 2016,

Andersen 2019).

Here, we used an energetic model combining size-dependent resources and risk of predation

(both derived from size-spectra) with size- and temperature-dependent physiological

requirements to predict the evolution of diverse fish life histories including growth, maximum

body size, reproductive investment, and longevity in different environments. We defined fitness

as expected lifetime reproductive success and determined its maximum value and the associated

life history decisions (allocation to maintenance, growth and reproduction) using stochastic

dynamic programming (Clark and Mangel, 2000, Houston et al.1988, Mangel, 2015). We used

this approach to examine general patterns in the life history evolution of fishes under different

scenarios of ecosystem productivity and environmental temperature.

We next investigated the evolution of growth, body sizes and reproductive patterns in tuna

species under several environmental scenarios with varying temperatures, ecosystem

productivities and seasonally varying conditions. Tunas (Family Scombridae) exhibit a wide

range of maximum body sizes (~ 40 cm - 400 cm), longevities (~ 4 - 41 years) and reproductive

patterns (Juan-Jorda et al. 2013, Horswill et al. 2019). Tunas are epipelagic species found in

temperate and tropical waters around the world's oceans with varying vertical, latitudinal and

seasonal distributions and movements. Confronting the predictions of our general model with

data on the life history diversity observed in tuna species in different environmental conditions

provided insight into the mechanisms underlying fish life histories. Where possible, our models

were parameterized with values for physiological processes measured for tunas, but we also

considered the sensitivity of our results to these parameters to ensure their generality across the

diverse life histories observed in fishes.

Methods

We used size-spectrum theory to infer mass-specific rates of prey encounter and mass-specific

rates of predation. The relationship between the numbers of organisms and individual mass w is

a power function with a scaling parameter κ and an exponent λ, which has been empirically

estimated to be close to -1 such that abundance is inversely proportional to mass (Sheldon et al.

1972, Trebilco et al. 2013, Sprules and Barth 2016, Hatton et al. 2021)

$$N(w) = \kappa w^{\lambda} \qquad (1)$$

The Sheldon size spectrum, which follows from Eq. 1, is the distribution of total ecosystem

biomass *B(w)* across body size classes, and is represented by $B(w) = N(w)w$. Based on

evidence from multiple ecosystems (Sprules and Barth 2016), the spectrum – as originally

conceived (Sheldon et al 1972) and recently confirmed (Hatton et al. 2021) – is nearly flat, or

may very slowly decrease as body size increases. Since $N(1) = \kappa$, we view κ as a metric of

ecosystem productivity. In a log-log plot of Eq. 1, the intercept is the $\log(\kappa)$.

This phenomenon of linear aquatic size spectra emerges from three size-dependent processes:

(1) the encounter rate of predators and prey; (2) the preference of predators for prey of a given

size; and (3) the limit to prey consumption imposed by the size of the predator's stomach (the

predator-prey mass ratio; Benoît and Rochet 2004, Blanchard et al. 2017, Andersen 2019).

Anderson (2019) derived a method to calculate the prey available to individuals of mass *w* by

relating the productivity of a spectrum to empirical estimates of prey encounter rates, predator-

prey mass ratios, and prey preferences. Following his rationale, we assumed that the prey

biomass available to an individual is a concave function of mass *w*, and scaled to be

approximately three times the size spectrum productivity κ (Andersen 2019). The empirical

estimates of physiological processes used to simplify this derivation are associated with a

significant amount of variability among species within a size class (Andersen 2019). Following

Andersen, as a baseline the per-period food available for an individual of mass *w* is represented

by

$$B_{prey}(w) = 3\kappa w^{0.05} \tag{2}$$

This equation approximates the integral over the foodscape of prey biomass (in kg) available to an individual of mass w each month (Fig. 1A), and includes a threefold scaling factor from empirical analyses (Andersen 2019). The foodscape is therefore the total biomass available to the predator, and the range of sizes of prey that it takes. To address variability across ecosystems in productivity, we chose to vary κ in subsequent analyses, representing potential differences among taxa or other ecosystem processes.

Second, to infer mass-specific risk of predation, we defined the fearscape as an individuals' instantaneous risk of predation, integrated across sizes of potential predators. For gape-limited taxa like fishes, a predator's prey field depends on its encounter rate with prey in its preferred size range. This encounter rate (the clearance rate) is typically measured in units of volume per time, as aquatic species forage in a three-dimensional habitat (Kiørboe and Hirst 2014). Again, we followed Andersen (2019), and in Eq. 3 (defined below) we used an empirically estimated constant of 0.07 to characterize the scaling of prey vulnerability with its mass, which is based on estimates of preference windows of predators and the volume of water each clears per month that come from empirical distributions of prey sizes in predator guts (Ursin 1973). We also define a coefficient h, which modifies the probability of consumption - how likely a predator is to capture the focal individual (based on its hunger level or capture efficiency) - and used the reported allometric exponent of -0.25 to represent how predation mortality scales with body size (Figure 1B; Eq. 2.22 in Andersen 2019). From these, Anderson (2019) integrated over the Sheldon

213   spectrum to produce a relationship predicting the scaling of mortality risk for prey $\mu_p(w)$ per

214   month:

$$\mu_p(w) = 0.07hw^{-0.25} \tag{3}$$

218   We convert this instantaneous rate to the probability of escaping from predators during each time

219   interval (Hilborn and Mangel 1997, Mangel 2006), so we can represent the probability an

220   individual survives $\gamma_{pred}(w)$ as:

$$\gamma_{pred}(w) = e^{-\mu_p(w)} \tag{4}$$

The resulting mass-specific survival through each month is plotted for different values of h in

Fig. 1B.

Defining physiological costs that depend on temperature and body mass

Our energetic model includes physiological costs which depend on temperature and body mass.

We assumed that metabolic costs increase with body mass and environmental temperature

(Clarke 2006). We modeedl individual costs $C(w, \tau)$ (in joules) as a function of temperature τ in

Kelvin, following the general form introduced in the MTE (Gillooly et al. 2001):

$$C(w, \tau) = cw^\theta e^{\left(-E/k_B\tau\right)} \tag{5}$$

Evidence for the MTE suggests the activation energy E (the energy required for the reactions of

respiration and other metabolic processes) does not vary appreciably among taxa (Bernhardt et

al. 2018); Boltzmann's constant k_B also does not vary. The normalization coefficient c accounts

for differences among taxonomic groups in the intercepts of the linear relationship that arise

from second-order effects such as stoichiometry or respiratory surface areas (Bigman et al 2021).

The slope of this relationship in log space, θ, is strikingly similar among taxa (Brown et al.

2004). We used a value for θ estimated from physiological studies on tunas (Table 1) but varied

it in sensitivity analyses. Note that τ in Eq. 5 is traditionally in Kelvin. However, we rescale all

values in our results in units of Celsius. We use this general formula for scaling of metabolic

costs at different temperatures τ to describe the monthly energetic expenses of an individual of

mass w (Fig 1C). Defining the relationships in Eqs. 2, 4, and 5 allowed us to specify mass-

dependent survival and energy dynamics and therefore examine the variables influencing growth

and reproduction in a common currency of individual fitness. All parameters are defined and

their values reported in Table 1.

Figure 1. Examples of allometric relationships between individual mass and (A) income (in kg per month) for differing ecosystem productivities κ (Eq. 2); (B) the scaling of survival with size and its interaction with predator efficiency (or hunger) *h* (Eq. 4), such that higher values of *h* represent more efficient predators; and (C) metabolic costs (Eq. 5), which also scale with temperature τ (on the graph, it is presented in degrees Celsius, °C; values in Kelvin are 285, 290, 295 and 300K). Note that these curves represent a subset of values considered in our results.

Defining fitness and determining the optimal allocation strategy

To find the optimal life history strategy in different scenarios of ecosystem productivity and

temperature conditions, including seasonal variation in the scenarios, we developed a model of

an individual's energy budget, tracking two physiological state variables, the body length l and

energy (lipid) stores s of an individual, which vary dynamically over an individual's lifetime

(Fig. 2). Following the conventions for dynamic state-variable models, we denoted the state

variables l and s are lowercase in the dynamic programming equation, representing the fact they

are iterated values; potential future states are denoted as l' and s'. Later, when we refer to specific

values of the state variables, we use uppercase L and S.

We define fitness as expected lifetime reproduction, averaged over the stochastic process of

survival, which we calculated numerically using stochastic dynamic programming (Houston et

al. 1988; Clark and Mangel 2000, Mangel, 2015). This method allowed us to consider how

individual age and physiological state (energy stores and body length) affect the optimal trade-

off between growth and reproduction in the context of lifetime fitness. We assumed that an

individual allocates a proportion of its budget to growth and reproduction on monthly time

intervals. Our choice to model allocation to growth and reproduction as proportions of an

individual's energy budget builds on prior dynamic state-variable models of fish growth

(Jørgensen and Fiksen 2006, Chapman et al. 2011).

Given specific values of $L(t)$ and $S(t)$, representing individual length and energy stores at the

start of month t, structural mass $W(t)$ is related to length with a standard cubic function (Froese

2006):

Figure 2. Conceptual overview of the optimization algorithm calculating the two dynamically varying state variables, body length $l(t)$ and energy stores $s(t)$ within each month t, as well as current and expected fitness. Arrows represent the flow of energy or decisions. Round shapes represent energetic input (green) and outputs (yellow); rectangular shapes are model states and outputs (fitness and fate). Note that $l(t)$ and $w(t)$ are related through Eq. 6. Both B_{prey} and μ_p are determined by the size spectrum.

$$W(t) = aL(t)^3 \tag{6}$$

We assumed that only structural mass is relevant for size-dependent gains (B_{prey}) and costs (C)

(Fig. 2). We convert from mass to units of energy (joules) and back using a conversion factor

estimated for tunas ($\rho = 4.2 \times 10^6$ J/kg; Chapman et al 2011) but that could take other values

for other species. Each month, an individual acquires energy from food, determined by their

structural mass w via Eq. 2. We then allowed the individual to allocate proportions of its energy

budget (which includes income and any stored energy) to growth (g) and reproduction (r).

Given an individual with specific values for length $L(t)$ and energy stores $S(t)$, we calculated

the increment of growth expected for each possible proportional allocation g by converting the

fraction of lipid stores from joules to the equivalent mass ($g\frac{S(t)}{\rho}$), and then adding it to existing

mass, such that $W(t + 1) = W(t) + g\frac{S(t)}{\rho}$. We then calculated the new length by rearranging

the mass-length relationship $W(t) = aL(t)^3$:

$$L(t + 1) = \left(L(t)^3 + g\frac{S(t)}{a\rho}\right)^{1/3} \tag{7}$$

294 We repeated this for every possible combination of values of the state variables, l and s. Values

295 for proportional allocation of lipid stores to reproduction r, along with g, combine to determine

296 the dynamics of energy stores and length from one month to the next:

$$S(t + 1) = B_{prey}(W(t)) - C(W(t), \tau) - (r + g) \cdot S(t) \tag{8}$$

We assumed that both stored energy and reproduction were limited by an individual's structural

mass (which in turn depends on its length). These constraints represent limits on both the amount

of lipid can be stored, and the mass of gametes that can be produced, given the capacity of the

body cavity. If the proportions of energy allocated to reproduction and growth were less than

100% (meaning $r + g < 1$), the remaining energy was assumed to be stored for future use, given

that total reserves did not exceed 60% of structural mass. This value was arbitrary, but

exploratory analyses suggest it did not have a strong effect on the results presented here because

in practice individuals do not store their energy long enough to come close to exceeding this

limit. Reproductive output (in units of kg) was similarly constrained so that it cannot exceed a

fixed proportion of φ of structural mass $w(t)$, such that the following condition must be met:

311
$$\frac{rS(t)}{\rho} \leq \varphi w(l, t) \tag{9}$$

This size-based limit on total reproduction is used in the calculation of current fitness.

Expected future fitness

At any age, expected fitness was the sum of current reproduction and accumulated future

reproduction, which was calculated assuming the individual behaved optimally for all future

ages. This required calculation of the future states (length and lipid stores) given each

combination of allocation to growth and reproduction. We denoted potential values of future

states as l' and s'. These are

322
$$l'(l, s, g) = l + \left(l(t)^3 + g\frac{s}{a\rho} \right)^{1/3}$$

323
$$s'(l, s, r, g) = B_{prey}\big(w(l)\big) - C\big(w(l), \tau\big) - (r + g)s \tag{10}$$

325   where $w(l)$ is the structural mass of an individual of length l. At the end of the month, if an

326   individual's lipid stores fell below the critical threshold for its mass (its expenditures have

327   exceeded its energy budget), it starved (Supp. Fig. 2). We let $\gamma_s(s, l)$ denote the probability of

328   avoiding starvation in the current time and modeled it as

330
$$\gamma_s(s, l) = \frac{1}{1 + e^{-q(s - vw(l)\rho)}} \tag{11}$$

involving two new parameters v and q. The parameter v determined the level of stores at which

starvation begins; q is a shape parameter. When $s = vw(l)\rho$, the right-hand side of Eq. 11 is

always ½. When q is large, then the right-hand side of Eq. 11 is approximately one when $s >$

$vw(l)\rho$ and zero otherwise. Thus, this function specified that if combined allocation to growth

and reproduction caused an individual's lipid stores to decrease below the critical value for its

mass, its probability of survival decreased smoothly toward zero (See Supp. Fig. 1 for more

details).

Equations 7-11 characterize changes in individual state and in fitness, given all possible monthly

allocation strategies (r and s). With these functions in place, we can find the allocation strategy

that maximize current and future fitness at every age until the age of senescence or maximum

lifespan, T, is reached (for all scenarios, we assumed the maximum lifespan of $T = 216$ months

or 18 years). We define $V(l, s, t)$ as the maximum expected accumulated reproduction between

time t and T, given size $L(t) = l$ and lipid stores $S(t) = s$. Since there can be no

accumulated reproduction after T, we assumed $V(l, s, T) = 0$. Expected future fitness at every

age $t < T$ was found by solving the stochastic dynamic programming equation

$$V(l, s, t+1) = \max_{g, r}\left[R(r, l, s, t) + \gamma_{pred}(w) \cdot \gamma_s(s, l) \cdot V\big((l'(l, s, g), s'(l, s, r, g), t\big) \right] \tag{12}$$

The first term of the on the right-hand side of Eq. 12 represents reproduction in month t. The

second term represents expected future reproduction, discounted by the probability of surviving

predation γ_{pred} (w) and starvation $\gamma_s(s, l)$, When these are combined, we can obtain expected

lifetime reproduction from time t onwards, given that size $L(t) = l$ and lipid stores $S(t) = s$.

The dynamic programming algorithm (Mangel and Clark 1988; Houston and MacNamara 1999)

iterates over all viable combinations of l and s, at each time t, and stores the fitness of each

allocation strategy. The optimal strategy (marked with an asterisk) at time t is the combination

g^* and r^* associated with the greatest current and future fitness. Further details of the

optimization algorithm are given in the Appendix. In Supp. Fig 3 we illustrate the array for both

allocation strategies ($g^*(l,s,t)$ and $r^*(l,s,t)$) at two ages, for all possible combinations of length

and lipid stores.

Calculating the fate of individuals allocating optimally

We used a Monte Carlo simulation to calculate the accumulating probability of mortality as an

individual ages, following the allocation rules generated by the solution of Eq. 12, which

provides optimal allocation rules for the range of state values specified in our numerical

algorithm. Some combinations of states (length, lipid stores and age) will not arise naturally and

others are inviable (the dark blue regions of Supp. Fig. 3). Therefore, to predict the relationships

between length, lipid stores, and age that will be observed in nature, we simulated the expected

fates of an individual that is born at a given size, and allocates energy according to the optimal

growth and reproduction fractions given by the solution to Eq. 12. We recorded the body length

and reproductive output in each subsequent month of an individual's life, and calculated its

probability of survival, given both the risk of starvation and the risk of predation. The expected

lifespan was defined as the age past which the cumulative chance of survival is less than 3%,

which we used as an arbitrary definition of lifespan. For simplicity, we considered reproductive

output in units of energy (joules) rather than considering allocation to offspring size and number

(Kindsvater et al. 2010); this can include migration costs. We did not build in any assumptions

about age at maturation, but rather let maturation patterns, along with natural trade-offs between

growth and reproduction (Supp. Fig. 3) emerge from patterns of allocation.

Scenarios for environmental variation

We used our general energetic model to predict a range of fish life histories (growth,

reproduction, and survival) that are evolutionarily advantageous across different scenarios of

ecosystem productivity and environmental temperatures. To do this, we developed different

productivity and temperature scenarios corresponding to different conditions commonly found in

the marine environment and impose different metabolic costs to individuals according to Eq. 5.

We solved for the optimal life histories under different temperatures (τ in Eq. 5 in units of

Kelvin), converted to Celsius and ranging from 11.85 to 26.85 ℃ (285-300 K) in five-degree

intervals. We considered these temperature ranges in different scenarios for ecosystem

productivity, for which κ ranged from 0.25 to five. We considered factorial combinations of

these parameters to understand the implications of their interactions for the evolution of fish life

histories.

Case study predicting life history diversity of tunas (Scombridae).

We developed a case study for tunas to examine if our general energetic model predicted the

general patterns of life history variation observed in this group of species adapted to different

environmental niches. There are 15 species of tunas within the Family Scombridae, from five

genera: *Allothunnus*, *Auxis*, *Euthynnus*, *Katsuwonus*, and *Thunnus* (Collette et al. 2001). These

species inhabit a wide range of environmental conditions in marine ecosystems. Paleo-

oceanographic evidence suggests that ancestral tunas evolved in a tropical environment

approximately 60 million years ago (Monsch 2000) and over time they have diversified and

evolved a suite of morphological and physiological adaptations that have allowed them to

expand their distributions into more temperate environments or deeper colder waters where they

can encounter higher prey densities to support their high somatic and gonadal growth rates

(Dickson and Graham 2004). Currently, tunas can be found in coastal and oceanic pelagic

waters, and have wide geographic distributions, ranging from the tropics to higher temperate

latitudes with some degree of habitat partitioning by depth. Tropical tunas can spawn throughout

the year, while the subtropical and temperate tunas undergo seasonal migrations returning form

cool high latitude feeding grounds back to warm waters for spawning (Juan-Jordá et al 2013,

Horswill et al 2019). Reflecting their tropical ancestor, all tunas (except for the slender tuna

Allothunnus fallai) spawn in warm waters, requiring a sea surface temperature of at least 24 ℃

(Schaefer 2001). This is a key aspect of their reproductive biology that we include in our model

scenarios.

To connect our general energetic model more explicitly with the observed patterns of life history

variation in tunas, we followed a proposed categorization of tunas into three ecological lifestyles

(Bernal et al. 2017). The three general lifestyles are based on species-specific vertical,

latitudinal, and temporal (seasonal) distributions and movement patterns of tunas (Figure 3;

Bernal et al. 2009, Bernal 2011, Bernal et al. 2017). The first ecological lifestyle represents a

tuna species that largely remains within the warmer and well-oxygenated surface layer above the

thermocline (generally above 20 °C) during both day and night. These tuna species have limited

vertical movements as they do not descend below the thermocline (Figure 3). Coastal species,

such as the tropical frigate tuna (*Auxis thazard*) may typify this group. The second ecological

lifestyle represents tuna species that spend the majority of the time above the thermocline

(generally above 20 °C) but also visit depths below the thermocline for foraging (Figure 4). The

oceanic species of yellowfin tuna (*Thunnus albacares*) with year-round tropical distributions

typifies this group. Its vertical movement exposes this species to a wider range of temperatures

and to less-oxygenated waters at depth but only for short periods of time because this species is

not hypoxia tolerant (Schaefer et al. 2009). The third ecological lifestyle characterizes tuna

species that are exposed to a wider range of environmental conditions and spend significant

periods of time in colder waters (Figure 3). The oceanic and temperate Atlantic bluefin tuna

Figure 3. Three representative ecological lifestyles of tunas and their distribution patterns based on their latitudinal and vertical movements. Characteristic spawning and foraging grounds are shown for each lifestyle; The three ecological lifestyles illustrate a tropical shallow-diving frigate tuna *Auxis thazard*, a tropical deep-diving yellowfin tuna *Thunnus albacares,* and a temperate deep-diving Atlantic bluefin tuna *Thunnus thynnus*. Fish silhouettes represent the depth distribution where species spend most of their time. The thermocline is defined as the depth range within which the water temperature changes rapidly and separates the water column into the upper well-mixed surface layer (water above 20 ℃) and the deeper waters (waters below 15 ℃). Figure modified from Bernal et al. 2009 and Bernal et al 2017.

(*Thunnus thynnus*) are one species that typifies this group, spending most of the year at higher

latitudes in colder and more productive waters between the upper mixed later and the cooler deep

waters below the thermocline, and migrating to warmer waters for spawning (Bernal et al. 2017).

To examine whether our general model could predict these three broad ecological lifestyles of

tunas, we modelled environmental scenarios corresponding to each lifestyle in terms of

temperature, ecosystem productivity, and seasonality (Table 2). We chose parameters based on

the results of our general model, but also added seasonal fluctuations in temperature to represent

the bluefin tuna migration from temperate, productive waters to their tropical spawning grounds.

We then asked whether the life-history traits emerging in these scenarios matched the range of

reported sizes of three representative species corresponding to each lifestyle (Juan-Jordá et al.

2016).

Sensitivity analyses

We ran a series of tests to examine how our choices of parameters in the size spectrum influence

model predictions. Specifically, we varied values of h (representing efficiency in prey capture or

hunger levels) in the predation function (Eq. 3) and φ (the fraction of body mass that can be

devoted to reproduction) in the reproductive constraint (Eq. 9). We also conducted sensitivity

analyses in which we varied the scale and shape coefficients in the metabolic cost function (c and

θ, respectively, in Eq. 5). Motivated by our tuna case study, we also varied seasonality in

resources, temperature, and spawning. We considered seasonal environments in which only

resources and only temperature varied, to understand how these factors alone contributed to

observed variation in body size and reproductive output. We additionally considered growth and

reproductive patterns with an extended warm season (where individuals could spawn in warm

temperatures for six months of the year, instead of three). Finally, we determined in preliminary

analyses that the maximum lifespan T did not strongly affect model results, because most

individuals reach a maximum body size well before the final time T.

Results

An asymptotic growth pattern naturally emerged from the model, after a period of exponential

growth early in life (Fig. 4). Our model predicted age-specific relationships between body size

(length, from which we calculate mass using Eq. 6) and reproduction that correspond to a range

of fish life histories (Fig 4). In general, we found that individuals allocated energy to growth

early in life and shifted this allocation to reproduction in later life. Ecosystem productivity (κ)

alone generated a range of maximum body sizes, from less than 100 cm at low levels of

productivity to well over 350 cm (Fig. 4A). Across scenarios of ecosystem productivity, the

predicted trajectories for individual growth were identical in early life (before age 2); subsequent

growth slowed earlier in lower productivity environments (Fig. 4A). For all life histories,

allocation to reproduction began at very low levels sometime during the individual's second year

(Fig. 4B) and increased steadily as the individual aged. Both the rate at which reproduction

increased with size and the maximum reproductive output were consistently greater with

increased ecosystem productivity (Fig. 4B). We also found the pattern of age-specific mortality

was very similar in early life for all productivities κ, but that lifespan increased predictably as

asymptotic body size increased (Supp. Fig. 5).

Figure 4. (A) Growth (length at age) and (B) Expected reproductive output of individuals in different productivity κ scenarios. Each curve ends when an individual's expected survival chances are less than 3%. Temperature in every scenario was constant at 16.85 °C, and h was constant at 8. The reproductive output was smoothed using the loess tool in R; data points can be found in Supp. Fig. 4.

When we compared the interacting effects of ecosystem productivity (κ) and environmental

temperature (τ) on patterns of growth and reproductive allocation, we found the effects of

temperature on body size and reproductive output were relatively small (Fig. 5). The biggest

differences emerged in highly productive environments with dramatic increases in temperature.

Specifically, when average temperature increased by 10 degrees (e.g., from 11.85 to 21.85 °C or

from 16.85 to 26.85 °C), maximum body sizes were 3-10% smaller, with the largest differences

occurring when $\kappa = 5$ (specific values in Fig. 5 are reported in Supp. Table 1). At the same time,

these increases in temperature led to concomitant reductions in lifetime reproductive output that

ranged from 3-10% (Fig 5B). However, it is important to note that comparing thermal increases

in 5-degree intervals only generated minor (3-5%) decreases in growth and reproduction, unless

both predators and prey were scarce ($\kappa \leq 0.5$, Supp. Table 1).

Figure 5. (A) Maximum body size and (B) Expected lifetime reproductive output (in kg) of individuals plotted against a range of productivity scenarios. Color of points corresponds to differences in average annual temperature in °C.

The interacting effects of temperature and productivity on body size showed that when both prey

and predators were abundant, the benefits of growing large outweighed associated increases in

metabolic costs. Remaining smaller in thermally taxing conditions (to minimize energetic

requirements for maintenance), equated to an increased risk of predation and thus shorter

lifespans. In other words, the effect of temperature on survival was indirectly expressed through

body size, i.e., where temperature influenced asymptotic body size (Fig. 5A), it had a predictable

effect on lifespan.

To understand the robustness of the main results in Figs. 4 and 5, we conducted sensitivity

analyses. We varied the parameter h, representing predator efficiency or hunger level (Eq. 3;

Andersen 2019), to change the allometric properties of the risk of predation (Supp. Fig. 6).

However, while varying this parameter affected the realized lifespans, it did not dramatically

change individual growth trajectories or patterns of size-specific reproduction (Supp. Fig. 6). By

contrast, the constraint on the amount of lipid stores that can be spent each month on

reproduction, φ, was of more importance for growth because it depended directly on the

individual's structural mass. This parameter contributes biological realism to the state dynamics,

but is difficult to measure or estimate directly, especially for fish that spawn multiple times per

year, because often little is known about the frequency of spawning. In sensitivity analyses, we

found that making this constraint more stringent (e.g., decreasing φ from 0.2 to 0.1) led to

increased body sizes, and this pattern held at both high and low productivity values (Supp. Fig.

7). Finally, we used estimates of parameters for the metabolic cost function that were measured

experimentally for tuna species, because of our interest in explaining diversity in tuna growth

patterns (Clarke and Johnston 1999). However, to understand if the relatively small effect of

increased temperatures on body size in Figure 5 was robust to differences in physiological costs,

we conducted sensitivity analyses in which the shape and allometric scaling of costs varied

(Supp. Figs. 8 and 9). We confirmed that our choice of parameters for this function had relatively

minor influences on growth and reproduction.

In the environmental scenarios representing the three ecological lifestyles of tunas (Table 2) our

model predicted growth, body size, and reproductive patterns that are representative of typical

species of each of the three tuna ecological lifestyles (Figure 6). With the environmental scenario

representing the tropical shallow-diving lifestyle, we predicted a maximum body size of 70 cm,

longevity of 6-7 years of age, and continuous reproduction at relatively low levels (left column,

Figure 6, Table 2). These predicted traits are similar to those of frigate tuna *Auxis thazard,* a

tropical shallow-diving tuna species with small body size. With the environmental scenario

matching the tropical deep-diving lifestyle, we predicted growth to sizes well over 200 cm,

lifespans of less than 15 years, and continuous reproductive output that increases over the

individual's life (middle column, Figure 6, Table 2). While we believe this is broadly consistent

Figure 6. Growth (top row) and reproduction (bottom row) predicted by our general energy-based model for the three characteristic ecological lifestyles of tunas using three representative environmental scenarios (columns). Details of each scenario and species are in Table 2. With the exception of C and F, all curves end at the age when less than 3% of the population is expected to survive, representing the cohort lifespan. A and D represent tropical shallow-diving tuna species such as the frigate tuna *Auxis thazard*. B and E represent tropical deep-diving tuna species such as yellowfin tuna *Thunnus albacares*. C and F represent temperate deep-diving species such as Atlantic bluefin tuna *Thunnus thynnus*, which migrate seasonally from higher latitudes with colder and productive waters to less productive and warmer waters. At age 18, more than 10% of individuals in the temperate deep-diving lifestyle were still alive and are likely to live much longer (C, F). As in Fig. 4, the reproductive output in panels D-E was smoothed using the loess function.

530   t

with the life histories of tropical deep-diving tuna species such as yellowfin tuna *Thunnus*

albacares, lifetime reproductive patterns of these batch-spawning species are not well known

(Horswill et al. 2019). In the environmental scenario matching the lifestyle of temperate deep-

diving tunas, such as Atlantic bluefin tuna, *Thunnus thunnus,* we predicted the maximum body

size exceeded 350 cm a longevity that was longer than 18 years (right column, Figure 6, Table

2). Spawning was constrained to be seasonal in tunas in the third ecological lifestyle (i.e.,

spawning could only occur in temperatures above 24 °C, which we specified individuals

experienced for three months of the year).

Analysis of seasonality in temperature and spectrum productivity

The results of the tuna case study yielded an unexpected pattern: when spawning activity,

environmental temperature and ecosystem productivity varied seasonally, i.e., in the third

ecological lifestyle of temperate deep-diving tunas, individuals grew to be substantially larger

than the size observed under constant environment conditions that are otherwise comparable in

terms of productivity (κ) and temperature (τ).

To investigate this pattern further, we made several follow-up comparisons of the effects of

seasonality in spawning season length, temperature (in which species spawned for three months

in waters that are 9 degrees warmer than their winter foraging grounds) and productivity (which

increased threefold during the winter months spent on the foraging grounds). In Figure 7, for a

range of ecosystem productivities ($\kappa = 0.5$ - 2.5 when averaged across the 12-month period), we

compared the maximum lengths predicted in constant environments with scenarios in which

reproductive season length varied (represented by the dash length in Fig 7), and/or temperature

and productivity fluctuated over the year, for a total of 6 contrasting scenarios: (i) the most basic

constant scenario when temperature, productivity (κ), and spawning were constant year round

Figure 7. Maximum body length across values of κ in multiple scenarios varying seasonality in food, temperature, and spawning season duration. As described in the text, solid lines represent scenarios where spawning is possible every month of the year, while the dashed lines represent scenarios where spawning is restricted to three months (short-dashed) and six months (yellow long-dashed) of the year. Colors correspond to different scenarios in which temperature and productivity fluctuate temporally.

(solid aqua line in Figure 7, which is identical to the $\tau = 16.85\ °C$ scenarios shown in Fig. 5); (ii)

a seasonal scenario when temperature increased during the three month spawning season but

productivity κ did not change throughout the year (purple short-dashed line in Fig. 7), (iii) a

seasonal scenario with a threefold increase in productivity for nine months to represent time

spent in foraging grounds, but with temperature kept constant throughout the year (navy blue

short-dashed line in Fig. 7); (iv) the full seasonal scenario where productivity and temperature

varied during the three-month summer when spawning was possible (salmon pink short-dashed

line in Fig 7; note this scenario is similar to Fig. 6C, but with lower values of κ); (v) a full

seasonal scenario where the warm, low-productivity season lasted for six months instead of three

(yellow long-dashed line in Fig. 7); and (vi) a seasonal scenario where productivity and

temperature varied during the three-month summer but spawning was possible during the full

twelve months (solid orange line in Fig. 7).

These comparisons clarified that the restriction of spawning to three months of the year has the

largest effect on maximum body size across a range of ecosystem productivities (because the

short-dashed lines are substantially greater than solid or long-dashed lines in Fig 7). When

spawning was restricted to three months, the effects of temperature and productivity alone were

comparable to those when both effects are combined (in Fig. 7, the three short-dashed lines are

similar). When spawning was restricted to 6 months of the year, and temperature and

productivity fluctuated seasonally, individuals grew to intermediate sizes. Smaller sizes were

predicted when spawning was possible during every month of the year, especially when

temperature and productivity fluctuated seasonally (the solid orange line in Fig. 7). From this set

of follow-up analyses, we inferred that resources and temperature effects on energetic budgets

were not the primary drivers of larger body sizes in our model. Rather, the compression of

reproductive opportunities into three months of the year, and the temporal separation of

reproduction from the most valuable foraging opportunities, favors increased allocation to

growth.

Discussion

We modeled the evolution of growth and reproduction in fish in differing environments, accounting for food web interactions represented by size spectra, and the effects of seasonality in metabolic demands and spawning opportunities. With our general energetic model we also were also able to predict observed life history traits of species representing the three main ecological lifestyles of tunas. By unpacking the effects of temporal variation in spawning opportunities, metabolic demands, and foraging opportunities, we predicted that seasonal cycles in resource intake and expenditures have a greater effect on the evolution of body size than that of temperature on metabolic rate.

Changes in environmental temperature and ecosystem productivity have been occurring for decades and have been shown to affect fish recruitment, growth, and fishery productivity (Free et al. 2019; Oke et al. 2020). There have been several mechanisms proposed to explain these changes (Fernandes et al. 2020). Early models based on differences in resource intake and metabolic demands predicted that increased temperature will lead to reduced growth and smaller body sizes, i.e., 'shrinking fishes' (Cheung et al. 2008), but experimental evidence shows growth is less affected by temperature than reproductive allocation (Wootton et al. 2022), and that metabolic rate can adapt to increased temperatures (Pilakouta et al. 2020). Critically, such physiological models and experimental observations necessarily minimize confounding effects of environmental productivity, seasonality, and ecological interactions. Our model is fundamentally different, as we incorporated all these extrinsic abiotic and ecological factors that previous work has had to ignore. Our general energetic model used insights from size-sspectrum theory to parametrize the availability of prey (foodscape) and the risk of predation (fearscape) for

individuals of different sizes into a model of individual fitness. The minor effect of temperature

on our model results implies that temperature is relatively unimportant compared to foraging and

surviving in the real world.

A key difference between our general energetic model and other models of life history evolution

of fishes in different environmental conditions is that we consider the mass-specific scaling of

metabolic costs with temperature in addition to differences in primary productivity, which drives

food-web interactions. Across a range of temperatures, our model predicted the benefits of

growing large through the foodscape and fearscape outweighed the increase in metabolic costs

until very large sizes and/or very dramatic increases in temperature. For example, although our

model predicted smaller body sizes and decreased lifetime reproduction at higher temperatures

when both predators and prey were scarce, these differences were relatively small until average

temperatures increased by 10-15 °C. Ecosystem productivity κ had a much stronger influence on

our results than any of our other parameters; consistent with expectations, our model predicted

larger maximum body sizes, greater reproductive output, and longer lifespans in more productive

ecosystems. One exception came when estimating age-specific mortality of each environment,

which was more sensitive to our choice of the value representing variation due to predator

satiation or efficiency in prey capture (h, Eq. 3). However, this process did not substantially

change growth trajectories.

Our results provide a novel explanation for asymptotic growth, which does not invoke metabolic

mechanisms or trade-offs between growth and reproduction. Prior work has explained

asymptotic growth patterns with physiological arguments for intrinsic geometric constraints (von

Bertalanffy 1960, Pauly 2010), or by specifying that increased allocation to reproduction limits

allocation to growth, leading to a slowdown in growth rate with age (Jusup et al. 2011). In our

model, multiple (mainly extrinsic) ecological mechanisms led to slowing of growth and

emergence of an asymptotic size in each environment. Unlike previous research, in our model

the onset of reproduction alone does not correspond immediately to decreasing growth. Instead,

the availability of prey, the physical and temporal limits on resource intake, the risk of predation,

and constraints on gonad size and reproductive timing all increase the advantage of growing to

larger body sizes; these advantages are balanced against metabolic demands and the actual

energetic requirements of growth.

We used variation among the ecological lifestyles of tunas as a motivation for our model

parameterizations and comparisons, while also aiming to provide general insights into the

evolution of fish life histories. We successfully predicted growth and reproductive patterns that

are consistent with species representing each ecological lifestyle of tunas (Fig. 6). Unexpectedly,

we found an effect of seasonal variation on body size. Further investigation revealed this effect is

robust to fluctuations in temperature or in productivity alone, and is largely determined by the

spawning season duration. These findings rule out explanations for spatial gradients in body size

based on metabolic costs, and strongly supports the conclusion that growth is driven by the

limited time available for reproduction, coupled with the opportunity to forage and store energy

in productive ecosystems the rest of the year. These results could explain giantism of fishes in

highly seasonal polar environments. For fishes like tunas, which often exhibit a combination of

capital and income breeding, and for which reproduction is determined by temperature, we

expect that as climate change generates warmer conditions, it will eventually allow spawning at

higher latitudes for longer periods. Yet our results suggest that cascading effects on their growth

are not easily predicted and will depend on how warming waters reverberate through the size

spectrum.

Our model illustrates that predicting responses in growth and reproduction in changing

environments is complex, because optimal strategies responding to selective pressures from

many factors, including intrinsic constraints on storage and reproduction, physiology, and the

nature of ecosystem fearscapes and foodscapes. We have made several simplifying assumptions

that could be explored in further investigations. For example, in the analyses presented here, we

have assumed a static relationship between length and mass, and the conversion of mass to joules

(energy density of tissue). Keeping these relationships constant ensured a common currency

linking individual energetic budgets to ecological changes in predation risk and foraging success.

Exploring the consequences of these relationships for differences among clades would be a

natural follow-up to our study. Furthermore, while we did not directly address the interaction

between of fisheries induced selection and growth, it is a rich area of further investigation

(Audzijonyte et al. 2016). Plastic or evolutionary responses to fishing potentially include faster

growth, earlier maturation rates, and smaller maximum body sizes, but the consequences for

marine food webs are complex (Hočevar and & Kuparinen 2021). The framework introduced

here potentially could capture how eco-evolutionary feedbacks between fishing mortality,

growth and mass relationships, and ecosystem size spectra interactively affect trends in body

sizes of focal species. Studies of the interaction between fishing and size spectra have been

focused on lake systems where food webs have been studied in greater detail (Perälä

and Kuparinen 2020). Understanding how these physiological and ecological processes are

mechanistically linked is necessary to understand how species will respond to different

environmental changes and climate change scenarios in the ocean.

In summary, our findings suggest that when predicting future growth patterns under projected

changes in climate, ecological and physiological factors - primary production, prey size and

availability, and predation risk - will all play a greater role than temperature in determining

trends in body size. We appreciate that our conclusion that increasing model complexity reveals

a limited effect of temperature on growth may be unpopular. Nevertheless, we offer our model as

a step toward models of fish growth with higher fidelity to nature that incorporate not only

physiology and energetic allocation budgets but are also embedded in mass-specific foodscapes

and fearscapes represented by size spectra. We hope that further exploration of our approach

may lead to reconciliation of divergent results regarding the effect of climate change on fish

body sizes.

Appendix. Optimization details.

Dynamic programming equations (Eq. 12) are constrained optimization algorithms with the

purpose of determining the optimal set of behaviors or decisions that maximize a quantitative

metric over time, such as lifetime expected reproductive success. A key property is that the

decisions at one time affect the state variables at the next time. When solving Eq. 12, we

consider how the allocation decisions an individual makes during one month of its life affect its

future size, energetic reserves (lipid stores) and chances of survival, and find the set of allocation

to growth and reproduction that maximizes its fitness in a given environment. We are able to

calculate lifetime fitness by solving the equation using backwards iteration. By starting at $t\ =$

$T - 1$, and assuming there is no possibility of future fitness (i.e., the second term of Eq. 12 is

equal to zero for all possible values of length and lipid stores) we can populate the array with the

fitness of all possible combinations of length, lipid stores, and allocation strategies at $t = T - 1$;

this fitness will be the first term on the right side of Eq. 12. This process is repeated, working

backwards, until $t = 1$. We can thereby determine the combination of g and r that maximizes

both current fitness at t, and the fitness expected from $t + 1$ until T, given the chance of survival,

length, and lipid stores resulting from that particular strategy. Solving Eq. 12 in this way

produces an array storing the proportional allocation to growth and reproduction that leads to the

highest lifetime fitness, for all possible combinations of size and lipid stores (l and s) for every

month until the final time T.

Solving a dynamic programming equation with two state variables, such as Eq. 12, is

computationally expensive, and we employed a number of techniques to make the iteration more

efficient and to approximate a smooth fitness surface. First, we constrained the parameter space

that was evaluated. Specifically, we used an integer index I (with a maximum of I_{max}) to

represent lipid stores s, converting the index to values in joules in the dynamic loop. We then

related the range of s we explored to each value of l (because lipid stores are constrained by l),

by setting

716
$$s = \frac{0.6 \cdot I^3 \rho a l_{max}{}^3}{I_{max}{}^3}$$

In other words, we adjusted the numerical step size for possible values of s to be finer for smaller

individuals. Furthermore, we used linear interpolation of state values (Clark and Mangel 2000)

when computing expected future fitness in Eq. 12 to minimize discontinuities on the fitness

landscape arising from the integer index of energetic state. We did not interpolate length, since

its unit (centimeters) was sufficiently fine-grained that there were minimal effects of

discontinuities.

We found that sharp transitions in the fitness landscape, such as at threshold values representing

constraints, hindered our optimization algorithm. Therefore, we used an asymptotic function for

the fitness increment associated with a given level of reproductive output. We let

$R(r, s, l, t)$ denote the potential increment in fitness in month t when $L(t) = l$, $S(t) = s$, and a

fraction of stores r is allocated to current reproduction, and assumed that R increased smoothly

toward the maximum possible for the given length l:

$$R(r, s, l) = \frac{rs}{1 + (rs/\varphi w(l))f}$$

The asymptotic value of this function depends on the value of f in the denominator, which

controls the abruptness of the constraint on current fitness (Supp. Fig. 2). In other words, the

steepness of the multivariate landscape around the fitness optimum is modulated by f. For

simplicity, we assume $f = 1$ for all results presented hereafter. In that case, when $\varphi w(l) >>$

$\frac{rs}{\rho}$, $R(r, s, l) \sim rs$ and when $rs >> \varphi w(l)$, then $R(r, s, l) \sim \varphi w(l)$.

All programming was done using R 4.1.1. We ran the code to solve the dynamic programming

equation and simulate the individual life histories using Rscript commands from the Linux shell

of a 2019 MacPro with 16 cores and 96 GB of RAM. Jobs were run in parallel and the runtime of

each job was between 60 and 120 minutes. The results of each job are presented in the figures

summarizing growth, reproduction, and survival data; these analyses and figures were produced

in the Rstudio IDE.

References

Álvarez-Noriega M., White C.R., Kozłowski J., Day T., Marshall D.J. 2023. Life history optimisation drives latitudinal gradients and responses to global change in marine fishes. PLoS Biol 21(5): e3002114.

Andersen, K. H. 2019. *Fish Ecology, Evolution, and Exploitation: A New Theoretical Synthesis.* Princeton University Press, Princeton.

Andersen, K. H., Jacobsen, N. S., Farnsworth, K. D., and J. Baum. 2016. The theoretical foundations for size spectrum models of fish communities. *Canadian Journal of Fisheries and Aquatic Sciences* 73:575–588.

Audzijonyte, A., Fulton, E., Haddon, M., Helidoniotis, F., Hobday, A.J., Kuparinen, A., Morrongiello, J., Smith, A.D., Upston, J. and Waples, R.S., 2016. Trends and management implications of human-influenced life-history changes in marine ectotherms. *Fish and Fisheries*, *17*(4), pp.1005-1028.

Audzijonyte, A., Richards, S.A., Stuart-Smith, R.D., Pecl, G., Edgar, G.J., Barrett, N.S., Payne, N. and Blanchard, J.L., 2020. Fish body sizes change with temperature but not all species shrink with warming. *Nature Ecology & Evolution*. 4:809-814.

Audzijonyte, A. and Richards, S.A., 2018. The energetic cost of reproduction and its effect on optimal life-history strategies. *American Naturalist*. 192:E150-E162.

Audzijonyte, A., Jakubavičiūtė, E., Lindmark, M., and Richards S.A. 2022. Mechanistic Temperature-Size Rule Explanation Should Reconcile Physiological and Mortality Responses to Temperature. *The Biological Bulletin* 2022 243:2, 220-238

Barnes, W., Maxwell, D., Reuman, D.C., and Jennings, S., 2010. Global patterns in predator-prey size relationships reveal size dependency of trophic transfer efficiency. *Ecology* 91:222-232.

Benoît, E., and M. J. Rochet. 2004. A continuous model of biomass size spectra governed by predation and the effects of fishing on them. *Journal of Theoretical Biology* 226:9-21.

Bernal, D. 2011. Physiological specializations of different fish groups. Pelagic fishes. Pages 1887-1902 *in* Farrell AP (ed) *Encyclopedia of Fish Physiology: From Genome to Environment,* Vol 3. Academic Press, San Diego.

Bernal, D., Sepulveda C., Musyl, M., Brill, R. (2009) The eco- physiology of swimming and movement patterns of tunas, billfishes, and large pelagic sharks. Pages 436–483 *in* Domenici P, Kapoor BG (eds) *Fish Locomotion: an Etho-Ecological Approach*. Science Publishers, Enfield.

788   Bernal, D., Brill, R.W., Dickson, K.A. and H.A. Shiels. 2017. Sharing the water column:
789   physiological mechanisms underlying species-specific habitat use in tunas. *Reviews in Fish*
790   *Biology and Fisheries* 27:843-880.
791

792   Bernhardt, J.R., Sunday, J.M. and M.I. O'Connor. 2018. Metabolic theory and the temperature-
793   size rule explain the temperature dependence of population carrying capacity. *American*
794   *Naturalist.* 192:687-697.
795

796   Beverton, R. J. H., and S. J. Holt. 1959. A review of the lifespans and mortality rates of fish in
797   nature, and their relation to growth and other physiological characteristics. Pages 142–177 *in*
798   CIBA Foundation Colloquium on Ageing.
799

800   Bigman, J.S., M'Gonigle, L., Wegner, N.C. and Dulvy, N.K. 2021. Respiratory capacity is twice
801   as important as temperature in driving patterns of metabolic rate across the vertebrate tree of life.
802   *Science Advances* 7: eabe5163.
803

804   Blanchard, J. L., Heneghan, R. F., Everett, J. D. , Trebilco, R. and A. J. Richardson. 2017. From
805   bacteria to whales: using functional size spectra to model marine ecosystems. *Trends in Ecology*
806   *& Evolution.* 32:174-186.
807

808   Blanchard, J. L., Jennings, S., Law, R., Castle, M.D., McCloghrie, P., Rochet, M.J., and E.
809   Benoît. 2009. How does abundance scale with body size in coupled size-structured food webs?
810   *Journal of Animal Ecology.* 78:270-280.
811

812   Blanchard, J.L., Jennings, S., Holmes, R., Harle, J., Merino, G., Allen, J.I., Holt, J., Dulvy, N.K.
813   and M. Barange. 2012. Potential consequences of climate change for primary production and fish
814   production in large marine ecosystems. *Phil. Trans. Roy. Soc. B* 367:2979-2989.
815

816   Brown, J. H., J. F. Gillooly, A. P. Allen, V. M. Savage, and G. B. West. 2004. Toward a
817   metabolic theory of ecology. *Ecology.* 85:1771-1789.
818

819   Chapman, E. W., Kørgensen, C., and M. E. Lutcavage. 2011. Atlantic bluefin tuna (*Thunnus*
820   *thynnus*): a state-dependent energy allocation model for growth, maturation, and reproductive
821   investment. *Canadian Journal of Fisheries and Aquatic Sciences* 68:1934–1951.
822

823   Charnov, E. L., Berrigan, D., and R. J. H. Beverton. 1991. Dimensionless numbers and the
824   assembly rules for life histories. *Phil. Trans. Roy. Soc. B* 332:41-48.
825

826   Charnov, E. L., and J. R. Krebs. 1974. On clutch-size and fitness. *Ibis.* 116:217-219.
827

828   Cheung, W.W., Close, C., Lam, V., Watson, R., and D. Pauly 2008. Application of
829   macroecological theory to predict effects of climate change on global fisheries potential. *Marine*
830   *Ecology Progress Series.* 365:187-197.
831

Christensen, A. and Andersen, K.H., 2011. General classification of maturation reaction-norm
shape from size-based processes. Bulletin of Mathematical Biology, 73(5): 1004-1027.

Cichoń, M. and Kozłowski, J., 2000. Ageing and typical survivorship curves result from optimal
resource allocation. Evolutionary Ecology Research, 2(7): 857-870.

Clark, C. W., and M. Mangel. 2000. Dynamic state variable models in ecology: methods and
applications. Oxford University Press, Oxford.

Clarke, A. 2006. Temperature and the metabolic theory of ecology. *Functional Ecology* 20:405-
412.

Clarke, A., and N. M. Johnston. 1999. Scaling of metabolic rate with body mass and temperature
in teleost fish. *Journal of Animal Ecology*. 68: 893-905.

Collette, B.B., Reeb, C. and Block, B.A., 2001. Systematics of the tunas and mackerels
(Scombridae). *Fish Physiology*, 19, pp.1-33.

Conover, D. O., and S. B. Munch. 2002. Sustaining fisheries yields over evolutionary timescales.
Science 297:94–96.

Daufresne, M, Lengfellner, K., and U. Sommer. 2009. Global warming benefits the small in
aquatic ecosystems. *Proc. Nat. Acad. Sci.* 106:12788–12793.

Dickson, K.A., and Graham, J.B. 2004. Evolution and consequences of endothermy in fishes
Physiological and Biochemical Zoology 77: 998-1018.

Ejsmond, M.J., Czarnołęski, M., Kapustka, F. and Kozłowski, J., 2010. How to time growth and
reproduction during the vegetative season: an evolutionary choice for indeterminate growers in
seasonal environments. *The American Naturalist* 175(5): 551-563.

Fernandes, J.A., Rutterford, L., Simpson, S.D., Butenschön, M., Frölicher, T.L., Yool, A.,
Cheung, W.W. and Grant, A., 2020. Can we project changes in fish abundance and distribution
in response to climate? *Global Change Biology.* 26:3891-3905.

Free, C.M., Thorson, J.T., Pinsky, M.L., Oken, K.L., Wiedenmann, J. and Jensen, O.P. 2019.
Impacts of historical warming on marine fisheries production. *Science.* 363: 979-983.

Froese R. 2006. Cube law, condition factor and weight-length relationships: history, meta-
analysis and recommendations. *Journal of Applied Ichthyology* 22: 241-253

Gadgil, M., and W. H. Bossert. 1970. Life historical consequences of natural selection. *American
Naturalist* 104:1-24.

Giacomini, H.C., Shuter, B.J. and Lester, N.P., 2013. Predator bioenergetics and the prey size
spectrum: Do foraging costs determine fish production? *Journal of Theoretical Biology* 332:
249-260.

Gillooly, J. F., Brown, J. H., West, G. B., Savage, V. M., and E. L. Charnov. 2001. Effects of
size and temperature on metabolic rate. *Science.* 293:2248–2251.

Gislason, H., Daan, N. Rice, J. C., and J. G. Pope. 2010. Size, growth, temperature and the
natural mortality of marine fish. *Fish and Fisheries* 11:149–158.

Hilborn, R. and Mangel, M. 1997. *The Ecological Detective.* Princeton University Press,
Princeton, NJ.

Hočevar, S., & Kuparinen, A. (2021). Marine food web perspective to fisheries-induced
evolution. *Evolutionary Applications*, 14, 2378– 2391.

Horswill, C., H. K. Kindsvater, M. J. Juan-Jordá, N. K. Dulvy, M. Mangel, and J. Matthiopoulos.
2019. Global reconstruction of life-history strategies: A case study using tunas. *Journal of
Applied Ecology* 56:855-865.

Houston, A., C. Clark, J. McNamara, and M. Mangel. 1988. Dynamic Models in Behavioural
and Evolutionary Ecology. *Nature.* 332:29-34

Houston, A., and J. McNamara. 1999. *Models of Adaptive Behaviour*. Cambridge University
Press.

Hulthén, K., Hill, J.S., Jenkins, M.R. and Langerhans, R.B., 2021. Predation and resource
availability interact to drive life-history evolution in an adaptive radiation of livebearing
fish. *Frontiers in Ecology and Evolution*, 9, p.619277.

Jørgensen, C., K. Enberg, and M. Mangel. 2016. Modelling and interpreting fish bioenergetics: a
role for behaviour, life-history traits and survival trade-offs. *Journal of Fish Biology* 88:389–
402.

Jørgensen, C., and Ø. Fiksen. 2006. State-dependent energy allocation in cod (*Gadus morhua*).
Canadian Journal of Fisheries and Aquatic Sciences 63:186–199.

Juan-Jordá, M. J., I. Mosqueira, J. Freire, and N. K. Dulvy. 2013. The Conservation and
Management of Tunas and Their Relatives: Setting Life History Research Priorities. *PLoS ONE*
8:e70405.

Juan-Jordá, M.J., Mosqueira, I., Freire, J., Ferrer-Jordá, E. and Dulvy, N.K. 2016. Global
scombrid life history data set. *Ecology*, 97: 809-809.

Jusup, M., Klanjscek, T., Matsuda, H., Kooijman, S.A.L.M. 2011. A Full Lifecycle Bioenergetic
Model for Bluefin Tuna. *PLoS ONE* 6(7): e21903.
923

Kindsvater, H.K., Alonzo, S.H., Mangel, M. and Bonsall, M.B., 2010. Effects of age-and state-
dependent allocation on offspring size and number. *Evolutionary Ecology Research*, 12(3),
pp.327-346.

Kingsolver, J. G., and R. B. Huey. 2008. Size, temperature , and fitness: three rules. *Evolutionary*
Ecology Research 10:251–268.

Kiørboe, T., and A. G. Hirst. 2014. Shifts in mass scaling of respiration, feeding, and growth
rates across life-form transitions in marine pelagic organisms. *American Naturalist* 183:E118-
E130.

Kitchell, J.F., Neill, W.H., Dizon, A.E., and J.J. Magnuson. 1978. Bioenergetic Spectra of
Skipjack and Yellofin Tunas. In:Sharp, G.D., Dizon, A.A. (Eds.), *The Physiological Ecology of*
Tunas. Academic Press.

Kozłowski, J., 1992. Optimal allocation of resources to growth and reproduction: implications
for age and size at maturity. *Trends in Ecology and Evolution*, 7(1):15-19.

Kozlowski, J. 1996. Optimal allocation of resources explains interspecific life-history patterns in
animals with indeterminate growth *Proc. R. Soc. B.* 263: 559-566.

Kozłowski, J. and Teriokhin, A.T., 1999. Allocation of energy between growth and reproduction:
the Pontryagin Maximum Principle solution for the case of age-and season-dependent mortality.
Evolutionary Ecology Research 1(4): 423-441

Law, R., M. J. Plank, A. James, and J. L. Blanchard. 2009. Size-spectra dynamics from
stochastic predation and growth of individuals. *Ecology* 90:802-811.

Lika, K. and Kooijman, S.A., 2003. Life history implications of allocation to growth versus
reproduction in dynamic energy budgets. Bulletin of Mathematical Biology, 65(5), pp.809-834.

Oke, K.B., Cunningham, C.J., Westley, P.A.H., Baskett, M.L., Carlson, S.M., Clark, J., Hendry,
A.P., Karatayev, V.A., Kendall, N.W., Kibele, J. ,Kindsvater, H.K., Kobayashi, K.M., Lewis, B.,
Munch, S., Reynolds, J.D., Vick, G.D., and E.P. Palkovacs. 2020. Recent declines in salmon
body size impact ecosystems and fisheries. *Nature communications*, 11:1-13.

Perälä, T, Kuparinen, A. Eco-evolutionary dynamics driven by fishing: From single species
models to dynamic evolution within complex food webs. *Evol Appl*. 2020; 13: 2507– 2520.

Perrin, N. and Sibly, R.M., 1993. Dynamic models of energy allocation and investment. Annual
Review of Ecology and Systematics, 24: 379-410.

Perrin, N., Sibly, R.M. and Nichols, N.K., 1993. Optimal growth strategies when mortality and
production rates are size-dependent. Evolutionary Ecology, 7(6), pp.576-592.

Mangel, M. 2006. The Theoretical Biologist's Toolbox : Quantitative Methods for Ecology and
Evolutionary Biology. Cambridge University Press.
971
972 Mangel, M. 2015. Stochastic dynamic programming illuminates the link between environment,
973 physiology, and evolution. *Bulletin of Mathematical Biology* 77:857-877.
974
975 Mangel, M. and C. W. Clark. 1988. *Dynamic Modelling in Behavioral Ecology.* Princeton
976 University Press, Princeton.
977
978 Monsch, K.A. 2000 *The Phylogeny of the Scombroid Fishes.* PhD thesis, University of Bristol p
979 272.
980
981 Neubauer, P., and K. H. Andersen. 2019. Thermal performance of fish is explained by an
982 interplay between physiology, behaviour and ecology. *Conservation Physiology* 7:coz025
983
984 Pauly D. 2010. Gasping fish and panting squids: oxygen, temperature and the growth of water-
985 breathing animals. International Ecology Institute, Germany.
986
987 Pilakouta, N, Killen, SS, Kristjánsson, BK, et al. 2020. Multigenerational exposure to elevated
988 temperatures leads to a reduction in standard metabolic rate in the wild. *Funct Ecol*. 34:1205-
989 1214.
990
991 Pignalosa, P., Pappalardo, L., Gioacchini, G., Carnevali, O. 2020. Length-weight relationships
992 and a new length conversion factor for Atlantic Bluefin tuna (*Thunnus thynnus L.*) caught in the
993 Mediterranean Sea. *Collect. Vol. Sci. Pap. ICCAT,* 76(2): 138-146.
994
995 Quince, C., Abrams, P.A., Shuter, B.J. and Lester, N.P., 2008. Biphasic growth in fish I:
996 theoretical foundations. *Journal of Theoretical Biology*, 254(2): 197-206.
997
998 Schaefer, K.M. 2001. Reproductive biology of tunas. *Fish Physiology.* 19: 225-270.
999
1000 Schaefer K.M., Fuller D.W., and B.A. Block 2009. Vertical movements and habitat utilization of
1001 skipjack (*Katsuwonus pelamis*), yellowfin (*Thunnus albacares*), and bigeye (*Thunnus obesus*)
1002 tunas in the Equatorial Eastern Pacific Ocean, ascertained through archival tag data. Pages 121-
1003 144 *in* Nielsen, J.L., Arrizabalaga, H., Fragoso, N., Hobday, A., Lutcavage, M., Sibert, J. (Eds.),
1004 *Tagging and Tracking of Marine Animals with Electronic Devices*. Springer, New York.
1005
1006 Sheldon, R.W., Prakash, A. and Sutcliffe Jr., W.H. 1972. The distribution of particles in the
1007 ocean. *Limnology and Oceanography.* 17:327-340.
1008
1009 Sheldon, R.W., Sutcliffe Jr., W.H., and M.A. Paranjape. 1977. Structure of pelagic food chain
1010 and relationship between plankton and fish production. *Journal of the Fisheries Research Board
1011 of Canada*. 34: 2344-2353.
1012

Shuter, B.J., Giacomini, H.C., de Kerckhove, D. and Vascotto, K., 2016. Fish life history
dynamics: shifts in prey size structure evoke shifts in predator maturation traits. *Canadian*
Journal of Fisheries and Aquatic Sciences. 73(4): 693-708.

Southwood, T.R.E. 1977. Habitat, the Templet for Ecological Strategies? *Journal of Animal*
Ecology. 46:337-365.

Sprules, W. G., and L. E. Barth. 2016. Surfing the biomass size spectrum: some remarks on
history, theory, and application. *Canadian Journal of Fisheries and Aquatic Sciences* 73:477-
495.

Thygesen, U.H., Farnsworth, K.D., Andersen, K.H., Beyer, J.E. 2005. How optimal life history
changes with the community size-spectrum. *Proc. R. Soc. B*. 272:1323-1331.

Trebilco, R., Baum, J. K., Salomon, A. K., and N. K. Dulvy. 2013. Ecosystem ecology: Size-
based constraints on the pyramids of life. *Trends in Ecology & Evolution*. 28:423-431.

Ursin, E. 1973. On the prey size preferences of cod and dab. *Meddelelser*
fra Danmarks Fiskeri- og Havundersøgelser 7: 85–98.

Varpe, Ø., 2017. Life history adaptations to seasonality. *Integrative and comparative*
biology, *57*(5), pp.943-960.

von Bertalanffy L. 1960. Principles and theory of growth. Pages 137-259 *in* Nowinski, W.W.,
Ed., *Fundamental Aspects of Normal and Malignant Growth*, Elsevier, Amsterdam.

Walsh, M. R., and D. N. Reznick. 2009. Phenotypic diversification across an environmental
gradient: a role for predators and resource availability on the evolution of life histories. *Evolution*
63:3201–3213.

White, C.R., Alton, L.A., Bywater, C.L., Lombardi, E.J. and Marshall, D.J., 2022. Metabolic
scaling is the product of life-history optimization. *Science*, *377*(6608), pp.834-839.

Wong, S., Bigman, J.S., and N.K. Dulvy. 2021. The metabolic pace of life histories across fishes.
Proc. R. Soc. B. 288:20210910.

Wootton, H.F., Morrongiello, J.R., Schmitt, T. & Audzijonyte, A. 2022. Smaller adult fish size
in warmer water is not explained by elevated metabolism. Ecology Letters, 25, 1177– 1188.

Yanco, S.W., Pierce, A.K. and Wunder, M.B. 2022. Life history diversity in terrestrial animals is
associated with metabolic response to seasonally fluctuating resources. Ecography. 2022:e05900
1054

1055   **Table 1**. Description of parameters and variables

Parameter or Variable	Description	Value
w	Body mass in kg	-
N	Number of individuals in a size category (or trophic level) when considering a Sheldon size spectrum	-
B	Absolute biomass in a size category (or trophic level) when considering a Sheldon size spectrum	-
κ	The spectrum parameter, which defines the total biomass of organisms of the smallest body size w in a given ecosystem; Andersen (2019) gives an estimate of 10 gained by averaging over all Predator-Prey Mass Ratio estimates measured from gut contents. We vary it to represent ecosystem differences in overall ecosystem productivity	Varies
$Bprey$	Biomass of prey expected by a focal individual	-
μ_p	Instantaneous risk of mortality due to predation, which depends on body mass and position in the size spectrum	-
h	Predation risk, comprised of predator satiation estimates (estimated from gut contents) and predator preference (or effectiveness) for consuming prey of a given mass (Andersen 2019)	4, 8
τ	Temperature of the environment (in degrees Kelvin except where noted)	285-300
C	Metabolic requirements (costs) that scale with mass and temperature	-
c	Normalization constant scaling metabolic costs (in J), based on metabolic rate data from tunas (Kitchell et al. 1978)	5×10^{16}
k_B	Boltzmann constant, relating particle energy to temperature in units of m^2 kg s^{-2} K^{-1}	1.3×10^{-23}
E	The average activation energy for the rate limiting enzymes in metabolism in units of joules; from the metabolic theory of ecology (Gillooly et al. 2001).	1.04×10^{-19}
θ	Metabolic scaling exponent; values vary among clade, here we use a value reported for tunas (Clarke and Johnston 1999)	0.66
ρ	The energy density of tuna body mass in our model in J/kg (estimated empirically and reported in Chapman et al. 2011)	4.2×10^6
t	Time in monthly time steps in the dynamic model	-
T_{max}	Maximum lifespan in months	216
$l, L(t)$	Body length (in cm) This is a dynamic state variable but can only increase with time. The maximum value possible is 400 cm. For a specific value, we use the capital letter notation.	-
$s, S(t)$	Lipid stores (in joules) – this is a dynamic state variable representing energy stores that can be used for metabolism, growth, and reproduction.	-
a	Scale coefficient relating length to structural mass, similar to values estimated empirically for bluefin tuna and reported in Pignalosa et al. (2020).	1.0×10^{-5}
$w(t)$	Structural mass of the individual (in kg) at time t – this depends on $L(t)$	-
υ	The fraction of structural mass that determines the critical threshold of energetic mass needed for survival	0.1
φ	The fraction of structural mass that determines the maximum possible reproductive output in a monthly time step	0.2
γ_{pred}	Survival from predation from one month to the next	
γ_s	Survival, given that the individual has sufficient energy reserves to meet metabolic requirements and avoid starvation	
g	Proportion of lipid stores allocated to growth (this allocation decision is optimized by the dynamic programming equation); can take values between 0 and 1	
r	Proportion of lipid stores allocated to reproduction (this allocation decision is optimized by the dynamic programming equation); can take values between 0 and 1 and the sum of g and r cannot exceed one.	
$R(l, s, t)$	Current fitness, the product of lipid stores and the optimal allocation to reproduction, r	
$V(l, s, t)$	Expected accumulated lifetime reproduction from time t onwards, given that $L(t) = l$ and $S(t) = s$	

1056
1057
1058
1059

1060 **Table 2**

1061

1062 Parameter values for environmental scenarios corresponding to the three ecological lifestyles of
1063 tunas described in Fig. 3, with corresponding predicted body size and maximum observed fork
1064 length (cm) for three representative species (data from Juan-Jordá et al. 2016).

1065

Lifestyle	Environment	Baseline or winter temp	Yearly mean κ	*h*	Predicted body size	Representative species	Reported range of body sizes
Tropical shallow	Constant	26.85 C	0.1	8	62 cm	Frigate tuna	40-60 cm
Tropical deep diving	Constant	21.85 C	1	8	214 cm	Yellowfin tuna	125-231 cm
Temperate deep diving	Seasonal	11.85	2.5	12	361 cm	Bluefin tuna	203-427 cm

1066
1067

1068   **Supplemental Table 1.** Model outputs corresponding to points in main text Figure 5. All
1069   scenarios had constant environments.

Temperature τ in °C	κ	Lifetime reproductive output (kg)	Maximum body length (cm)
11.85	0.25	2.15	109.5
11.85	0.5	5.87	173
11.85	1	15.51	224
11.85	2	38.94	289.1
11.85	5	123.77	390.4
16.85	0.25	2.08	101.5
16.85	0.5	5.9	159
16.85	1	15.2	222
16.85	2	37.97	289
16.85	5	121.95	384.3
21.85	0.25	2.02	98.4
21.85	0.5	5.65	159
21.85	1	14.73	214
21.85	2	36.93	282
21.85	5	119.28	380.3
26.85	0.25	1.85	92.4
26.85	0.5	5.33	151
26.85	1	13.85	202
26.85	2	35.15	274
26.85	5	115.73	373.4

1070

Supplementary Figure 1. Main panel: The sigmoid relationship between lipid stores (in joules) and survival for individuals of different lengths. The threshold amount of energy required for metabolic maintenance is a fixed proportion of body mass. The probability of avoiding starvation $\gamma_s(s, l)$ is a logistic function of mass given in Eq. 11. Inset: closeup of the x-axis in the region encompassing the smaller threshold.

Supplementary Figure 2. The fitness increment expected with different amounts of lipid reserves (energy stores) Left panel: Fitness increment for an individual of 100 cm. Right panel: Fitness increment for an individual of 250 cm. We show the shape of the function for three values of f but for all analyses here we used $f = 1$.

1095
1096
1097
1098 **Supplementary Figure 3.** Example rules for allocation to growth and reproduction for ages 1 and 16. The inviable region (dark blue) represents parameter space where remaining stores are insufficient for survival, given the individual's length (in other words, energetic requirements for maintenance are not met). Comparing panels within rows shows differences between allocation to growth and reproduction as a function of state and length. Comparing panels within columns shows how the tradeoff between growth and reproduction changes with age. Note that the striations come from the discretization of allocation values of g and r in Eq. 12.

1099

1100
1101

Supplementary Figure 4. Top: The emergent pattern of survival to age at different levels of κ. Bottom: The natural log of survival to age. Dashed horizonal line shows the cutoff past which less than 3% of individuals will survive. For all results here, $h = 8$ and $\tau = 16.85°C$

1102
1103
1104

Supplementary Figure 5. The emergent pattern of ln(survival) to age at different levels of κ. Dashed horizonal line shows the cutoff past which less than 3% of individuals will survive. For all results here, $h = 8$ and $\tau = 16.85°C$.

Age (years)

Supplementary Figure 6.
The emergent patterns of growth, reproduction, and survival that emerge when h varies. We contrast two values of κ to highlight that it is still the main driver of variation in body size, although h affects lifespan. In all scenarios, $\tau = 16.85°C$. Note that for results presented elsewhere, $h = 8$ except where noted.

1108

Supplementary Figure 7. Growth curves showing the effect of changes in the reproductive capacity φ, for two values of κ. As the reproductive constraint increases (the fraction of the body cavity that can be devoted to gonad decreases from 0.2 to 0.1), selection favors the evolution of larger body sizes. In all scenarios, $\tau = 16.85°C$. Note that for all results presented elsewhere, $\varphi = 0.2$.

1119
1120

A

Supplementary Figure 8.
Body size and lifetime reproduction for several values of κ and θ (the allometric exponent in the metabolic cost function). For all scenarios, temperature is constant year-round at 21.85°C.

B

1129
1130
1131

Supplementary Figure 9. Body size and lifetime reproduction for several values of κ and c (the normalization constant in the metabolic cost function). The estimate of c used elsewhere (5×10^{16}) comes from metabolic rate data from tunas (Clarke and Johnston 1999). For all scenarios, temperature is constant year-round at 21.85°C.