Long-term effect of forest harvesting on boreal species assemblages under climate change

Ilhem Bouderbala[1,2,4,]*, Guillemette Labadie[2], Jean-Michel Béland[3], Junior A. Tremblay [5],
Yan Boulanger[3], Christian Hébert[3], Patrick Desrosiers[1,4,6], Antoine Allard[1,4], Daniel
Fortin[2,4]

August 21, 2022

[1] Département de physique, de génie physique et d'optique, Université Laval, Québec (Qc), Canada

[2] Département de biologie, Université Laval, Québec (Qc), Canada

[3] Centre de foresterie des Laurentides, Service canadien des forêts, Québec (Qc), Canada

[4] Centre interdisciplinaire en modélisation mathématique de l'Université Laval, Québec (Qc), Canada

[5] Direction générale des sciences et de la technologie, Environnement et Changement Climatique Canada, Québec
(Qc), Canada

[6] Centre de recherche CERVO, Québec (Qc), Canada

1   **Abstract**

Habitat changes due to human exploitation of natural resources impact ecosystems worldwide, notably

by causing pronounced changes in abiotic and biotic conditions, which modify biological diversity at

the taxonomic, phylogenetic, and functional levels. In forest ecosystems, logging is the main human

disturbance impacting biodiversity. The interplay between forest harvesting and biodiversity varies,

however, with climate change through complex relationships that remain poorly documented. Our aim

was to study the expected long-term variations in the assemblage of bird and beetle communities following

modifications in forest management under different climate change scenarios. We develop empirical

models predicting the distribution of 104 animal species in eastern Canadian boreal forest over the next

century. We predict that forest harvesting will alter the diversity of both taxa by increasing assemblage

*Corresponding author

E-mail: ilhem.bouderbala.1@ulaval.ca

dissimilarity under all the studied climate scenarios, mainly due to species turnover. Regionally, an overall increase in occupancy probability is projected for bird species, whereas a decrease is predicted for beetles, a variation that could reflect differences in ecological traits between taxa. Our results further predict a decrease in the number of species that increase their occupancy after harvest under the most severe climatic scenario for both taxa. We anticipate that under severe climate change, increasing forest disturbance will modify species habitat associated with both young and old habitats.

Keywords: Harvest disturbance, Biodiversity, Climate change, habitat loss, assemblage dissimilarity, boreal forests

1   Introduction

Biodiversity is facing an unprecedented rate of change due to human actions (Sala et al., 2000; Newbold et al., 2015). For terrestrial ecosystems, land-use changes through forest harvesting, agriculture or urbanization are considered the main causes of biodiversity losses and ecosystem services reductions (Sala et al., 2000; Naeem et al., 2012; Bichet et al., 2016; Matuoka et al., 2020). Human exploitation of natural resources can influence species through their mortality, force displacement and biotic interactions, modify their life-history traits, change their morphology and physiology, and even their polymorphic forms (Sergio et al., 2018). Moreover, the exploitation of natural habitat for human activities generate habitat loss and biome fragmentation, which constrains the size of natural populations (Coelho et al., 2020). For instance, forest harvesting activities bring back the stands to an early successional stage and leads to an alteration in local animal communities by favoring species associated with early-succession forests at the expense of late-succession species (Cyr et al., 2009; Fortin et al., 2011; Boucher et al., 2017).

Climate change is the second most important driver of change in biodiversity after land use, for the terrestrial ecosystems (Sala et al., 2000; Dobson et al., 2021). Climate change may impact species indirectly by triggering natural disturbances such as wildfire, which strongly alter forest structure and composition. It may also impact species directly by changing species occurrence or abundance, when climatic conditions go out of their tolerance limits, due to modifications in temperature and precipitation conditions (Micheletti et al., 2021; Bouderbala et al., 2022), which might reshape community composition by species turnover (Bouderbala et al., 2022). For example, the displacement of species to higher elevations is one of the predicted consequences of climate change to overcome the potential increase in temperature (Davis and Shaw, 2001). However, climate change could have implications on the abundance of species with low dispersal capacity

that are unable to follow the climate conditions to which they are currently adapted (Davis and Shaw, 2001; Muluneh, 2021) and could lead to increase in extinction risks (Thomas et al., 2004).

Although modeling biodiversity changes is becoming essential to predict the effects of human actions on the global environment, many challenges remain because there are many related issues that need to be integrated (Ceaușu et al., 2021). To achieve conservation objectives, we want to identify the most appropriate forest harvesting scenario minimizing biodiversity changes. This would help identify more targeted conservation actions under climate change such as vegetation restoration.

Despite the number of studies made to better understand the complex implications of the different types of disturbances on biodiversity (Sala et al., 2000), there are considerable uncertainties over the magnitude of changes in future animal communities Almeida-Rocha et al. (2020). The cumulative effect of anthropogenic disturbances could provoke complex effects on biodiversity through the adjustment in species' ecological traits, which will have cascading repercussions from individual changes to community-level responses (Sergio et al., 2018). However, most literature evoking the modeling of biodiversity changes considered the assumption of no interactions among the various drivers (Sala et al., 2000). In this work, we investigate how the interaction between forest harvesting and climate change would impact biodiversity. Specifically, we examine future implications of forest harvesting levels on the variation of bird and beetle assemblages in Québec's boreal forest under different climate change scenarios. We aim first to quantify the extent of assemblages' alteration by increasing harvest levels under different climate change scenarios and second to determine the main drivers. Moreover, we study how species associated with different habitat classes will be affected by the combined anthropogenic and natural disturbances.

2   Materials and methods

2.1   Study area and occurrence data

The study area was located in the Côte-Nord region of Québec, Canada ($48°N$ to $53°N$, $65°W$ to $71°W$) within an area of 114,118 km^2 (Fig. 1). This region is part of the eastern spruce-moss subdomain and is dominated by black spruce (*Picea mariana* [Mill.] BSP) with balsam fir (*Abies balsamea* [L.] Miller). Forest harvesting had been the main source of anthropogenic disturbances since the late 1990s (Bouchard and Pothier, 2011). Moreover, This area is mainly affected by two major natural disturbances: spruce budworm outbreak (SBW) (*Choristoneura fumiferana*) and frequent wildfires (Boucher et al., 2017; Labadie et al., 2021).

Figure 1: Study area with the presence-absence distribution of the two taxa.

We used presence-absence data collected between 2004 and 2018 to model species distribution of two taxa: birds and beetles. For birds, we used the data from the Second Atlas of Breeding Birds of Québec (Atlas des oiseaux nicheurs du Québec, 2018), based on species occurrence detected using unlimited distance 5-minute point counts (Bibby et al., 2000) collected during the breeding season, from late May to mid-July. For beetles, we merged different databases collected in 2004, 2005, 2007, 2011 (Janssen et al., 2009; Légaré et al., 2011; Bichet et al., 2016). We added 54 sites sampled in 2018 in the northern part of the territory, along the northeast principal road going to Labrador. Insect sampling involved one multidirectional flight-interception trap per site to capture the flying beetles, and four meshed pitfall traps per site to sample beetles that move on the soil surface (Janssen et al., 2009; Bichet et al., 2016). Data from the sites heavily impacted by the SBW outbreak (Ministère des Forêts de la Faune et des Parcs, 2018) were removed to focus only on the effect of the wildfires. We used for this purpose a cumulative index of defoliation severity (Labadie et al., 2021) (we removed 4% and 7% of sampling sites for beetle and bird databases, respectively).

2.2   The forest management scenarios

Three harvest levels were considered: no harvest activity (NoHarvest), 8% of the areas harvested over 10 years (Harvest) – the typical harvest rate for the study area – and half this level (i.e., 4%, $Harvest_{0.5}$). The harvest levels were combined with three climate scenarios: baseline level, radiative forcing associated with the Representative Concentration Pathway scenario (RCP) 4.5, which involved a 3°C increase in mean temperature from 2000 to 2100; and RCP 8.5 associated with a 7.5°C increase over the same period. We

87 compared the response of species occurrence and occupancy between different harvest intensities under

88 three climate scenarios, which explains why we had three reference scenarios (Table 1). Also, we took into

89 consideration in our simulations the northern limit of territorial forest attributes of harvest activities. Beyond

90 this limit, no forest management was authorized by the Ministry of Forests, Wildlife, and Parks (Ministère

91 des Forêts de la Faune et des Parcs, 2018).

Table 1: The studied scenarios. We considered three of them for reference (Climate=Baseline, Harvest=NoHarvest; Climate=RCP 4.5, Harvest=NoHarvest; Climate=RCP 8.5, Harvest=NoHarvest) and compared them to six other scenarios. Under each climate scenario (Baseline, RCP 4.5 and RCP 8.5) we compared NoHarvest to Harvest$_{0.5}$ and Harvest, which makes six comparisons in total.

Reference scenario		Simulated scenario		Compared landscapes
Climate	Harvest	Climate	Harvest	(Reference-Simulated)
Baseline	NoHarvest	Baseline	Harvest$_{0.5}$	Baseline (NoHarvest-Harvest$_{0.5}$)
Baseline	NoHarvest	Baseline	Harvest	Baseline (NoHarvest-Harvest)
RCP 4.5	NoHarvest	RCP 4.5	Harvest$_{0.5}$	RCP 4.5 (NoHarvest-Harvest$_{0.5}$)
RCP 4.5	NoHarvest	RCP 4.5	Harvest	RCP 4.5 (NoHarvest-Harvest)
RCP 8.5	NoHarvest	RCP 8.5	Harvest$_{0.5}$	RCP 8.5 (NoHarvest- Harvest$_{0.5}$)
RCP 8.5	NoHarvest	RCP 8.5	Harvest	RCP 8.5 (NoHarvest-Harvest)

2.3   Forest landscapes simulation

We simulated forest succession across landscapes using LANDIS-II (Scheller and Mladenoff, 2004), a forest

landscape model that captures forest succession across landscapes as an emergent property of both stand and

landscape-scale processes (Scheller et al., 2007). Specifically, we used the LANDIS-II Biomass Succession

extension v5.0, Base Fire v4.0, BDA v4.0, together with the Biomass Harvest v5.0 (Scheller and Mladenoff,

2004) to simulate forest succession, fire (Scheller and Domingo, 2005) and spruce budworm outbreak distur-

bances as well as harvesting in each 250-m cell at a 10-year time step. Outbreaks of SBW were simulated

using the LANDIS-II Biological Disturbance Agent (BDA) extension v3.0 (Sturtevant et al., 2004, 2012).

The Biomass Succession extension simulates modifications in cohort aboveground biomass (AGB) over time

by taking into consideration cohort age, life-history traits, and land type responses of individual tree species.

A random forest model was used to predict crown closure covariate by using Canadian National Forest

Inventory (NFI) forest attribute maps (Beaudoin et al., 2014). These maps are a k-Nearest Neighbours in-

terpolation of the NFI photoplot data acquired in 2001 and are depicting over 130 forest attributes including

species-specific biomass, stand age and crown closure at a 250-m resolution (see Beaudoin et al., 2014). We

therefore build a random forest model predicting cell-level crown closure in NFI product from NFI species-

specific biomass as well as stand age. This model had very high goodness of fit ($R^2 = 0.86$). The model was

then applied on LANDIS-II outputs to predict crown closure all along the simulation, for each cell by using

simulated species specific biomass and stand age (Labadie et al., 2022). Using species group and predicted

¹¹⁰ crown closure, we created five land cover classes from the Earth Observation for Sustainable Development

¹¹¹ of Forests (EOSD) Land Cover Classification Legend (Beaubien et al., 1999; see Table 2). See Labadie et al.

¹¹² (2022) for more details regarding the spatially-explicit forest simulation model with LANDIS-II.

Table 2: Natural habitats used (if less than 50% of the pixel's surface is in disturbance).

Habitat	Condition
Dense conifer	- vegetation exceeds more than 50% of the surface
	- canopy density is higher than 60% of the surface
	- coniferous stands occupy more than 75% of the surface
Open conifer	- vegetation exceeds more than 50% of the surface
	- canopy density is less than 60% of the surface
	- coniferous stands occupy more than 75% of the surface
Mixed-wood	- vegetation exceeds more than 50% of the surface
	- neither coniferous nor broad-leaf trees account for more than 75% of the surface
	- the vegetation surface with trees is greater than the surface without trees
Open habitat	- vegetation exceeds more than 50% of the surface
	- the vegetation surface with trees is less than the surface without trees
Other	- Non vegetation (trees, shrubs, grasses, bryophytes) that exceeds more than 50% of the surface

2.4   The predictor variables

¹¹⁴ Land cover associated with sampling sites was determined from the Canadian National Forest Inventory

¹¹⁵ referenced in the year 2001 (Beaudoin et al., 2014), updated yearly for subsequent fires and cuts. The

¹¹⁶ landscape was split into 11 land cover types. Specifically, we considered land covers with stand age greater

¹¹⁷ than 50 years, i.e., closed-canopy conifer forest, open-canopy mature conifer forest, mixed forest, and open

¹¹⁸ area (see Table 2), together with six disturbance classes, i.e, fire and forest harvesting subdivided by age

¹¹⁹ classes: 0–10, 11–20 and 21–50 years (Labadie et al., 2021). The category of the other types of habitat was

¹²⁰ not included in the models to avoid collinearity. We used 20 pixels (231 m resolution) centered on the focal

¹²¹ pixel (i.e., 21 pixels in total) to calculate the frequencies of the land cover classes with a radius of 2.5 pixels

¹²² (~0.58 km). This radius was chosen based on Bichet et al. (2016) and Zhao et al. (2013) who reported

¹²³ that the influence of landscape composition varied between 400 m for beetles to 1 km for birds. We also

¹²⁴ considered the distance to nearest burned areas as well as the stand age as potential predictor variables.

2.5   Association between bird species and habitat type

¹²⁶ We considered three land cover classes to calculate species winning conditional probability (WCP), i.e., an

¹²⁷ increase in the probability of species occupancy from uncut to post-harvest in 2100, conditional to species

¹²⁸ habitat (see Eq. 2). We calculated WCP for (1) generalist bird species; (2) species associated with early-to-

¹²⁹ mid succession forest (EMSF), which included the following land covers: wetland, deciduous, mixed-wood

¹³⁰ (young to mid-age), and coniferous habitat (young to mid-age); and (3) species associated with late succession

forest (LSF), which included the following land covers: mixed-wood (mature to old), deciduous (mature to old) and coniferous habitat (mature to old). Stand age was classified as follows: young (stand age < 30 years), mid-age (stand aged between 30 and 60 years), mature (stand aged between 60 and 80 years), and old (stand age greater than 80 years). This classification was used only for birds because information regarding beetles' associated land cover was unavailable at the moment of the study. Bird-associated land cover has been based on habitat associations from Gauthier and Aubry (1995) and Robert (2019).

2.6   Modelling framework

We developed species distribution models (SDMs) (Guisan and Zimmermann, 2000; Guisan and Thuiller, 2005; Peterson et al., 2011) to estimate single-species occurrence probability based on extensive field surveys of 104 species (77 beetle and 27 bird species). We partitioned the modeling strategy into several steps starting from data pre-processing to the species occurrence probability projections (see Bouderbala et al. 2022 for further details about the approach). SDMs were estimated only for species recorded at least at 1% and 5% of sampling sites for birds and beetles, respectively. We used land cover variables extracted from LANDIS forest landscape simulations for the simulated scenarios. We also standardized the predictor variables just before any model calibration to facilitate the convergence of the models (MacKenzie et al., 2017). We removed highly correlated variables ($|r| > 0.60$, where r represented pairwise Pearson correlation coefficient) and kept the 5 most important potential predictor variables according to the Akaike information criterion (AIC) calculated on a univariate generalized linear model (GLM) with linear and quadratic terms for each variable (Zurell et al., 2020; Bouderbala et al., 2022). We used generalized linear mixed models to predict species occurrence probability (package 'lme4'; Bates et al., 2015). A random intercept was added to include any differences between sampling years based on three full potential models deferring only in the fixed effects of the models. The best models were selected for each full model based on a 10-fold cross-validation procedure by minimizing the AIC criterion (package 'MuMIn'; Barton, 2015). For each species, we kept only the models with $AUC \geq 0.7$ (Araújo et al., 2005; Hosmer et al., 2013). In addition, we analyzed the assemblage structure based on continuous occurrence probabilities instead of binarizing the outputs (Guisan and Thuiller, 2005; Grenié et al., 2020).

We calculated the percentage change in the regional occupancy probability (ROP) between the reference (Ref) and the simulated (Sim) scenarios $\Delta\overline{ROP}_{Sim,Ref}$ (Bichet et al., 2016) as follows:

$$\Delta\overline{ROP}_{Sim,Ref} = \left(\frac{\overline{ROP}_{Sim} - \overline{ROP}_{Ref}}{\overline{ROP}_{Ref}} \right) \times 100 \ , \tag{1}$$

where $\overline{ROP}_s$ denotes the average of the ROP over all the species under the scenario s and is given by

$$\overline{ROP}_s = \frac{1}{N_{species}} \sum_{i=1}^{N_{species}} ROP_{s,i}$$

with

$$ROP_{s,i} = \frac{1}{N_{pixel}} \sum_{j=1}^{N_{pixel}} P_{s,i,j}$$

159 and where $N_{species}$, N_{pixel} and $P_{s,i,j}$ respectively represent the number of species, the number of pixels in

160 the study area, and the probability of occurrence of the species i for the scenario s at the cell j.

161 Winning conditional to habitat probability (WCP) was calculated as

$$\hat{P}\left(W_{Sim,Ref}|H\right) = \frac{\sum_{i=1}^{N_{species}} \mathbb{1}\left(\Delta ROP_{Sim,Ref,i} > 0, H_i \in H\right)}{\sum_{i'=1}^{N_{species}} \mathbb{1}\left(H_{i'} \in H\right)}, \qquad H \in \{G, EMSF, LSF\}, \qquad (2)$$

162 where $\mathbb{1}$ is the indicator function and H_i is the associated habitat for species i. In addition, we used

163 the regional Jaccard dissimilarity index $(RJ_{Sim,Ref})$ to measure the degree of dissimilarity in assemblage

164 composition between the compared scenarios (Baselga, 2010, 2013; Legendre, 2014; Bichet et al., 2016;

165 Belliard et al., 2018; Scherrer et al., 2020). The $RJ_{Sim,Ref}$ was defined as

$$RJ_{Sim,Ref} = \frac{2RBC_{Sim,Ref}}{1 + RBC_{Sim,Ref}}, \qquad (3)$$

where

$$RBC_{Sim,Ref} = \sum_{i=1}^{N_{species}} \frac{|ROP_{Sim,i} - ROP_{Ref,i}|}{\sum_{i'=1}^{N_{species}} (ROP_{Sim,i'} + ROP_{Ref,i'})}$$

is the regional Bray-Curtis dissimilarity between the scenarios Sim and Ref. We also used the Bray-Curtis dissimilarity index which we split into two additive components:

$$BC_{Sim,Ref} = BC\_grad_{Sim,Ref} + BC\_bal_{Sim,Ref},$$

where $BC\_grad_{Sim,Ref}$ is the occupancy gradient and $BC\_bal_{Sim,Ref}$ is the species balanced variation in occupancy. The latter was defined as

$$BC\_bal_{Sim,Ref} = \frac{\min(B_{Sim,Ref}, C_{Sim,Ref})}{A_{Sim,Ref} + \min(B_{Sim,Ref}, C_{Sim,Ref})},$$

where

$$B_{Sim,Ref} = \sum_{i=1}^{N_{species}} \left(ROP_{Sim,i} - \min(ROP_{Sim,i}, ROP_{Ref,i}) \right),$$

$$C_{Sim,Ref} = \sum_{i=1}^{N_{species}} \left(ROP_{Ref,i} - \min(ROP_{Sim,i}, ROP_{Ref,i}) \right),$$

$$A_{Sim,Ref} = \sum_{i=1}^{N_{species}} \min(ROP_{Sim,i}, ROP_{Ref,i})$$

To quantify the main source of assemblage dissimilarity, we performed a decomposition of the regional Bray-Curtis dissimilarity, based on the probability of occurrence, into *balanced variation in species occupancy* and *occupancy gradient*, which are the generalizations of species turnover and the nestedness respectively in the context of continuous output (Baselga, 2013) and occurrence probabilities in our case (Bouderbala et al., 2022). This decomposition of the dissimilarity measure computed based on scenarios helped us to characterize the mechanisms that drive variation in species composition. Hereafter the terms species turnover and nestedness will be used as equivalents of *balanced variation in species occupancy* and *occupancy gradient* respectively. We then computed the ratio

$$BC\_ratio_{Sim,Ref} = \frac{BC\_grad_{Sim,Ref}}{BC_{Sim,Ref}} \ . \tag{4}$$

The case $BC\_ratio_{Sim,Ref} > 0.5$ then indicated that the community change was mostly caused by nestedness, whereas $BC\_ratio_{Sim,Ref} < 0.5$ indicated the dominance of species turnover in the assemblage dissimilarity. We used the package 'betapart' (Baselga and Orme, 2012) for the assemblage analysis.

3   Results

Of 62 and 169 of the candidate species of birds and beetles, 27 and 77 species were selected for the projection with $AUC \geq 0.7$ (Fig. 2a-b) respectively. Stand age was the most frequently selected predictor variable for both taxa (Fig. 2c).

3.1   Changes in disturbance level following climate change

Changing climate from the baseline to RCP 8.5 scenarios in 2100 increased the proportion of burned stands at each harvest scenario. For example, the proportion of burned stands was 17.2% under baseline-Harvest

Figure 2: Performance metrics (area under the curve (AUC), sensitivity (Sens), specificity (Spec) and the true skill statistic (TSS)) of the selected models (**a, b**). Percentage of the predictor variables that were included in the species regressions for the two taxa (**c**). Abbreviations: distance to the nearest fire (dist_pert_Fire), stand age (Age), frequency of conifer dense (FreqConDens), frequency of conifer open (FreqConOpen), frequency of mixed wood (FreqMixed), frequency of open habitat (FreqOpen), frequency of disturbance by fire (FreqFire), frequency of disturbance by cut (FreqCut).

184 scenario, 24.7% under RCP 4.5-Harvest scenario, and 31.6% under RCP 8.5-Harvest scenario (Table. 3).

185 Furthermore, the increase either in wildfire or in harvest levels decreased the proportion of coniferous forests

186 (dense, open and mixed wood). Moreover, climate change as well as forest harvesting decreased significantly

187 stand age. For instance, under RCP 4.5 the mean stand age decreased from 90 to 46 years from NoHarvest

188 to Harvest scenarios. Also, the mean stand age under Harvest$_{0.5}$ went from 67 to 49 years from baseline to

RCP 8.5 (Table 3).

Table 3: Percentage of the area occupied by the different land covers in 2100 (stand age average in years). Abbreviations: conifer dense (ConDens), conifer open (ConOpen), mixed wood (Mixed), open habitat (Open), Other, disturbance by fire (Fire), disturbance by cut (Cut) and Fire+Cut (Disturbance).

Landscape	ConDens	ConOpen	Mixed	Open	Other	Fire	Cut	Total	Disturbance
Baseline-NoHarvest	10.83(156.91)	18.93(136.15)	30.97(142.48)	0.96(0)	23.18(31.66)	15.13(24.17)	0(–)	100(97.89)	15.13
Baseline-Harvest$_{0.5}$	6.91(149.04)	11.5(123.48)	22.7(120.13)	0.96(0)	23.87(28.84)	17.5(24.88)	16.56(22.36)	100(66.7)	34.06
Baseline-Harvest	4.83(146.46)	8.1(116.74)	17.83(109.52)	0.96(0)	23.87(27.87)	17.17(20.58)	27.25(23.25)	100(52.57)	44.42
RCP 4.5-NoHarvest	10.74(161.74)	17.87(142.92)	28.59(140.72)	0.96(0)	18.9(14.35)	22.94(17.35)	0(–)	100(89.83)	22.94
RCP 4.5-Harvest$_{0.5}$	6.66(154.74)	10.92(130.48)	21.43(117.39)	0.96(0)	20.06(15.83)	23.76(17.28)	16.21(23.64)	100(60.82)	39.97
RCP 4.5-Harvest	4.17(152.53)	6.74(120.14)	17.01(105.03)	0.96(0)	19.85(15.32)	24.65(19.05)	26.6(22.49)	100(46.06)	51.25
RCP 8.5-NoHarvest	6.67(160.38)	11.58(144.26)	24.68(144.53)	0.96(0)	20.36(21.31)	35.76(15.17)	0(–)	100(72.83)	35.76
RCP 8.5-Harvest$_{0.5}$	4.24(152.07)	7.12(133.56)	17.24(122.12)	0.96(0)	20.48(19.13)	36.32(14.77)	13.65(21.82)	100(49.27)	49.97
RCP 8.5-Harvest	2.94(151.72)	5.3(126.17)	14.65(113.17)	0.96(0)	21.14(19.21)	31.63(15.31)	23.38(21.48)	100(41.66)	55.01

3.2   Assemblage analysis

Species assemblage dissimilarity was higher in NoHarvest-Harvest than in NoHarvest-Harvest$_{0.5}$, under the three climate scenarios in 2100 (Fig. 3). Under RCP 4.5, Jaccard dissimilarity increased from 0.16 to 0.22 for birds and 0.13 to 0.20 for beetles when comparing NoHarvest-Harvest$_{0.5}$ to NoHarvest-Harvest scenarios. Compared to beetles, birds had slightly higher dissimilarity in assemblages regardless of climate scenarios (Fig. 3). However, the dissimilarity in species assemblage over harvest gradient decreased with climate change (see the difference in Jaccard dissimilarity between baseline and RCP 8.5 in Fig. 3).

Figure 3: Regional Jaccard dissimilarity measures (Eq. 3) computed based on the difference in species regional occupancy between reference and simulated scenario. At each climate change scenario (Baseline, RCP 4.5, RCP 8.5), we computed the dissimilarity between Harvest$_{0.5}$ and Harvest vs. NoHarvest in 2100. Under Baseline we compared respectively Baseline-NoHarvest to Baseline-Harvest$_{0.5}$ and Baseline-NoHarvest to Baseline-Harvest. Under RCP 4.5 we compared respectively RCP 4.5-NoHarvest to RCP 4.5-Harvest$_{0.5}$ and RCP 4.5-NoHarvest to RCP 4.5-Harvest. Finally, under RCP 8.5 we compared respectively RCP 8.5-NoHarvest to RCP 8.5-Harvest$_{0.5}$ and RCP 8.5-NoHarvest to RCP 8.5-Harvest.

3.3   Occupancy analysis

We observed a change in the regional occupancy between the scenarios of forest harvesting (Harvest$_{0.5}$ and Harvest vs. NoHarvest) under all three climate change scenarios. The increase in the overall occupancy of birds, and the decrease for beetles from NoHarvest to Harvest cases was visible over 114,118 km^2 of the study area (Fig. 4) and also regionally (Fig. 5a). Post-harvesting changes in species assemblages were mainly caused by species turnover ($BC\_ratio_{Sim,Ref} < 0.5$) for both taxa (Fig. 5a). This dominance of species turnover was associated with an increase in regional occupancy of birds contrary to beetles in which species turnover was associated with a decrease in the regional occupancy. Under each climate scenario, the occupancy of species associated with young habitats together with generalist species increased significantly contrary to species associated with mature and old forests, when we compared harvested to unharvested

Figure 4: The average and standard deviation maps of the potential occurrence distribution for each taxon based on the four scenarios **BaselineNoHarvest**, **BaselineHarvest**, **RCP 8.5NoHarvest** and **RCP 8.5Harvest**. The black line represents the northern limit for forest harvesting activities.

scenarios (Fig. 5b). Under baseline and RCP 4.5, only 40% of species associated with late succession forest increased in occupancy from no harvested to harvested scenarios (Fig. 6), whereas 89% of species associated with early to mid succession forest increased in occupancy (see Fig. S1 for the variation at the species scale). By comparing RCP 8.5 to RCP 4.5, the proportion of winner species (from non harvested to harvested landscapes) that were associated with early to mid succession forest decreased from 89% to 83%. Under the same conditions, species associated with late succession forest increased from 40% to 60% (Fig. 6).

Figure 5: **(a)** Regional Bray-Curtis dissimilarity ratio (BC_ratio) with the percentage of change in the regional occupancy probability ($\Delta\overline{ROP}$). At each climate change scenario (Baseline, RCP 4.5, RCP 8.5), we computed the change in the ROP (Eq. 1) and Bray-Curtis ratio (Eq. 4) between Harvest$_{0.5}$ and Harvest vs. NoHarvest in 2100. **(b)** Percentage of change in the regional occupancy probability between Harvest and Noharvest under the baseline and RCP 8.5 for bird species. **Abbreviations**: late succession forest (LSF), early-to-mid succession forest ($EMSF$), generalist species (G).

4   Discussion

Forest harvesting can impact significantly forest biota and may have lasting effects on biodiversity (Lindenmayer and Franklin, 2002). We aimed to quantify how the interaction between forest harvesting and climate change would impact biodiversity. We analyzed how species assemblages can be expected to change under different forest management scenarios and climate conditions. Our models suggest that forest harvesting will cause pronounced changes in species assemblages for birds and beetles by 2100 under all climate change scenarios. Our result is supported by many empirical studies (Baker et al., 2016; Bichet et al., 2016) evoking the noticeable effect of harvesting on the assemblage of common species and predicted that the effect could be stronger by including rare species (Bichet et al., 2016). Moreover, Tremblay et al. (2018) projects a drastic reduction in the habitat of the Black-backed Woodpecker (*Picoides arcticus*), a focal species representative of deadwood and old-growth biodiversity in eastern Canada (Tremblay et al., 2020), due to cumulative impacts of forest harvesting and climate change. The simulated change in assemblage dissimilarity was mainly dominated by species turnover. This implies that the change in the assemblage composition from NoHarvest to Harvest scenarios was dominated by an increase in occupancy for a subset of species that was balanced mostly by a decrease in occupancy for the other subset of species.

The comparison between climate scenarios must be taken with caution, however. We quantified the extent of change in species occurrence by modifying forest harvest magnitude under each climate situation. So, the difference between the less disturbed scenario (baseline-NoHarvest, i.e. 15% of disturbance) and

Figure 6: Conditional winning probability (Eq. 2) of bird species among land covers when the landscape changes from NoHarvest to "Harvest", under the three climates scenarios. Under Baseline we compared Baseline-NoHarvest to Baseline-Harvest. Under RCP 4.5 we compared RCP 4.5-NoHarvest to RCP 4.5-Harvest. Finally, under RCP 8.5 we compared RCP 8.5-NoHarvest to RCP 8.5-Harvest.

the most disturbed scenario (RCP8.5-Harvest, i.e. 55% of disturbance) on species occurrence could be different from our results based on comparing different harvest levels within the same climate scenario. Our results predict an increase in occupancy in the majority of bird species associated with Early-to-Mid succession forest contrary to species associated with Late succession forest, by comparing harvested to unharvested scenarios, under baseline and medium climate scenarios. This result is coherent with previous studies (Fortin et al., 2011; Bichet et al., 2016). However, under the extreme climate scenario, there is one more late-succession species increasing in occupancy when we compared harvested to unharvested scenarios, while this is the opposite for early-to-mid succession species which decreased by one species. This difference could be explained by the proportion of burned stands on the harvestable stands under each climate scenario. Indeed, we observed a decrease in the harvested stands with climate change contrary to the burned stands. Those interactions between forest harvesting and future climate change could impact the occupancy of species that depend on disturbed stands. For example, on the one hand, we observed that *Yellow Warbler* (YEWA, *Setophaga petechia*), a bird species associated with early-to-mid succession forest and negatively affected by both harvested forest and stand age regressed under RCP 8.5 contrary to the baseline climate scenario. On the other hand, *Swainson's Thrush* (SWTH, *Catharus ustulatus*) that was associated with late succession forest and negatively affected with burned forests increased under RCP 8.5 (see Fig. S1). The increase in wildfires under the extreme climate scenario could affect negatively species preferring harvested stands.

Our results were based only on the most common species, the response of rare specialist and threatened species might decrease the regional occupancy with both harvest activities and climate conditions (Cadieux et al., 2019). Therefore, it will be interesting to analyze the response of the rare species to disturbance. One possible approach could be through the development of a probabilistic model which combines the occurrence

probability of rare species with indicator species.

In conclusion, forest harvesting activities are predicted to modify the composition of beetle and bird
species assemblages under all climate scenarios. We also indicated that the consequences of maintaining the
current level of forest harvesting under the worst climate change scenario could modify the species-habitat
association. Ideally, we aim to generate a situation where both early and late successional stages can coexist
by managing forests. Interestingly, we observed opposite responses in occupancy between birds, which was
generally positive, and beetle, which was negative, in our study area. The perfect separation between the
two taxonomic groups regarding the change in occupancy could reflect the impact of ecological traits such
as movement capacity and diet differences between boreal species. In any case, forest management must be
adapted to achieve conservation objectives under all climate scenarios. Our simulations show that harvest
activities must be reduced. Even when it maximizes biodiversity, harvest could be harmful to taxa with low
moving capacity (beetles in our case).

264 Acknowledgements

This work was supported by the Sentinel North program of Laval University, funded by the Canada First
Research Excellence Fund. A.A., D.F., C.H., and P.D. were also supported by The Natural Sciences and
Engineering Research Council of Canada (NSERC). We acknowledge Calcul Québec and Compute Canada
for their technical support and computing infrastructures. We also thank the Québec Breeding Bird Atlas
for supplying data. We would also like to thank the following partners: Regroupement QuébecOiseaux,
Environment and Climate Change Canada, and Birds Canada, as well as all of the volunteer participants who
gathered data for the project. We are grateful to Louis-Paul Rivest for his valuable statistical suggestions.
Special thanks are due to Alexandre Terrigeol for his valuable efforts regarding data collection, as well as to
Georges Pelletier and Nicolas Bédard for beetle identification.

274 References

Almeida-Rocha, J. M., Soares, L. A., Andrade, E. R., Gaiotto, F. A., and Cazetta, E. (2020). The impact of
anthropogenic disturbances on the genetic diversity of terrestrial species: A global meta-analysis. *Molecular*
Ecology, 29(24):4812–4822.

Araújo, M. B., Pearson, R. G., Thuiller, W., and Erhard, M. (2005). Validation of species–climate impact
models under climate change. *Global Change Biology*, 11(9):1504–1513.

Atlas des oiseaux nicheurs du Québec (2018). Données obtenues en réponse à une demande présentée aux

bureaux de l'Atlas (www.atlas-oiseaux.qc.ca). Regroupement QuébecOiseaux; Service canadien de la faune

d'Environnement Canada; Études d'Oiseaux Canada. Québec, QC, Canada.

Baker, S. C., Halpern, C. B., Wardlaw, T. J., Kern, C., Edgar, G. J., Thomson, R. J., Bigley, R. E., Franklin,

J. F., Gandhi, K. J. K., Gustafsson, L., Johnson, S., Palik, B. J., Spies, T. A., Steel, E. A., Weslien, J.,

and Strengbom, J. (2016). A cross-continental comparison of plant and beetle responses to retention of

forest patches during timber harvest. *Ecological Applications*, 26(8):2495–2506.

Barton, K. (2015). Package 'MuMIn'. https://cran.r-project.org/web/packages/MuMIn.

Baselga, A. (2010). Partitioning the turnover and nestedness components of beta diversity. *Global Ecology*

and Biogeography, 19(1):134–143.

Baselga, A. (2013). Separating the two components of abundance-based dissimilarity: balanced changes in

abundance vs. abundance gradients. *Methods in Ecology and Evolution*, 4(6):552–557.

Baselga, A. and Orme, C. D. L. (2012). betapart: an R package for the study of beta diversity. *Methods in*

Ecology and Evolution, 3(5):808–812.

Bates, D., Mächler, M., Bolker, B., and Walker, S. (2015). Fitting linear mixed-effects models using lme4.

https://cran.r-project.org/web/packages/lme4.

Beaubien, J., Cihlar, J., Simard, G., and Latifovic, R. (1999). Land cover from multiple thematic mapper

scenes using a new enhancement-classification methodology. *Journal of Geophysical Research: Atmo-*

spheres, 104(D22):27909–27920.

Beaudoin, A., Bernier, P., Guindon, L., Villemaire, P., Guo, X., Stinson, G., Bergeron, T., Magnussen, S.,

and Hall, R. (2014). Mapping attributes of Canada's forests at moderate resolution through kNN and

MODIS imagery. *Canadian Journal of Forest Research*, 44(5):521–532.

Belliard, J., Beslagic, S., Delaigue, O., and Tales, E. (2018). Reconstructing long-term trajectories of

fish assemblages using historical data: the Seine River Basin (France) during the last two centuries.

Environmental Science and Pollution Research, 25(24):23430–23450.

Bibby, C. J., Burgess, N. D., Hill, D. A., Hillis, D. M., and Mustoe, S. (2000). *Bird census techniques.*

Elsevier.

307 Bichet, O., Dupuch, A., Hébert, C., Le Borgne, H., and Fortin, D. (2016). Maintaining animal assemblages
308 through single-species management: the case of threatened caribou in boreal forest. *Ecological Applications*,
309 26(2):612–623.

310 Bouchard, M. and Pothier, D. (2011). Long-term influence of fire and harvesting on boreal forest age structure
311 and forest composition in eastern Québec. *Forest Ecology and Management*, 261(4):811–820.

312 Boucher, Y., Perrault-Hébert, M., Fournier, R., Drapeau, P., and Auger, I. (2017). Cumulative patterns
313 of logging and fire (1940–2009): consequences on the structure of the eastern Canadian boreal forest.
314 *Landscape Ecology*, 32(2):361–375.

315 Bouderbala, I., Labadie, G., Béland, J.-M., Boulanger, Y., Hébert, C., Desrosiers, P., Allard, A., and Fortin,
316 D. (2022). Effects of global change on animal biodiversity in boreal forest landscape: an assemblage
317 dissimilarity analysis. *bioRxiv*.

318 Cadieux, P., Boulanger, Y., Cyr, D., Taylor, A. R., Price, D. T., and Tremblay, J. A. (2019). Spatially
319 explicit climate change projections for the recovery planning of threatened species: The Bicknell's Thrush
320 (Catharus Bicknelli) as a case study. *Global Ecology and Conservation*, 17:e00530.

321 Ceaușu, S., Apaza-Quevedo, A., Schmid, M., Martín-López, B., Cortés-Avizanda, A., Maes, J., Brotons, L.,
322 Queiroz, C., and Pereira, H. M. (2021). Ecosystem service mapping needs to capture more effectively the
323 biodiversity important for service supply. *Ecosystem Services*, 48:101259.

324 Coelho, A. J. P., Magnago, L. F. S., Matos, F. A. R., Mota, N. M., Diniz, É. S., and Meira-Neto, J. A. A.
325 (2020). Effects of anthropogenic disturbances on biodiversity and biomass stock of Cerrado, the Brazilian
326 savanna. *Biodiversity and Conservation*, 29(11):3151–3168.

327 Cyr, D., Gauthier, S., Bergeron, Y., and Carcaillet, C. (2009). Forest management is driving the eastern north
328 american boreal forest outside its natural range of variability. *Frontiers in Ecology and the Environment*,
329 7(10):519–524.

330 Davis, M. B. and Shaw, R. G. (2001). Range shifts and adaptive responses to quaternary climate change.
331 *Science*, 292(5517):673–679.

332 Dobson, A., Rowe, Z., Berger, J., Wholey, P., and Caro, T. (2021). Biodiversity loss due to more than
333 climate change. *Science*, 374(6568):699–700.

334 Fortin, D., Hébert, C., Légaré, J.-P., Courbin, N., Swiston, K., Hodson, J., LeBlanc, M.-L., Dussault, C.,
335 Pothier, D., Ruel, J., et al. (2011). Partial harvesting in old-growth boreal forests and the preservation of

animal diversity from ants to woodland caribou. *E.B. Wallace (Ed.), Woodlands: Ecology, Management*
and Conservation, Nova Science Publishers, Inc, Hauppauge, NY, USA.

Gauthier, J. and Aubry, Y. (1995). *Les oiseaux nicheurs du Québec: Atlas des oiseaux nicheurs du Québec*
méridional. Association québécoise des groupes d'ornithologues; Société québécoise de protection des
oiseaux; Service canadien de la faune; Environnement Canada, région du Québec.

Grenié, M., Violle, C., and Munoz, F. (2020). Is prediction of species richness from stacked species distribution
models biased by habitat saturation? *Ecological Indicators*, 111:105970.

Guisan, A. and Thuiller, W. (2005). Predicting species distribution: offering more than simple habitat
models. *Ecology Letters*, 8(9):993–1009.

Guisan, A. and Zimmermann, N. E. (2000). Predictive habitat distribution models in ecology. *Ecological*
Modelling, 135(2):147–186.

Hosmer, D. W., Lemeshow, S., and Sturdivant, R. X. (2013). *Applied logistic regression, third edition.* John
Wiley & Sons.

Janssen, P., Fortin, D., and Hébert, C. (2009). Beetle diversity in a matrix of old-growth boreal forest:
influence of habitat heterogeneity at multiple scales. *Ecography*, 32(3):423–432.

Labadie, G., Hardy, C., Boulanger, Y., Vanlandeghem, V., Hebblewhite, M., and Fortin, D. (2022). Global
change impacts a threatened species through alteration of predator-prey dynamics. Research Square.

Labadie, G., McLoughlin, P. D., Hebblewhite, M., and Fortin, D. (2021). Insect-mediated apparent compe-
tition between mammals in a boreal food web. *Proceedings of the National Academy of Sciences of the*
United States of America, 118(30).

Légaré, J.-P., Hébert, C., and Ruel, J.-C. (2011). Alternative silvicultural practices in irregular boreal forests:
Response of beetle assemblages. *Silva Fennica*, 45(5):937–956.

Legendre, P. (2014). Interpreting the replacement and richness difference components of beta diversity.
Global Ecology and Biogeography, 23(11):1324–1334.

Lindenmayer, D. B. and Franklin, J. F. (2002). *Conserving forest biodiversity: a comprehensive multiscaled*
approach. Island Press.

MacKenzie, D. I., Nichols, J. D., Royle, J. A., Pollock, K. H., Bailey, L., and Hines, J. E. (2017). *Occupancy*
estimation and modeling: inferring patterns and dynamics of species occurrence. Elsevier.

Matuoka, M. A., Benchimol, M., de Almeida-Rocha, J. M., and Morante-Filho, J. C. (2020). Effects of
anthropogenic disturbances on bird functional diversity: A global meta-analysis. *Ecological Indicators*,
116:106471.

Micheletti, T., Stewart, F. E. C., Cumming, S. G., Haché, S., Stralberg, D., Tremblay, J. A., Barros, C.,
Eddy, I. M. S., Chubaty, A. M., Leblond, M., Pankratz, R. F., Mahon, C. L., Van Wilgenburg, S. L.,
Bayne, E. M., Schmiegelow, F., and McIntire, E. J. B. (2021). Assessing pathways of climate change
effects in SpaDES: An application to bBoreal Landbirds of Northwest Territories Canada. *Frontiers in*
Ecology and Evolution, 9:654.

Ministère des Forêts de la Faune et des Parcs (2018). Aires infestées par la tordeuse des bourgeons de
l'épinette au québec en 2018, 20 p. Technical report, Québec, gouvernement du Québec, Direction de la
protection des forêts.

Muluneh, M. G. (2021). Impact of climate change on biodiversity and food security: a global perspective—a
review article. *Agriculture & Food Security*, 10(1):1–25.

Naeem, S., Duffy, J. E., and Zavaleta, E. (2012). The functions of biological diversity in an age of extinction.
Science, 336(6087):1401–1406.

Newbold, T., Hudson, L. N., Hill, S. L., Contu, S., Lysenko, I., Senior, R. A., Börger, L., Bennett, D. J.,
Choimes, A., Collen, B., et al. (2015). Global effects of land use on local terrestrial biodiversity. *Nature*,
520(7545):45–50.

Peterson, A. T., Soberón, J., Pearson, R. G., Anderson, R. P., Martínez-Meyer, E., Nakamura, M., and
Araújo, M. B. (2011). *Ecological niches and geographic distributions (MPB-49)*, volume 49. Princeton
University Press.

Robert, M. (2019). *Deuxième atlas des oiseaux nicheurs du Québec méridional*. Regroupement Québec-
Oiseaux; Environnement et changement climatique Canada; Études d'oiseaux Canada.

Sala, O. E., Chapin, F. S., null III, Armesto, J. J., Berlow, E., Bloomfield, J., Dirzo, R., Huber-Sanwald,
E., Huenneke, L. F., Jackson, R. B., Kinzig, A., Leemans, R., Lodge, D. M., Mooney, H. A., Oesterheld,
M., Poff, N. L., Sykes, M. T., Walker, B. H., Walker, M., and Wall, D. H. (2000). Global biodiversity
scenarios for the year 2100. *Science*, 287(5459):1770–1774.

Scheller, R. M. and Domingo, J. B. (2005). LANDIS-II Base Fire Extension (v1. 0) User Guide.

Scheller, R. M., Domingo, J. B., Sturtevant, B. R., Williams, J. S., Rudy, A., Gustafson, E. J., and Mladenoff,
D. J. (2007). Design, development, and application of LANDIS-II, a spatial landscape simulation model
with flexible temporal and spatial resolution. *Ecological Modelling*, 201(3):409–419.

Scheller, R. M. and Mladenoff, D. J. (2004). A forest growth and biomass module for a landscape simulation
model, LANDIS: design, validation, and application. *Ecological Modelling*, 180(1):211–229. Modelling
disturbance and succession in forest landscapes using LANDIS.

Scherrer, D., Mod, H. K., and Guisan, A. (2020). How to evaluate community predictions without thresh-
olding? *Methods in Ecology and Evolution*, 11(1):51–63.

Sergio, F., Blas, J., and Hiraldo, F. (2018). Animal responses to natural disturbance and climate extremes:
a review. *Global and Planetary Change*, 161:28–40.

Sturtevant, B. R., Gustafson, E. J., Li, W., and He, H. S. (2004). Modeling biological disturbances in LAN-
DIS: a module description and demonstration using spruce budworm. *Ecological Modelling*, 180(1):153–174.

Sturtevant, B. R., Miranda, B. R., Shinneman, D. J., Gustafson, E. J., and Wolter, P. T. (2012). Comparing
modern and presettlement forest dynamics of a subboreal wilderness: Does spruce budworm enhance fire
risk? *Ecological Applications*, 22(4):1278–1296.

Thomas, C. D., Cameron, A., Green, R. E., Bakkenes, M., Beaumont, L. J., Collingham, Y. C., Erasmus,
B. F., De Siqueira, M. F., Grainger, A., Hannah, L., et al. (2004). Extinction risk from climate change.
Nature, 427(6970):145–148.

Tremblay, J. A., Boulanger, Y., Cyr, D., Taylor, A. R., Price, D. T., and St-Laurent, M.-H. (2018). Harvesting
interacts with climate change to affect future habitat quality of a focal species in eastern Canada's boreal
forest. *PLoS ONE*, 13(2):1–25.

Tremblay, J. A., Dixon, R. D., Saab, V. A., Pyle, P., and Patten, M. A. (2020). Black-backed Woodpecker
(Picoides arcticus), version 1.0. In Birds of the World (P. G. Rodewald, Editor). Cornell Lab of Ornithology,
Ithaca, NY, USA.

Zhao, Q., Azeria, E. T., Le Blanc, M.-L., Lemaître, J., and Fortin, D. (2013). Landscape-scale disturbances
modified bird community dynamics in successional forest environment. *PLoS ONE*, 8(11):1–11.

Zurell, D., Zimmermann, N. E., Gross, H., Baltensweiler, A., Sattler, T., and Wüest, R. O. (2020). Testing
species assemblage predictions from stacked and joint species distribution models. *Journal of Biogeography*,
47(1):101–113.

(a) Change in occupancy VS BC_ratio

(b) Change in occupancy by habitat