The Ecosystem Integrity Index: a novel measure of terrestrial ecosystem integrity

Authors: Hill, S.L.L.*, Harrison, M.L.K, Maney, C., Fajardo, J., Harris, M., Ash, N., Bedford, J., Danks, F.S., Guaras, D., Hughes, J., Jones, M., Mason, T., Burgess, N. [1]

Address: All authors: United Nations Environment Programme World Conservation Monitoring Centre (UNEP-WCMC), 219 Huntington Road, Cambridge, UK. [1]Also at the Centre for Macroecology, Evolution and Climate, GLOBE Institute, University of Copenhagen, Denmark

Abstract

While the importance of ecosystem integrity has long been recognised (Leopold, 1949), conservation science has tended to focus on measuring and monitoring species and habitats, avoiding the complexities of working at the ecosystem level. Ecosystems are highly dynamic, defined by both living and non-living components as well as their interactions (CBD, 1992), making it difficult to assess baseline levels of integrity. We present a novel index that represents the integrity of all terrestrial ecosystems globally at 1km^2 resolution: the Ecosystem Integrity Index (EII). The index provides a simple, yet scientifically robust, way of measuring, monitoring and reporting on ecosystem integrity. It is formed of three components; structural, compositional and functional integrity, and measured against a natural (current potential) baseline on a scale of 0 to 1. We find that ecosystem integrity is severely impacted in terrestrial areas across the globe with approximately one fifth of all ecosystems and one quarter of all ecoregions having lost, on average, over half of their ecosystem integrity. At a national scale, we estimate similar challenges with 115 nations or territories having lost, on average, over half of their ecosystem integrity. This presents a significant threat for humanity as such levels of degradation are likely to be linked to substantial declines in the ecosystem services on which humanity is reliant. The EII has been developed principally to help national governments measure and report on Goal A of the Kunming-Montreal Global Biodiversity Framework (GBF) (CBD, 2022a), for which it has been listed as a Component Indicator. The EII will also be useful in helping non-state actors measure and report their contributions to the GBF and is listed as an indicator by both the Taskforce for Nature-Related Financial Disclosures (TNFD) (TNFD, 2023) and the Science Based Targets Network (SBTN) (SBTN, 2023). The EII aims to enable these actors to make informed decisions on the conservation, restoration and sustainable use of ecosystems for which they are wholly or partly responsible. We propose that with sufficient effort, ecosystem integrity can be restored and contribute towards the GBF's vision of living in harmony with nature, resulting in the safeguarding of the ecosystem services on which humanity depends.

Introduction

Ecosystem integrity encompasses the full complexity of an ecosystem, including the physical, biological and functional components, together with their interactions, and measures these against a 'natural' (i.e., current potential) reference level (Carter *et al.,* 2019). Ecosystem integrity is fundamental to the stability of Earth systems on which humanity depends. For instance, natural areas containing ecosystems with higher integrity have greater potential to provide services such as carbon sequestration (Lewis *et al.,* 2009), maintenance of water quality (Mello *et al.,* 2018), climate

regulation (Bonan, 2008), pest control (Bianchi *et al.,* 2006), and pollination (Carvalheiro *et al.,* 2010)
– as well as supporting higher levels of biodiversity (Gibson, 2011; Barlow *et al.,* 2016).

The importance of healthy ecosystems (ecosystems with high integrity) is recognised in the three Rio
Conventions: the Convention on Biological Diversity (CBD), the UN Framework Convention on
Climate Change and the UN Convention to Combat Desertification.

The Kunming-Montreal Global Biodiversity Framework (GBF), negotiated under the CBD, includes
ambitions for the maintenance, restoration or enhancement of ecosystem integrity within Goal A
(CBD, 2022a). This index has been designed to help parties to the Convention to directly monitor
ecosystem integrity and is listed as a Component Indicator within the adopted Monitoring
Framework for the GBF (CBD, 2022b).

Meeting the goals and targets of the GBF will require action not only at the national level (by the 196
parties to the CBD), but also by a range of other actors including businesses. The Ecosystem Integrity
Index (EII) provides a means of alignment between the goals and targets of the GBF and also; 1)
target guidance under development by the Science Based Targets Network (SBTN); 2) the risk
management and disclosure framework of the Taskforce on Nature-related Financial Disclosures
(TNFD). Both TNFD and SBTN have identified the need for a scalable metric of ecosystem integrity,
linked to the GBF, which is readily usable by businesses and financial institutions for evidence-based
target setting, monitoring and disclosure (TNFD, 2022; SBTN, 2020).

Conceptualisation of ecosystem integrity

The notion of ecosystem integrity first emerged in the work of Leopold (1949), who used the term to
encompass the basic criteria necessary for biotic communities to remain stable. The concept was
elaborated upon by Parrish *et al*. (2003), who proposed that 'an ecological system has integrity when
its dominant ecological characteristics (e.g., elements of structure, composition, function, and
ecological processes) occur within their natural ranges of variation'. Carter *et al.* (2019), simplified
this to define ecosystem integrity as 'the extent to which the composition, structure, and function of
an ecosystem fall within their natural range of variation'. Structural integrity comprises the three-
dimensional aspect of ecosystems – the biotic and abiotic elements that form the heterogeneous
matrix supporting composition and functioning. Composition refers to the biotic constitution of
ecosystems – the pattern of the makeup of species communities. Functioning describes the
ecological processes and ecosystem services provided by the ecosystem. It follows that any
measurement of ecosystem integrity should encompass structural, compositional, and functional
components. However, it should be noted that the three components are interdependent and are
likely to covary with varying pressures on the system.

Ecosystem integrity has generated considerable interest in recent years with the development of
several indices designed to assess changes in ecological factors relating to integrity at a global level
or within specific habitat types. Recent work has focussed on measuring ecosystem extent (there
remains little scientific consensus on the best way to do this (Sayre *et al.*, 2020; IUCN-CEM, 2022)),
and on aspects of ecosystem integrity (Beyer *et al.*, 2019; Blumetto *et al.*, 2019; Grantham *et al.,*
2020; Hansen *et al.*, 2021; Mora, 2017; Perkl, 2017). For instance, Grantham et al. (2020) created the
Forest Landscape Integrity Index (FLII) integrating data on forest extent, observed and inferred
human pressures, and changes in forest connectivity, and an index of ecosystem integrity – focussing
on agricultural systems – was developed by Blumetto *et al.* (2019) to monitor integrity at the farm
level. (For a longer discussion of relevant work in this area, see SI – methodology).

Despite these various efforts, there is currently no index that matches the definition of ecosystem
integrity, in that it brings together all three components of ecosystem integrity, being structure,
composition and functioning, measured against a natural baseline, that can be applied globally.

Introducing the Ecosystem Integrity Index

Our Ecosystem Integrity Index (EII) quantifies degradation to ecosystem integrity based on an
aggregation of all three components: ecosystem structure, composition, and functioning. This
framing is consistent with the definitions provided by Carter *et al.* (2019) and Parrish *et al.* (2003).
Our index provides a measure of a fundamental property of ecosystems, ecosystem integrity, at a
fine scale, $1km^2$, but can be aggregated to other levels, increasing its utility to provide support to
monitoring and spatial planning at local to global scales. The ability to aggregate EII at different
scales allows it to be applied in multiple use-cases. For example, it is compatible with all levels of the
IUCN Global Ecosystem Typology (Keith *et al.,* 2020), which forms the basis for the biomes used
within the TNFD Recommended Disclosures (TNFD, 2023), or, as we demonstrate in this analysis, it
can be used with the World Ecosystems typology (Sayre *et al., 2020)* or ecoregions (Dinerstein *et al.,*
2017), but it is also compatible with national-level ecosystem typologies and so could support
governments engaged in spatial planning.

The first component of EII – structure – is correlated with habitat intactness, the area of contiguous
habitat, and fragmentation. This component is calculated by inputting the Human Modification Index
(HMI) (Kennedy *et al.,* 2019) into the algorithm described in Beyer *et al.* (2019) that accounts for
habitat intactness within a grid cell as well as the influence of habitat quality within neighboring
cells. This method ensures that our final structure layer encompasses impacts of landscape
fragmentation as well as local intactness.

The second component captures ecosystem composition, which refers to the identity and variety of
life within an area. The metric chosen for this layer is the Biodiversity Intactness Index (BII), which
summarizes change in the make-up of local ecological communities in response to human pressures
(Newbold *et al.,* 2016, Hill *et al.,* 2019). The BII is calculated using two models fitted to species
community data recorded in the PREDICTS database (Hudson *et al*., 2017). The first model estimates
the impact of human pressures on the total abundance of species within a community and the
second analyses the similarity between the relative abundance of each of the species in degraded
areas against those of the same communities found within natural areas. The product of the two
models, projected onto maps of human pressures, results in the BII.

The third component captures ecosystem function. The functioning component is estimated using
the difference between potential natural and current net primary productivity (NPP) within each
$1km^2$ grid cell. Current NPP values are taken from remote sensed geospatial layers (Running and
Zhao, 2019). Potential natural NPP values are modelled using environmental input data (for model
details see Methods).

The three component layers are aggregated to form the EII. We used a modified minimum value
approach, in which we take the lowest score of the structure, composition and functioning
components as the EII value for a pixel and then down-weight the value based upon the integrity of
the other two components. This method was chosen with the reasoning that the integrity of an
ecosystem cannot be higher than the minimum score from any of the three components. The
derived EII, as well as the three components, ranges between 0 and 1, with the most degraded areas
having a score approaching zero.

Results

Ecosystem integrity is highly degraded across much of the world with only remote areas in higher
latitudes (note that Antarctica was excluded from the analysis), remote areas of rainforest across the
tropics, and extremely arid areas across the planet estimated to have extensive areas where
ecosystem integrity has been retained (Figure 1A).

Mean ecosystem integrity across all ecosystems (ecosystem boundaries as defined by Sayre *et al.,*
2020) was 0.73 (±0.22), with 74 of 422 ecosystems having, on average, lost at least half of their
ecosystem integrity. Ecosystems within human settlement areas in warm temperate regions were
found to have the lowest levels of ecosystem integrity, followed by ecosystems found within
cropland areas (Figure 1B). At broader scales, approximately one quarter (198 of 824) of ecoregions
have, on average, lost at least half of their ecosystem integrity (mean EII <0.5). Dry tropical broadleaf
forests are more degraded than moist, and temperate grasslands, temperate broadleaf forest and
Mediterranean forests, woodlands and scrub all have high levels of degradation (Figure 1C).

Figure 1. Global ecosystem integrity. A) The Ecosystem Integrity Index for all terrestrial areas. Areas with high levels of ecosystem integrity are shown in dark blue, whereas those with highly degraded ecosystem integrity are shown in pale yellow. Note: the boundaries and names shown and the designations used on this map do not imply official endorsement or acceptance by the United Nations. B) Mean ecosystem integrity per ecosystem (Sayre *et al.,* 2020) grouped by landcover types.

Colours indicate climate regions. C) Frequency of ecosystem integrity index values within terrestrial
biomes (WWF 2017).

Ecosystem degradation is widespread at the national scale (Figure 2). Across all 255 countries and
territories assessed, the mean EII value was 0.55 (SD:0.23). When comparing between continents,
North American countries and territories were found to have, on average, the lowest EII scores with
a mean value of 0.46 (SD:0.22), and those in West Asia had the highest EII scores with a mean value
of 0.68 (SD:0.17).

Figure 2. Ecosystem integrity within countries and territories grouped by UN Regions. Each dot
represents the values for a single country. Box plots indicate mean ±SE.

Discussion

The EII provides a unique insight into the extent that terrestrial ecosystems have been degraded
across the world. We have used ecological theory and the best available data to create an index at
operational scales (1km^2) globally. Our results show that ecosystem integrity has been highly
degraded across the terrestrial world with nearly one fifth of all ecosystems, and one quarter of all
ecoregions, having lost, on average, over half of their ecosystem integrity. Such dramatic losses of
ecosystem integrity are likely to have had severe consequences for species persistence and provision
of ecosystem services, but further work is needed to understand how the spatial distribution of
ecosystem integrity interacts with these factors. Such knowledge will be required to enable effective
spatial planning, land management options and restoration interventions to help achieve the goals
and targets of the Kunming-Montreal GBF.

We developed this index for two main use cases. The first is for use at the national and sub-national
scales by countries seeking ways to monitor progress against their national goals and targets within
the context of the Kunming-Montreal GBF. In particular, the index will facilitate tracking of progress
towards national contributions to Goal A, maintenance, enhancement and restoration of ecosystem
integrity, as well as targets 1, 2 and 10 covering spatial planning, restoration and sustainable use.
The global EII layer will be made available through assessment tools such as the UN Biodiversity Lab
(https://unbiodiversitylab.org/) to support countries in planning for the delivery of these targets and
monitoring progress in terms of ecological outcomes.

Our second use case is for businesses and financial institutions which are starting to include
biodiversity measurement in their risk assessment and performance reporting. There is an urgent
need for trackable targets using an appropriate set of metrics to enable the private sector to move
towards positive outcomes for biodiversity (Locke *et al.,* 2022; Milner-Gulland, 2022). Such
outcomes are reliant upon the ability of a business to measure their biodiversity baseline, assess
their exposure to biodiversity risk within operations, and to drive innovation towards positive
outcomes on the ground through a quantified understanding of the pressure-response relationship.
Initiatives such as the SBTN and the TNFD are working with businesses and financial institutions,
helping them contribute to nature-positive outcomes. They are developing guidance for businesses
to set science-based targets, and through recommendations for disclosures when reporting on
impacts and dependencies on nature. In both these initiatives, participating businesses and financial
institutions will need to understand where they interact with ecosystems and identify priority
locations, for example, in the 'Locate' step in the TNFD LEAP (Locate, Evaluate, Assess, Prepare)
approach (TNFD, 2023) and within the 'Assess' and 'Interpret and Prioritize' steps of the SBTN's 5-
step approach (SBTN, 2020). The TNFD, requires under 'Strategy Disclosure D' (TNFD, 2023) that the
location of assets or activities are disclosed for 'priority locations', defined as, inter alia, those which
have 'high ecosystem integrity' and those with 'rapid decline in ecosystem integrity'. When
implementing actions, and reporting on results, businesses and financial institutions will also need to
understand their potential impacts and dependencies on ecosystems before going on to monitor
and disclose how the integrity of these ecosystems is increasing or decreasing over time. The EII can
support all of these use cases.

In the case of both state and non-state actor users, the EII should be seen as complementary to, and
ideally used in conjunction with, measures of other facets of biodiversity. Complementary species
metrics, for instance, include measures of species importance and threat across the world, such as
the Species Threat Abatement and Restoration (STAR) metric based on the International Union for
Conservation of Nature (IUCN) Red List (Mair *et al*., 2021), or range rarity metrics such as those
presented in Hill et al. (2019). The Red List of Ecosystems (RLE) provides a framework for assessment
and prioritisation of ecosystems analogous to the species threat assessments used to inform the Red
List of Species (Rodriguez *et al.*, 2015). The EII can be paired with the RLE to provide standardised
information on ecosystem integrity to guide RLE assessments, or it can be used to complement the
information provided by the RLE by providing spatially explicit information on ecosystem integrity
across different ecosystems that is regularly updated.

The index has been designed to utilise datasets that track trend data, allowing annual updates as
well as the potential to calculate historical trendlines. The EII framework also has the flexibility to
allow exploration of different management practices at a local level by businesses, or pathways to

sustainability with the alignment of the EII framework to national or global scenarios. The index can
be disaggregated into its component layers and this may be informative when determining
underlying causes of low ecosystem integrity; however, we encourage its use as an aggregated
index, with actions focussed on improving integrity as a whole, rather than just one aspect of
ecosystem health. Although many nature-positive actions will be beneficial for all three components
of integrity, it should be recognised that components may respond at different rates in terms of
recovery towards a more natural state. These temporal lags should be considered when planning
actions and in the tracking of progress towards targets for nature.

The EII has limitations in that it does not yet directly quantify degradation to ecosystem integrity
caused by climate change (although recent climate change impacts may contribute to lower scores
through changes to NPP as reflected in the functioning layer). We note that many of the high
latitude areas that appear unimpacted in this analysis may be at particular risk from climate change
(Beaumont *et al.* 2011; Asamoah *et al.* 2021). Another limitation is that the composition and
structural components are derived from models that assume the same pressure response
relationships across all terrestrial ecosystems (as per the layers from which these components are
derived). Further work will target these limitations to examine the sensitivity of the components to
these assumptions.

Research currently in progress aims to 1) calculate EII for freshwater and marine habitats, 2) link EII
with risk to business operations through loss of ecosystem services, and 3) assess business
counterfactuals for EII. Future development of EII could be enhanced by linking the models used
here to emerging temporal 10m land cover products (for instance, the Sentinel-2 10m Land
Use/Land Cover Time Series; Karra *et al.,* 2021 or the Dynamic World layer; Brown *et al*., 2022) or to
ground-based biodiversity monitoring to provide higher resolution information on biodiversity
change at the local scale, enabling more regular updates of the EII.

The EII shows dramatic degradation to ecosystem integrity over much of the terrestrial area globally.
It has been designed to be flexible, responsive and updateable, and useful at multiple scales. As a
metric it provides an important means for countries and businesses to measure ecosystem integrity
within areas of interest, to set targets and to take action to enhance ecosystem integrity at
appropriate geographical scales.

Methods
250

Structural integrity
251

An ideal measurement of structural integrity should capture both complexity and the multi-
dimensionality of an ecosystem. This should include a measure of how intact the vertical or internal
structure of a given location is, as well as the intactness of such features in the surrounding
landscape (Carter *et al*., 2019; Grantham *et al*., 2020; Ehbrecht *et al.*, 2021; Rayfield *et al.*, 2011).

A single range of measurable characteristics is unsuitable for measuring ecosystem structural
integrity at the global scale because the 'natural' state of ecosystem structure varies greatly by
realm, ecoregion, and habitat type (e.g., Tierney *et al*., 2009). We therefore focus the structural
integrity component on capturing the landscape level intactness through a knowledge of where
human pressures are assumed to cumulatively degrade structural integrity (e.g., Venter *et al*., 2016;

Kennedy *et al*., 2019), combined with an adaptation of the methods established by Beyer *et al*.
(2019) to produce a layer that is sensitive to habitat patch size, fragmentation, and connectivity. This
is achieved by comparing each location on a map not just to a natural reference for that point, but
the whole neighbourhood around that point to a counterfactual landscape in which no pressures
occur, computed at a scale in agreement with other connectivity metrics (e.g., Saura *et al*., 2018).

To measure structural integrity, we used the Human Modification Index (HMI) layer (Kennedy *et al.,*
2019) that is based upon the following pressure layers: croplands, pasturelands, rangelands,
plantations, built-up areas, human population density, roads, rails, quarries and mining, wind
turbines, electrical infrastructure and powerlines (SI). We used the same pressure intensities
described in Kennedy *et al.* (2019). Second, we inverted the human modification layer to transform it
in an equivalent of the habitat quality and extent layer in Beyer *et al.* (2019). Finally, we applied the
Beyer *et al.* (2019) method to incorporate landscape quality and fragmentation through the use of a
moving window that compares the quality of a grid cell with that of cells in their neighbourhood,
which we defined as cells within a 27km distance (i.e., grid cells in a 55km x 55km box were used to
estimate the structural integrity of the target cell at the centre of the box). All spatial processing was
done in Google Earth Engine with a resolution of 30 arcsec.

Our layer differs from the one produced in the Beyer *et al.* (2019) paper due to the source 'habitat
quality' layer. Where the original authors use the Human Footprint Index (HFI) (Venter *et al*., 2016),
rescaled with an exponential function, as their layer, we opt for a recreation of the HMI as described
by Kennedy *et al.* (2019). This approach has the benefit of meaningful interpretation: an area with a
value of 0.5 has twice the 'intactness' of an area with a value of 0.25. It also avoids the double use of
neighbourhood functions. For example, the HFI takes roads and applies a buffer to estimate the area
of their impact; when the landscape-level effects are calculated this area is then further expanded,
which may overestimate the spatial extent of these threats to ecosystem structure. In contrast, the
HMI uses the physical footprint of these threats, preventing this issue when comparing at the
landscape scale.

Compositional integrity

A measure of compositional intactness should incorporate elements of both species' abundance and
community composition (Haase *et al.*, 2018; Carter *et al*., 2019). The Biodiversity Intactness Index
(BII) has been suggested as a suitable metric to track global planetary boundaries due to its
relevance to ecosystem health and functioning (Steffen *et al*., 2015; Mace *et al*., 2018). The BII
estimates the change in the makeup of species communities relative to an 'unimpacted' baseline
(Newbold *et al*., 2016). The BII tracks changes in the relative abundance of species within a
community (excluding novel species) due to land use changes as well as other human pressures. The
BII is based upon modelled relationships estimated using a large taxonomically and geographically
representative database, the PREDICTS database (Hudson *et al.,* 2017). Data representing the
communities of sites from many land-uses and many taxonomic groups are used to construct
hierarchical models for the two elements of the BII, abundance and compositional similarity. These
modelled relationships are then projected onto global pressure maps to form the BII layer.

Mixed effects models were fitted using land use and density of human population as fixed effects
(for full modelling details see SI) and block nested within study as random effects. A novel land use
map was constructed using 1) land use data from the Historic Land Dynamics Assessment+ (HILDA+)
project (Winkler *et al.,* 2021), 2) the Spatial Database of Planted Trees (Harris *et al.,* 2019), 3) the

Global Land Analysis and Discovery (GLAD) database (Potapov *et al.,* 2022) and 4) the Potential
Threats to Soil Biodiversity data from the Global Soil Biodiversity Atlas (JRC, 2016). The HILDA
dataset provides land use details denoting natural areas, cropland, pasture/rangeland and urban
from 1899 to present. We supplemented this data as follows: 1) within current day natural areas we
tracked transitions and noted time since transition back to natural to distinguish primary vegetation,
mature secondary vegetation, intermediate secondary vegetation and young secondary vegetation,
2) within agricultural areas we distinguish use intensity using levels of soil threat, 3) we distinguish
between pasture and rangeland using overlap with the GLAD database, and 4) we distinguish
plantation forests using the SDPT. (For further details on how the land use map was constructed see
SI).

Functional integrity

Attempts at mapping ecosystem functioning have explored changes in the values of specific
ecosystem function variables. One example comes from the framework assessing Human
Appropriation of Net Primary Productivity (HANPP), which explores departures from potential NPP
values to investigate where human production systems have modified ecosystem functioning and
the extent of this change (Mayer *et al.,* 2021). Other authors have taken a very different approach to
functioning through mapping patterns in the diversity of functional traits. For instance, Faurby and
Svenning (2015) estimated changes to the richness of mammalian functional traits. However, no
single metric has yet been developed that is considered to represent all aspects of ecosystem
functioning.

In the absence of current alternative more comprehensive methods, we opted for adapting the
approach to focus on one key ecosystem function, NPP, because of its well-documented association
with ecosystem functioning (Malhi *et al*., 2011; Mayer *et al*., 2021), its advantages in terms of spatial
resolution (layers are available in raster format at 1km^2 resolution or higher) and update frequency
(e.g., MODIS releases a global new NPP layer annually). Indeed, within the planetary boundaries
framework, HANPP, as a fraction of Holocene levels of NPP, is the metric of choice to measure
functional biosphere integrity (Richardson *et al*., 2023). NPP is also known to be closely associated
with the provision of key ecosystem services such as mass flow regulation and atmospheric
regulation, pest control and water purification (Harrison *et al*., 2014).

We developed a global layer of potential NPP used as a 'natural' reference and compared it with a
current-day NPP layer derived from remote sensing to map changes in NPP. Natural levels of NPP
were modelled per grid cell using environmental variables trained on mean NPP levels (Running and
Zhao, 2015) measured within strictly managed protected areas (IUCN categories I and II) between
2001 and 2021 (UNEP-WCMC and IUCN, 2017). We used a generalised linear model framework, with
a Gamma distribution for the response variable and a log-link. Variables were selected for testing
based on a literature review of likely predictors of NPP and included: latitude, bioclimatic variables
(Brun et al. 2022); mean, min and max solar radiation (Fick and Hijmans, 2017); aridity (Global Aridity
Index, Zomer and Trabucco, 2022); total nitrogen, cation exchange capacity, predicted sand
concentration, pH of water in soil (Poggio *et al.,* 2021); continuous heat-insolation load index (CHILI,
Theobald *et al*., 2015); roughness of terrain, slope, topographic position index, terrain ruggedness
index (Amatulli *et al.,* 2018); and landforms (Sayre *et al.,* 2020). As the selection process did not
result in a single top model, spatial projections of the two best models were used as inputs to the
NPP calculation, along with two previously published potential NPP layers (Brun *et al.* 2022; Mayer
et al. 2021).

As the four potential NPP models provide a range of values, we included two types of uncertainty in
our calculations. Firstly, for each grid cell, we calculated the range of possible values using the
lowest value and highest value observed between all models. All measures were calculated as
movements away from this range (i.e., where NPP was estimated to be lost, the value was compared
to the lowest potential value, and where NPP was estimated to be gained, the value was compared
to the highest potential value). Secondly, we accounted for inter-seasonal variation by modelling the
coefficient of variation of annual measures between 2001 and 2021 following the same
methodology as that undertaken for the potential NPP. This provided an estimation of what natural
level of variation may be expected within each grid cell and allowed us to adjust our calculations to
account for movements outside the natural levels of variation.

Ecosystem functioning occurs on different scales. To assess changes to ecosystem functioning on a
broader scale the absolute loss or gain of NPP per grid cell was calculated; however, to account for
changes to ecosystem functioning on a local scale the proportion of NPP measured versus potential
per grid cell was calculated. The mean of the proportional change and the decile value of the
absolute change provided the final functional integrity score.

Aggregation to derive the global Ecosystem Integrity Index

To calculate the Ecosystem Integrity Index (EII) value per grid cell we first extracted the lowest
scoring of the three ecosystem components within that grid cell. This calculation follows the logic
that all components of integrity are integral and therefore the overall condition of the ecosystem is
limited by the lowest component. This emphasizes the interconnected nature of ecosystem integrity
and avoids issues that may have been caused by an averaging/additive approach. For example,
phenomena such as extinction debt (Kuussaari *et al.,* 2009) suggest it would be prudent to refrain
from allowing moderate values where any one component has been significantly weakened.
However, simply taking the lowest value cannot distinguish between an area where all three
components are degraded versus an area where only a single component is degraded. We therefore
adjusted the EII value using fuzzy algebraic summing (Zadeh, 1965, see SI) to downgrade the EII
score based on the level of degradation of the two components that were not providing the lowest
score.

Mean EII scores were calculated at biome and ecoregion levels using terrestrial biomes and
ecoregions spatial data (Dinerstein *et al.,* 2019), and ecosystem levels using Sayre *et al*., (2020).

Acknowledgements

We would like to thank Mike Harfoot, Sharon Brooks, Sebastian Bekker, Maria Julia Oliva, Tom
Evans, Tom Brooks, Emily Nicholson, David Keith, Malcolm Starkey, Craig Beatty, Marco Daldos Pirri,
Varsha Vijay and Pamela Collins for their comments and review of the EII methodology. This output
has been funded in whole or part by the UK Research and Innovation's Global Challenges Research
Fund under the Trade, Development and the Environment Hub project (project number
ES/S008160/1) and by the Science Based Targets Network.

References

Amatulli, G., Domisch, S., Tuanmu, M.N., Parmentier, B., Ranipeta, A., Malczyk, J. and Jetz, W. (2018).
A suite of global, cross-scale topographic variables for environmental and biodiversity modelling.
Scientific data, 5(1), pp.1-15.

Asamoah, E., Di Marco, M., Watson, J., Beaumont, L., Venter, O., and Maina, J. (2021). Tracing
climate and land-use instability reveals new insights into the future of Earth's remaining wilderness.
Preprint: DOI: 10.21203/rs.3.rs-620309/v1

Barlow, J., Lennox, G.D., Ferreira, J., Berenguer, E., Lees, A.C., Nally, R.M., Thomson, J.R., Ferraz,
S.F.D.B., Louzada, J., Oliveira, V.H.F. and Parry, L. (2016). Anthropogenic disturbance in tropical
forests can double biodiversity loss from deforestation. *Nature*, 535(7610), pp.144-147.

Bianchi, F.J., Booij, C.J.H. and Tscharntke, T. (2006). Sustainable pest regulation in agricultural
landscapes: a review on landscape composition, biodiversity and natural pest control. *Proceedings of*
the Royal Society B: Biological Sciences, 273(1595), pp.1715-1727.

Beaumont, L.J., Pitman, A., Perkins, S., Zimmermann, N.E., Yoccoz, N.G. and Thuiller, W. (2011).
Impacts of climate change on the world's most exceptional ecoregions. P*roceedings of the National*
Academy of Sciences, 108(6), pp.2306-2311.

Beyer, H. L., Venter, O., Grantham, H. S., & Watson, J. E. (2020). Substantial losses in ecoregion
intactness highlight urgency of globally coordinated action. *Conservation Letters*, *13*(2), e12692.
https://doi.org/10.1111/conl.12692

Blumetto, O., Castagna, A., Cardozo, G., García, F., Tiscornia, G., Ruggia, A., Scarlato, S., Albicette,
M.M., Aguerre, V. and Albin, A. (2019). Ecosystem Integrity Index, an innovative environmental
evaluation tool for agricultural production systems. *Ecological indicators*, 101, pp.725-733.

Bonan, G. B. (2008). Forests and climate change: forcings, feedbacks, and the climate benefits of
forests. *Science* 320, pp.1444–1449.

Brown, C. F., Brumby, S. P., Guzder-Williams, B., Birch, T., Hyde, S. B., Mazzariello, J., Czerwinski, W.,
Pasquarella, V.J., Haertel, R., Ilyushchenko, S., Schwehr, K. Weisse, M., Stolle, F., Hanson, C., Guinan,
O., Moore, R. & Tait, A. M. (2022). Dynamic World, Near real-time global 10 m land use land cover
mapping. *Scientific Data*, 9(1), pp. 1-17.

Burgess, N.D., D'Amico Hales, J., Ricketts, T. & Dinerstein, E. (2006). Factoring species, non-species
values and threats into biodiversity priority-setting across the ecoregions of Africa and its islands.
Biological Conservation, 127, 383-401

Carter, S.K., Fleishman, E., Leinwand, I.I., Flather, C.H., Carr, N.B., Fogarty, F.A., Leu, M., Noon, B.R.,
Wohlfeil, M.E. and Wood, D.J. (2019). Quantifying ecological integrity of terrestrial systems to inform
management of multiple-use public lands in the United States. *Environmental management*, *64*(1),
pp.1-19. https://doi.org/10.1007/s00267-019-01163-w

Carvalheiro, L.G., Seymour, C.L., Veldtman, R. and Nicolson, S.W. (2010). Pollination services decline
with distance from natural habitat even in biodiversity-rich areas. *Journal of Applied Ecology*, 47(4),
pp.810-820.

CBD (Convention on Biological Diversity). (1992). Convention on Biological Diversity: text and
annexes. https://www.cbd.int/convention/text/

CBD (Convention on Biological Diversity). (2000). Decision V/6 Ecosystem approach. Fifth meeting of
the Conference of the Parties to the Convention on Biological Diversity, Nairobi.

CBD (Convention on Biological Diversity). (2022a). Decision 15/4 Kunming-Montreal Global Biodiversity Framework. Fifteenth meeting of the Conference of the Parties to the Convention on Biological Diversity, Montreal.

CBD (Convention on Biological Diversity). (2022b). Decision 15/5 Monitoring Framework for the Kunming-Montreal Global Biodiversity Framework. Fifteenth meeting of the Conference of the Parties to the Convention on Biological Diversity, Montreal.

Dinerstein, E., Olson, D., Joshi, A., Vynne, C., Burgess, N.D., Wikramanayake, E., Hahn, N., Palminteri, S., Hedao, P., Noss, R. and Hansen, M. (2017). An ecoregion-based approach to protecting half the terrestrial realm. *BioScience*, 67(6), pp.534-545.

Ehbrecht, M., Seidel, D., Annighöfer, P., Kreft, H., Köhler, M., Zemp, D.C., Puettmann, K., Nilus, R., Babweteera, F., Willim, K. and Stiers, M. (2021). Global patterns and climatic controls of forest structural complexity. *Nature communications*, 12(1), pp.1-12.

Faurby, S. and Svenning, J.C. (2015). Historic and prehistoric human-driven extinctions have reshaped global mammal diversity patterns. *Diversity and Distributions*, 21(10), pp.1155-1166.

Fick, S.E. and Hijmans, R.J. (2017). WorldClim 2: new 1km spatial resolution climate surfaces for global land areas. *International Journal of Climatology* 37 (12): 4302-4315

Grantham, H.S., Duncan, A., Evans, T.D., Jones, K.R., Beyer, H.L., Schuster, R., Walston, J., Ray, J.C., Robinson, J.G., Callow, M. and Clements, T. (2020). Anthropogenic modification of forests means only 40% of remaining forests have high ecosystem integrity. *Nature communications*, 11(1), pp.1-10.

Gibson, L. (2011). Primary forests are irreplaceable for sustaining tropical biodiversity. *Nature* 478, https://doi.org/10.1038/nature10425.

Haase, P., Tonkin, J.D., Stoll, S., Burkhard, B., Frenzel, M., Geijzendorffer, I.R., Häuser, C., Klotz, S., Kühn, I., McDowell, W.H. and Mirtl, M. (2018). The next generation of site-based long-term ecological monitoring: Linking essential biodiversity variables and ecosystem integrity. *Science of the Total Environment*, 613, pp.1376-1384.

Hansen, A.J., Noble, B.P., Veneros, J., East, A., Goetz, S.J., Supples, C., Watson, J.E., Jantz, P.A., Pillay, R., Jetz, W. and Ferrier, S. (2021). Toward monitoring forest ecosystem integrity within the post-2020 Global Biodiversity Framework. *Conservation Letters*, *14*(4), p.e12822.

Harris, N., Goldman, E.D. and Gibbes, S. (2019). Spatial Database of Planted Trees (SDPT Version 1.0)

Harrison, P.A., Berry, P.M., Simpson, G., Haslett, J.R., Blicharska, M., Bucur, M., Dunford, R., Egoh, B., Garcia-Llorente, M., Geamănă, N. and Geertsema, W. (2014). Linkages between biodiversity attributes and ecosystem services: A systematic review. *Ecosystem services*, 9, pp.191-203.

Hill, S.L., Gonzalez, R., Sanchez-Ortiz, K., Caton, E., Espinoza, F., Newbold, T., Tylianakis, J., Scharlemann, J.P., De Palma, A. and Purvis, A. (2018). Worldwide impacts of past and projected future land-use change on local species richness and the Biodiversity Intactness Index. *BioRxiv*, p.311787.

Hill, S.L., Arnell, A., Maney, C., Butchart, S.H., Hilton-Taylor, C., Ciciarelli, C., Davis, C., Dinerstein, E., Purvis, A. and Burgess, N.D. (2019). Measuring forest biodiversity status and changes globally. *Frontiers in Forests and Global Change*, *2*, p.70. https://doi.org/10.3389/ffgc.2019.00070

Hudson, L.N., Newbold, T., Contu, S., Hill, S.L., Lysenko, I., De Palma, A., Phillips, H.R., Alhusseini, T.I.,
Bedford, F.E., Bennett, D.J. and Booth, H. (2017). The database of the PREDICTS (projecting
responses of ecological diversity in changing terrestrial systems) project. *Ecology and evolution*, 7(1),
pp.145-188.

IUCN and UNEP-WCMC. (2022). The World Database on Protected Areas (WDPA). April. Cambridge,
UK: UNEP-WCMC. www.protectedplanet.net. Accessed through Resource Watch, (accessed 04-Mar-
2022). www.resourcewatch.org.

IUCN-CEM. (2022). The IUCN Red List of Ecosystems. Version 2022-1. http://iucnrle.org.

Karra, K., Kontgis, C., Statman-Weil, Z., Mazzariello, J. C., Mathis, M., and Brumby, S. P. (2021).
Global land use/land cover with Sentinel 2 and deep learning. *In 2021 IEEE international geoscience
and remote sensing symposium IGARSS,* pp. 4704-4707. IEEE.

Keith, D. A., Ferrer-Paris, J. R., Nicholson, E., Kingsford, R. (2020), IUCN Global Ecosystem Typology
2.0. IUCN, Swizerland.

Kennedy, C.M., Oakleaf, J.R., Theobald, D.M., Baruch-Mordo, S. and Kiesecker, J., (2019). Managing
the middle: A shift in conservation priorities based on the global human modification
gradient. *Global Change Biology*, *25*(3), pp.811-826. https://doi.org/10.1111/gcb.14549

Kuussaari, M., Bommarco, R., Heikkinen, R.K., Helm, A., Krauss, J., Lindborg, R., Öckinger, E., Pärtel,
M., Pino, J., Rodà, F. and Stefanescu, C. (2009). Extinction debt: a challenge for biodiversity
conservation. *Trends in ecology & evolution*, 24(10), pp.564-571.

Leopold, A. (1949). *A Sand County almanac*. Oxford University Press, USA, Oxford and New York

Lewis, S.L., Lopez-Gonzalez, G., Sonké, B., Affum-Baffoe, K., Baker, T.R., Ojo, L.O., Phillips, O.L.,
Reitsma, J.M., White, L., Comiskey, J.A. and Ewango, C.E. (2009). Increasing carbon storage in intact
African tropical forests. *Nature*, *457*(7232), pp.1003-1006.

Locke, H., Rockstrom, J., Bakker, P., Bapna, M., Gough, M., Hilty, J., Lambertini, M., Morris, J.,
Polman, P., Rodriguez, C.M., Samper, C., Snjayan, M., Zabey, E., and Zurita, P. (2022) A nature-
positive world: the global goal for nature.

Mace, G.M., Barrett, M., Burgess, N.D., Cornell, S.E., Freeman, R., Grooten, M. and Purvis, A. (2018).
Aiming higher to bend the curve of biodiversity loss. *Nature Sustainability*, 1(9), pp.448-451.

Mayer, A., Kaufmann, L., Kalt, G., Matej, S., Theurl, M.C., Morais, T.G., Leip, A. and Erb, K.H. (2021).
Applying the Human Appropriation of Net Primary Production framework to map provisioning
ecosystem services and their relation to ecosystem functioning across the European Union.
Ecosystem services, 51, pp.101344.

Mair, L., Bennun, L.A., Brooks, T.M., Butchart, S.H., Bolam, F.C., Burgess, N.D., Ekstrom, J.M., Milner-
Gulland, E.J., Hoffmann, M., Ma, K. and Macfarlane, N.B. (2021). A metric for spatially explicit
contributions to science-based species targets. *Nature Ecology & Evolution*, 5(6), pp.836-844.

Malhi, Y., Doughty, C., and Galbraith, D. (2011). The allocation of ecosystem net primary productivity
in tropical forests. *Phil. Trans R. Soc. B.*, 366, pp. 3225-3245.

Mello, K. D., Valente, R. A., Randhir, T. O. & Vettorazzi, C. A. (2018). Impacts of tropical forest cover
on water quality in agricultural watersheds in southeastern Brazil. *Ecological Indicators*. 93, 1293–
1301.

Milner-Gulland, E. J. (2022). Don't dilute the term Nature Positive. *Nature Ecology & Evolution*, 1-2.

Mora, F. (2017). A structural equation modeling approach for formalizing and evaluating ecological
integrity in terrestrial ecosystems. *Ecological Informatics*, 41, pp.74-90.

Newbold, T., Hudson, L.N., Arnell, A.P., Contu, S., De Palma, A., Ferrier, S., Hill, S.L., Hoskins, A.J.,
Lysenko, I., Phillips, H.R. and Burton, V.J. (2016). Has land use pushed terrestrial biodiversity beyond
the planetary boundary? A global assessment. *Science*, 353(6296), pp.288-291.
https://doi.org/10.1126/science.aaf2201

Parrish, J. D., Braun, D. P. & Unnasch, R. S. (2003). Are we conserving what we say we are?
Measuring ecological integrity within protected areas. *Bioscience* 53, 851–860.

Perkl, R. (2017). Measuring landscape integrity (LI): development of a hybrid methodology for
planning applications. *Journal of Environmental Planning and Management*, 60(1), pp. 92-114.

Poggio, L., De Sousa, L.M., Batjes, N.H., Heuvelink, G., Kempen, B., Ribeiro, E. and Rossiter, D. (2021).
SoilGrids 2.0: producing soil information for the globe with quantified spatial uncertainty. *Soil*, 7(1),
pp.217-240.

Potapov P., Hansen M.C., Pickens A., Hernandez-Serna A., Tyukavina A., Turubanova S., Zalles V., Li
X., Khan A., Stolle F., Harris N., Song X.-P., Baggett A., Kommareddy I., Kommareddy A. (2022) The
global 2000-2020 land cover and land use change dataset derived from the Landsat archive: first
results. Frontiers in Remote Sensing [https://doi.org/10.3389/frsen.2022.856903] [Open PDF]

Rayfield, B., Fortin, M.J. and Fall, A. (2011). Connectivity for conservation: a framework to classify
network measures. *Ecology*, 92(4), pp.847-858.

Running, S. W., and Zhao, M. (2019). Daily GPP and Annual NPP (MOD17A2H/A3H) and Year-End
Gap- Filled (MOD17A2HGF/A3HGF) Products NASA Earth Observing System MODIS Land Algorithm
(For Collection 6), MODIS Land Team, Version 4.2, pp.35.

Running, S.W. and Zhao, M. (2015). Daily GPP and annual NPP (MOD17A2/A3) products NASA Earth
Observing System MODIS land algorithm. *MOD17 User's Guide*, 2015, pp.1-28.

Richardson, K., Steffen, W., Lucht, W., Bendtsen, J., Cornell, S.E., Donges, J.F., Drüke, M., Fetzer, I.,
Bala, G., von Bloh, W. and Feulner, G. (2023). Earth beyond six of nine planetary boundaries. *Science
Advances*, *9*(37), p.eadh2458.

Saura, S., Bertzky, B., Bastin, L., Battistella, L., Mandrici, A. and Dubois, G. (2018). Protected area
connectivity: Shortfalls in global targets and country-level priorities. *Biological conservation*, 219,
pp.53-67.

Sayre, R., Karagulle, D., Frye, C., Boucher, T., Wolff, N.H., Breyer, S., Wright, D., Martin, M., Butler, K.,
Van Graafeiland, K. and Touval, J. (2020). An assessment of the representation of ecosystems in
global protected areas using new maps of World Climate Regions and World Ecosystems. *Global
Ecology and Conservation*, 21, p.e00860.

SBTN (2020). SCIENCE-BASED TARGETS for NATURE: Initial Guidance for Business

TNFD (2022). The TNFD Nature-Related Risk and Opportunity Management and Disclosure
Framework. Beta v0.2.

TNFD (2023a). Task force on nature-related financial disclosures. Guidance on the identification and
assessment of nature-related issues: the LEAP approach.

TNFD (2023b). Recommendations of the Taskforce on Nature-Related Financial Disclosures.

Theobald, D. M., Harrison-Atlas, D., Monahan, W. B., and Albano, C. M. (2015). Ecologically-Relevant
Maps of Landforms and Physiographic Diversity for Climate Adaptation Planning. *PLOS ONE* 10, no.
12: e0143619. https://doi.org/10.1371/journal.pone.0143619.

Steffen, W., Richardson, K., Rockström, J., Cornell, S.E., Fetzer, I., Bennett, E.M., Biggs, R., Carpenter,
S.R., De Vries, W., De Wit, C.A. and Folke, C. (2015). Planetary boundaries: Guiding human
development on a changing planet. *Science*, 347(6223), p.1259855.

Tierney, G.L., Faber-Langendoen, D., Mitchell, B.R., Shriver, W.G. and Gibbs, J.P. (2009). Monitoring
and evaluating the ecological integrity of forest ecosystems. *Frontiers in Ecology and the*
Environment, 7(6), pp.308-316.

UNCCD, (2022). Summary for Decision Makers. Global Land Outlook, second edition. United Nations
Convention to Combat Desertification, Bonn.

Venter, O., Sanderson, E.W., Magrach, A., Allan, J.R., Beher, J., Jones, K.R., Possingham, H.P.,
Laurance, W.F., Wood, P., Fekete, B.M. and Levy, M.A. (2016). Sixteen years of change in the global
terrestrial human footprint and implications for biodiversity conservation. *Nature communications*,
7(1), pp.1-11.

Zadeh, L.A. (1965). Fuzzy sets. *Information and control*, 8(3), pp. 338-353.

Zomer, R.J., Xu, J. and Trabuco, A. (2022). Version 3 of the Global Aridity Index and Potential
Evapotranspiration Database. *Scientific Data* 9, 409. https://www.nature.com/articles/s41597-022-
01493-1

Zuur, A. F., Ieno, E. N. and Saveliev A. A. (2009). *Mixed Effects Models and Extensions in Ecology with*
R. Springer, New York

573