Mechanistic Deep Learning Framework on Cell Traits Derived from Single-Cell Mass Cytometry Data

Bo Wang[1], Sandra S. Zinkel[2,3], and Eric R. Gamazon[1, 4, 5, 6]

[1]Division of Genetic Medicine, Department of Medicine, Vanderbilt University Medical Center, Nashville, TN, USA

[2]Department of Cell and Developmental Biology, Vanderbilt University, Nashville, TN, USA

[3]Department of Medicine, Vanderbilt University Medical Center, Nashville, TN, USA

[4]Vanderbit Genetics Institute, Vanderbilt University Medical Center, Nashville, TN, USA

[5]Data Science Institute, Vanderbilt University Medical Center, Nashville, TN

[6]Clare Hall, University of Cambridge, Cambridge, United Kingdom

Send correspondence to:

Eric R. Gamazon <eric.gamazon@vumc.org> or Sandra Zinkel <sandra.zinkel@vumc.org>

Abstract

Germline genetic variations can alter cellular differentiation, potentially impacting the response of immune cells to inflammatory challenges. Current variant- and gene- based methods in mouse and human models have established associations with disease phenotypes; however, the underlying mechanisms at the cellular level are less well-understood. Immunophenotyping by multi-parameter flow cytometry, and more recently mass cytometry, has allowed high-resolution identification and characterization of hematopoietic cells. The obtained characterization yields increased dimensionality; however, conventional analysis workflows have been inefficient, incomplete, or unreliable. In this work, we develop a comprehensive machine learning framework – MDL4Cyto – that is tailored to the analysis of mass cytometry data, incorporating statistical, unsupervised learning, and supervised learning models. The statistical modeling can be used to illuminate cell fate decision and cell-type dynamics. The unsupervised learning models along with complementary marker enrichment analyses highlight genetic perturbations that are significantly associated with alterations in cell populations in the hematopoietic system. Furthermore, our supervised learning models, including deep learning and tree-based algorithms, address the bottleneck to data pre-processing that characterizes conventional workflows and generate inferences (e.g., on marker/cell-type interactions) from raw experimental characterization. Notably, we reveal a close relationship among network design, prediction performance, and the underlying biological context. We show that the network architecture extracted from the differentiation cascade of the investigated biological system yields enhanced prediction performance. The presented methodology will enable new insights into hematopoietic differentiation at baseline and following perturbation.

Highlights

- Analysis pipeline on mass cytometry data with high-performance implementation of statistical, unsupervised learning, and supervised learning models

- Concordance of machine learning results with biological contexts

- Biologically-informed neural network designs enhance prediction performance

Graphical Abstract

Keywords

Machine Learning, Mass Cytometry, CyTOF, Hematopoiesis, Cellular Differentiation, Regulatory Cascade, Clonal Hematopoiesis of Indeterminate Potential

Introduction

Hematopoiesis is a hierarchical and dynamic process. Proper blood cell development requires

highly coordinated and tightly regulated cytokine signals, affecting cell and stage-appropriate gene

expression. Genetically modified mouse models are an invaluable model system for studying

hematopoiesis, due to their defined genetic background, wealth of reagents to precisely define

hematopoietic cell populations, ability to undergo precise genetic modification, and the wealth of data

and assays arising from decades of study. Cytometry tools have proved to be essential in simultaneously

analyzing labeled molecules at the single-cell level and, consequently, have been widely implemented in

basic and clinical research[1]. While traditional fluorescence cytometers can detect up to 35 markers, the

technique is limited by spectral overlap of fluorophores. Single-cell mass cytometry (Cytometry by Time-

Of-Flight [CyTOF]) enables multiple-parameter tracking by the use of metal-isotope-tagged antibodies,

overcoming the inherent limitations of spectral overlap. Therefore, a more comprehensive panel of

cellular functions (achieved from significantly expanded feature capacity[2]) along with intracellular

signaling states can be explored to identify disease-associated changes or functionally-relevant states;

differential expression under external perturbation or intracellular response is more easily elucidated[3].

Insights from cytometry characterization are traditionally extracted from manual gating.

Automated workflows such as FlowCore[4], PhenoGraph[5] or the recently reported Tracking Responders

EXpanding (T-REX)[6] have been introduced with significantly enhanced processing efficiency. However,

these pipelines are still limited by expert intervention into (manual) data pre-processing, where key

thresholds for transformation, gating or cell type clustering are tailored to the investigated systems.

Recently, machine learning algorithms have attracted significant attention for their extensive application

to the analysis of cytometry data[6–8]. Specifically, beyond functioning as embedded algorithms in

pipelines, machine learning tools provide "end-to-end" solutions, generating inferences or predictions

directly from raw experimental characterization, with potential to uncover novel associations that are

obscured by conventional gating strategies.

Hematopoiesis has served as a paradigm of stem cell biology and cellular differentiation[9].

Genetic perturbations within cells (in particular, Hematopoietic Stem and Progenitor Cells [HSPCs])

induce challenges to cellular functions, thus leading to various diseases or conditions, e.g., Clonal

Hematopoiesis of Indeterminate Potential (CHIP), Myelodysplastic Syndromes (MDS), and Acute Myeloid

Leukemia (AML). Tet2 inactivating mutations are among the early mutations that initiate clonal

expansion (CHIP). As the most expressed Tet gene in the hematopoietic tissue (especially in HSCs), Tet2

functions as a tumor suppressor whose haplo-insufficiency initiates myeloid and lymphoid

transformations[8]. Tet2 has been shown to regulate resolution of inflammatory responses. Tet2 KO mice

and cell lines display prolonged and enhanced inflammatory cytokine signaling, which impacts HSPC

differentiation.

Hematopoietic homeostasis is further controlled through tight regulation of programmed cell

death. The two major types of cell death, apoptosis (programmed cell death) and necrosis (uncontrolled

cell death), maintain homeostasis by downregulating detrimental inflammation and prevent

transformation by allowing efficient removal of damaged cells. It has been widely recognized that the

genes in the Bcl-2 family are critical in regulating apoptosis[10,11], comprising of both pro-survival

members and pro-apoptotic members. The functions of BID (pro-apoptotic member and multidomain

effector) were investigated and confirmed in our previous genomic studies[12]. Specifically, an association

between decreased genetically determined BID expression and myocardial infarction susceptibility was

identified in a discovery cohort using a gene-based association method[13], and validated by two

independent genome-wide association studies (GWAS). In the follow-up studies, we further tested the

functionalities of proteins in the family, including multidomain effectors Bax, Bak and the BH3-only

protein, Bid, in murine models[14]. We found that deletion of Bax/Bak inhibits both bone marrow

apoptosis and bone marrow necroptosis. Further deletion of BH3-only member Bid (TKO) leads to robust

activation of necroptosis, which results in a phenotype characterized by dysplasia and cytopenia,

resembling the bone marrow failure disorder MDS (myelodysplastic syndrome)[14].

To address the limitations of conventional cytometry data analysis pipelines[5–8,15], we developed

a comprehensive *Mechanistic Deep Learning for Cytometry* (MDL4Cyto) framework, implementing

statistical, unsupervised learning, and supervised learning models. The framework was evaluated against

bone marrow samples of murine models that have been genetically perturbed in the hematopoietic

system, including Wild Type (WT), *VavCreBaxBakBid* triple-knockout (TKO), and *VavCreTet2* knock-out

(Tet2 KO) mice, and validated with biological observations. Notably, building on our previous work[16,17],

we designed the architectures of the neural networks using the intrinsic hierarchy of the hematopoietic

system as a guide, establishing a *biological feature – network architecture – phenotype* relationship. We

show that biologically-informed network designs could function as probes on the regulatory cascade

with prediction performance as the evaluation metric. Our framework may inform future machine

learning implementations to address fundamental questions in cell biology.

Results

Overview of mass cytometry data and MDL4Cyto framework

Mice from three genetic murine models – WT, Tet2 and TKO – were treated with Phosphate-

buffered saline (PBS) or lipopolysaccharide (LPS). LPS is a major surface membrane polysaccharide found

on Gram-negative bacteria and is a typical pathogen-associated molecular pattern (PAMP).

Inflammation triggered by LPS stimulation activates a robust innate immune response and can induce

the proliferation of hematopoietic progenitor cells[18] and the production of cytokines such as TNFα,

interleukin and IL-2. Bone marrow from these mice was subsequently harvested and processed for

cytometry experiments. Besides genetic perturbations, all the samples challenged by LPS stimulation

were compared to the PBS vehicle control (Figure 1A). Across different types of samples (WT-PBS, WT-

LPS, Tet2-PBS, Tet2-LPS, TKO-PBS, and TKO-LPS), lineage marker groups (Stem Cells, Pan Leukocyte

Markers, T Lymphocytes, Mature Myeloid Cells, B cells, and Erythrocytes) and functional marker groups

(Cytokines, Cell death, and Proliferation), were tagged and monitored (Table S1). As sample data profiles

obtained from the mass cytometry measurements varied in length, we imputed data points at each of

the six sample data profiles via binning methods with 30 seconds.

We developed MDL4Cyto, an analysis pipeline for mass cytometry data (Figure 1B and

Methods), implementing statistical, unsupervised, and supervised learning models. Statistical tests on

the cell count of key lineage and functional markers were performed, aiming for a glimpse of the

underlying biology. As both genetic perturbations and LPS stimulation may induce inflammatory changes

in hematopoiesis, we applied Generalized Mixed-Effects Model (GMM) estimation to determine the

extent to which cell count was statistically altered by these modifications. Two established workflows,

namely PhenoGraph[5] and T-REX[6], were benchmarked as established pipelines capable of annotating cell

populations or capturing cell responses. In addition, supervised learning models were explicitly designed

to probe marker/cell-type interactions within the regulatory cascade, as well as to predict sample type

directly from the raw mass cytometry characterization. Nine machine learning models were evaluated,

including Linear Regression (LR), Dense Neural Network (DNN), Boosted Tree (BT), Random Forest (RF),

stacked Dense Neural Network - Random Forest (DNN-RF), Convolutional Neural Network in One

Dimension (CNN-1D), Convolutional Neural Network in Two Dimensions (CNN-2D), Recurrent Neural

Network with Long Short-Term Memory cell (RNN-LSTM), and Recurrent Neural Network with Gated

Recurrent Units cell (RNN-GRU).

147
Figure 1. MDL4Cyto pipeline for analysis of single-cell mass cytometry data. Bone marrow samples (**A**)
were harvested from genetically perturbed murine models (Wild Type [WT], *VavCreBaxBakBid* triple-
knockout [TKO], and *VavCreTet2* knock-out [Tet2 KO]), which had been treated with LPS stimulation or
PBS vehicle control. Statistical, unsupervised, and supervised learning models were implemented in our
Mechanistic Deep Learning for Cytometry (MDL4Cyto) framework. The analysis pipeline (**B**) was
benchmarked, using the reference dataset obtained from mass cytometry characterization, for the tasks
of differential abundance analysis, cell response detection, and sample type classification.

Statistical modeling of genetically-determined or condition-dependent effects

The cell count summary statistic, which was calculated on the reference dataset, was found to

be altered by genetic perturbations and/or LPS stimulation (Figure S1). Clear differences were observed

for all markers. As reported in our previous work[14] and other studies[19], the TKO sample leads to necrosis

with increasing production of cytokines. Enhanced inflammatory signaling was also observed in Tet2-

deficient samples, as has also been previously described[19]. We observed a statistically significant

difference in cell count for Sca1 (LSK cells), CD45 (leukocyte), TNFα (cytokines), and Caspase 3 (cell

death indicator) with respect to genetic perturbation (Figure 2A). Markers of cytokines were found to be

highly correlated with other surface markers, especially under LPS stimulation (Figure 2B).

Figure 2. Impact of genetic perturbations on cell populations and correlations among the markers. A statistically significant change in cell count was observed for Sca1 (LSK cells), CD45 (leukocyte), cytokines

(TNFα), and Caspase 3 and (cell death indicator) in both the PBS vehicle control and LPS stimulation
condition, demonstrating genetic perturbations altering cell subpopulations **(A)**. Cytokines from genetic
knock-out samples were highly correlated with other surface markers, in particular under LPS
stimulation **(B)**, which is consistent with the biological context. (Wilcoxon test was performed for the
pairwise comparisons; Pearson correlation was calculated for the markers along with 95% confidence
interval. *p<.05, **p<0.01, ***p<0.001, ****p<0.0001).

The data profiles obtained from cytometry characterizations are dynamical panels, i.e., have a

time component. To estimate the overall effects of genotype and condition, we analyzed the

progression of marker Intensity for Ki67 from the aggregated data panel with 1 hour binning window

(Figure 3 and **Methods**). Ki67 is a proliferation protein that is increased in the S phase of the cell cycle

and indicates proliferating cells. We observed progression divergence according to genotype (Figure 3A)

and differentiated marker intensity between conditions (PBS vs LPS, Figure 3B). This is consistent with

our understanding that genetic perturbations and LPS stimulation impose pressures on differentiation

and cell cycle as dynamical processes. To achieve a better quantitative estimate of the contributions

arising from genotype and condition, we implemented GMM estimation. Besides the fixed and/or

interaction terms (**Methods**), the time term was considered in the model (Figure 3C). We applied

corrected Akaike's Information Criterion (AICc) and Bayesian Information Criterion (BIC) for model

evaluation and Marginal R squared (R2m) for variance coverage. Models built from genotype

demonstrated the best performance, which corroborates the fact that genetic perturbations have

significant impact on cellular differentiation and function. We note that the statistical modeling

implemented in MDL4Cyto (Figure 1B and **Methods**) is broadly applicable, which can facilitate the study

of cell fate decision and cell-type dynamics.

Figure 3. Analysis of dynamic data panel. Progression (%Events over the course of time) **(A)** and
histogram **(B)** of cell count of proliferation marker Ki67 diverged with genetic background and shifted
with condition. We implemented Generalized Mixed-Effects Model (GMM) estimation that considered
the time term in addition to a fixed term and interaction term. While using genotype and condition as
main parameters, several models were constructed, as represented by the equations in **(C)**. The
performance of each model was further evaluated using the corrected Akaike's Information Criterion
(AICc), Bayesian Information Criterion (BIC), and Marginal R squared (R2m). Genotype was identified as
an important feature.

Quantifying cell response and population alteration via unsupervised learning

Identifying cell response from cell populations has been a key interest in mass cytometry

analysis. Hematopoietic responses, induced either by genetic perturbations or by pathogen-like

stimulation (LPS), were analyzed via pairwise comparisons, including WT-PBS vs TKO-PBS, WT-LPS vs

TKO-LPS, WT-PBS vs Tet2-PBS, WT-LPS vs Tet2-LPS. Two established unsupervised learning algorithms, T-

REX and PhenoGraph, were benchmarked in clustering our reference dataset (see Table S3 for the

hyperparameters). Both algorithms apply k-nearest neighbors (kNN) for clustering and Uniform Manifold

Approximation and Projection (UMAP) for dimension reduction and visualization. As illustrated in the

UMAP plot, genetic perturbations induced alterations in cell subpopulations (Figure 4A&4B). Notably,

differential cell count of markers under a genetic perturbation, including Sca1, CD45, TNFα, and Caspase

3, could be seen in the mapping of cell count to the clusters (Figure S2, T-REX algorithm). These results

are consistent with the comparisons from our statistical tests (Figure 2A). Compared to conventional

methods, unsupervised learning showed a unique advantage in efficiently capturing changes in cell

populations. In our framework, cell subpopulations within the clusters were further investigated in two

ways. While reference datasets trained here contained only a single sample for each type, we created

two pseudo-replicates from a single sample (**Methods**) for the PhenoGraph algorithm, thus enabling

differential abundance analysis for the markers within each cluster (Figure 4C). In addition, populations

from the T-REX algorithm were screened and enriched through Density-Based Spatial Clustering and

Application with Noise (DBSCAN) and Marker Enrichment Modeling (MEM)[20] (Figure 4D). Strikingly, in

contrast to the T-REX method, our framework uncovered a strong signal from the cytokines, i.e., IL6 and

TNFα, from the differential abundance analysis (Figure 4C). This observation could well be attributed to

the known LPS-driven inflammation, as demonstrated by the enrichment analysis from MEM (Figure

4D). In addition, B cells and T cells are derived from the myeloid lineage, providing a broad and general

defense against foreign pathogens. Collectively, the TNFα and IL6 differential abundance results are

consistent with myeloid-driven or innate inflammation.

Figure 4. Cell response identified by the unsupervised learning. Clustering from Phenograph **(A)** and T-REX **(B)** algorithms were performed on the paired genetically perturbed samples. PBS vehicle control and LPS stimulation were separately explored and compared. Using both algorithms, we found that the genetic perturbations altered the cell subpopulations. Clusters of significant fold changes identified from PhenoGraph were further explored for differential abundance of markers. Markers with significant responses that were captured in more than two clusters were identified **(C).** On the other hand, cell subpopulations obtained from T-REX were screened for those with significant changes and enriched through DBSCAN and MEM **(D).**

Probing the regulatory cascade with biologically-informed neural networks

The hematopoietic system has classically been represented as a rigid hierarchical tree,

illustrating HSC progression through multipotent, oligopotent, and bipotent stages. Evaluating a

"dynamic" model with complex interactions, we interrogated the marker/cell-type interactions within

the regulatory cascade from supervised learning models, considering both the network architecture and

prediction performance as probes. LPS stimulation impacts all stages of HSC differentiation, thus having

a global impact on all surface markers[18]. Therefore, we assigned PBS vehicle control and LPS stimulation

as predicted labels to the samples (**Methods**). Twenty-two lineage/functional makers from nine

different cell types (Table S1) were encoded as essential numerical features, while genotype was loaded

as an external categorical/global feature.

Considering each node as a marker/cell type, their interactions could be represented as network

connections. Specifically, a linear regression model reflects the additive effect from markers/cell types; a

dense neural network models the complex interactions among markers/cell types; and a tree model

considers the binary relations between levels within the hierarchy (Figure 5A). LPS stimulation on the

cells (the prediction label) could be treated as a perturbation on the neural network. In this model,

HSPCs will be differentiated to the common lymphoid progenitors, unipotent precursor cells as well as

megakaryocyte-erythroid progenitor cells, eventually producing mature cells. Informed by the

differentiation hierarchy, we implemented Boosted Tree (BT) and Random Forest (RF) models along with

two reference models, Linear Regression (LR) and Deep Neural Network (DNN). The four models varied

in performance: Linear Regression (LR) achieved a prediction accuracy of 64.8%; the DNN model with

three-hidden layers enhanced the accuracy to 68.1%; the Boosted Tree (BT) model further increased the

prediction performance to 72.7%; and the Random Forest (RF) models topped the accuracy at 77.13%

(Figure 5B). This overall pattern of prediction accuracy (LR < DNN < BT < RF) suggests a non-linear,

hierarchical relationship among the features, consistent with the classical hematopoietic hierarchical

tree. Interestingly, when we stacked a Deep Neural Network on top of the Random Forest model, where

DNN functioned as a denoising head[21], the new stacked model showed degraded performance in

comparison to the RF model alone.

Figure 5. Mechanistic insights extracted from biologically-informed neural networks. Our neural network designs were guided by the biological context, in which we considered each node as a marker. The marker/cell-type interactions could therefore be represented as a network **(A)**, e.g., a decision tree or neural network could be linked to the regulatory cascade. With PBS vehicle control and LPS stimulation as labels, tree-based models (Boosted Tree [BT] and Random Forest [RF]) showed improved performance compared to Linear Regression (LR) and Deep Neural Network (DNN) models as well as DNN-RF stacked models **(B),** consistent with the hematopoietic tree hierarchy (i.e., representing a sequential binary branching decision process) among the markers. Feature importance of individual markers in the BT model, which implements gain-based feature importance **(C),** and RF model, which implements mean decrease in AUC **(D),** were examined. The group of markers with significant contributions would imply an overall cell response under inflammatory (LPS), e.g., pathogen-induced, challenges.

We further considered the feature contributions of individual markers using gain-based feature importance (Figure 5C) and mean decrease in AUC (Figure 5D). Gain-based feature importance reflects the relative contribution of a given feature to the model. Mean decrease in AUC indicates the decline in performance once the corresponding feature has been removed from the training dataset. We hypothesized that markers/features with the top feature contributions are biologically meaningful.

Indeed, the single most important contribution came from the strongest perturbation – genotype.

Individual markers, including Caspase3, Gr1, IgM, CD11b, and CD25, that contribute substantially to the

prediction of LPS stimulation, are highly biologically relevant in the full differentiation cycle of

hematopoiesis. Caspase 3 is an important player in the regulated cell death. Gr1 and CD11b reflect a

response from myeloid cells. And Ter119 indicates a strong immune response from the erythrocytes

under inflammatory challenges.

Applying machine learning in sample classification

While conventional methods/pipelines are limited by domain knowledge or manual

intervention, a predictive model that bypasses cell clustering or manual gating may provide an end-to-

end approach to the analysis of mass cytometry data. Here, we benchmarked a group of supervised

learning models, including CNN-1D, CNN-2D, RNN-LSTM, and RNN-GRU, that directly handle the raw

cytometry profiles for sample classification, i.e., prediction of clinical or biological outcome (Figure 6A).

We established multi-class output labels corresponding to six different sample types, including WT-PBS,

WT-LPS, Tet2-PBS, Tet2-LPS, TKO-PBS, and TKO-LPS. We used the curated ensemble of (1 hour)

"segments" from the initial bulk samples in evaluating the model performance. We found that CNN and

RNN-derived models demonstrated good prediction performance with at least 95% accuracy (Figure 6B).

We argue that convolutional layers and recurrent cells are efficient in learning the relevant features.

Algorithms such as CNN and LSTM as applied here, with cell count as the trait, captured with greater

fidelity the landscape of marker/cell-type interactions than the tree-based discrete models evaluated

above.

Model	Train Acc	Val Acc	Test Acc
CNN-1D	0.99783	0.97400	0.95887
CNN-2D	0.97580	0.95233	0.96537
RNN-LSTM	0.99819	0.96425	0.97186
RNN-GRU	1.000	0.98375	0.98052

Figure 6. Supervised learning identified sample type with high prediction performance. Convolutional Neural Network (CNN) and Recurrent Neural Network (RNN) derived models (CNN-1D, CNN-2D, RNN-LSTM, RNN-GRU) were benchmarked for the task of sample type classification. The specifics of the architectures are shown in **(A)**. All tested models achieved high performance with over 95% in accuracy **(B),** demonstrating the importance of learning the complex interactions and high-order features.

Discussion

The current single-cell genomic revolution has enabled entire systems, such as the

hematopoietic system, to be interrogated comprehensively at an unprecedented resolution, revealing a

previously unappreciated degree of functional and phenotypic variation within cell compartments. In

this work, bone marrow samples were harvested from mice and further characterized in mass cytometry

– the baseline murine model (WT) was compared to genetic perturbations, including *Tet2*-knockout

(*VavCreTet2*) and *VavCreBaxBakBid* triple-knockout (TKO). LPS served as an external stimulus or an

innate immune response, simulating pathogen infection. Statistical modeling and unsupervised learning

revealed the direct effect of genetic perturbations. Markers of interest that are typically processed

individually could be combined in unsupervised learning models. Importantly, we revealed a close

connection among the intrinsic hierarchy of HSPC differentiation, neural network architecture, and

prediction performance. Specifically, Boosted Tree (BT) and Random Forest (RF), which closely resemble

the stepwise binary branching decisions of the classical hierarchical model of hematopoiesis, showed

superior performance in comparison with Linear Regression (LR), Deep Neural Network (DNN) or even

the DNN-RF stacked model. The prediction performance was substantially enhanced relative to neural

network models, as the model reflected the cell differentiation cascade. However, as recent studies with

functional assays, non-invasive lineage tracing, and omics studies at single-cell resolution have shown[22],

hematopoiesis is a continuous process in which obvious boundaries are lacking between cell populations

located at different levels of the hierarchy. Consequently, given their architecture, tree-based models

may have an inherent limitation in capturing transition states, even with extensive optimization,

constraining their ability to further improve prediction performance. From this perspective, the complex

relationships among the cell populations (including feed-back loops in pathways) may be better

elucidated by neural networks.

Nevertheless, results of the analyses from the machine learning models were consistent with

biological insights such as myeloid-driven or innate inflammation (responses from TNFα and IL6). In

summary, the presented framework provides a unifying methodological framework for the analysis of

cytometry data, combining data-driven and model-driven approaches. We constructed interpretable

models that are anchored in biological context, with implications for ongoing and future studies in

single-cell biology.

Limitations of the study

Machine learning models, including graphical representation learning and tree-based

algorithms, were trained in datasets obtained from mass cytometry (CyTOF) characterization. While we

designed the neural network architectures with biological information as guide, assumptions of cell

types as nodes and edges as their interactions within a graph-based framework are based on the classic

discrete cell differentiation model. Cell differentiation is a continuous process, with intermediate states

among various levels leading to much more complex networks.

Methods

Mass cytometry and murine samples

Three generic murine models were included in this study, namely, wild type (WT), Tet2 knockout

(Tet2)[23], and *VavCreBaxBakBid* triple-knockout (TKO)[14]. Prior to bone marrow isolation, the mice had

been treated intraperitoneally with 1.5 mg/kg LPS (L4391; Millipore-Sigma) or PBS for 8 hours. The

obtained samples for the six genotype-condition types were subsequently characterized using mass

cytometry, according to previously established protocols[24]. Mass cytometry combines inductively

coupled plasma time-of-flight mass spectrometry and cytometry. Specifically, mass cytometry staining

was performed on whole bone marrow with antibodies conjugated to heavy metals. Cell responses from

9 different cell type and functional categories (stem cells, pan leukocyte, T lymphocytes cells, mature

myeloid cells, B cells, erythrocytes, cytokines, cell death, and proliferation) were recapitulated by 16

surface/lineage markers and 6 function/intracellular markers. Markers from pre-gating channels (DNA-1

and DNA-2) and instrument channels (Time/Event) were also considered (Table S1). The CyTOF data

generated from the six samples consisted of a maximum of 293,778 cells and a minimum of 121,124

cells.

Data pre-processing

Initial cell count data profiles obtained from the mass cytometry characterization were filtered

out for debris, doublets, and dead cells, thus reducing noise. They are represented as dynamical panels

on which the *arcsinh* transformation with co-factor = 5 and a binning window of 30 seconds was applied.

We sought to analyze the data in a way that maximizes robustness. The processed profiles were sliced

into ensembles of "segments" to achieve an optimized tradeoff between computational efficiency and

prediction accuracy. We evaluated an array of sizes ([2, 10, 30, 60, 120, 240, 480]), corresponding to

2*30S=1M, 10 * 30S = 5M, 30 * 30S = 15M, 120 * 30S = 1H, 240 * 30S = 2H, & 480 * 30S = 4H sampling

intervals (M: Minute, H: Hour), respectively. The obtained pile of "segments" was completely shuffled

and re-assembled into new ensembles. We implemented an RNN-LSTM model for this special

"hyperparameter search" (see prediction accuracy in Table S2) to determine the optimal segment size,

which was 1H. Signals of select markers, including Sca1, CD45, TNFα, Caspase 3, and Ki67, were

aggregated and averaged into 1 hour binning window for the statistical test and GMM estimation.

Mechanistic Deep Learning Framework

Statistical test and model

Wilcoxon test was performed on pairwise comparisons of marker intensity. Pearson correlation

coefficient was calculated for inter-marker association. We implemented a GMM on the marker Ki67.

We considered fixed effects and interactions among time, genotype, and condition (PBS control or LPS

stimulation). After testing the random slope of a sample, $(1 + \text{Time} \mid \text{Sample})$, we considered only the

random intercept from a sample, i.e., $(1 \mid \text{Sample})$. The models were evaluated using corrected Akaike's

Information Criterion (AICc), Bayesian Information Criterion (BIC), and Marginal R squared (R2M). The

optimal model was found to be given by the following expression:

$$\text{Marker Intensity} = \text{Time} + \text{Genotype} + \text{Condition} + \text{Genotype:Time} + \text{Condition:Time} + (1 \mid \text{Sample}) + \varepsilon$$

Unsupervised Learning models

We analyzed two established unsupervised learning pipelines, namely Phenograph[5] and T-REX[6],

which were applied to the preprocessed CyTOF dataset with binning window 30S. For both pipelines,

Uniform Manifold Approximation and Projection (UMAP)[25] was used for dimensionality reduction; k-

nearest neighbors (kNN)[26] for clustering; Marker Enrichment Modeling (MEM) for generating the

enriched heatmap. The applied hyper-parameters are presented in Table S3. For comparison, we

screened and enriched the cell populations derived from the T-REX algorithm through Density-Based

Spatial Clustering and Application with Noise (DBSCAN) and MEM.

Supervised Learning models

All the machine learning models were trained using the *TensorFlow (TF)* ecosystem[27], which we

applied to the preprocessed CyTOF dataset (with binning window size of 30S) to perform model training

and evaluation. Linear Regression (LR), Deep Neural Network (DNN), Boosted Tree (BT), and Random

Forest (RF) models were implemented on the dataset with train and test split of 8:2. The DNN was

implemented as a multilayer perceptron. A stacked model of Dense Neural Network and Random Forest

(DNN-RF) was constructed following the same splitting. Models implementing a Convolutional Neural

Network (CNN) or a Recurrent Neural Network (RNN) in the framework include CNN-1D, CNN-2D, RNN-

LSTM, and RNN-GRU. For these four architectures, pre-processed datasets were divided into train, valid,

and test data according to 6:2:2 ratio.

We trained the neural networks using mean squared error (as the loss function), *Adam* (an

efficient stochastic optimization method), and Leaky ReLU (as the activation function), building on our

previous work[16,17]. To reduce the possibility of overfitting, we implemented *dropout* assuming a dropout

rate of 20%.

Acknowledgments

This research is supported by National Institutes of Health (NIH) grants NHGRI R35HG010718,

NHGRI R01HG011138, NIA AG068026, and NHLBI R01HL133559.

Author contributions

B.W., S.S.Z., and E.R.G. conceived the project and designed the study. B.W. performed the

analyses and drafted the article. B.W., S.S.Z., and E.R.G. revised the article. S.S.Z. and E.R.G. supervised

the project and acquired funding for the study.

References

1. Davis, M. M., Tato, C. M. & Furman, D. Systems immunology: just getting started. *Nat. Immunol.* **18**,

725–732 (2017).

2. Gadalla, R. *et al.* Validation of CyTOF Against Flow Cytometry for Immunological Studies and

Monitoring of Human Cancer Clinical Trials. *Front. Oncol.* **9**, (2019).

3. Bendall, S. C. *et al.* Single-Cell Mass Cytometry of Differential Immune and Drug Responses Across a

Human Hematopoietic Continuum. *Science* **332**, 687–696 (2011).

4. Hahne, F. *et al.* flowCore: a Bioconductor package for high throughput flow cytometry. *BMC*

Bioinformatics **10**, 106 (2009).

5. Levine, J. H. *et al.* Data-Driven Phenotypic Dissection of AML Reveals Progenitor-like Cells that

Correlate with Prognosis. *Cell* **162**, 184–197 (2015).

6. Barone, S. M. *et al.* Unsupervised machine learning reveals key immune cell subsets in COVID-19,

rhinovirus infection, and cancer therapy. *eLife* **10**, e64653 (2021).

7. Hu, Z., Tang, A., Singh, J., Bhattacharya, S. & Butte, A. J. A robust and interpretable end-to-end deep

learning model for cytometry data. *Proc. Natl. Acad. Sci.* **117**, 21373–21380 (2020).

8. Arvaniti, E. & Claassen, M. Sensitive detection of rare disease-associated cell subsets via

representation learning. *Nat. Commun.* **8**, 14825 (2017).

9. Liggett, L. A. & Sankaran, V. G. Unraveling Hematopoiesis through the Lens of Genomics. *Cell* **182**,

1384–1400 (2020).

10.   Domen, J., Cheshier, S. H. & Weissman, I. L. The role of apoptosis in the regulation of

hematopoietic stem cells: Overexpression of Bcl-2 increases both their number and repopulation

potential. *J. Exp. Med.* **191**, 253–264 (2000).

11.   Kale, J., Osterlund, E. J. & Andrews, D. W. BCL-2 family proteins: changing partners in the dance

towards death. *Cell Death Differ.* **25**, 65–80 (2018).

12.   Salisbury-Ruf, C. T. *et al.* Bid maintains mitochondrial cristae structure and function and protects

against cardiac disease in an integrative genomics study. *eLife* **7**, e40907 (2018).

13.   Gamazon, E. R. *et al.* A gene-based association method for mapping traits using reference

transcriptome data. *Nat. Genet.* **47**, 1091–1098 (2015).

14.   Wagner, P. N. *et al.* Increased Ripk1-mediated bone marrow necroptosis leads to myelodysplasia

and bone marrow failure in mice. *Blood* **133**, 107–120 (2019).

15.   Lun, A. T. L., Richard, A. C. & Marioni, J. C. Testing for differential abundance in mass cytometry

data. *Nat. Methods* **14**, 707–709 (2017).

16.   Wang, B. & Gamazon, E. R. *Towards mechanistic models of mutational effects: Deep Learning on*

Alzheimer's Aβ peptide. 2021.12.19.473403

https://www.biorxiv.org/content/10.1101/2021.12.19.473403v2 (2021)

doi:10.1101/2021.12.19.473403.

17.   Wang, B. & Gamazon, E. R. Modeling mutational effects on biochemical phenotypes using

convolutional neural networks: application to SARS-CoV-2. *iScience* **7**, 104500 (2022).

18.   King, K. Y. & Goodell, M. A. Inflammatory modulation of HSCs: viewing the HSC as a foundation

for the immune response. *Nat. Rev. Immunol.* **11**, 685–692 (2011).

19.   Cai, Z. *et al.* Inhibition of Inflammatory Signaling in Tet2 Mutant Preleukemic Cells Mitigates

Stress-Induced Abnormalities and Clonal Hematopoiesis. *Cell Stem Cell* **23**, 833-849.e5 (2018).

20.   Diggins, K. E., Greenplate, A. R., Leelatian, N., Wogsland, C. E. & Irish, J. M. Characterizing cell

subsets using marker enrichment modeling. *Nat. Methods* **14**, 275–278 (2017).

21.   Xie, J., Xu, L. & Chen, E. Image Denoising and Inpainting with Deep Neural Networks. in *Advances*

in Neural Information Processing Systems vol. 25 (Curran Associates, Inc., 2012).

22.   Zhang, Y., Gao, S., Xia, J. & Liu, F. Hematopoietic Hierarchy – An Updated Roadmap. *Trends Cell*

Biol. **28**, 976–986 (2018).

23.   Moran-Crusio, K. *et al.* Tet2 loss leads to increased hematopoietic stem cell self-renewal and

myeloid transformation. *Cancer Cell* **20**, 11–24 (2011).

24.   Zinkel, S. S. *et al.* Proapoptotic BID is required for myeloid homeostasis and tumor suppression.

Genes Dev. **17**, 229–239 (2003).

25.   McInnes, L., Healy, J. & Melville, J. UMAP: Uniform Manifold Approximation and Projection for

Dimension Reduction. *ArXiv180203426 Cs Stat* (2020).

26.   Mucherino, A., Papajorgji, P. J. & Pardalos, P. M. k-Nearest Neighbor Classification. in *Data*

Mining in Agriculture (eds. Mucherino, A., Papajorgji, P. J. & Pardalos, P. M.) 83–106 (Springer, 2009).

doi:10.1007/978-0-387-88615-2_4.

27.   Abadi, M. *et al.* TensorFlow: Large-Scale Machine Learning on Heterogeneous Distributed

Systems. 19.

Supplementary Information

Figure S1. Summary statistics on the cell count. Initial sample data profiles (WT-PBS, WT-LPS, Tet2-PBS, Tet2-LPS, TKO-PBS, and TKO-LPS) were obtained from mass cytometry characterization, where summary statistics on the transferred cell count were computed. Lineage and functional markers are grouped based on cell type.

Figure S2. Differential cell count illustrated from the unsupervised learning. Markers of interest under genetic perturbation, including Sca1 (LSK cells), CD45 (Leukocyte), TNFα (Cytokines) and Caspase 3 (Cell Death Indicator), were mapped to the clustering results from the T-REX algorithm.

Table S1. Cell markers investigated in this study

Marker Type	Cell Type	Marker Names
Lineage	Stem Cells	CD41, CD48, CD150, cKit, Sca1
Lineage	Pan Leukocyte	CD45
Lineage	T Lymphocytes	CD3c, CD8a, CD4, CD25
Lineage	Mature Myeloid	CD11b, Gr1
Lineage	B Cells	CD43, IgM, CD45R
Lineage	Erythrocytes	Ter119

Functional	Cytokines	TNF alpha, IL2, IL6
Functional	Cell Death	Caspase3, Granzyme B
Functional	Proliferation	Ki67
Pre-gating		DNA1, DNA2
Instrument		Cell Length, Time

495 **Table S2**. Searching for the optimal size "hyperparameter" via RNN-LSTM model architecture

Micro Sample Size	Train Acc	Val Acc	Test Acc
2	0.73653	0.71342	0.71033
10	0.93403	0.88667	0.87375
30	0.99791	0.95369	0.95125
60	**0.99499**	**0.9325**	**0.9575**
120	1.0	0.9598	0.975
240	1.0	0.93	0.93
480	1.0	0.77551	0.88

497 **Table S3.** Hyperparameters applied in PhenoGraph and T-REX

Key components in pipeline	Phenograph	T-REX
UMAP	ret_nn = TRUE	
kNN	k = 30 for single sample, k = 60 for paired sample	