Natriuretic peptides protect against apoptosis and increase cGMP around

cardiomyocyte mitochondria

Gaia Calamera[1], Bernadin Ndongson-Dongmo[1], Dulasi Arunthavarajah[1], Mette Ovesen[1],

Choel Kim[2, 3], Finn Olav Levy[1†], Kjetil Wessel Andressen[1*] and Lise Román Moltzau[1*].

[1]Department of Pharmacology, Institute of Clinical Medicine, University of Oslo and Oslo

University Hospital, P.O. Box 1057 Blindern, 0316 Oslo, Norway

[2]Department of Pharmacology and Chemical Biology, Baylor College of Medicine, Houston,

TX, USA

[3]Verna and Marrs McLean Department of Biochemistry and Molecular Biology, Baylor

College of Medicine, Houston, TX, USA

*shared senior author

† Corresponding Author: Finn Olav Levy, Department of Pharmacology, Faculty of

Medicine, University of Oslo, Sognsvannsveien 20, P.O.Box 1057 Blindern, N-0316 Oslo,

Norway. Tel: +47 22840237. Fax: +47 22840202. E-mail: f.o.levy@medisin.uio.no

Abstract

Natriuretic peptides (NPs) increase cGMP, show beneficial cardiovascular effects and has been

shown to regulate energy metabolism in other tissues. However, little is known about their

direct effect on cardiac mitochondria and cardiomyocyte apoptosis. Here, we examined

whether NPs increase cGMP around mitochondria and alter apoptosis in cardiomyocytes.

Stimulating with ANP or CNP reduced apoptosis, together with reduced caspase 9 activation

and cytochrome c release, suggesting that NPs decrease apoptosis through the intrinsic pathway

that involves mitochondria. We engineered a novel FRET-based biosensor with high selectivity

towards cGMP and targeted this to the outer mitochondrial membrane (OMM) and found that

ANP and CNP increase cGMP at the OMM. Moreover, ANP and CNP increased

phosphorylation of the pro-apoptotic protein Drp1 and CNP prevented fragmentation of

mitochondria. We suggest that cGMP increase in the OMM microdomain inhibits the pro-

apoptotic protein Drp1, leading to reduced mitochondrial fragmentation that inhibits apoptosis.

Introduction

Heart failure is associated with an imbalance in cell energy, changes in the energy source and oxidative stress[1-3]. In addition to providing ATP for the contraction mechanism in cardiomyocytes, mitochondria are also important in regulation of cellular metabolism, cell differentiation, motility, intracellular Ca^{2+} signaling and apoptosis[4,5]. The signaling molecule cGMP has been shown to induce mitochondrial potassium channel opening, mediating a beneficial effect in ischemia-reperfusion injury[6-10]. Natriuretic peptides (NP) increase cGMP by activation of the particulate guanylyl cyclase (pGCs). Atrial (ANP) and brain NP (BNP) both activate the NP receptor GC-A while C-type NP (CNP) activates GC-B. NPs can mediate several beneficial effects in the cardiovascular system such as diuresis, natriuresis, vasodilation and antihypertrophic effects[11]. Consequently, recent attention has been given to drugs that increase NP levels as potential treatment of heart failure. Sacubitril-valsartan, a neprilysin (NEP) inhibitor in combination with an AT_1 angiotensin II receptor antagonist, was shown to be superior to the angiotensin converting enzyme inhibitor enalapril in the treatment of heart failure with reduced ejection fraction (HFrEF)[12]. The role of NPs in heart disease still needs to be better understood to further improve treatments. Natriuretic peptides have been shown to regulate cell metabolism and be beneficial against metabolic disease[13], but little is known about direct effects of NPs on mitochondrial function in the heart.

Hence, our aim was to investigate whether NPs can modulate mitochondrial function in the heart and whether NPs specifically increase cGMP in nanodomains surrounding the mitochondria. In the present study, we show that both CNP and ANP protect from apoptosis, confirmed by reduced Poly (ADP-ribose) polymerase (PARP) cleavage, and we show reduced cytochrome c release from mitochondria and reduced caspase 9 activation, suggesting that the apoptosis involves the mitochondrial pathway. We also show that NPs can increase phosphorylation of Drp1 and CNP can prevent mitochondrial fragmentation, which can mediate an anti-apoptotic effect. Finally, we developed a novel FRET-based biosensor selective for cGMP and confined this to the outer mitochondrial membrane (OMM) that allowed us to detect local cGMP increases from both ANP and CNP in real time in intact cells.

Materials and Methods

Materials. Human C-Type natriuretic peptide (CNP) and rat atrial natriuretic peptide (ANP) were from GenScript Corp. (Piscataway, NJ, USA). Mitotracker Deep Red, Mitotracker Green, Dulbecco's modified Eagle's medium (DMEM) with GlutaMAX, sodium pyruvate and MEM NEAA were from ThermoFisher Scientific. All other chemicals and media were from Sigma-Aldrich (Sigma-Aldrich, St.Louis, MO), unless otherwise specified.

Animals. This investigation was approved by the Norwegian Food Safety Authority (application number 8683 and 23286) and conforms to the research animal Directive 2010/63/EU, the Guide for the Care and Use of Laboratory Animals (NIH publication No. 85-23, revised 2011, US), in accordance with the ARRIVE guidelines[14]. Animals were anaesthetized in 4-5 % isoflurane and anesthesia was confirmed by abolished pain reflexes. Animals were euthanized by cervical dislocation and the chest was then opened to extract the heart.

Isolation of cardiomyocytes. Primary culture of cardiac ventricular myocytes from male Wistar rats were prepared as previously described[15].

LDH measurements. Cell death was measured by the LDH-Glo™ Cytotoxicity Assay (Promega, Mannheim, Germany). Cardiomyocytes cultured for 20 h were stimulated with CNP (1 µM), ANP (1 µM) or vehicle (H_2O) at time 0. Medium was sampled at the indicated time (0, 3, 6, 12 and 24 h) and added to LDH storage buffer (200 mM Tris-HCl pH 7.4, 10% glycerol, 1% BSA). Measurements were performed according to manufacturer's instructions.

ATP assay. Cardiomyocyte viability was measured as changes in cellular ATP levels using the CellTiter-Glo® 2.0 Cell Viability Assay (Promega). Cardiomyocytes cultured for 20 h were stimulated with CNP (1 µM), ANP (1 µM) or vehicle (H_2O). After 3 and 24 h from stimulation, cardiomyocytes were washed with 50 mM Tris-HCl pH 7.4, 150 mM NaCl, 2 mM EDTA and lysed using Solobuffer cell lysis kit (Fabgennix, Frisco, TX 75033, US). Measurements of ATP in the samples were performed according to manufacturer's instructions. Change in cellular ATP content after each stimulation time was calculated as (ATP in treated cells)/(ATP in vehicle-treated).

TUNEL assay. After 20 h from plating, adult cardiomyocytes were stimulated with either CNP (1 µM), ANP (1 µM) or vehicle (H_2O) for 3 h. To detect apoptotic cells, the ApopTag Plus Fluorescein In Situ Apoptosis Detection Kit (Merck, Darmstadt, Germany) was used according to the manufacturer's protocol. DAPI (5µg/ml) was added for nuclear staining. Cells were

visualized under an Olympus BX53 (Olympus, Japan), using an air objective (10x, 0.30NA,

Olympus), and images taken with a XC50 color camera and cellSens Entry Software (Olympus,

Japan). Images were acquired and analysed by a blinded operator.

Immunoblot analysis. Cardiomyocytes cultured for 20 h were stimulated with CNP (1 μM),

ANP (1 μM) or vehicle (H_2O). After 3 h stimulation, protein was extracted on ice using lysis

buffer (Tris 50 mM, EDTA 1 mM, EGTA 1 mM, NaF 50 mM, DTT 1 mM, $Na_4P_2O_7$ 10 mM,

Na_3VO_4 1 mM, Triton X- 100 0.01%, SDS 0.05% and deoxycholat 0.5 %) supplemented with

protease inhibitor cocktail and PMSF (1 mM). Samples were solubilized in loading buffer (62.5

mM Tris-HCl, glycerol 10%, bromophenol blue 1% wt/vol, SDS 2%, mercaptoethanol 2.5%)

and denatured for 5 min at 95°C. Protein samples were separated by SDS-PAGE using 6%,

7.5% or 15% polyacrylamide gels and then transferred to a PVDF membrane. Membranes were

blocked with 3% non-fat dry milk in Tris-buffered saline containing Tween-20 (0.1%) for 2 h

and then incubated with the indicated primary antibodies overnight at 4°C, followed by

incubation with appropriate secondary antibodies for 1 h at room temperature. Drp1 expression

was quantified using a nitrocellulose membrane previously probed with pDrp1 S637. The

membrane was stripped for 10 min in stripping solution (Glycine 0.15% w/v, SDS 0.1% w/v,

Tween-20 1%, pH 2.2), then washed with phosphate-buffered saline and Tris-buffered saline

containing Tween-20 (0.1%). The membrane was then blocked and incubated with Drp1

primary antibody followed by appropriate secondary antibody as described above. Blots were

developed by enhanced chemiluminescence reaction using an Epi Chemi II Darkroom with

Labworks Software (UVP Laboratory Products, Cambridge, UK). Protein expression or protein

phosphorylation was quantified as measured intensity of area under the curve in ImageJ

software. The intensity of each band was normalized to the intensity of the corresponding

loading control (vinculin) except for cleaved PARP that was normalised to total PARP

(cleaved+uncleaved). Antibodies: Vinculin (1:5000), PARP (1:1000), Cleaved PARP (1:800),

Caspase 9 (1:1000), Phospho Drp1 S637 (1:100), Drp1 (1:1000), all from Cell Signaling.

Cytoplasmic cytochrome c release. Adult cardiomyocytes were cultured for 20 h and then

treated with CNP (1 μM), ANP (1 μM) or vehicle (H_2O) for 3 h, washed twice with 50 mM

Tris-HCl pH 7.4, 150 mM NaCl, 2 mM EDTA, incubated on ice with "cytoplasmic extraction

buffer" from Subcellular Protein Fractionation Kit (Thermo Scientific, Waltham, USA) for 15

min, cells were then scraped and samples collected and centrifuged at 4 °C for 5 min at 500 x

g. The supernatant (cytoplasmic fraction) was further centrifuged at 13000 x g for 10 min to

separate the mitochondria (and other organelles). The supernatant was then collected and

cytochrome c measured using cytochrome c ELISA Kit (MyoBiosource, San Diego, USA)

following the manufacturer's instructions.

Construction of FRET-based biosensors for cGMP. The sequences encoding cGi-500V and

ScGI sensors were synthesized by GenScript and inserted into the vector pcDNA3.1(-). cGi-

500V sequence was from *Bos Taurus* PKG I (Gln^{79}-Tyr^{345}) flanked by CFP and Venus. The

ScGI sequence differs only by the mutation Cys158Arg. The OMM-targeting sequence

AKAP1 (1-30), was synthetized by GenScript, opened by *Eco*RI/*Xba*I and inserted N-

terminally into the ScGI sequence preceded by a linker A(EAAAK)^{5}A to separate the fusion

proteins and keep the flexibility[16]. The nucleotide sequences for ScGI has been deposited in

GenBank under the accession number OP221137. ScGI and OMM-ScGI biosensors were then

packaged in adenovirus type 5, amplified and titer measured by VectorBiolabs (Malvern, PA).

Transfection of HEK293 cells and *in vitro* FRET assay. HEK293 cells were cultured in

Dulbecco's modified Eagle's medium (DMEM) with GlutaMAX supplemented with 10% fetal

bovine serum (FBS) and 100 U penicillin/streptomycin at 37ºC in a humidified atmosphere of

5% CO_2 in air. Cells were transiently transfected with the indicated plasmids using

LipofectAMINE 2000 (ThermoFischer) according to the manufacturer's protocol. After

transfection, cells were cultured for 48 h, then homogenized with Ultra-Turrax (Franke &

Kunkel KG, Staufen, Germany) for 30 s at 4°C in Buffer A (HEPES 10 mM pH 7.3, NaCl 137

mM, KCl 5.4 mM, $CaCl_2$ 2 mM, $MgCl_2$ 1 mM). Homogenates were stimulated with increasing

concentrations of cGMP or/and cAMP in the presence of IBMX (100 μM). The donor

fluorophore CFP was excited at 430 nm (430/24 filter) and the fluorescence emission (F) of

Venus and CFP were measured (535/25 and 470/24 nm, respectively) in an EnVision plate

reader (PerkinElmer, USA, NY, Melville). Venus emission was corrected for spillover of CFP

emission into the Venus channel (72.3%). The FRET response was calculated as changes in

F_{Venus}/F_{CFP}. Affinities (pEC_{50}) were calculated by GraphPad Prism 9 for Windows (GraphPad

Software, San Diego, CA) where the lowest FRET was set as the highest concentration of

cGMP.

Confocal microscopy. *Evaluation of the FRET biosensors localization at mitochondria:* Rat

heart derived H9c2 cardiac myoblasts (Sigma-Aldrich, St.Louis, MO) were cultured in

Dulbecco's modified Eagle's medium (DMEM) with GlutaMAX supplemented with 10% fetal

bovine serum (FBS), 100 U penicillin/streptomycin, 100 mM sodium pyruvate and 100 mM

MEM NEAA at 37ºC in a humidified 5% CO_2–95% air atmosphere. Cells were transiently

transfected with the indicated plasmids using LipofectAMINE 2000 (ThermoFisher) according to the manufacturer's protocol and cultured on poly-lysin coated glass cover-slides. Rat adult cardiomyocytes were plated in laminin coated cover-slides and transduced with adenovirus containing the biosensor as previously described[17,18]. After 48 h transfection (H9c2) and transduction (cardiomyocytes), cells were incubated with 50 nM Mitotracker Deep Red (ThermoFisher) for 30 min, washed and incubated in Buffer B (NaCl 144 mM, KCl 1.97 mM, $CaCl_2$ 1 mM, $MgCl_2$ 1 mM, KH_2PO_4 0.43 mM, K_2HPO_4 1.5 mM, glucose 10 mM) to be visualized under an Olympus FV1000/BX61 confocal microscope using a water immersion objective (60x 1.1 NA). Cells were sequentially excited with a 488 nm laser for CFP and a 633 nm laser for the Mitotracker Deep Red and the emission was measured by Olympus Fluoview (Olympus, Tokyo, Japan) at 510/25 nm for CFP and 664/20 nm for Mitotracker Deep Red.

Evaluation of mitochondria morphology: H9c2 cells were cultured on poly-lysin coated glass cover-slides and after 24 h stimulated as indicated and incubated with 50 nM Mitotracker Green for 15 min, washed and incubated in Buffer B. Cells were visualized under an Olympus FV1000/BX61 confocal microscope using a water immersion objective (60x 1.1 NA). Cells were excited with a 488 nm laser and the emission was measured by Olympus Fluoview 1000 at 510/25 nm. Images were analysed by Mito-Morphology ImageJ macro[19], which was modified for this measurements to apply a Gaussian filter. In brief, the macro extracts the green channel (stained with Mitotracker Green) to grayscale, inverts to a binary mask and thresholds to resolve mitochondria; these are analysed as particles and different morphology parameters are generated. Among them the ratio between the major and minor axis for each mitochondria was employed as index of mitochondria elongation. For each cell the median of these values was calculated.

FRET imaging. H9c2 cells and cardiomyocytes expressing the indicated biosensors (as described above) were placed into a watertight cell imaging chamber (Attofluor, ThermoFisher) at room temperature in Buffer B. Cells were visualized through a motorized digital inverted fluorescent microscope (iMIC; FEI, Graefelfing, Germany) with an oil objective (40x, 1.35 NA; Olympus). Fluorescent molecules were illuminated by a monochromator with fast-switching wavelengths (Polychrome V; FEI) for 20-80 ms at a frequency of 0.1-1 Hz. Cells were excited at 436±10 nm for 20-80 ms and emission was split using a dichrotome iMIC Dual Emission Module with a DCLP 505 filter that separates the two fluorophores onto a camera (EVOLVE 512; Photometrics, Tucson, USA). Fluorescence was measured at 530±15 nm for Venus and at 470±12 nm for CFP emission. Images were acquired by Live Acquisition browser

(FEI) and FRET was calculated using Offline Analysis software (FEI). FRET was measured

as F_{Venus}/F_{CFP}. Venus emission was corrected for spillover of CFP emission into the Venus

channel (63%). FRET was adjusted for photobleaching. In a subset of experiments, cells were

visualized through a motorized digital inverted microscope (IX81; Olympus) with an oil

objective (40x, 1.35 NA; Olympus). Fluorescent molecules were excited at 436nm for 8-20ms

(CoolLed pE-4000) and emission split with a 495 nm filter using an Optosplit II (Cairn

research, Kent, UK) separating two fluorophores onto a camera (Prime 95B; Photometrics).

Fluorescence was measured at 535±15 nm for Venus and at 470±12 nm for CFP emission.

Images were acquired by Micro-Manager. FRET was measured as F_{Venus}/F_{CFP} through custom-

built scripts in ImageJ. Representative experiments were run through a Savitzky–Golay filter

using GraphPad Prism 9 for Windows for clarity.

Total cGMP measurements. H9c2 cells and isolated ventricular cardiomyocytes plated for

20h were stimulated with ANP (1µM) or CNP (1µM) for 10 min, 30 min, 1, 3, 6, 12 and 24 h.

Cyclic GMP levels were measured using a cGMP enzyme immunoassay kit (Cayman Chemical

Company, Ann Arbor, MI, USA) as previously described [20].

Statistics. Statistical significance was determined by one-way ANOVA with Bonferroni's,

Dunett's, Tukey's and Sidak's multiple comparisons test, two-way ANOVA with Sidak's

multiple comparisons test as indicated using GraphPad Prism 9 for Windows. $P<0.05$ was

considered statistically significant.

Results

Natriuretic peptides increase cell viability

In order to study the effects of NPs on cell survival, we first measured the degree of cell viability of cardiomyocytes cultured for 20 h in a medium deprived of insulin and triiodothyronine (T3) to induce stress to the cells. During cell death, LDH is released to the medium, and this was measured over the next 24 h and progressively showed a time-dependent increase in cell death (Figure 1a). To determine if NPs could modify cell viability, we stimulated cardiomyocytes with CNP (1μM) or ANP (1μM) and found that both natriuretic peptides significantly reduced LDH levels in the medium compared to vehicle over time (Figure 1a). This indicates a protective effect of CNP and ANP against cardiomyocyte death. Further, we assessed the production of ATP as an indicator of metabolically active cells. A 3 h stimulation with CNP or ANP increased ATP levels (CNP:1.6 ± 0.1 and ANP:1.8 ± 0.2 fold increase *vs.* control). In addition, 24 h stimulation with either CNP or ANP also increased ATP levels (CNP:1.4 ± 0.1 and ANP:1.4 ± 0.1 fold *vs.* control; Figure 1b) suggesting that NPs increase cell viability, in line with our results on LDH measurements. To determine whether NPs increase cGMP over such prolonged periods (24h), cardiomyocytes were stimulated with CNP or ANP for up to 24 h and we found that CNP-stimulation increased cGMP over the entire 24 h showing peak levels around 1 h, while ANP stimulation increased cGMP with a peak around 30 min and gave an earlier decrease in cGMP levels than CNP (Supplementary Fig. 1). These data suggest that cGMP increases are sustained and can mediate downstream effects on cell viability during prolonged NP stimulation.

NPs protect against apoptosis through the mitochondrial intrinsic pathway

LDH release from damaged cells could result from either necrosis or apoptosis. We therefore wanted to explore if NPs could alter apoptosis and performed TUNEL assay on adult cardiomyocytes. Stimulating cardiomyocytes for 3 h with either CNP or ANP reduced the number of apoptotic cells ($76.1\pm5.3\%$ and $65.9\pm8.3\%$ compared to control, respectively; Figure 2a). To verify the effects of NPs on apoptosis, we measured the cleavage of PARP, which will prevent PARP from repairing DNA damage and therefore induce apoptosis. Stimulation with CNP or ANP for 3 h significantly decreased PARP cleavage (Figure 2b). Taken together, these data suggest that both CNP and ANP reduce apoptotic cell death.

To investigate if the NP-mediated protective effect against apoptosis was through the intrinsic apoptotic pathway, we measured the activation of caspase 9, which depends on cytochrome c release from the mitochondria. CNP- and ANP-stimulation for 3 h significantly reduced the levels of cleaved caspase 9 compared to control (Figure 2c). In addition, CNP and ANP significantly reduced levels of cytochrome c in the cytosol compared to the control (82.7% ± 5.3 (CNP) and 76.7% $\pm$ 8.3 (ANP) of control, Figure 2d). Together, these data show that both NPs prevent cardiomyocyte apoptosis through the intrinsic mitochondrial pathway.

Development of selective FRET-based cGMP biosensors targeted to mitochondria

Since NPs reduced mitochondria-induced apoptosis, we wanted to explore if NPs increased cGMP levels around mitochondria.

We have previously shown that cGMP increases in different cellular microdomains by NP-stimulation, using both untargeted and localized FRET-based biosensors in adult cardiomyocytes [21,22] and others have shown that stimulation of soluble guanylyl cyclase (sGC) increases cGMP at the OMM in neonatal cardiomyocytes using an OMM-localized FRET-based biosensor[23]. To investigate whether NPs increase cGMP at the mitochondria, we developed FRET-based cGMP biosensors that measure cGMP in the cytosol and at the outer mitochondrial membrane. We first constructed a potentially brighter variant of the previously published cGi-500 biosensor[24] by replacing the acceptor fluorophore with Venus and termed this new biosensor cGi-500V (Figure 3a). We then expressed the biosensor in HEK293 cells and determined its affinity for cGMP by incubating cell homogenates with increasing concentrations of cGMP. The resulting biosensor had an affinity (EC_{50}) of 510 ± 42 nM (Figure 3a) for cGMP with a maximal change in FRET of 53.2 ± 1.4 % (Figure 3d).

Cyclic AMP is structurally similar to cGMP and many biosensors only display moderate selectivity between these nucleotides[21,25]. Since cGMP can directly and indirectly modulate cellular concentrations of cAMP through inhibition or activation of certain PDEs[26,27], biosensors with low cyclic nucleotide selectivity can indiscriminately bind either cyclic nucleotides. It is therefore essential to engineer a biosensor with high cGMP selectivity. First we measured FRET of cGi-500V in the presence of cAMP. FRET *increased* by 23.7 ± 2.4 % at low cAMP concentrations (EC_{50High} 12.1 ± 1.8 nM), while FRET decreased at higher concentrations of cAMP (EC_{50Low} 106 ± 28 μM) (Figure 3a and Table 1) demonstrating that cAMP can bind cGi-500V and low level of cAMP can interfere with the FRET responses for

cGMP. cGi-500V is based on the tandem cyclic nucleotide binding (CNB) domains of cGMP-dependent protein kinase Iα with different cyclic nucleotide affinities. Only the C-terminal CNB-domain (CNB-B) is selective for cGMP and its co-crystal structure with cGMP showed that an arginine residue at β5 is responsible for the enzyme selectivity[28]. The N-terminal CNB domain (CNB-A) binds both cGMP and cAMP with similarly high affinities with little cyclic nucleotide selectivity. It contains a cysteine at the corresponding position in β5 instead of arginine[28]. We therefore mutated this cysteine (C158) in the CNBD-A to an arginine and termed the mutant biosensor ScGI (Selective cyclic GMP kinase I sensor). Cyclic AMP did not show high affinity increase in FRET (as that obtained with the cGi-500V) and only a reduction of FRET was observed for the higher cAMP concentrations (EC$_{50}$ of 125±23 µM; Figure 3b and Table 1). The ScGI biosensor also displayed a ~ 2 fold *increased* affinity for cGMP (216±18 nM) compared to the cGi-500V resulting in ~500 fold cGMP selectivity over cAMP. However, the mutation caused a reduction in the maximal FRET observed (34.0±3.6% *vs.* 53.2±1.4% for the cGi-500V, Figure 3d).

Since the ScGI with Cys158Arg no longer responded to the high affinity cAMP binding, and the affinity for cGMP closely matched previously published affinity for the isolated CNB-B[28], we hypothesized that the cGMP-induced FRET change in both cGI-500V and ScGI was only due to binding of cGMP to the CNBD-B. To test this, we determined if the presence of cAMP influenced cGMP-binding to the cGI-500V and ScGI biosensors in an *in vitro* competition-binding assay. Adding a low concentration (1µM) of cAMP increased FRET of the cGI-500V by 25.0±1.7 %, but did not alter the affinity for cGMP (EC$_{50}$ of 321±61 nM; Table 1; Supplementary Fig. 2). Adding a high concentration of cAMP (100µM) slightly reduced FRET (9.2±2.1 % reduction) and significantly decreased the potency of cGMP by ~ 2 fold (EC$_{50}$ 1173±202 nM) (Supplementary Fig. 2). For the ScGI biosensor, 1 µM cAMP did not affect the FRET response (7.3±3.6% increase, p=0.08) and did not modify the potency of cGMP (EC$_{50}$ 294±116 nM). Adding 100µM cAMP significantly reduced FRET (37.2±7.3 % reduction) and reduced the cGMP potency by ~ 3 fold (EC$_{50}$ 683±174 nM) (Supplementary Fig. 2). Taken together, these data suggest that cAMP at low concentrations binds to the CNBD-A of cGi-500V, which is not responsible for determining the cGMP affinity but rather induces a conformational change of the biosensor and therefore increases FRET. Cyclic AMP at high concentrations, on the other hand, competes with cGMP for binding to the CNB-B in both cGi-500V and ScGI or the Cys158Arg mutation also alters CNB-A affinity for cGMP. In addition,

this confirms that the mutation in the CNB-A of the ScGI prevents the high affinity binding of
cAMP displayed by the cGi-500V.

Next, we constructed a mitochondria-localized ScGI biosensor using the N-terminal
targeting sequence of AKAP1, an anchoring protein known to be localized at the OMM and
directing PKA signaling[29-31]. This biosensor was termed OMM-ScGI and displayed an affinity
for cGMP similar to the untargeted ScGI biosensor (EC_{50} 225.7±32.3 nM) with ~200 fold
selectivity *vs*. cAMP (EC_{50} 52±11 µM) (Figure 3c and Table 1). However, adding this tag to
the ScGI biosensor decreased the maximal change in FRET significantly compared to the
untargeted sensor (14.6±0.4%, Figure 3d).

Real time measurement of cGMP around mitochondria

To determine if the OMM-ScGI biosensor displayed mitochondrial targeting, we expressed the
ScGI and OMM-ScGI in H9c2 cells and in adult rat ventricular cardiomyocytes and determined
if these co-stained with the mitochondrial marker Mitotracker Deep Red. The OMM-ScGI
biosensor co-localized with Mitotracker Deep Red while the untargeted biosensor did not
(Figure 4a; Supplementary Fig. 3a), suggesting that the OMM-ScGI localizes to the
mitochondria. To assess if NPs increased cGMP around this localized biosensor, we stimulated
single cells expressing either the ScGI or the OMM-ScGI biosensors with CNP (300 nM)
followed by the non-selective PDE inhibitor IBMX (100 µM) to obtain maximal biosensor
activation and measured FRET, reflecting cGMP levels (Figure 4b and d; Supplementary Fig.
3b and d). Stimulating ScGI-expressing cells with CNP decreased FRET (increase in cGMP)
by 90.5±2.0% of max in H9c2 and by 68.8±3.1% of max in cardiomyocytes, while in OMM-
ScGI expressing cells, CNP decreased FRET by 81.1±4.3% of maximum in H9c2 and by
66.6±4.3% of maximum in cardiomyocytes (Figure 4 b-d; Supplementary Fig. 3b-d). This
suggests that GC-B stimulation causes an increase in cGMP in the cytosol, as previously
shown[21,22,32] and at the OMM; moreover IBMX gave a further increase in cGMP in both the
cytosol and at the OMM, indicating that PDEs regulate these pools of cGMP (Figure 4d;
Supplementary Fig. 3d).

Considering that both CNP and ANP protected against apoptosis, we also wanted to
measure the ANP-stimulated cGMP pool. Stimulation of ScGI-expressing cells with ANP
(1µM) gave a FRET response of 2.4±1.5% of maximum in H9c2 and of 2.2±1.3% of max in
cardiomyocytes (right panels in Figure 4b and Supplementary Fig. 3b). Further stimulation

with IBMX gave a FRET response of 5.1±1.1 % in H9c2 and 29.2±3.8% in cardiomyocytes

(Figure 4d and Supplementary Fig. 3d). Stimulation of OMM-ScGI-expressing cells with ANP

generated a larger response compared to the untargeted sensor, with 21.1±6.7 % of maximum

in H9c2 and 15.2±1% of maximum in cardiomyocytes (right panels in Figure 4c and

Supplementary Fig. 3c). IBMX gave a further FRET response of 16.1± 2.4% in H9c2 and

39.1±2.9% in cardiomyocytes (Figure 4d and Supplementary Fig. 3d). Both biosensors

displayed an ANP-induced cGMP increase significantly lower than that induced by CNP, but

ANP at the OMM gave a significantly larger response than in the cytosol in adult

cardiomyocytes. These data taken together suggest that CNP induces a greater cGMP pool both

in the cytosol and at the OMM, while ANP induces a smaller cGMP pool, which instead is

restricted to the OMM in adult cardiomyocytes.

NPs modulate Drp1 phosphorylation and prevent mitochondrial fragmentation

Mitochondrial function is modulated by phosphorylation/dephosphorylation of several proteins

involved in mitochondrial dynamics. Among these, phosphorylation of Drp1 at Ser637 was

found to protect against apoptosis and mitochondrial fragmentation [33-35]. We therefore

investigated whether NPs modulate phosphorylation of Drp1 in adult cardiomyocytes and

found that both CNP and ANP increased phosphorylation at Ser637, with CNP giving a

significantly greater phosphorylation compared to ANP (Figure 5a) without affecting the total

protein expression (Supplementary Fig. 4). Therefore we tested whether NPs can prevent

mitochondrial fragmentation in H9c2 cells undergoing H_2O_2-mediated apoptosis. Firstly, we

measured CNP- and ANP-mediated cGMP increase in H9c2 cells over 24 h and found that

cGMP levels reach a peak already at 10 min and with an increase above basal maintained

troughout 12 h for CNP and ANP (Supplementary Fig. 5). Therefore, we stimulated H9c2

cells with H_2O_2 (100µM) for 3h and showed that mitochondria are significantly fragmented

compared to control cells (Figure 5b) and that concomitant CNP stimulation for 3 h

significantly inhibited mitochondrial fragmentation induced by H_2O_2 (Figure 5b). ANP, on the

other hand, did not modify H_2O_2-induced mitochondrial fragmentation (Figure 5b), even

though the proportion of cells with elongated mitochondria was slightly increased (30.4% with

H_2O_2 + ANP *vs* 18.5% with H_2O_2; Table 2; not significantly different). All together, these data

suggest that NPs prevent cell apoptosis through the instrinsic mitochondrial pathway, possibly

involving Drp1 phosphorylation and reduced mitochondrial fragmentation.

Discussion

In this study, we show that the natriuretic peptides CNP and ANP protect against apoptosis of cardiomyocytes, with reduced PARP cleavage and increase in cell viability. Moreover, we show that the natriuretic peptides reduce both activation of caspase 9 and release of cytochrome c from mitochondria to the cytosol, suggesting that the protective effects involve the mitochondrial intrinsic pathway. Further, by constructing a selective biosensor and targeting this to the OMM, we also show that ANP and CNP increase cGMP at the outer mitochondrial membrane in cardiomyocytes. In addition, both CNP and ANP increase Drp1 phosphorylation, with CNP also preventing mitochondrial fragmentation, likely protecting mitochondria from apoptosis.

Natriuretic peptide effects on apoptosis

Here we demonstrate increased survival of adult cardiomyocytes and protection from apoptosis by NP stimulation. Previous studies determining the effects of NPs on regulation of apoptosis show conflicting results: some studies found that stimulating GC-A and/or GC-B is protective against apoptosis, mostly in other cell types[36-39], but also in cardiomyocytes and the heart[40,41]. However, these studies used neonatal cardiomyocytes, which have a lower degree of differentiation, and focussed on ANP signalling specifically at the nuclei. Others showed an increase in apoptosis by NPs in cardiomyocytes[42-45] and in other tissues[46]. However, the different results between the studies showing increased apoptosis and the data presented in this study could potentially be due to study of adult (this study) *vs.* neonatal cardiomyocytes[42,44,45] or normal (this study) *vs.* a heart failure model[43].

NPs/cGMP signalling involved in mitochondrial function

Our results show that ANP and CNP increase cGMP at the OMM. Cyclic GMP has previously been found in cardiac mitochondria, where mtNOS and sGC have been reported to be located in the matrix[47-49] and sGC-generated cGMP has also been found at the OMM [23]. However, those studies did not examine natriuretic peptide signalling.

The exact mechanism of how NPs regulate mitochondrial function and prevent apoptosis remains elusive. Some studies show that NPs promote mitochondrial biogenesis by upregulation of PGC-1α and PPARδ that could protect cardiomyocytes against mitochondria-

mediated apoptosis[50,51]. Moreover, PKG I seems to protect against apoptosis by increasing

expression of Bcl-2 and decreasing Bax in adult rat cardiomyocytes subjected to

ischemia/reoxygenation stress[52]. Further, ANP- and BNP-stimulation has been shown to induce

mitoK$_{ATP}$ opening and prevent mitochondrial permeability transition pore opening, where both

mechanisms are protective against apoptosis[41,53-55]. A third pathway for NP regulation of

mitochondrial function could be through regulation of Drp1. It has earlier been shown that

PKA can phosphorylate Drp1 at S637, reducing its activity [33], and this has been associated with

prevention of apoptosis [34,35]. PKG-mediated phosphorylation of Drp1 and its protective effect

has been shown via the NO/cGMP pathway in myogenic precursor cells [56]. Thus, there are

several possible pathways for how natriuretic peptides can regulate apoptosis, involving either

gene regulation or direct regulation of apoptotic pathways. In the present study, we show that

NPs increase phosphorylation of Drp1 at Ser637. This could prevent Drp1 activation and

initiation of mitochondrial fragmentation, as shown by CNP stimulation (Figure 5b), and thus

possibly cause prevention of apoptosis[57]. However, we cannot rule out that other pathways are

also involved. Our results show comparable effects of CNP and ANP on LDH activity in the

medium, ATP production, PARP cleavage, cleaved caspase 9 and cytochrome c release in

addition to the prevention of apoptosis. However, compared to CNP, ANP gave significantly

lower cGMP levels at the OMM and also lower Drp1 phosphorylation that did not translate

into reduced H$_2$O$_2$-induced mitochondrial fragmentation. This might indicate that different

pathways are involved in the anti-apoptotic effects seen after ANP and CNP stimulation.

Alternatively, the magnitude of Drp1 phosphorylation seen after ANP stimulation is sufficient

to inhibit the mitochondrial intrinsic pathway.

FRET-based biosensors detect cGMP increase at the mitochondrial outer membrane

The larger phosphorylation of Drp1 and reduced mitochondria fragmentation by CNP,

compared to ANP is in line with the differential cGMP levels measured by FRET biosensor at

the OMM (Figure 4 and 5). There are some indications for the presence and effect of cGMP

inside the mitochondrial matrix and at the OMM after stimulation of sGC[23,41,47-49,53-55].

However, the role of NP-stimulated cGMP at the OMM shown herein has not been examined

previously. More is known about the role of cAMP in the mitochondria, which was found both

in the matrix[58,59] and at the OMM, where PKA and PDE2A activity were also present [35,60]. In

these studies a mitochondrial-localized FRET-based biosensor for cAMP was used[35,59], and

recently also a FRET-based biosensor for cGMP localized at the OMM was developed [23].

However, that biosensor used the cGi-500 biosensor which did not show high selectivity *vs.*

cAMP in our hands (Figure 3; Table 1). Therefore, we constructed cGMP-selective biosensors

localized to the OMM to investigate the presence of cGMP around mitochondria.

Several FRET-based biosensors have been developed with different fluorescent

proteins, cGMP binding domains and affinity/selectivity profiles. Among the most sensitive

biosensors, we find the red-cGES-DE5, Yellow *Pf*PKG and Red *Pf*PKG biosensors[21,32], with

cGMP affinity in the nM range, and the cGi-500 biosensor[24], which is >10 times less sensitive.

The untargeted ScGI and the OMM-ScGI biosensors constructed in this study presented a mid-

range affinity with an EC_{50} of around 200 nM, and were shown to be sensitive enough to detect

similar increases of cGMP by CNP stimulation compared to the other biosensors (Figure 4d,

Supplementary Fig. 3d). ANP stimulation, however, gave a significant increase in cGMP only

at the OMM and not with the untargeted biosensor in cardiomyocytes (Figure 4d), indicating

that GC-A stimulation locally increases cGMP around the OMM. CNP has previously been

shown to stimulate GC-B both in the crest and T-tubules and produce cGMP that reaches

remote targets in cardiomyocytes [22,61], therefore the cGMP increase at the OMM could reflect

the widespread CNP-induced cGMP increase. ANP, on the other hand, has been reported to

produce a small cGMP pool confined near the plasma membrane by activation of GC-A only

in the T-tubules of cardiomyocytes [61]. Such local cGMP increase might not be detected by an

untargeted biosensor due to widespread distribution and local signals being "diluted" across

the entire cell. In fact, even using the ~10 times more sensitive untargeted Yellow *Pf*PKG

biosensor we could only detect a modest cGMP increase after GC-A stimulation[21]. Since ANP

increases cGMP considerably more at the OMM nanodomain compared to that detected by the

untargeted biosensors in cardiomyocytes (Figure 4d), the GC-A receptor and therefore its

cGMP pool at the T-tubules might be in close vicinity of the OMM and form a nanodomain.

Given that mitochondria are tightly distributed between the sarcomeres but also in the sub-

sarcolemmal region[62], either the membranal localization of the GCs or the different ability of

cGMP from the GC-A and GC-B to reach different targets, could help explain the higher levels

of cGMP detected in the OMM by CNP compared to ANP.

CNB domains *responsible for inducing changes in FRET in the cGI biosensors*

Given that cAMP is structurally similar to cGMP and that basal cAMP concentration in

cardiomyocytes are higher (1µM) than cGMP (10nM)[32,63,64], it is important to determine the

selectivity of cGMP biosensors and to ensure that the response is representative of increases in

cGMP and not cAMP. In our hands, the cGi-500V showed a poor selectivity profile, where

cAMP bound to the biosensor at low concentrations resulting in *increased* FRET (Figure 3a).

However, these concentrations of cAMP did not compete with the ability of cGMP to alter

FRET (Supplementary Fig. 2a). In addition, when mutating the CNB-A domain (in the ScGI),

the increase in FRET by low cAMP concentrations was prevented, whereas the cGMP potency

was only slightly *increased* (Figure 3b and Supplementary Fig. 2b). It has previously been

shown that the CNB-A domain displays similar high affinity for both cGMP and cAMP[65],

whereas the CNB-B domain has a slightly lower affinity for cGMP (~215nM), but the affinity

for cAMP is much lower (EC_{50}=52µM) and the CNB-B domain is therefore responsible for the

cGMP selectivity of PKG I[28]. Considering that the ScGI has similar affinity for cGMP as the

isolated CNB-B domain, this domain could also be solely responsible for the biosensors

sensitivity or both CNB domains are responsible, but the response is indistinguishable as both

domains now binds cGMP with similar affinity. Taken together, this indicates that 1) the FRET

increase seen with low concentrations of cAMP (in the cGi-500V) is possibly due to non-

selective cAMP binding to the CNB-A domain, 2) cAMP at high concentrations binds to the

CNB-B domain of cGi-500V and ScGI that reduces FRET, and 3) either cGMP binding to the

CNB-A does not alter the conformation of PKG to trigger a change in FRET, or cGMP binds

to both the CNB-A and –B domains with similar affinity to trigger an increase in the distance

between the CFP and Venus, thus reducing FRET. The ScGI biosensor will thus be a valuable

tool for determining cGMP, particularly in complex biological systems where cAMP can act

as a confounding factor.

Conclusion

489   In conclusion, we present evidence for a protective effect of natriuretic peptides against

490   cardiomyocyte apoptosis, mediated by the intrinsic apoptotic pathway, which also involves

491   phosphorylation of Drp1 and reduced mitochondrial fragmentation. Further, we constructed a

492   novel FRET-based biosensor with high cGMP selectivity in order to detect cGMP at the

493   mitochondria and show that both CNP and ANP increase cGMP at the OMM.

Acknowledgements

496   The authors wish to thank Iwona Gutowska Schiander (University of Oslo) and Soheil Naderi

497   (Oslo University Hospital) for excellent technical assistance.

Funding

This work was supported by the South-Eastern Norway Regional Health Authority (grant # 2015036 and 2019051), the Norwegian Council on Cardiovascular Diseases, The Research Council of Norway (grant # 205167 and 303490), The Anders Jahre Foundation for the Promotion of Science, The Family Blix Foundation, The Simon Fougner Hartmann Family Foundation, and grants from the University of Oslo

Ethics declarations

The authors declare no competing interests.

Author Contribution

GC, BDN, CK, KWA and LRM designed research; GC, BDN, DA and MO performed research. GC, BDN, DA, MO, FOL, KWA and LRM analyzed data; GC, KWA and LRM wrote the manuscript. All authors revised and approved the manuscript.

Data availability

Nucleotide sequence for ScGI and OMM-ScGI have been deposited in GenBank. The data that support the findings of this study are available from the corresponding author upon request. The authors declare that the data supporting the findings of this study are available within the paper and its supplementary information files.

References

Munzel, T. *et al.* Impact of Oxidative Stress on the Heart and Vasculature: Part 2 of a 3-Part Series. *J. Am. Coll. Cardiol.* **70**, 212-229, doi:10.1016/j.jacc.2017.05.035 (2017).

Stanley, W. C., Recchia, F. A. & Lopaschuk, G. D. Myocardial Substrate Metabolism in the Normal and Failing Heart. *Physiol. Rev.* **85**, 1093-1129, doi:10.1152/physrev.00006.2004 (2005).

Bertero, E. & Maack, C. Metabolic remodelling in heart failure. *Nat. Rev. Cardiol.* **15**, 457-470, doi:10.1038/s41569-018-0044-6 (2018).

Braschi, E. & McBride, H. M. Mitochondria and the culture of the Borg: understanding the integration of mitochondrial function within the reticulum, the cell, and the organism. *Bioessays* **32**, 958-966, doi:10.1002/bies.201000073 (2010).

Kasahara, A. & Scorrano, L. Mitochondria: from cell death executioners to regulators of cell differentiation. *Trends Cell Biol.* **24**, 761-770, doi:10.1016/j.tcb.2014.08.005 (2014).

Lee, K. H. *et al.* Cardioprotective effects of PKG activation by soluble GC activator, BAY 60-2770, in ischemia-reperfusion-injured rat hearts. *PLoS One* **12**, e0180207, doi:10.1371/journal.pone.0180207 (2017).

Frankenreiter, S. *et al.* cGMP-Elevating Compounds and Ischemic Conditioning Provide Cardioprotection Against Ischemia and Reperfusion Injury via Cardiomyocyte-Specific BK Channels. *Circulation* **136**, 2337-2355, doi:doi:10.1161/CIRCULATIONAHA.117.028723 (2017).

Bork, N. I. *et al.* Rise of cGMP by partial phosphodiesterase-3A degradation enhances cardioprotection during hypoxia. *Redox Biology* **48**, 102179, doi:https://doi.org/10.1016/j.redox.2021.102179 (2021).

Behmenburg, F. *et al.* Impact of Mitochondrial Ca^{2+}-Sensitive Potassium (mBKCa) Channels in Sildenafil-Induced Cardioprotection in Rats. *PLoS One* **10**, e0144737, doi:10.1371/journal.pone.0144737 (2015).

Frankenreiter, S. *et al.* Cardioprotection by ischemic postconditioning and cGMP-elevating agents involves cardiomyocyte nitric oxide-sensitive guanylyl cyclase. *Cardiovasc. Res.* (2018).

Volpe, M., Rubattu, S. & Burnett, J., Jr. Natriuretic peptides in cardiovascular diseases: current use and perspectives. *Eur. Heart J.* **35**, 419-425, doi:10.1093/eurheartj/eht466 (2014).

McMurray, J. J. *et al.* Angiotensin-neprilysin inhibition versus enalapril in heart failure. *N. Engl. J. Med.* **371**, 993-1004, doi:10.1056/NEJMoa1409077 (2014).

Miyashita, K. *et al.* Natriuretic Peptides/cGMP/cGMP-Dependent Protein Kinase Cascades Promote Muscle Mitochondrial Biogenesis and Prevent Obesity. *Diabetes* **58**, 2880-2892, doi:10.2337/db09-0393 (2009).

551	14	Kilkenny, C., Browne, W. J., Cuthill, I. C., Emerson, M. & Altman, D. G. Improving bioscience
552		research reporting: the ARRIVE guidelines for reporting animal research. *PLoS Biol.* **8**,
553		e1000412, doi:10.1371/journal.pbio.1000412 (2010).

14 Kilkenny, C., Browne, W. J., Cuthill, I. C., Emerson, M. & Altman, D. G. Improving bioscience research reporting: the ARRIVE guidelines for reporting animal research. *PLoS Biol.* **8**, e1000412, doi:10.1371/journal.pbio.1000412 (2010).

15 Moltzau, L. R. *et al.* Different Compartmentation of Responses to Brain Natriuretic Peptide and C-Type Natriuretic Peptide in Failing Rat Ventricle. *J. Pharmacol. Exp. Ther.* **350**, 681-690, doi:10.1124/jpet.114.214882 (2014).

16 Arai, R., Ueda, H., Kitayama, A., Kamiya, N. & Nagamune, T. Design of the linkers which effectively separate domains of a bifunctional fusion protein. *Protein Eng.* **14**, 529-532 (2001).

17 Meier, S. *et al.* PDE3 inhibition by C-type natriuretic peptide-induced cGMP enhances cAMP-mediated signaling in both non-failing and failing hearts. *Eur. J. Pharmacol.* **812**, 174-183, doi:10.1016/j.ejphar.2017.07.014 (2017).

18 Orstavik, O. *et al.* The inotropic effect of the active metabolite of levosimendan, OR-1896, is mediated through inhibition of PDE3 in rat ventricular myocardium. *PLoS One* **10**, e0115547, doi:10.1371/journal.pone.0115547 (2015).

19 Dagda, R. K. *et al.* Loss of PINK1 function promotes mitophagy through effects on oxidative stress and mitochondrial fission. *J. Biol. Chem.* **284**, 13843-13855, doi:10.1074/jbc.M808515200 (2009).

20 Moltzau, L. R. *et al.* SERCA2 activity is involved in the CNP-mediated functional responses in failing rat myocardium. *Br. J. Pharmacol.* **170**, 366-379, doi:10.1111/bph.12282 (2013).

21 Calamera, G. *et al.* FRET-based cyclic GMP biosensors measure low cGMP concentrations in cardiomyocytes and neurons. *Commun Biol* **2**, 394, doi:10.1038/s42003-019-0641-x (2019).

22 Manfra, O. *et al.* CNP regulates cardiac contractility and increases cGMP near both SERCA and TnI - difference from BNP visualized by targeted cGMP biosensors. *Cardiovasc. Res.*, doi:10.1093/cvr/cvab167 (2021).

23 Brescia, M., Chao, Y. C., Koschinski, A., Tomek, J. & Zaccolo, M. Multi-Compartment, Early Disruption of cGMP and cAMP Signalling in Cardiac Myocytes from the mdx Model of Duchenne Muscular Dystrophy. *Int J Mol Sci* **21**, doi:10.3390/ijms21197056 (2020).

24 Russwurm, M. *et al.* Design of fluorescence resonance energy transfer (FRET)-based cGMP indicators: a systematic approach. *Biochem. J.* **407**, 69-77, doi:10.1042/BJ20070348 (2007).

25 Russwurm, M. & Koesling, D. Measurement of cGMP-generating and -degrading activities and cGMP levels in cells and tissues: Focus on FRET-based cGMP indicators. *Nitric Oxide* **77**, 44-52, doi:10.1016/j.niox.2018.04.006 (2018).

26 Omori, K. & Kotera, J. Overview of PDEs and their regulation. *Circ. Res.* **100**, 309-327, doi:10.1161/01.RES.0000256354.95791.f1 (2007).

27 Rascon, A. *et al.* Purification and properties of the cGMP-inhibited cAMP phosphodiesterase from bovine aortic smooth muscle. *Biochim. Biophys. Acta* **1134**, 149-156, doi:10.1016/0167-4889(92)90038-d (1992).

28 Huang, G. Y. *et al.* Structural basis for cyclic-nucleotide selectivity and cGMP-selective
activation of PKG I. *Structure* **22**, 116-124, doi:10.1016/j.str.2013.09.021 (2014).

29 Huang, L. J. *et al.* NH2-Terminal targeting motifs direct dual specificity A-kinase-anchoring
protein 1 (D-AKAP1) to either mitochondria or endoplasmic reticulum. *J. Cell Biol.* **145**, 951-
959, doi:10.1083/jcb.145.5.951 (1999).

30 Kim, H. *et al.* Fine-tuning of Drp1/Fis1 availability by AKAP121/Siah2 regulates
mitochondrial adaptation to hypoxia. *Mol. Cell* **44**, 532-544 (2011).

31 Merrill, R. A. *et al.* Mechanism of Neuroprotective Mitochondrial Remodeling by
PKA/AKAP1. *PLoS Biol.* **9**, e1000612, doi:10.1371/journal.pbio.1000612 (2011).

32 Götz, K. R. *et al.* Transgenic mice for real-time visualization of cGMP in intact adult
cardiomyocytes. *Circ. Res.* **114**, 1235-1245, doi:10.1161/CIRCRESAHA.114.302437 (2014).

33 Chang, C.-R. & Blackstone, C. Cyclic AMP-dependent Protein Kinase Phosphorylation of
Drp1 Regulates Its GTPase Activity and Mitochondrial Morphology. *J. Biol. Chem.* **282**,
21583-21587, doi:10.1074/jbc.C700083200 (2007).

34 Cribbs, J. T. & Strack, S. Reversible phosphorylation of Drp1 by cyclic AMP-dependent protein
kinase and calcineurin regulates mitochondrial fission and cell death. *EMBO Rep* **8**, 939-944,
doi:10.1038/sj.embor.7401062 (2007).

35 Monterisi, S. *et al.* PDE2A2 regulates mitochondria morphology and apoptotic cell death via
local modulation of cAMP/PKA signalling. *Elife* **6**, doi:10.7554/eLife.21374 (2017).

36 Fiscus, R. R., Tu, A. W. & Chew, S. B. C. Natriuretic peptides inhibit apoptosis and prolong
the survival of serum-deprived PC12 cells. *Neuroreport* **12**, 185-189 (2001).

37 Fiscus, R. R. Involvement of cyclic GMP and protein kinase G in the regulation of apoptosis
and survival in neural cells. *Neurosignals* **11**, 175-190 (2002).

38 Jin, X., Zhang, Y., Li, X., Zhang, J. & Xu, D. C-type natriuretic peptide ameliorates
ischemia/reperfusion-induced acute kidney injury by inhibiting apoptosis and oxidative stress
in rats. *Life Sci.* **117**, 40-45 (2014).

39 Kulhanek-Heinze, S., Gerbes, A. L., Gerwig, T., Vollmar, A. M. & Kiemer, A. K. Protein
kinase A dependent signalling mediates anti-apoptotic effects of the atrial natriuretic peptide in
ischemic livers. *J. Hepatol.* **41**, 414-420 (2004).

40 Kato, T. *et al.* Atrial natriuretic peptide promotes cardiomyocyte survival by cGMP-dependent
nuclear accumulation of zyxin and Akt. *The Journal of clinical investigation* **115**, 2716-2730
(2005).

41 Sun, Y., Zhang, Y., Yan, M., Wu, Y. & Zheng, X. B-type natriuretic peptide-induced
cardioprotection against reperfusion is associated with attenuation of mitochondrial
permeability transition. *Biol. Pharm. Bull.* **32**, 1545-1551 (2009).

623	42	Wang, T.-N., Ge, Y.-K., Li, J.-Y., Zeng, X.-H. & Zheng, X.-X. B-type natriuretic peptide enhances mild hypoxia-induced apoptotic cell death in cardiomyocytes. *Biol. Pharm. Bull.* **30**, 1084-1090 (2007).
626	43	Qvigstad, E. *et al.* Natriuretic peptides increase beta1-adrenoceptor signalling in failing hearts through phosphodiesterase 3 inhibition. *Cardiovasc. Res.* **85**, 763-772, doi:10.1093/cvr/cvp364 (2010).
629	44	Han, B. *et al.* The opposing effects of endothelin-1 and C-type natriuretic peptide on apoptosis of neonatal rat cardiac myocytes. *Eur. J. Pharmacol.* **474**, 15-20 (2003).
631	45	Wu, C.-F., Bishopric, N. H. & Pratt, R. E. Atrial Natriuretic Peptide Induces Apoptosis in Neonatal Rat Cardiac Myocytes. *J. Biol. Chem.* **272**, 14860-14866 (1997).
633	46	Suenobu, N., Shichiri, M., Iwashina, M., Marumo, F. & Hirata, Y. Natriuretic peptides and nitric oxide induce endothelial apoptosis via a cGMP–dependent mechanism. *Arterioscler. Thromb. Vasc. Biol.* **19**, 140-146 (1999).
636	47	Costa, L. E. *et al.* Long-term angiotensin II inhibition increases mitochondrial nitric oxide synthase and not antioxidant enzyme activities in rat heart. *J. Hypertens.* **20**, 2487-2494 (2002).
638	48	Boveris, A., D'Amico, G., Lores-Arnaiz, S. & Costa, L. E. Enalapril increases mitochondrial nitric oxide synthase activity in heart and liver. *Antioxidants and Redox Signaling* **5**, 691-697 (2003).
641	49	Pozdniakova, S. *et al.* 17β-Estradiol reduces mitochondrial cAMP content and cytochrome oxidase activity in a phosphodiesterase 2-dependent manner. *Br. J. Pharmacol.* **175**, 3876-3890 (2018).
644	50	Mitsuishi, M., Miyashita, K. & Itoh, H. cGMP rescues mitochondrial dysfunction induced by glucose and insulin in myocytes. *Biochem. Biophys. Res. Commun.* **367**, 840-845, doi:https://doi.org/10.1016/j.bbrc.2008.01.017 (2008).
647	51	Engeli, S. *et al.* Natriuretic peptides enhance the oxidative capacity of human skeletal muscle. *The Journal of clinical investigation* **122**, 4675-4679 (2012).
649	52	Das, A., Smolenski, A., Lohmann, S. M. & Kukreja, R. C. Cyclic GMP-dependent protein kinase Iα attenuates necrosis and apoptosis following ischemia/reoxygenation in adult cardiomyocyte. *J. Biol. Chem.* **281**, 38644-38652 (2006).
652	53	D'Souza, S. P. *et al.* B-type natriuretic peptide limits infarct size in rat isolated hearts via KATP channel opening. *American Journal of Physiology-Heart and Circulatory Physiology* **284**, H1592-H1600, doi:10.1152/ajpheart.00902.2002 (2003).
655	54	Yang, X.-M., Philipp, S., Downey, J. M. & Cohen, M. V. Atrial natriuretic peptide administered just prior to reperfusion limits infarction in rabbit hearts. **101**, 311-318, doi:10.1007/s00395-006-0587-2 (2006).

55   Hong, L. *et al.* Atrial natriuretic peptide prevents the mitochondrial permeability transition pore
opening by inactivating glycogen synthase kinase 3β via PKG and PI3K in cardiac H9c2 cells.
Eur. J. Pharmacol. **695**, 13-19, doi:https://doi.org/10.1016/j.ejphar.2012.07.053 (2012).

56   De Palma, C. *et al.* Nitric oxide inhibition of Drp1-mediated mitochondrial fission is critical
for myogenic differentiation. *Cell Death Differ.* **17**, 1684-1696, doi:10.1038/cdd.2010.48
(2010).

57   Hu, C., Huang, Y. & Li, L. Drp1-Dependent Mitochondrial Fission Plays Critical Roles in
Physiological and Pathological Progresses in Mammals. *International Journal of Molecular
Sciences* **18**, 144 (2017).

58   Acin-Perez, R. *et al.* Cyclic AMP produced inside mitochondria regulates oxidative
phosphorylation. *Cell Metab* **9**, 265-276, doi:10.1016/j.cmet.2009.01.012 (2009).

59   DiPilato, L. M., Cheng, X. & Zhang, J. Fluorescent indicators of cAMP and Epac activation
reveal differential dynamics of cAMP signaling within discrete subcellular compartments. *Proc.
Natl. Acad. Sci. U. S. A.* **101**, 16513, doi:10.1073/pnas.0405973101 (2004).

60   Lefkimmiatis, K., Leronni, D. & Hofer, A. M. The inner and outer compartments of
mitochondria are sites of distinct cAMP/PKA signaling dynamics. *J. Cell Biol.* **202**, 453-462,
doi:10.1083/jcb.201303159 (2013).

61   Subramanian, H. *et al.* Distinct submembrane localisation compartmentalises cardiac NPR1
and NPR2 signalling to cGMP. *Nat Commun* **9**, 2446, doi:10.1038/s41467-018-04891-5 (2018).

62   Piquereau, J. *et al.* Mitochondrial dynamics in the adult cardiomyocytes: which roles for a
highly specialized cell? *Front. Physiol.* **4**, 102, doi:10.3389/fphys.2013.00102 (2013).

63   Sprenger, J. U. *et al.* In vivo model with targeted cAMP biosensor reveals changes in receptor-
microdomain communication in cardiac disease. *Nat Commun* **6**, 6965,
doi:10.1038/ncomms7965 (2015).

64   Koschinski, A. & Zaccolo, M. Activation of PKA in cell requires higher concentration of cAMP
than in vitro: implications for compartmentalization of cAMP signalling. *Sci. Rep.* **7**, 14090,
doi:10.1038/s41598-017-13021-y (2017).

65   Kim, J. J. *et al.* Co-Crystal Structures of PKG Iβ (92–227) with cGMP and cAMP Reveal the
Molecular Details of Cyclic-Nucleotide Binding. *PLoS One* **6**, e18413,
doi:10.1371/journal.pone.0018413 (2011).

Tables

Table 1.

Sensitivity and dynamic range of the FRET biosensors.

HEK293 cells expressing the indicated biosensor were homogenized and incubated with increasing concentration of cGMP or cAMP (where indicated) and fluorescence measured as described in Materials and Methods. The maximal FRET change was determined at the highest cGMP concentrations. EC_{50} values shown in µM were calculated from the average pEC_{50} values. Results shown are mean±SEM of 5-12 experiments. *p<0.001 and †p<0.0001 *vs*. cGi-500V; ‡p<0.01 for cGMP EC_{50} *vs*. cGMP EC_{50} +100µM cAMP (One-way ANOVA, with Sidak's post hoc test).

Sensor	cGMP		cAMP		cAMP		cGMP +1µM cAMP		cGMP +100µM cAMP		Max FRET change (%)
	pEC_{50}	EC_{50} (µM)	pEC_{50High}	EC_{50} (µM)	pEC_{50Low}	EC_{50} (µM)	pEC_{50}	EC_{50} (µM)	pEC_{50}	EC_{50} (µM)	
cGi-500V	6.31±0.04	0.5	7.93±0.06	12	4.04±0.13	106	6.52±0.08	0.3	5.94±0.06‡	1.1	53.24±1.43
ScGI	6.68±0.03*	0.2			3.97±0.03	125	6.65±0.16	0.2	6.24±0.14‡	0.6	34.08±3.62†
OMM-ScGI	6.66±0.06*	0.2			4.32±0.12	52					14.62±0.41†

Table 2.

Fraction of fragmented *vs.* elongated mitochondria.

Data from Figure 5 were distributed in two different fractions per each group, where cells

with mitochondrial elongation (axis ratio) smaller than 1.75 were considered as having

fragmented mitochondria and cells with mitochondria larger than 1.75 were considered as

having elongated mitochondria. The threshold was chosen based on mitochondria size

distribution in Ctr group. Fractions are indicated as per cent of total number of cells per each

group.

Mitochondrial elongation	Ctr	CNP	ANP	H_2O_2	H_2O_2+CNP	H_2O_2 +ANP
<1.75 (fragmented)	30.9%	16.8%	38.8%	81.5%	56.9%	69.6%
>1.75 (elongated)	69.1%	83.2%	61.2%	18.5%	43.1%	30.4%

Figures

Figure 1

Figure 1. NPs reduce cell death and increase ATP levels compared to the control. a) Cardiomyocytes were stimulated with Vehicle (H_2O), CNP (1µM) or ANP (1µM) for 24 h and sampling of the medium performed at the indicated time. LDH activity in the medium was determined as described in Materials and Methods. LDH activity at 24 h from Vehicle was used as control. Data shown are mean±SEM of 8 independent experiments. *p <0.05 compared to vehicle (two-way ANOVA with Dunnett's multiple comparisons test). **b)** Cardiomyocytes were stimulated with vehicle (control), CNP (1µM) or ANP (1µM) for the indicated times and ATP levels measured as described in Materials and Methods and normalized to control ($44\pm14x10^3$ RLU/µg protein at 3h and $26\pm10x10^3$ RLU/µg protein at 24h). Data shown are mean±SEM of 5 independent experiments. *p<0.05 and **p<0.01 compared to control (one-way ANOVA with Sidak's test).

Figure 2

Figure 2. NPs protect against apoptosis through the mitochondrial intrinsic pathway. Cardiomyocytes were incubated with vehicle (H$_2$O, control), CNP (1µM) or ANP (1µM) for 3 h. **a)** Cells were fixed in 1% paraformaldehyde and fragmented DNA was stained by anti-digoxigenin antibody conjugated with fluorescein. Total number of cells and TUNEL positive cells were counted and shown as a per cent of vehicle treated cells (29.4 % ±3.2). Data shown are mean±SEM of 7 independent experiments. **b)** Cells were lysed and the amount of uncleaved and cleaved PARP and vinculin (loading control) were determined as described in Materials and Methods. Insert shows a representative blot. Data are mean±SEM of 5 independent experiments. **c)** Cells were lysed and active caspase 9 was determined as described in Materials and Methods and normalized to vinculin. Insert shows a representative blot. Quantification of cleaved caspase 9 as a fraction of the loading control (vinculin) and are shown as mean±SEM of 4 independent experiments. **d)** Cytochrome c levels were measured in the cytosolic fraction as described in Material and Methods. Values shown are mean±SEM of 5 independent experiments. *p<0.05 and **p<0.01 *vs*. Ctr (one-way ANOVA with Sidak's (**a**) or Dunnett's multiple comparisons (**b-d**) test. a.u.: Arbitrary units.

Figure 3

Figure 3. Development of FRET-based biosensor selective for cGMP and targeted to mitochondria. a-c) Upper panels: schematic of the indicated biosensor consisting of the fluorescent proteins CFP and Venus flanking the two cyclic nucleotide binding domains from PKG I without (cGi-500V) or with (ScGI) the C158R mutation. The OMM-ScGI contains an AKAP1 targeting sequence selective for the outer mitochondrial membrane. Lower panels: homogenates of HEK293 cells transfected with the indicated biosensor were incubated with increasing concentrations of either cGMP or cAMP. FRET (F_{Venus}/F_{CFP}) was measured as described in Materials and Methods and normalized to FRET in the absence of cGMP and at the highest cGMP concentration. Shown are representative of 10 (cGMP) and 5 (cAMP) independent experiments in **a**, 15 (cGMP) and 6 (cAMP) in **b**, 5 (cGMP) and 5 (cAMP) in **c. d)** Concentration-response curves generated by increasing concentrations of cGMP in homogenates of HEK293 cells expressing the indicated biosensors, as shown in a, b and c. EC_{50} values (marked as black filled symbols) and maximal change in FRET (marked as red filled symbols) are shown as mean±SEM from 12, 17 and 5 independent experiments for cGi-500V, ScGI and OMM-ScGI, respectively. *p<0.001 for pEC_{50} of ScGI or OMM-ScGI vs. cGI-500V; †p< 0.0001 for maximal FRET change of ScGI or OMM-ScGI vs. cGi-500V (one-way ANOVA, Sidak's test).

Figure 4

Figure 4. CNP and ANP increase cGMP at the outer mitochondrial membrane in cardiomyocytes. a) Cells expressing the ScGI (upper panel) and OMM-ScGI (lower panel),

were stained with Mitotracker Deep Red, washed and visualized under a confocal microscope as described in Materials and Methods. Scale bar: 10μm. Right panels show colocalization by intensity overlay of both fluorescent signals (Mitotracker and sensors) from indicated dotted bar. a.u.: Arbitrary units **b and c)** Representative FRET ratio (Venus/CFP) recordings in single cells expressing the indicated biosensor and stimulated with CNP (300nM) and IBMX (100μM) (left panels) or ANP (1μM) followed by maximal activation of the biosensor (IBMX 100μM + CNP 300 nM) (right panels). The inserts in the left panels show corresponding traces of separate Venus and CFP intensities upon stimulation with CNP (300 nM). Shown are traces representative of 10-14 cells from 5-8 independent experimtents **d)** Quantification of FRET responses from stimulation with CNP, ANP or combined response of IBMX + ANP. **p<0.01 and ***p<0.001 (one-way ANOVA, Sidak's test).

Figure 5

Figure 5. NPs modulate Drp1 phosphorylation and prevent mitochondrial fragmentation.
a) Adult rat cardiomyocytes were stimulated with vehicle (H_2O; Ctr), CNP (1 μM) or ANP (1 μM) for 3 h and lysed. Phosphorylation of Drp1 at Ser637 (pDrp1^{Ser637}) was determined as described in Materials and Methods. The insert shows rrepresentative western blots of 5 independent experiments. The bands were quantified and normalized to loading control (vinculin) and Ctr (mean±SEM, n=5). *$p < 0.05$ and **$p<0.01$ *vs*. Ctr; †$p<0.01$ CNP *vs*. ANP (one-way ANOVA with Tukey's test). **b)** H9c2 cells were stained with Mitotracker Green, washed and visualized under a confocal microscope and pictures processed by ImageJ to obtain the mitochondrial elongation as described in Materials and Methods. Data are shown as mean±SEM of 102-158 cells from 4 independent experiments. *$p<0.05$, ***$p<0.001$ (one-way ANOVA, Sidak's multiple comparisons test).

Figure 6. Natriuretic peptides reduce cardiomyocyte apoptosis through the mitochondrial intrinsic pathway by increasing cGMP at the OMM and increasing Drp1 phosphorylation. GC-A and GC-B stimulation by ANP and CNP respectively, generates cGMP differently in the cytosol and at the OMM, with the CNP-stimulated cGMP pool being greater than the ANP-stimulated cGMP pool. cGMP can activate PKG and mediate Drp1 phosphorylation, preventing its activation and its pro-apoptotic function at the OMM. ANP and CNP reduce cytochrome c release from the mitochondria, decreasing caspase 9 activation and the progression in apoptosis (PARP cleavage), while only CNP can prevent mitochondrial fragmentation. Dotted lines indicate suggested pathways. This figure was produced using Biorender.