1 **Sex, stress and UCP2:**
2 **mitochondria illuminate microglial identity beyond the dichotomy**
3

4 **Authors**
5 Margaret E Maes[1], Gloria Colombo[1], Felix Sternberg [2], Elena E Pohl[2], Sandra Siegert[1†]
6

Affiliations
[1] Institute of Science and Technology Austria (ISTA), Am Campus 1, 3400 Klosterneuburg,
Austria
[2] Institute of Physiology, Pathophysiology and Biophysics, University of Veterinary
Medicine, Veterinärplatz 1, 1210 Vienna, Austria
† Corresponding author: ssiegert@ist.ac.at
13

Abstract
Environmental cues influence microglial interaction with their neuronal landscape, often
causing changes in microglial functional identity. This process demands rapid energy
production, which is supported by an optimized mitochondrial network. Whether
mitochondrial adaptations define a microglial functional state is unknown due to limitations
in mitochondrial visualization and manipulation within their microenvironment.
Here, we investigate the mitochondrial network of individual microglia in physiological and
dysfunctional conditions during a naïve or injury-induced mouse retinal environment. After
optic nerve crush (ONC) injury, we identify mitochondrial and microglial phenotypic
remodeling that is distinct from the naïve environment. Remarkably, knockout of uncoupling
protein 2 (UCP2) causes transient, stress-induced mitochondrial hyperfusion only in males
after ONC, even though both sexes suffer from increased intrinsic stress levels. When we
compare key metabolic genes between sexes in physiological conditions, only males exhibit a
metabolic transition towards glycolysis in the injury-induced environment, which attenuates
in UCP2 knockout resulting in acutely accelerated retinal ganglion cell death. Our data shows
that the cell-extrinsic ONC environment and cell-intrinsic UCP2[KO] manipulation triggers
several microglial states distinguishable by mitochondrial networks that are heavily
influenced by sex and injury-induced stress.
32

33 **Key words**
34 Microglia, mitochondria, UCP2, retina, sex, stress, hyperfusion, optic nerve crush

Introduction

Microglia transiently and dynamically respond to a myriad of cues within their microenvironment, ranging from local surveillance to active engagement in removing cellular components (Hanisch and Kettenmann, 2007; Salter and Stevens, 2017). An unresolved challenge is how to define this spectrum of microglial functional states (Paolicelli et al., 2022), which has been described by single cell transcriptomes and morphological analyses (Colombo et al., 2021; Masuda et al., 2020; Stogsdill et al., 2022). Moreover, systemic factors such as sex, age, and genetic background influence the microglial functional state. Whereas -omics studies are often limited in their readout to a sole factor *e.g.,* gene transcripts, a multifaceted readout for microglia within their microenvironment would be desirable.

A cell's energetic demands predominantly rely on mitochondrial function, where ATP production is optimized through the dynamic processes of organelle fission and fusion (Mishra and Chan, 2016; Rambold and Pearce, 2018). In physiological conditions, these mitochondrial dynamics allow for exchange of matrix content or elimination of dysfunctional organelles, creating a balanced, homeostatic mitochondrial network (Youle and van der Bliek, 2012). Under high energy demand, this equilibrium is disrupted, resulting in a more fragmented mitochondrial network as the cell shifts from oxidative phosphorylation (OXPHOS) to favor glycolysis (Buck et al., 2016). In contrast, mitochondria can also take on a hyperfused morphology, which is reflected by a filamentous, highly connected mitochondrial network (Gomes et al., 2011; Tondera et al., 2009). This hyperfused network serves as a short-term cell survival mechanism to increase membrane potential and calcium accumulation enhancing ATP levels in stressed environments (Das and Chakrabarti, 2020; Gomes et al., 2011; Mitra et al., 2009; Tondera et al., 2009; Wai and Langer, 2016). Therefore, mitochondrial network connectivity is a critical parameter to discriminate the intrinsic cellular energetic demands.

Transcriptomic studies frequently identify differentially expressed metabolism-associated genes in microglia during development and disease progression (Baik et al., 2019; Bernier et al., 2020; He et al., 2022; Ulland et al., 2017), that is paralleled in macrophages outside the CNS (Borst et al., 2019; Jha et al., 2015). One such gene is uncoupling protein 2 (UCP2, Slc25a8) which is found in immune cells (Arsenijevic et al., 2000; Pecqueur et al., 2001; Rupprecht et al., 2012), including microglia (Siegert et al., 2012), as well as in cells that

favor glycolysis such as, rapidly proliferating cells or stem cells (Horimoto et al., 2004;

Rupprecht et al., 2014, 2019; Zhang et al., 2011). The two main transport functions of UCP2

are the fatty acids-induced import of protons to bypass ATP synthetase (Ježek et al., 2018)

and export of Krebs cycle metabolites to limit the oxidation of acetyl-CoA producing

substrates (Vozza et al., 2014). Loss of UCP2 has recently been reported to inhibit

mitochondrial dynamics, ATP production, and microglial inflammation in the hypothalamus

after high fat diet (Kim et al., 2019; Toda et al., 2016). This makes UCP2 a promising

candidate to manipulate mitochondria and determine their impact on the microglial functional

state.

One major limitation in assessing mitochondrial networks *in vivo* is targeting cell type-

specific mitochondria, which can easily be achieved *in vitro*. However, studying microglial

properties *in vitro* or *ex vivo* leads to significant morphological and transcriptomic alterations

that do not recapitulate the *in vivo* environment (Galatro et al., 2017; Gosselin et al., 2017;

Hammond et al., 2019; Matcovitch-Natan et al., 2016). Moreover, sex hormones have been

shown to influence microglial function (Villa et al., 2018), further emphasizing the need to

investigate microglia within their microenvironment. Thus, we must address the following

two challenges: first, their selective visualization and second, their targeted manipulation

with minimal external interference.

Here, we generated a mouse model that labels mitochondria exclusively within microglia,

allowing us to perform multifaceted analysis on single microglia in their microenvironment.

We focused on the retina where microglial populations are distinctly separated in the outer-

and inner plexiform layers (OPL and IPL, respectively). Then, we selectively knocked out

UCP2 to establish a dysfunctional mitochondrial condition. Finally, we performed optic

nerve crush (ONC), which damages the axonal projections of retinal ganglion cells (RGCs) in

the optic nerve but leaves the dura and the blood vessels intact (Williams et al., 2020). As a

consequence, microglia in the inner plexiform layer (IPL) respond to RGCs undergoing

apoptosis within the first week (Liu et al., 2012; Mac Nair et al., 2016; Wohl et al., 2010).

Remarkably, our multifaceted mitochondria and microglial analysis *in vivo* revealed sexual-

dimorphic microglia functional states based on the ONC and the loss of UCP2. This

underscores the necessity of evaluating sex in microglia in neuropathological environments.

Results

Resolving mitochondrial networks of microglia in naïve environments

To selectively visualize mitochondria in microglia, we took advantage of the mouse reporter line PhAM$^{fl/fl}$, which labels mitochondria with a Dendra2-tag (Pham et al., 2012). We crossed this mouse with the $Cx3cr1^{CreERT2}$ mouse model (Yona et al., 2013) (**Figure 1A**). Upon tamoxifen induction, only microglia detected with IBA1 (ionized calcium binding adaptor molecule 1) expressed the Dendra2-tag in the retina (**Figure 1B-C**). Throughout this study, we will refer to this tamoxifen-induced mouse model $Cx3cr1^{CreERT2/+}$/PhAM$^{fl/fl}$ as µG^{mito} mice. To perform targeted mitochondrial manipulation, we revisited our retinal transcriptome dataset for mitochondria-associated genes and confirmed high expression of uncoupling protein 2 (UCP2) in retinal microglia (**Figure S1A-B**) (Siegert et al., 2012). Since recent single cell RNA-sequencing studies also supported UCP2 transcript enrichment in retinal microglia (Hoang et al., 2020; O'Koren et al., 2019), we generated a tamoxifen-inducible microglia-specific UCP2-knockout mouse model by crossing the $Ucp2^{fl/fl}$ mouse (Kong et al., 2010) with µG^{mito} (**Figure 1D**). This $Cx3cr1^{CreERT2/+}$/Pham$^{fl/fl}$/$Ucp2^{fl/fl}$ mouse model, hereafter referred to µG^{mito}UCP2^{KO}, also showed microglia-selective mitochondrial labeling (**Figure 1E-F**). To verify UCP2 depletion, we isolated retinal microglia with fluorescence-activated cell sorting (FACS) using their Dendra2-tag (**Figure S1C**). We found that the $Ucp2$ transcript was undetectable three weeks after tamoxifen induction (**Figure S1D**). Similarly, UCP2 protein expression decreased after 4-OHT (4-hydroxytamoxifen) exposure in murine primary microglia culture (**Figure S1E-G**).

After establishing both mouse models, we profiled the mitochondrial network of microglia at single-cell resolution (**Figure 1C, F**). We first observed that the mitochondria of both conditions intersect at the soma and extend through the primary microglial processes. To further evaluate these networks, we assessed three parameters: mitochondrial connectivity based on the median organelle volume and total number of organelles per cell (**Figure 1G**), the percentage of mitochondria in microglia as an estimation of mitochondrial biogenesis (Ron-Harel et al., 2016) (**Figure 1H**), and perinuclear mitochondrial localization to gain insights into transcriptional activity (Al-Mehdi et al., 2012) (**Figure 1I**). All mitochondrial parameters were indistinguishable between µG^{mito} and µG^{mito}UCP2^{KO} microglia in the homeostatic environment (**Figure 1G-I**). Thus, we established the balanced mitochondrial networks of microglia within their homeostatic microenvironment, which are not critically dependent on UCP2 in physiological conditions.

Optic nerve crush upregulates microglial UCP2 and alters the mitochondrial network

Next, we were interested to trigger an injury-induced environment in the retina. For this, we took advantage of the optic nerve crush (ONC) model (Levkovitch-Verbin et al., 2000), where the retinal ganglion cell (RGC) axons posterior to the optic disc are mechanically injured (**Figure 2A**). This induces RGC apoptosis (**Figure S2A**) and microglial reactivity (**Figure 2B**) (Li et al., 1999; Wohl et al., 2010). To determine the time of robust microglial response after ONC, we collected retinas 1, 5, and 14 days after either ONC or sham injury and compared their morphological changes using Sholl analysis (**Figure S2B-C**). The most pronounced adaptation occurred at 5 days after ONC, whereas the sham surgery-induced response was no longer visible (**Figure S2C**). In the injury-induced environment, microglia upregulate the endosomal-lysosomal marker CD68 (Bauer et al., 1994; Damoiseaux et al., 1994). Thus, we assessed the percentage volume of CD68 per microglia volume and confirmed the Sholl analysis identified response pattern at 5 days (**Figure S2D-E**). Microglia in the outer plexiform layer (OPL) reflected transient morphological adaptation at 5 days but without changes in CD68 volume (**Figure S2F-I**). When we analyzed the *Ucp2* transcript level in μG^{mito} microglia from untreated (naïve) mice or mice 5 days after ONC, the *Ucp2* transcript was significantly upregulated (**Figure 2C**) indicating *Ucp2* involvement in the ONC microglial response at 5 days after injury.

Next, we performed the single microglia analysis and compared microglial morphology, CD68, and the mitochondrial network in naïve (**Figure 2D-F**) and 5 days after ONC in μG^{mito} mice (**Figure 2G-I**). Microglia reduced their branching complexity shown by an increased number of Sholl intersections closer to the soma (**Figure 2J**). Furthermore, they expressed a greater percentage volume of CD68, with increased CD68-vesicle localization in the peripheral microglial processes (**Figure 2K-L**). When we looked at the mitochondrial profile, the median mitochondrial volume was reduced while the total number of organelles increased (**Figure 2M**). This indicates a more fragmented mitochondrial network and suggests changes in metabolic demand (Buck et al., 2016). We also found mitochondrial content (**Figure 2N**) and localization to the perinuclear region (**Figure 2O**) significantly increased. This data shows that microglia upregulate UCP2 and adapt their mitochondrial network in the injury-induced microenvironment.

Loss of UCP2 elicits a sexually dimorphic microglial response after ONC

Previous studies in the arcuate nucleus of the hypothalamus suggested that loss of UCP2 affects microglial reactivity and prevented mitochondrial alterations after high fat-diet (Kim et al., 2019). Surprisingly, when we performed ONC in the UCP2^{KO} condition (**Figure 3**), mitochondria were more fragmented (**Figure 3H-J**), showed an overall increased volume per microglia (**Figure 3K**), and localized to the perinuclear space (**Figure 3L**). This is in line with microglial adaptations that we observed in **Figure 2**, which recapitulated the reduced microglia branching complexity (**Figure 3A-C**), and increased CD68 volume (**Figure 3D-G**).

To resolve this and obtain unbiased insights into the relationship between the parameters, we applied principal component analysis (PCA) across experimental groups (μG^{mito}_{naive}, μG^{mito}-$_{ONC}$, $\mu G^{mito}UCP2^{KO}_{naive}$, $\mu G^{mito}UCP2^{KO}_{ONC}$, see Methods for parameter descriptions). In the principal component (PC) space, each dot represents a microglia profile, which contains all parameters for a single-cell within a defined condition and environment. In the homeostatic environment, the profiles of naïve μG^{mito} and $\mu G^{mito}UCP2^{KO}$ intermingled within the same PC space (**Figure 4A**), indicating that UCP2 loss does not affect the microglial profile at baseline conditions as indicated in **Figure 1G-I**. Also, we did not find any polarization in the naïve clusters analyzed at homeostatic conditions that would point to sexual differences of the microglial profiles (**Figure 4B-C**).

In the ONC-induced environment, individual microglial profiles transitioned away from their naïve counterparts along the first PC (PC1, **Figure 4D**). Here, we observed a difference between μG^{mito} and $\mu G^{mito}UCP2^{KO}$ conditions, in which the microglia profiles from the UCP2^{KO} were positioned between μG^{mito}_{naive} and μG^{mito}_{ONC} profiles. Moreover, they showed an increased spread along the second PC (PC2) indicating that UCP2^{KO} alters the microglia profile. Strikingly, sex explained the segregation of $\mu G^{mito}UCP2^{KO}_{ONC}$ along PC2. While μG^{mito}_{ONC} ♂ and ♀ profiles remained indistinguishable (**Figure 4E**), there was distinct separation between ♂ and ♀ microglia in $\mu G^{mito}UCP2^{KO}_{ONC}$ (**Figure 4F**). Contrary to previous reports, our data suggests that UCP2 loss causes a sex-specific microglial profile after ONC injury.

Male UCP2^{KO} microglia exhibit hyperfused mitochondria after ONC

To identify which of the analyzed parameters in the microglia profile strongly contributed to the sex-specific differences of $\mu G^{mito}UCP2^{KO}_{ONC}$ (**Figure 4F**), we looked at the PC loadings.

The top contributing parameters in PC1 were the Sholl index and CD68, whereas in PC2 they were mitochondria-derived (**Figure S3A**). To determine whether the sex-dependency persisted with a subset of parameters, we repeated the PCA separately on the mitochondrial (**Figure 5A-C**), or the Sholl index and CD68 parameters (**Figure S3B-D**). The Sholl index and CD68 volume PCA showed no evidence of sex-dependency (**Figure S3C-D**). In contrast, the mitochondrial parameters recapitulated the segregation (**Figure 5C**) as illustrated for naïve and ONC counterparts (**Figure 4D**). The microglial profiles from the μG^{mito} spread along PC1 and importantly, the sex-specific segregation of $\mu G^{mito}UCP2^{KO}_{ONC}$ persisted along PC2 (**Figure 5C**). The median mitochondrial volume and the number of mitochondria were the influential factors for PC1 (**Figure S3E**). These parameters suggested a more fragmented mitochondrial network in $\male$ and $\female \mu G^{mito}_{ONC}$ microglia (**Figure S3F-G**) – an anticipated mitochondrial network change after injury. Interestingly, $\female \mu G^{mito}UCP2^{KO}_{ONC}$ behaved similar to μG^{mito}_{ONC} microglia and intermingled with their profiles (**Figure 5A-C**). In contrast, $\male \mu G^{mito}UCP2^{KO}_{ONC}$ lacked this fragmentation (**Figure S3F-G**) and instead significantly increased the percentage of mitochondrial volume per microglia (**Figure S3G**). This parameter also explained the variance along PC2 (**Figure S3E**). Together, these three parameters suggested a hyperfused mitochondria network (**Figure S3F-H**). Therefore, we closely re-examined the mitochondria within the microglia and found the mitochondrial network was highly connected in $\male \mu G^{mito}UCP2^{KO}_{ONC}$ microglia (**Figure 5D-E**). When we plotted the volume of each organelle within a cell, rather than the median mitochondrial volume, we found that $\male \mu G^{mito}UCP2^{KO}_{ONC}$ microglia had significantly more organelles with large volumes ($> 20\mu m^3$) compared to females (**Figure 5D-E**). These data illustrate that UCP2 loss induced a hyperfused mitochondrial network in male microglia after ONC.

Sexually dimorphic metabolic stress response elucidates mitochondrial hyperfusion

Mitochondrial hyperfusion has been described *in vitro* as a transient phenotype that protects cellular and mitochondrial health in stress-induced environments (Gomes et al., 2011; Tondera et al., 2009). To assess whether the hyperfused mitochondrial network is transient, we included individual microglia analyzed at day 14 after ONC in the mitochondrial PCA analysis. At 14 days, all microglial profiles formed a single cluster in the PC space (**Figure 6A, S4A**) including the $\male \mu G^{mito}UCP2^{KO}_{ONC}$. Accordingly, the number of large-volume mitochondria was reduced (**Figure 6B**), suggesting that the hyperfused phenotype at day 5 is transient after injury, reflecting a unique microglial profile.

Since the hyperfused mitochondrial network is an adaptive mechanism to manage cellular

stress (Tondera et al., 2009), we interrogated the relative transcript expression of key

regulators of metabolic pathways and cellular stress response (**Figure 6C**): *Pgc1-a*

(proliferator-activated receptor gamma coactivator 1- alpha) and *Nrf1* (nuclear respiratory

factor 1) for oxidative phosphorylation (OXPHOS); *Hif1-a* (hypoxia inducible factor 1 alpha)

for glycolytic regulation (Finck and Kelly, 2006); the oxidative stress indicator *Sod1*

(superoxide dismutase 1); and *Cd68* (cluster of differentiation 68) as microglial cell stress

markers (Bauer et al., 1994; Spanier et al., 2009). After RT-qPCR of FACSorted microglia,

we performed PCA to identify whether conditions separate in the PC space and which

variables contribute to this segregation. The PC loadings indicated that metabolic genes

Pgc1-a and *Nrf1* strongly influenced PC1, whereas the stress genes affected PC2 (**Figure**

S4B). In the naïve environment, the sex of both conditions shared the same PC space (**Figure**

S4C). As expected, $UCP2^{KO}$ condition separated from μG^{mito} along PC2 for both sexes

(**Figure 6D**), suggesting a more stressed environment already in the naïve environment.

Surprisingly, upon ONC we observed a prominent difference between male and female

microglia in both μG^{mito} and $\mu G^{mito}UCP2^{KO}$. Whereas $\female\mu G^{mito}_{ONC}$ microglia occupied the

same PC space as in naïve condition, $\male\mu G^{mito}_{ONC}$ microglia moved along the metabolism-

associated PC1 (**Figure 6E**). When we referenced the individual genes, $\male\mu G^{mito}_{ONC}$

microglia significantly shifted towards a more glycolytic metabolism with decreased *Pgc1-α*

and *Nrf1* and some change in *Hif1-α* transcript level (**Figure S4D**). In contrast, $\female\mu G^{mito}_{ONC}$

microglia transcript levels were unaffected (**Figure S4E**). Upon $UCP2^{KO}$ in the injury-

induced environment, this metabolic shift was less pronounced in males compared to

$\male\mu G^{mito}_{ONC}$ (**Figure 6F**) suggesting that male microglia response was attenuated. In fact,

$\male\mu G^{mito}UCP2^{KO}_{ONC}$ had greater *Sod1* upregulation than $\female\mu G^{mito}UCP2^{KO}_{ONC}$ or either sex in

μG^{mito} conditions (**Figure S4F**).

Finally, to obtain insights whether increased metabolic stress in $\male\mu G^{mito}UCP2^{KO}_{ONC}$

microglia impacts retinal ganglion cell (RGC) survival after injury, we assessed RGC

numbers using the marker RBPMS (RNA binding protein with multiple splicing) (Rodriguez

et al., 2014). $\male\mu G^{mito}UCP2^{KO}_{ONC}$ had significantly more RGC loss at day 5 after ONC

compared to females (**Figure 6G**). However, this difference was transient as the loss of

RGCs was similar between sexes by day 14 (**Figure 6G**).

Overall, this data indicates that the hyperfused mitochondrial network in $\male\mu G^{mito}UCP2^{KO}_{ONC}$

is due to greater intrinsic stress of $\male$microglia, resulting in an acute adverse effect in the

microenvironment.

<u>Discussion</u>

Extensive *in vitro* studies have shown that cellular energy demands of immune cells critically

depend on an optimized equilibrium of mitochondrial fission and fusion (Rambold and

Pearce, 2018). Whether this holds true for microglia *in vivo* is unknown due to the technical

limitations of selective mitochondrial visualization and manipulation, two difficulties that we

overcome here. We demonstrate that distinct mitochondrial networks reflect *in vivo* microglia

functional states dependent on the mitochondrial condition (μG^{mito} or $\mu G^{mito}UCP2^{KO}$) and the

local environment (naïve or ONC).

In the naïve environment, mitochondria exhibit a balanced network with organelles spanning

the primary microglial processes (**Figure 1C**, **F-I**). However, after ONC injury, mitochondria

of μG^{mito} accumulated in the perinuclear region (**Figure 2I**, **O**), increased their content

(**Figure 2N**) and became more fragmented throughout the microglial processes and soma

(**Figure 2M**). This corroborates reports of mitochondrial fragmentation after

lipopolysaccharide stimulation of microglia *in vitro* and of increased mitochondrial mass in

immature microglia of germ-free environment *ex vivo* (Erny et al., 2021; Nair et al., 2019;

Park et al., 2013). We also found unexpected mitochondrial alterations in the $\mu G^{mito}UCP2^{KO}$

condition (**Figure 3I**, **J-L**) (Toda et al., 2016), which we further elucidated using an unbiased

PCA approach (**Figure 4**). PCA indicated that the mitochondrial parameters explained the

sex-dependent variance among the ONC-induced $\mu G^{mito}UCP2^{KO}$ microglia profiles.

Remarkably, this sex-specific difference persisted even if we performed the PCA for the

mitochondrial parameters alone (**Figure 5C**) and pointed towards the hyperfused

mitochondrial network present exclusively in males (**Figure 5D**). Overall, our analysis

supports that mitochondrial networks elucidate microglial functional states in their well-

defined microenvironments with sex as a confounding factor.

Mitochondrial hyperfusion has predominantly been described in mouse embryonic fibroblast

cell lines (Gomes et al., 2011; Tondera et al., 2009). Here we report that this phenotype also

occurs *in vivo*. Until now, UCP2 has been implicated in cell metabolism (Rupprecht et al.,

2019) and stress responses (Smorodchenko et al., 2017) even though its function is not fully

resolved (Arsenijevic et al., 2000; Faas et al., 2021; Vozza et al., 2014). Our data shows that

microglia lacking UCP2 have increased intrinsic stress in the naïve environment (**Figure**

S4F), aligning with previous literature (Kim et al., 2019; Yasumoto et al., 2021). However,

the added extrinsic stressor of the injury-induced environment provoked a hyperfused

mitochondrial network, a hallmark of a pro-survival mechanism to manage extensive

intracellular stress (Wai and Langer, 2016). Interestingly, recent studies reported that

UCP2^{KO} in microglia prevents mitochondrial fission (Kim et al., 2019; Toda et al., 2016),

which could be one interpretation of our results. However, with a whole-cell representation of

the mitochondrial network, combined with the contrasting sex-dependent phenotypes, we can

conclude that UCP2 does not directly mediate fission, but rather affects stress-induced

mitochondrial network dynamics.

The hyperfused mitochondrial phenotype in male UCP2^{KO} was striking, but transient,

therefore we cannot exclude that females exhibit a similar phenotype at different injury time

points. However, we already observed a differing metabolic gene response after ONC in the

physiological mitochondria condition (**Figure 6**), emphasizing an impact of sex prior to

additional stressors. Sex hormones have been suggested to drive microglial dimorphisms by

epigenetically shaping microglia during development and continuing to influence them

during adulthood (Baker et al., 2004; Thion et al., 2018; Villa et al., 2018; Yang et al., 2004).

For example, estrogens are implicated to be protective under higher cell-stress conditions

(Colombo et al., 2021; Man et al., 2003; Nakazawa et al., 2006; Nuzzi and Caselgrandi,

2022; Quigley and Broman, 2006). Whereas in UCP2^{KO}, both sexes showed increased

intrinsic stress response in the naïve environment (**Figure 6D**), only the males exhibited

acute acceleration of RGC death 5 days after ONC (**Figure 6G-H**), a time point of robust

microglia response and peak RGC apoptotic commitment (Galindo-Romero et al., 2013;

Maes et al., 2017). Such sex-specific susceptibility has only been reported in long-term RGC

degenerative models (Libby et al., 2005). Our data suggests that microglia play an early

protective role after ONC, even if short-lived, which is contrary to previous reports (Hilla et

al., 2017).

In summary, we demonstrate that mitochondrial networks provide another modality to

interrogate the microglial functional state and highlights the importance of sex. This will

have substantial implications as we begin to address immunometabolism-related targets for

therapy in neurodegenerative diseases.

Author contributions

M.M. and S.S. conceived, developed experimental design and analysis, and wrote the manuscript. M.M., F.S. and E.E.P. conceived and verified UCP2 protein knockout. M.M. performed experiments, data and statistical analysis, and created figures. G.C. performed FACS, RT-qPCR experiments, and data analysis. All authors provided comment and discussion on the manuscript and approved the final version.

Acknowledgements

We thank the Scientific Service Units (SSU) of ISTA through resources provided by the Imaging and Optics Facility (IOF), the Lab Support Facility (LSF), and the Pre-Clinical Facility (PCF) team, specifically Sonja Haslinger and Michael Schunn for excellent mouse colony management and support. This research was partly supported by the FWF Sonderforschungsbereich F83 (to E.E.P). We thank Bálint Nagy, Alessandro Venturino, Ryan John A. Cubero, Marco Benevento and all members of the Siegert group for constant feedback on the project and manuscript.

<u>Figure legends</u>

Figure 1. Mitochondrial networks in retinal microglia of the inner plexiform layer (IPL).

(**A-F**) Mouse models for mitochondrial visualization (**A-C**) and UCP2 (uncoupling protein 2) knockout (**D-F**). (**A, D**) Schematic of mouse models. $Cx3cr1^{CreERT2}$/Pham$^{fl/fl}$ for selective mitochondria-labeling in microglia (**A**) or combined with UCP2 knockout ($Cx3cr1^{CreERT2/+}$/Pham$^{fl/fl}$/UCP2$^{fl/fl}$) (**D**). Upon tamoxifen injection for three consecutive days in adult mice, $Cx3cr1^{CreERT2}$-expressing microglia excise the stop cassette in Pham$^{fl/fl}$ (**A**, μG^{mito}) or in addition, exon 3 and 4 of UCP2 in Pham$^{fl/fl}$/UCP2$^{fl/fl}$ (**D**, $\mu G^{mito}UCP2^{KO}$) to induce mito-Dendra2 (green) expression and UCP2 knockout. (**B, E**) Overview image of immunostained microglia (IBA1, magenta) in the inner plexiform layer (IPL) expressing the mitochondrial label (Dendra2, green) in retinal wholemounts for μG^{mito} (**B**) and $\mu G^{mito}UCP2^{KO}$ (**E**). Scale bar: 50 µm. (**C, F**) Zoom-in (dashed orange line) of a single cell acquisition of immunostained IPL microglia (IBA1, magenta) expressing mitochondria label Dendra2 from μG^{mito} (**C**) and $\mu G^{mito}UCP2^{KO}$ (**F**) retinas. Below: Zoom-out of the corresponding 3D-surface rendering of IBA1 microglia (magenta outline) and Dendra2 mitochondria (green). Dashed blue circle: Location of cell soma. Scale bar: zoom-in: 5 µm,

3D-surface: 10 μm. (**G-I**) Mitochondrial parameters. Boxplot minimum and maximum:

InterQuartile Range (IQR) around median (center line). Whiskers: 1.5 IQRs. Black diamond:

outliers outside of 1.5 IQRs. Each overlaid point represents data point of a single microglia.

Filled circle: female, filled diamond: male. μG^{mito} naïve: n = 18 cells, 5 retinas;

$\mu G^{mito}UCP2^{KO}$: n = 21 cells, 4 retinas. (**G**) Mitochondrial network connectivity determined

by median mitochondrial volume of all organelles (left, Student's t-test: $p = 0.7710$) and

number of organelles within a single cell (right, Student's t-test: $p = 0.7187$).

(**H**) Percentage of mitochondrial volume per microglial volume. Wilcoxon rank sum test: $p =$

0.6121. (**I**) Percentage of total mitochondrial volume localized within the perinuclear region

(10 μm radius from center soma point). Student's t-test: $p = 0.4908$.

[n.s.] $p > 0.05$: not significant.

Figure 2. *Ucp2* is upregulated upon ONC and accompanied by mitochondrial network

alterations in responsive μG^{mito} microglia.

(**A**) Schematic of optic nerve crush (ONC). Forceps induces unilateral damage (red) to retinal

ganglion cell's (RGC) axonal projections (orange) posterior to the optic disc. (**B**) Left, retinal

side-view schematic. OPL, outer plexiform layer. IPL, inner plexiform layer. GCL, ganglion

cell layer. Right, overview confocal image of immunostained IPL microglia (IBA1, magenta)

of C57BL6/J in naïve (left) and 5 days after ONC (right) condition. Scale bar: 30 μm. (**C**)

Boxplot of relative *Ucp2* transcript expression from Dendra2$^+$-FACSorted retinal microglia

in naïve or 5 days ONC condition. Boxplot minimum and maximum: InterQuartile Range

(IQR) around median (center line). Whiskers: 1.5 IQRs. Black diamond: outliers outside of

1.5 IQRs. Filled circle: female, filled diamond: male. Naïve: grey. ONC: blue. Wilcoxon rank

sum test: $p < 0.0458$. (**D-O**) Comparison of microglial and mitochondrial response in μG^{mito}

in naïve (grey) and 5 days after ONC (blue). (**D- I**) A representative IBA1-immunostained

microglia for naïve (**D-F**) or 5 days ONC (**G-I**) depicting 3D-traced filaments (**D, G**) or 3D-

surface renderings of IBA1 (magenta outline) and CD68 (blue **E, H**) or mitochondria (green,

F, I). Dashed blue circle: Location of cell soma. Scale bar: 10 μm. (**J**) Line plot for mean

number of Sholl intersections per radial distant from the soma (in μm) with 95% confidence

interval band. Linear regression analysis: $p < 0.0001$. (**K-O**) Boxplot minimum and

maximum: InterQuartile Range (IQR) around median (center line). Whiskers: 1.5 IQRs.

Black diamond points: outliers outside 1.5 IQRs. Each overlaid point represents data point of

a single microglia. Filled circle: female, filled diamond: male. $\mu G^{mito}_{naïve}$, n= 18 cells, 5

retinas; μG^{mito}_{ONC}, n = 14 cells, 6 retinas. (**K**) Percentage of total CD68 volume per microglia volume. Wilcoxon rank sum test: $p < 0.0001$.

(**L**) Percentage of total CD68 volume localized in the peripheral processes of microglia (>10 µm radius from center soma point). Wilcoxon rank sum test: $p = 0.0021$. (**M**) Mitochondiral network connectivity determined by median mitochondrial volume of all organelles (left, Welch's t-test: $p < 0.0001$) and number of organelles within a single cell (right, Welch's t-test: $p = 0.0004$). (**N**) Percentage of mitochondrial volume per microglial volume. Student's t-test: $p = 0.0126$. (**O**) Percentage of total mitochondrial volume localized within the perinuclear region (<10 µm radius from center soma point). Wilcoxon rank sum test: $p < 0.0001$. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

Figure 3. $\mu G^{mito}UCP2^{KO}$ microglia exhibit ONC-induced mitochondrial and microglial alterations

(**A-L**) Comparison of microglial and mitochondrial response in $\mu G^{mito}UCP2^{KO}$ in naïve (grey/sand) and 5 days after optic nerve crush (ONC, gold). (**A-I**) A representative IBA1-immunostained microglia for naïve (**A**, **D**, **H**) or 5 days ONC (**B**, **E**, **I**) depicting 3D-traced filaments (**A**, **B**) or 3D-surface renderings of IBA1 (magenta outline) and CD68 (blue **D**, **E**) or mitochondria (green, **H**, **I**). Dashed blue circle: location of cell soma. Scale bar: 10 µm. (**C**) Line plot for mean number of Sholl intersections per radial distant from the soma (in µm) with 95% confidence interval band. Linear regression analysis: $p < 0.0001$. (**F-L**) Boxplot minimum and maximum: InterQuartile Range (IQR) around median (center line). Whiskers: 1.5 IQRs. Black diamond points: outliers outside 1.5 IQRs. Each overlaid point represents data point of a single microglia. Filled circle: female, filled diamond: male. $\mu G^{mito}UCP2^{KO}_{naïve}$, n = 21 cells, 4 retinas; $\mu G^{mito}UCP2^{KO}_{ONC}$, n = 22 cells, 4 retinas. (**F**) Percentage of total CD68 volume per microglia volume. Wilcoxon rank sum test: $p = 0.0004$. (**G**) Percentage of total CD68 volume localized in the peripheral processes of microglia (>10 µm radius from center soma point). Student's t-test: $p = 0.9277$. (**J**) Mitochondiral network connectivity determined by median mitochondrial volume of all organelles (left, Welch's t-test: $p = 0.0008$) and number of organelles within a single cell (right, Welch's t-test: $p = 0.0693$). (**K**) Percentage of mitochondrial volume per microglial volume. Wilcoxon Rank Sum test: $p = 0.0093$.

(**L**) Percentage of total mitochondrial volume localized within the perinuclear region (<10 µm radius from center soma point). Student's t-test: $p < 0.0001$. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, [n.s.] $p > 0.05$: not significant.

Figure 4. Sex-specific microglial response in $\mu G^{mito}UCP2^{KO}$ after ONC

Principle component analysis (PCA) of microglia from μG^{mito} and $\mu G^{mito}UCP2^{KO}$ condition in naïve (**A-C**) and 5 days after ONC (**D-F**). The first two principle components (PC) are visualized. Each grey dot represents a single microglial profile defined by the following seven parameters: microglial Sholl index, CD68 volume, CD68 localization, median mitochondrial volume, number of mitochondria, % mitochondrial volume per microglia, % mitochondrial volume perinuclear obtained from aforementioned condition and environment. PC1 and PC2 describe 45% and 22% of the explained variance across the population, respectively. Each subpanel highlights the following comparisons: (**A**) Naïve microglial profiles for μG^{mito} (grey) and $\mu G^{mito}UCP2^{KO}$ (sand). (**B-C**) Naïve microglial profiles in male (green) and female (magenta) for μG^{mito} (**B**) and in sand for $\mu G^{mito}UCP2^{KO}$ (**C**). Dashed outline: Reference of distribution shown in (**A**). (**D**) ONC microglial profiles for μG^{mito} (blue) and $\mu G^{mito}UCP2^{KO}$ (gold). (**E**) Naïve microglial profiles in male (green) and female (magenta) for μG^{mito} (**E**) and $\mu G^{mito}UCP2^{KO}$ (**F**). Dashed outline: Reference of distribution shown in (**D**).

Figure 5. Male $\mu G^{mito}UCP2^{KO}$ microglia exhibit hyperfused mitochondrial network after ONC

(**A-C**) Principle component analysis (PCA) of microglia using mitochondrial parameters from μG^{mito} (blue) and $\mu G^{mito}UCP2^{KO}$ (gold) in ONC environment. The first two principle components (PC) are visualized. Each grey dot represents a single microglial profile defined by the following four parameters: median mitochondrial volume, number of mitochondria, % mitochondrial volume per microglia, % mitochondrial volume perinuclear. PC1 and PC2 describe 46% and 33% of the explained variance across the population, respectively. Each subpanel highlights the following comparisons: (**A**) ONC microglial profiles for μG^{mito} (blue) and $\mu G^{mito}UCP2^{KO}$ (gold). (**B-C**) ONC microglial profiles in male (green) and female (magenta) for μG^{mito} (**B**) and for $\mu G^{mito}UCP2^{KO}$. Dashed outline: Reference of distribution shown in (**A**). (**D-E**) Representative 3D-surface renderings of IBA1 (magenta outline) and mitochondria (statistics-based volume spectral coloring) from ONC-induced μG^{mito} (top) and $\mu G^{mito}UCP2^{KO}$ (bottom) in male (**D**) or female (**E**) microglia. Stripplot of the entire population of mitochondrial organelles in each cell for all conditions and environments. Each dot represents one organelle. Each hue: One microglia within a condition. Volume spectrum: $0 \ \mu m^3$ (blue) – $10 \ \mu m^3$ (red).

Figure 6. Sex-specific, transient microglial stress response affects RGC susceptibility in UCP2^{KO} mice

(**A**) Principle component analysis (PCA) of microglia using mitochondrial parameters from µG^{mito} and µG^{mito}UCP2^{KO} in ONC condition. The first two principle components (PC) are visualized. Each dot (grey) represents a single microglial profile defined by the following four mitochondrial parameters: median mitochondrial volume, number of mitochondria, % mitochondrial volume per microglia, % mitochondrial volume perinuclear. PC1 and PC2 describe 41% and 32% of the explained variance across the population, respectively. Each subpanel highlights the following comparisons: (**A**) ONC microglia profiles for male (green) and female (magenta) at 5 days (open circles) and 14 days (filled circles) after injury in ONC microglia profiles for µG^{mito}UCP2^{KO} (left, gold) or µG^{mito} (right, blue). (**B**) Stripplot of all mitochondrial organelles in each cell for all conditions and environments. Each dot represents one organelle. Each hue: separate microglia within a condition. (**C**) Schematic of genes evaluated for RT-qPCR analysis. OXPHOS: oxidative phosphorylation. (**D-F**) Principle component analysis (PCA) of microglia using metabolic and stress genes from µG^{mito} and µG^{mito}UCP2^{KO} conditions in naïve or ONC environments from male (top) or female (bottom). The first two principle components (PC) are visualized. Each dot (filled diamond: male, filled circle: female, grey) represents 100 FACSorted microglia from one retina defined by the ΔCq values of *Nrf1, Pgc1-α, Hif1-α, Sod1, Cd68*. PC1 and PC2 describe 42% and 25% of the explained variance across the population, respectively. An enlarged summary PC point for each condition and environment is overlaid on the plot, with green (male) or magenta (female) arrows depicting the overall shift between the following highlighted conditions compared in each subpanel: (**D-F**) Male (top, filled diamond) and female (bottom, filled circle) profiles for naive µG^{mito} (grey) or µG^{mito}UCP2^{KO} (sand) (**D**), naive µG^{mito} (grey) or ONC µG^{mito} (blue) (**E**), and µG^{mito}UCP2^{KO} (sand) or ONC µG^{mito}UCP2^{KO} (gold) (**F**). (**G-H**) Percentage of retinal ganglion cells (RGC) remaining (% of naïve control) after ONC in µG^{mito} at 5 days (blue) or 14 days (dark blue) and µG^{mito}UCP2^{KO} at 5 days (gold) or 14 days (orange) in for male (**G**) or female (**H**). n, number of retinas. **G**, ♂µG$_{naïve}$, n = 5; day 5, n = 4; day 14, n = 3. ♂UCP2$^{KO}_{naïve}$, n = 6; day 5, n = 4; day 14, n = 3. One-way ANOVA, $p <$ 0.0001. Selected Tukey's post-hoc comparison: Day 5 ONC ♂µG^{mito} *vs.* ♂µG^{mito}UCP2^{KO} $p =$ 0.0081; Day 14 ONC ♂µG^{mito} *vs.* ♂µG^{mito}UCP2^{KO} $p =$ 0.9000. **H**, ♀µG$_{naïve}$, n = 6; day 5, n = 4; day 14 = 2. ♀UCP2$^{KO}_{naïve}$, n = 6; day 5, n = 4: day 14 = 2 retinas. Kruskal-Wallis, $p =$ 0.044. Selected Dunn's post-hoc comparison with Bonferonni p-adjustment: Day 5 ONC

♀μG^{mito} *vs.* ♀μG^{mito}UCP2^{KO} $p = 1.000$; Day 14 ONC ♀μG^{mito} *vs.* ♀μG^{mito}UCP2^{KO} $p = 1.000$.

** $p < 0.01$. $^{n.s.}$ $p > 0.05$: not significant

Figure S1. Selective expression and knockout of UCP2 in retinal microglia

(**A-B**) Relative fold change expression level of mitochondrial-associated genes (**A**) and *Ucp2*

(**B**) among retinal cell types. Plotted from available data in *Siegert et. al.* (Siegert et al.,

2012). Purple and blue: high and low normalized expression value, respectively. PR:

photoreceptors, H: horizontal cells, BP: bipolar cells, Am: amacrine cells, RGC: retinal

ganglion cells, μG: microglia. (**C**) Timeline for tamoxifen administration to induce mito-

Dendra2 expression in microglia used for FACSorting. FACS: fluorescence activated cell

sorting. (**D**) Relative *Ucp2* transcript expression from tamoxifen-induced μG^{mito} or

μG^{mito}UCP2^{KO} FACSorted- microglia from the retina. Wilcoxon rank sum test: $p = 0.0004$.

*** $p < 0.001$. (**E**) Experimental timeline for primary murine mixed glia cultures to obtain

microglia-enriched cultures for protein analysis. P0-P3: postnatal day 0 to 3. 4-OHT: 4-

hydroxytamoxifen. Next: IBA1-immunostained mixed glial culture at day 10 before

trypsination. Scale bar: 50 μm. (**F**) Western blot analysis of UCP2 protein expression (top) or

β-actin loading control (bottom) in primary microglia of either *Cx3cr1*$^{CreERT2/+}$ or

Cx3cr1$^{CreERT2/+}$/UCP2$^{fl/fl}$ vehicle or 4-OHT (n = 1). kDa, kilodaltons. (**G**) Quantification of

UCP2 band intensity.

Figure S2. Time course of optic nerve crush (ONC)-induced retinal ganglion cell death

and microglial response in C57BL6/J mice

(**A**) Representative retinal ganglion cell distribution in wholemount retina immunostained for

RBPMS (RNA binding protein with multiple splicing) at naïve, or 1, 5, 14 days after ONC.

Scale bar: 30 μm. (**B-I**) Response of microglia after ONC in the inner (IPL, **B-E**) and the

outer plexiform layer (OPL, **F-I**). (**B, F**) Representative images of immunostained microglia

(magenta, IBA1) in naïve, or 1, 5, and 14 days after either ONC (top) or sham (bottom)

surgery. Scale bar: 30 μm. (**C, G**) Lineplots of the mean number of Sholl intersections at

increasing radii from the microglia soma (in μm). 95% confidence interval band. Naïve: grey.

Sham: green. ONC: blue. n = cells/retina. **C**, IPL: naïve, n = 161/8; ONC day 1, n = 26/3;

ONC day 5, n = 152/4; ONC day 14, n = 90/4; Sham day 1, n = 58/2; Sham day 5, n = 44/4;

Sham day 14, n = 15/2. **G**, OPL: naïve, n= 159/8; ONC day 1, n = 29/2; ONC day 5, n =

47/3; ONC day 14, n = 66/4; Sham day 1, n = 45/4; Sham day 5, n = 42/3; Sham day 14, n =

12/2 retinas. (**D, H**) Representative images of CD68 immunostaining (white) in microglia

from naïve or 5 days after ONC highlighted with dashed, colored borders in (**B**, **F**). Arrows:

CD68 labeling. Scale bar: 10 μm. (**E**, **I**) Percent CD68 volume in total microglia volume.

Naïve: grey. Sham: green. ONC: blue. Boxplot minimum and maximum: InterQuartile Range

(IQR) around median (center line). Whiskers: 1.5 IQRs. Black diamond: outliers outside of

1.5 IQRs. Filled circle: one image. n= 3-4 retinas, 1 image per retina. **E**, IPL: Naïve *vs.* ONC

Day 5, Student's t-test: $p = 0.0353$. Naïve *vs.* Sham Day 5, Student's t-test: $p = 0.9438$. **I**,

OPL: One-way ANOVA: $p = 0.6282$. * $p < 0.05$. [n.s.] $p > 0.05$: not significant.

Figure S3. Individual mitochondrial parameters and PC loadings

(**A**) Table of principal component (PC) loadings using mitochondrial, CD68, and Sholl index

parameters from **Figure 4**. Orange outline: Highest loadings for PC1. Green outline: Highest

loadings for PC2. (**B-D**) Principle component analysis (PCA) of microglia using Sholl index

and CD68 parameters from μG^{mito} (blue) and $\mu G^{mito}UCP2^{KO}$ (gold) in ONC environment. The

first two principle components (PC) are visualized. Each dot (grey) represents a single

microglial profile defined by the following three parameters: microglial Sholl index, CD68

volume, CD68 localization. PC1 and PC2 describe 58% and 29% of the explained variance

across the population, respectively. Each subpanel highlights the following comparisons:

(**B**) ONC microglial profiles for μG^{mito} (blue) and $\mu G^{mito}UCP2^{KO}$ (gold). (**C-D**) ONC

microglial profiles in male (green) and female (magenta) for μG^{mito} (**C**) and $\mu G^{mito}UCP2^{KO}$

(**D**) Dashed outline: reference of distribution shown in (**B**). (**E**) Table of principal component

(PC) loadings from PCA using four mitochondrial parameters (median mitochondrial

volume, number of mitochondria, % mitochondrial volume per microglia, % mitochondrial

volume perinuclear) from **Figure 5A-C**. Green outline: Highest PC loading in PC2.

(**F-H**) Mitochondrial parameters. Boxplot minimum and maximum: InterQuartile Range

(IQR) around median (center line). Whiskers: 1.5 IQRs. Black diamond: outliers outside 1.5

IQRs. Each overlaid point represents data point of a single microglia. n = cells/retina.

♂$\mu G^{mito}_{naïve}$ (grey), n = 11/2; ♀$\mu G^{mito}_{naïve}$ (grey), n = 7/3; ♂μG^{mito}_{ONC} (blue), n = 7/4;

♀μG^{mito}_{ONC} (blue), n = 7/2; ♂$\mu G^{mito}UCP2^{KO}_{naïve}$ (sand), n = 10/2; ♀$\mu G^{mito}UCP2^{KO}_{naïve}$ (sand),

n = 11/2; ♂$\mu G^{mito}UCP2^{KO}_{ONC}$ (gold), n = 10/2; ♀$\mu G^{mito}UCP2^{KO}_{ONC}$ (gold), n = 12/2.

(**F**) Mitochondrial network connectivity determined by median mitochondrial volume of all

organelles (left, one-way ANOVA: $p < 0.0001$, Selected Tukey's post-hoc comparisons:

♂$\mu G^{mito}UCP2^{KO}_{naïve}$ *vs.* ♂$\mu G^{mito}UCP2^{KO}_{ONC}$, $p = 0.4917$; ♀$\mu G^{mito}UCP2^{KO}_{naïve}$ *vs.*

♀$\mu G^{mito}UCP2^{KO}_{ONC}$, $p = 0.0094$) and number of organelles within a single cell (right,

Kruskal-wallis test: $p = 0.0004$. Selected Dunn's post-hoc comparisons with Bonferonni p-

adjustment: $\male\mu$G^{mito}UCP2$^{KO}_{naïve}$ *vs.* $\male\mu$G^{mito}UCP2$^{KO}_{ONC}$, $p = 1.0$; $\female\mu$G^{mito}UCP2$^{KO}_{naïve}$ *vs.*

$\female\mu$G^{mito}UCP2$^{KO}_{ONC}$, $p = 1.0$). (**G**) Percentage of mitochondrial volume per microglial

volume. Kruskal-wallis test: $p = 0.0006$. Selected Dunn's post-hoc comparisons with

Bonferonni p-adjustment: $\male\mu$G$^{mito}_{naïve}$ *vs.* $\male\mu$G^{mito}UCP2$^{KO}_{ONC}$, $p = 0.0011$;

$\female\mu$G^{mito}UCP2$^{KO}_{navie}$ *vs.* $\male\mu$G^{mito}UCP2$^{KO}_{ONC}$, $p = 0.0005$. (**H**) Percentage of total

mitochondrial volume localized within the perinuclear region (10 μm radius from center

soma point). One-way ANOVA: $p < 0.0001$. Selected Tukey's post-hoc comparisons:

$\male\mu$G$^{mito}_{naive}$ *vs.* $\male\mu$G$^{mito}_{ONC}$ $p = 0.0032$. ** $p < 0.01$, *** $p < 0.001$, [n.s.] $p > 0.05$: not

significant. For other post-hoc comparisons see **Table S3**.

Figure S4. PCA comparison for naïve environment and relative transcript expression

(**A**) PCA from **Figure 6A** highlighting comparison between naïve conditions (left) for in

μG^{mito} (grey) or μG^{mito}UCP2^{KO} (sand) and days 5 and 14 after ONC condition (right) in

μG^{mito} (blue) or μG^{mito}UCP2^{KO} (gold). (**B**) Table of principal component (PC) loadings from

PCA using the metabolic (*Nrf1, Pgc1α, Hif1α*) and stress (*Sod1, Cd68*) genes in **Figure 6**.

(**C**) PCA from **Figure 6** highlighting comparison of male (filled diamond) and female (filled

circle) profiles for naïve μG^{mito} (left, grey) or naïve μG^{mito}UCP2^{KO} (right, sand). (**D-E**)

Boxplot minimum and maximum: InterQuartile Range (IQR) around median (center line).

Whiskers: 1.5 IQRs. Black diamond points: outliers outside 1.5 IQRs. Each overlaid point

represents data point of a single microglia. Filled circle: female, filled diamond: male. n,

number of retinas. μG$^{mito}_{naïve}$: n= $\male$9, $\female$9; μG$^{mito}_{ONC}$: n = $\male$4, $\female$3; μG^{mito}UCP2$^{KO}_{naïve}$: n = $\male$3,

$\female$4; μG^{mito}UCP2$^{KO}_{ONC}$: n = $\male$3, $\female$5. (**D-E**) Fold change of *Pgc1α, Nrf1, Hif1α* in conditions

and environments for males (**D**) and females (**E**) normalized to their corresponding naïve

μG^{mito}. $\male$*Pgc1α*: Kruskal-Wallis test, $p = 0.0075$ with shown selected Dunn's post-hoc

comparison with Bonferonni p-adjustment: μG$^{mito}_{naive}$ *vs.* μG$^{mito}_{ONC}$ $p = 0.0393$; μG$^{mito}_{naive}$ *vs.*

μG^{mito}UCP2$^{KO}_{ONC}$ $p = 0.0322$. $\male$*Nrf1*: Kruskal-Wallis test, $p = 0.0078$ with shown selected

Dunn's post-hoc comparison with Bonferonni p-adjustment: μG$^{mito}_{naive}$ *vs.* μG$^{mito}_{ONC}$ $p =$

0.0071. $\male$*Hif1α*: One-way ANOVA, $p = 0.2400$. $\female$*Pgc1α*: Kruskal-Wallis test, $p = 0.0706$.

$\female$*Nrf1*: Kruskal-Wallis test, $p = 0.1620$. $\female$*Hif1-α*: One-way ANOVA, $p = 0.2323$. (**F**) Fold

change of *Sod1* (left) and *Cd68* (right) across sexes. Left, *Sod1*: One-way ANOVA, $p =$

0.0181 with selected Tukey's post-hoc comparison: μG$^{mito}_{naive}$ *vs.* μG^{mito}UCP2$^{KO}_{ONC}$ $p =$

0.0398. Right, *Cd68*: Kruskal-Wallis test: $p = 0.0001$ with selected Dunn's post-hoc

comparison with Bonferonni p-adjustment: $\male\mu$G$^{mito}_{naive}$ *vs.* $\male\mu$G$^{mito}_{ONC}$ $p = 0.0205$;

$\male$ μG^{mito}_{naive} *vs.* $\male$ $\mu G^{mito}UCP2^{KO}_{ONC}$ $p = 0.0184$. * $p < 0.05$, ** $p < 0.05$, [n.s.] $p > 0.05$: not

significant. For other post-hoc comparisons see **Table S3**.

Table S1. Primer sequences.

Table S2. Antibody table.

Table S3. Statistical analysis.

<u>Materials and Methods</u>

Animals

Animal housing and procedures were approved by the "Bundesministerium für Wissenschaft,

Forschung und Wirtschaft (bmwfw) Tierversuchsgesetz 2012, BGBI. I Nr. 114/2012 (TVG

2012) under the number GZ: 2021-0.607.460. Mice were housed in a 12 hour light-dark cycle

in individually ventilated cages in a controlled environment (room temperature 22 ± 1 °C;

relative humidity 55 ± 10 %) with access to standard diet (rat/mouse maintenance diet

(V1534-300) or mouse breeding diet (V1124-300), ssniff Spezialitäten GmbH) and

autoclaved water provided *ad libitum* in the ISTA Preclinical facility. Founder mouse strains

were obtained from Jackson Laboratories and backcrossed to C57Bl6/J (#000664)

background for at least 10 generations: $Cx3cr1^{CreERT2}$ (#020940)*,* PhAM$^{fl/fl}$ (#018385),

UCP2$^{fl/fl}$ (#022394). Throughout the manuscript μG^{mito} refers to $Cx3cr1^{CreERT2/+}$/Pham$^{fl/fl}$,

while $\mu G^{mito}UCP2^{KO}$ refers to $Cx3cr1^{CreERT2/+}$/Pham$^{fl/fl}$/UCP2$^{fl/fl}$. For UCP2 protein knockout

primary microglia culture experiments (**Figure S1E-G**) and retinal ganglion cell counting

experiments (**Figure 6C**), mice did not contain the Dendra2 mitochondria label, and instead

were $Cx3cr1^{CreERT2/+}$/ and $Cx3cr1^{CreERT2/+}$/UCP2$^{fl/fl}$.

Tamoxifen administration for Cre-induced recombination

Tamoxifen (Sigma Aldrich, T5648-5G) was dissolved in corn oil (Sigma Aldrich, C8267-

500ML) at a concentration of 20 mg/mL and sonicated for 40 minutes. Freshly prepared

tamoxifen was administered via intraperitoneal injection daily (150mg/kg) for three days.

The animals did not show any abnormal behavior or discomfort after gene knockout. All

experimental groups (μG^{mito} and $\mu G^{mito}UCP2^{KO}$, naïve and ONC) were tamoxifen-injected for

two reasons; first, expression of the Dendra2 mitochondria label is only possible after Cre

recombinase excision of the stop cassette and second, $Cx3Cr1^{CreERT2}$ reporter lines have been

shown to include some 'leakiness', *i.e.* tamoxifen-independent *Cre* recombination (Van Hove

et al., 2020). *Cx3Cr1*[CreERT2] mice also express CreERT2 in monocyte populations, therefore

to ensure no interference from UCP2[KO] circulating monocytes, ONC experiments were

performed at least three weeks after tamoxifen induction when monocyte populations have

turned over (Goldmann et al., 2013). This also provided sufficient time for adaptation of

microglia to the UCP2 loss conditions and clearance of all tamoxifen metabolites (Jahn et al.,

2018).

Optic nerve crush (ONC) procedure

Mice were anesthetized in an induction chamber with 5% (v/v) isoflurane (Zoetis) supplied

with oxygen at a flow rate of 0.6 L/min. After lack of a foot pinch reflex, mice were

maintained at 2.5% (v/v) isoflurane applied through a nose cone while on a heating pad to

maintain body temperature at 37°C. Proparacaine hydrochloride 0.5% ophthalmic eye drops

(Ursapharm Arzneimittel GmbH) were applied to numb the eyes, and subcutaneous injection

of 5mg/kg Metacam alleviated pain (Meloxacam, Boehringer Ingelheim). The lateral canthus

was de-vascularized by clamping with a hemostat (Fine Science Tools) for 10 seconds. Using

a Leica dissection microscope, a lateral canthotomy allowed visualization of the posterior

pole. While firmly holding the conjunctiva with a jeweler forceps, the conjunctiva was cut

perpendicular to the posterior pole. The surrounding muscle was carefully dissected as to not

puncture the vascular plexus. The optic nerve was pinched 1mm from the posterior pole for 4

seconds using a curved N7 self-closing forceps (Dumont). Triple antibiotic ointment was

applied to the eye directly after the surgery to prevent infection.

Primary glial culture

Mixed glia culture were prepared from the published protocol by *Bronstein et al*. (Bronstein

et al., 2013). Cortices from 3-5 murine pups aged P0-P2 were dissected under a sterile

ventilation hood in ice-cold Hank's buffered saline (ThermoFisher, #14025-050) and digested

in 0.05% (v/v) Trypsin + 1X EDTA (ThermoFisher, #25300-054) for 15 minutes at 37°C.

The trypsinization was stopped with the addition of serum-containing medium (DMEM, 10%

(v/v) heat-inactivated FBS, 1% (v/v) Penicillin/Streptomycin, 1% (v/v) Non-essential amino

acids (Sigma-Aldrich, M7145-100ML)). Cells were pelleted at 500xg for 5 min, washed one

time with serum-containing medium, then passed through a 40μm cell strainer (Szabo-

Scandic, #352340). For Western blotting, the resulting cell suspension was plated onto T75

cell culture flasks (TPP, #Z707503). The culture medium was replaced at day 3 and day 10.

Primary microglia isolation

Two T75 flasks containing mixed primary glia cultures prepared in tandem were treated with 1µM 4-hydroxy tamoxifen (Sigma-Aldrich, SML1666-1ML) or vehicle at day 10. Three days later, microglia were isolated using mild trypsinization (0.8mM $CaCl_2$ in 1xTrypsin-EDTA, ThermoFisher #25300-054) is added to the flask after a washing step with pre-warmed 1X PBS to remove inhibiting serum, then incubate at 37ºC for 10-20 minutes, or until the non-microglial cells have just lifted from the flask forming a floating layer. The floating cells were collected with the supernatant, spun down at 10,000xg, washed once with 1X PBS, pelleted and snap-frozen. The microglia remaining on the flask were overlaid with 5ml 1X PBS, removed using a cell scraper, pelleted, and snap-frozen.

Western blotting

Total cellular protein isolation, production of recombinant UCP2 in *E. coli*, and western blot (WB) was performed as described previously (Rupprecht et al., 2012, 2014). For WB analyses 10 µg of total protein were loaded on the SDS-gel, 1 ng of recombinant UCP2 was used as positive control for anti-UCP2 antibody detection. To ensure blotting homogeneity to nitrocellulose membranes, Ponceau S staining was performed. To relate UCP2 expression to loading control, the membranes were stripped, re-blocked and incubated with β-actin. The affinity purified antibody against the N-terminal sequence of mouse UCP2 (VGFKATDVPPTATVKF) was designed and evaluated in the Pohl laboratory (Rupprecht et al., 2012). Peptide synthesis, rabbit immunization and affinity purification were performed by PINEDA Antibody-service GmbH (Berlin, Germany). The production of recombinant murine UCP2, used as positive control, was performed as described in (Žuna et al., 2021).

Microglia isolation and sample preparation for fluorescence-activated cell sorting (FACS)

Adult animals expressing the mito-Dendra2 tag were anesthetized with isoflurane and decapitated. Retinas were immediately explanted and dissected in 1X PBS on ice. Each retina was transferred to a 1.5ml Eppendorf low-adhesion tube filled with the 800µl digestion buffer (1:8:1 Cysteine/EDTA solution (2.5mM Cysteine, 0.5mM EDTA (ethylenediaminetetraacetic acid) in HBSS, 10mM HEPES (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid) in HBSS, and 10mg/ml Papain, Roche #10108014001), incubated at 37ºC for 10 minutes, then centrifuged for 2.5 min at 240xg. Samples were transferred on ice, the supernatant was discarded and 800µl 1mM EDTA in HBSS + 2% (v/v) FBS added. After two washes with

1mM EDTA in HBSS + 2% (v/v) FBS, the digested tissue was triturated 10-15 times with a

pulled glass pipette, then filtered through a 70 μm strainer.

Fluorescent activated cell sorting (FACS)

FACS was performed using a SONY SH800SFP equipped with a 100 μm nozzle and sorting

speed of approximately 8,000 events per second in normal purity mode. Microglia were gated

from the population by setting the forward and side scatter and by the fluorescence of the

mito-Dendra2 tag. From each retina, 100 microglia were sorted and immediately collected in

one well of a 96-well plate (Eppendorf) filled with 5 μl cold lysis buffer from the NEBNext®

Single Cell/Low Input cDNA Synthesis & Amplification Module (New England Bio Labs,

#E6421L). After the sorting, the plate was shortly spun down to ensure all cells were

collected at the bottom of the well and was immediately processed for cDNA synthesis.

cDNA synthesis

Sorted microglia were processed with the NEBNext® Single Cell/Low Input cDNA

Synthesis & Amplification Module (New England Bio Labs, #E6421L) according to the

manufacturer's protocol.

Reverse transcription quantitative real-time PCR and gene expression analysis

Primers (**Table S1**) were designed with the free PrimerQuest Tool from Integrated DNA

Technologies (https://eu.idtdna.com/PrimerQuest/Home/Index). To ensure the target

sequence of each primer, primers were blasted (https://www.ncbi.nlm.nih.gov/tools/primer-

blast/). Their self-complementarity and folding probability were investigated using the

UNAFold Web server (http://www.unafold.org/). Primer efficiencies were validated from the

slope of four to five serial 1:4 dilutions of cDNA template according to equation (1). Primers

with efficiencies between 90-110% were used.

$$(1)\ Efficiency = ((2^{-\frac{1}{Slope}}) - 1) * 100$$

For gene expression analysis, RT-qPCR (Luna® Universal qPCR Master Mix; New England

BioLabs; M3003L) was performed in 384-well plates (Bio-Rad; HSR4805) on a Roche

Lightcycler 480 according to the manufacturer's manual. Total reaction volume was 10μl

containing 1μl of 1:10 diluted cDNA as template and a final concentration of 0.25μM for

each primer. Cycle conditions were 60 seconds at 95°C for initial denaturation, followed by

40 cycles of denaturation (15 seconds; 95°C) and annealing/extension (30 seconds; 60°C).

Each run was completed with a melting curve analysis to confirm amplification of only one

amplicon. Each PCR reaction was run in triplicate from which a mean Cq value was

calculated and used for further analysis.

Analysis of RT-qPCR results

Fold change differences between each condition and μG^{mito}_{naive} were calculated according to

delta-delta Ct method (Schmittgen and Livak, 2008). dCq values were obtained by

normalizing mean Cq values to the reference housekeeping gene (GAPDH) measured within

the same experiment (Equation 2). ddCq values were then calculated by normalizing dCq

values to the respective control condition (μG^{mito}_{naive}) within each experimental repetition

(Equation 3). Fold changes were obtained by transforming ddCq values from log2-scale to

linear scale (Equation 4). These fold changes were used for data visualization. Exclusion

criteria were based on IBA1 and GFAP fold change expression to determine purity of cell

isolation, which would indicate substantial contamination from astrocytes and müller glia.

$$(2)\ dCq = Cq_{\text{reference gene}} - Cq_{\text{gene of interest}}$$

$$(3)\ ddCq = dCq - dCq_{\text{control condition}}$$

$$(4)\ Fold\ change = 2^{\wedge(ddCq)}$$

Retina dissection and fixation

Animals were briefly anesthetized with isoflurane (Zoetis) until breathing slowed, then

cervical dislocation was performed. Retinas were immediately dissected from the enucleated

eyes in 1X phosphate buffered saline (PBS) and fixed in 4% (w/v) paraformaldehyde for 30

minutes. After 3x PBS washes, retinas were put in 30% (w/v) sucrose in 1X PBS overnight at

4°C for either -80°C freezer storage or immunostaining.

Immunostaining

Staining was carried out in 24-well tissue culture plates (TPP, #92024). Retinas saturated in

sucrose (30% sucrose (w/v) in 1X PBS) underwent three freeze/thaw cycles, then were

washed 3x with 1X PBS. Next, a blocking step using blocking solution (1% (w/v) bovine

serum albumin (Sigma A9418), 5% (v/v) Triton X-100 (Sigma T8787), 0.01% (v/v) sodium

azide (VWR 786-299), and 10% (v/v) goat (Millipore S26) or donkey (Millipore S30) serum) was carried out on an orbital shaker for 1 hour at room temperature. Primary antibodies (**Table S2**) were diluted in antibody solution (1% (w/v) bovine serum albumin, 5% (v/v) triton X-100, 0.01% (v/v) sodium azide, 3% (v/v) goat or donkey serum) and incubated for three nights at 4°C on an orbital shaker. Following 3x PBS washes for 30 mins each, secondary solutions were prepared in the antibody solution and incubated at room temperature for 2 hours. After three washes, Hoechst 334422 (1:5,000, Thermo Fisher Scientific, H3570) for 10 mins, and three more 1X PBS washes. Retinas were mounted using an antifade solution (10% (v/v) mowiol (Sigma, 81381), 26% (v/v) glycerol (Sigma, G7757), 0.2M tris buffer pH 8, 2.5% (w/v) Dabco (Sigma, D27802)) on glass microscope slides with parafilm spacers to prevent tissue distortion and overlaid with coverslips (VWR, #631-0147).

Confocal microscopy

Single cell images for deconvolution were acquired on Zeiss LSM800 microscope using a Plan-Apochromat 40X objective NA 1.4 (#420762-9900) at 1.5x digital zoom to acquire a pixel size of 0.05 μm in X and Y resolution and 0.150 μm in Z. Images were acquired from the peripheral region of the retina, in two opposing quadrants, and inner plexiform layer microglia were selected by setting the z-stack between the ganglion cell and inner nuclear layer Hoechst staining. Microglia in the ganglion cell layer were excluded due to their low population in naïve environment, and known migration from the optic nerve head to the nerve fiber layer after ONC (Heuss et al., 2018). After IPL selection, the IBA1 channel was visualized and the nearest IBA1$^+$ cell was selected for imaging, thereby reducing bias in cell selection and blinded to the mitochondrial architecture within the cell. Tile scans (2x2) were necessary to acquire the area of single IPL microglia in naïve condition, resulting in images of dimension 202 x 202 μm. For each animal, 2-3 images for from two opposing quadrants were acquired. Images were stitched using Zen 2.3 or 3.5 desktop version.

Tiled (2x2) images for C57Bl6/J timecourse ONC experiments (Figure S2) were acquired using a Plan-Apochromat 40X objective NA 1.4 (#420762-9900) with pixel size of 0.156 μm in X and Y and 0.21 μm in Z plane for a region measuring 303 x303 μm. Tiled images (2x2) for microglia and retinal ganglion cell counting were acquired using a Plan-Apochromat 20X objective NA 0.8 (#420650-9901) with pixel resolution 0.312 x 0.312 x 0.5 μm to obtain a region measuring 605 x 605 μm in four peripheral quadrants of each retinal wholemount.

Image processing

SVI Huygens Professional v21.10 (https://svi.nl/HomePage) was used for batch

deconvolution processing of stitched confocal images. Microscope parameters for batch

settings were as follows: 50 nm sampling intervals for X and Y, 150 nm in Z. The refractive

index and embedding medium were set to 1.515 and 1.49, respectively. Objective quality was

listed as 'good.' For channel settings, the back projected pinhole was set to 250 nm, and the

excitation fill factor was 2. Excitation (ex) and emission (em) settings for the four channels

were as follows: 405/420 nm, 488/520 nm, 561/602 nm, 633/650 nm. Deconvolution

parameters were the default settings for the 'confocal low template' for deconvolution using

the classic MLE (maximum likelihood estimation) algorithm with maximum 30 iterations.

Images were output in the Imaris file format.

Filament tracing

Microglia morphology was traced in 3-dimensions using the *Filament* wizard in IMARIS.

The starting point was detected with diameter 12 µm and 0.6 µm was used to set seeding

points and the disconnected segment filtering smoothness. Final filament traces were

manually edited to remove incorrect segments from the semi-automatic *Filament* tracing.

These settings were used for both single cell traces and C57Bl6/J traces from multiple cells in

a larger field of view.

CD68 volume rendering for multiple cells in field of view

Quantification of CD68 volume per microglia in Figure S2 was performed using Imaris v9.3

using the *Surface* wizard. 3d-surface renderings were generated for the CD68 channel using

surface detail 0.2 µm and IBA1 channel using 0.5 µm. The *surface-surface coloc* plugin was

used to determine the CD68 surfaces within microglia IBA1 surface. The co-localized CD68

surfaces total volume was extracted and expressed as a ratio of the total microglia volume.

The volumes and ratios were calculated from the entire image.

Mitochondria and CD68 surfaces in single microglia

Surface rendering analysis was performed using IMARIS v9.1-9.3 (Bitplane, Imaris).

Individual cells were cropped from the original tiled image when necessary to reduce

processing time. Then, 3-dimensional surfaces were created for microglia, mitochondria and

CD68 vesicles using the *Surface* wizard with surface details settings of 0.5 µm, 0.2 µm, and

0.4 µm, respectively. Each surface was manually edited to resolve any inconsistencies in

surface creation. Mitochondrial surfaces were carefully edited to verify connectivity

throughout the entirety of the network, often requiring manual splitting of surfaces which

were joined by the thresholding within the wizard.

Mitochondrial organelle and CD68 vesicle localization in single microglia

Completed *Surfaces* for either mitochondria or CD68 were 'split' using the Xtension *Split*

Surfaces available via Imaris bitplane (https://imaris.oxinst.com/open/). This generates a *Split*

Surface group where each individual organelle/vesicle becomes a new uniquely named

surface nested within the *Split Surface* group. A *spot* 5 µm in diameter was manually added

to identify the nucleus. Then, the *Spots and Surfaces distance* XTension calculates the

distance between the nucleus *spot* and the center point of each individual *surface* in the *Split*

surface group. The Xtension was modified from the original version to allow for up to 100

distance calculations and to output a *csv* file.

Data compilation and metric calculations

Python version 3.7 with packages *pandas* (Mckinney, 2010), *matplotlib* (Hunter, 2007) and

seaborn (Waskom, 2021) for data compilation and visualization. From Imaris, mitochondria,

microglia, and CD68 surfaces, microglia filament tracings, and mitochondrial and CD68

localization *Statistics* files for each cell were exported to comma separated values file format

with a unique identifier for ease in compiling cell-specific data for downstream analysis.

Metric calculations for each surface type are detailed below:

<u>Filament tracing</u>

The 'Number of Filament Sholl intersections' from the *Statistics* file was used to plot the

number of intersections per increasing radii of 1 µm (Sholl, 1953). Plotted Sholl curves were

truncated at 60 µm.

<u>Median mitochondrial volume and number</u>

From the *Statistics* file, the volume of each mitochondrial *Surface* within a cell was used to

determine the median mitochondrial volume per cell. Here, the total number of organelles per

cell was also be extracted.

<u>Percentage of CD68 volume</u>

From the *Statistics* file, the volume from each CD68 *Surface* within a cell was summed.

Additionally, the sum volume of the microglia *Surface* was extracted from the microglia

Statistics file. The percentage of CD68 volume was calculated from total CD68 volume per

total microglia volume for each cell.

<u>Percentage of mitochondrial volume</u>

From the *Statistics* file, the volume from each mitochondrial *Surface* within a cell was

summed. Additionally, the sum volume of the microglia *Surface* was extracted from the

microglia *Statistics* file. The percentage of mitochondrial volume was calculated from total

mitochondria volume per total microglia volume for each cell.

<u>Mitochondrial perinuclear localization</u>

The *Statistics* file for localization data includes the distance of each organelle from the cell

soma (µm) and a unique *Surface ID* that is derived from the original mitochondrial *Surface*.

From the *Statistics* file of mitochondrial volume, the *Surface ID* can be used to compile

distance and volume for each unique organelle within the cell.

The percentage of mitochondrial volume in the perinuclear region was determined by

calculating the sum of organelle volumes within a 10 µm radius of the cell soma, and

dividing by the total mitochondria volume in that cell. The microglia soma ranges from 5-10

µm in diameter due to its non-uniform shape, therefore a 10 µm radius from the center of the

soma *Spot* in IMARIS includes ~2.5-5 µm of soma, allowing an estimation of mitochondrial

localized in the perinuclear space to be at maximum 5-7 µm from the nuclear membrane.

We corroborated this perinuclear boundary distance by principal component analysis of the

mitochondrial distances. Mitochondrial distances were binned in increments of 10 µm, and

the sum volume in each bin was normalized to total mitochondria volume within a cell. PC

loadings of the normalized values confirmed that the two bins contributing the most to

principal component loadings for distances between 0-10 µm, and 71-80 µm.

<u>CD68 peripheral localization</u>

CD68 localization data was determined in a similar manner, except that the percentage of

vesicles outside the perinuclear region were reported as the sum volume of these organelles

divided by the total volume of vesicles in the cell. PCA of binned and normalized CD68

volumes resulted in greatest PC loadings in the 0-10 µm bin.

Identifying Sholl radii for PCA analysis

The Sholl curve (see Method: <u>Filament tracing</u>) is a high-dimensional, vector representation of the microglial morphology. To incorporate filament tracing into the PCA without overpowering the other scalar parameters, we first determined which Sholl radius contributed most to the differences in Sholl intersection curves. Here, Sholl intersections with 10 μm radial distances were normalized for each cell. We observed the greatest PC loadings in the 11-20 μm bin. We repeated this analysis with 5 μm radial distances and confirmed the greatest PC loadings in 11-15 μm bin. As the PC loading was greater for the 10 μm radial increments, we implemented the normalized ratio of Sholl intersections at 11-20 μm for each cell and refer to this as the Sholl index.

Principal Component Analysis (PCA)

PCA analysis was performed using scikit-learn in Python (Pedregosa et al., 2011). Prior to PCA, a correlation analysis was performed to determine whether any of the individual microglial or mitochondrial metrics derived from the same surface were highly correlated with one another. PCA was then implemented on the metrics or groups of metrics as indicated in Figure legends. In Figure 4, the following seven metrics were used from mitochondria and microglia analysis to serve as components for PCA including; median mitochondria volume, number of mitochondria, percentage of mitochondrial volume per microglia and percentage of CD68 volume per microglia volume were included as mean values for each cell. Mitochondria and CD68 localization values were included as percentage values. Figure 5 used only mitochondrial parameters, while Figure S4 used the Sholl index and CD68 parameters as denoted within the text. In Figure 6, PCA was performed using dCq values of metabolic and stress genes. If a transcript was undetected, a mean Cq value was of 40 was used.

Cell counting

RBPMS⁺-cells were counted using the *Spots* function of Imaris v9.3, with additional manual editing. Each of these metrics was quantified from four peripheral quadrants of each analyzed retina and represented as a median value.

Statistics

Statistical tests were performed using the Python module *statsmodels* (Seabold and Perktold, 2010) as indicated in the figure legends. Groups for comparison were first tested for normal

distribution and equal variance using the Shapiro test and Levene test, respectively. Where

applicable, post-hoc tests were performed using either Tukey's multiple comparison of means

or Dunn's test in Python *statsmodels* 'multicomp'. Significance levels are indicated using the

following notation ($^{n.s}$ $p > 0.05$; *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$). For Sholl analysis

comparisons, linear regression models from *lme4* package (Bates et al., 2015) were

performed in R. The default contrast for unordered variables was changed to 'contr.sum',

allowing type III Anova comparison. Statistical analysis for all figure panels are found in

Table S3.

<u>**References**</u>

Al-Mehdi, A.B., Pastukh, V.M., Swiger, B.M., Reed, D.J., Patel, M.R., Bardwell, G.C.,

Pastukh, V. V., Alexeyev, M.F., and Gillespie, M.N. (2012). Perinuclear mitochondrial

clustering creates an oxidant-rich nuclear domain required for hypoxia-induced transcription.

Sci. Signal. *5*.

Arsenijevic, D., Onuma, H., Pecqueur, C., Raimbault, S., Manning, B.S., Miroux, B.,

Couplan, E., Alves-Guerra, M.-C., Goubern, M., Surwit, R., et al. (2000). Disruption of the

uncoupling protein-2 gene in mice reveals a role in immunity and reactive oxygen species

production. Nat. Genet. *26*, 435–439.

Baik, S.H., Kang, S., Lee, W., Choi, H., Chung, S., Kim, J. Il, and Mook-Jung, I. (2019). A

Breakdown in Metabolic Reprogramming Causes Microglia Dysfunction in Alzheimer's

Disease. Cell Metab. *30*, 493-507.e6.

Baker, A.E., Brautigam, V.M., and Watters, J.J. (2004). Estrogen Modulates Microglial

Inflammatory Mediator Production via Interactions with Estrogen Receptor β. Endocrinology

145, 5021–5032.

Bates, D., Mächler, M., Bolker, B.M., and Walker, S.C. (2015). Fitting Linear Mixed-Effects

Models Using lme4. J. Stat. Softw. *67*, 1–48.

Bauer, J., Sminia, T., Wouterlood, F.G., and Dijkstra, C.D. (1994). Phagocytic activity of

macrophages and microglial cells during the course of acute and chronic relapsing

experimental autoimmune encephalomyelitis. J. Neurosci. Res. *38*, 365–375.

Bernier, L.P., York, E.M., and MacVicar, B.A. (2020). Immunometabolism in the Brain:

How Metabolism Shapes Microglial Function. Trends Neurosci. *43*, 854–869.

Borst, K., Schwabenland, M., and Prinz, M. (2019). Microglia metabolism in health and

disease. Neurochem. Int. *130*, 104331.

Bronstein, R., Torres, L., Nissen, J.C., and Tsirka, S.E. (2013). Culturing microglia from the

neonatal and adult central nervous system. J. Vis. Exp.

Buck, M.D., O'Sullivan, D., Klein Geltink, R.I., Curtis, J.D., Chang, C.-H., Sanin, D.E., Qiu,

J., Kretz, O., Braas, D., van der Windt, G.J.W., et al. (2016). Mitochondrial Dynamics

Controls T Cell Fate through Metabolic Programming. Cell *166*, 63–76.

Colombo, G., Cubero, R.J.A., Kanari, L., Venturino, A., Schulz, R., Scolamiero, M.,

Agerberg, J., Mathys, H., Tsai, L.-H., Chachólski, W., et al. (2021). Microglial MorphOMICs

unravel region- and sex-dependent morphological phenotypes from postnatal development to

degeneration. BioRxiv 2021.11.30.470610.

Damoiseaux, J.G., Döpp, E.A., Calame, W., Chao, D., MacPherson, G.G., and Dijkstra, C.D.

(1994). Rat macrophage lysosomal membrane antigen recognized by monoclonal antibody

ED1. Immunology *83*, 140–147.

Das, R., and Chakrabarti, O. (2020). Mitochondrial hyperfusion: a friend or a foe. Biochem.

Soc. Trans. *48*, 631–644.

Erny, D., Dokalis, N., Mezö, C., Castoldi, A., Mossad, O., Staszewski, O., Frosch, M., Villa,

M., Fuchs, V., Mayer, A., et al. (2021). Microbiota-derived acetate enables the metabolic

fitness of the brain innate immune system during health and disease. Cell Metab. *33*, 2260-

2276.e7.

Faas, M., Ipseiz, N., Ackermann, J., Culemann, S., Grüneboom, A., Schröder, F., Rothe, T.,

Scholtysek, C., Eberhardt, M., Böttcher, M., et al. (2021). IL-33-induced metabolic

reprogramming controls the differentiation of alternatively activated macrophages and the

resolution of inflammation. Immunity.

Finck, B.N., and Kelly, D.P. (2006). PGC-1 coactivators: inducible regulators of energy

metabolism in health and disease. J. Clin. Invest. *116*, 615.

Galatro, T.F., Holtman, I.R., Lerario, A.M., Vainchtein, I.D., Brouwer, N., Sola, P.R., Veras,

M.M., Pereira, T.F., Leite, R.E.P., M?ller, T., et al. (2017). Transcriptomic analysis of

purified human cortical microglia reveals age-associated changes. Nat. Neurosci.

Galindo-Romero, C., Valiente-Soriano, F.J., Jiménez-López, M., García-Ayuso, D., Villegas-

Pérez, M.P., Vidal-Sanz, M., and Agudo-Barriuso, M. (2013). Effect of Brain-Derived

Neurotrophic Factor on Mouse Axotomized Retinal Ganglion Cells and Phagocytic

Microglia. Invest. Ophthalmol. Vis. Sci. *54*, 974–985.

Goldmann, T., Wieghofer, P., Müller, P.F., Wolf, Y., Varol, D., Yona, S., Brendecke, S.M.,

Kierdorf, K., Staszewski, O., Datta, M., et al. (2013). A new type of microglia gene targeting

shows TAK1 to be pivotal in CNS autoimmune inflammation. Nat. Neurosci. 2013 1611 *16*, 1618–1626.

Gomes, L.C., Benedetto, G. Di, and Scorrano, L. (2011). During autophagy mitochondria elongate, are spared from degradation and sustain cell viability. Nat. Cell Biol. 2011 135 *13*, 589–598.

Gosselin, D., Skola, D., Coufal, N.G., Holtman, I.R., Schlachetzki, J.C.M.M., Sajti, E., Jaeger, B.N., O'Connor, C., Fitzpatrick, C., Pasillas, M.P., et al. (2017). An environment-dependent transcriptional network specifies human microglia identity. Science (80-.). *356*, 1248–1259.

Hammond, T.R., Dufort, C., Dissing-Olesen, L., Giera, S., Young, A., Wysoker, A., Walker, A.J., Gergits, F., Segel, M., Nemesh, J., et al. (2019). Single-Cell RNA Sequencing of Microglia throughout the Mouse Lifespan and in the Injured Brain Reveals Complex Cell-State Changes. Immunity.

Hanisch, U.-K., and Kettenmann, H. (2007). Microglia: active sensor and versatile effector cells in the normal and pathologic brain. Nat. Neurosci. *10*, 1387–1394.

He, D., Xu, H., Zhang, H., Tang, R., Lan, Y., Xing, R., Li, S., Christian, E., Hou, Y., Lorello, P., et al. (2022). Disruption of the IL-33-ST2-AKT signaling axis impairs neurodevelopment by inhibiting microglial metabolic adaptation and phagocytic function. Immunity *55*, 159-173.e9.

Heuss, N.D., Pierson, M.J., Roehrich, H., McPherson, S.W., Gram, A.L., Li, L., and Gregerson, D.S. (2018). Optic nerve as a source of activated retinal microglia post-injury. Acta Neuropathol. Commun. *6*, 66.

Hilla, A.M., Diekmann, H., and Fischer, D. (2017). Microglia Are Irrelevant for Neuronal Degeneration and Axon Regeneration after Acute Injury. J. Neurosci. *37*, 6113–6124.

Hoang, T., Wang, J., Boyd, P., Wang, F., Santiago, C., Jiang, L., Yoo, S., Lahne, M., Todd, L.J., Jia, M., et al. (2020). Gene regulatory networks controlling vertebrate retinal regeneration. Science (80-.). *934*, eabb8598.

Horimoto, M., Resnick, M.B., Konkin, T.A., Routhier, J., Wands, J.R., and Baffy, G. (2004). Expression of uncoupling protein-2 in human colon cancer. Clin. Cancer Res. *10*, 6203–6207.

Van Hove, H., Antunes, A.R.P., De Vlaminck, K., Scheyltjens, I., Van Ginderachter, J.A., and Movahedi, K. (2020). Identifying the variables that drive tamoxifen-independent CreERT2 recombination: Implications for microglial fate mapping and gene deletions. Eur. J. Immunol. *50*, 459–463.

Hunter, J.D. (2007). Matplotlib: A 2D graphics environment. Comput. Sci. Eng. *9*, 90–95.

Jahn, H.M., Kasakow, C. V., Helfer, A., Michely, J., Verkhratsky, A., Maurer, H.H.,

Scheller, A., and Kirchhoff, F. (2018). Refined protocols of tamoxifen injection for inducible

DNA recombination in mouse astroglia. Sci. Reports 2018 81 *8*, 1–11.

Ježek, P., Holendová, B., Garlid, K.D., and Jabůrek, M. (2018). Mitochondrial Uncoupling

Proteins: Subtle Regulators of Cellular Redox Signaling. Antioxid. Redox Signal. *29*, 667–

714.

Jha, A.K., Huang, S.C.C., Sergushichev, A., Lampropoulou, V., Ivanova, Y., Loginicheva,

E., Chmielewski, K., Stewart, K.M., Ashall, J., Everts, B., et al. (2015). Network integration

of parallel metabolic and transcriptional data reveals metabolic modules that regulate

macrophage polarization. Immunity *42*, 419–430.

Kim, J.D., Yoon, N.A., Jin, S., and Diano, S. (2019). Microglial UCP2 Mediates

Inflammation and Obesity Induced by High-Fat Feeding. Cell Metab. *30*, 952-962.e5.

Kong, D., Vong, L., Parton, L.E., Ye, C., Tong, Q., Hu, X., Choi, B., Brüning, J.C., and

Lowell, B.B. (2010). Glucose Stimulation of Hypothalamic MCH Neurons Involves KATP

Channels, Is Modulated by UCP2, and Regulates Peripheral Glucose Homeostasis. Cell

Metab. *12*, 545–552.

Levkovitch-Verbin, H., Harris-Cerruti, C., Groner, Y., Wheeler, L.A., Schwartz, M., and

Yoles, E. (2000). RGC death in mice after optic nerve crush injury: Oxidative stress and

neuroprotection. Investig. Ophthalmol. Vis. Sci. *41*, 4169–4174.

Li, Y., Schlamp, C.L., and Nickells, R.W. (1999). Experimental induction of retinal ganglion

cell death in adult mice. Invest. Ophthalmol. Vis. Sci. *40*, 1004–1008.

Libby, R.T., Anderson, M.G., Pang, I.H., Robinson, Z.H., Savinova, O. V., Cosma, I.M.,

Snow, A., Wilson, L.A., Smith, R.S., Clark, A.F., et al. (2005). Inherited glaucoma in

DBA/2J mice: Pertinent disease features for studying the neurodegeneration. Vis. Neurosci.

22, 637–648.

Liu, S., Li, Z.W., Weinreb, R.N., Xu, G., Lindsey, J.D., Ye, C., Yung, W.H., Pang, C.P.,

Lam, D.S.C., and Leung, C.K.S. (2012). Tracking retinal microgliosis in models of retinal

ganglion cell damage. Investig. Ophthalmol. Vis. Sci. *53*, 6254–6262.

Maes, M.E., Schlamp, C.L., and Nickells, R.W. (2017). BAX to basics: How the BCL2 gene

family controls the death of retinal ganglion cells. Prog. Retin. Eye Res. *57*, 1–25.

Man, P.Y.W., Griffiths, P.G., Brown, D.T., Howell, N., Turnbull, D.M., and Chinnery, P.F.

(2003). The Epidemiology of Leber Hereditary Optic Neuropathy in the North East of

England. Am. J. Hum. Genet. *72*, 333–339.

Masuda, T., Sankowski, R., Staszewski, O., and Prinz, M. (2020). Microglia Heterogeneity in

the Single-Cell Era. CellReports *30*, 1271–1281.

Matcovitch-Natan, O., Winter, D.R., Giladi, A., Vargas Aguilar, S., Spinrad, A., Sarrazin, S.,

Ben-Yehuda, H., David, E., Zelada González, F., Perrin, P., et al. (2016). Microglia

development follows a stepwise program to regulate brain homeostasis. Science *353*,

aad8670.

Mckinney, W. (2010). Data Structures for Statistical Computing in Python.

Mishra, P., and Chan, D.C. (2016). Metabolic regulation of mitochondrial dynamics. J. Cell

Biol. *212*, 379–387.

Mitra, K., Wunder, C., Roysam, B., Lin, G., and Lippincott-Schwartz, J. (2009). A

hyperfused mitochondrial state achieved at G1-S regulates cyclin E buildup and entry into S

phase. Proc. Natl. Acad. Sci. U. S. A. *106*, 11960–11965.

Nair, S., Sobotka, K.S., Joshi, P., Gressens, P., Fleiss, B., Thornton, C., Mallard, C., and

Hagberg, H. (2019). Lipopolysaccharide-induced alteration of mitochondrial morphology

induces a metabolic shift in microglia modulating the inflammatory response in vitro and in

vivo. Glia *67*, 1047–1061.

Mac Nair, C.E., Schlamp, C.L., Montgomery, A.D., Shestopalov, V.I., and Nickells, R.W.

(2016). Retinal glial responses to optic nerve crush are attenuated in Bax-deficient mice and

modulated by purinergic signaling pathways. J. Neuroinflammation *13*, 93.

Nakazawa, T., Takahashi, H., and Shimura, M. (2006). Estrogen has a neuroprotective effect

on axotomized RGCs through ERK signal transduction pathway. Brain Res. *1093*, 141–149.

Nuzzi, R., and Caselgrandi, P. (2022). Sex Hormones and Their Effects on Ocular Disorders

and Pathophysiology: Current Aspects and Our Experience. Int. J. Mol. Sci. *23*.

O'Koren, E.G., Yu, C., Klingeborn, M., Wong, A.Y.W., Prigge, C.L., Mathew, R., Kalnitsky,

J., Msallam, R.A., Silvin, A., Kay, J.N., et al. (2019). Microglial Function Is Distinct in

Different Anatomical Locations during Retinal Homeostasis and Degeneration. Immunity *50*,

723-737.e7.

Paolicelli, R., Sierra, A., Stevens, B., Tremblay, M.-E., Aguzzi, A., Ajami, B., Amit, I.,

Audinat, E., Bechmann, I., Bennett, M., et al. (2022). Defining Microglial States and

Nomenclature: A Roadmap to 2030. SSRN Electron. J.

Park, J., Choi, H., Min, J.-S., Park, S.-J., Kim, J.-H., Park, H.-J., Kim, B., Chae, J.-I., Yim,

M., and Lee, D.-S. (2013). Mitochondrial dynamics modulate the expression of pro-

inflammatory mediators in microglial cells. J. Neurochem. *127*, 221–232.

Pecqueur, C., Alves-Guerra, M.-C., Gelly, C., Lévi-Meyrueis, C., Couplan, E., Collins, S.,

Ricquier, D., Bouillaud, F., and Miroux, B. (2001). Uncoupling Protein 2, *in Vivo*

Distribution, Induction upon Oxidative Stress, and Evidence for Translational Regulation. J. Biol. Chem. *276*, 8705–8712.

Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., Blondel, M., Prettenhofer, P., Weiss, R., Dubourg, V., et al. (2011). Scikit-learn: Machine Learning in Python. J. Mach. Learn. Res. *12*, 2825–2830.

Pham, A.H., McCaffery, J.M., and Chan, D.C. (2012). Mouse lines with photo-activatable mitochondria to study mitochondrial dynamics. Genesis *50*, 833–843.

Quigley, H., and Broman, A.T. (2006). The number of people with glaucoma worldwide in 2010 and 2020. Br. J. Ophthalmol. *90*, 262–267.

Rambold, A.S., and Pearce, E.L. (2018). Mitochondrial Dynamics at the Interface of Immune Cell Metabolism and Function. Trends Immunol. *39*, 6–18.

Rodriguez, A.R., de Sevilla Müller, L.P., and Brecha, N.C. (2014). The RNA binding protein RBPMS is a selective marker of ganglion cells in the mammalian retina. J. Comp. Neurol. *522*, 1411–1443.

Ron-Harel, N., Santos, D., Ghergurovich, J.M., Sage, P.T., Reddy, A., Lovitch, S.B., Dephoure, N., Satterstrom, F.K., Sheffer, M., Spinelli, J.B., et al. (2016). Mitochondrial biogenesis and proteome remodeling promotes one carbon metabolism for T cell activation. Cell Metab. *24*, 104.

Rupprecht, A., Bräuer, A.U., Smorodchenko, A., Goyn, J., Hilse, K.E., Shabalina, I.G., Infante-Duarte, C., and Pohl, E.E. (2012). Quantification of Uncoupling Protein 2 Reveals Its Main Expression in Immune Cells and Selective Up-Regulation during T-Cell Proliferation. PLoS One *7*, e41406.

Rupprecht, A., Sittner, D., Smorodchenko, A., Hilse, K.E., Goyn, J., Moldzio, R., Seiler, A.E.M., Bräuer, A.U., and Pohl, E.E. (2014). Uncoupling Protein 2 and 4 Expression Pattern during Stem Cell Differentiation Provides New Insight into Their Putative Function. PLoS One *9*, e88474.

Rupprecht, A., Moldzio, R., Mödl, B., and Pohl, E.E. (2019). Glutamine regulates mitochondrial uncoupling protein 2 to promote glutaminolysis in neuroblastoma cells. Biochim. Biophys. Acta. Bioenerg. *1860*, 391–401.

Salter, M.W., and Stevens, B. (2017). Microglia emerge as central players in brain disease. Nat. Med. *23*, 1018–1027.

Schmittgen, T.D., and Livak, K.J. (2008). Analyzing real-time PCR data by the comparative CT method. Nat. Protoc. 2008 36 *3*, 1101–1108.

Seabold, S., and Perktold, J. (2010). Statsmodels: Econometric and Statistical Modeling with

Python. Proc. 9th Python Sci. Conf.

Sholl, D.A. (1953). Dendritic organization in the neurons of the visual and motor cortices of
the cat. *87*.

Siegert, S., Cabuy, E., Scherf, B.G., Kohler, H., Panda, S., Le, Y.Z., Fehling, H.J., Gaidatzis,
D., Stadler, M.B., and Roska, B. (2012). Transcriptional code and disease map for adult
retinal cell types. Nat. Neurosci. *15*, 487–495.

Smorodchenko, A., Schneider, S., Rupprecht, A., Hilse, K., Sasgary, S., Zeitz, U., Erben,
R.G., and Pohl, E.E. (2017). UCP2 up-regulation within the course of autoimmune
encephalomyelitis correlates with T-lymphocyte activation. Biochim. Biophys. Acta. Mol.
Basis Dis. *1863*, 1002–1012.

Spanier, G., Xu, H., Xia, N., Tobias, S., Deng, S., Wojnowski, L., Forstermann, U., and Li,
H. (2009). Resveratrol reduces endothelial oxidative stress by modulating the gene
expression of superoxide dismutase 1 (SOD1), glutathione peroxidase 1 (GPx1) and NADPH
oxidase subunit (Nox4). J. Physiol. Pharmacol. *60 Suppl 4*, 111–116.

Stogsdill, J.A., Kim, K., Binan, L., Farhi, S.L., Levin, J.Z., and Arlotta, P. (2022). Pyramidal
neuron subtype diversity governs microglia states in the neocortex. Nat. 2022 1–7.

Thion, M.S., Low, D., Silvin, A., Chen, J., Grisel, P., Schulte-Schrepping, J., Blecher, R.,
Ulas, T., Squarzoni, P., Hoeffel, G., et al. (2018). Microbiome Influences Prenatal and Adult
Microglia in a Sex-Specific Manner. Cell *172*, 500-516.e16.

Toda, C., Kim, J.D., Impellizzeri, D., Cuzzocrea, S., Liu, Z.-W., and Diano, S. (2016). UCP2
Regulates Mitochondrial Fission and Ventromedial Nucleus Control of Glucose
Responsiveness. Cell *164*, 872–883.

Tondera, D., Grandemange, S., Jourdain, A., Karbowski, M., Mattenberger, Y., Herzig, S.,
Da Cruz, S., Clerc, P., Raschke, I., Merkwirth, C., et al. (2009). SLP-2 is required for stress-
induced mitochondrial hyperfusion. EMBO J. *28*, 1589–1600.

Ulland, T.K., Song, W.M., Huang, S.C.-C., Ulrich, J.D., Sergushichev, A., Beatty, W.L.,
Loboda, A.A., Zhou, Y., Cairns, N.J., Kambal, A., et al. (2017). TREM2 Maintains
Microglial Metabolic Fitness in Alzheimer's Disease. Cell *170*, 649-663.e13.

Villa, A., Gelosa, P., Castiglioni, L., Cimino, M., Rizzi, N., Pepe, G., Lolli, F., Marcello, E.,
Sironi, L., Vegeto, E., et al. (2018). Sex-Specific Features of Microglia from Adult Mice.
Cell Rep. *23*, 3501–3511.

Vozza, A., Parisi, G., Leonardis, F. De, Lasorsa, F.M., and Castegna, A. (2014). UCP2
transports C4 metabolites out of mitochondria, regulating glucose and glutamine oxidation.
111, 960–965.

Wai, T., and Langer, T. (2016). Mitochondrial Dynamics and Metabolic Regulation. Trends

Endocrinol. Metab. *27*, 105–117.

Waskom, M.L. (2021). seaborn: statistical data visualization. J. Open Source Softw. *6*, 3021.

Williams, P.R., Benowitz, L.I., Goldberg, J.L., and He, Z. (2020). Axon Regeneration in the

Mammalian Optic Nerve. Annu. Rev. Vis. Sci. *6*, 195–213.

Wohl, S.G., Schmeer, C.W., Witte, O.W., and Isenmann, S. (2010). Proliferative response of

microglia and macrophages in the adult mouse eye after optic nerve lesion. Investig.

Ophthalmol. Vis. Sci. *51*, 2686–2696.

Yang, S.-H., Liu, R., Perez, E.J., Wen, Y., Stevens, S.M.†, Valencia, T., Brun-Zinkernagel,

A.-M., Prokai, L., Will, Y., Dykens, J., et al. (2004). Mitochondrial localization of estrogen

receptor.

Yasumoto, Y., Stoiljkovic, M., Kim, J.D., Sestan-Pesa, M., Gao, X.-B., Diano, S., and

Horvath, T.L. (2021). Ucp2-dependent microglia-neuronal coupling controls ventral

hippocampal circuit function and anxiety-like behavior. Mol. Psychiatry 2021 1–13.

Yona, S., Kim, K.-W., Wolf, Y., Mildner, A., Varol, D., Breker, M., Strauss-Ayali, D.,

Viukov, S., Guilliams, M., Misharin, A., et al. (2013). Fate Mapping Reveals Origins and

Dynamics of Monocytes and Tissue Macrophages under Homeostasis. Immunity *38*, 79–91.

Youle, R.J., and van der Bliek, A.M. (2012). Mitochondrial Fission, Fusion, and Stress.

Science (80-.). *337*, 1062–1065.

Zhang, J., Khvorostov, I., Hong, J.S., Oktay, Y., Vergnes, L., Nuebel, E., Wahjudi, P.N.,

Setoguchi, K., Wang, G., Do, A., et al. (2011). UCP2 regulates energy metabolism and

differentiation potential of human pluripotent stem cells. EMBO J. *30*, 4860–4873.

Žuna, K., Jovanović, O., Khailova, L.S., Škulj, S., Brkljača, Z., Kreiter, J., Kotova, E.A.,

Vazdar, M., Antonenko, Y.N., and Pohl, E.E. (2021). Mitochondrial uncoupling proteins

(UCP1-UCP3) and adenine nucleotide translocase (ANT1) enhance the protonophoric action

of 2,4-dinitrophenol in mitochondria and planar bilayer membranes. Biomolecules *11*.

1204

Figure 1.

A

Mouse model for mitochondrial visualization

Cx3Cr1[CreERT2/+]

Cx3cr1 — CreERT2 — 2

Pham[fl/fl]

Rosa26 — loxP — STOP — loxP — mito-Dendra2

→ μG[mito] mouse + Tamoxifen

Rosa26 — mito-Dendra2

D

Mouse model for mitochondrial visualization and manipulation

Cx3Cr1[CreERT2/+]

Cx3cr1 — CreERT2 — 2

Pham[fl/fl]

Rosa26 — loxP — STOP — loxP — mito-Dendra2

UCP2[fl/fl]

Ucp2 — 1 — 2 — loxP — 3 — 4 — loxP — 5 — 6 — 7 — 8

→ μG[mito]UCP2[KO] mouse + Tamoxifen

Rosa26 — mito-Dendra2

Ucp2 — 1 — 2 — 5 — 6 — 7 — 8

B

Microglia-specific mitochondrial label in μG[mito]

mito-Dendra2 | IBA1 | Merge

E

Microglia-specific mitochondria label in μG[mito]UCP2[KO]

mito-Dendra2 | IBA1 | Merge

C

μG[mito] single-cell mitochondrial analysis

mito-Dendra2 | mito-Dendra2 IBA1

Zoom-in

3D-Surface

F

μG[mito]UCP2[KO] single-cell mitochondrial analysis

mito-Dendra2 | mito-Dendra2 IBA1

Zoom-in

3D-Surface

G

Mitochondrial network connectivity

n.s. — Median mitochondrial volume

n.s. — Number of organelles

H

Mitochondrial content

n.s. — % Mitochondria volume/ microglia

I

Mitochondrial localization

n.s. — % Mitochondria perinuclear

Naive μG[mito]

Naive μG[mito]UCP2[KO]

Figure 2.

A *Optic nerve crush (ONC)*

Optic nerve
Retina
Lens
Forceps
RGC axons
Cornea

B *ONC elicits IPL microglia response 5 days post-injury*

Naive   ONC   IBA1

OPL
IPL
GCL

C *Microglial Ucp2 transcript increases upon ONC*

Ucp2

Naive
ONC

Fold change (2^{ddCq})

*

μG^{mito} Naive   **μG^{mito} ONC**   **Analysis**

D (3D-trace)

G

J *Sholl intersections*

Naive
ONC

Sholl intersections
Radius (μm)

***

E (CD68 surface)

H   CD68
IBA1

K *CD68 volume*

% CD68 volume

***

L *CD68 localization*

% CD68 volume in periphery

***

F (Mitochondria surface)

I   mito-Dendra2
IBA1

M *Mitochondrial network connectivity*

Median mitochondrial volume

***

Number of organelles

***

N *Mitochondrial content*

% Mitochondria volume/ microglia

*

O *Mitochondrial localization*

% Mitochondria perinuclear

***

Figure 3.

μG^mitoUCP2^KO Naive **μG^mitoUCP2^KO ONC** **Analysis**

3D-trace

C *Sholl intersections*

CD68 surface

CD68
IBA1

F *CD68 volume*   **G** *CD68 localization*

Mitochondria surface

mito-Dendra2
IBA1

J *Mitochondrial network connectivity*

K *Mitochondrial content*   **L** *Mitochondrial localization*

Figure 4.

Figure 5.

Mitochondrial parameter PCA

Separated by condition

Conditions separated by sex

Figure 6.

A *Mitochondrial parameter PCA*

μG^mito UCP2^KO ONC

μG^mito ONC

♂ Day 5
♂ Day 14
♀ Day 5
♀ Day 14

B *Male μG^mito UCP2^KO mitochondrial hyperfusion*

Organelles per condition

C *Metabolic and stress markers*

OXPHOS — Glycolysis — Stress

Pgc1a   Nrf1   Hif1a   Sod1   Cd68

D μG^mito vs μG^mito UCP2^KO

Naive μG^mito
Naive μG^mito UCP2^KO

E μG^mito

Naive
ONC

F μG^mito UCP2^KO

Naive
ONC

G *RGC loss*

♂

**
n.s.

μG^mito   μG^mito UCP2^KO

Day 5
Day 14

H *RGC loss*

♀

n.s.
n.s.

μG^mito   μG^mito UCP2^KO

Figure S1

A
Candidate mitochondrial genes enriched in retinal microglia

Myc
Tmem173
Rab32
Ucp2
Bcl2a1b
Bcl2a1a
Bcl2a1d
Coro1a

PR   H   BP   Am   RGC   μG

Normalized expression value
1 0.8 0.6 0.4 0.2 0

B
Relative Ucp2 expression among retinal cell types

Ucp2

Relative expression

PR   H   BP   Am   RGC   μG

C
Tamoxifen injection timeline

*Cx3Cr1*CreERT2/+/Phamfl/fl or
*Cx3Cr1*CreERT2/+/Phamfl/fl/UCP2fl/fl

Tamoxifen
1 2 3   > 3 weeks
Day
Adult

μG^mito

μG^mitoUCP2^KO

FACS for Dendra2^+ microglia

D
Ucp2 transcript level by RT-qPCR

***

μG^mito
μG^mitoUCP2^KO

Fold change (2^ddCq)

E
Experimental design for primary culture

*Cx3Cr1*CreERT2/+ or
*Cx3Cr1*CreERT2/+/UCP2fl/fl

P0-P3

4-OHT or vehicle

Mild trypsinization

10 days   3 days

Microglia → Protein

3-5 cortices

Mixed glial culture
IBA1

F
Western blot analysis

kDA

← UCP2
33.4 kDa

← βactin
42 kDa

Microglia

Recombinant UCP2
Cx3Cr1^CreERT2/+
Cx3Cr1^CreERT2/+/UCP2^fl/fl
Cx3Cr1^CreERT2/+
Cx3Cr1^CreERT2/+/UCP2^fl/fl

Vehicle   4-OHT

G
Quantification of Western blot bands

Microglia

Relative UCP2/βactin
1
0.75
0.5
0.25
0

Vehicle   4-OHT

Cx3Cr1^CreERT2/+
Cx3Cr1^CreERT2/+/UCP2^fl/fl
Cx3Cr1^CreERT2/+
Cx3Cr1^CreERT2/+/UCP2^fl/fl

Figure S2

A *Progressive retinal ganglion cell loss after optic nerve crush (ONC)*

Naive | Day 1 | Day 5 | Day 14
ONC — RBPMS

B *IPL microglia after ONC or sham surgery*

Naive | Day 1 | Day 5 | Day 14
ONC — IBA1
Sham

C *IPL microglia Sholl intersections*

Naive | Day 1 | Day 5 | Day 14
Intersections — Radius (µm)

Naive | ONC | Sham

D *Representative CD68 immunostaining*

Naive | ONC Day 5
IBA1 CD68

E *CD68 volume in IPL microglia*

%CD68 volume
Naive | ONC | Sham
Naive — 1 5 14 — 1 5 14 days

F *OPL microglia after ONC or sham surgery*

Naive | Day 1 | Day 5 | Day 14
ONC — IBA1
Sham

G *OPL microglia Sholl intersections*

Naive | Day 1 | Day 5 | Day 14
Intersections — Radius (µm)

Naive | ONC | Sham

H *Representative CD68 immunostaining*

Naive | ONC Day 5
IBA1 CD68

I *CD68 volume in OPL microglia*

%CD68 volume
n.s.
Naive | ONC | Sham
Naive — 1 5 14 — 1 5 14 days

Figure S3

A

PC loadings from seven parameters

	PC 1	PC 2
% Mitochondria volume/microglia	0.220362	0.629862
Median mitochondria volume	-0.34296	0.514091
% CD68 peripheral	0.13551	-0.36167
Number of organells	0.378888	-0.29817
% Mitochondria perinuclear	0.456013	0.255364
Sholl index	0.470127	0.227415
% CD68 volume	0.492882	-0.04842

Sholl index and CD68 parameter PCA

B *Separated by condition*

C *Conditions separated by sex*

D

E

PC loadings from mitochondria PCA

	PC 1	PC 2
% Mitochondria volume/microglia	0.123798	0.7945075
Median mitochondrial volume	-0.576187	0.4279728
% Mitochondria perinuclear	0.5185627	0.4166628
Number of organelles	0.6194963	-0.109495

F *Mitochondrial connectivity*

Legend:
- Naive µG^mito (gray)
- ONC µG^mito (blue)
- Naive µG^mito UCP2^KO (light yellow)
- ONC µG^mito UCP2^KO (gold)

G *Mitochondrial content*

H *Mitochondrial localization*

Figure S4

Mitochondrial parameter PCA

A

B

PC loadings for metabolic and stress genes

	PC 1	PC 2
Pgc1α	0.60227	0.10366
Nrf1	0.50238	0.06933
Hif1α	0.48479	0.4826
Sod1	-0.2558	0.67006
Cd68	-0.2906	0.55005

C

Naive microglia similar between sexes in μG^mito and μG^mitoUCP2^KO

D

Relative expression of male microglia metabolic genes

E

Relative expression of female microglia metabolic genes

F

Male μG^mitoUCP2^KO ONC microglia greater stress markers

Table S1. Primer sequences.

Gene	Primer	Sequence	Product length
GAPDH	FW	ACAGCAACTCCCACTCTTC	131 bp
	REV	CATTGTCATACCAGGAAATGAGC	
IBA1	FW	ATGAGGATCTGCCGTCCAAAC	114 bp
	REV	CAAGTTTCTCCAGCATTCGCTTC	
GFAP	FW	ATGAGGATCTGCCGTCCAAAC	243 bp
	REV	CAAGTTTCTCCAGCATTCGCTTC	
UCP2	FW	AAGTGTTTCGTCTCCCAGCC	188 bp
	REV	GGGACCTTCAATCGGCAAGA	
PGC1a	FW	CGTGTTCCCGATCACCATATTC	102 bp
	REV	GGTGTCTGTAGTGGCTTGATTC	
NRF1	FW	GGTGAAATAAGCCTCCCGATAG	110 bp
	REV	TGAGGCAGTTTAGACAGAATGG	
CD68	FW	CCTCTGTTCCTTGGGCTATAAG	105 bp
	REV	ATTGAGGAAGGAACTGGTGTAG	
HIF1a	FW	AGGATGAGTTCTGAACGTCGAAAAG	208 bp
	REV	CACTGTCTAGACCACCGGCA	
SOD1	FW	CACTTCGAGCAGAAGGCAAGC	186 bp
	REV	ATGCCTCTCTTCATCCGCCG	

Table S2. Antibody table.

Antibody	Dilution	Source	Catalog No.	RRID
Rabbit anti-IBA1	1:500	GeneTex	GTX100042	AB_1240434
Goat anti-IBA1	IHC: 1:250 WB: 1µg/ml	Abcam	ab 5076	AB_2224402
Rat anti-CD68	1:500	Bio-Rad	MCA1957	AB_322219
Rabbit anti-RBPMS	1:200	Abcam	ab194213	AB_2920590
UCP2	1:1000	Pohl Lab, purified at Pineda	NT-Tier2	n/a
β-actin	1:5000	Sigma	A5441	AB_476744
Goat anti-rabbit IgG (H+L) Alexa Fluor 568	1:2000	ThermoFisher	A11036	AB_10563566
Goat anti- rat IgG (H+L) Alexa Fluor 647	1:2000	ThermoFisher	A21247	AB_141778
Donkey Anti-Guinea Pig IgG (H+L) Alexa Fluor 647	1:1000	Jackson ImmunoResearch	706-605-148	AB_2340476
Donkey anti-rabbit IgG (H+L) Alexa Fluor 568	1:2000	ThermoFisher	A10042	AB_2534017
Donkey anti-goat IgG (H+L)	1:2000	ThermoFisher	A11057	AB_2534104
Donkey anti-goat Alexa Fluor 568 IgG (H+L) Alexa Fluor 488	1:2000	ThermoFisher	A11055	AB_2534102
Donkey anti-rabbit IgG (H+L) Alexa Fluor 488	1:2000	ThermoFisher	A21206	AB_2535792
Donkey anti-Rat IgG (H+L) Antibody DyLight 650	1:2000	ThermoFisher	SA5-10029	AB_2556609
Donkey anti-Rat IgG (H+L) Alexa Fluor 488	1:2000	ThermoFisher	A21208	AB_2535794
Donkey anti-rabbit IgG (H+L) Alexa Fluor 647	1:2000	ThermoFisher	A31573	AB_2536183